A Meta-analysis of Perceptual Pseudoneglect reveals the Functional Anatomy of Perceptual

Judgements

Abbreviated title: The Functional Anatomy of Perceptual Pseudoneglect

Authors: Mura Abdul-Nabi[1], Matthias Niemeier[1,2]
[1] Department of Psychology Scarborough
University of Toronto
Scarborough, ON, Canada, M1C1A4
[2] Centre for Vision Research
York University
Toronto, ON, Canada, M4N3M6
13
14   Correspondence concerning this article should be addressed to Matthias Niemeier at
15   m.niemeier@utoronto.ca

17   **Number of pages:** 28
18   **Number of figures:** 4
19   **Number of words:** Abstract: 210; Introduction: 999; Discussion: 1615
20
21   **Conflict of interest statement:** The authors declare no competing financial interests.
22
23   **Acknowledgement:** We thank Kathleen Pirog Revill for making the imaging data available to be
24   included in this study.
25
26   **Keywords:** spatial neglect, attentional bias, line bisection, landmark task, greyscales task, grating-scales
27   task, fMRI

28
29
30

Abstract

Major evidence for a right-hemisphere dominance of the brain in spatial and/or attentional tasks comes from lesion studies in patients with spatial neglect. However, the neuroanatomy of the different forms of neglect remains a matter of debate, and it remains unclear how dysfunctions in neglect relate to intact processes. In the healthy brain, perceptual pseudoneglect has been considered to be a phenomenon complementary to specific subtypes of neglect as observed in paradigms such as the line bisection task. Therefore, the current study investigated the intact functional anatomy of perceptual pseudoneglect using a meta-analysis to compensate for some of the limitations of individual imaging studies. We collated the data from 24 articles that tested 952 participants with a range of paradigms (landmark task, line bisection, grating-scales task, and number line task) obtaining 337 foci. Using Activation Likelihood Estimation (ALE) we identified a right-hemisphere biased network of cortical areas, including superior and intraparietal regions, the intraoccipital sulcus together with other occipital regions, as well as inferior frontal areas that were associated with perceptual pseudoneglect in partial agreement with lesion studies in patients with neglect. Our study is the first meta-analysis on the mechanisms underlying perceptual judgments which have been shown to give rise to perceptual pseudoneglect.

Introduction

Spatial neglect is a set of spatial cognitive deficits that have been summarized as inabilities to report, respond, or orient to stimuli contralateral to the side of a brain damage (e.g., Heilman et al., 1993). Neglect is more commonly observed after right than left brain lesions, and thus it is an important indicator for a right-hemisphere dominance of the neural mechanisms underlying visual attention and spatial awareness. As such neglect has garnered significant interest in the literature. Yet, several questions remain unresolved. Crucially, it is unclear how neglect symptoms map onto brain damage.

A recent meta-analysis of lesion studies in neglect patients collated a list of best-practice criteria for lesion-symptom mapping (Moore et al., 2023; also see Bates et al., 2003; de Haan and Karnath, 2018; Sperber and Karnath, 2017; Thiebaut de Schotten et al., 2014) showing none of the relevant studies attained a perfect score. Also, depending on the score, different lesion patterns emerged with varying core lesions in the supramarginal gyrus, the superior longitudinal fasciculus, the insular cortex, or the postcentral gyrus. Furthermore, the analysis found substantial differences in functional anatomy between different subtypes of neglect. Also consistent with the heterogeneity of neglect, lesion patterns have been found to vary as a function of diagnostic tests (Molenberghs et al., 2012).

To complicate matters functional deficits and brain plasticity after brain lesions interact in a complex and difficult-to-observe manner (Pascual-Leone et al., 2005). For example, extinction, a deficit akin to neglect that affects contralesional stimuli presented together with competing ipsilesional stimuli, does not occur immediately after simulated brain lesions induced by inhibitory continuous theta-burst stimulation (Vesia et al., 2015). Instead, the authors found

extinction to arise several minutes after stimulation, arguably resulting from short-term plasticity

due to interactions between the two hemispheres. Interhemispheric interactions also seem to play

an essential role in neglect in general (Koch et al., 2008; also see Corbetta & Shulman, 2011) as

well as in related asymmetries in the intact brain (Le et al., 2015). This suggests that behavioural

impairments and the site of damage do not necessarily relate to one another in a straight-forward

manner, contrary to one of the central assumptions of lesion studies (Caramazza, 1984).

Given these inherent challenges of lesion studies, a comprehensive picture of the

functional anatomy of neglect calls for comparisons between research in patients with imaging

studies in healthy participants to estimate to which extent lesion patterns in neglect agree with

the anatomy of intact processes that can be assumed to be affected in spatial neglect. One such

set of functions involve perceptual judgements about visuo-spatial features of horizontal lines or

bars such as length (e.g., Harvey & Milner, 1995), luminance (Mattingley et al., 1994), or spatial

frequency (Niemeier et al., 2007). These tasks have been shown to yield comparable visual field

differences in the intact brain called perceptual pseudoneglect (Chen et al., 2020) or attentional

bias (Bultitude et al., 2006; McCourt et al., 2005; Singh et al., 2011). Furthermore, the

differences seem to reflect approximately complementary, although much less severe,

asymmetries compared to deficits found in neglect patients (Bowers & Heilman, 1980; McCourt

& Jewell, 1999) when performing line length judgements in the manual line bisection task that is

commonly used in clinical assessments (e.g., Jewell & McCourt, 2000).

Line bisection and other perceptual judgment tasks have been employed in several

imaging studies with a variety of designs. However, most of these studies used simple contrasts

with only one experimental and one control condition (see Table 1) thereby bearing a greater risk

of false positive and false negative errors in the identification of relevant brain areas. This is

particularly an issue for perceptual judgment paradigms given that they permit test participants to

use a variety of tasks strategies.

A better approach is to design a control condition with physically identical stimuli that

are used with different task instructions (e.g., Revill et al., 2011). However, as a potential

limitation of this approach, it is possible that brain areas that are critical for some function or task

A, might nevertheless yield a stronger BOLD signal in some other task B. This way, subtracting

task B activity from task A activity would potentially remove an essential area from the results.

For example, Filimon and colleagues (2009) have found that the frontal eye fields activate more

during reach tasks than eye movement tasks. Thus, subtracting BOLD signal associated with

reaching from the BOLD signal associated with eye movements could obscure eye movement-

related activation in the frontal eye fields even though they are critically involved in eye

movements.

A third approach then is to use more complex experimental designs for better

experimental control. Benwell et al. (2014) used a factorial design for an EEG experiment with

landmark tasks with long and short lines and with a control task that required bisections to be

detected. To our knowledge a factorial design has not been used in an fMRI experiment on the

attentional bias. Chen et al. (2020) used partial-least squares to model a cognitive vector based

on four tasks that were incrementally more likely to involve brain regions to do with perceptual

judgments that yield pseudoneglect biases. However, not all perceptual judgment paradigms lend

themselves to such a set of tasks. The arguably simplest way to attain better control would be a

conjunction analysis where multiple contrasts of experimental and control tasks are looked at

together such that confounds of one contrast are compensated for by other contrasts (Price &

Friston, 1997). To our knowledge to date no pseudoneglect study has used conjunction analysis.

However, it can be assumed that a meta-analysis, although different from a conjunction

analysis, collates multiple contrasts from potentially different paradigms (depending on the

paradigms of the included studies) and this way potentially discards some of the suspected false

positive and false negative errors of the included studies. Therefore, the aim of the present study

was to conduct a meta-analysis of perceptual judgments tasks that are known to yield perceptual

pseudoneglect to improve overall validity. Furthermore, the question was whether the resulting

functional neuroanatomy would be mirrored in patients exhibiting line bisection deficits after

lesions along the intraparietal and intraoccipital sulci (IPS and IOS), medial occipital cortex and

anterior portions of the middle frontal gyrus (MFG; see meta-analysis in Molenberghs et al.,

2012).

132   **Materials and Methods**

To ensure a comprehensive article list was obtained, imaging studies using perceptual judgments

to measure attentional bias / pseudoneglect published on or prior to March 10, 2025 were

obtained by searching the PubMed (https://pubmed.ncbi.nlm.nih.gov/), PsycInfo

(https://www.apa.org/pubs/databases/psycinfo), and Web of Science

(https://www.webofknowledge.com/) databases using the following keyword search: "(line

bisection OR landmark task OR pseudoneglect OR greyscales task OR grayscales task OR

grating-scales task) AND (fMRI OR PET)." Reference lists were also checked to avoid missing

relevant articles. A repetition of the search on November 29, 2023, yielded no additional relevant

publications.

Articles were selected according to the following inclusion criteria: 1) studies were

published in English, 2) participants were healthy, neurotypical adults (18-65 years old), 3)

studies assessed attentional biases using either horizontal line bisection, landmark, greyscales, or

gratingscales tasks, 4) fMRI or PET was used to assess neural activity (one study using SPECT

was included as well; additional searches revealed that no other relevant study used this method),

and 5) Montreal Neurological Institute (MNI) or Talairach standardized coordinates of activation

were provided. The exclusion criteria were as follows: 1) participants with a history of brain

damage or other clinical populations, 2) clinical or drug related interventions, 3) studies on

children (< 18 years old) or seniors (65+ years old), and 4) meta-analyses or literature reviews as

evidenced based on inspecting the articles' titles or content. Additionally, we only included those

contrasts that compared task activation to a control or baseline condition and that imaged the

entire brain. Using these criteria, we selected 24 articles consisting of 952 subjects and 337 foci

of activation to be included in this meta-analysis. An inter-rater agreement of r = 0.93 was

obtained between the authors' selections of studies to be included. Any discrepancy was then

resolved through closer inspection of the articles in question. Figure 1 details our literature

search and selection process.

Figure 1. Outline of the literature search process.

Upon finalizing the list of studies, we converted all coordinates of activation reported in

Talairach standardized space to MNI space using the mni2tal transform

(https://bioimagesuiteweb.github.io/webapp/mni2tal.html). These data were then submitted to an

omnibus meta-analysis (24 articles consisting of 952 subjects and 337 foci). Next, we

categorized studies based on the paradigms used to assess pseudoneglect to allow for comparison

among tasks. These included: landmark tasks, physical line bisection tasks, and number line

bisection tasks. The meta-analysis for the Landmark task included 16 studies involving 810

subjects and 173 foci. Here, we selected contrasts that employed baseline/control conditions. ~~The~~

~~meta-analysis for the Line Bisection task included 6 studies involving 94 subjects and 44 foci.~~

~~Lastly, the exploratory meta-analysis for the Number Line task included 2 studies of 58 subjects~~

~~and 52 foci.~~ Multiple contrasts from a single experiment were pooled together to avoid greater

weight being given to multiple analyses with the same set of participants (Müller et al., 2018). A

complete list of studies and their attributes can be found in Table 1.

For each meta-analysis, data were analyzed using the Activation Likelihood Estimation

(ALE) method outlined in Brainmap GingerALE Version 3 (http://brainmap.org/ale/). The

algorithm accounts for spatial uncertainty by treating individual foci as centers of 3D Gaussian

probability distributions with a width that depends on the sample size of the respective study

(i.e., the width is inversely related to the square root of the sample size). Summing over the

Gaussians then results in a probability distribution that is superimposed with a grey matter mask.

Next, a permutation procedure computes nonparametrically a null-distribution of random spatial

association between experiments to identify statistically significant convergence of activation

foci. Here, we employed a cluster level approach with an uncorrected P value of $p < 0.001$ and a

cluster-level family-wise error (FWE) set at $p < 0.05$ for the omnibus analysis. An additional

meta-analysis for the landmark data alone was slightly below the number of studies

recommended for ALE (Eickhoff et al., 2016). To avoid unstable results we chose a more

conservative p-value of 0.001 for the cluster-level FWE. Threshold permutations were set at

1000, as per recommendations outlined in Eickhoff et al. (2012). Results of the ALE analyses

were visualized using MRIcron (http://www.cabiatl.com/mricro/mricron/) and BrainNet Viewer

(https://www.nitrc.org/projects/bnv/; Xia et al., 2013). BrainNet Viewer was also used to

visualize activation foci for the line bisection (6 studies) and the number line task (2 studies) and

plotted together with the activation maps for grating-scales task (Chen et al., 2020). Given the

small numbers of these studies only qualitative syntheses and comparisons instead of meta-

analyses were used.

Table 1. Studies included in the meta-analysis.

Publication	N	Contrast	Foci	Analysis
Badzakova-Trajkov et al. (2010)	84	LM > Bisector Detection	7	Omnibus; Landmark
Badzakova-Trajkov et al. (2011)	199	LM > Bisector Detection	9	Omnibus; Landmark
Cai et al. (2013)	31	LM > Bisector Detection (includes L and R language dominance)	19	Omnibus; Landmark
Cavezian et al. (2011)	9	LM w/ greyscles > Bisector Detection	27	Omnibus; Landmark
		LB w/ greyscales > Saccade Task	14	Omnibus; Line Bisection
Chen et al. (2020)	21	Latent variable modelling a continuum of the grating-scales task/high spatial frequency vs. low spatial frequency vs. frame condition vs. fixation	21	Omnibus
Cicek et al. (2009)	11	LM > Bisector Detection	6	Omnibus; Landmark
		LB > Move cursor to edge	3	Omnibus; Line Bisection
Fink et al. (2000a)	12	LM > Bisector Detection	7	Omnibus; Landmark
Fink et al. (2000b)	12	LM > Bisector Detection	5	Omnibus; Landmark
Fink et al. (2001)	11	LMV + LM > Bisector Detection	13	Omnibus
Fink et al. (2002)	12	LM + LL > Instructions	12	Omnibus
Fink et al. (2003)	12	LM (eye movements) > Bisector Detection	9	Omnibus; Landmark
Flöel et al. (2005)	7	LM > Overlapping waveform line detection	26	Omnibus; Landmark
Hurwitz et al. (2011)	16	DB > Saccade task	3	Omnibus; Line Bisection
Liu et al. (2019)	34	Number LB > Number Recognition Task	37	Omnibus; Number Line
		LB > Dot Position Recognition Task	11	Omnibus; Line Bisection
Marshall et al. (1997)*	9	LM > Line Flicker Task	1	Omnibus; Landmark
Plewan et al. (2012)	24	LM Mueller Lyer > Rest	5	Omnibus; Landmark

Revill et al. (2011)	26	LM > Rest	15	Omnibus; Landmark
		LM > search task	13	Omnibus
Sahan et al. (2019)	24	Number LB > Shape Recognition Task (Encoding & Decision-Making Phases)	26	Omnibus; Number Line
Schuster et al. (2017)	15	LM (Version C) > Bisector Detection (Session 1 & 2)	31	Omnibus; Landmark
Weidner et al. (2007)	15	LM Mueller Lyer > Luminance Detection	2	Omnibus; Landmark
Weiss et al. (2000)	12	Near + Far LB > Pointing Task	6	Omnibus; Line Bisection
Weiss et al. (2003)	12	Near + Far LM > Near + Far LB	16	Omnibus
		Near + Far LB > Near + Far LM	7	Omnibus; Line Bisection
Zago et al. (2016)	293	LM > Saccade Task	11	Omnibus; Landmark
Zago et al. (2017)	51	LM > Saccade Task	10	Omnibus; Landmark

196   LM: Landmark task; LB: Line Bisection task; LMV: Vertical Landmark Task; LL: Line Length

197   Task; DB: Dynamic Bisection Task. *The study employed SPECT.

199   **Results**

The functional anatomy of the perceptual attentional bias: omnibus analysis

Figure 2 shows the results of the omnibus meta-analysis, including all 24 studies. Significant

overlap in activation was found in posterior regions, including the anterior and posterior

intraparietal sulcus (IPS) that was more prominent in the right than the left hemisphere and

reached into the superior parietal lobule (SPL). Exclusively right hemisphere overlap was found

in the intraoccipital sulcus (IOS). Likewise, exclusive right-brain overlap was found in frontal

regions for the precentral sulcus at the intersection with the superior frontal sulcus, probably the

frontal eye field (FEF) and at the level of the inferior frontal cortex (IFC). In addition, there was

right dorsolateral prefrontal cortex (DLPFC) activity and dorso-medial activity near the pre-

supplementary motor area. Overlap in the anterior insula (aIns) was bilateral. Notably, several

studies reported foci in the cerebellum but there was no significant overlap. We confirmed these

observations statistically using the AAL3 atlas (Rolls et al., 2020) to create independently

defined masks. In terms of posterior regions, as defined by the atlas, we found significantly more

(or exclusive) right-brain involvement for the SPL, IPS (i.e., the inferior parietal lobule, not

including supramarginal and angular gyrus), as well as the superior and middle occipital gyrus

(t's≥10.85, p<0.001) with spurious involvement of the right but not left postcentral and angular

gyrus (t's≥3.84, p's≤0.001). As for AAL3-defined frontal regions we found exclusive right

hemisphere involvement of the superior frontal gyrus, middle frontal gyrus, and precentral gyrus

(t's≥4.27, p's≤0.001) and right-dominant involvement for the SMA as well as the opercular and

(spuriously) triangular part of the inferior frontal gyrus, both of which were reaching into the

aIns (t's≥4.50, p's≤0.001). The remainder of the insula showed no significant left- vs. right-side

difference (t=0.77, p=0.441).

Figure 2. Results of the omnibus meta-analysis. A. Horizontal slices. B. Lateral, dorsal, and
posterior views.
227

Influence of task on the cortical network of the perceptual attentional bias

Perceptual forms of attentional biases have been captured with several different kinds of tasks,

although not all these tasks have been frequently used in imaging studies such that more

commonly used paradigms have a stronger influence on our omnibus results. To examine the

effect of task on brain activation we inspected the data for specific tasks, respectively. Only for

the landmark task there were sufficient numbers of studies to conduct a separate ALE analysis

(although with a stricter cluster criterion). For the other paradigms we examined the results in a

qualitative manner with the caveat that the robustness the respective activation patters has yet to

be confirmed. Figure 3A shows significant overlap for studies using the landmark task. The

resulting cortical network included no activation of right FEF and DLPF cortex and left aIns, in

part due to our more conservative criterion for cluster-level FEW. Otherwise, the pattern was

quite similar to that found with the omnibus analysis. While that similarity was expected given

that the landmark experiments constituted the largest portion of the omnibus analysis, what was

important, the landmark network was remarkably similar to the activation pattern for grating-

scales task, i.e., for the only other existing study that used a purely perceptual paradigm (Chen et

al., 2019; Fig. 3B).

244
245 **Figure 3.** Task-specific analyses. A. Meta-analysis of the landmark task data. B. Partial-least
246 square analysis of the grating-scales task (Chen et al., 2020). C. Overlay plot of the line bisection
247 foci (cerebellar foci not shown). D. Overlay plot of the mental number line foci (calcarine focus
248 not shown).
249

250   In addition, to qualitatively examine the influence of action vs. imagery we plotted the

251 foci reported in the six line bisection studies (Fig.3C) and in the two mental number line studies

252 (Fig.3D), respectively. This suggested more bilateral activation with relatively more overlap in

253 the FEFs for manual line bisection as well as more overlap in SPL and DLPFC for mental

254 number lines.

256   **Discussion**

With the current meta-analysis, we intended to investigate the intact functional anatomy

underlying perceptual judgment tasks that have been found to give rise to pseudoneglect. We

found that across several paradigms (landmark task, line bisection, grating-scales task, and

number line task) significant right hemisphere-biased overlap emerged for posterior regions: the

IPS, the SPL, and the IOS. Furthermore, we observed right-biased overlap in the FEF, preSMA,

DLPFC, caudal IFC and the anterior insula of the right hemisphere. As we will argue these

results add to our understanding of right-dominant functions of perceptual judgments that give

rise to perceptual pseudoneglect as well as subtypes of spatial neglect.

However, as one caveat it should be noted that the present meta-analysis is likely

imperfect. Specifically, the study included far more studies using the landmark task than other

paradigms. Such a bias in existing data could result in imbalanced results that would be avoided

in a systematically planned conjunction analysis. Despite this imbalance we found that results for

the latter type of studies alone were in good agreement with results from a multivariate study

using the grating-scales task (Chen et al., 2020; Fig. 3A vs. B).

This consistency also argues against the possibility that the present results merely

reflected unspecific activation of the "task-positive" network (e.g., Fox et al., 2005). That is, the

pattern of activity reported by Chen and colleagues (2020) was associated with perceptual

judgments about the high spatial frequency components of their stimuli relative to a very similar

task pertaining to low spatial frequencies. In addition, the present results show a clear asymmetry

that was not reported for the task-positive network. In sum, we argue that the current results do

serve as a reasonably accurate approximation of the true functional network of brain areas

associated with perceptual pseudoneglect tasks. Therefore, in the following paragraphs we will

proceed to comparing the current results with those of other methods to then propose a new

framework of the neuro-computations underlying perceptual pseudoneglect.

The results of our meta-analysis are partially consistent with the lesion patterns in stroke

patients exhibiting line bisection deficits, i.e., lesion sites in the IPS and IOS, medial occipital

cortex, and the MFG as reported by Molenberghs et al. (2012).

In contrast to Molenbergh et al. (2012), the current study found the SPL, IFC and anterior

insula to be associated with line bisection deficits. This could indicate that these areas merely

coactive in imaging studies without playing a critical role in perceptual judgments. On the other

hand, it is possible that lesion studies overlook critical areas, either because these studies are

administratively challenging and, thus, often have to deal with methodological imperfections

(Moore et al., 2023), or because areas that are critically involved in perceptual judgments in the

intact brain are rapidly substituted after lesions (e.g., Pascual-Leone et al., 2005). E.g., some

evidence suggests that such neural plasticity starts to play out within minutes of a brain damage

(Vesia et al., 2015) which would be outside the temporal scope of lesion studies, potentially

resulting in false negative errors.

In sum, we can argue that the current findings offer a largely valid view of the areas that

are critically involved in perceptual judgments. Furthermore, this activation pattern can be

approximately matched to source localization results from an electroencephalography (EEG)

study: Le et al. (2015) observed that activation differences between grating-scales conditions that

either do or do not capture pseudoneglect first emerge within roughly bilateral parieto-occipital

cortex starting some 74 to 148 ms after stimulus onset and again from 210 to 242 ms, stretching

across late stages of the visually evoked P1 component and past the P2, respectively. These

sources of activation might be associated with IOS in both hemispheres for the grating-scales

task (Fig. 3B) but could involve other visual areas in other tasks (e.g., lateral occipital cortex in

the landmark task, Fig. 3A). From 242 to 280 ms then Le et al. (2015) observed clearly right

lateralized activation over right posterior cortex (n.b., Le et al., 2015, located the source in the

temporo-parietal junction, probably due to imprecise EEG source localization). Based on the

current results we believe that this signal originates from right IPS and SPL. The same regions

are co-active together with additional sources in interior frontal regions, arguably IFC and

anterior insula, from 280 to 340 ms before activation becomes confined to those frontal regions

from 340 to 394 ms (Le et al., 2015; for the role of the second component of the superior

longitudinal fasciculus and its parieto-frontal connectivity during the line bisection task also see

Thiebaut de Schotten et al., 2005; Vallar et al., 2014).

This putative temporo-spatial pattern suggests a network that first activates posterior

portions of the dorsal attention network before involving anterior portions of the ventral

attentional network (Corbetta & Shulman, 2002; Vossel et al., 2014). As such it could be viewed

as consistent with the activation-orientation hypothesis of pseudoneglect (Bultitude & Aimola

Davies, 2006; Reuter-Lorenz et al., 1990; see Anderson, 1996, for a different account of an

underlying attentional bias) according to which a right hemisphere dominance for perceptual

judgment tasks leads to greater activation of the right brain which then causes attention to orient

to the left visual field (although see Chen et al. 2014; Chen et al. 2019; Karnath et al., 1998).

Even so, it remains unclear what the right-hemisphere dominance entails and what causes

leftward oriented attention to lead to biased perceptual judgments. Instead, here we suggest a

more detailed framework informed by the current results.

A framework for perceptual judgments and pseudoneglect

Considering the functional network observed in the current study together with the EEG data

obtained for the grating-scales task (Le et al., 2015) we tentatively propose the following neuro-

computational steps that might underlie perceptual pseudoneglect. What is crucial in this context,

we argue that these steps should follow a process of perceptual decision making given the fact

that perceptual pseudoneglect is based on perceptual judgments:

1. Visual analysis. Perceptual judgment tasks trigger visual analysis of the stimulus

material in approximately bilaterally activated striate and extrastriate areas.

2. Integration and magnitude estimation. Next, retinotopic areas in IPS and SPL

activate especially in the right hemisphere (presumably after interhemispheric competition; Le et

al., 2015). Importantly, right IPS and SPL areas are known to respond not only to the

contralateral visual half field but also to larger portions of the ipsilateral half field (Sheremata &

Silver, 2015). As such they would be suitable to form integrated representations of the stimulus

(although with imperfect isometry of the left and right visual field) as a whole while preserving

its topography.

2a. Feedback. It appears reasonable to assume that these parietal representations yield

activation in striate and extrastriata areas thereby utilizing their high spatial resolution (e.g.,

Williams et al., 2008; Chambers et al., 2013; Monaco et al., 2017).

3. Magnitude comparison. The integrated representation of the stimulus then serves to

compute magnitudes of the task-relevant feature of the stimulus (length of landmark stimuli,

luminance of greyscales, spatial frequency of grating-scales etc.; Walsh, 2003) and to compare

that magnitude for the two sides of the stimulus along its horizontal extent within IPS and SPL in

the right hemisphere.

4. Perceptual decision and response implementation. Finally, a perceptual decision about the

left vs. the right side of the stimulus is made involving the anterior insula, given that IFC and

insula are associated with forming an action-oriented representation in preparation of a response

(e.g., Demanet et al., 2016; Myers et al., 2017). This decision would also explain activity in the

preSMA that is known to be involved in action selection (e.g., Coull et al., 2016; Mueller et al.,

2007) and in the FEF that is associated with shifting attention (e.g., Corbetta & Shulman, 2002).

Although this framework is merely provisional requiring further investigation, it is

consistent with perceptual judgment processes. As such we argue that the framework can explain

multiple central properties of perceptual pseudoneglect. For one, stimulus analysis in visual areas

contributing to perceptual pseudoneglect (step 1) is consistent with the recent finding that visual

stimuli impoverished by pixel noise yield stronger biases in a variety of perceptual judgment

tasks (Chen et al., 2019; Chen & Niemeier, 2014), perhaps due to stronger activation of earlier

visual, in particular parvocellular processes (Niemeier et al., 2008) at the expense of higher-order

visual information (e.g., Kayser et al., 2003; Murray et al., 2002). Next, we argue that forming

integrated representations of visual stimuli in right parietal cortex (step 2) is the core reason for

right-biased brain activation in perceptual judgment tasks and that within these areas a more

detailed or salient representation of the contralateral visual field then leads to right-biased

behavioural biases (step 3). The fact that these parietal areas are also involved in attentional

cueing paradigms (Corbetta & Shulman, 2002) further explains why attentional cues modulate

perceptual pseudoneglect (Bultitude et al., 2006; McCourt et al., 2005; Singh et al., 2011; Toba

et al., 2011). Finally, feedback as proposed to occur as step 2a might also occur in judgment

tasks that involve imagery such as mental number line tasks.

In conclusion, in the present study we used a meta-analysis to identify the functional

network involved in perceptual pseudoneglect to reduce the risk of false positive and false

negative errors of previous imaging studies. Given the resulting set of brain areas matched

against EEG data for their temporal activation pattern we propose a new framework of neuro-

computational steps that explain pseudoneglect as the result of interhemispheric integration

afforded by perceptual judgments that, as a consequence, results in asymmetric brain activation

and biased attention rather than attentional biases causing asymmetric judgments. This

framework and intact functional anatomy of perceptual pseudoneglect sheds new light on

perceptual forms of spatial neglect after brain damage.

References

Anderson B (1996). A mathematical model of line bisection behaviour in neglect. Brain 119(Pt
3):841-850. https://doi:10.1093/brain/119.3.841

Badzakova-Trajkov G, Häberling IS, Corballis MC (2011) Magical ideation, creativity,
handedness, and cerebral asymmetries: a combined behavioural and fMRI study.
Neuropsychologia 49(10):2896-2903.
https://doi.org/10.1016/j.neuropsychologia.2011.06.016

Badzakova-Trajkov G, Häberling IS, Roberts RP, Corballis MC (2010) Cerebral asymmetries:
complementary and independent processes. PLoS One 5(3):e9682.
https://doi.org/10.1371/journal.pone.0009682

Bates E, Wilson SM, Saygin AP, Dick F, Sereno MI, Knight RT, Dronkers NF (2003) Voxel-
based lesion-symptom mapping. Nat Neurosci 6(5):448-450.
https://doi.org/10.1038/nn1050

Benwell CS, Harvey M, Thut G (2014) On the neural origin of pseudoneglect: EEG-correlates of
shifts in line bisection performance with manipulation of line length. Neuroimage
86(100):370-380. https://doi.org/10.1016/j.neuroimage.2013.10.014

Bowers D, Heilman KM (1980) Pseudoneglect: effects of hemispace on a tactile line bisection
task. Neuropsychologia 18(4-5):491-498. https://doi.org/10.1016/0028-3932(80)90151-7

Bultitude JH, Aimola Davies AM (2006) Putting attention on the line: investigating the
activation-orientation hypothesis of pseudoneglect. Neuropsychologia 44(10):1849-1858.
https://doi.org/10.1016/j.neuropsychologia.2006.03.001

Cai Q, Van der Haegen L, Brysbaert M (2013) Complementary hemispheric specialization for

language production and visuospatial attention. Proc Natl Acad Sci U S A 110(4):E322-

E330. https://doi.org/10.1073/pnas.1212956110

Caramazza A (1984) The logic of neuropsychological research and the problem of patient

classification in aphasia. Brain Lang 21(1):9-20. https://doi.org/10.1016/0093-

934x(84)90032-4

Cavézian C, Valadao D, Hurwitz M, Saoud M, Danckert J (2012) Finding centre: ocular and

fMRI investigations of bisection and landmark task performance. Brain Res 1437:89-103.

https://doi.org/10.1016/j.brainres.2011.12.002

Chambers CD, Allen CP, Maizey L, Williams M A (2013) Is delayed foveal feedback critical for

extra-foveal perception? Cortex 49(1):327-335.

https://doi.org/10.1016/j.cortex.2012.03.007

Chen J, Kaur J, Abbas H, Wu M, Luo W, Osman S, Niemeier M (2019) Evidence for a common

mechanism of spatial attention and visual awareness: Towards construct validity of

pseudoneglect. PLoS One 14(3):e0212998. https://doi.org/10.1371/journal.pone.0212998

Chen J, Lee ACH, O'Neil EB, Abdul-Nabi M, Niemeier M (2020) Mapping the anatomy of

perceptual pseudoneglect: A multivariate approach. Neuroimage 207:116402.

https://doi.org/10.1016/j.neuroimage.2019.116402

Chen J, Niemeier M (2014) Distractor removal amplifies spatial frequency-specific crossover of

the attentional bias: A psychophysical and Monte Carlo simulation study. Exp Brain Res

232(12):4001-4019. https://doi.org/10.1007/s00221-014-4082-y

Ciçek M, Deouell LY, Knight RT (2009) Brain activity during landmark and line bisection tasks.

Front Hum Neurosci 3:7. https://doi.org/10.3389/neuro.09.007.2009

Corbetta M, Shulman GL (2011) Spatial neglect and attention networks. Annu Rev Neurosci

34:569-599. https://doi.org/10.1146/annurev-neuro-061010-113731

Corbetta M, Shulman GL (2002) Control of goal-directed and stimulus-driven attention in the

brain. Nat Rev Neurosci 3(3):201-215. https://doi.org/10.1038/nrn755

Coull JT, Vidal F, Burle B (2016). When to act, or not to act: that's the SMA's question. Curr

Opin Behav Scie 8:14-21. http://dx.doi.org/10.1016/j.cobeha.2016.01.003

de Haan B, Karnath HO (2018) A hitchhiker's guide to lesion-behaviour mapping.

Neuropsychologia 115:5-16. https://doi.org/10.1016/j.neuropsychologia.2017.10.021

Demanet J, Liefooghe B, Hartstra E, Wenke D, De Houwer J, Brass M (2016) There is more into

'doing' than 'knowing': The function of the right inferior frontal sulcus is specific for

implementing versus memorising verbal instructions. Neuroimage 141:350-356.

https://doi.org/10.1016/j.neuroimage.2016.07.059

Eickhoff SB, Bzdok D, Laird AR, Kurth F, Fox PT (2012). Activation likelihood estimation

meta-analysis revisited. Neuroimage 59(3):2349-2361.

https://doi:10.1016/j.neuroimage.2011.09.017

Filimon F, Nelson JD, Huang RS, Sereno MI (2009) Multiple parietal reach regions in humans:

Cortical representations for visual and proprioceptive feedback during on-line reaching. J

Neurosci 29(9):2961-2971. https://doi.org/10.1523/JNEUROSCI.3211-08.2009

Fink GR, Marshall JC, Shah NJ, Weiss PH, Halligan PW, Grosse-Ruyken M, ... Freund HJ

(2000) Line bisection judgments implicate right parietal cortex and cerebellum as

assessed by fMRI. Neurology 54(6):1324-1331. https://doi.org/10.1212/wnl.54.6.1324

Fink GR, Marshall JC, Weiss PH, Shah NJ, Toni I, Halligan PW, Zilles K (2000) 'Where'

depends on 'what': A differential functional anatomy for position discrimination in one-

versus two-dimensions. Neuropsychologia 38(13):1741-1748.

https://doi.org/10.1016/s0028-3932(00)00078-6

Fink GR, Marshall JC, Weiss PH, Stephan T, Grefkes C, Shah NJ, ... Dieterich M (2003)

Performing allocentric visuospatial judgments with induced distortion of the egocentric

reference frame: An fMRI study with clinical implications. Neuroimage 20(3):1505-

1517. https://doi.org/10.1016/j.neuroimage.2003.07.006

Fink GR, Marshall JC, Weiss PH, Toni I, Zilles K (2002) Task instructions influence the

cognitive strategies involved in line bisection judgments: Evidence from modulated

neural mechanisms revealed by fMRI. Neuropsychologia 40(2):119-130.

https://doi.org/10.1016/s0028-3932(01)00087-2

Fink GR, Marshall JC, Weiss PH, Zilles K (2001) The neural basis of vertical and horizontal line

bisection judgments: An fMRI study of normal volunteers. Neuroimage 14(1 Pt 2):S59-

S67. https://doi.org/10.1006/nimg.2001.0819

Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME (2005). The human

brain is intrinsically organized into dynamic, anticorrelated functional networks. Proc

Natl Acad Sci U S A 102(27):9673-9678. https://doi.org10.1073/pnas.0504136102

Flöel A, Buyx A, Breitenstein C, Lohmann H, Knecht S (2005) Hemispheric lateralization of

spatial attention in right- and left-hemispheric language dominance. Behavioural Brain

Research 158(2): 269-275. https://doi.org/10.1016/j.bbr.2004.09.016

Harvey M, Milner AD, Roberts RC (1995) An investigation of hemispatial neglect using the

Landmark Task. Brain Cogn 27(1):59-78. https://doi.org/10.1006/brcg.1995.1004

Heilman, K., Watson, R., Valenstein, E. (1993) Neglect and related disorders. In Heilman K

Valenstein E (eds) Clinical Neuropsychology, Oxford University Press, New York, pp

279–336.

Hurwitz M, Valadao D, Danckert J (2011) Functional MRI of dynamic judgments of spatial

extent. Exp Brain Res 214(1):61-72. https://doi.org/10.1007/s00221-011-2806-9

Jewell G, McCourt ME (2000) Pseudoneglect: A review and meta-analysis of performance

factors in line bisection tasks. Neuropsychologia 38(1):93-110.

https://doi.org/10.1016/s0028-3932(99)00045-7

Karnath HO, Niemeier M, Dichgans J (1998) Space exploration in neglect. Brain 121(Pt

12):2357-2367. https://doi.org/10.1093/brain/121.12.2357

Kayser C, Salazar RF, Konig P (2003) Responses to natural scenes in cat V1. J Neurophysiol

90(3):1910-1920. https://doi.org/10.1152/jn.00195.2003

Koch G, Oliveri M, Cheeran B, Ruge D, Lo Gerfo E, Salerno S, ... Caltagirone C (2008)

Hyperexcitability of parietal-motor functional connections in the intact left-hemisphere of

patients with neglect. Brain 131(Pt 12):3147-3155. https://doi.org/10.1093/brain/awn273

Le A, Stojanoski BB, Khan S, Keough M, Niemeier M (2015) A toggle switch of visual

awareness? Cortex 64:169-178. https://doi.org/10.1016/j.cortex.2014.09.015

Liu D, Zhou D, Li M, Li M, Dong W, Verguts T, Chen Q (2019) The Neural Mechanism of

Number Line Bisection: A fMRI study. Neuropsychologia 129:37-46.

https://doi.org/10.1016/j.neuropsychologia.2019.03.007

Marshall RS, Lazar RM, Van Heertum, RL, Esser PD, Perera GM, Mohr JP (1997) Changes in

regional cerebral blood flow related to line bisection discrimination and visual attention

using HMPAO-SPECT. Neuroimage 6(2):139-144.

https://doi.org/10.1006/nimg.1997.0283

Mattingley JB, Berberovic N, Corben L, Slavin MJ, Nicholls ME, Bradshaw JL (2004) The

greyscales task: a perceptual measure of attentional bias following unilateral hemispheric

damage. Neuropsychologia 42(3):387-394.

https://doi.org/10.1016/j.neuropsychologia.2003.07.007

McCourt ME, Garlinghouse M, Reuter-Lorenz PA (2005) Unilateral visual cueing and

asymmetric line geometry share a common attentional origin in the modulation of

pseudoneglect. Cortex 41(4):499-511. https://doi.org/10.1016/s0010-9452(08)70190-4

McCourt ME, Jewell G (1999) Visuospatial attention in line bisection: stimulus modulation of

pseudoneglect. Neuropsychologia 37(7):843-855. https://doi.org/10.1016/s0028-

3932(98)00140-7

Molenberghs P, Sale MV, Mattingley JB (2012). Is there a critical lesion site for unilateral

spatial neglect? A meta-analysis using activation likelihood estimation. Front Hum

Neurosci 6:78. https://doi.org/10.3389/fnhum.2012.00078

Monaco S, Gallivan JP, Figley TD, Singhal A, Culham JC (2017) Recruitment of Foveal

Retinotopic Cortex During Haptic Exploration of Shapes and Actions in the Dark. J

Neurosci, 37(48):11572-11591. https://doi.org/10.1523/JNEUROSCI.2428-16.2017

Moore MJ, Milosevich E, Mattingley JB, Demeyere N (2023) The neuroanatomy of visuospatial

neglect: A systematic review and analysis of lesion-mapping methodology.

Neuropsychologia 180:108470. https://doi.org/10.1016/j.neuropsychologia.2023.108470

Mueller VA, Brass M, Waszak F, Prinz W (2007). The role of the preSMA and the rostral

513   cingulate zone in internally selected actions. Neuroimage 37(4):1354-1361.

514   https://doi.org10.1016/j.neuroimage.2007.06.018

515   Müller VI, Cieslik EC, Laird AR, Fox PT, Radua J, Mataix-Cols D, ... Eickhoff SB (2018) Ten

516   simple rules for neuroimaging meta-analysis. Neurosci Biobehav Rev 84:151-161.

517   https://doi.org/10.1016/j.neubiorev.2017.11.012

518   Murray SO, Kersten D, Olshausen BA, Schrater P, Woods DL (2002) Shape perception reduces

519   activity in human primary visual cortex. Proc Natl Acad Sci U S A99(23):15164-15169.

520   https://doi.org/10.1073/pnas.192579399

521   Myers NE, Stokes MG, Nobre AC (2017) Prioritizing Information during Working Memory:

522   Beyond Sustained Internal Attention. Trends Cogn Sci 21(6):449-461.

523   https://doi.org/10.1016/j.tics.2017.03.010

524   Niemeier M, Singh VV, Keough M, Akbar N (2008) The perceptual consequences of the

525   attentional bias: evidence for distractor removal. Exp Brain Res 189(4):411-420.

526   https://doi.org/10.1007/s00221-008-1438-1

527   Niemeier M, Stojanoski B, Greco AL (2007) Influences of time and spatial frequency on the

528   perceptual bias: evidence for competition between hemispheres. Neuropsychologia

529   45(5):1029-1040. https://doi.org/10.1016/j.neuropsychologia.2006.09.006

530   Pascual-Leone A, Amedi A, Fregni F, Merabet LB (2005) The plastic human brain cortex. Annu

531   Rev Neurosci 28:377-401. https://doi.org/10.1146/annurev.neuro.27.070203.144216

532   Plewan T, Weidner R, Eickhoff SB, Fink GR (2012) Ventral and dorsal stream interactions

533   during the perception of the Müller-Lyer illusion: evidence derived from fMRI and

dynamic causal modeling. J Cogn Neurosci 24(10):2015-2029.

https://doi.org/10.1162/jocn_a_00258

Price CJ, Friston KJ (1997) Cognitive conjunction: a new approach to brain activation

experiments. Neuroimage 5(4 Pt 1):261-270. https://doi.org/10.1006/nimg.1997.0269

Revill KP, Karnath HO, Rorden C (2011) Distinct anatomy for visual search and bisection: a

neuroimaging study. Neuroimage 57(2):476-481.

https://doi.org/10.1016/j.neuroimage.2011.04.066

Sahan MI, Majerus S, Andres M, Fias W (2019) Functionally distinct contributions of parietal

cortex to a numerical landmark task: An fMRI study. Cortex 114:28-40.

https://doi.org/10.1016/j.cortex.2018.11.005

Schuster V, Herholz P, Zimmermann KM, Westermann S, Frässle S, Jansen A (2017)

Comparison of fMRI paradigms assessing visuospatial processing: Robustness and

reproducibility. PLoS One 12(10):e0186344.

https://doi.org/10.1371/journal.pone.0186344

Sheremata SL, Silver MA (2015) Hemisphere-dependent attentional modulation of human

parietal visual field representations. J Neurosci 35(2):508-517.

https://doi.org/10.1523/JNEUROSCI.2378-14.2015

Singh VVW, Stojanoski B, Le A, Niemeier M (2011) Spatial frequency-specific effects on the

attentional bias: Evidence for two attentional systems. Cortex 47(5):547–556.

https://doi.org/10.1016/j.cortex.2010.03.006

Sperber C, Karnath HO (2017) Impact of correction factors in human brain lesion-behavior

inference. Hum Brain Mapp 38(3):1692-1701. https://doi.org/10.1002/hbm.23490

Szczepanski SM, Kastner S (2013) Shifting attentional priorities: Control of spatial attention

through hemispheric competition. J Neurosci 33(12):5411-5421.

https://doi.org/10.1523/JNEUROSCI.4089-12.2013

Thiebaut de Schotten M, Urbanski M, Duffau H, Volle E, Lévy R, Dubois B, Bartolomeo P

(2005). Direct evidence for a parietal-frontal pathway subserving spatial awareness in

humans. Science 309(5744):2226-2228. https://doi:10.1126/science.1116251

Thiebaut de Schotten M, Tomaiuolo F, Aiello M, Merola S, Silvetti M, Lecce F, ... Doricchi F

(2014) Damage to white matter pathways in subacute and chronic spatial neglect: a group

study and 2 single-case studies with complete virtual "in vivo" tractography dissection.

Cereb Cortex 24(3):691-706. https://doi.org/10.1093/cercor/bhs351

Toba MN, Cavanagh P, Bartolomeo P (2011). Attention biases the perceived midpoint of

horizontal lines. Neuropsychologia 49(2):238-246.

https://doi:10.1016/j.neuropsychologia.2010.11.022

Vallar G, Bello L, Bricolo E, Castellano A, Casarotti A, Falini A, Riva M, Fava E, Papagno C

(2014). Cerebral correlates of visuospatial neglect: a direct cerebral stimulation study.

Hum Brain Mapp 35(4):1334-1350. https://doi:10.1002/hbm.22257

Vesia M, Niemeier M, Black SE, Staines WR (2015) The time course for visual extinction after a

'virtual' lesion of right posterior parietal cortex. Brain Cogn 98:27-34.

https://doi.org/10.1016/j.bandc.2015.05.003

Vossel S, Geng JJ, Fink GR (2014) Dorsal and ventral attention systems: distinct neural circuits

but collaborative roles. Neuroscientist 20(2):150-159.

https://doi.org/10.1177/1073858413494269

Walsh V (2003) A theory of magnitude: Common cortical metrics of time, space and quantity.

Trends Cogn Sci 7(11):483-488. https://doi.org/10.1016/j.tics.2003.09.002

Weidner R, Fink GR (2007) The neural mechanisms underlying the Müller-Lyer illusion and its

interaction with visuospatial judgments. Cereb Cortex 17(4):878-884.

https://doi.org/10.1093/cercor/bhk042

Weiss PH, Marshall JC, Wunderlich G, Tellmann L, Halligan PW, Freund HJ, ... Fink GR (2000)

Neural consequences of acting in near versus far space: a physiological basis for clinical

dissociations. Brain 123(Pt 12):2531-2541. https://doi.org/10.1093/brain/123.12.2531

Weiss PH, Marshall JC, Zilles K, Fink, GR (2003) Are action and perception in near and far

space additive or interactive factors? Neuroimage 18(4):837-846.

https://doi.org/10.1016/s1053-8119(03)00018-1

Williams MA, Baker CI, Op de Beeck, HP, Shim WM, Dang S, Triantafyllou C, Kanwisher N

(2008) Feedback of visual object information to foveal retinotopic cortex. Nat Neurosci

11(12):1439-1445. https://doi.org/10.1038/nn.2218

Zago L, Petit L, Jobard G, Hay J, Mazoyer B, Tzourio-Mazoyer N, ... Mellet E (2017)

Pseudoneglect in line bisection judgement is associated with a modulation of right

hemispheric spatial attention dominance in right-handers. Neuropsychologia 94:75-83.

https://doi.org/10.1016/j.neuropsychologia.2016.11.024

Zago L, Petit L, Mellet E, Jobard G, Crivello F, Joliot M, ... Tzourio-Mazoyer N (2016) The

association between hemispheric specialization for language production and for spatial

attention depends on left-hand preference strength. Neuropsychologia 93(Pt B):394-406.

https://doi.org/10.1016/j.neuropsychologia.2015.11.018

601