1   Diploidy confers genomic instability in

2   *Schizosaccharomyces pombe*

4   Joshua M. Park, Daniel F. Pinski & Susan L. Forsburg[*]

5   Section of Molecular & Computational Biology

6   University of Southern California

7   1050 Childs Way, RRI 108

8   Los Angeles CA 90089

[*]corresponding author, forsburg@usc.edu

Abstract

Whole genome duplication, or polyploidy, has been implicated in driving genome instability and
tumorigenesis. Recent studies suggest that polyploidy in tumors promotes cancer genome
evolution, progression, and chemoresistance resulting in worse prognosis of survival. The
mechanisms by which whole genome duplications confer genome instability are not yet fully
understood. In this study, we use *Schizosaccharomyces pombe* (fission yeast) diploids to
investigate how whole genome duplication affects genome maintenance and response to stress.
We find that *S. pombe* diploids are sensitive to replication stress and DNA damage, exhibit high
levels of loss of heterozygosity, and become dependent on a group of ploidy-specific lethal genes
for viability. These findings are observed in other eukaryotic models suggesting conserved
consequences of polyploidy. We further investigate ploidy-specific lethal genes by depleting
them using an auxin-inducible degron system to elucidate the mechanisms of genome
maintenance in diploids. Overall, this work provides new insights on how whole genome
duplications lead to genome instability.

Introduction

Recent studies have estimated that ~36% of tumors undergo a whole genome duplication event
(Zack *et al.* 2013; Quinton *et al.* 2021). Whole genome duplications (polyploidy) have been
increasingly associated with genome instability and tumorigenesis in various eukaryotic models
(Fujiwara, *et al.* 2005; Storchová *et al.* 2006; Gemble *et al.* 2022). Human tetraploid cells show
signs of increased DNA damage and chromosomal instability (Kuznetsova *et al.* 2015; Nano *et*
al. 2019) that may promote cancer progression (Dewhurst *et al.* 2014). Polyploidy has also been
linked to increased resistance of cancer cells to chemotherapeutic drugs (Mirzayans *et al.* 2018;
Bharadwaj and Mandal 2020) and worse prognosis of survival in cancer patients (Dewhurst *et al.*
2014; Bielski *et al.* 2018).

Another hallmark of cancer is loss of heterozygosity (LOH; Steele *et al.* 2022). LOH is a
phenomenon where a heterozygous region of the genome becomes homozygous, usually as a
result of mitotic recombination (reviewed in Heil 2023). Crossover events or break-induced
replication can result in long tracts of the genome losing heterozygosity, driving tumorigenesis
through loss of tumor suppressor function (Had ija *et al.* 2001; Boulay *et al.* 2009; Lourenço *et*
al. 2014; Ciani *et al.* 2022). Interestingly, cancer vulnerabilities can arise when non-driver genes
undergo LOH, providing potential therapeutic targets (Nichols *et al.* 2020; Zhang and Sjöblom
2021). Recent research has suggested that cancer cells will undergo WGD as a means to combat
the negative effects of LOH while still driving tumorigenesis (López *et al.* 2020; Archetti 2022).
Although WGDs have been shown to play a role in genome instability and cancer progression,
the underlying mechanisms require further investigation.

Yeasts provide a single celled model to study genome duplication. The budding yeast *S.*
cerevisiae normally alternates between haploid and diploid states (Knop 2006), although higher
degrees of polyploidy have been isolated and are associated with genome instability and LOH
(Mayer and Aguilera 1990; Storchová *et al.* 2006; Storchová 2014; Sui *et al.* 2020; Dutta *et al.*
2022). In contrast, fission yeast is typically haploid, with a transient diploid zygote phase

(Forsburg and Nurse 1991). In this organism, both diploids and higher level polyploids have
been isolated but are unstable (Molnar and Sipiczki 1993).

In this study, we use the fission yeast to investigate the consequences of WGD, and the response
to genomic stress. We compare *S. pombe* haploids to diploids and find that diploids exhibit
increased sensitivity to genotoxins and experience elevated levels of LOH in the absence of
exogenous stress. We show that this effect is independent of mating type heterozygosity. We also
identify ploidy-specific lethal (PSL) genes that are conserved in budding yeast and humans
(Storchová *et al.* 2006; Quinton *et al.* 2021). Using an auxin-inducible degron system, we
investigate the mechanisms of cell death in PSL depleted diploids. We find that *S. pombe*
diploids exhibit similar genomic instability phenotypes as human cells and tumors that have
undergone WGD.

Materials and Methods

Yeast strains and growth

S. pombe strains used are listed in Table S1. *S. pombe* cells were cultured following standard
protocols and methods (Forsburg and Rhind 2006).

AID tagged strain construction

Auxin-inducible degron (AID) strains were constructed as described in Watson *et al.* 2021.
pAW8-aid-5myc-Turg1-kanMX6 plasmid (Addgene 169356) was used as a template for PCR.
Two rounds of PCR were performed to add sequence to both ends of the AID cassette that were
homologous to the genomic target locus for C-terminus tagging. This PCR product was
transformed using electroporation (Sabatinos and Forsburg 2010) into a wild type strain, then
crossed with a strain with *OsTIR1F74A* integrated into the genome. Integration was validated by
genomic PCR and Sanger sequencing. Rad52-AID-V5 strains were constructed by crosses using
a strain provided by Dr. Antony Carr (Watson *et al.* 2021).

Diploid strain generation

Mating type heterozygous diploid strains (sporulating and non-sporulating) were constructed
using *ade6-M210/ade6-M216* intragenic complementation (Ekwall and Thon 2017). Mating type
homozygous diploid strains were constructed by protoplast fusion (Park and Forsburg 2024).
Diploid strains were validated by flow cytometery on the BD Accuri C6 Plus platform.

Serial dilution growth assay

Liquid cultures were grown overnight in yeast extract +supplements (YES) media. Cultures were
diluted to equal concentrations, then diluted five-fold serially. Dilutions were spotted onto YES
or minimal glutamate media (PMG) plates with indicated supplements and drug concentrations.
Plates were incubated at 32ºC for the time indicated in figure legends.

Acute genotoxic stress treatment survival

Liquid cultures were grown to mid log phase. Each culture was split in half for no treatment and
treatment group. For irradiation (IR), cells were irradiated at the indicated doses using a 160 kV

X-ray source (X-RAD iR-160, Precision X-Ray). For methyl methanesulfonate (MMS)

treatment, MMS was added to the indicated doses and liquid cultures were incubated at 32⁰C

with shaking for 4 hours. After treatment, cells were washed twice and resuspended in water.

Cells were then counted on a Bright-Line Hemacytometer (Hausser Scientific) and 500 cells

were plated onto YES + phloxine B plates. Plates were incubated at 32⁰C for 3 days, then

colonies were counted and ratioed to untreated samples. Statistical significance was calculated

using Mann-Whitney U-test (p-value > 0.05 ns, p-value < 0.05 *, p-value < 0.01 **, p-value <

0.001 ***, p-value < 0.0001 ****).

<u>Live cell fluorescent imaging</u>

Fluorescent imaging was performed as described in Jones and Forsburg 2021. Briefly, liquid

cultures were grown to mid-log phase in PMG supplemented with histidine, uracil, leucine,

adenine, lysine, and arginine at 225 mg/L each (PMG-HULALA). MMS (0.0075%) or 5-

Adamantyl-IAA (5'a-IAA, 100 nM) were added to the culture and incubated at 32⁰C with

shaking. Samples were taken at the indicated timepoints for fluorescent imaging. Cells

concentrated by a brief microfuge spin were applied to 2% agarose pads made from PMG +

HULA and prepared on glass slides (Green *et al.* 2009). Images were acquired on a DeltaVision

microscope with softWoRx v4.1 (GE, Issaquah, WA) using a 60x (NA1.4 PlanApo) lens, solid-

state illuminator and 12-bit CCD camera. Images were acquired in twenty 0.2µm z-sections, then

deconvolved and Maximum Intensity Projected (softWoRx, default settings). Images for

publication were contrast adjusted using an equivalent histogram stretch on all samples. Color

balance was adjusted and scale bars were added in Fiji (ImageJ 2.0). Statistical significance was

calculated using Mann-Whitney U-test.

<u>Fixed cell imaging</u>

Fixed cell imaging by 4′,6-diamidino-2-phenylindole (DAPI) staining was performed as

described in Luche and Forsburg 2009. Cells were cultured in PMG-HULALA and samples were

collected at the indicated time points. Samples were ethanol fixed in ice-cold 70% ethanol and

rehydrated in water. 5 µL of rehydrated sample was placed on a microscope slide and dried on a

hot plate (70⁰C). 5 µL of mounting solution (50% glycerol, 1 mg/mL p-Phenylenediamine, 1

µg/mL DAPI) was added on top of the dried sample and a cover slip placed on top. Images were

acquired as described above.

<u>Induced mutagenesis and LOH assay</u>

Liquid cultures were grown to log phase in YES media. Mutagenesis was induced by 0.0025%

MMS treatment for one hour at 32⁰C with shaking. Cultures were unperturbed for the LOH

assay. Cells were washed twice in water then counted on a hemacytometer. 500 cells were plated

onto YES + phloxine B plates and 1×10^5 cells were plated onto PMG supplemented with

histidine and leucine (225 mg/L each), uracil (50 mg/L), and 5-Fluoroorotic acid (5-FOA, 1

mg/mL) plates (150 mm x 15 mm). YES + phloxine B plates were incubated for 3 days and 5-

FOA plates for 4 days at 32⁰C before counting colonies. Haploidization was measured by

counting lighter pink colonies on the phloxine B plates. 5-FOA resistant colonies were ratioed to

YES colonies minus those that had haploidized. 5-FOA plates were replica plated onto YES

plates containing nourseothricin or zeocin and incubated at 32ºC overnight. Resistant colonies
were ratioed to total 5-FOA resistant colonies. Statistical significance was calculated using
Mann-Whitney U-test.

<u>LOH sequencing analysis</u>

Hybrid diploid strains were constructed by mating a *mat2-102 S. pombe* strain containing a
ura4+ cassette with a *h- S. pombe var. kambucha* strain. Colonies that had undergone LOH were
selected for using 5-FOA as detailed above. A control colony and 10 5-FOA resistant colonies
were cultured in PMG supplemented with histidine, leucine, adenine, and uracil (225 mg/L
each). 20 mL of mid log phase culture was processed for genomic DNA extraction using a
Monarch HMW DNA Extraction Kit for Tissue (New England Biolabs). Library preparation was
performed using a Native Barcoding Kit 24 V14 and samples were sequenced on a MinION
Mk1C platform (Oxford Nanopore Technologies). Raw pod5 files were basecalled using Dorado
v7.1.4 dna_r10.4.1_e8.2_400bps_5khz_sup model. Reads were aligned to the *S. pombe var.*
kambucha reference genome and single nucleotide polymorphism (SNP) analysis was performed
using the Geneious Prime 2024.0.7 software. SNP density plots were generated on R 4.3.1 with
the package CMPlot (Yin *et al.* 2021).

<u>Protein extraction and immunoblot</u>

Liquid cultures were grown to log phase, then treated with either 0.01% MMS (Sigma-Aldrich)
or 10 nM 5'a-IAA (TCI America). Samples were collected at time points indicated and were
washed twice in PBS. Sample pellets were then frozen in liquid nitrogen and all time point
samples were processed together. Protein was extracted by trichloroacetic acid (TCA)
precipitation (Grallert and Hagan 2017). Protein samples were quantified using Pierce BCA
Protein Assay Kits (Thermo Scientific) and 25-50 µg protein was loaded. Samples were run on a
7.5% SDS-PAGE gel and transferred onto a nitrocellulose membrane. The following antibodies
and dilutions were used for immunoblotting: α-*cdc2* (Novus Biologicals NB100-2716; 1:2000),
α-V5 (Abcam ab27671; 1:1000), α-myc (Abcam ab9106 and Novus Biologicals NBP2-52636;
1:1000), α -Alpha Tubulin (Millipore Sigma T5168; 1:1000), α-Rabbit Alexa Fluor 488 (Thermo
Scientific A32790; 1:1000), and α-Mouse Alexa Fluor 488 (Thermo Scientific A28175; 1:1000).
Blots were visualized on the Amersham Typhoon platform and quantified using ImageQuant TL
software (GE healthcare).

<u>Results</u>

To study how diploidy affects genome stability in *S. pombe*, and to eliminate any contribution of
mating type heterozygosity, we generated three kinds of diploids. First, we constructed a mating-
type heterozygous h^+/h^- diploid that is sporulation competent. The second diploid h$^-$ /*mat2-102*
lacks *matP*-information required for sporulation (Willer *et al.* 1995) but otherwise maintains
mating type heterozygosity. The third strain is mating type homozygous, constructed by
protoplast fusion. Initially, we examined h^+/ h^+ homozygous strains, but to eliminate any
potential reversion to h^{90}, we also examined *h$^-$ smt-0* / *h$^-$ smt-0* strains that lack the site for the

mating type switching imprint (Styrkársdóttir *et al.* 1993). These configurations control for any
possible effects from mating type heterozygosity and isolate ploidy specific effects.

<u>Diploids are more sensitive to genotoxins</u>

We observed that all three diploid strains exhibited increased sensitivity to a variety of genotoxic
agents compared to the haploids (Figure 1A). All four diploid strains had similar levels of
sensitivity, suggesting that mating heterozygosity did not have any affect, and h^+/h^+ and h^- *smt-*
0/h^- smt-0 diploids behaved similarly (Figure 1A). We also observed increased sensitivity to
irradiation in diploids compared to haploids across a range of exposure (Figure 1B). Because the
h^+/h^- strain is prone to enter meiosis, we continued with the two non-sporulating strains for
further characterization.

We tested diploid response to acute treatment of the alkylating agent methyl methanesulfonate
(MMS) and observed that diploids exhibited increased sensitivity across a range of
concentrations (Figure 1C). We visualized DNA damage/repair foci using fluorescently tagged
Ssb1 (RPA), Rad52, and Rad54 (Lisby *et al.* 2004; Sabatinos *et al.* 2012). We found that the
percentage of cells with RPA and Rad54 foci did not significantly differ in haploid and diploid
strains in unperturbed conditions (Figure 1D and 1F). After treatment with MMS, diploid cells
showed an increased number of cells with RPA and Rad54 foci (Figure 1D and 1F). Interestingly,
Rad52 foci were slightly elevated in diploids compared to haploids in unperturbed conditions but
were at similar levels after MMS treatment (Figure 1E).

<u>Diploids have loss of heterozygosity</u>

To further characterize the response of diploids to MMS, we compared the induced mutagenesis
rate between haploids and diploids. We performed a forward mutation assay in *ura4+* strains by
determining the frequency of 5-Fluoroorotic acid (5-FOA) resistance with and without MMS
treatment (Liu *et al.* 1999; Dolan *et al.* 2010). We constructed diploid strains that were
homozygous or heterozygous for the *ura4+* gene at its endogenous locus and treated haploid and
diploid strains with 0.0025% MMS for one hour. To our surprise, the *ura4+* heterozygous
strains had significantly elevated levels of 5-FOA resistance in both the untreated and treated
conditions (Fig. 2A). We confirmed by PCR that nearly all the diploid samples tested had lost the
entire *ura4+* gene (Table S2), resulting in a loss of heterozygosity (LOH) rather than point
mutation(s).

We were concerned that the endogenous *ura4+* locus's proximity to the telomeres and rDNA
array on chromosome III may have influenced the results, so we inserted the *ura4+* cassette into
chromosome I and II. Again, we observed elevated levels of LOH in these diploids even in the
absence of MMS (Fig. 2B). To assess whether this was a limited gene conversion, or loss of a
chromosome, we constructed strains with the *ura4+* cassette and two drug markers *natMX* and
kanMX (conferring resistance to nourseothricin and G418, respectively) equally spaced along the
left arm of chromosome I and another marker *bleMX* (conferring resistance to bleomycin and
related drugs) on the right arm. We observed that the rate of *ura4+* loss was similar in strains
with the *ura4+* cassette inserted in the left and middle arm of the chromosome (Fig. 2C).

However, the *ura4+* loss was significantly reduced in the strain that had the *ura4+* cassette
inserted near the centromere (Fig. 2C).

Next, we measured the loss of the other markers in cells that had lost the *ura4+* cassette. In the
strain with the *ura4+* telomere proximal, the middle marker was also lost in ~70% of the
population while the centromere proximal and right arm marker were lost in ~10% (Fig. 2D). In
the strain with the *ura4+* at the middle position, the telomere proximal maker was lost in ~100%
(Fig. 2E). However, in the limited population of colonies that had lost the centromere proximal
ura4+ cassette, ~90% of the telomere proximal and middle markers were lost as well as ~80% of
the right arm marker (Fig. 2F). We suggest the high rates of LOH of the telomere-proximal and
middle marker reflect an extended recombination or break induced replication event, while the
loss of the centromere proximal *ura4+* reflects a chromosome mis-segregation event that results
in homozygosis. Remarkably, this happens in strains that are otherwise unperturbed by
genotoxic stress.

<u>Mapping LOH breakpoints in hybrid *S. pombe* diploids</u>

The high rate of LOH associated with replication stress led us to wonder if there were
recombination hotspots linked to fragile sites in the genome. To investigate recombination
breakpoints, we used the *S. pombe variant kambucha* to generate a hybrid diploid with our *S.*
pombe lab strain. The *kambucha* variant has less than 1% nucleotide divergence from *S. pombe*
(Rhind *et al.* 2011) but provides enough single nucleotide polymorphisms (SNPs) to measure
LOH in the hybrid diploid strain. We inserted the *ura4+* cassette in the left arm of Chromosome
I in the *S. pombe* strain (Fig. 3A) and measured loss of the marker using 5-FOA as described
above. We found that the hybrid diploid has slightly lower levels of LOH compared to an
isogenic *S. pombe* diploid (Fig. S1A). We sequenced a control colony and 10 5-FOA resistant
colonies, aligned the reads to the *S. pombe var. kambucha* reference genome, and performed a
SNP analysis. Loss of the *ura4+* cassette on the *S. pombe* strain should be the result of
recombination which leads to homozygosis of a region of the *S. pombe* chromosome and loss of
SNPs in reference to the *S. pombe var. kambucha* genome. Indeed, we find that the 5-FOA
resistant colonies have significantly reduced number of SNPs on the left arm of chromosome I
(Fig. 3B) resulting in loss of heterozygosity. There were two classes of LOH in the 10 samples
we analyzed: 7 of the 10 samples experienced a distinct break at ~1.1 Mb and the remaining 3
samples experienced more extensive LOH (>2 Mb; Fig. 3B). The 1.1 Mb breakpoint coincides
with a long terminal repeat element (LTR; SPLTRA.22) and an early firing origin 22 kb upstream
of the breakpoint (Fig. 3C, S1B).

<u>LOH increases in mutant diploids</u>

To investigate the pathways that may be involved with the LOH phenotype, we took a candidate
approach. We selected a panel of strains with deletion mutations in genes that are implicated in
genome maintenance pathways. We constructed homozygous deletion diploids with a single
telomere proximal *ura4+* marker on the left chromosome arm, a marker in the middle of the left
chromosome arm, and a marker on the right chromosome arm and assessed their effect on LOH
frequency (Fig. 4A, Table 1). We found that deletion of checkpoint kinases *tel1Δ* and *chk1Δ* did

not change the frequency of *ura4+* loss, but *rad3Δ* and *cds1Δ* diploids had a significant increase
(1.4 and 3.2 fold increase, respectively). Deletion of *mrc1*, a replication checkpoint mediator and
interactor of the fork protection complex (FPC; Tanaka and Russell 2001; Tanaka *et al.* 2010),
also increased the loss of *ura4+* at a similar rate as *cds1Δ*. This suggests that an active
replication checkpoint reduces the frequency of mitotic recombination in wild type cells.

Next, we looked at a component of non-homologous end joining (NHEJ) *pku70*. We find that
pku70Δ increases *ura4+* loss rate (3.2 fold increase)*,* similar to *cds1Δ* and *mrc1Δ*. Deletion of
rad8, an ubiquitin ligase (Frampton *et al.* 2006; Ding and Forsburg 2014), increased the rate of
ura4+ loss by 1.6 times compared to wild type (WT). Next, we looked at deletion of *mgs1*, a
Werner helicase-interacting protein 1 (WRNIP1) homolog that is implicated in genome
maintenance and DNA repair (Hishida *et al.* 2001; Saugar *et al.* 2012). Surprisingly, *mgs1Δ*
resulted in a small but significant reduction in *ura4+* loss rate compared to the WT diploid (0.6
times lower).

We assayed HDR associated genes including *exo1Δ, srs2Δ, smc6-74.* Deletion of *exo1*, a long
range resection exonuclease (Mimitou and Symington 2008), and *smc6-74*, a loss of function
mutation in a structural maintenance of chromosome (SMC) gene involved in HDR (Ampatzidou
et al. 2006; Irmisch *et al.* 2009), both increased the rate of loss of *ura4+* by 2.2 and 2.4 fold,
respectively. Deletion of *srs2*, a helicase with anti-recombinogenic properties (Doe and Whitby
2004), induced the highest increase in *ura4+* loss among all the mutants tested at 4.4 fold.
Finally, we looked at deletion of *clr4*, a lysine methyltransferase responsible for depositing
histone H3K9 methylation at the pericentromeric heterochromatin (reviewed in Allshire and
Ekwall 2015). Although we find that *clr4Δ* reduces loss of *ura4+*, this may be skewed by the
high rate of haploidization in this strain probably reflecting chromosome loss due to centromere
disruption (Fig. S2). Overall, we find a trend that genes involved in response to replication stress
and DNA repair tend to increase the loss of the *ura4+* marker.

In the population of diploids that had lost the *ura4+* marker, we also probed for the loss of the
other markers on the left and on the right arm of the chromosome. We observed that *mrc1Δ*,
pku70Δ, and *srs2Δ* have reduced rate of loss of the additional markers compared to WT, while
smc6-74 and *clr4Δ* have increased rates (Fig. 4B). For the marker on the right arm of the
chromosome, we found that *smc6-74* and *clr4Δ* had the largest increase in rate of loss (Fig. 4C).
This is consistent with *clr4* and *smc6-74* impacting loss of the entire chromosome, which may
reflect whole chromosome mis-segragation. Notably, *srs2Δ* had several outliers in both directions
of rate change suggesting it may impact multiple pathways.

Next, we tested MMS sensitivity for haploid and diploids mutants used in the LOH assay. We
found that *exo1Δ* and *smc6-74* diploids were more sensitive to MMS than wild type diploids and
their haploid counterparts (Fig. S3). The remaining diploid mutants were either just as sensitive
to MMS as their haploid counterparts, or comparable to the wild type.

<u>Mutants inviable as diploids</u>

Previous studies suggested that deletions of DNA homologous recombination genes *rad51,*
rad52, and *rad54* could not be constructed as diploids (Catlett and Forsburg 2003; Grishchuk *et*

al. 2004; Octobre et al. 2008). Similar results were observed for *bub1* (Bernard *et al.* 1998). In the process of constructing diploid mutants for the LOH assay, we validated and identified new ploidy specific lethal genes in *S. pombe* (Table 1). We confirmed that *rad51Δ, rad52Δ,* and *rad54Δ* diploids could not be stably isolated by crossing or protoplast fusion. Additionally, we found *mre11Δ, fbh1Δ, mus81Δ,* and *crb2Δ* diploids could not be constructed, adding evidence to the homologous recombination pathway being essential in *S. pombe* diploids. We also found that deletion of topoisomerase I *top1* and FPC component *swi3* were inviable, suggesting that responding to replication stress is essential as ploidy increases. We observed that spindle assembly checkpoint genes *bub1, mad2,* and *mad3* are also essential in diploids. For certain deletions, we were initially able to isolate diploids under selection but were unable to maintain them as they haploidized in culture. High frequency of haploidization is linked to chromosome loss (Bodi *et al.* 1991). These included additional DNA repair and FPC genes (*rqh1Δ, smc6-x, cdc27-D1,* and *swi1Δ*) and cell cycle mutants generally used for synchrony (*cdc10-V50, cdc25-22,* and *cdc2-asM17*) (Table 1).

Seeing as these group of genes were essential for diploid viability, we were interested if overexpression of these genes could rescue diploid sensitivity to genome stress. We tested overexpression of the DNA damage and replication stress genes in response to MMS and the spindle assembly checkpoint genes in response to the microtubule destabilizing agent thiabendazole (TBZ). We found that overexpression of the diploid essential genes did not rescue diploid sensitivity to MMS or TBZ, and in some cases resulted in decreased viability in both haploids and diploids regardless of genotoxic stress (Fig. S4).

Ploidy-specific lethal protein depletion in diploids

To investigate why some genes are necessary for survival in *S. pombe* diploids, we used the auxin-inducible degron (AID) system (Watson *et al.* 2021). Not all genes were amenable to this degron fusion but we successfully constructed diploid strains with both copies of *rad52* and *mad2* tagged with an AID cassette. In the haploid strains, the addition of the auxin analog 5'adamantyl-IAA (5'a-IAA) leads to rapid degradation of the tagged protein within 30 minutes (Watson *et al.* 2021, Fig. 5C). In the diploid strains, protein degradation was ablated for Mad2 but not Rad52 (Fig. 5A and 5C). However, both diploid strains showed significant decrease in viability when plated on media with 5'a-IAA (Fig. 5B and 5E) suggesting that the level of protein depletion was sufficient for loss of viability. The Rad52-AID diploids showed hypersensitivity to low levels of HU compared to their haploid counterpart when Rad52 was depleted (Fig. 5B). The AID cassette did increase sensitivity to MMS in Rad52 tagged haploid and diploid strains in the absence of 5'a-IAA, suggesting impairment of protein function (Fig. S5). Similarly, Mad2-AID diploids exhibited increased sensitivity to low levels of TBZ (Fig. 5E). These results highlight the effectiveness of using the AID system in *S. pombe* diploids and provides a tool for investigating the mechanisms on how depletion of PSL proteins leads to cell death.

Characterizing Rad52 depletion in diploids

We investigated the phenotype of Rad52 depletion, beginning with accumulation of RPA repair
foci. Both haploids and diploids had increased number of cells with RPA foci at 4 hours of
depletion, but the diploids had about 1.6 time more (39% and 65%, respectively; Fig. 6A). The
number of cells with RPA foci steadily increased in both haploids and diploids with time but
maintained a similar ratio. Diploids accumulated increasing levels of multi-foci RPA compared
to haploids (Fig. S6A). While the haploids continued to divide and increase in cell number,
diploids mainly grew in size (coinciding with increase in OD) and divided very slowly (Fig. 6B
and 6C). We see that Chk1 is persistently activated when Rad52 is depleted in both haploid and
diploid strains (Fig 6D and 6E). The Chk1 activation is strongest at the 4 hour mark, but is not as
strong compared to an MMS treated sample.

We used a fluorescently labeled histone to measure any abnormal nuclear morphology and
mitotic events (e.g. lagging chromosomes, uneven segregation, micronuclei; Fig. S6B and S6C).
Most of the diploids showed single nuclei consistent with cell cycle arrest. Diploids had a small
increase in nuclear abnormalities before treatment with 5a'-IAA, and the difference grew
throughout the time course (Fig. 6F). We observed that at the 12 and 24 hour marks the diploids
exhibited aberrant septation attempts such as multiple septa (Fig. S6D). These phenotypes are
consistent with prolonged cell cycle arrest. Next, we deleted Chk1 in Rad52-AID strains and
visualized DNA by DAPI staining in fixed cells as we were unable to isolate *chk1Δ* Rad52-AID
histone tagged strains. We find a similar pattern of increasing levels of abnormal mitotic events
as time progresses, but to a higher degree compared to the *Chk1$^+$* Rad52-AID strains (Fig. 6G).
Interestingly, *chk1Δ* Rad52-AID diploids do not arrest at the 12 hour mark like the *Chk1$^+$* Rad52-
AID diploids do but eventually seem to arrest at the 24 hour mark (Fig. S6E). These results
suggest that DNA damage and Chk1 activation upon Rad52 depletion passes the threshold for
cell cycle arrest in diploids but not haploids.

Discussion

In this study, we investigate how diploidy affects genome stability in the fission yeast
Schizosaccharomyces pombe. This organism is typically a haploid, and generally only forms
diploids as brief zygotes before entering meiosis. However, vegetative diploids can be recovered
in the laboratory using complementing markers. This is in contrast to budding yeast, which
alternates between diploid and haploid states.

In budding yeast, mating type heterozygosity affects DNA repair in diploids (Kadyk and
Hartwell 1992; Heude and Fabre 1993; Barbour and Xiao 2006). We compared *S. pombe*
diploids with different configurations of mating type loci from fully heterozygous, to fully
homozygous, and observed no difference in growth or sensitivity to genotoxins. All *S. pombe*
diploids exhibit increased sensitivity to a wide range of genotoxins regardless of mating type
heterozygosity (Fig. 1A). This result was unexpected as we hypothesized that having another set
of chromosomes would provide an additional template for homology directed repair and act as a
buffer for recessive mutations that would be deleterious in a haploid context. Early work in *S.
cerevisiae* showed that diploids exhibited increased resistance to irradiation compared to

haploids, but further increasing ploidy resulted in decreased resistance (Mortimer 1958). However, more recent work in budding yeast and humans found that tetraploids have increased sensitivity to replication stressors and double-strand break inducing agents (Storchová *et al.* 2006; Wangsa *et al.* 2018). Our results suggest that increase in ploidy from the natural genome state confers genomic instability in *S. pombe* as well, although in this case it is from haploid to diploid, rather than diploids to tetraploid as in budding yeast and humans.

We examined diploid response to the genotoxin methyl methanesulfonate (MMS). MMS is an alkylating agent that creates DNA adducts which leads to replication stalling (Szyjka *et al.* 2008; Iyer and Rhind 2017) and induces a replication stress transcriptional response in both haploids and diploids (Park and Forsburg 2024). However, the degree of transcriptional induction of damage responsive genes is somewhat attenuated in diploids (Park and Forsburg 2024). We found that an acute treatment of MMS reduces viability in diploids (Fig. 1C), consistent with an impaired damage response. When we look at DNA damage markers, we find that diploids have a slight increase in cells with Rad52 foci compared to haploids (Fig. 1E) but not RPA or Rad54 (Fig. 1D and 1F) in unperturbed conditions. This is in contrast with *S. cerevisiae* and human tetraploids that have increased DNA damage markers (Storchová *et al.* 2006; Gemble *et al.* 2022). Upon treatment of MMS, *S. pombe* diploids have increased DNA damage markers compared to untreated diploids (Fig. 1D-F). These results, taken together with the increased sensitivity of diploids to MMS, are consistent with *S. pombe* diploids having an impaired repair response and increased damage in MMS.

When yeast cells are treated with low levels of MMS, the cells will repair some of the lesions using an error-prone pathway involving translesion polymerases that can be measured using a forward mutation assay (Stelter and Ulrich 2003; Dolan *et al.* 2010). We observed no change in induced mutagenesis in diploids, but to our surprise, we found that our diploid strains exhibited high levels of loss of *ura4+* function even in untreated cells (Fig. 2A). We determined the cause to be partial loss of heterozygosity (Table S2). The LOH phenomenon was observed for all three *S. pombe* chromosomes (Fig. 2B) and occurred at similar rates when the markers were inserted telomere proximal and in the middle of the chromosome arm (Fig. 2C) suggesting a global effect. *S. cerevisiae* diploids have also been shown to experience spontaneous LOH using similar systems, but at rates that are 2-4 magnitudes lower than our results (Chen *et al.* 1999; Ohnishi *et al.* 2004; Storchová *et al.* 2006). LOH in *S. cerevisiae* diploids were a result of crossover and break-induced replication, non-crossover gene conversion events, and chromosome loss (Ohnishi *et al.* 2004; Suetomi *et al.* 2010; Chumki *et al.* 2016). The majority of the LOH events in our assay resulted in the loss of the telomere proximal marker but maintained the opposite arm of the chromosome, with only a small fraction of events reflecting whole chromosome mis-segregation (Fig. 2E). A recent study in diploid fission yeast showed interactions between homologous chromosomes after an induced break that was Rad51 dependent (Vines *et al.* 2022). This suggests that HDR uses homologous chromosomes, in addition to sister chromatids, for repair in fission yeast. We infer that the LOH events reflect recombination or BIR and speculate that these may be driven by fragile sites that are prone to replication stress.

We created *S. pombe*/*S. pombe var. kambucha* diploids, in which the chromosomes can be
distinguished by SNPs. We find there are two classes of LOH: a clear breakpoint at ~1.1 Mb
position and a more diffuse site closer to the centromere (Fig. 3B). The ~1.1 Mb breakpoint
coincides with a long terminal repeat in 7 of the 10 samples which may suggest a potential
fragile site that is prone to recombination (Fig. 3C, S1B). This may be similar in nature to the
known fragile site FS2 in *S. cerevisiae* associated with Ty1 elements that causes mitotic
recombination in diploid budding yeast (Rosen *et al.* 2013). The other three samples exhibited
much longer tracts of LOH, but did not share similar breakpoints between them. All three
samples did have LTR elements in the vicinity of the breakpoint. These potential fragile sites
may play a role in the increased endogenous replication stress that require HDR protein for
repair. However, further investigation is required to determine if these LTR elements are fragile
sites and if there are other sites in the *S. pombe* genome.

We see LOH rates increase in strains that lack replication checkpoint responders such as Mrc1 or
Cds1. Deletion or mutation of genes involved in DNA repair (*rad8Δ, rad16Δ, exo1Δ, srs2Δ,*
smc6-74) also increased LOH to varying degrees. Rad8 has been shown in *S. pombe* to play a
role in responding to stalled replication forks through post replication repair (PRR) pathways
(Frampton *et al.* 2006; Ding and Forsburg 2014). Knock out of Rad8 potentially drives repair
through HDR, resulting in the increase in LOH observed. Similarly, deletion of Srs2 and Exo1
have increased rates of mitotic recombination in *S. cerevisiae* and *S. pombe* resulting in LOH
(Wang *et al.* 2001; Ira *et al.* 2003; Lydeard *et al.* 2010; Marrero and Symington 2010).
Interestingly, *smc6-x* mutants were inviable diploids, but using the *smc6-74* allele, we could
construct diploids and observed elevated levels of LOH. The Smc5/6 complex has been
implicated in processing stalled replication forks (Morikawa *et al.* 2004; Pebernard *et al.* 2006)
and the difference in the two mutants may reflect a separation of function (Harvey *et al.* 2004)
that is emphasized in diploids. Deletion of the NHEJ protein Ku70 also resulted in increased
LOH. The ablation of NHEJ may drive increased repair by HDR resulting in mitotic
recombination and LOH (Maruyama *et al.* 2015; Li *et al.* 2018). *mgs1Δ* resulted in a slight
decrease of *ura4+* loss in our assay. Deletion of *MGS1* in *S. cerevisiae* haploids and diploids
caused an increase in recombination (Hishida *et al.* 2001) in contrast to our results. Studies on
Mgs1 in *S. pombe* are lacking and further investigation is required to elucidate the differences
observed between budding and fission yeast. We have generally found that genes involved in
responding to replication stress are either essential or lead to higher levels of LOH when deleted.
This suggests the *S. pombe* diploids experience increased endogenous replication stress that lead
to mitotic recombination and cell death if core HDR pathways are compromised.

As we were constructing the mutants for the LOH assay, we found that a subset of mutants could
be isolated but not maintained as diploids due to rapid haploidization which typically occurs
following whole chromosome loss (Bodi *et al.* 1991). These include a BIR-specific allele of
DNA polymerase delta (*cdc27-D1*; Tinline-Purvis *et al.* 2009), a helicase that dissolves HR
intermediates (*rqh1Δ*; Wu *et al.* 2006; Hope *et al.* 2007), and a member of the fork protection
complex (*swi1Δ*). The histone methyltransferase Clr4, which is required for efficient
chromosome segregation (Ekwall *et al.* 1996), also experienced high levels of haploidization but
maintained enough diploids for the experimental assay (Fig. S2). Interestingly, cell cycle mutants

cdc10-V50, cdc25-22, and *cdc2-asM17* could not be maintained as diploids even at permissive
temperatures, suggesting *S. pombe* diploids are very sensitive to cell cycle perturbations.

We did not find any mutants that dramatically reduced the rate of LOH. We suggest this is related
to the requirement of some genes for viability in diploids. The Rad52 epistasis group required for
homologous recombination, as well as spindle assembly checkpoint mutants, have previously
been reported to be inviable as diploids in fission yeast (Bernard *et al.* 1998; Catlett and
Forsburg 2003; Grishchuk *et al.* 2004; Octobre *et al.* 2008). Interestingly, screens for ploidy-
specific lethal genes in both budding yeast and human tetraploid cells have also identified HR
components, kinetochore, and SAC genes (Storchová *et al.* 2006; Quinton *et al.* 2021). These
studies suggested that kinetochore functions are particularly important in tetraploids and may not
scale effectively, but the role of the HR proteins has not been investigated. We speculate that the
LOH event is an essential process requiring HR proteins.

Using a candidate approach, we found that, in addition to the previously identified genes,
additional fission yeast PSL mutants include topoisomerase I (*top1Δ*), the structure specific
endonuclease *mu81Δ,* the helicase *fbh1Δ,* and the fork protection component *swi3Δ*. These are all
involved in resolving replication fork collapse (Shyian *et al.* 2020; Grabarczyk 2022; reviewed in
Chakraborty *et al.* 2023) and again suggest that diploid fission yeast have increased susceptibility
to replication stress. We propose that the LOH reflects diploids-specific fragile regions on the
chromosome.

Using an auxin-degron fusion to Rad52, we find that depletion of Rad52 leads to loss of viability,
with increased RPA foci and elongated cells, most with a single nucleus, consistent with cell
cycle arrest (Fig. 5B, 6, and S6). Although the haploids also exhibited persistent RPA foci, they
did not show any signs of cell cycle arrest and had a minor increase in abnormal mitosis. These
results suggest that haploids are capable of responding to the increased damage without inducing
arrest when Rad52 is depleted, but the diploids pass the threshold of DNA damage that is
manageable resulting in cell death.

Our previous transcriptome analysis suggested that most transcripts scale up ~2x in diploids
relative to haploids, but there may be modestly reduced expression of a few genes including the
polymerase subunit *cdc1* (Park and Forsburg 2024). It is possible that the transcription doesn't
lead to an appropriately scaled response of all replication factors. Overexpression of individual
diploid essential genes in fission yeast did not rescue MMS sensitivity (Fig. S4), suggesting a
more global increase is required to see a response. A screen in fission yeast diploids found that
heterozygous deletion of some replication factors, including *cdc1*, resulted in haploinsufficient
growth (Kim *et al.* 2010). Indeed, a recent report from human cells suggests that the first cell
cycle following tetraploidization is extremely prone to replication stress, due to insufficient
dosage of some replication factors (Gemble *et al.* 2022). We speculate that fission yeast diploids
may model this state.

Taken together, our data suggest that fission yeast diploidy has the same deleterious effects
observed in budding yeast or human tetraploidy (Storchová *et al.* 2006; Quinton *et al.* 2021;
Gemble *et al.* 2022). This suggests there is a conserved response to whole genome duplication

above normal ploidy. Since diploids are genetically more straightforward in analysis, this
establishes *S. pombe* as a useful model for polyploidy.

<u>Data Availability</u>

The sequencing data discussed in this publication have been deposited in NCBI's Sequence Read
Archive under BioProject ID PRJNA1216252. Strain list is available in Table S1. All strains and
reagents are available upon request.

Acknowledgements

We thank Jiping Yuan and Kassandra Martinez for their technical assistance and members of the
lab for helpful comments. We thank Dr. Nicholas Rhind and Dr. Sarah Zanders for their help and
generosity with strains and resources for the *S. pombe var. kambucha* project. We thank Dr.
Antony for providing strains for the auxin-inducible degron project. We thank Dr. Irene Chiolo
was allowing us to use the X-RAD iR-160 machine.

Competing interests

The authors declare no competing or financial interests.

Funding

This research was supported by the National Institute of General Medical Sciences [R35-
GM118109] (S.L.F) and University of Southern California Student Opportunities for Academic
Research (D.F.P).

<u>References</u>

Allshire, R. C., & Ekwall, K. (2015). Epigenetic Regulation of Chromatin States in
Schizosaccharomyces pombe. *Cold Spring Harbor perspectives in biology*, *7*(7), a018770.
https://doi.org/10.1101/cshperspect.a018770

Ampatzidou, E., Irmisch, A., O'Connell, M. J., & Murray, J. M. (2006). Smc5/6 is required for
repair at collapsed replication forks. *Molecular and cellular biology, 26*(24), 9387–9401.
https://doi.org/10.1128/MCB.01335-06

Archetti M. (2022). Polyploidy as an Adaptation against Loss of Heterozygosity in Cancer.
International journal of molecular sciences, *23*(15), 8528.
https://doi.org/10.3390/ijms23158528

Barbour, L., & Xiao, W. (2006). Mating type regulation of cellular tolerance to DNA damage is
specific to the DNA post-replication repair and mutagenesis pathway. *Molecular*
microbiology, *59*(2), 637–650. https://doi.org/10.1111/j.1365-2958.2005.04965.x

Bernard, P., Hardwick, K., & Javerzat, J. P. (1998). Fission yeast bub1 is a mitotic centromere
protein essential for the spindle checkpoint and the preservation of correct ploidy through

mitosis. *The Journal of cell biology, 143*(7), 1775–1787.
https://doi.org/10.1083/jcb.143.7.1775

Bharadwaj, D., & Mandal, M. (2020). Senescence in polyploid giant cancer cells: A road that
leads to chemoresistance. *Cytokine & growth factor reviews*, *52*, 68–75.
https://doi.org/10.1016/j.cytogfr.2019.11.002

Bielski, C. M., Zehir, A., Penson, A. V., Donoghue, M. T. A., Chatila, W., Armenia, J., Chang, M.
T., Schram, A. M., Jonsson, P., Bandlamudi, C., Razavi, P., Iyer, G., Robson, M. E., Stadler,
Z. K., Schultz, N., Baselga, J., Solit, D. B., Hyman, D. M., Berger, M. F., & Taylor, B. S.
(2018). Genome doubling shapes the evolution and prognosis of advanced cancers. *Nature*
genetics, *50*(8), 1189–1195. https://doi.org/10.1038/s41588-018-0165-1

Bodi, Z., Gysler-Junker, A., & Kohli, J. (1991). A quantitative assay to measure chromosome
stability in Schizosaccharomyces pombe. *Molecular & general genetics : MGG*, *229*(1),
77–80. https://doi.org/10.1007/BF00264215

Boulay, J. L., Stiefel, U., Taylor, E., Dolder, B., Merlo, A., & Hirth, F. (2009). Loss of
heterozygosity of TRIM3 in malignant gliomas. *BMC cancer*, *9*, 71.
https://doi.org/10.1186/1471-2407-9-71

Catlett, M. G., & Forsburg, S. L. (2003). Schizosaccharomyces pombe Rdh54 (TID1) acts with
Rhp54 (RAD54) to repair meiotic double-strand breaks. *Molecular biology of the*
cell, 14(11), 4707–4720. https://doi.org/10.1091/mbc.e03-05-0288

Chakraborty, S., Schirmeisen, K., & Lambert, S. A. (2023). The multifaceted functions of
homologous recombination in dealing with replication-associated DNA damages. *DNA*
repair, *129*, 103548. https://doi.org/10.1016/j.dnarep.2023.103548

Chen, C., Merrill, B. J., Lau, P. J., Holm, C., & Kolodner, R. D. (1999). Saccharomyces
cerevisiae pol30 (proliferating cell nuclear antigen) mutations impair replication fidelity
and mismatch repair. *Molecular and cellular biology*, *19*(11), 7801–7815.
https://doi.org/10.1128/MCB.19.11.7801

Chumki, S. A., Dunn, M. K., Coates, T. F., Mishler, J. D., Younkin, E. M., & Casper, A. M.
(2016). Remarkably Long-Tract Gene Conversion Induced by Fragile Site Instability in
Saccharomyces cerevisiae. *Genetics*, *204*(1), 115–128.
https://doi.org/10.1534/genetics.116.191205

Ciani, Y., Fedrizzi, T., Prandi, D., Lorenzin, F., Locallo, A., Gasperini, P., Franceschini, G. M.,
Benelli, M., Elemento, O., Fava, L. L., Inga, A., & Demichelis, F. (2022). Allele-specific
genomic data elucidate the role of somatic gain and copy-number neutral loss of
heterozygosity in cancer. *Cell systems, 13*(2), 183–193.e7.
https://doi.org/10.1016/j.cels.2021.10.001

Dewhurst, S. M., McGranahan, N., Burrell, R. A., Rowan, A. J., Grönroos, E., Endesfelder, D.,
Joshi, T., Mouradov, D., Gibbs, P., Ward, R. L., Hawkins, N. J., Szallasi, Z., Sieber, O. M.,
& Swanton, C. (2014). Tolerance of whole-genome doubling propagates chromosomal

560   instability and accelerates cancer genome evolution. *Cancer discovery*, *4*(2), 175–185.
561   https://doi.org/10.1158/2159-8290.CD-13-0285

562   Ding, L., & Forsburg, S. L. (2014). Essential domains of Schizosaccharomyces pombe Rad8
563   required for DNA damage response. *G3 (Bethesda, Md.)*, *4*(8), 1373–1384.
564   https://doi.org/10.1534/g3.114.011346

565   Doe, C. L., & Whitby, M. C. (2004). The involvement of Srs2 in post-replication repair and
566   homologous recombination in fission yeast. *Nucleic acids research*, *32*(4), 1480–1491.
567   https://doi.org/10.1093/nar/gkh317

568   Dolan, W. P., Le, A. H., Schmidt, H., Yuan, J. P., Green, M., & Forsburg, S. L. (2010). Fission
569   yeast Hsk1 (Cdc7) kinase is required after replication initiation for induced mutagenesis
570   and proper response to DNA alkylation damage. *Genetics, 185*(1), 39–53.
571   https://doi.org/10.1534/genetics.109.112284

572   Dutta, A., Dutreux, F., & Schacherer, J. (2022). Loss of Heterozygosity Spectrum Depends on
573   Ploidy Level in Natural Yeast Populations. *Molecular biology and evolution*, *39*(11),
574   msac214. https://doi.org/10.1093/molbev/msac214

575   Ekwall, K., Nimmo, E. R., Javerzat, J. P., Borgstrøm, B., Egel, R., Cranston, G., & Allshire, R.
576   (1996). Mutations in the fission yeast silencing factors clr4+ and rik1+ disrupt the
577   localisation of the chromo domain protein Swi6p and impair centromere function. *Journal*
578   *of cell science*, *109 (Pt 11)*, 2637–2648. https://doi.org/10.1242/jcs.109.11.2637

579   Ekwall, K., & Thon, G. (2017). Selecting *Schizosaccharomyces pombe* Diploids. *Cold Spring*
580   *Harbor protocols*, *2017*(7), pdb.prot091702. https://doi.org/10.1101/pdb.prot091702

581   Forsburg, S. L., & Nurse, P. (1991). Cell cycle regulation in the yeasts Saccharomyces cerevisiae
582   and Schizosaccharomyces pombe. *Annual review of cell biology*, *7*, 227–256.
583   https://doi.org/10.1146/annurev.cb.07.110191.001303

584   Forsburg, S. L., & Rhind, N. (2006). Basic methods for fission yeast. *Yeast (Chichester,*
585   *England)*, *23*(3), 173–183. https://doi.org/10.1002/yea.1347

586   Frampton, J., Irmisch, A., Green, C. M., Neiss, A., Trickey, M., Ulrich, H. D., Furuya, K., Watts,
587   F. Z., Carr, A. M., & Lehmann, A. R. (2006). Postreplication repair and PCNA modification
588   in Schizosaccharomyces pombe. *Molecular biology of the cell*, *17*(7), 2976–2985.
589   https://doi.org/10.1091/mbc.e05-11-1008

590   Fujiwara, T., Bandi, M., Nitta, M., Ivanova, E. V., Bronson, R. T., & Pellman, D. (2005).
591   Cytokinesis failure generating tetraploids promotes tumorigenesis in p53-null
592   cells. *Nature, 437*(7061), 1043–1047. https://doi.org/10.1038/nature04217

593   Gemble, S., Wardenaar, R., Keuper, K., Srivastava, N., Nano, M., Macé, A. S., Tijhuis, A. E.,
594   Bernhard, S. V., Spierings, D. C. J., Simon, A., Goundiam, O., Hochegger, H., Piel, M.,
595   Foijer, F., Storchová, Z., & Basto, R. (2022). Genetic instability from a single S phase after

whole-genome duplication. *Nature, 604*(7904), 146–151. https://doi.org/10.1038/s41586-022-04578-4

Grabarczyk D. B. (2022). The Fork Protection Complex: A Regulatory Hub at the Head of the Replisome. *Sub-cellular biochemistry*, *99*, 83–107. https://doi.org/10.1007/978-3-031-00793-4_3

Grallert, A., & Hagan, I. M. (2017). Preparation of Protein Extracts from Schizosaccharomyces pombe Using Trichloroacetic Acid Precipitation. *Cold Spring Harbor protocols*, 2017(2), 10.1101/pdb.prot091579. https://doi.org/10.1101/pdb.prot091579

Green, M. D., Sabatinos, S. A., & Forsburg, S. L. (2009). Microscopy techniques to examine DNA replication in fission yeast. Methods in molecular biology (Clifton, N.J.), 521, 463–482. https://doi.org/10.1007/978-1-60327-815-7_26

Grishchuk, A. L., Kraehenbuehl, R., Molnar, M., Fleck, O., & Kohli, J. (2004). Genetic and cytological characterization of the RecA-homologous proteins Rad51 and Dmc1 of Schizosaccharomyces pombe. *Current genetics, 44*(6), 317–328. https://doi.org/10.1007/s00294-003-0439-7

Harvey, S. H., Sheedy, D. M., Cuddihy, A. R., & O'Connell, M. J. (2004). Coordination of DNA damage responses via the Smc5/Smc6 complex. *Molecular and cellular biology*, *24*(2), 662–674. https://doi.org/10.1128/MCB.24.2.662-674.2004

Had ija, M. P., Kapitanović, S., Radosević, S., Cacev, T., Mirt, M., Kovacević, D., Cacev, T., Hadzija, M., Spaventi, R., & Pavelić, K. (2001). Loss of heterozygosity of DPC4 tumor suppressor gene in human sporadic colon cancer. *Journal of molecular medicine (Berlin, Germany)*, *79*(2-3), 128–132. https://doi.org/10.1007/s001090000179

Heil, CS. (2023). Loss of Heterozygosity and Its Importance in Evolution. *Journal of molecular evolution*, *91*(3), 369–377. https://doi.org/10.1007/s00239-022-10088-8

Heude, M., & Fabre, F. (1993). a/alpha-control of DNA repair in the yeast Saccharomyces cerevisiae: genetic and physiological aspects. *Genetics*, *133*(3), 489–498. https://doi.org/10.1093/genetics/133.3.489

Hishida, T., Iwasaki, H., Ohno, T., Morishita, T., & Shinagawa, H. (2001). A yeast gene, MGS1, encoding a DNA-dependent AAA(+) ATPase is required to maintain genome stability. *Proceedings of the National Academy of Sciences of the United States of America, 98*(15), 8283–8289. https://doi.org/10.1073/pnas.121009098

Hope, J. C., Cruzata, L. D., Duvshani, A., Mitsumoto, J., Maftahi, M., & Freyer, G. A. (2007). Mus81-Eme1-dependent and -independent crossovers form in mitotic cells during double-strand break repair in Schizosaccharomyces pombe. *Molecular and cellular biology*, *27*(10), 3828–3838. https://doi.org/10.1128/MCB.01596-06

Ira, G., Malkova, A., Liberi, G., Foiani, M., & Haber, J. E. (2003). Srs2 and Sgs1-Top3 suppress
crossovers during double-strand break repair in yeast. *Cell*, *115*(4), 401–411.
https://doi.org/10.1016/s0092-8674(03)00886-9

Irmisch, A., Ampatzidou, E., Mizuno, K., O'Connell, M. J., & Murray, J. M. (2009). Smc5/6
maintains stalled replication forks in a recombination-competent conformation. *The*
EMBO journal, 28(2), 144–155. https://doi.org/10.1038/emboj.2008.273

Iyer, D. R., & Rhind, N. (2017). Replication fork slowing and stalling are distinct, checkpoint-
independent consequences of replicating damaged DNA. *PLoS genetics*, *13*(8),
e1006958. https://doi.org/10.1371/journal.pgen.1006958

Jones, C. E., & Forsburg, S. L. (2021). Monitoring *Schizosaccharomyces pombe* genome stress
by visualizing end-binding protein Ku. *Biology open*, *10*(2), bio054346.
https://doi.org/10.1242/bio.054346

Kadyk, L. C., & Hartwell, L. H. (1992). Sister chromatids are preferred over homologs as
substrates for recombinational repair in Saccharomyces cerevisiae. *Genetics*, *132*(2),
387–402. https://doi.org/10.1093/genetics/132.2.387

Kim, D. U., Hayles, J., Kim, D., Wood, V., Park, H. O., Won, M., Yoo, H. S., Duhig, T., Nam,
M., Palmer, G., Han, S., Jeffery, L., Baek, S. T., Lee, H., Shim, Y. S., Lee, M., Kim, L.,
Heo, K. S., Noh, E. J., Lee, A. R., … Hoe, K. L. (2010). Analysis of a genome-wide set
of gene deletions in the fission yeast Schizosaccharomyces pombe. Nature biotechnology,
28(6), 617–623. https://doi.org/10.1038/nbt.1628

Knop M. (2006). Evolution of the hemiascomycete yeasts: on life styles and the importance of
inbreeding. *BioEssays : news and reviews in molecular, cellular and developmental*
biology, *28*(7), 696–708. https://doi.org/10.1002/bies.20435

Kuznetsova, A. Y., Seget, K., Moeller, G. K., de Pagter, M. S., de Roos, J. A., Dürrbaum, M.,
Kuffer, C., Müller, S., Zaman, G. J., Kloosterman, W. P., & Storchová, Z. (2015).
Chromosomal instability, tolerance of mitotic errors and multidrug resistance are
promoted by tetraploidization in human cells. *Cell cycle (Georgetown, Tex.)*, *14*(17),
2810–2820. https://doi.org/10.1080/15384101.2015.1068482

Li, G., Liu, D., Zhang, X., Quan, R., Zhong, C., Mo, J., Huang, Y., Wang, H., Ruan, X., Xu, Z.,
Zheng, E., Gu, T., Hong, L., Li, Z., Wu, Z., & Yang, H. (2018). Suppressing Ku70/Ku80
expression elevates homology-directed repair efficiency in primary fibroblasts. *The*
international journal of biochemistry & cell biology, *99*, 154–160.
https://doi.org/10.1016/j.biocel.2018.04.011

Lisby, M., Barlow, J. H., Burgess, R. C., & Rothstein, R. (2004). Choreography of the DNA
damage response: spatiotemporal relationships among checkpoint and repair
proteins. *Cell, 118*(6), 699–713. https://doi.org/10.1016/j.cell.2004.08.015

Liu, V. F., Bhaumik, D., & Wang, T. S. (1999). Mutator phenotype induced by aberrant
668   replication. *Molecular and cellular biology, 19*(2), 1126–1135.
669   https://doi.org/10.1128/MCB.19.2.1126

670   López, S., Lim, E. L., Horswell, S., Haase, K., Huebner, A., Dietzen, M., Mourikis, T. P.,
671   Watkins, T. B. K., Rowan, A., Dewhurst, S. M., Birkbak, N. J., Wilson, G. A., Van Loo,
672   P., Jamal-Hanjani, M., TRACERx Consortium, Swanton, C., & McGranahan, N. (2020).
673   Interplay between whole-genome doubling and the accumulation of deleterious
674   alterations in cancer evolution. *Nature genetics, 52*(3), 283–293.
675   https://doi.org/10.1038/s41588-020-0584-7

676   Lourenço, N., Hélias-Rodzewicz, Z., Bachet, J. B., Brahimi-Adouane, S., Jardin, F., Tran van
677   Nhieu, J., Peschaud, F., Martin, E., Beauchet, A., Chibon, F., & Emile, J. F. (2014). Copy-
678   neutral loss of heterozygosity and chromosome gains and losses are frequent in
679   gastrointestinal stromal tumors. *Molecular cancer*, *13*, 246. https://doi.org/10.1186/1476-
680   4598-13-246

681   Luche, D. D., & Forsburg, S. L. (2009). Cell-cycle synchrony for analysis of S. pombe DNA
682   replication. *Methods in molecular biology (Clifton, N.J.)*, *521*, 437–448.
683   https://doi.org/10.1007/978-1-60327-815-7_24

684   Lydeard, J. R., Lipkin-Moore, Z., Jain, S., Eapen, V. V., & Haber, J. E. (2010). Sgs1 and exo1
685   redundantly inhibit break-induced replication and de novo telomere addition at broken
686   chromosome ends. *PLoS genetics*, *6*(5), e1000973.
687   https://doi.org/10.1371/journal.pgen.1000973

688   Marrero, V. A., & Symington, L. S. (2010). Extensive DNA end processing by exo1 and sgs1
689   inhibits break-induced replication. *PLoS genetics*, *6*(7), e1001007.
690   https://doi.org/10.1371/journal.pgen.1001007

691   Maruyama, T., Dougan, S. K., Truttmann, M. C., Bilate, A. M., Ingram, J. R., & Ploegh, H. L.
692   (2015). Increasing the efficiency of precise genome editing with CRISPR-Cas9 by
693   inhibition of nonhomologous end joining. *Nature biotechnology, 33*(5), 538–542.
694   https://doi.org/10.1038/nbt.3190

695   Mayer, V. W., & Aguilera, A. (1990). High levels of chromosome instability in polyploids of
696   Saccharomyces cerevisiae. *Mutation research*, *231*(2), 177–186.
697   https://doi.org/10.1016/0027-5107(90)90024-x

698   Mimitou, E. P., & Symington, L. S. (2008). Sae2, Exo1 and Sgs1 collaborate in DNA double-
699   strand break processing. *Nature, 455*(7214), 770–774.
700   https://doi.org/10.1038/nature07312

701   Mirzayans, R., Andrais, B., & Murray, D. (2018). Roles of Polyploid/Multinucleated Giant
702   Cancer Cells in Metastasis and Disease Relapse Following Anticancer Treatment.
703   *Cancers*, *10*(4), 118. https://doi.org/10.3390/cancers10040118

Molnar, M., & Sipiczki, M. (1993). Polyploidy in the haplontic yeast Schizosaccharomyces
pombe: construction and analysis of strains. *Current genetics*, *24*(1-2), 45–52.
https://doi.org/10.1007/BF00324664

Morikawa, H., Morishita, T., Kawane, S., Iwasaki, H., Carr, A. M., & Shinagawa, H. (2004).
Rad62 protein functionally and physically associates with the smc5/smc6 protein
complex and is required for chromosome integrity and recombination repair in fission
yeast. *Molecular and cellular biology*, *24*(21), 9401–9413.
https://doi.org/10.1128/MCB.24.21.9401-9413.2004

Mortimer R. K. (1958). Radiobiological and genetic studies on a polyploid series (haploid to
hexaploid) of Saccharomyces cerevisiae. *Radiation research*, *9*(3), 312–326.

Nano, M., Gemble, S., Simon, A., Pennetier, C., Fraisier, V., Marthiens, V., & Basto, R. (2019).
Cell-Cycle Asynchrony Generates DNA Damage at Mitotic Entry in Polyploid
Cells. *Current biology : CB*, *29*(22), 3937–3945.e7.
https://doi.org/10.1016/j.cub.2019.09.041

Nichols, C. A., Gibson, W. J., Brown, M. S., Kosmicki, J. A., Busanovich, J. P., Wei, H.,
Urbanski, L. M., Curimjee, N., Berger, A. C., Gao, G. F., Cherniack, A. D., Dhe-Paganon,
S., Paolella, B. R., & Beroukhim, R. (2020). Loss of heterozygosity of essential genes
represents a widespread class of potential cancer vulnerabilities. *Nature communications*,
11(1), 2517. https://doi.org/10.1038/s41467-020-16399-y

Octobre, G., Lorenz, A., Loidl, J., & Kohli, J. (2008). The Rad52 homologs Rad22 and Rti1 of
Schizosaccharomyces pombe are not essential for meiotic interhomolog recombination,
but are required for meiotic intrachromosomal recombination and mating-type-related
DNA repair. *Genetics, 178*(4), 2399–2412. https://doi.org/10.1534/genetics.107.085696

Ohnishi, G., Endo, K., Doi, A., Fujita, A., Daigaku, Y., Nunoshiba, T., & Yamamoto, K. (2004).
Spontaneous mutagenesis in haploid and diploid Saccharomyces cerevisiae. *Biochemical*
and biophysical research communications, *325*(3), 928–933.
https://doi.org/10.1016/j.bbrc.2004.10.120

Park, J. M., & Forsburg, S. L. (2024). Analysis of transcriptional response in haploid and diploid
Schizosaccharomyces pombe under genotoxic stress. *G3 (Bethesda, Md.)*, *14*(9), jkae177.
https://doi.org/10.1093/g3journal/jkae177

Pebernard, S., Wohlschlegel, J., McDonald, W. H., Yates, J. R., 3rd, & Boddy, M. N. (2006). The
Nse5-Nse6 dimer mediates DNA repair roles of the Smc5-Smc6 complex. *Molecular and*
cellular biology, *26*(5), 1617–1630. https://doi.org/10.1128/MCB.26.5.1617-1630.2006

Quinton, R. J., DiDomizio, A., Vittoria, M. A., Kotýnková, K., Ticas, C. J., Patel, S., Koga, Y.,
Vakhshoorzadeh, J., Hermance, N., Kuroda, T. S., Parulekar, N., Taylor, A. M., Manning,
A. L., Campbell, J. D., & Ganem, N. J. (2021). Whole-genome doubling confers unique
genetic vulnerabilities on tumour cells. *Nature, 590*(7846), 492–497.
https://doi.org/10.1038/s41586-020-03133-3

Rhind, N., Chen, Z., Yassour, M., Thompson, D. A., Haas, B. J., Habib, N., Wapinski, I., Roy, S., Lin, M. F., Heiman, D. I., Young, S. K., Furuya, K., Guo, Y., Pidoux, A., Chen, H. M., Robbertse, B., Goldberg, J. M., Aoki, K., Bayne, E. H., Berlin, A. M., … Nusbaum, C. (2011). Comparative functional genomics of the fission yeasts. Science (New York, N.Y.), 332(6032), 930–936. https://doi.org/10.1126/science.1203357

Rosen, D. M., Younkin, E. M., Miller, S. D., & Casper, A. M. (2013). Fragile site instability in Saccharomyces cerevisiae causes loss of heterozygosity by mitotic crossovers and break-induced replication. *PLoS genetics*, *9*(9), e1003817. https://doi.org/10.1371/journal.pgen.1003817

Sabatinos, S. A., & Forsburg, S. L. (2010). Molecular genetics of Schizosaccharomyces pombe. *Methods in enzymology, 470*, 759–795. https://doi.org/10.1016/S0076-6879(10)70032-X

Sabatinos, S. A., Green, M. D., & Forsburg, S. L. (2012). Continued DNA synthesis in replication checkpoint mutants leads to fork collapse. *Molecular and cellular biology*, *32*(24), 4986–4997. https://doi.org/10.1128/MCB.01060-12

Saugar, I., Parker, J. L., Zhao, S., & Ulrich, H. D. (2012). The genome maintenance factor Mgs1 is targeted to sites of replication stress by ubiquitylated PCNA. *Nucleic acids research, 40*(1), 245–257. https://doi.org/10.1093/nar/gkr738

Shyian, M., Albert, B., Zupan, A. M., Ivanitsa, V., Charbonnet, G., Dilg, D., & Shore, D. (2020). Fork pausing complex engages topoisomerases at the replisome. *Genes & development*, *34*(1-2), 87–98. https://doi.org/10.1101/gad.331868.119

Steele, C. D., Abbasi, A., Islam, S. M. A., Bowes, A. L., Khandekar, A., Haase, K., Hames-Fathi, S., Ajayi, D., Verfaillie, A., Dhami, P., McLatchie, A., Lechner, M., Light, N., Shlien, A., Malkin, D., Feber, A., Proszek, P., Lesluyes, T., Mertens, F., Flanagan, A. M., … Pillay, N. (2022). Signatures of copy number alterations in human cancer. *Nature*, *606*(7916), 984–991. https://doi.org/10.1038/s41586-022-04738-6

Stelter, P., & Ulrich, H. D. (2003). Control of spontaneous and damage-induced mutagenesis by SUMO and ubiquitin conjugation. *Nature*, *425*(6954), 188–191. https://doi.org/10.1038/nature01965

Storchová, Z., Breneman, A., Cande, J., Dunn, J., Burbank, K., O'Toole, E., & Pellman, D. (2006). Genome-wide genetic analysis of polyploidy in yeast. *Nature*, *443*(7111), 541–547. https://doi.org/10.1038/nature05178

Storchova Z. (2014). Ploidy changes and genome stability in yeast. Yeast (Chichester, England), 31(11), 421–430. https://doi.org/10.1002/yea.3037

Styrkársdóttir, U., Egel, R., & Nielsen, O. (1993). The smt-0 mutation which abolishes mating-type switching in fission yeast is a deletion. *Current genetics*, *23*(2), 184–186. https://doi.org/10.1007/BF00352020

Suetomi, K., Mochizuki, M., Suzuki, S., Yamamoto, H., & Yamamoto, K. (2010). Effects of
Saccharomyces cerevisiae mec1, tel1, and mre11 mutations on spontaneous and
methylmethane sulfonate-induced genome instability. *Genes & genetic systems*, *85*(1), 1–
8. https://doi.org/10.1266/ggs.85.1

Sui, Y., Qi, L., Wu, J. K., Wen, X. P., Tang, X. X., Ma, Z. J., Wu, X. C., Zhang, K., Kokoska, R.
J., Zheng, D. Q., & Petes, T. D. (2020). Genome-wide mapping of spontaneous genetic
alterations in diploid yeast cells. *Proceedings of the National Academy of Sciences of the
United States of America*, *117*(45), 28191–28200.
https://doi.org/10.1073/pnas.2018633117

Szyjka, S. J., Aparicio, J. G., Viggiani, C. J., Knott, S., Xu, W., Tavaré, S., & Aparicio, O. M.
(2008). Rad53 regulates replication fork restart after DNA damage in Saccharomyces
cerevisiae. *Genes & development*, *22*(14), 1906–1920.
https://doi.org/10.1101/gad.1660408

Tanaka, K., & Russell, P. (2001). Mrc1 channels the DNA replication arrest signal to checkpoint
kinase Cds1. *Nature cell biology*, *3*(11), 966–972. https://doi.org/10.1038/ncb1101-966

Tanaka, T., Yokoyama, M., Matsumoto, S., Fukatsu, R., You, Z., & Masai, H. (2010). Fission
yeast Swi1-Swi3 complex facilitates DNA binding of Mrc1. *The Journal of biological
chemistry*, *285*(51), 39609–39622. https://doi.org/10.1074/jbc.M110.173344

Tinline-Purvis, H., Savory, A. P., Cullen, J. K., Davé, A., Moss, J., Bridge, W. L., Marguerat, S.,
Bähler, J., Ragoussis, J., Mott, R., Walker, C. A., & Humphrey, T. C. (2009). Failed gene
conversion leads to extensive end processing and chromosomal rearrangements in fission
yeast. *The EMBO journal*, *28*(21), 3400–3412. https://doi.org/10.1038/emboj.2009.265

Vines, A. J., Cox, K., Leland, B. A., & King, M. C. (2022). Homology-directed repair involves
multiple strand invasion cycles in fission yeast. *Molecular biology of the cell*, *33*(4), ar30.
https://doi.org/10.1091/mbc.E20-07-0433

Wang, S. W., Goodwin, A., Hickson, I. D., & Norbury, C. J. (2001). Involvement of
Schizosaccharomyces pombe Srs2 in cellular responses to DNA damage. *Nucleic acids
research*, *29*(14), 2963–2972. https://doi.org/10.1093/nar/29.14.2963

Wangsa, D., Quintanilla, I., Torabi, K., Vila-Casadesús, M., Ercilla, A., Klus, G., Yuce, Z.,
Galofré, C., Cuatrecasas, M., Lozano, J. J., Agell, N., Cimini, D., Castells, A., Ried, T., &
Camps, J. (2018). Near-tetraploid cancer cells show chromosome instability triggered by
replication stress and exhibit enhanced invasiveness. *FASEB journal : official publication
of the Federation of American Societies for Experimental Biology*, *32*(7), 3502–3517.
https://doi.org/10.1096/fj.201700247RR

Watson, A. T., Hassell-Hart, S., Spencer, J., & Carr, A. M. (2021). Rice (Oryza sativa) TIR1 and
5'adamantyl-IAA Significantly Improve the Auxin-Inducible Degron System
in Schizosaccharomyces pombe. *Genes, 12*(6), 882. https://doi.org/10.3390/genes12060882

Willer, M., Hoffmann, L., Styrkársdóttir, U., Egel, R., Davey, J., & Nielsen, O. (1995). Two-step
activation of meiosis by the mat1 locus in Schizosaccharomyces pombe. *Molecular and*
cellular biology, *15*(9), 4964–4970. https://doi.org/10.1128/MCB.15.9.4964

Wu, L., Bachrati, C. Z., Ou, J., Xu, C., Yin, J., Chang, M., Wang, W., Li, L., Brown, G. W., &
Hickson, I. D. (2006). BLAP75/RMI1 promotes the BLM-dependent dissolution of
homologous recombination intermediates. *Proceedings of the National Academy of*
Sciences of the United States of America, *103*(11), 4068–4073.
https://doi.org/10.1073/pnas.0508295103

Yin, L., Zhang, H., Tang, Z., Xu, J., Yin, D., Zhang, Z., Yuan, X., Zhu, M., Zhao, S., Li, X., &
Liu, X. (2021). rMVP: A Memory-efficient, Visualization-enhanced, and Parallel-
accelerated Tool for Genome-wide Association Study. *Genomics, proteomics &*
bioinformatics, *19*(4), 619–628. https://doi.org/10.1016/j.gpb.2020.10.007

Zack, T. I., Schumacher, S. E., Carter, S. L., Cherniack, A. D., Saksena, G., Tabak, B., Lawrence,
M. S., Zhsng, C. Z., Wala, J., Mermel, C. H., Sougnez, C., Gabriel, S. B., Hernandez, B.,
Shen, H., Laird, P. W., Getz, G., Meyerson, M., & Beroukhim, R. (2013). Pan-cancer
patterns of somatic copy number alteration. *Nature genetics*, *45*(10), 1134–1140.
https://doi.org/10.1038/ng.2760

Zhang, X., & Sjöblom, T. (2021). Targeting Loss of Heterozygosity: A Novel Paradigm for
Cancer Therapy. *Pharmaceuticals (Basel, Switzerland)*, *14*(1), 57.
https://doi.org/10.3390/ph14010057

Table 1	List of genes tested for ploidy-specific lethality	
	Gene	**Function**
Diploid - Essential		
	Rad51	Homologous recombination
	Rad52	Homologous recombination
	Rad54	Homologous recombination
	Mre11	Resection
	Fbh1	Helicase
	Mus81	Structure specific endonuclease
	Crb2	DNA end binding protein; checkpoint
	Top1	Topoisomerase
	Swi3 (TIPIN)	Fork protection complex
	Bub1	Spindle assembly checkpoint
	Mad2	Spindle assembly checkpoint
	Mad3	Spindle assembly checkpoint
Diploid - Unstable		
	Rqh1 (BLM)	Helicase
	cdc27-D1	Polymerase delta subunit required for BIR
	smc6-x	DNA repair; checkpoint
	Swi1(TIMELESS)	Fork protection complex
	cdc2-asM17	Cell cycle;
	cdc10-V50	Cell cycle; MluI binding factor
	cdc25-22	Cell cycle; phosphatase
Diploid - Dispensable		
	Rad3 (ATR)	Checkpoint
	Cds1	Checkpoint
	Tel1 (ATM)	Checkpoint
	Chk1	Checkpoint
	Mrc1 (CLSPN)	Fork protection complex
	Ku70	Non-homologous end joining
	Lig4	Non-homologous end joining
	Rad8	HLTF helicase; ubiquitin ligase
	Rad16 (XPF)	Structure specific endonuclease
	Mgs1 (WRNIP1)	DNA-dependent ATPase
	Exo1	Long range resection
	Srs2	Helicase; Rad51 regulator
	smc6-74	DNA repair; checkpoint
	Clr4	Lysine methyltransferase
	*Fml1 (FANCM)**	Helicase
	*Fml1/Fml2**	Helicase
	*top2-191**	Topoisomerase
	*Rev3**	Translesion synthesis polymerase
	*Eso1/Kpa1/Rev1/Rev3**	Translesion synthesis polymerases
	*Mus7**	DNA repair

List of genes that were tested for ploidy-specific lethality in *S.pombe* diploids. Diploid – essential genes were genes
whose deletions or mutations were not able to be isolated as homozygous diploids by crosses or protoplast fusion.
Diploid – unstable genes were genes whose deletions or mutations were able to be isolated as homozygous diploids,
but failed to maintain diploidy over time (i.e. overnight culturing or freezing to add to collection resulted in
haploidy). Diploid – dispensable genes were genes that were able to form stable diploids as homozygous deletions
or mutations. Genes in the Diploid – dispensable category with asterisks (*) were not included in LOH assay.

Table S2	Percent loss of *ura4+* in diploids	
Strain	**Treatment**	**% *ura4Δ***
Haploid	0.0025% MMS	0% (0/12)
Mat2-102 het.	No treatment	87% (20/23)
Mat2-102 het.	0.0025% MMS	100% (22/22)
Proto. het.	No treatment	96% (23/24)
Proto. het.	0.0025% MMS	100% (24/24)

Percent of 5-FOA resistant colonies that had completely lost the *ura4+* marker resulting in loss of heterozygosity.

Colonies that maintained heterozygosity of *ura4+* have presumably lost *ura4+* function by mutation.

Figure 1. Diploids are sensitive to genotoxic stress.

A) Serial dilution growth assay of haploid and diploid *S. pombe* strains. Diploid strains
constructed by mating were crossed fresh before plating. Strains were spotted onto YES plates
containing drugs as noted in the figure (HU=hydroxyurea, Phleo=phleomycin,
CPT=camptothecin, MMS= methane methylsulfonate). Plates were incubated at 32ºC for 3 days
then imaged. B) Graph of haploid and diploid survival after irradiation. Statistical analysis was
performed as mentioned in Materials and Methods. C) Graph of haploid and diploid survival
after acute MMS treatment. h^+/h^- diploid was not included in this assay and future assays as there
were no phenotypic differences with the Mat2-102 diploid, which is sufficient to control for
mating type heterozygosity. D) Graph of RPA foci (*rad11-Cerulean fluorescent protein* [CFP])
after treatment with 0.0075% MMS, six replicates. E) Graph of Rad52 foci (*rad22-CFP*) after
treatment with MMS, six replicates. F) Graph of Rad54 foci (*rad54-Green fluorescent protein*
[GFP]) after treatment with MMS, nine replicates. Mat2-102 diploid was not included in this
assay as there were no phenotypic differences with the protoplast fusion diploid.

Figure 2. Diploids experience LOH in the absence of exogenous stress.

A) Graph of induced mutagenesis in diploid strains that were heterozygous or homozygous for
endogenous *ura4+* marker, six replicates. B) Graph of *ura4+* loss in diploid strains with *ura4+*
inserted in chromosome I or II as heterozygous. *ura4+* was inserted ~1.1 Mb downstream of the
telomere of each chromosome compared to ~150 kb downstream of the telomere on the
endogenous chromosome III. C) Graph comparing *ura4+* loss when the marker was placed
telomere proximal (tel.), middle (mid), or centromere proximal (cen.) on chromosome I, nine
replicates. Middle marker was inserted 1.4 Mb downstream of the telomere proximal marker and
the centromere proximal marker was inserted 1.2 Mb downstream of the middle marker. Marker
on the right arm of chromosome I was inserted 800 kb downstream of the centromere. D-F)
Graph showing loss of the other markers in the population that had lost the *ura4+* marker, nine
replicates.

Figure 3. Sequencing of LOH samples show two distinct breakpoint sites

A) Schematic of hybrid strain construction. B) SNP density plots depicting number of SNPs per
2.5 kb window across Chromosome I. C) Enlargement of Chromosome I at the 1.1 Mb
breakpoint showing relevant gene features and SNPs of sample 10.

Figure 4. LOH in diploids is influenced by replication stress response and DNA repair genes.

A) Bar graph showing rate of *ura4+* loss in homozygous deletion and mutant diploid strains.
Bars represent mean and error bars represent standard deviation. Wild type (WT) had fifty four
replicates. Mutant strains had seven to fourteen replicates due to picking colonies that already
experienced LOH (both loss and homozygosing of the *ura4+* marker). B) Bar graph depicting
rate of *natMX* loss (middle marker). C) Bar graph depicting rate of *bleMX* loss (right arm
marker) indicating whole chromosome loss.

Figure 5. Depletion of Rad52 and Mad2 using the AID system results in loss of viability in
diploids.

A) Immunoblot and quantification of Rad52 degradation in Rad52-AID homozygous diploids.
Cells were treated with 100 nM 5'a-IAA and sampled at the indicated time points. Cdc2 was
used as the loading control. B) Control and Rad52-AID strains were spotted onto YES plates
with and without 100 nM 5'a-IAA and/or 2 mM HU. Plates were incubated at 32ºC for 3 days
then imaged. C-D) Immunoblot and quantification of Mad2 degradation in Mad2-AID haploids
and Mad2-AID homozygous diploids. Conditions were the same as the Rad52 assay. E) Control
and Mad2-AID strains were spotted onto YES plates with and without 100 nM 5'a-IAA and/or
10 µg/mL thiabendazole (TBZ). Plates were incubated at 32ºC for 3 days then imaged.

Figure 6. Depletion of Rad52 leads to increased genome instability in diploids.

A) Graph of RPA foci (*ssb2-mCherry*) in haploid and diploid Rad52-AID strains after 5'a-IAA
treatment. Samples were collected at the time points indicated and imaged by fluorescent
microscopy, nine replicates per time point. B) Graph showing percent increase in optical density
(OD; 595 nm) relative to the time point before 5'a-IAA addition. Time points were taken every
hour for ten hours, three replicates. C) Graph showing percent increase in cell count relative to
time point before 5'a-IAA addition. Samples were from the same batch used for OD reading. D-
E) Immunoblot of Chk1 activation in Rad52-AID haploids and Rad52-AID homozygous
diploids. Cells were treated with 100 nm 5'a-IAA and sampled at the indicated times. Alpha
tubulin was used as the loading control. F) Graph showing percent increase in abnormal mitotic
events measured using a histone marker tagged with a red fluorescent protein (*hht1-RFP*).
Samples were collected at the time points indicated and imaged by fluorescent microscopy, nine
replicates per time point. G) Graph showing percent increase in abnormal mitotic events
measured using DAPI stain on fixed cell samples collected at the time points indicated. Nine
replicates per time point.

Figure S1. LOH rate of *S. pombe/S. pombe var. kambucha* hybrid strain

A) Graph showing the *Ura4+* loss rate in isogenic *S. pombe* diploids and hybrid *S. pombe/S.*
pombe var. kambucha diploids. B) Enlargement of Chromosome I at the breakpoints of samples
1-9 showing relevant gene features and SNPs.

Figure S2. Haploidization rates of mutant diploid strains.

Bar graph showing haploidization rates in the mutant diploid strains used in the LOH assay.
Haploids were measured as lighter pink colonies compared to the darker pink colonies when
plated on phloxine B plates. Lighter colonies were ratioed to total number of colonies counted.

Figure S3. MMS sensitivity assay of the LOH mutant panel.

Serial dilution growth assay of mutant haploid and diploid strains used in the LOH assay. Strains
were spotted onto pombe minimal glutamate (PMG) media supplemented with histidine, uracil,
leucine, and adenine without MMS or with MMS at the concentrations as noted in the figure.
Plates were incubated at 32ºC for 3 days then imaged.

Figure S4. Overexpression of diploid essential genes does not rescue MMS sensitivity

Serial dilution growth assay of haploid and diploid strains overexpressing diploid essential genes
listed in Table 1. Yeast strains were transformed with a "no message in thiamine" (NMT)
promoter vector controlling the expression of the gene of interest or an empty vector (EV) as
control. Transformed yeast strains were spotted onto PMG media supplemented with histidine
and leucine to maintain the vector (uracil selection) and ploidy (adenine selection). Thiamine
was added to a final concentration of 5 μg/mL to repress expression or omitted to induce
expression of the gene of interest. MMS was added at the concentrations as noted in the figure.
Plates were incubated at 32°C for 3 days then imaged.

Figure S5. Rad52-AID strains exhibit MMS sensitivity in the absence of degradation.

Serial dilution growth assay of Rad52-AID haploid and diploids. Strains were spotted onto YES
containing MMS at the indicated concentrations without 5'a-IAA. Plates were incubated at 32°C
for 3 days then imaged.

Figure S6. Diploids exhibit abnormal nuclear and cell phenotypes when Rad52 is depleted.

A) Stacked bar graph showing distribution of single, multiple, and large RPA (*ssb2-mCherry*)
foci ratios in haploid and diploid Rad52-AID strains after 5'a-IAA addition. Two or more RPA
foci were categorized as multiple foci and RPA foci that were amorphous and larger on average
were categorized as large foci. B) Representative image of diploid cell with RPA coated bridge at
eight hours after 5'a-IAA addition. C) Representative image of diploid cells with abnormal
segregation of DNA twelve hours after 5'a-IAA addition. D) Representative brightfield image of
diploid cells with abnormal septation patterns twenty four hours after 5'a-IAA addition. E)
Representative brightfield images of *Chk1*$^+$ and *chk1Δ* Rad52-AID cells at the time points
indicated.

A

Chromosome I

Ura4+ ●———— *S. pombe*

X

●———— *S. pombe var. kambucha*

B

Chromosome I

0Mb 0.6Mb 1.2Mb 1.8Mb 2.4Mb 3Mb 3.6Mb 4.2Mb 4.8Mb 5.4Mb

Control

Sample 1

Sample 2

Sample 3

Sample 4

Sample 5

Sample 6

Sample 7

Sample 8

Sample 9

Sample 10

Number of SNPs per
2.5-kb window

0
≤ 10
11-29
≥ 30

C

1.09Mb   1.13Mb

Sample 10

Origin of replication

Coding DNA sequence

Long terminal repeat

Single nucleotide polymorphism

A

B

C

A ssb2-mCherry Rad52-AID

B Rad52-AID

C Rad52-AID

F hht1-mRFP Rad52-AID

D Haploid Rad52-AID-V5
Chk1-Myc

G *chk1Δ* Rad52-AID

E Diploid Rad52-AID-V5/Rad52-AID-V5
Chk1-Myc/Chk1-Myc