Effective *in vivo* RNA base editing *via* engineered cytidine deaminase APOBECs

fused with PUF proteins

Wenjian Han [1, #, *], Bo Yuan [2, #], Xiaojuan Fan [3], Weike Li [4], Yiting Yuan [1], Yuefang

Zhang[1], Shu Wang [1], Shifang Shan [4], Markus Hafner [3], Zefeng Wang [5, 6, *] & Zilong Qiu

[1, 4, *]

[1] Department of Neurology, Songjiang Research Institute, Songjiang Hospital & MOE-

Shanghai Key Laboratory for Children's Environmental Health, Shanghai Jiao Tong

University School of Medicine, Shanghai, China

[2] Institute of Medical Genetics and Genomics, Fudan University, Shanghai, China

[3] RNA Molecular Biology Laboratory, National Institute of Arthritis and

Musculoskeletal and Skin Disease, Bethesda, MD, USA.

[4] Institute of Neuroscience, Center for Excellence in Brain Science and Intelligence

Technology, Chinese Academy of Sciences, Shanghai, China

[5] Shanghai Institute of Nutrition and Health, Chinese Academy of Sciences, Shanghai,

China

[6] Southern University of Science and Technology, Shenzhen, Guangdong, China

[#] These authors contributed equally to this work.

*Co-corresponding authors: hanwenjian@shsmu.edu.cn (W.H.),

wangzf@sustech.edu.cn (Z.W.), qiuzilong@shsmu.edu.cn (Z.Q.)

Summary

Base editing stands at the forefront of genetic engineering, heralding precise genetic modifications with broad implications. While CRISPR-based DNA and RNA base editing systems capitalize on sgRNA-guided specificity and diverse deaminase functionalities, the pursuit of efficient C-to-U RNA editing has been hampered by the inherent constraints of cytidine deaminases. Here, we report an RNA base editing platform by refining cytidine deaminases, termed professional APOBECs (ProAPOBECs), through systematic enhancements and AI-driven protein engineering. ProAPOBECs demonstrate unprecedented catalytic versatility, particularly fused with RNA-recognizing Pumilio and FBF (PUF) proteins. We demonstrated that *in vivo* RNA base editing of *Pcsk9* using ProAPOBECs effectively lowers cholesterol levels in mice. Additionally, AAV-mediated RNA base editing with ProAPOBECs in the brain of an autism mouse model not only corrected point mutations in *Mef2c* mRNAs but also significantly alleviated disease-associated phenotypes. This work introduces a pioneering collection of RNA base editing instruments, emphasizing their therapeutic potential in combatting genetic disorders.

Introduction

CRISPR-based base editing offers a potent method for editing genetic material by directly modifying single base pair in DNA or RNA. This approach, involving C-to-U (cytidine-to-uridine) and A-to-I (adenosine-to-inosine) transitions mediated by deaminases, avoids the strand breaks typically induced by the CRISPR/Cas system [1,2]. Initially developed for DNA modification, CRISPR/Cas-based targeting has become an effective tool in genetic research and holds potential for therapeutic applications, such as correcting disease-associated mutations in animal models [3-9].

Unlike the permanent modifications made by DNA base editing [10-12], RNA base editing offers a reversible method for *in vivo* base alternations, providing a relatively safer avenue for therapeutic strategies [13-17]. The A-to-I transition in RNA, mediated by Adenosine Deaminases Acting on RNA (ADAR), has been effectively demonstrated *in vivo* [18-20]. Two evolved ADAR2-based editing systems, RESCUE-S and xCBE, have been developed to achieve both A-to-I and C-to-U RNA editing [21,22]. However, RESCUE-S is strongly biased against editing G<u>C</u> and C<u>C</u> motifs, limiting its versatility [23]. To address this limitation, Stafforst's group introduced the SNAP-CDAR-S system, a SNAP-tag-based tool designed to edit C<u>C</u> motifs in specific RNA targets [24]. While promising, evolved ADAR2-mediated base editing systems, including SNAP-CDAR-S, exhibit simultaneous C-to-U and A-to-I off-target editing, imposing constraints on their broader application [21-24]. Engineering Apolipoprotein B mRNA editing enzyme catalytic polypeptide (APOBEC) family proteins offers a potential solution. These

proteins are specifically tailored for precise C-to-U base editing, effectively eliminating

the A-to-I off-target effects and reducing undesired side effects. This approach

represents a significant advancement in RNA base editing technology.

More recently, an RNA base editing system called CURE has been developed

utilizing the dCasRx protein and a native APOBEC3A, to achieve limited yet specific

U<u>C</u> transitions [23]. However, the specific Cas13-based C-to-U transition, has not

achieved for efficiently correcting disease-causing mutations in mouse models [23,25].

One potential reason is the strong preference of APOBEC for single-stranded RNA,

conflicting with the double-stranded structures formed by target mRNA and small guide

RNA used in the CRISPR/Cas system [26].

To address existing limitations, the gRNA-free system, REWIRE (RNA editing

with individual RNA-binding enzyme), was developed by exploiting the programmable

RNA targeting capabilities of PUF proteins [27]. PUF proteins feature 8- or 10- repeat

motifs, each of which can be programmed to specifically bind any RNA base *via*

interaction with the Watson–Crick edge [28-32]. Various groups have developed PUF-

based systems incorporating different functional domains to manipulate RNA splicing

[33], translation [34], degradation [35] and methylation [36]. Given the versatility of PUF domain

in recognizing nearly any short 8- or 10- nucleotide RNA sequences, its combination

with ADAR or APOBEC enzymes allows REWIRE to perform precise and efficient A-

to-I or C-to-U editing on specific nucleosides of RNA targets in cultured cells [27].

Notably, the use of mammalian-derived proteins in the REWIRE minimizes the

potential immune responses often triggered by bacterial Cas enzymes in therapeutic

base editing scenarios [37].

Current C-to-U RNA base editing systems rely on the enzymatic activity of APOBEC

to hydrolytically deaminate cytidine [13,27]. Natural APOBECs, however, often exhibit

deamination activity in a sequence context-dependent manner, primarily with single-

strand DNA or RNA [38]. For instance, Human APOBEC3A preferentially edits cytidine

within UC motifs in mRNA [23,27]. These context preferences limited further application

of RNA base editors.

APOBEC enzymes, recognized for their critical role in inducing mutations within

the genomes of retroviruses during infection, constitute a significant group of antiviral

genes. Current NCBI database has documented over 8,231 eukaryotic genes

homologous to the AID/APOBEC family, highlighting their extensive presence and

diversity in eukaryotic organisms [39]. These AID/APOBEC proteins share a common

domain configuration, with a highly conserved catalytic domain centered between the

variable N-terminal (NTD) and C-terminal domain (CTD) [39-41], which may enable a

rational design of novel cytidine deaminases using artificial intelligence-assisted

protein engineering.

In this study, we use AlphaFold2-mediated structural engineering to develop

Professional APOBECs (ProAPOBECs) with greatly expanded C-to-U editing capability. When integrated within the REWIRE system, ProAPOBECs demonstrate improved specificity and efficiency in multiple sequence contexts, including G<u>C</u>, C<u>C</u>, A<u>C</u>, and U<u>C</u>.

Importantly, we achieved effective *in vivo* C-to-U RNA editing in the liver and brain of mice using the CU-REWIRE5s with ProAPOBECs delivered *via* adeno-associated virus (AAV). This *in vivo* RNA editing mediated by CU-REWIRE successfully reducing cholesterol levels in mice by targeting Pcsk9 mRNA and alleviated autistic-like behaviors by correcting a point mutation in Mef2c mRNA in an autism spectrum disorder (ASD) mouse model [8]. In conclusion, this study demonstrated that the AI-assisted protein engineering can be applied to refine the activity of RNA base editors, enabling the correction of autistic-like phenotype in mouse model and paving the way for the practical applications of RNA base editing in the realm of gene therapy.

Results

Enhanced Stability and Efficacy of CU-REWIRE *via* Structural Optimization of PUF Domain

Our initial efforts focused on improving CU-REWIRE3.0, which combined a 10-repeat PUF domain (PUF10) with the cytidine deaminase enzyme APOBEC3A. Structural analysis revealed high similarity between the fourth repeat (R4) of PUF10 in CU-REWIRE3.0 and the fifth repeat (R5) of the native Pumilio 2 protein. Importantly, R4

lacked the Leucine-Proline (LP) peptide present in R5, a potentially crucial element for

structural flexibility as suggested by the previous research [42]. This insight led us to

engineer an enhanced PUF10 (ePUF10) with the LP peptide integrated into R4,

resulting in the development of CU-REWIRE4.0 (Fig. 1a and Supplementary Fig. 1a).

Comparative analysis showed that CU-REWIRE4.0 exhibited higher expression

levels than its predecessor, suggesting improved stability due to the LP insertion (Fig.

1b). We then assessed the editing efficacy of CU-REWIRE4.0 by targeting the C-to-U

transition on the C_{459} site in the Enhanced Green Fluorescent Protein (EGFP) mRNA.

As judged by mRNA sequencing (mRNA-seq), the CU-REWIRE4.0 achieved a

substantial increase in editing efficiency, with an 82.3% success rate compared to 69.7%

for CU-REWIRE3.0 (Supplementary Fig. 1b).

To evaluate off-target effects, we conducted RNA-seq analysis with 50X

transcriptome coverage (See Methods). Four independent sample sets (APOBEC3A,

ePUF10, CU-REWIRE3.0, and CU-REWIRE4.0) were analyzed with mRNA-seq,

revealing 224 potential off-target events for CU-REWIRE3.0 and 731 for CU-

REWIRE4.0 (Supplementary Fig. 1c-e). Notably, none of the off-target sites were

within 20-nt downstream of ePUF10-binding sequences (Supplementary Fig. 1e),

suggesting a high precision of ePUF10 in recognizing target and a significantly lower

off-target rate compared to the earlier CU-REWIREs [27]. However, off-target events

were largely attributed to the basal activity of APOBECs, highlighting the need for

further optimization.

While showing promise in editing efficiency, the original CU-REWIRE3.0 was

restricted to C-to-U editing activity within the UC context [27]. Similar to its predecessors,

we found that the CU-REWIRE4.0 maintained the preference for editing cytidines

within the UC consensus motif (Supplementary Fig. 1f). Its precision was further

improved in that the C-to-U conversions predominantly occurred at the second position

downstream of the ePUF10 binding site (Supplementary Fig. 1g). We also applied CU-

REWIRE4.0 to specifically targeting the C_{832} site on the report mRNA of the mouse

Pcsk9 gene. Our results revealed a marked improvement in C-to-U editing efficiency at

C_{832} with CU-REWIRE4.0 compared to CU-REWIRE3.0 (Supplementary Fig. 1h).

This underscores the enhanced specificity and efficiency of CU-REWIRE4.0.

Structure-Based Optimization of APOBEC3A Minimizes Off-Target Effects

APOBEC3A proteins tend to form dimers that stabilize their interaction with single-

stranded DNA or RNA [38]. While this interaction is key in inducing mutations in viral

genomes, it also contributes to bystander editing and off-target effects in non-target

genomic regions during base editing. Therefore, our next objective was to reduce

APOBEC3A dimerization.

By investigating the crystalline structure of APOBEC3A and building upon previous

research, we identified two amino acid sites (H11, C171) critical for dimerization [38].

The mutation of these sites could reduce the dimer formation and the association with

nucleic acids of APOBEC3A (Fig. 1c). Additionally, two amino acids (K30, H56) were

found to be involved in substrate RNA recognition by APOBEC3A [38]. Thus we

speculate that mutations at these sites may diminish RNA recognition and, consequently,

reduce off-target effects (Fig. 1d).

To test this possibility, we introduced various point mutations into human

APOBEC3A (Fig. 1e). We then fused these mutated APOBEC3A versions with the

ePUF10 domain, creating multiple CU-REWIRE4 variants that target C_{459} of EGFP

mRNA expressed in HEK293T cells (Fig. 1f). After co-transfecting these constructs

with the EGFP expressing plasmid, we evaluated the C-to-U editing efficiency at the

C_{459} site of EGFP mRNA by these CU-REWIRE4 variants (Fig. 1g).

One variant, CU-REWIRE 4.1, with a C171A substitution in the C-terminal domain

(CTD) of APOBEC3A, showed high editing efficiency while significantly reducing off-

target effects (Fig. 1g, h). Interestingly, the CU-REWIRE 4.2, 4.4, and 4.5, each

containing point mutations in the NTD, showed moderate editing efficiency but

substantially minimized off-target events (Fig. 1g, h). These findings indicate that

strategically designed point mutations in the NTD and CTD of APOBEC3A can

significantly decrease off-target effects while maintaining sufficient editing efficiency

(Fig. 1h). We further examined the on-target editing rates and off-target editing events

of each CU-REWIRE variant, and found that these newly engineered variants provide

optimal ratios of on- *vs.* off-target editing rates, which may offer versatile options for

base editing applications (Fig.1i).

Like the original CU-REWIREs, the modified CU-REWIRE4 variants

predominantly edited cytidines within the UC motif (Supplementary Fig. 2a), reflecting

the intrinsic target preference of APOBEC3A [13,27]. To broaden the application scope of

RNA base editing, we explored whether the APOBEC-derived base editors can achieve

C-to-U editing in other context (i.e., CC, GC, and AC motifs) by systematically

screening different AID/APOBEC enzymes that are integrated in the CU-REWIRE

platform (Supplementary Fig. 2b). We found that only the APOBEC3A-ePUF10

configuration showed detectable base editing near the ePUF10 binding site

(Supplementary Fig. 2c), and therefore select APOBEC3A as the chassis to introduce

additional modifications that may expand the editing capabilities of new generation

CU-REWIREs.

AI-Assisted Engineering of Professional APOBECs (ProAPOBECs)

Our analysis of AID/APOBEC proteins revealed that while the deaminase domain is

relatively conserved, the NTD and CTD are prone to evolutionary changes [39-41]. This

implies that the deaminase domain is crucial for catalyzing the C-to-U transition, while

the NTD and CTD may specialize in recognizing different DNA/RNA targets [39].

Consistent with this notion, previous studies suggested that various APOBEC proteins

might exhibit C-to-U activity outside of the UC context [13], leading us to hypothesize

that the core deaminase domain of APOBEC proteins may potentially edit cytosine in

diverse contexts when combined with distinct NTD or CTD domains.

To test this hypothesis, we utilized AlphaFold2 to analyze the structures of

APOBEC3 and APOBEC1 from both human and rodent origins (Fig. 2a), similar to a

previously used approach that analyzes deaminases across different species [43]. These

two enzymes are the only cytosine deaminases that are currently effective as active

domains in DNA and RNA base editors [13]. We analyzed the phylogenetic tree of

AID/APOBEC family using 1160 full length eukaryotic AID/APOBEC-related genes

documented in the NCBI database, and found that the APOBEC3 and APOBEC1

belong to the closely related clade of this tree (Fig. 2b). There is striking similarity in

the core deaminase domains of APOBEC3s and APOBEC1s, suggesting that functional

modules may be interchangeable among these proteins. Therefore we combined the

core deaminase domain of different APOBECs with the NTD and CTD of human

APOBEC3A. These hybrid deaminases were termed ProAPOBECs, potentially

applicable for both DNA and RNA base editing (Fig. 2c). Together, over thousands of

functional ProAPOBEC proteins could be predicted (Fig. 2d).

We then applied AlphaFold2 to predict structures of ProAPOBECs derived from human and rodent deaminase domains (Fig. 2e and Supplementary Fig. 2d). These ProAPOBECs showed structural conservation with human APOBEC3A, suggesting their potential functionality for base editing. We subsequently fused various ProAPOBECs-C171A with the ePUF10 domain, creating a new generation of CU-REWIREs (version 5.1 to 5.17) (Fig. 2f,g and Supplementary Table 1). Upon expression analysis, most of the CU-REWIRE5 variants (CU5s) demonstrated stable expression and various editing activity towards C_{459} site of EGFP mRNA expressed in the HEK293T cells (Fig. 2h,i). This result demonstrated that we have significantly expanded the AID/APOBEC protein repertoire, leading to the development of novel RNA base editors with a broader targeting context for C-to-U editing. Next, we aim to characterize the editing properties and specificities of these newly engineered CU5s in additional target sites.

Enhanced Targeting Range and Specificity in RNA Base Editing with CU5s Toolkit

We evaluated the RNA editing activity of CU5s by testing their C-to-U editing capabilities at the artificially modified sites of EGFP mRNA, where the $U_{458}\underline{C}_{459}$ was altered to G$\underline{C}$, C$\underline{C}$, and A$\underline{C}$ to test the sequence preference of CU5s (Fig. 3a). Surprisingly, multiple CU5s base editor containing ProAPOBECs exhibited C-to-U transition under G$\underline{C}$, C$\underline{C}$, and A$\underline{C}$ contexts (Fig. 3b). We found that while some new enzymes (like CU5.7 and 5.13) maintain the original preference of U$\underline{C}$ context for their editing activity (Fig. 3b and Supplementary Fig. 3a,b). The CU5.16, which demonstrates a particular interest in the novel C-to-U editors with A$\underline{C}$ context

(Supplementary Fig. 3c-f).

Importantly, new CU5s have high sequence specificity distinct from the original U<u>C</u>

specificity [21,27]. For example, the CU5.1 and 5.8, containing the deaminase domain of

human APOBEC1 and APOBEC3H respectively, demonstrated significant C-to-U

editing activity within G<u>C</u> or A<u>C</u> motifs (Fig. 3b and Supplementary Fig. 4a,b). Our

finding underscores that Pro.hAPOBEC1 editing capabilities significantly differ from

natural human APOBEC 3A (only editing U<u>C</u> motifs) and human APOBEC1 (editing

U<u>C</u> motifs with low efficiency), indicating the emergence of novel cytidine editing

activities in engineered ProAPOBECs. In addition, the CU5.15, containing the

deaminase domain of mouse APOBEC3, demonstrated significant C-to-U editing

activity within C<u>C</u> or U<u>C</u> motifs (Fig. 3b and Supplementary Fig. 4c). The other

enzymes can edit cytosine at various sequence context, including two enzymes (CU5.3

and 5.17, containing the deaminase domain of mouse APOBEC1 and rat APOBEC1

respectively.) that can edit the cytosine in any context (i.e., N<u>C</u> motifs), suggesting they

may function as a base editor with broad target range (Fig. 3b and Supplementary Fig.

4d,e). Such expansion of specificity is particularly useful in their application as a

therapeutic protein (Fig. 3b, Supplementary Fig. 3 and 4).

We further used RNA-seq to assess the bystander editing effects of these new CU5s,

with particular interest in the novel C-to-U editors with different sequence specificity

(like G<u>C</u> editors CU5.1 and 5.8 (Fig. 3c, top), the C<u>C</u> editor CU5.15 (Fig. 3c, middle),

and the N<u>C</u> editors CU5.3 and 5.17 (Fig. 3c, below).

The RNA-seq experiments also allow us to assess the global off-target effects of

CU5s across the entire transcriptome (Fig. 3d). Our results demonstrated that the CU5.1,

5.17, and 5.16 showed higher off-targeting editing rates compared to CU4.1, while

certain CU5s have lower number of off-target editing sites (Fig. 3d). There seems to be

correlation between off-target editing rates and the on-target editing efficiency (i.e.,

more activity CU5s have a larger number of off-target edit sites), suggesting that we

have to balance these two properties of CU5s in the practical applications. We further

analyzed the off-target editing sites for their consensus sequence motifs (Fig. 3e), and

found that these motifs are largely consistent with the sequence preference observed

using C-to-U editing on a single target site (C_{459}) of EGFP mRNA (Fig. 3b and

Supplementary Fig. 4).

Editing Windows and *in vitro* Application of CU5s

To elucidate the editing characteristics of CU5s on the target RNA, we conducted a

comprehensive evaluation of the editing windows associated with selected CU5s. Three

variants, CU5.1, CU5.15 and CU5.17, were selected for this analysis, because they can

edit cytosine in the new sequence context other that the original U<u>C</u> motifs. We found

that CU5.1 (editing at G<u>C</u> motif) demonstrated the highest editing efficiency at

positions 2- or 4-nucleotides (nt) downstream of the ePUF10 binding site in EGFP

mRNA, as shown by different target reporters containing G<u>C</u> dinucleotides at different

positions downstream of the ePUF10 binding site (Fig. 4a).

We further evaluated the efficacy of CU5.1 in reporter genes carrying disease-

causing mutations. For instance, the T143C mutation in the *SOD1* allele, which results

in the p.V48A substitution in the SOD1 protein, is the most common mutation associated with ALS in Southeastern China [44]. We designed two CU5.1 enzymes with ePUF10 domains targeting sequences at positions U_{132}–U_{141} and A_{131}–A_{140} (i.e., 2 or 3 nucleotides upstream of C_{143}, respectively). The CU5.1 enzyme binding 2 nucleotides upstream of the mutated site successfully converted C_{143} to U_{143} with an editing efficiency of ~85%, effectively transforming the mutant *SOD1* allele into the benign form [45] (Fig. 4b).

In another example, the T526C mutation in the APOE4 allele, causing p.R176C in APOE4 protein, was found to be associated with Alzheimer's disease [46,47]. We designed CU5.1 binding to the APOE4 sequences, and found that the CU5.1 binding at 2-nt upstream of the mutated site effectively converted C_{526} to U_{526} with editing efficiency at ~90%, potentially transforming APOE4 into the benign APOE1 allele [48] (Fig. 4c). In addition, the c.68C>A (p.P23H) mutation in the human Rhodopsin gene, known to cause Retinitis Pigmentosa (RP) [49] can be effectively edited by CU5.1 that bind to 2-nt upstream of the mutated site (Figure 4c). As a result of editing C_{76} to U_{76}, we introduced a stop codon in the mutated mRNA, which should abolish the toxic protein mutant (Fig. 4d). Combining the results across different target mRNAs, we summarized the editing window for CU5.1 as 2~5-nt downstream of the ePUF10 binding site (Fig. 4e).

Furthermore, CU5.15 demonstrated the highest editing efficiency 2-nt downstream of the ePUF10 binding site in EGFP mRNA (Fig. 4f). We adopted similar approach

to examine the editing window of CU5.15 using reporters containing C<u>C</u> at different distance from the designed binding site (Fig. 4f), or using multiple CU5.15 targeting the c.104T>C (p.L35P) mutation in MEF2C gene (Fig. 4g) [8]. We found that CU5.15 effectively edited cytidine 2-nt or 3-nt downstream of the ePUF10 binding site, resulting in an editing window within 2~4-nt in various target mRNAs (Fig. 4h).

Finally, CU5.17, with editing activity in "N<u>C</u>" context, demonstrated a high efficiency at positions 2-nt downstream of the ePUF10 binding site as judged by EGFP reporter (Fig. 4i). Importantly, CU5.17 was highly efficient in editing the T1541C site of the human DDX3X mRNA linked to neurodevelopmental disorders [50], particularly when the CU5.17 binding to 2-nt or 3-nt upstream of the mutated site (Fig. 4j). Although bystander editing at C_{1542} occurred, this did not affect the amino acid sequence of DDX3X (Fig. 4j), indicating the editing window for CU5.17 is within 2~5-nt after the ePUF10 binding site across different mRNAs (Fig. 4k).

Improving the CU-REWIRE5 system with AI-Assisted Design of ProAPOBECs utilizing cytosine Deaminase derived from Mammals

We also explored the possibility of using ProAPOBECs derived from deaminase of other mammalian species using similar approach (Supplementary Fig. 5 and Supplementary Table 1). AlphaFold2-based predictions revealed that the deaminase domains from many mammalian APOBEC1s and APOBEC3s may form functional structures with the NTD and CTD of the human APOBEC3A (Supplementary Fig. 5a).

We found that both Z1 and Z2 domains of mammalian APOBEC3s could form

functional ProAPOBECs when combined with NTD and CTD of human APOBEC3A

(Supplementary Fig. 5b,c).

Furthermore, we integrated these newly engineered ProAPOBECs into CU-

REWIRE (i.e., by fusing with ePUF10 domain) and tested their editing activity on

EGFP mRNA. We found significant C-to-U editing activity in GC, AC, and CC contexts

by different ProAPOBECs (Supplementary Fig. 5d,e). ProAPOBECs with deaminase

domains from APOBEC1 showed strong activity in GC contexts, while some with

APOBEC3-Z1 domains demonstrated activity in CC and AC contexts (Supplementary

Fig. 5f-h). These results highlight the versatility of engineered ProAPOBECs for C-to-

U base editing, suggesting an expansion of specificity is particularly useful in potential

therapeutic applications.

***In Vivo* RNA Editing of PCSK9 in Mice reduces cholesterol levels**

To demonstrate *in vivo* applications, we applied CU-REWIREs to edit *Pcsk9*, a

therapeutic target gene, by introducing a premature stop codon to repress PCSK9

protein expression [51]. Inhibiting PCSK9 protein reduces serum low-density lipoprotein

cholesterol (LDL-C) levels, which is associated with a lower risk of cardiovascular

disease [52]. Our aim was to achieve precise RNA editing of *PCSK9* without alternations

of genome DNA, addressing limitations commonly associated with DNA editing

strategies [53-55].

First, we assessed the editing efficiency of various CU-REWIREs and the Cas13-

based C-to-U editors (CURE [23], xCBE [22]) in generating premature stop codons at the

$C_{832}AG$ position within a mouse *Pcsk9* reporter in HEK293T cells (Fig. 5a). While no

significant editing was observed with CURE or xCBE, CU4.1 (CU-REWIRE4.1) and

CU5-A3Z1-1 (CU-REWIRE5 featuring ProAPOBEC from *Ailuropoda melanoleuca*)

effectively converted $C_{832}AG$ to $U_{832}AG$ when ePUF10 domains bound to sequences

spanning U_{821}–G_{830} (Fig. 5a). Notably, CU5-A3Z1-1 demonstrated a remarkable 1.6-

fold increase in editing efficiency compared to CU4.1, achieving 96% versus 37%,

respectively (Fig. 5a). This highlights the superior application potential of ProAPOBEC

in specific contexts compared to native APOBEC proteins.

Next, we packaged CU5-A3Z1-1 into an AAV8 delivery system (Fig. 5b), leveraging

its high liver tropism for in vivo base editing. Mice injected with CU5-A3Z1-1 were

analyzed for changes in *PCSK9* mRNA, protein levels, and total cholesterol four weeks

post-administration. By introducing a premature stop codon, the edited *PCSK9* mRNA

was subjected to degradation via nonsense-mediated mRNA decay [56]. Quantitative PCR

revealed that CU5-A3Z1-1 (2×10^{12} viral genomes (vg)) reduced *PCSK9* mRNA levels

by over 70% compared to the EGFP control group (Fig. 5c). Consistently, this led to

significant reductions in both PCSK9 protein levels (Fig. 5d) and cholesterol levels (Fig.

5e), without affecting serum glyceride or total protein levels (Fig. 5f,g).

As expected, the mice showed no abnormalities in growth following CU5-A3Z1-1

administration (Fig. 5h). Transcriptome-wide RNA sequencing identified

approximately 400 C-to-U editing events associated with CU5-A3Z1-1 treatment (Fig.

5i). These findings demonstrate that CU5-A3Z1-1 enables efficient and precise in vivo

editing of *PCSK9*, offering therapeutic benefits with minimal transcriptome-wide off-

target effects.

***In Vivo* RNA Editing in an ASD Mouse Model Using CU5.15**

Effective RNA base editing in the brain has been a challenging endeavor [20]. To explore

the potential of CU5s, powered by ProAPOBECs, for *in vivo* RNA editing, we utilized

an ASD mouse model with L35P (c.104T>C) point mutation in MEF2C gene, which

was reported to cause severe ASD in human [8]. An APOBEC-embedded cytosine base

editor for DNA has successfully performed C-to-T conversion at the C_{104} site of the

Mef2c gene in the Mef2c L35P mutant mice, effectively reversing the ASD-like

phenotypes in this mouse model [8]. However, the previous methods need to co-inject

two AAV vectors for in vivo DNA base editing, posing practical challenges for its

application in clinical settings [6,8,57]. Packaging the base editing system into a single

AAV vector could significantly reduce costs and enhance delivery efficiency in clinical

applications.

We first examined the C-to-U base editing efficiency of the CU-REWIRE system as

compared with Cas13-based RNA editing tools on the C_{104} site on the Mouse Mef2c

(c.104) mRNA reporter [23,27] (Supplementary Fig. 6). We observed that CU5s exhibited

significantly higher editing efficiency compared to CRISPR/Cas-based tools.

Specifically, CU5.15 achieved an editing efficiency of 43%, and CU1.15 reached 18%

(Supplementary Fig. 6a-b), whereas the CRISPR/Cas-based tool CURE-X-

Pro.mAPOBEC3-1 showed only 5% efficiency (Supplementary Fig. 6c-d). Notably, the

CURE-X system utilizing native APOBEC proteins failed to edit the target *Mef2c*

mRNA. However, replacing the native APOBEC with ProAPOBEC in the CURE-X-

Pro.mAPOBEC3-1 system resulted in effective editing (Supplementary Fig. 6c-e),

underscoring the therapeutic potential of ProAPOBECs in Cas13-based RNA base

editors.

We further packaged CU5.15, featuring the ePUF10 module targeting the A_{92}-A_{101}

region of mouse Mef2c mRNA, into brain blood barrier (BBB)-crossing AAV-PHP.eB

vectors (Fig. 6a). These vectors, along with an EGFP reporter driven by the human

synapsin 1 gene promoter (hSyn), were injected into the tail vein of 4-week-old Mef2c

WT and $L35P^{+/-}$ mice [8]. Robust EGFP expression was observed in the cortical and

hippocampal regions 4- and 16- weeks post-injection (Fig. 6b, c), confirming the

efficient gene delivery in brains.

Sanger sequencing and RNA-sequencing of prefrontal cortical and hippocampus

tissue samples from the mice revealed that CU5.15 induced C-to-U editing at the C_{104}

site in the Mef2c $L35P^{+/-}$ mice injected with CU5.15 (Supplementary Fig. 7). RNA-seq

experiments showed a significant increase in the percentage of U_{104} in the prefrontal

cortex (61.3%) and hippocampus (68%) of Mef2c L35P$^{+/-}$ mice injected with CU5.15,

compared to those injected with ePUF10 (Fig. 6d,e). This data confirmed the induction

of C-to-U editing events by CU5.15. Additionally, no bystander editing events were

observed in the C nucleotides from positions C_{65}-C_{150} of Mef2c mRNA, demonstrating

the precision of CU5.15-mediated RNA editing (Figure 6d). C-to-U editing rates across

various brain regions ranged from 30-50% (Fig. 6e). Considering that the brain tissues

collected for RNA-seq are mixture of neurons and glial cells, the exact C-to-U editing

of the Mef2c L35P site in neurons would be higher. Transcriptome-wide RNA

sequencing revealed approximately 800 C-to-U editing events associated with CU5.15

(Fig. 6f).

Notably, MEF2C protein levels were fully restored in the cortical and hippocampal

regions of Mef2c L35P$^{+/-}$ mice injected with CU5.15, as compared to those injected

with ePUF10 and to Mef2c WT mice (Fig. 6g). This restoration was further validated

by immunostaining using an anti-MEF2C antibody (Fig. 7a,b). Remarkably, the

lifespans of Mef2c L35P$^{+/-}$ mice injected with CU5.15 were close to WT group (Fig.

7c), which was not observed in previous DNA editing work [8], suggesting that RNA base

editing facilitated by CU5.15 may exert a broader impact in this mouse model.

We subsequently examined the influence of CU5.15 on autistic-like behaviors in

Mef2c L35P mice using a series of social interaction tests. The investigation began with

the three-chamber test, comprising both sociability and social novelty tests. In the

sociability test, mice were given the choice between interacting with another mouse or

an empty cage (Fig. 7d). This was followed by the social novelty test, where the mice

had to choose between a familiar partner and a novel partner. Mice treated with AAV-

CU5.15 demonstrated a significant improvement in social interaction performance

during the social novelty test [8] (Fig. 7e,f).

The study progressed to analyzing social interaction behaviors using the social

intruder test, consisting of six trials (T0-T5) (Fig. 7g). After three days of social

isolation (T0), the mice encountered the same partner for four consecutive trials (T1-

T4, each lasting 5 minutes), and then a new partner in the fifth trial (T5). The duration

of sniffing time between the subject mice and their partners was recorded. Notably, the

aberrant social behaviors observed in Mef2c L35P mice during the initial four trials

(T1-T4) with intruder partners were entirely mitigated following treatment with

CU5.15 (Fig. 7g-j). These findings suggest that in vivo RNA base editing facilitated by

CU5.15 effectively restores normal MEF2C protein expression and alleviates the social

impairments associated with the Mef2c L35P mutation.

In summary, our study demonstrates that RNA base editing, as facilitated by CU5.15,

provides a potent tool for correcting genetic mutations in the brain and has potential

therapeutic applications for ASD and other genetic disorders.

Discussion

APOBEC3A is a highly active deaminase frequently used in DNA base editors. However, its application has been hindered by some intrinsic limitations of this enzyme, including off-target effects and RNA binding issues [11,12]. Although point mutations in APOBEC3A have reduced off-target effects in DNA base editing, its application in RNA base editing has remained largely unexplored [58,59]. We hypothesized that the high off-targeting rates of APOBEC3A might be largely due to its close interaction with target mRNA. Structurally, the dimerization of APOBEC3A plays a key role in stabilizing its mRNA binding. Therefore, our initial strategy was to modulate dimerization to minimize its off-target effects and decrease the strong associations with mRNA. Indeed, APOBEC3A mutants in the NTD or CTD, exhibiting reduced mRNA interaction and dimerization, showed fewer off-target effects. This provided a range of APOBEC3A variants for precise RNA base editing in conjunction with the REWIRE system (Fig. 1c-i). However, such variants were limited to editing cytidines in UC contexts (Supplementary Fig. 2a), indicating the need for a more fundamental protein engineering of cytosine base editors with broader targeting scopes.

Utilizing AlphaFold2-assisted protein structure analysis, we discovered that the deaminase domains of APOBECs could act as interchangeable modules (Fig. 2). This suggested that the NTD and CTD may contribute to the recognition of viral genetic materials in various organisms. Consequently, we anticipated that novel deaminase activities could emerge from new combinations of deaminases with NTDs and CTDs.

This led to the identification of a series of APOBECs with novel cytidine-editing

activity under various contexts, significantly broadening the application of REWIRE-

mediated RNA base editing.

Compared to Cas13-based C-to-U base editors, the CU5s with installed

ProAPOBECs demonstrated an impressive capacity for effecting C-to-U transitions.

Notably, our study revealed that the CU 5.7 and 5.13 are capable of editing cytosine in

UC motifs, akin to the CasR13-based RNA editing system that utilizes human

APOBEC3A or evolved ADAR2 enzyme [21,23], the CU5.15 enzyme can edit cytosine

across CC sequence context, similar to the SNAP-CDAR-S system that employs a

evolved ADAR2 enzyme [24]. Meanwhile, the CU5.3 and 5.17 enzyme can edit cytosine

across NC sequence context, suggesting their potential as broad-spectrum editing tools.

Importantly, multiple CU5s base editor have shown remarkable C-to-U transitions

capabilities within GC, AC, and CC motifs, exhibiting high sequence specificity distinct

from traditional RNA base editors (Fig. 3). For example, CU5.1 has shown significant

C-to-U editing activity within GC motifs, albeit with lower efficiency in AC motifs.

This enzyme notably reduces bystander editing effects and enhances editing specificity

in the target region. Such expansion of specificity is particularly useful in their

application as a therapeutic protein (Fig. 3, Supplementary Fig. 3 and 4).

While Cas13-mediated A-to-I RNA editing has been reported in mouse models of

human genetic disorders [18,20], effective C- to-U RNA editing had not been achieved due

to the intrinsic properties of APOBECs. With the new REWIRE system, we demonstrated that ProAPOBEC can effectively perform RNA base editing in the liver and brain (Fig. 5,6). With an editing rate of nearly 50% *in vivo*, Mef2c mutant mice showed restored protein levels and correction of autistic-like behaviors. Furthermore, effective RNA base editing prolonged the lifespan of Mef2c mutant mice, a result not observed with the Mef2c L35P DNA editing [8]. The effects of *in vivo* RNA base editing in our study appear to be long-lasting for over 6 months (Fig. 7c).

ProAPOBECs also have potential applications in the field of RNA epigenetics. The elucidation of epigenetic modifications in DNA and RNA has become crucial for understanding their diverse biological functions [60]. RNA modifications add a new layer to gene regulation, giving rise to "RNA epigenetics." Techniques such as DART-seq (deamination adjacent to RNA modification targets) [61], ACE-Seq (APOBEC-coupled epigenetic sequencing) [62], and AMD-Seq (APOBEC3A-mediated deamination sequencing) [63], all employing APOBECs manipulation, have been developed for comprehensive mapping of RNA epigenetics. Additionally, RNA-binding proteins (RBPs) are pivotal in gene expression and RNA processing. To elucidate RBP dynamics at the single-cell level, the Gene lab developed STAMP (Surveying Targets by APOBEC-Mediated Profiling), enhancing the detection of RBP-RNA interactions [64].

The ProAPOBECs engineered in this study have enhanced editing efficiency and specificity over their natural APOBEC counterparts, emerging as significant

improvement to the current repertoire of tools available for genetic modification. Our objective is to refine the ProAPOBECs' activity and expand their application, facilitating the development of streamlined and accurate methods for conducting both DNA and RNA edits in vivo. In summary, the integration of REWIRE technology with ProAPOBEC-mediated base editing heralds a promising new era in both fundamental research and applied therapeutic strategies. This work not only enriches the existing collection of tools for genetic engineering but also paves the way for novel approaches to treat and manage genetic disorders, marking a significant step forward in the field of genomics and molecular medicine.

Acknowledgments

We extend our gratitude to Dr. Li Yang of Fudan University for the gift of native human/mouse APOBECs, to Dr. Tian Chi of ShanghaiTech University for the CURE plasmids, and to Dr. Chunlong Xu of Lingang Laboratory for the xCBE plasmids. This work is supported by Fundamental Research Funds for the Central Universities (Z.Q), NSFC grants (82430046 to Z.Q., 32030064 to Z.W., 82201632 to W.H.); Project of Medical Technology Research and Transformation supported by Shanghai Municipal Health Commission (2024ZZ1007) (Z.Q.); the National Key Research and Development Program of China (2021YFA1300503), the Starry Night Science Fund at Shanghai Institute for Advanced Study of Zhejiang University (SN-ZJU-SIAS-009) (Z.W.). W.H. is supported by the Fellowship of China Postdoctoral Science Foundation (2022M723244) and the Shanghai Postdoctoral Excellence Program (2022777).

Author contributions statement

Conceptualization, W.H., Z.Q., Z.W.; Biochemical and cell culture experiments, W.H.;

NGS data analysis, B.Y.; AI-Assisted protein structure predictions, X.F., M.H.; Animal

experiments, W.H., W.L., Y.Y., Y.Z., S.W. and S.S.; Writing, W.H., Z.W. and Z.Q.

Data and code availability

The original RNA sequencing datasets have been deposited in the NCBI Gene

Expression Omnibus (accession code PRJNA1068267). Custom computer code is

available upon request. Accession numbers of RNA-seq data are provided in Table S2.

Competing interests:

Authors declare no competing interests.

Reference

1   Rees, H. A. & Liu, D. R. Base editing: precision chemistry on the genome and transcriptome
of living cells. *Nat Rev Genet* **19**, 770-788 (2018). https://doi.org:10.1038/s41576-018-0059-1

2   Porto, E. M., Komor, A. C., Slaymaker, I. M. & Yeo, G. W. Base editing: advances and
therapeutic opportunities. *Nat Rev Drug Discov* **19**, 839-859 (2020).
https://doi.org:10.1038/s41573-020-0084-6

3   Komor, A. C., Kim, Y. B., Packer, M. S., Zuris, J. A. & Liu, D. R. Programmable editing of a
target base in genomic DNA without double-stranded DNA cleavage. *Nature* **533**, 420-424
(2016). https://doi.org:10.1038/nature17946

4   Nishida, K. *et al.* Targeted nucleotide editing using hybrid prokaryotic and vertebrate adaptive
immune systems. *Science* **353**, aaf8727 (2016). https://doi.org:10.1126/science.aaf8729

5   Gaudelli, N. M. *et al.* Programmable base editing of A*T to G*C in genomic DNA without
DNA cleavage. *Nature* **551**, 464-471 (2017). https://doi.org:10.1038/nature24644

6   Koblan, L. W. *et al.* In vivo base editing rescues Hutchinson-Gilford progeria syndrome in
mice. *Nature* **589**, 608-614 (2021). https://doi.org:10.1038/s41586-020-03086-7

7   Mandana Arbab, Z. M., Kaitlyn M. Kray, Ailing Du, Gregory A. Newby, Anton J. Blatnik,
Aditya Raguram, Michelle F. Richter, Kevin T. Zhao, Jonathan M. Levy, Max W. Shen, W.
David Arnold, Dan Wang, Jun Xie, Guangping Gao, Arthur H. M. Burghes, David R. Liu.
Base editing rescue of spinal muscular atrophy in cells and in mice. *science* **380** (2023).
https://doi.org:doi: 10.1126/science.adg6518.

8   Li, W. K. *et al.* Whole-brain in vivo base editing reverses behavioral changes in Mef2c-
mutant mice. *Nat Neurosci* **27**, 116-128 (2024). https://doi.org:10.1038/s41593-023-01499-x

9   Wang, X. *et al.* Efficient base editing in methylated regions with a human APOBEC3A-Cas9
fusion. *Nat Biotechnol* **36**, 946-949 (2018). https://doi.org:10.1038/nbt.4198

10   Kim, D. *et al.* Genome-wide target specificities of CRISPR RNA-guided programmable
deaminases. *Nat Biotechnol* **35**, 475-480 (2017). https://doi.org:10.1038/nbt.3852

11   Erwei Zuo, Y. S., Wu Wei, Tanglong Yuan, Wenqin Ying, Hao Sun, Liyun Yuan, Lars M.
Steinmetz, Yixue Li, Hui Yang. Cytosine base editor generates substantial off-target single
nucleotide variants in mouse embryos. *science* **364**, 289–292 (2019).
https://doi.org:10.1126/science.aav9973

12   Shuai Jin et al. Cytosine, but not adenine, base editors induce genome-wide off-target
mutations in rice. *Science* **364**, 292–295 (2019). https://doi.org:10.1126/science.aaw7166

13   Pecori, R., Di Giorgio, S., Paulo Lorenzo, J. & Nina Papavasiliou, F. Functions and
consequences of AID/APOBEC-mediated DNA and RNA deamination. *Nat Rev Genet* **23**,
505-518 (2022). https://doi.org:10.1038/s41576-022-00459-8

14   Stafforst, T., Schneider, M. F. An RNA-deaminase conjugate selectively repairs point
mutations. *Angew Chem Int Ed Engl* **51**, 11166-11169 (2012).
https://doi.org:10.1002/anie.201206489

15   David B.T. Cox, J. S. G., Omar O. Abudayyeh, Brian Franklin,Max J. Kellner,Julia Joung,
Feng Zhang. RNA editing with CRISPR-Cas13. *Science* **358**, 1019-1027 (2017).
https://doi.org:DOI: 10.1126/science.aaq0180

16   Vogel, P. *et al.* Efficient and precise editing of endogenous transcripts with SNAP-tagged
ADARs. *Nat Methods* **15**, 535-538 (2018). https://doi.org:10.1038/s41592-018-0017-z

17   Katrekar, D. *et al.* In vivo RNA editing of point mutations via RNA-guided adenosine
deaminases. *Nat Methods* **16**, 239-242 (2019). https://doi.org:10.1038/s41592-019-0323-0

18   Qingquan Xiao, Z. X., Yuanyuan Xue, Chunlong Xu, Lei Han, Yuanhua Liu, Fang Wang,
Runze Zhang, Shuang Han, Xing Wang, Geng-Lin Li, Huawei Li, Hui Yang, Yilai Shu.
Rescue of autosomal dominant hearing loss by in vivo delivery of mini dCas13X-derived
RNA base editor. *Sci Transl Med* **14** (2022). https://doi.org:10.1126/scitranslmed.abn0449

19   Yi, Z. *et al.* Engineered circular ADAR-recruiting RNAs increase the efficiency and fidelity
of RNA editing in vitro and in vivo. *Nat Biotechnol* **40**, 946-955 (2022).
https://doi.org:10.1038/s41587-021-01180-3

20   Sinnamon, J. R. *et al.* Targeted RNA editing in brainstem alleviates respiratory dysfunction in
a mouse model of Rett syndrome. *Proc Natl Acad Sci U S A* **119**, e2206053119 (2022).
https://doi.org:10.1073/pnas.2206053119

21   Omar O Abudayyeh, J. S. G., Brian Franklin, Jeremy Koob, Max J Kellner, Alim Ladha, Julia
Joung, Paul Kirchgatterer, David B T Cox, Feng Zhang. A cytosine deaminase for
programmable single-base RNA editing. *Science* **353**, aaf5573 (2019).
https://doi.org:10.1126/science.aaf5573

22   Xu, C. *et al.* Programmable RNA editing with compact CRISPR-Cas13 systems from
uncultivated microbes. *Nat Methods* **18**, 499-506 (2021). https://doi.org:10.1038/s41592-021-
01124-4

23   Huang, X. *et al.* Programmable C-to-U RNA editing using the human APOBEC3A
deaminase. *EMBO J* **39**, e104741 (2020). https://doi.org:10.15252/embj.2020104741

24   Latifi, N., Mack, A. M., Tellioglu, I., Di Giorgio, S. & Stafforst, T. Precise and efficient C-to-
U RNA base editing with SNAP-CDAR-S. *Nucleic Acids Res* **51**, e84 (2023).
https://doi.org:10.1093/nar/gkad598

25   Stroppel, A. S. *et al.* Harnessing self-labeling enzymes for selective and concurrent A-to-I and
C-to-U RNA base editing. *Nucleic Acids Res* **49**, e95 (2021).
https://doi.org:10.1093/nar/gkab541

26   Salter, J. D., Bennett, R. P. & Smith, H. C. The APOBEC Protein Family: United by
Structure, Divergent in Function. *Trends Biochem Sci* **41**, 578-594 (2016).
https://doi.org:10.1016/j.tibs.2016.05.001

27   Han, W. *et al.* Programmable RNA base editing with a single gRNA-free enzyme. *Nucleic
Acids Res* **50**, 9580–9595 (2022). https://doi.org:10.1093/nar/gkac713

28   Hall, C.-G. C. a. T. M. T. Engineering RNA sequence specificity of Pumilio repeats. *Proc
Natl Acad Sci USA* **103**, 13635–13639 (2006). https://doi.org:10.1073 pnas.0606294103

29   Dong, S. *et al.* Specific and modular binding code for cytosine recognition in Pumilio/FBF
(PUF) RNA-binding domains. *J Biol Chem* **286**, 26732-26742 (2011).
https://doi.org:10.1074/jbc.M111.244889

30   Filipovska, A., Razif, M. F., Nygard, K. K. & Rackham, O. A universal code for RNA
recognition by PUF proteins. *Nat Chem Biol* **7**, 425-427 (2011).
https://doi.org:10.1038/nchembio.577

31   Zhao, Y. Y. *et al.* Expanding RNA binding specificity and affinity of engineered PUF
domains. *Nucleic Acids Res* **46**, 4771-4782 (2018). https://doi.org:10.1093/nar/gky134

32   Jarmoskaite, I. *et al.* A Quantitative and Predictive Model for RNA Binding by Human
Pumilio Proteins. *Mol Cell* **74**, 966-981 e918 (2019).
https://doi.org:10.1016/j.molcel.2019.04.012

33   Wang, Y., Cheong, C. G., Hall, T. M. & Wang, Z. Engineering splicing factors with designed
specificities. *Nat Methods* **6**, 825-830 (2009). https://doi.org:10.1038/nmeth.1379

34   Cooke, A., Prigge, A., Opperman, L. & Wickens, M. Targeted translational regulation using
the PUF protein family scaffold. *Proc Natl Acad Sci U S A* **108**, 15870-15875 (2011).
https://doi.org:10.1073/pnas.1105151108

35   Choudhury, R., Tsai, Y. S., Dominguez, D., Wang, Y. & Wang, Z. Engineering RNA
endonucleases with customized sequence specificities. *Nat Commun* **3**, 1147 (2012).
https://doi.org:10.1038/ncomms2154

36   Shinoda, K., Suda, A., Otonari, K., Futaki, S. & Imanishi, M. Programmable RNA
methylation and demethylation using PUF RNA binding proteins. *Chem Commun (Camb)* **56**,
1365-1368 (2020). https://doi.org:10.1039/c9cc09298f

37   Gough, V. & Gersbach, C. A. Immunity to Cas9 as an Obstacle to Persistent Genome Editing.
Mol Ther **28**, 1389-1391 (2020). https://doi.org:10.1016/j.ymthe.2020.05.007

38   Bohn, M. F. *et al.* The ssDNA Mutator APOBEC3A Is Regulated by Cooperative
Dimerization. *Structure* **23**, 903-911 (2015). https://doi.org:10.1016/j.str.2015.03.016

39   Ito, J., Gifford, R. J. & Sato, K. Retroviruses drive the rapid evolution of mammalian
APOBEC3 genes. *Proc Natl Acad Sci USA* **117**, 610-618 (2020).
https://doi.org:10.1073/pnas.1914183116

40   Conticello, S. G., Thomas, C. J., Petersen-Mahrt, S. K. & Neuberger, M. S. Evolution of the
AID/APOBEC family of polynucleotide (deoxy)cytidine deaminases. *Mol Biol Evol* **22**, 367-
377 (2005). https://doi.org:10.1093/molbev/msi026

41   Wenqiang Wang, G.-Z. H. A Long-Running Arms Race between APOBEC1 Genes and
Retroviruses in Tetrapods. *Journal of   Virology* **97** (2023). https://doi.org:10.1128/jvi.01795-
22

42   Lu, G. & Hall, T. M. Alternate modes of cognate RNA recognition by human PUMILIO
proteins. *Structure* **19**, 361-367 (2011). https://doi.org:10.1016/j.str.2010.12.019

43   Huang, J. *et al.* Discovery of deaminase functions by structure-based protein clustering. *Cell*
186, 3182-3195 e3114 (2023). https://doi.org:10.1016/j.cell.2023.05.041

44   Chen, L. X. *et al.* SOD1 Mutation Spectrum and Natural History of ALS Patients in a 15-Year
Cohort in Southeastern China. *Front Genet* **12**, 746060 (2021).
https://doi.org:10.3389/fgene.2021.746060

45   Wang, P. S. *et al.* Clinical characterization and founder effect analysis in Chinese amyotrophic
lateral sclerosis patients with SOD1 common variants. *Ann Med* **56**, 2407522 (2024).
https://doi.org:10.1080/07853890.2024.2407522

46   E. H. Corder, A. M. S., W. J. Strittmatter, D. E. Schmechel, P. C. Gaskell, G. W. Small, A. D.
Roses, J. L. Haines and M. A. Gene Dose of Apolipoprotein E Type 4 Allele and the Risk of
Alzheimer's Disease in Late Onset Families. *Science* **261**, 921-923 (1993).
https://doi.org:10.1126/science.8346443.

47   Blanchard, J. W. *et al.* APOE4 impairs myelination via cholesterol dysregulation in
oligodendrocytes. *Nature* **611**, 769-779 (2022). https://doi.org:10.1038/s41586-022-05439-w

48   Ordovas, J. M., Litwack-Klein, L., Wilson, P. W., Schaefer, M. M. & Schaefer, E. J.
Apolipoprotein E isoform phenotyping methodology and population frequency with
identification of apoE1 and apoE5 isoforms. *Journal of Lipid Research* **28**, 371-380 (1987).
https://doi.org:10.1016/s0022-2275(20)38690-9

49   Thaddeus P. Dryja, T. M., Elias Reichel, Lauri B. Hahn, GlennS. Cowley, David W. Yandell,
Michael A. Sandberg & Eliot L. Berson*. A point mutation of the rhodopsin gene in one form
of retinitis pigmentosa. *Nature* **343**, 364–366 (1990). https://doi.org:10.1038/343364a0

50   Lennox, A. L. *et al.* Pathogenic DDX3X Mutations Impair RNA Metabolism and
Neurogenesis during Fetal Cortical Development. *Neuron* **106**, 404-420 e408 (2020).
https://doi.org:10.1016/j.neuron.2020.01.042

51   Wang, L. *et al.* Eliminating base-editor-induced genome-wide and transcriptome-wide off-
target mutations. *Nat Cell Biol* **23**, 552-563 (2021). https://doi.org:10.1038/s41556-021-
00671-4

52   Horton, J. D., Cohen, J. C. & Hobbs, H. H. Molecular biology of PCSK9: its role in LDL
metabolism. *Trends Biochem Sci* **32**, 71-77 (2007). https://doi.org:10.1016/j.tibs.2006.12.008

53   Rothgangl, T. *et al.* In vivo adenine base editing of PCSK9 in macaques reduces LDL
cholesterol levels. *Nat Biotechnol* **39**, 949-957 (2021). https://doi.org:10.1038/s41587-021-
00933-4

54   Musunuru, K. *et al.* In vivo CRISPR base editing of PCSK9 durably lowers cholesterol in
primates. *Nature* **593**, 429-434 (2021). https://doi.org:10.1038/s41586-021-03534-y

55   Ran, F. A. *et al.* In vivo genome editing using Staphylococcus aureus Cas9. *Nature* **520**, 186-
191 (2015). https://doi.org:10.1038/nature14299

56   Lykke-Andersen, S. & Jensen, T. H. Nonsense-mediated mRNA decay: an intricate machinery
that shapes transcriptomes. *Nat Rev Mol Cell Biol* **16**, 665-677 (2015).
https://doi.org:10.1038/nrm4063

57 Qi M, M. S., Liu J, Liu X, Wei J, Lu WJ, Zhang S, Chang Y, Zhang Y, Zhong K, Yan Y, Zhu
M, Song Y, Chen Y, Hao G, Wang J, Wang L, Lee AS, Chen X, Wang Y, Lan F. . In Vivo
Base Editing of Scn5a Rescues Type 3 Long QT Syndrome in Mice. *Circulation* **149**, 317-329
(2024). https://doi.org:10.1161/CIRCULATIONAHA.123.065624

58 Zhou, C. *et al.* Off-target RNA mutation induced by DNA base editing and its elimination by
mutagenesis. *Nature* **571**, 275-278 (2019). https://doi.org:10.1038/s41586-019-1314-0

59 Grunewald, J. *et al.* Transcriptome-wide off-target RNA editing induced by CRISPR-guided
DNA base editors. *Nature* **569**, 433-437 (2019). https://doi.org:10.1038/s41586-019-1161-z

60 Zhao, L. Y., Song, J., Liu, Y., Song, C. X. & Yi, C. Mapping the epigenetic modifications of
DNA and RNA. *Protein Cell* **11**, 792-808 (2020). https://doi.org:10.1007/s13238-020-00733-
7

61 Meyer, K. D. DART-seq: an antibody-free method for global m(6)A detection. *Nat Methods*
16, 1275-1280 (2019). https://doi.org:10.1038/s41592-019-0570-0

62 Emily K Schutsky, J. E. D., Peng Hu, Monica Yun Liu, Christopher S Nabel, Emily B
Fabyanic, Young Hwang, Frederic D Bushman, Hao Wu & Rahul M Kohli. Nondestructive,
base-resolution sequencing of 5-hydroxymethylcytosine using a DNA deaminase. *Bature
Biotechnology* **36**, 1083–1090 (2018). https://doi.org:10.1038/nbt.42041084

63 Li, Q. Y., Xie, N. B., Xiong, J., Yuan, B. F. & Feng, Y. Q. Single-Nucleotide Resolution
Analysis of 5-Hydroxymethylcytosine in DNA by Enzyme-Mediated Deamination in
Combination with Sequencing. *Anal Chem* **90**, 14622-14628 (2018).
https://doi.org:10.1021/acs.analchem.8b04833

64 Brannan, K. W. *et al.* Robust single-cell discovery of RNA targets of RNA-binding proteins
and ribosomes. *Nat Methods* **18**, 507-519 (2021). https://doi.org:10.1038/s41592-021-01128-0

Figure legends

Figure 1

758 **Fig. 1| Structure-based optimization of APOBEC3A enhances editing efficacy and**

reduces off-target effects of REWIRE-mediated RNA base editing.

a, Schematic design of REWIREs with PUF variants. Top: Original CU-REWIRE 3.0 with PUF10, targeting 10-nt sequences. Bottom: Upgraded CU-REWIRE 4.0 with ePUF10.

b, Expression levels of CU-REWIREs in cultured cells. Representative Western blot images using anti-Flag antibody, related to Supplementary Fig. 1**b**.

c, Crystal structure of Human APOBEC3A (PDB: 4XXO) highlighting putative dimerization interactions. Amino acids involved in dimer formation are shown in spherical forms.

d, Crystal structure of Human APOBEC3A (PDB: 5KEG) in complex with single-stranded DNA. Amino acids responsible for RNA recognition are labeled in spherical forms. Zn symbolizes the catalytic core.

e, Schematic illustration of APOBEC3A with its functional domains.

f, Schematic showing interaction between target EGFP mRNA and CU-REWIRE4.0 and 4.X with modified APOBEC3A variants.

g, Base editing efficiency of EGFP mRNA by CU-REWIRE4.X. The PUF binding site in EGFP is underlined in blue, with adjacent cytosines marked in red (top). Heatmap displays editing rates of cytosines near the on-target site C_{459} for each CU-REWIRE and control (ePUF10 alone), with C_{459} editing rate detailed on the right. Editing rates were measured by RNA-seq with triplicates (refer to methods).

h, Global editing rates of transcriptome-wide RNA base editing in samples treated with different CU-REWIRE4.X (related to data from panel **g**, labels and cutoff as in Supplementary Fig. 1**e**). Orange rhombuses indicate the on-target site EGFP-C_{459}, and the mean number of off-target editing events is noted beside the dot plots. n represents the total number of edited cytosines detected.

i, Correlation between on-target editing rates (Y-axis) and transcriptome-wide C-to-U off-target editing events (X-axis) for various CU-REWIRE4s.

Fig. 2 | AI-Assisted design of programmable APOBEC platform for engineering ProAPOBEC enzymes.

a, Crystal structure of Human APOBEC3A (PDB: 4XXO), AlphaFold2-predicted structures of native APOBEC proteins: Mouse APOBEC3, Human APOBEC1, Mouse APOBEC1, and Rat APOBEC1, showcasing the NTD, C-to-U deaminase domain, and CTD.

b, Phylogenetic tree of the AID/APOBEC family, with major clades highlighted in different colors.

c, Schematic representation of the assembly of ProAPOBECs.

d, Estimated number of potential ProAPOBEC candidates based on AI-predicted AID/APOBEC deaminase domains.

e, AlphaFold2-predicted structures of AID/APOBEC proteins. Top: AI-predicted structures of native AID/APOBECs. Bottom: AI-predicted structures of ProAPOBECs.

f, The programmable CU-REWIRE5 system with ProAPOBEC variants. Top: Upgraded CU-REWIRE5, featuring integration of ProAPOBEC variants for targeted C-to-U editing. Middle: Schematic depiction of the modular assembly process of CU-REWIRE5. The ProAPOBECs domain, constructed from a selection of natural APOBEC deaminase domains (APOBEC$_{DD}$) derived from various sources, is engineered for specific C-to-U base editing. Accompanying this, the ePUF10 domain includes 10 RNA recognition units, each designed for the specific detection of a unique RNA base type. Bottom: Illustration of the precise engagement of CU-REWIRE5 with the target RNA, showcasing the accurately binding and editing C$_{459}$ of EGFP mRNA.

g, Illustrations depicting the components of CU-REWIRE5, ProAPOBECs, and their deaminase donors.

h, Various editing rates on C$_{459}$ of EGFP mRNA of CU-REWIRE5s, assembled with ProAPOBECs in panel (**i**) and Supplementary Fig. 2**d**.

i, Protein expression levels of CU-REWIRE5s in HEK293T cells. The CU-REWIREs were Flag-tagged, anti-Flag antibody was used to examine CU-REWIREs and GAPDH was used as a loading control.

Figure 3

Fig.3 | Enhanced targeting range and specificity in RNA base editing with CU5s.

a, Schematic illustration demonstrating the targeting of the C_{459} site in EGFP mRNA

by CU5s. The native $U_{458}C$ context was manually altered to $A_{458}C$, $G_{458}C$, and $C_{458}C$ to

test editing preferences.

b, Heatmap depicting the percent editing levels of CU5s at the C_{459} site of EGFP

transcript in AC, UC, GC, and CC contexts. The PUF10 binding sequence and on-target

site were showed by panel (**a**).

c, Editing efficacy of CU5s at the on-target and bystander site of EGFP transcript in

AC, UC, GC, and CC contexts. A heatmap displays editing rates of all cytosines near

the on-target site C_{459} for each CU5s and control. Editing rates of C_{459} are measured by

RNA-seq with triplicates.

d, Global editing rates of transcriptome-wide RNA-seq by CU5s (related to data from

panel **b**, with labels and cutoffs the same as in Fig. 1**g**). Orange rhombuses indicate the

on-target EGFP sites, and the average efficiency of total off-target editing events is

noted on the right. n numbers represent edited cytosines detected.

e, Sequence motif logos derived from cytosines edited by different CU5s, based on

RNA-seq data (related to panel **d**).

Figure 4

Fig. 4 | Editing windows and application of CU5s.

a, Determination of editing windows for CU5.1 on C_{459} of EGFP mRNA. The graph

shows on-target editing rates by CU5.1 at various distances from the PUF binding site.

b-d, Editing efficacy of CU5.1 on C_{143} of SOD1 reporter mRNA (**b**), C_{526} of APOE4

mRNA (**c**) and C_{76} of Rhodopsin reporter mRNA (**d**) expressed in HEK293T cells. The

PUF-binding sites are marked in blue.

e, Summary of the editing window for CU5.1 across EGFP, APOE4, and Rhodopsin

reporter mRNAs. The x-axis represents the distance from the editing site to the PUF

binding site; the y-axis shows the C-to-U editing rate. Editing rates were acquired by

Sanger sequencing.

f, Determination of editing windows for CU5.15 on C_{459} of EGFP mRNA.

g, Editing efficacy of CU5.15 on C_{104} of Mef2c reporter mRNA expressed in HEK293T

cells.

h, Summary of the cytosine editing window mediated by CU5.15 around the target sites

of EGFP and Mef2c reporter mRNAs.

i, Determination of editing windows for CU5.17 on C_{459} of EGFP mRNA.

j, Editing efficacy of CU5.17 on C_{1541} of DDX3X reporter mRNA expressed in

HEK293T cells.

k, Summary of the cytosine editing window mediated by CU5.17 around the target sites

of EGFP and DDX3X reporter mRNAs. Values represent mean ± SEM (n=3).

Figure 5

Fig. 5 | *In Vivo* RNA Editing of PCSK9 in Mice reduces cholesterol levels

a, Efficient editing of C_{832} in PCSK9 mRNA by CU-REWIREs and Cas13-based editor. The PUF binding site in PCSK9 is underlined in blue, with on-target site marked in red. Editing rates of C_{832} are measured by Sanger-sequencing with triplicates.

b, Schematic illustration of CU5-A3Z1-1 vectors delivered into mice through intravenous injection of AAV8 virus (top) and the key experimental steps (bottom).

c, In vivo knockdown frequencies induced by CU5-A3Z1-1. Unpaired two-sided Student's t-test was used, statistical values represent the mean ± s.e.m. *P < 0.05. n = 5 mice.

d-g, The levels of serum PCSK9 protein, cholesterol, glyceride and total protein levels in the mice injected with CU5-A3Z1-1. Unpaired two-sided Student's t-test was used, statistical values represent the mean ± s.e.m. *P < 0.05. n = 5 mice.

h, Statistical analysis of weight changes in mice following administration in the CU5-A3Z1-1 and EGFP control. n = 5 mice.

i, Scatter plots showing transcriptome-wide C-to-U RNA base editing in liver samples

treated with CU5-AEZ1-1 and EGFP control in mice. Values represent mean ± s.e.m.

(n = 3).

Fig. 6 | *In vivo* RNA base editing in Mef2c L35P$^{+/-}$ mice corrects genetic mutations

and protein expression.

a, Schematic illustration of AAV-CU-REWIRE5.15 (CU5.15) vectors delivered into

Mef2c L35P$^{+/-}$ mice through intravenous injection of AAV-PHP.eB virus in the tail vein.

b, Immunofluorescence images showing CU5.15 (green) and DAPI (blue) in the

cortical and hippocampal brain regions, 4-weeks post-viral injection. Scale bar: 200μm.

c, Immunofluorescence images of CU5.15 (green) and DAPI (blue) in the cortical and

hippocampal regions, 16-weeks post-viral injection.

d, Editing efficiency of C$_{104}$ in endogenous mouse Mef2c mRNA by CU5.15. The

ePUF10 binding site is underlined in blue, C$_{104}$ marked in red (top). Lower panel:

Editing rates of cytosines around the on-target site. Rates were measured by RNA-seq

with triplicates. Values represent mean ± s.e.m. (n = 3).

e, Percentage of U_{104} in prefrontal cortex and hippocampus tissues of Mef2c WT or

$L35P^{+/-}$ mice edited by ePUF10 or CU5.15, measured by RNA-seq with triplicates

(Left). Editing efficiency of C-to-U of the Mef2c C_{104} site by CU5.15 in various groups

(Right). Editing rate formula: $C_{edited} = (U_{CU5.15} - U_{ePUF10}) / C_{total} (1 - U_{ePUF10})$.

f, Scatter plots showing transcriptome-wide C-to-U RNA base editing in prefrontal

cortex or hippocampus samples treated with CU5.15 and ePUF10 control in mice brain

samples. Values represent mean ± s.e.m. (n = 3).

g, MEF2C protein levels in the prefrontal cortex or hippocampus of Mef2c WT or

$L35P^{+/-}$ mice treated with ePUF10 or CU5.15. anti-MEF2C antibody was used.

GAPDH used as a loading control (Left). Quantification of MEF2C protein level

(Right). ePUF10 vs. CU5.15 in prefrontal cortex, P = 0.0203; ePUF10 vs. CU5.15 in

hippocampus, P = 0.0423), Unpaired two-sided Student's t-test was used, statistical

values represent the mean ± s.e.m. *P < 0.05. n = 3 mice.

Figure 7

Fig. 7 | *In vivo* RNA base editing in Mef2c L35P$^{+/-}$ mice rescued abnormal social behaviors.

a, Immunohistochemical staining for MEF2C (red) and DAPI (blue) in the retro splenial cortex (RSC) and hippocampus of Mef2c WT and L35P$^{+/-}$ mice, indicating viral injections.

b, Left: Quantification of MEF2C protein fluorescence density in the RSC (ePUF10 vs. CU5.15, P = 0.0040). Right: In the hippocampus (ePUF10 vs. CU5.15, P < 0.0001. *P < 0.01. n = 6 brain slices from three mice of each group).

c, Lifespan of Mef2c L35P$^{+/-}$ mice with ePUF10 or CU5.15. Kaplan-Meier curve statistical analysis (L35P$^{+/-}$(ePUF10) versus L35P$^{+/-}$(CU5.15), Chi square=25.99, P < 0.0001). Kaplan-Meier curve statistical analysis, Log-rank (Mantel-Cox) test was used. WT (ePUF10) n = 23 mice, L35P$^{+/-}$ (ePUF10) n = 17 mice, L35P$^{+/-}$ (CU5.15) n = 23

912 mice).

913 **d,** Heatmaps representing locomotion of Mef2c WT or L35P$^{+/-}$ mice in the sociability

914 and social novelty sessions of the three-chamber test.

915 **e,** Cumulative interaction duration with mouse and empty cage in different groups of

916 Mef2c mice (stranger vs. empty, WT+ePUF10, $P < 0.0001$, L35P$^{+/-}$+ePUF10, $P <$

917 0.0001; L35P$^{+/-}$+ CU5.15, $P < 0.0001$. n = 14 mice).

918 **f,** Cumulative interaction duration with stranger and familiar mouse in different groups

919 of Mef2c mice (stranger vs. familiar, WT+ePUF10, $P = 0.0007$; L35P$^{+/-}$+ePUF10, $P =$

920 0.9130; L35P$^{+/-}$+ CU5.15, $P < 0.0001$. Significantly different, $P < 0.01$. n = 14 mice).

921 **g,** Diagram of the home cage experiment. T0: social isolation for three days. T1-T4:

922 Social interaction trials with one same partner mouse. T5: Social interaction with a

923 novel partner mouse. Recording sniffing time within a 2-minute period.

924 **h,** Sniffing time in social intruder test for Mef2c WT +ePUF10 (T1 vs. T2, $P < 0.0001$;

925 T1 vs. T3, $P < 0.0001$; T1 vs. T4, $P < 0.0001$; T4 vs. T5, $P < 0.0001$. n = 12 mice).

926 **i,** Sniffing time in the social intruder test for Mef2c L35P$^{+/-}$ +ePUF10 (T1 vs. T2, $P =$

927 0.0153; T1 vs. T3, $P = 0.0614$; T1 vs. T4, $P = 0.0026$; T4 vs. T5, $P < 0.0001$. n = 12

928 mice).

929 **j,** Sniffing time in the social intruder test for Mef2c L35P$^{+/-}$ +CU5.15 (T1 vs. T2, $P <$

930 0.0001; T1 vs. T3, $P < 0.0001$; T1 vs. T4, $P < 0.0001$; T4 vs. T5, $P < 0.0001$. n = 12

931 mice).

932 Paired two-sided Student's t-test was used, statistical values represent the mean $\pm$ s.e.m.

933 * $P < 0.01$, ** $P < 0.001$, *** $P < 0.0001$.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

a

b

c CU-REWIRE5.15 expression in brain

d Endogenous, Mef2c mRNA

e

f CU-REWIRE5.15

g

Figure 7