Metabolic constraints modulate the likelihood and predictability of global epistasis.

Pavithra Venkataraman[#], Supreet Saini[#]

Department of Chemical Engineering, Indian Institute of Technology Bombay, India 400076

[#]Corresponding Authors: PV (pavithrav@iitb.ac.in), SS (saini@che.iitb.ac.in);
+91 22 2576 4235

Running Title: *A mechanistic explanation of global epistasis*

Abstract

Epistasis, where a mutation's effect depends on its genetic background, is a key driver of evolution. A key pattern, global epistasis, suggests that mutation fitness effects can be predicted based on background fitness. However, no mechanistic explanation is known for this statistical trend.

Here, we present a first-principles model demonstrating that global epistasis emerges naturally from metabolic constraints on cell growth and division. Using a prokaryotic growth model based on an adder-sizer metabolic principle, we show that: (i) global epistasis becomes less strong as genetic variation increases, and (ii) mutations in rate-limiting terminal reaction module of an enzymatic pathway exhibit the most predictable fitness effects. These findings establish a direct link between metabolism, genetic interactions, and evolutionary predictability.

By uncovering the mechanistic basis of global epistasis, our study advances the understanding of evolutionary constraints and has implications for antibiotic resistance, synthetic biology, and microbial adaptation in dynamic environments.

Keywords: Epistasis; Global epistasis; Mutation Effects; Metabolism.

Introduction

Predicting evolution is a fundamental challenge in biology, with implications on biodiversity, genetics, human health, biogeochemical processes, and agriculture[1–6]. A key obstacle is epistasis, where mutation effects depend on genetic background, shaping evolutionary trajectories in complex ways[7,8]. One striking pattern observed in experimental evolution is global epistasis, where the fitness effects of mutations systematically decline as background fitness increases[9–12]. While this trend has been widely documented, there are also exceptions[13,14]. The mechanistic origins of these patterns, however, remain unclear[15].

Most studies of epistasis rely on statistical modeling approaches[16,17], or empirically characterizing individual mutation effects across backgrounds and analyzing fitness effect distributions (DFEs)[12,18–27]. However, the vast number of possible mutations and genotypes makes full experimental characterization impractical[28,29]. Although metabolic networks and protein interactions have been implicated in epistasis[30–38], no first-principles model explains its statistical patterns or why fitness effects follow predictable trajectories despite the complexity of genetic and metabolic interactions.

Here, we present a mechanistic framework demonstrating that global epistasis arises naturally from metabolic constraints on cell growth and division. Using a quantitative prokaryotic metabolism model[39], we show that mutations in rate-limiting terminal enzymatic steps exhibit strong negative global epistasis, while mutations in interconnected steps display less predictable effects. Furthermore, as background genetic variability increases, epistasis' predictability declines, marking a transition from deterministic to idiosyncratic fitness effects.

Our findings have broad implications for microbial adaptation, antibiotic resistance evolution, and biotechnological strain engineering. Additionally, our model generates testable predictions, making it a valuable theoretical tool for guiding future empirical studies. By moving beyond statistical correlations to mechanistic insights, this study provides a foundational framework for understanding how metabolic constraints shape evolutionary dynamics.

Results

Mutational Fitness Effects Depend on Genetic Background and Exhibit Global Epistasis/

We modelled a prokaryotic cell with three rate-limiting metabolic pathways, each containing two enzymatic reactions (**Figure 1A**). Cell division occurs once predefined threshold amounts of all three metabolites (P_1, P_2, and P_3) are accumulated. The time taken to reach this threshold (t_{div}) determines division timing, with fitness defined as its reciprocal ($1/t_{div}$).

We call reach reaction step a 'module', numbered from one to six in the cell. Each module is defined by four parameters. Two of these parameters, t_{ON} and t_{OFF}, define the promoter activity; while the other two, k_{cat} and K_M, define the enzyme properties.

To study epistasis, we introduced a mutation by altering a parameter (t_{ON1}) in module 1, which dictates the production of intermediate I_1 and directly regulates flux through the other two pathways by drawing up the common substrate (indicated by a triangle in **Figure 1A**). The mutation of interest in this study, unless specified otherwise, is t_{ON1} changing from 0.5 to 1. We assess how this mutation's fitness effect varied across distinct genetic backgrounds (See Methods).

Figure 1B illustrates the mutation's fitness effect in four representative cases, where cells differed in one, two, three, or four background parameters. The distributions across all conditions are shown in **Supplementary Figure 1**. Our results demonstrate that mutation effects are not uniform but vary with genetic background, a hallmark of epistasis. This trend persisted even when the number of threshold metabolites was altered (**Supplementary Figure 2**).

A key signature of global epistasis is that the fitness effect of a mutation declines predictably as background fitness increases. To test whether the mutation changing t_{ON1} from 0.5 to 1 in module 1 exhibited this property, we plotted fitness effects of this mutation against background fitness. The strength of this relationship is quantified by R^2 of a linear regression, with higher values indicating stronger global epistasis[40]. Experimental studies have observed epistasis of varying globality in microbial evolution, with R^2 ranging from 0.4 to 0.9[9,10,40–42]. **Figure 1C** presents representative regression analyses, confirming that our model captures global epistasis. A full set of 300 regressions across all background conditions is provided in **Supplementary Figure 3**.

Figure 1. Mutational fitness effects depend on genetic background and vary in predictability. (A) Modeled cells divide upon accumulating threshold amounts of three bottleneck metabolites (P_1, P_2, and P_3), each produced via enzymatic cascades from a common substrate. **(B)** The distribution of fitness effects for a fixed mutation (t_{ON1} changes from 0.5 to 1) is shown across genetic backgrounds differing in one (k_{cat1}), two (k_{cat1}, t_{ON2}), three (k_{cat1}, t_{ON2}, k_{cat3}), or four (k_{cat1}, t_{ON2}, k_{cat3}, t_{OFF6}) parameters. Fitness effects are non-uniform, confirming their dependence on genetic background. **(C)** Regression plots of fitness effects against background fitness for mutations occurring in backgrounds differing by one, two, three, or four parameters. The fitness effect of a mutation is predictable to varying degrees based on background fitness, demonstrating global epistasis.

Stronger Global Epistasis is Less Likely with Increased Background Variation

Global epistasis can be characterized by the slope of the regression between fitness effects and background fitness. A negative slope indicates diminishing-returns and increasing-costs epistasis, while a positive slope reflects increasing-returns and diminishing-costs epistasis[40]. To determine whether global epistasis was predominantly negative or positive, we analyzed the

slopes of regression lines (**Supplementary Figure 3**). As shown in **Figure 2A**, negative global epistasis (NGE) remained the dominant pattern across all levels of background variation.

Despite NGE being common, the skewness of the distribution of R^2 values increased as the extent of background variation increased (*1-site*: -0.5864; *2-site*: -0.365; *3-site*: -0.951; *4-site*: 0.2385), though the mean values remained statistically equal (**Figure 2B**, two-tailed *t*-test, $p > 0.05$; see **Supplementary Figure 4** for full distributions across fifteen background variation types).

We defined strong negative global epistasis as cases where $R^2 > 0.5$, distinguishing them from weaker instances. Using this threshold, we quantified how often strong NGE occurred at different levels of background variation. As shown in **Figure 2A**, the likelihood of strong negative global epistasis decreases as background variation increases.

Figure 2. Strong negative global epistasis (NGE) becomes less frequent as background variation increases. (A) The sign of global epistasis was determined from the slope of the regression between mutation fitness effects and background fitness. Negative global epistasis (NGE; squares) remained dominant across all levels of background variation. However, the frequency of strong NGE ($R^2 > 0.5$; circles) decreased as background variation increased. **(B)** Distribution of R^2 values for NGE cases across different background variation levels. While R^2 values varied, their mean remained statistically unchanged ($p > 0.05$, two-tailed *t*-test). See **Supplementary Figure 5** for full distributions across all 15 background variation types.

Reaction Speeds Predict the Strength of Global Epistasis

We next examined whether the strength of global epistasis can be predicted based on the relative speeds of the enzymatic modules within the cell. To test this, we constructed four types

of simplified ("toy") cells where six enzymatic modules were predefined as either fast or slow (see Methods). As shown in **Figure 3A**, we created the following variants:

1. 1A – One slow first-step module, with all others fast.
2. 1B – One fast first-step module, with all others slow.
3. 2A – One slow second-step module, with all others fast.
4. 2B – One fast second-step module, with all others slow.

Thus, each cell type contains one outlier reaction step compared to the rest. We simulated twenty-five cells of each type and introduced a hundred random mutations in the outlier step to assess how global epistasis varies.

Our results (**Figure 3B**) show that only mutations in slow second-step modules (2A) exhibit the strongest global epistasis, with fitness effects most strongly correlated with background fitness. In contrast, mutations in other reaction types (1A, 1B, and 2B) show weaker global epistasis. These findings hold even when metabolic network architecture is altered, such that the cell has two threshold metabolites each produced by a cascade of three enzymatic reactions (**Supplementary Figure 5**). Additionally, when we introduced strong metabolic feedback— shut down a reaction arm once a threshold amount of metabolite was produced—we found that mutations in the slowest reaction steps consistently exhibit strong global epistasis (**Supplementary Figure 5**).

Recent studies suggest that as background fitness increases, beneficial mutations become rarer while deleterious mutations increase, reinforcing that global epistasis strongly determines the sign of mutational effects[17,42]. Our findings provide a mechanistic basis for this observation.

Using our dataset of a hundred mutations across twenty-five genotypes in each of the four cell types, we analyzed the mean and variance of these distributions of fitness effects (DFEs). While mean DFE values consistently declined with increasing background fitness (**Supplementary Figure 6**), this decline was most predictable in case 2A, where mutations affected a slow second-step module (**Figure 3C**). This result holds when the metabolic architecture is altered, and when feedback is introduced (**Supplementary Figure 7**). On the other hand, variance of the DFE did not follow a clear pattern, showing no consistent correlation with background fitness (**Figure 3C**, **Supplementary Figure 8**).

Figure 3. Predictability of mutational fitness effects depends on reaction speed and metabolic control. (A) We constructed four types of simplified ("toy") cells, each with six enzymatic modules pre-defined as either slow (red arrows) or fast (green arrows). The outlier module in each cell type is indicated by bold arrows: 1A: One slow first-step module, with all others fast; 1B: One fast first-step module, with all others slow; 2A: One slow second-step module, with all others fast; and 2B: One fast second-step module, with all others slow. **(B)** To assess global epistasis, we introduced a hundred mutations in the outlier step of twenty-five genetic backgrounds for each cell type and performed a linear regression of mutation fitness effects against background fitness. R^2 values for each mutation are plotted, with mean trends shown by a connecting line. Mutations in slow second-step modules (2A) exhibit the strongest global epistasis, with fitness effects most predictable using background fitness. **(C)** The strength of global epistasis was further quantified by performing linear regressions of DFE mean and variance against background fitness for each cell type. While the mean fitness effect declines predictably in case 2A, variance does not follow a consistent trend.

Discussion

In this study, we developed a mechanistic cell growth and division model that captures epistasis and its globality feature, revealing how metabolic constraints shape mutational effects and drive declining adaptability. Our model defines fitness based on the time required to accumulate threshold levels of key metabolites before division. In living organisms, metabolic networks are complex and reaction arms are mostly interconnected, except terminal steps that lead to the production of metabolites[43]. While the cell that we model does not encode an intricate network, we have two types of reaction modules – ones that control the flux through the other paths, and ones that do not exert control over other paths. This metabolic structure in the cell allows us to comment on the macroscopic effects of coding and non-coding mutations in both types of reaction modules that are part of real cellular networks.

Empirical studies show that negative global epistasis (NGE)—where beneficial mutation's effects diminish as background fitness increases—is common but not universal[13,14,44]. Our model reproduces this trend and predicts that strong NGE becomes less frequent as genetic variability increases. Additionally, we also show that mutations in the slowest, least-controlling reaction modules exhibit strong NGE. As these slow steps improve, beneficial mutations become rarer, contributing to declining adaptability over time[23,26,45].

Conversely, positive global epistasis (increasing-returns epistasis), which implies synergistic mutational effects[46–49], is predicted to occur in highly interconnected modules, where a single mutation can yield a significant fitness gain.

Beyond its theoretical insights, this framework has broad implications for understanding how metabolic complexity influences epistasis. Future work could extend this model by incorporating cellular regulation layers and environmental changes, to explore the role of these factors in dictating epistasis' predictability[40,50]. Additionally, our findings could help in engineering microbial strains with predictable evolutionary trajectories, impacting biotechnology and antimicrobial resistance research[21,51].

Our model provides a quantitative framework for global epistasis. Importantly, it also makes clear testable predictions. In the metabolic map of the cell, several key metabolites feed branches of enzymatic reactions. For instance, when *E. coli* is grown in anaerobic conditions, acetyl-CoA feeds into the Krebs cycle, fatty acid biosynthesis, and mixed acid fermentation[52–54]. Similarly, glutamate is a substrate for the synthesis of glutamine, proline and arginine[55]. Our

model predicts the reaction modules in these pathways in which mutations, under appropriate

conditions, would exhibit strong global epistasis. Our model also predicts sign-epistasis

between promoter and gene mutations, which is likely to constrain evolutionary trajectories

through mechanisms which are not fully understood yet (**Supplementary Figure 9**)[50,56].

Ultimately, this work provides a quantitative foundation for predicting fitness gains in evolving

populations. Future studies will be crucial in testing and refining these insights, to further our

understanding of the role of cellular processes in dictating evolutionary dynamics.

Funding

This work was funded by a DBT/Wellcome Trust (India Alliance) grant (Award no. IA/S/19/2/504632) to SS. PV is supported by the Prime Minister's Research Fellowship (PMRF ID 1302050).

Acknowledgments

We are grateful to Deepa Agashe, Sunil Laxman, Sandeep Krishna, Christian Landry, Sergey Kryazhimskiy, and Kirti Jain for their critical insights and feedback. Their inputs helped refine arguments and strengthen the manuscript's clarity.

References

1. Wang, L. & Wang, H. Marine Biogeochemical Processes and Ecosystem Evolution: Observational and Predictive Approaches. *ijms* (2024) doi:10.5376/ijms.2024.14.0034.
2. Omenn, G. S. Evolution and public health. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 1702–1709 (2010).
3. Thrall, P. H. *et al.* Evolution in agriculture: the application of evolutionary approaches to the management of biotic interactions in agro-ecosystems. *Evolutionary Applications* **4**, 200–215 (2011).
4. Mackay, T. F. & Moore, J. H. Why epistasis is important for tackling complex human disease genetics. *Genome Med* **6**, 42 (2014).
5. Tang, D., Freudenberg, J. & Dahl, A. Factorizing polygenic epistasis improves prediction and uncovers biological pathways in complex traits. *The American Journal of Human Genetics* **110**, 1875–1887 (2023).
6. Borrell, S. & Gagneux, S. Strain diversity, epistasis and the evolution of drug resistance in Mycobacterium tuberculosis. *Clin Microbiol Infect* **17**, 815–820 (2011).
7. Bateson, W., Mendel, G. & Leighton, A. G. *Mendel's Principles of Heredity, by W. Bateson.* (University Press, Cambridge [Eng.], 1909). doi:10.5962/bhl.title.1057.
8. Østman, B., Hintze, A. & Adami, C. Impact of epistasis and pleiotropy on evolutionary adaptation. *Proceedings of the Royal Society B: Biological Sciences* **279**, 247–256 (2012).
9. Kryazhimskiy, S., Rice, D. P., Jerison, E. R. & Desai, M. M. Microbial evolution. Global epistasis makes adaptation predictable despite sequence-level stochasticity. *Science* **344**, 1519–1522 (2014).
10. Johnson, M. S., Martsul, A., Kryazhimskiy, S. & Desai, M. M. Higher-fitness yeast genotypes are less robust to deleterious mutations. *Science* **366**, 490–493 (2019).
11. Couce, A. & Tenaillon, O. A. The rule of declining adaptability in microbial evolution experiments. *Front. Genet.* **6**, 99 (2015).
12. Wiser, M. J., Ribeck, N. & Lenski, R. E. Long-Term Dynamics of Adaptation in Asexual Populations. *Science* **342**, 1364–1367 (2013).
13. Rojas Echenique, J. I., Kryazhimskiy, S., Nguyen Ba, A. N. & Desai, M. M. Modular epistasis and the compensatory evolution of gene deletion mutants. *PLoS Genet* **15**, e1007958 (2019).
14. Smith, C. E., Smith, A. N. H., Cooper, T. F. & Moore, F. B.-G. Fitness of evolving bacterial populations is contingent on deep and shallow history but only shallow history creates predictable patterns. *Proc Biol Sci* **289**, 20221292 (2022).
15. Johnson, M. S., Reddy, G. & Desai, M. M. Epistasis and evolution: recent advances and an outlook for prediction. *BMC Biol* **21**, 120 (2023).
16. Diaz-Colunga, J. *et al.* Global epistasis on fitness landscapes. *Philosophical Transactions of the Royal Society B* **378**, 20220053 (2023).
17. Reddy, G. & Desai, M. M. Global epistasis emerges from a generic model of a complex trait. *eLife* **10**, e64740 (2021).
18. Bank, C. Epistasis and Adaptation on Fitness Landscapes. *Annu. Rev. Ecol. Evol. Syst.* **53**, 457–479 (2022).
19. Pokusaeva, V. O. *et al.* An experimental assay of the interactions of amino acids from orthologous sequences shaping a complex fitness landscape. *PLoS Genet* **15**, e1008079 (2019).
20. Jayaraman, V., Toledo-Patiño, S., Noda-García, L. & Laurino, P. Mechanisms of protein evolution. *Protein Sci* **31**, e4362 (2022).
21. Weinreich, D. M., Delaney, N. F., DePristo, M. A. & Hartl, D. L. Darwinian Evolution Can Follow Only Very Few Mutational Paths to Fitter Proteins. *Science* **312**, 111–114 (2006).
22. Good, B. H. & Desai, M. M. The Impact of Macroscopic Epistasis on Long-Term Evolutionary Dynamics. *Genetics* **199**, 177–190 (2015).
23. Kassen, R. & Bataillon, T. Distribution of fitness effects among beneficial mutations before selection in experimental populations of bacteria. *Nat Genet* **38**, 484–488 (2006).
24. Levy, S. F. *et al.* Quantitative evolutionary dynamics using high-resolution lineage tracking. *Nature* **519**, 181–186 (2015).

25. Domingo-Calap, P., Cuevas, J. M. & Sanjuán, R. The Fitness Effects of Random Mutations in
Single-Stranded DNA and RNA Bacteriophages. *PLoS Genet* **5**, e1000742 (2009).

26. Aggeli, D., Li, Y. & Sherlock, G. Changes in the distribution of fitness effects and adaptive
mutational spectra following a single first step towards adaptation. *Nat Commun* **12**, 5193 (2021).

27. Sane, M., Diwan, G. D., Bhat, B. A., Wahl, L. M. & Agashe, D. Shifts in mutation spectra
enhance access to beneficial mutations. *Proc. Natl. Acad. Sci. U.S.A.* **120**, e2207355120 (2023).

28. Lässig, M., Mustonen, V. & Walczak, A. M. Predicting evolution. *Nat Ecol Evol* **1**, 0077 (2017).

29. Bank, C., Matuszewski, S., Hietpas, R. T. & Jensen, J. D. On the (un)predictability of a large
intragenic fitness landscape. *Proc. Natl. Acad. Sci. U.S.A.* **113**, 14085–14090 (2016).

30. Segrè, A. V., Murray, A. W. & Leu, J.-Y. High-Resolution Mutation Mapping Reveals Parallel
Experimental Evolution in Yeast. *PLoS Biol* **4**, e256 (2006).

31. Szathmáry, E. Do deleterious mutations act synergistically? Metabolic control theory provides a
partial answer. *Genetics* **133**, 127–132 (1993).

32. Kryazhimskiy, S. Emergence and propagation of epistasis in metabolic networks. *eLife* **10**,
e60200 (2021).

33. Maclean, R. C. Predicting epistasis: an experimental test of metabolic control theory with
bacterial transcription and translation. *Journal of Evolutionary Biology* **23**, 488–493 (2010).

34. Ortlund, E. A., Bridgham, J. T., Redinbo, M. R. & Thornton, J. W. Crystal Structure of an Ancient
Protein: Evolution by Conformational Epistasis. *Science* **317**, 1544–1548 (2007).

35. Sethi, A., Eargle, J., Black, A. A. & Luthey-Schulten, Z. Dynamical networks in tRNA:protein
complexes. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 6620–6625 (2009).

36. Campbell, E. C. *et al.* Laboratory evolution of protein conformational dynamics. *Current Opinion*
in Structural Biology **50**, 49–57 (2018).

37. Jiménez-Osés, G. *et al.* The role of distant mutations and allosteric regulation on LovD active site
dynamics. *Nat Chem Biol* **10**, 431–436 (2014).

38. Otten, R. *et al.* How directed evolution reshapes the energy landscape in an enzyme to boost
catalysis. *Science* **370**, 1442–1446 (2020).

39. Dutta, D. & Saini, S. Cell Growth Model with Stochastic Gene Expression Helps Understand the
Growth Advantage of Metabolic Exchange and Auxotrophy. *mSystems* **6**,
10.1128/msystems.00448-21 (2021).

40. Diaz-Colunga, J., Sanchez, A. & Ogbunugafor, C. B. Environmental modulation of global
epistasis in a drug resistance fitness landscape. *Nat Commun* **14**, 8055 (2023).

41. Sailer, Z. R. & Harms, M. J. High-order epistasis shapes evolutionary trajectories. *PLoS Comput*
Biol **13**, e1005541 (2017).

42. Ardell, S. M., Martsul, A., Johnson, M. S. & Kryazhimskiy, S. Environment-independent
distribution of mutational effects emerges from microscopic epistasis. *Science* **386**, 87–92 (2024).

43. Judge, A. & Dodd, M. S. Metabolism. *Essays Biochem* **64**, 607–647 (2020).

44. Baheti, S., Raj, N. & Saini, S. Variations and predictability of epistasis on an intragenic fitness
landscape. Preprint at https://doi.org/10.7554/eLife.104848.1 (2025).

45. Venkataram, S., Monasky, R., Sikaroodi, S. H., Kryazhimskiy, S. & Kacar, B. Evolutionary
stalling and a limit on the power of natural selection to improve a cellular module. *Proc. Natl.*
Acad. Sci. U.S.A. **117**, 18582–18590 (2020).

46. Chou, H.-H., Berthet, J. & Marx, C. J. Fast growth increases the selective advantage of a
mutation arising recurrently during evolution under metal limitation. *PLoS Genet* **5**, e1000652
(2009).

47. Hsieh, Y.-Y. P., Makrantoni, V., Robertson, D., Marston, A. L. & Murray, A. W. Evolutionary
repair: Changes in multiple functional modules allow meiotic cohesin to support mitosis. *PLoS*
Biol **18**, e3000635 (2020).

48. Blount, Z. D., Borland, C. Z. & Lenski, R. E. Historical contingency and the evolution of a key
innovation in an experimental population of Escherichia coli. *Proc Natl Acad Sci U S A* **105**,
7899–7906 (2008).

49. He, X., Qian, W., Wang, Z., Li, Y. & Zhang, J. Prevalent positive epistasis in Escherichia coli and
Saccharomyces cerevisiae metabolic networks. *Nat Genet* **42**, 272–276 (2010).

50. Cisneros, A. F. *et al.* Epistasis between promoter activity and coding mutations shapes gene
evolvability. *Sci Adv* **9**, eadd9109 (2023).

51. Abbate, E. *et al.* Optimizing the strain engineering process for industrial-scale production of bio-
based molecules. *Journal of Industrial Microbiology and Biotechnology* **50**, kuad025 (2023).
52. Krivoruchko, A., Zhang, Y., Siewers, V., Chen, Y. & Nielsen, J. Microbial acetyl-CoA metabolism
and metabolic engineering. *Metabolic Engineering* **28**, 28–42 (2015).
53. Janßen, H. & Steinbüchel, A. Fatty acid synthesis in Escherichia coli and its applications towards
the production of fatty acid based biofuels. *Biotechnol Biofuels* **7**, 7 (2014).
54. El-Mansi, E. M. T. & Holms, W. H. Control of Carbon Flux to Acetate Excretion During Growth
of Escherichia coli in Batch and Continuous Cultures. *Microbiology* **135**, 2875–2883 (1989).
55. Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new perspectives on
genomes, pathways, diseases and drugs. *Nucleic Acids Res* **45**, D353–D361 (2017).
56. Chou, H.-H., Delaney, N. F., Draghi, J. A. & Marx, C. J. Mapping the Fitness Landscape of Gene
Expression Uncovers the Cause of Antagonism and Sign Epistasis between Adaptive Mutations.
PLoS Genet **10**, e1004149 (2014).
342