Environmental and anthropogenic impacts on key fish and mollusc species of the Mediterranean Sea from the Late Pleistocene until the Industrial Revolution

Daniela Leal[1,2*], Konstantina Agiadi[1,3], Marta Coll[1,4], Maria Bas[1]

[1]Institut de Ciències del Mar (ICM-CSIC), Passeig Marítim de la Barceloneta 37-49, 08003 Barcelona, Spain

[2]University of Algarve, Campus de Gambelas, 8005-139 Faro, Portugal

[3]Department of Geology, University of Vienna, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

[4]Ecopath International Initiative (EII), Barcelona, Spain

*Corresponding author: dani.leal2000@gmail.com; +351 96 048 52 94

Keywords:

Atlantic bluefin tuna

Gilthead sea bream

Banded-dye murex

Environmental variability

Overexploitation

Holocene

ABSTRACT

Historical baselines are essential for evaluating the cumulative impacts on modern marine ecosystems, particularly in regions such as the Mediterranean, where human activities have been intensive for millennia and climate change is accelerating. However, quantitative evidence of historical impacts remains fragmented. In this study, we investigate changes in the presence, abundance and body size of the Atlantic bluefin tuna, gilthead sea bream, banded-dye murex and turbinate monodont across the Mediterranean Sea from 130,000 years Before Present until the Industrial Revolution (1850 AD), using geological, archaeological, and historical records. Our results reveal significant temporal shifts in the abundance and body size of the banded-dye murex, Atlantic bluefin tuna and gilthead sea bream. Environmental factors, particularly seawater temperature, were the primary drivers of abundance and size in Atlantic bluefin tuna in the past. Human activities, especially exploitation, influenced the abundance of banded-dye murex and Atlantic bluefin tuna, as well as the body size of gilthead sea bream. These findings underscore the importance of integrating long-term ecological data to better understand the interplay between climate, human pressures and ecosystem dynamics.

1. INTRODUCTION

Marine ecosystems are major components of the Earth system, supporting biodiversity, human societies and economies (Halpern *et al.* 2008). However, they are increasingly under threat. Understanding how multiple stressors affect marine biota requires long-term ecological records, yet continuous baselines remain scarce (Dietl *et al.* 2015). While most ecological studies rely on short-term datasets (Jackson 1997; Dayton *et al.* 1998), geological, archaeological and historical records offer valuable insights into ecosystems' long-term ecosystem responses to environmental and anthropogenic pressures (Jackson *et al.* 2001; Lotze *et al.* 2006; Lotze, Coll and Dunne 2011; Fortibuoni *et al.* 2010; Bas *et al.* 2019, 2023; Agiadi and Albano 2020; Agiadi *et al.* 2023, 2024a; Porz *et al.* 2024).

The Mediterranean Sea, a global biodiversity hotspot (IUCN-MED 2009; Coll *et al.* 2010), hosts emblematic species, such as the Atlantic bluefin tuna (*Thunnus thynnus*) (MacKenzie, Mosegaard and Rosenberg 2009) and the Mediterranean monk seal (*Monachus monachus*) (Reijnders, Verriopoulos and Brasseur 1997). Past changes in climate and ocean connectivity have shaped the Mediterranean ecosystems, influencing species migrations and distributions (Agiadi *et al.* 2011, 2018, 2024b). Since the Industrial Revolution, the Mediterranean has warmed faster than the global average (Salat *et al.* 2019; Artana *et al.* Submitted), with profound consequences for marine biodiversity and ecosystems functioning (Garrabou *et al.* 2009; Smale *et al.* 2019; Trisos, Merow and Pigot 2020; Ouled-Cheikh *et al.* 2022).

Marine resources have supported Mediterranean societies for millennia (Madariaga 1964; Mehvar *et al.* 2018). Shellfish exploitation dates back to the Lower Palaeolithic (e.g. ~300 ka B.P., de Lumley 1966), and by the Bronze Age (5-3 ka BP), extensive fisheries and trade networks were established (e.g., Van Neer *et al.* 2004; Guy *et al.* 2018a; Zohar and Artzy 2019), intensifying throughout the Classical period (Sáez Romero 2014; Mylona 2018). Several key marine species were extensively exploited and/or affected by past environmental changes (Leal, Agiadi and Bas 2025). Long-term exploitation combined with environmental variability, has contributed to the reshaping of marine ecosystems (Lotze *et al.* 2006; Lotze, Coll and Dunne 2011). Today, habitat loss, degradation, and overexploitation continue to alter species productivity, fitness and distribution, with cascading effects on

ecosystem structure and functioning (Halpern *et al.* 2008; Coll *et al.* 2010, 2012; Planque *et*

al. 2010; Howarth *et al.* 2014; Artana *et al.* Submitted).

In this study, we examine how environmental variability and human activities have

influenced commercially relevant species in the Mediterranean Sea over the past 130 kyrs

until the Industrial Revolution. Using regional data (Leal, Agiadi and Bas 2025) we analyse

Atlantic bluefin tuna (*Thunnus thynnus*), gilthead sea bream (*Sparus aurata*), banded-dye

murex (*Hexaplex trunculus*) and turbinate monodont (*Phorcus turbinatus*), testing whether:

1) temperature oscillations from the Late Pleistocene to the Late Holocene (Cacho, Grimalt

and Canals 2002; Marchal *et al.* 2002; Rohling *et al.* 2002; Cacho *et al.* 2006; Schmiedl *et al.*

2010) influenced species abundance and body size; and 2) human exploitation compounded

these effects over time.

2. MATERIALS AND METHODS

2.1 Study area

83   The Mediterranean Sea is a semi-enclosed basin (3,000,000 km^2, maximum depth ~5,300 m

84   (Robinson and Golnaraghi 1994). For analytical purposes, we divided the basin into three

85   sub-regions: Western, Eastern and Adriatic, defined by the straits of Sicily and Otranto

86   (Millot and Taupier-Letage 2005; Rio *et al.* 2007; Robinson *et al.* 2009) (Figure 1). The

87   Marmara and Black seas were included in the Eastern Mediterranean.

 Figure 1 - Map of the Mediterranean Sea showing the sub-regions (Western Mediterranean, Eastern

90   Mediterranean and Adriatic Sea) and their natural divisions (Strait of Otranto and Strait of Sicily),

91   with the data locations (Table S6). Variables used include the Number of Identified Specimens (NISP).

92   Minimum Number of Individuals (MNI), Total Length (TL) and Basal Diameter (BD).

2.2 Species and temporal framework

Species data were available across both spatial and temporal scales within the

Mediterranean sub-regions, although catch data were only available for the Atlantic bluefin

tuna.

We adopted the temporal divisions defined by the International Commission on Stratigraphy

(ICS; Cohen, Harper and Gibbard 2013) into the Late Pleistocene (130,000–11,650 years

Before Present (BP); Head *et al.* 2021), Early Holocene (11,650–8,186 years BP; Stuiver and

Reimer 1986), Middle Holocene (8,186–4,200 years BP; Stuiver and Reimer 1986) and Late

Holocene (4,200–0 years BP; Stuiver and Reimer 1986). Each period is characterised by

major climatic events (e.g., glacial-interglacial cycles, Lionello *et al.* 2023) and societal

transitions (e.g., from hunter-gatherers to sedentary communities, Isern *et al.* 2014; Valla

2018). The Late Pleistocene includes the Last Interglacial (130-116 kyrs BP; Kukla *et al.* 2002)

and the Younger Dryas (14,749–13,646 years BP; Benjamin *et al.* 2017). The Holocene

features notable temperature fluctuations, such as the Medieval Warm Period (MWP;

1,050–650 BP) and the Little Ice Age (LIA; 650–100 years BP) (Cisneros *et al.* 2016).

Mediterranean fisheries evolved significantly between the pre-industrial (130 kyrs - 1850

AD) to the post-industrial era (1850 AD – present day). Fishing practices transitioned from

manual and artisanal techniques (harpoons, gillnets and traps; Ruas 2005; Roberts *et al.*

2011; Horden and Purcell 2013; Pedergnana *et al.* 2021), to mechanised fishing and steam-

powered trawlers with the onset of industrial fishing (1880 AD) (Gabriel *et al.* 2005).

2.3 Database

The database compiled by Leal, Agiadi and Bas (2025a) was developed through a systematic

review following the PRISMA methodology (Page *et al.* 2021), integrating peer-reviewed and

secondary literature with geological, archaeological and historical sources (Table S7).

Presence and abundance data were quantified using two zooarchaeological metrics, which

are commonly used to infer patterns of human exploitation of faunal resources: the Number

of Identified Specimens (NISP; Grayson 1984) and Minimum Number of Individuals (MNI;

Lyman 1994).. The review also incorporated over 1,200 entries of Atlantic bluefin tuna catch

data from Italian tuna traps (1700–1936; Addis *et al.* 2008; Longo and Clark 2012; Polanco-

Martínez *et al.* 2018), which serve as long-term proxies for abundance (Ravier 2001).

Historical abundance trends were analysed in relation to Sea Surface Temperature (SST;

Jalali *et al.* 2016), human civilisations and fishing technologies. Body size (BS) data were

derived from multiple sources: a) Total Length (TL) for Atlantic bluefin tuna from Andrews *et*

al. (2022, 2023a, 2023c, 2023b); b) TL for gilthead sea bream from Western archaeological

sites (Fernández-López De Pablo and Gabriel 2016) and Eastern sites (Guy *et al.* 2018; Fuller

et al. 2020); and c) Basal Diameter (BD) of the turbinate monodont from Haua Fteah cave,

Libya (Hunt *et al.* 2011; Prendergast *et al.* 2016).

2.4 Data Analyses

Normality and homoscedasticity were tested using the Shapiro–Wilk and Levene's tests,

respectively (*car* R package; Fox and Weisberg 2019). For NISP and MNI, non-parametric

tests were applied: the Mann-Whitney U test for two-group comparisons and the Kruskal-

Wallis test for multi-group comparisons, followed by post-hoc pairwise Wilcoxon tests. For

TL, a two-sample t-test for Western Mediterranean tuna and a Mann-Whitney *U* test for the

Eastern Mediterranean tuna. For the gilthead sea bream, ANOVA with a Tukey's HSD post-

hoc test was used to compare three periods. For BD, ANOVA (mean values) and Welch

ANOVA (maximum values) were applied to multi-period comparisons. Catch data trends

were analysed using the Mann-Kendall test (*trend* R package; Pohlert 2015). Data

classification was performed using Jenks' natural breaks method (Jenks 1977) (*classInt* R

package; Bivand 2006). Inter-period differences were assessed with Kruskal-Wallis and

pairwise Wilcoxon tests. A significance threshold of $p \leq 0.05$ was applied, with Bonferroni

correction. Maps were produced in QGIS 3.34.13, and statistical analyses in R 4.4.3 (R Core

Team 2025).

3. RESULTS

3.1 Presence and abundance of targeted species

NISP data for gilthead sea bream and Atlantic bluefin tuna span the entire temporal framework. However, comparisons between periods within each region revealed no statistically significant differences (Table S2.1; Figures S3.1B–S3.4B). Among molluscs, only turbinate monodont presented sufficient data for temporal analysis across the Mediterranean Sea, but no significant differences were detected across periods (Table S2.1; Figures S3.5B and S3.6B).

MNI data for turbinate monodont were available for both the Western and Eastern Mediterranean, yet no significant differences were found between periods (Table S2.2, Figures S3.7 and S3.8). In contrast, banded-dye murex showed significant temporal variation in the MNI in the Western Mediterranean (H_2 = 15.792; p = 0.0004, Figure 2A), particularly between the Early and Late Holocene (p = 0.0013; Figure 2B; Table S2.2).

Figure 2 - Distribution of Minimum Number of Individuals (MNI) for banded-dye murex in the Western Mediterranean. (A) Temporal distribution of records, with major climatic events (purple rectangles) and human cultural periods (black dashed lines); (B) Boxplots of MNI data by geological period: Early Holocene (EH), Middle Holocene (MH), and Late Holocene (LH). Blue dots represent mean values and bars indicate standard deviation. Letters denote statistically significant differences. Sample sizes (n) are shown above each boxplot.

3.2 Catch records

Catch data (annual totals) were available only for Atlantic bluefin tuna from 250 to 14 years
BP, corresponding to the period between 1700 and 1936 AD (Figure 3). Recorded values
ranged from 0 to approximately 12,500 individuals per year along the coasts of Sardinia and
Sicily in the Western Mediterranean (Figure 1). Two major peaks were observed: around 200
years BP (~1750 AD) during the LIA with ~10,000 individuals, and between 75–45 years BP
(1875–1905 AD), following the Industrial Revolution (1850 AD) (~12,000 individuals).

Figure 3 – Atlantic bluefin tuna catch records from the Late Holocene (250–14 years BP;1700–1936
AD), based on tuna traps from the Western Mediterranean Sea. Black points represent pre-industrial
catches (250–100 years BP; 1700–1850 AD), grey points represent post-industrial catches (99–14
years BP; 1851–1936 AD). Red points indicate reconstructed Sea Surface Temperature (SST) data
from the Gulf of Lions (Jalali *et al.* 2016). The blue rectangle refers to the Little Ice Age (LIA). Vertical
lines indicate the use of tuna traps, the onset of industrial fisheries (70 BP; 1880 AD; von Brandt

1972; Gabriel *et al.* 2005) and WWI (World War I). Note: tuna trap technology remained largely

unchanged until the 1960s (Doumenge 1998).

A significant declining trend in catches was detected between 250-14 years BP (S = −3.98 e^5;

Kendall's tau coefficient = −2.39 e^{-1}; p < 2.2e^{-16}), with differences between specific intervals

(H_5 = 407.55; p < 2.2e^{-16}; Table S4; Figure S4).

3.3 Body size

In the Western Mediterranean, a significant increase in the TL of Atlantic bluefin was found

between the MWP and the LIA ($t_{15.141}$ = 2.967; p = 0.01; Figures 4 and S5.1; Table S2.4).

Figure 4 - Distribution of Total Length (TL) data for Atlantic bluefin tuna in the Western Mediterranean. (A) Temporal distribution of records, including major climatic events (purple rectangles) and anthropogenic periods (black dashed lines); (B) Late Holocene distribution of records, with vertical lines indicating the main fishing technologies used during the period.

195
196
197
198

In the Eastern Mediterranean, TL of Atlantic bluefin tuna also showed significant increases between the Roman Period (2,500–1,050 years BP) and the MWP (1,050–650 years BP), as well as within the MWP itself ($W = 8$; $p = 0.01035$) (Figures 5 and S5.2; Table S2.4).

Figure 5 - Distribution of Total Length (TL) data for Atlantic bluefin tuna in the Eastern Mediterranean. (A) Temporal distribution of records, including major climatic events (purple rectangles) and anthropogenic periods (black dashed lines); (B) Late Holocene distribution of records, with vertical lines indicating the main fishing technologies used during the period.

For gilthead sea bream, TL data were available for the Eastern Mediterranean (Figure 6).

Pairwise comparisons (F = 10.28; p = 7.54e^{-5}) revealed significant reductions in TL between

the Early and Late Holocene (p = 0.0061), the Middle and Late Holocene (p = 0.0010; Figure

6B), and between the Bronze and Iron Ages and the Byzantine Period (H_2 = 8.501; p = 0.014,

Table S2.4 and Figure S5.3).

Figure 6 - Distribution of Total Length (TL) data for gilthead sea bream in the Eastern Mediterranean.

(A) Temporal distribution of records, including major climatic events (purple rectangles) and

anthropogenic periods (black dashed lines); (B) Boxplots of TL data for each geological period: Early Holocene (EH), Middle Holocene (MH), and Late Holocene (LH). Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences. Sample sizes (n) are shown above each boxplot; (C) Late Holocene distribution of records, with vertical lines indicating the main fishing technologies used during the period.

For the turbinate monodont, BD data were available for the Eastern Mediterranean, including both mean and maximum values. No significant differences were found between the geological periods, for either metric (Table S2.5; Figures S5.4-S5.5).

4. DISCUSSION

4.1 Environmental Driven Changes in Atlantic Bluefin Tuna

Atlantic bluefin tuna is a highly migratory species that travels between cold feeding grounds in the North Atlantic and warm spawning areas in the Mediterranean Sea (Mather, Mason and Jones 1995; Block *et al.* 2001, 2005; Rooker *et al.* 2007). Climatic changes influence its growth, reproduction, population size, mortality and migration patterns (Pepin 1991; Graham and Dickson 2004). Our data show a peak in tuna catches during the late LIA, which may be linked to cooler conditions (Cisneros *et al.* 2016; Margaritelli *et al.* 2020) and enhanced nutrient availability and primary productivity in the North Atlantic (Pourmand *et al.* 2007), resulting in a greater number of mature individuals entering the Mediterranean. Additionally, tuna migration may have shifted toward the basin during the LIA, due to increased climatic instability and extreme events, leading to higher recruitment in the Western Mediterranean and greater concentration of schools near historical tuna traps (Ravier and Fromentin 2004, Ganzedo *et al.* 2009). Considering the 6–7 year lag between hatching and recruitment (Zorita *et al.* 2005), the increased landings observed in the later stages of the LIA may reflect changes in distribution and behaviour rather than an overall increase in population size.

Temperature also plays a key role in shaping plankton dynamics, influencing both production and distribution (e.g., McGowan, Cayan and Dorman 1998; Beaugrand *et al.* 2002). Our results suggest that cooler conditions during the late LIA enhanced trophic productivity,

increasing tuna availability and catch rates in the Western Mediterranean. Similar patterns have been observed for Pacific bluefin tuna, where SST is a stronger driver of recruitment than spawning stock biomass, likely due to its influence on juvenile habitat suitability (Muhling *et al.* 2018). Temperature-driven changes in plankton propagate through the food web, altering prey availability and influencing tuna migration (Mather, Mason and Jones 1995; Ravier and Fromentin 2004).

We also observed an increase in TL of Atlantic bluefin tuna during the late LIA, likely driven by cooler conditions, improved feeding opportunities, and altered migration patterns that concentrated larger individuals in the region. This aligns with the species' recruitment lag (Santamaria *et al.* 2009), explaining the delayed increase in adult abundance and TL. Although warmer conditions can enhance consumer-resource interactions (Allen, Gillooly and Brown 2005; López-Urrutia *et al.* 2006; O'Connor 2009), the increase in TL under cooler SST suggests indirect effects, such as enhanced trophic productivity or greater adult retention, played a key role. The species' broad dietary niche (Olesen and Jordano 2002; Pardo *et al.* 2025) likely supported adaptation to shifting prey fields, while cooler, more productive conditions may have improved larval growth and survival (García *et al.* 2006; Catalán *et al.* 2011; Reglero *et al.* 2018; Trueman *et al.* 2023). Consistent with other Mediterranean fish (Hattab *et al.* 2021), larger body sizes under colder temperatures indicate that long-lived top predators like the Atlantic bluefin tuna (Santamaria *et al.* 2009) may deviate from general patterns due to their complex ecological responses (Fromentin and Powers 2005).

4.2 Human Exploitation and Species Decline

This study highlights the long-term effects of human exploitation on species abundance and body size in the Mediterranean Sea, supported by historical and ethnographic records of traditional practices (Galili, Zemer and Rosen 2013; De Nicolò 2018; Lucchetti *et al.* 2023).

The banded-dye murex was intensively harvested in the Late Holocene, particularly in the Roman Period (Oliver 2015). Its intertidal habitat made it easily accessible to early communities (Bonanno *et al.* 2016; Fernández-López De Pablo and Gabriel 2016) and later enabled large-scale exploitation for purple dye production (Michel and McGovern 1987; Oliver 2015). Shellfish provided a stable, year-round food resource, supporting coastal

populations during periods of environmental stress or resource scarcity (Erlandson 1988; Colonese *et al.* 2011). However, this accessibility also made them vulnerable to overexploitation, especially with the rise of industrial dye production (Michel and McGovern 1987; Erlandson *et al.* 2008; Oliver 2015). These dynamics illustrate the dual role of shellfish as both resilient resources and indicators of human-induced ecosystem change.

The long-term decline in Atlantic bluefin tuna abundance reflects sustained fishing pressure (Polanco-Martínez *et al.* 2018; this study). Traditional tuna traps, once considered sustainable (De la Serna *et al.* 2012; Longo 2012), were eventually replaced by industrial technologies (e.g., iron anchors, steel cables; Lentini 2001), and steam-powered gears; Joseph 2003), which enabled large-scale exploitation and contributed to the stock collapse (Andrews *et al.* 2022). Removing top predators can destabilise food webs through top-down effects (Pauly *et al.* 1998; Scheffer, Carpenter and Young 2005). Although direct evidence of trophic cascades involving tuna in the Mediterranean Sea is limited, isotopic and archaeological data suggest shifts in its trophic ecology during historical periods (Andrews *et al.* 2022, 2023c), altering predator-prey dynamics. Ecopath models from the Catalan Sea confirm these findings, showing food web restructuring following the loss of top and intermediate trophic-level species (Forrestal *et al.* 2012), consistent with broader patterns in the Mediterranean (Stergiou, Tsikliras and Pauly 2009) and the North Sea (Mariani *et al.* 2017).

4.3 Changes in Body Size

Overexploitation may have contributed to the reduction in body size of gilthead sea bream from the Early to the Late Holocene, particularly between the Bronze and Iron Age and the Byzantine Period. In contrast, Atlantic bluefin tuna showed the opposite trend, with increasing body sizes from the Roman Period through the MWP and the LIA.

Historical records show declining tuna landings in the Gulf of Cadiz between the mid-16[th] and 18[th] centuries (Sarmiento 1757), likely due to juvenile overfishing and habitat degradation (García 2016). However, tuna body size increased during the MWP and LIA, possibly reflecting fisheries-induced evolution (FIE), where intense exploitation selects for early-maturing, faster-growing genotypes (Law 2000; Enberg *et al.* 2012). Similar long-term effects have been documented in Baltic cod, with larger individuals in the Neolithic (~4500

yrs BP) compared to modern specimens (Limburg *et al.* 2008). From the Iron Age onward, the adoption of fish traps (Mylona 2021) and later industrial gears such as purse seines marked a shift to large-scale fisheries, likely influencing tuna body size across the basin.

Gilthead sea bream showed signs of overexploitation in the Eastern Mediterranean during the Late Holocene, especially from the Bronze Age to the Byzantine period, with a marked decline in body size (Guy *et al.* 2018; this study). Similar trends have been observed in other sparids, such as *Archosargus probatocephalus,* under long-term fishing pressure (Guiry *et al.* 2021). Zooarchaeological records from Orkney (Scotland) also reveal offshore overfishing of large cod between the 11[th] and 13[th] centuries, followed by declines in abundance and size (Harland and Barrett 2012). The shift to industrial-scale, less selective fishing gear (Galili, Zemer and Rosen 2013) likely reinforced these reductions. Size-selective harvesting can alter trophic interactions and reduce ecosystem resilience (Shackell *et al.* 2010; Garcia *et al.* 2012), underscoring the long-term ecological consequences of sustained exploitation.

4.4 Limitations

While this study provides valuable insights, several limitations must be acknowledged. Our analyses focused on data-rich, historically and economically relevant species, potentially overlooking less-studied taxa and broader ecosystem dynamics. Data scarcity, particularly in the Adriatic Sea, reflects the uneven availability of published records (Leal, Agiadi and Bas 2025). Most data originate from coastal archaeological sites, which may bias the representation of past exploitation patterns (Galili and Rosen 2008; Leal, Agiadi and Bas 2025). Additional constraints include variability in dating, body size measurements, and taphonomic preservation, which limit the accuracy of past population reconstructions and inter-period comparability (Agiadi *et al.* 2024b, 2024a). Moreover, uncertainty regarding the fishing gear used and the exclusion of non-peer-reviewed sources, such as logbooks or diaries, introduce further biases (Agiadi *et al.* 2024a). Despite these limitations, our study advances the understanding of how environmental variability and human activities have shaped key Mediterranean marine species over long timescales. Emerging techniques, such as ancient (aDNA) and sedimentary ancient DNA (*seda*DNA), offer promising avenues for higher resolution reconstructions of paleoecological conditions and ecosystem dynamics (De Schepper *et al.* 2019; Monk *et al.* 2021; Kjær *et al.* 2022).

Conclusions

This study underscores the importance of adopting long-term perspectives to understand changes in the marine ecosystem. Our findings suggest that both environmental and human factors influenced Atlantic bluefin tuna, while banded-dye murex and gilthead sea bream were mainly affected by sustained human pressure. Understanding these historical trajectories helps clarify the mechanisms and magnitude of ecological shifts, contributing to the interpretation of "shifting baselines" (Pauly *et al.* 1998). By integrating palaeoecological, archaeological, and historical records, this research provides a robust framework for reconstructing historical baselines and informing future management of Mediterranean marine ecosystems.

ACKNOWLEDGEMENTS

This research contributes to the objectives of Q-MARE (a PAGES working group), and it is part of the Integrated Marine Ecosystem Assessments (iMARES) research group funded by Agència de Gestió d'Ajuts Universitaris i de Recerca (Generalitat de Catalunya).

AUTHOR CONTRIBUTIONS

Conceptualisation: DL, KA, and MB. Methodology: DL, KA, MC and MB. Data curation: DL. Formal analysis: DL. Writing—original draft: DL. Writing—review & editing: KA, MC and MB. Supervision: KA and MB.

DECLARATION OF INTERESTS

The authors declare that there are no conflicts of interest.

FUNDING

363 This work was supported by the Erasmus+ Program Grant no. 2022-1-PT01-KA131-HED-
364 000061155; MCIN/AEI/10.13039/501100011033 and NextGenerationEU/PRTR, Grant no.
365 FJC2020-043762-I; EU HORIZON-CL6-2021-BIODIV-01-04 project Ges4Seas (Grant Agreement
366 101059877); the "Severo Ochoa Centre of Excellence" accreditation grant CEX2024-001494-S
367 funded by AEI 10.13039/501100011033.

369 DATA AVAILABILITY

370 All data produced for this work are available within the manuscript and in the
371 supplementary material.

373 REFERENCES

Addis P, Dean JM, Pesci P *et al.* Effects of local scale perturbations in the Atlantic bluefin
tuna (Thunnus thynnus L.) trap fishery of Sardinia (W. Mediterranean). *Fish Res*
2008;**92**:242–54.

Agiadi K, Albano PG. Holocene fish assemblages provide baseline data for the rapidly
changing eastern Mediterranean. *The Holocene* 2020;**30**:1438–50.

Agiadi K, Caswell BA, Almeida R *et al.* Geohistorical insights into marine functional
connectivity. Selden R (ed.). *ICES J Mar Sci* 2024a;**81**:1884–911.

Agiadi K, Girone A, Koskeridou E *et al.* Pleistocene marine fish invasions and
paleoenvironmental reconstructions in the eastern Mediterranean. *Quat Sci Rev*
2018;**196**:80–99.

Agiadi K, Hohmann N, Gliozzi E *et al.* Late Miocene transformation of Mediterranean Sea
biodiversity. *Sci Adv* 2024b;**10**:eadp1134.

Agiadi K, Quillévéré F, Nawrot R *et al.* Palaeontological evidence for community-level

decrease in mesopelagic fish size during Pleistocene climate warming in the eastern

Mediterranean. *Proc R Soc B Biol Sci* 2023;**290**:20221994.

Agiadi K, Triantaphyllou M, Girone A *et al.* The early Quaternary palaeobiogeography of the

eastern Ionian deep-sea Teleost fauna: A novel palaeocirculation approach.

Palaeogeogr Palaeoclimatol Palaeoecol 2011;**306**:228–42.

Allen AP, Gillooly JF, Brown JH. Linking the global carbon cycle to individual metabolism.

Funct Ecol 2005;**19**:202–13.

Andrews AJ, Di Natale A, Addis P *et al.* Vertebrae reveal industrial-era increases in Atlantic

bluefin tuna catch-at-size and juvenile growth. Andrews AH (ed.). *ICES J Mar Sci*

2023a;**80**:836–47.

Andrews AJ, Di Natale A, Bernal-Casasola D *et al.* Exploitation history of Atlantic bluefin tuna

in the eastern Atlantic and Mediterranean—insights from ancient bones. Poulsen B

(ed.). *ICES J Mar Sci* 2022;**79**:247–62.

Andrews AJ, Orton D, Onar V *et al.* Isotopic life-history signatures are retained in modern

and ancient Atlantic bluefin tuna vertebrae. *J Fish Biol* 2023b;**103**:118–29.

Andrews AJ, Pampoulie C, Di Natale A *et al.* Exploitation shifted trophic ecology and habitat

preferences of Mediterranean and Black Sea bluefin tuna over centuries. *Fish Fish*

2023c;**24**:1067–83.

Artana C, Kaplan A, Ramírez F *et al.* Marine Heat Waves Are Transforming Western

Mediterranean Marine Ecosystems. Submitted.

Bas M, Briz I Godino I, Álvarez M *et al.* Back to the future? Late Holocene marine food web

structure in a warm climatic phase as a predictor of trophodynamics in a warmer

South-Western Atlantic Ocean. *Glob Change Biol* 2019;**25**:404–19.

Bas M, Tivoli AM, Briz I Godino I *et al.* Changing diets over time: knock-on effects of marine

megafauna overexploitation on their competitors in the southwestern Atlantic

Ocean. *Paleobiology* 2023;**49**:176–90.

Beaugrand G, Reid PC, Ibañez F *et al.* Reorganization of North Atlantic Marine Copepod

Biodiversity and Climate. *Science* 2002;**296**:1692–4.

Benjamin J, Rovere A, Fontana A *et al.* Late Quaternary sea-level changes and early human

societies in the central and eastern Mediterranean Basin: An interdisciplinary review.

Quat Int 2017;**449**:29–57.

Bivand   R.   classInt: Choose   Univariate   Class   Intervals.   2006,   DOI:

10.32614/cran.package.classint.

Block BA, Dewar H, Blackwell SB *et al.* Migratory Movements, Depth Preferences, and

Thermal Biology of Atlantic Bluefin Tuna. *Science* 2001;**293**:1310–4.

Block BA, Teo SLH, Walli A *et al.* Electronic tagging and population structure of Atlantic

bluefin tuna. *Nature* 2005;**434**:1121–7.

Bonanno A, Barra M, Basilone G *et al.* Environmental processes driving anchovy and sardine

distribution in a highly variable environment: the role of the coastal structure and

riverine input. *Fish Oceanogr* 2016;**25**:471–90.

von Brandt A. *Fish Catching Methods of the World*. Wiley-Blackwell, 1972.

Cacho I, Grimalt JO, Canals M. Response of the Western Mediterranean Sea to rapid climatic

variability during the last 50,000 years: a molecular biomarker approach. *J Mar Syst*

2002;**33–34**:253–72.

Cacho I, Shackleton N, Elderfield H *et al.* Glacial rapid variability in deep-water temperature

and δ18O from the Western Mediterranean Sea. *Quat Sci Rev* 2006;**25**:3294–311.

Catalán IA, Tejedor A, Alemany F *et al.* Trophic ecology of Atlantic bluefin tuna *Thunnus*

thynnus larvae. *J Fish Biol* 2011;**78**:1545–60.

Cisneros M, Cacho I, Frigola J *et al.* Sea surface temperature variability in the central-western

Mediterranean Sea during the last 2700 years: a multi-proxy and multi-record

approach. *Clim Past* 2016;**12**:849–69.

Cohen KM, Harper DAT, Gibbard PL. *ICS International Chronostratigraphic Chart*.

International Commission on Stratigraphy (ICS), IUGS, 2013.

Coll M, Piroddi C, Albouy C *et al.* The Mediterranean Sea under siege: spatial overlap

between marine biodiversity, cumulative threats and marine reserves. *Glob Ecol*

Biogeogr 2012;**21**:465–80.

Coll M, Piroddi C, Steenbeek J *et al.* The Biodiversity of the Mediterranean Sea: Estimates,

Patterns, and Threats. Bograd SJ (ed.). *PLoS ONE* 2010;**5**:e11842.

Colonese AC, Mannino MA, Bar-Yosef Mayer DE *et al.* Marine mollusc exploitation in

Mediterranean prehistory: An overview. *Quat Int* 2011;**239**:86–103.

Dayton PK, Tegner MJ, Edwards PB *et al.* SLIDING BASELINES, GHOSTS, AND REDUCED

EXPECTATIONS IN KELP FOREST COMMUNITIES. *Ecol Appl* 1998;**8**:309–22.

De la Serna JM, Macías D, de Urbina JO *et al.* Study on the eastern Atlantic and

Mediterranean bluefin tuna stock using the Spanish traps as scientific observatories.

Collect. Vol. Sci. Pap. ICCAT. 2012:331–43.

De Nicolò ML. Evolution of the gangamo (ganguy) fishing gear in the Mediterranean Sea

from Antiquity to the early Modern period and the debate on trawling sustainability.

Reg Stud Mar Sci 2018;**21**:79–85.

De Schepper S, Ray JL, Skaar KS *et al.* The potential of sedimentary ancient DNA for

reconstructing past sea ice evolution. *ISME J* 2019;**13**:2566–77.

Dietl GP, Kidwell SM, Brenner M *et al.* Conservation Paleobiology: Leveraging Knowledge of

the Past to Inform Conservation and Restoration. *Annu Rev Earth Planet Sci*

2015;**43**:79–103.

Doumenge F. L'histoire des pêches thonières. *ICCAT Sci Pap* 1998;**50**:753–802.

Erlandson JM. The Role of Shellfish in Prehistoric Economies: A Protein Perspective. *Am*

Antiq 1988;**53**:102–9.

Erlandson JM, Rick TC, Braje TJ *et al.* Human impacts on ancient shellfish: a 10,000 year

record from San Miguel Island, California. *J Archaeol Sci* 2008;**35**:2144–52.

Fernández-López De Pablo J, Gabriel S. El Collado shell midden and the exploitation patterns

of littoral resources during the Mesolithic in the Eastern Iberian Peninsula. *Quat Int*

2016;**407**:106–17.

Forrestal F, Coll M, Die D *et al.* Ecosystem effects of bluefin tuna Thunnus thynnus thynnus

aquaculture in the NW Mediterranean Sea. *Mar Ecol Prog Ser* 2012;**456**:215–31.

Fortibuoni T, Libralato S, Raicevich S *et al.* Coding Early Naturalists' Accounts into Long-Term

Fish Community Changes in the Adriatic Sea (1800–2000). Thrush S (ed.). *PLoS ONE*

2010;**5**:e15502.

Fox J, Weisberg S. *An R Companion to Applied Regression*. 3rd ed. Thousand Oaks, CA: Sage,

2019.

Fromentin J, Powers JE. Atlantic bluefin tuna: population dynamics, ecology, fisheries and

management. *Fish Fish* 2005;**6**:281–306.

Fuller BT, Van Neer W, Linseele V *et al.* Fish δ13C and δ15N results from two Bronze/Iron

Age sites (Tell Tweini & Sidon) along the Levantine coast. *J Archaeol Sci Rep*

2020;**29**:102066.

Gabriel O, Lange K, Dahm E *et al.* eds. *Fish Catching Methods of the World*. 1st ed. Wiley,

2005.

Galili E, Rosen B. Fishing Gear from a 7th-Century Shipwreck off Dor, Israel. *Int J Naut*

Archaeol 2008;**37**:67–76.

Galili E, Zemer A, Rosen B. Ancient fishing gear and associated artifacts from underwater

explorations in Israel–a comparative study. *Archaeofauna*. 2013:166.

García A, Cortés D, Ramírez T *et al.* First data on growth and nucleic acid and protein content

of field-captured Mediterranean bluefin (Thunnus thynnus) and albacore (Thunnus

alalunga) tuna larvae: a comparative study. *Sci Mar* 2006;**70**:67–78.

Garcia SM, Kolding J, Rice J *et al.* Reconsidering the Consequences of Selective Fisheries.

Science 2012;**335**:1045–7.

Garrabou J, Coma R, Bensoussan N *et al.* Mass mortality in Northwestern Mediterranean

rocky benthic communities: effects of the 2003 heat wave. *Glob Change Biol*

2009;**15**:1090–103.

Graham JB, Dickson KA. Tuna comparative physiology. *J Exp Biol* 2004;**207**:4015–24.

Grayson K. *Quantitative Zooarchaeology*. New York: Academic Press, 1984.

Guiry EJ, Kennedy JR, O'Connell MT *et al.* Early evidence for historical overfishing in the Gulf

of Mexico. *Sci Adv* 2021;**7**:eabh2525.

Guy S-V, Thomas T, Irit Z *et al.* Tooth oxygen isotopes reveal Late Bronze Age origin of

Mediterranean fish aquaculture and trade. *Sci Rep* 2018;**8**:14086.

Halpern BS, Walbridge S, Selkoe KA *et al.* A Global Map of Human Impact on Marine

Ecosystems. *Science* 2008;**319**:948–52.

Harland JF, Barrett JH. The maritime economy: Fish bone. *Being an Islander: Production and*

Identity at Quoygrew, Orkney, AD 900–1600. Cambridge: McDonald Institute for

Archaeological Research, 2012, 115–38.

Hattab T, Gucu A, Ventero A *et al.* Temperature strongly correlates with regional patterns of

body size variation in Mediterranean small pelagic fish species. *Mediterr Mar Sci*

2021;**22**:800.

Head MJ, Pillans B, Zalasiewicz JA *et al.* Formal ratification of subseries for the Pleistocene

Series of the Quaternary System. *Episodes* 2021;**44**:241–7.

Horden P, Purcell N. *The Corrupting Sea: A Study of Mediterranean History*. Nachdr. Malden,

Mass.: Blackwell, 2013.

Howarth LM, Roberts CM, Thurstan RH *et al.* The unintended consequences of simplifying

the sea: making the case for complexity. *Fish Fish* 2014;**15**:690–711.

Hunt CO, Reynolds TG, El-Rishi HA *et al.* Resource pressure and environmental change on the

North African littoral: Epipalaeolithic to Roman gastropods from Cyrenaica, Libya.

Quat Int 2011;**244**:15–26.

Isern N, Fort J, Carvalho AF *et al.* The Neolithic Transition in the Iberian Peninsula: Data

Analysis and Modeling. *J Archaeol Method Theory* 2014;**21**:447–60.

IUCN-MED. *IUCN Centre for Mediterranean Cooperation*. https://iucn.org/our-

work/region/mediterranean

Jackson JBC. Reefs since Columbus. *Coral Reefs* 1997;**16**:S23–32.

Jackson JBC, Kirby MX, Berger WH *et al.* Historical Overfishing and the Recent Collapse of

Coastal Ecosystems. *Science* 2001;**293**:629–37.

Jalali B, Sicre M-A, Bassetti M-A *et al.* NOAA/WDS Paleoclimatology - Gulf of Lion,

Mediterranean Sea 10,000 Year Biomarker Data and SST Reconstruction. 2016, DOI:

10.25921/6S87-Y952.

Joseph J. *Managing Fishing Capacity of the World Tuna Fleet*. Rome: Food and Agriculture

Organization (FAO), 2003.

Kjær KH, Winther Pedersen M, De Sanctis B *et al.* A 2-million-year-old ecosystem in

Greenland uncovered by environmental DNA. *Nature* 2022;**612**:283–91.

Kukla GJ, Bender ML, De Beaulieu J-L *et al.* Last Interglacial Climates. *Quat Res* 2002;**58**:2–13.

Leal D, Agiadi K, Bas M. Commercially relevant species in the Mediterranean Sea: A

perspective from Late Pleistocene to the Industrial Revolution. *J Archaeol Sci*

2025;**179**:106242.

Lentini R. Favignana nella Seconda Metà dell'800: Innovazioni e Mercato. *La Pesca in Italia*

Tra Età Moderna e Contemporanea: Produzione, Mercato, Consumo. Sassari, Italy:

Editrice Democratica Sarda, 2001, 507–20.

Limburg KE, Walther Y, Hong B *et al.* Prehistoric versus modern Baltic Sea cod fisheries:

selectivity across the millennia. *Proc R Soc B Biol Sci* 2008;**275**:2659–65.

Lionello P, Giorgi F, Rohling E *et al.* Mediterranean climate. *Oceanography of the*

Mediterranean Sea. Elsevier, 2023, 41–91.

Longo SB. Mediterranean Rift: Socio-Ecological Transformations in the Sicilian Bluefin Tuna

Fishery. *Crit Sociol* 2012;**38**:417–36.

Longo SB, Clark B. The Commodification of Bluefin Tuna: The Historical Transformation of

the Mediterranean Fishery. *J Agrar Change* 2012;**12**:204–26.

López-Urrutia Á, San Martin E, Harris RP *et al.* Scaling the metabolic balance of the oceans.

Proc Natl Acad Sci 2006;**103**:8739–44.

Lotze HK, Coll M, Dunne JA. Historical Changes in Marine Resources, Food-web Structure and

Ecosystem Functioning in the Adriatic Sea, Mediterranean. *Ecosystems* 2011;**14**:198–

222.

Lotze HK, Lenihan HS, Bourque BJ *et al.* Depletion, Degradation, and Recovery Potential of

Estuaries and Coastal Seas. *Science* 2006;**312**:1806–9.

Lucchetti A, Petetta A, Bdioui M *et al. Catalogue of Fishing Gear in the Mediterranean and*

Black Sea Region. FAO, 2023.

de Lumley H. Les fouilles de Terra Amata à Nice. Premiers résultats. *Bull Mus*

D'Anthropologie Préhistoire Monaco 1966;**13**:29–51.

Lyman R. *Vertebrate Taphonomy*. Cambridge: Cambridge University Press, 1994.

MacKenzie BR, Mosegaard H, Rosenberg AA. Impending collapse of bluefin tuna in the

northeast Atlantic and Mediterranean. *Conserv Lett* 2009;**2**:26–35.

Marchal O, Cacho I, Stocker TF *et al.* Apparent long-term cooling of the sea surface in the

northeast Atlantic and Mediterranean during the Holocene. *Quat Sci Rev*

2002;**21**:455–83.

Margaritelli G, Cacho I, Català A *et al.* Persistent warm Mediterranean surface waters during

the Roman period. *Sci Rep* 2020;**10**:10431.

Mariani P, Andersen KH, Lindegren M *et al.* Trophic impact of Atlantic bluefin tuna

migrations in the North Sea. Browman H (ed.). *ICES J Mar Sci* 2017;**74**:1552–60.

Mather FJ, Mason JM, Jones AC. *Life History and Fisheries of Atlantic Bluefin Tuna*. NMFS

(National Marine Fisheries Service); SEFSC (Southeast Fisheries Science Center),

1995:145–65.

McGowan JA, Cayan DR, Dorman LM. Climate-Ocean Variability and Ecosystem Response in

the Northeast Pacific. *Science* 1998;**281**:210–7.

Mehvar S, Filatova T, Dastgheib A *et al.* Quantifying Economic Value of Coastal Ecosystem

Services: A Review. *J Mar Sci Eng* 2018;**6**:5.

Michel RH, McGovern PE. The Chemical Processing of Royal Purple Dye: Addendum.

Archaeomaterials. 1987:93.

Millot C, Taupier-Letage I. Circulation in the Mediterranean Sea. In: Saliot A (ed.), *The*

Mediterranean Sea. Vol 5K. Berlin, Heidelberg: Springer Berlin Heidelberg, 2005, 29–

66.

Monk CT, Bekkevold D, Klefoth T *et al.* The battle between harvest and natural selection

creates small and shy fish. *Proc Natl Acad Sci* 2021;**118**:e2009451118.

Muhling BA, Tommasi D, Ohshimo S *et al.* Regional-scale surface temperature variability

allows prediction of Pacific bluefin tuna recruitment. Hidalgo M (ed.). *ICES J Mar Sci*

2018;**75**:1341–52.

Mylona D. Fish Processing in the Mediterranean: Varying Traditions, Technologies and Scales

of Production with Particular Reference to the Eastern Mediterranean. *J Marit*

Archaeol 2018;**13**:419–36.

Mylona D. Catching tuna in the Aegean: biological background of tuna fisheries and the

archaeological   implications.   *Anthropozoologica*   2021;**56**,   DOI:

10.5252/anthropozoologica2021v56a2.

O'Connor MI. Warming strengthens an herbivore–plant interaction. *Ecology* 2009;**90**:388–

98.

Olesen JM, Jordano P. GEOGRAPHIC PATTERNS IN PLANT–POLLINATOR MUTUALISTIC

NETWORKS. *Ecology* 2002;**83**:2416–24.

Oliver AV. An ancient fishery of Banded dye-murex (Hexaplex trunculus): zooarchaeological

evidence from the Roman city of Pollentia (Mallorca, Western Mediterranean). *J*

Archaeol Sci 2015;**54**:1–7.

Ouled-Cheikh J, Coll M, Cardona L *et al.* Fisheries-enhanced pressure on Mediterranean

regions and pelagic species already impacted by climate change. *Elem Sci Anthr*

2022;**10**:00028.

Page MJ, Moher D, Bossuyt PM *et al.* PRISMA 2020 explanation and elaboration: updated

guidance and exemplars for reporting systematic reviews. *BMJ* 2021:n160.

Pardo M, Fernández-Álvarez FÁ, Varela JL *et al.* Long-term trophic ecology of juvenile

Atlantic bluefin tuna in the western mediterranean sea. *Estuar Coast Shelf Sci*

2025;**325**:109471.

Pauly D, Christensen V, Dalsgaard J *et al.* Fishing Down Marine Food Webs. *Science*

1998;**279**:860–3.

Pedergnana A, Cristiani E, Munro N *et al.* Early line and hook fishing at the Epipaleolithic site

of Jordan River Dureijat (Northern Israel). Peresani M (ed.). *PLOS ONE*

2021;**16**:e0257710.

Pepin P. Effect of Temperature and Size on Development, Mortality, and Survival Rates of

the Pelagic Early Life History Stages of Marine Fish. *Can J Fish Aquat Sci* 1991;**48**:503–

18.

Planque B, Fromentin J-M, Cury P *et al.* How does fishing alter marine populations and

ecosystems sensitivity to climate? *J Mar Syst* 2010;**79**:403–17.

Pohlert T. trend: Non-Parametric Trend Tests and Change-Point Detection. 2015, DOI:

10.32614/cran.package.trend.

Polanco-Martínez JM, Caballero-Alfonso ÁM, Ganzedo U *et al.* A reconstructed database of

historic bluefin tuna captures in the Gibraltar Strait and Western Mediterranean.

Data Brief 2018;**16**:206–10.

Porz A, Zuschin M, Strotz L *et al.* Controls on long-term changes in bathyal bivalve biomass:

The Pleistocene glacial–interglacial record in the eastern Mediterranean. *Deep Sea*

Res Part Oceanogr Res Pap 2024;**203**:104224.

Pourmand A, Marcantonio F, Bianchi TS *et al.* A 28-ka history of sea surface temperature,

primary productivity and planktonic community variability in the western Arabian

Sea. *Paleoceanography* 2007;**22**:2007PA001502.

Prendergast AL, Stevens RE, O'Connell TC *et al.* Changing patterns of eastern Mediterranean

shellfish exploitation in the Late Glacial and Early Holocene: Oxygen isotope evidence

from gastropod in Epipaleolithic to Neolithic human occupation layers at the Haua

Fteah cave, Libya. *Quat Int* 2016;**407**:80–93.

R Core Team. R: A Language and Environment for Statistical Computing. 2025.

Ravier C. Long-term fluctuations in the eastern Atlantic and Mediterranean bluefin tuna

population. *ICES J Mar Sci* 2001;**58**:1299–317.

Ravier C, Fromentin J. Are the long-term fluctuations in Atlantic bluefin tuna (*Thunnus*

thynnus) population related to environmental changes? *Fish Oceanogr* 2004;**13**:145–

60.

Reglero P, Ortega A, Balbín R *et al.* Atlantic bluefin tuna spawn at suboptimal temperatures

for their offspring. *Proc R Soc B Biol Sci* 2018;**285**:20171405.

Reijnders PJH, Verriopoulos G, Brasseur SMJM. *Status of Pinnipeds Relevant to the European*

Union. Wageningen, The Netherlands: IBN-DLO, 1997:195.

Rio M-H, Poulain P-M, Pascual A *et al.* A Mean Dynamic Topography of the Mediterranean Sea computed from altimetric data, in-situ measurements and a general circulation model. *J Mar Syst* 2007;**65**:484–508.

Roberts N, Brayshaw D, Kuzucuoğlu C *et al.* The mid-Holocene climatic transition in the Mediterranean: Causes and consequences. *The Holocene* 2011;**21**:3–13.

Robinson AR, Golnaraghi M. The Physical and Dynamical Oceanography of the Mediterranean Sea. In: Malanotte-Rizzoli P, Robinson AR (eds.), *Ocean Processes in Climate Dynamics: Global and Mediterranean Examples*. Dordrecht: Springer Netherlands, 1994, 255–306.

Robinson AR, Leslie WG, Theocharis A *et al.* Mediterranean Sea circulation. *Ocean Currents*. Academic Press. 2009, 283–98.

Rohling EJ, Cane TR, Cooke S *et al.* African monsoon variability during the previous interglacial maximum. *Earth Planet Sci Lett* 2002;**202**:61–75.

Rooker JR, Alvarado Bremer JR, Block BA *et al.* Life History and Stock Structure of Atlantic Bluefin Tuna (*Thunnus thynnus*). *Rev Fish Sci* 2007;**15**:265–310.

Ruas M-P. Aspects of early medieval farming from sites in Mediterranean France. *Veg Hist Archaeobotany* 2005;**14**:400–15.

Sáez Romero AM. Fish processing and salted-fish trade in the Punic West: New archaeological data and historical evolution. In: Botte E, Leitch V (eds.), *Fish & Ships*. Aix-en-Provence: Publications du Centre Camille Jullian, 2014, 159–74.

Salat J, Pascual J, Flexas M *et al.* Forty-five years of oceanographic and meteorological observations at a coastal station in the NW Mediterranean: a ground truth for satellite observations. *Ocean Dyn* 2019;**69**:1067–84.

663   Santamaria N, Bello G, Corriero A *et al.* Age and growth of Atlantic bluefin tuna, *Thunnus*

664   *thynnus* (Osteichthyes: Thunnidae), in the Mediterranean Sea. *J Appl Ichthyol*

665   2009;**25**:38–45.

666   Sarmiento FM. *De Los Atunes y Sus Transmigraciones y Conjeturas Sobre La Decadencia de*

667   *Las Almadrabas; y Sobre Los Medios Para Restituirlas*., 1757.

668   Scheffer M, Carpenter S, Young B. Cascading effects of overfishing marine systems. *Trends*

669   *Ecol Evol* 2005;**20**:579–81.

670   Schmiedl G, Kuhnt T, Ehrmann W *et al.* Climatic forcing of eastern Mediterranean deep-

671   water formation and benthic ecosystems during the past 22 000 years. *Quat Sci Rev*

672   2010;**29**:3006–20.

673   Shackell NL, Frank KT, Fisher JAD *et al.* Decline in top predator body size and changing

674   climate alter trophic structure in an oceanic ecosystem. *Proc R Soc B Biol Sci*

675   2010;**277**:1353–60.

676   Smale DA, Wernberg T, Oliver ECJ *et al.* Marine heatwaves threaten global biodiversity and

677   the provision of ecosystem services. *Nat Clim Change* 2019;**9**:306–12.

678   Stergiou KI, Tsikliras AC, Pauly D. Farming up Mediterranean Food Webs. *Conserv Biol*

679   2009;**23**:230–2.

680   Stuiver M, Reimer PJ. A Computer Program for Radiocarbon Age Calibration. *Radiocarbon*

681   1986;**28**:1022–30.

682   Trisos CH, Merow C, Pigot AL. The projected timing of abrupt ecological disruption from

683   climate change. *Nature* 2020;**580**:496–501.

684   Trueman CN, Artetxe-Arrate I, Kerr LA *et al.* Thermal sensitivity of field metabolic rate

685   predicts differential futures for bluefin tuna juveniles across the Atlantic Ocean. *Nat*

686   *Commun* 2023;**14**:7379.

Valla FR. Sedentism, the "point of no return", and the Natufian issue. An historical
perspective. *Paléorient* 2018;**44**:19–34.

Van Neer W, Lernau O, Friedman R *et al.* Fish remains from archaeological sites as indicators
of former trade connections in the Eastern Mediterranean. *Paléorient* 2004;**30**:101–
47.

Zohar I, Artzy M. The role of preserved fish: Evidence of fish exploitation, processing and
long-term preservation in the Eastern Mediterranean during the Late Bronze Age
(14th–13th Century BCE). *J Archaeol Sci Rep* 2019;**23**:900–9.

Zorita E, González-Rouco JF, Von Storch H *et al.* Natural and anthropogenic modes of surface
temperature variations in the last thousand years. *Geophys Res Lett*
2005;**32**:2004GL021563.

Supplementary material

S1. List of environmental and cultural events

Table S1 – List of environmental and cultural events, with corresponding names, time intervals and references.

Type of Event	Name	Interval (years BP)	Reference
Climatic	Last Glacial Maximum (LGM)	32,000 - 21,000	Lionello *et al.* (2023)
	Younger Dryas	14,749 - 13,646	Benjamin *et al.* (2017)
	Holocene 1st cooling event (Event 1)	10,500 - 9,800	Lionello *et al.* (2023)
	Rapid Climate Change (RCC)	8,499 - 7,999	Krauß *et al.* (2018)
	Holocene 2nd cooling event (Event 2)	8,000 - 7,000	Lionello *et al.* (2023)
	Holocene 3rd cooling event (Event 3)	5,000 - 4,800	Lionello *et al.* (2023)
	Medieval Warm Period (MWP)	1,050 - 650	Cisneros *et al.* (2016)
	Little Ice Age (LIA)	650 - 100	Cisneros *et al.* (2016)
Cultural	Late Paleolithic	51,949 - 21,949	Stiner and Kuhn (2006)
	Mesolithic	21,949 - 11,949	Stiner and Kuhn (2006)
	Epipaleolithic	11,949 - 8,949	Tortosa (1995); Sánchez (2007)
	Neolithic (Eastern Mediterranean)	8,649 - 5,049	Andreou *et al.* (1996)
	Neolithic (Western Mediterranean)	7,000 - 5,000	Stuiver and Reimer (1986)
	Bronze Age (Eastern Mediterranean)	5,049 - 2,999	Andreou *et al.* (1996)
	Chalcolithic-Bronze (Western Mediterranean)	5,000 - 3,000	Stuiver and Reimer (1986)

Iron Age	3,000 - 2,200	Stuiver and Reimer (1986)
Hellenistic Period	2,349 - 1,950	Morand (2020)
Roman Period (Western Mediterranean)	2,249 - 1,400	Roberts *et al.* (2011)
Roman Period (Eastern Mediterranean)	1,950 - 1,350	Morand (2020)
Byzantine Period	1,626 - 497	Athanassopoulos (2004)
Dark Age	1,400 - 1,090	Roberts *et al.* (2011)

S2. Results from statistical analyses

Table S2.1 – Results from the statistical analyses performed regarding NISP data for the Atlantic bluefin tuna, the gilthead sea bream and the turbinate monodont by sub-regions of the Mediterranean Sea.

Tests	Shapiro-Wilk test		Levene's test		Kruskal-Wallis		Mann-Whitney U
Species/ Sub-region	Western	Eastern	Western	Eastern	Western	Eastern	Western
Atlantic bluefin tuna	$p = 2.2e^{-16}$	$p = 4.917e^{-15}$	$p = 0.917$	$p = 4.99e^{-15}$	$H_2 = 0.0504$ $p = 0.9751$	$H_2 = 8.1208$ $p = 001724$	
Gilthead sea bream	$p = 2.2e^{-16}$	$p = 2.2e^{-16}$	$p = 0.5199$	$p = 0.9085$		$H_2 = 1.636$ $p = 0.4413$	$W = 108.5$ $p = 0.866$
Turbinate monodont	$p = 9.22e^{-6}$	$p = 6.454e^{-5}$	$p = 0.6171$	$p = 0.6171$	$H_3 = 4.5617$ $p = 0.2069$	$H_2 = 0.0362$ $p = 0.982$	

Table S2.2 – Results from the statistical analyses performed regarding MNI data for the banded-dye murex for the Western Mediterranean. Numbers in bold represent statistical differences.

Tests	Shapiro-Wilk test		Levene's test		Kruskal-Wallis	
Species/ Sub-region	Western	Eastern	Western	Eastern	Western	Eastern
Banded-dye murex	$W = 0.480$ $p = 1.349e^{-7}$		$F = 4.270$ $p = 0.03$		$H_2 = 15.792$ **$p = 0.0004$**	
Turbinate monodont	$p = 2.247e^{-7}$	$p = 9.57e^{-5}$	$p = 0.5961$	$p = 0.3724$	$H_3 = 0.8223$ $p = 0.8441$	$H_2 = 1.667$ $p = 0.4346$

Table S2.3 – Results from the statistical analyses performed regarding catch data for the Atlantic bluefin tuna for the Western Mediterranean, for periods divided using the Jenks natural breaks classification method (*classInt* package; Bivand 2023).

Tests	Shapiro-Wilk test	Levene's test	Kruskal-Wallis test
Atlantic bluefin tuna	$W = 0.91514$ $p = 3.12e^{-12}$	$F = 33.624$ $p < 2.2e^{-16}$	$H_5 = 407.55$ **$p < 2.2e^{-16}$**

Table S2.4 – Results from the statistical analyses performed regarding total length (TL) data for the Atlantic bluefin tuna and the gilthead sea bream by sub-regions of the Mediterranean Sea.

Tests	Shapiro-Wilk test		Levene's test		Two-sample t-test	Mann-Whitney U test	one-way ANOVA
Species/ Sub-region	Western	Eastern	Western	Eastern	Western	Eastern	Eastern
Atlantic bluefin tuna	W = 0.9593 p = 0.1686	W = 0.9091 p = 0.0163	F = 2.103 p = 0.1554	F = 0.9167 p = 0.3468	t = 2.967 **p = 0.01**	W = 8 p = **0.01035**	
Gilthead sea bream		W = 0.9898 p = 0.4921		F = 1.5013 p = 0.227			F = 10.28 **p = 7.54e^{-05}**

Table S2.5 – Results from the statistical analyses performed regarding basal diameter (BD) data (mean and maximum values) for the Turbinate monodont for the Eastern Mediterranean.

Tests	Shapiro-Wilk test		Levene's test		one-way ANOVA	Welch ANOVA
Variable/ Sub-region	Mean	Maximum	Mean	Maximum	Mean	Maximum
Turbinate monodont	W = 0.9102 p = 0.245	W = 0.9794 p = 0.4367	F = 1.2519 p = 0.3364	F = 4.4836 p = 0.0071	F = 2.021 p = 0.195	F = 0.3701 p = 0.7767

S3. Presence and abundance of the targeted species

Figure S3.1 - Distribution of data for NISP on the Atlantic bluefin tuna in the Western
Mediterranean. (A) Temporal distribution of records, including major climatic events (purple
rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each
geological period. Blue dots represent mean values, and bars indicate standard deviation.
Letters denote statistically significant differences between periods. The number of data
samples (n) for each boxplot is shown above. The plot inside the second figure is a zoom-in
of the data, for better visualisation of the increase in NISP values towards the Late Holocene
(no outliers were excluded).

Figure S3.2 - Distribution of data for NISP on the Atlantic bluefin tuna in the Eastern
Mediterranean. (A) Temporal distribution of records, including major climatic events (purple
rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each
geological period. Blue dots represent mean values, and bars indicate standard deviation.
Letters denote statistically significant differences between periods. The number of data
samples (n) for each boxplot is shown above.

Figure S3.3 - Distribution of data for NISP on the gilthead sea bream in the Western
Mediterranean. (A) Temporal distribution of records, including major climatic events (purple
rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each
geological period. Blue dots represent mean values, and bars indicate standard deviation.
Letters denote statistically significant differences between periods. The number of data
samples (n) for each boxplot is shown above.

Figure S3.4 - Distribution of data for NISP on the gilthead sea bream in the Eastern Mediterranean. (A) Temporal distribution of records, including major climatic events (purple rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each geological period. Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences between periods. The number of data samples (n) for each boxplot is shown above.

Figure S3.5 - Distribution of data for NISP on the turbinate monodont in the Western Mediterranean. (A) Temporal distribution of records, including major climatic events (purple rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each geological period. Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences between periods. The number of data samples (n) for each boxplot is shown above.

Figure S3.6 - Distribution of data for NISP on the turbinate monodont in the Eastern Mediterranean. (A) Temporal distribution of records, including major climatic events (purple rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each geological period. Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences between periods. The number of data samples (n) for each boxplot is shown above.

Figure S3.7 - Distribution of data for MNI on the turbinate monodont in the Western
Mediterranean. (A) Temporal distribution of records, including major climatic events (purple
rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of MNI data for each
geological period. Blue dots represent mean values, and bars indicate standard deviation.
Letters denote statistically significant differences between periods. The number of data
samples (n) for each boxplot is shown above.

Figure S3.8 - Distribution of data for MNI on the turbinate monodont in the Eastern Mediterranean. (A) Temporal distribution of records, including major climatic events (purple rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of MNI data for each geological period. Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences between periods. The number of data samples (n) for each boxplot is shown above.

S4. Catch records

Figure S4 – Catch records of the Atlantic bluefin tuna are divided into six periods, using the
Jenks Natural Breaks Method, between 250–14 years BP. The blue dots, connected by the
blue line, are the mean values for each period. The number of samples (n) included in each
box plot is indicated above. The black dots are outliers.

Table S4 – Pairwise comparisons between intervals (years BP) for the Atlantic bluefin tuna
catch numbers. Numbers in bold represent the periods that differ significantly from each
other.

	[14,54]	(54, 95]	(95,136]	(136,175]	(175,212]
(54, 95]	**< 2e^{-16}**				
(95,136]	1	**< 2e^{-16}**			
(136,175]	1	**< 2e^{-16}**	1		
(175,212]	**< 2e^{-16}**	1	**< 2e^{-16}**	**< 2e^{-16}**	
(212,250]	**6e^{-04}**	**< 2e^{-16}**	**3e^{-04}**	**1.1e^{-05}**	**< 2e^{-16}**

S5. Body size

Figure S5.1 – Distribution of data for TL on the Atlantic bluefin tuna in the Western Mediterranean. (A) Boxplots of TL data for each geological period. Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences between periods. The number of data samples (n) for each boxplot is shown above. (B) Total length differences between the MWP (Medieval Warm Period) and LIA (Little Ice Age).

Figure S5.2 – Distribution of data for TL on the Atlantic bluefin tuna in the Eastern Mediterranean. (A) Boxplots of TL data for each geological period. Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences between periods. The number of data samples (n) for each boxplot is shown above. (B) differences between the before and during the MWP (Medieval Warm Period).

Figure S5.3 – Distribution of data for the TL on the Gilthead sea bream in the Eastern Mediterranean. Differences between the Bronze and Iron Ages and the Byzantine period.

Figure S5.4 - Distribution of data for mean BD on the turbinate monodont in the Eastern
Mediterranean. (A) Temporal distribution of records, including major climatic events (purple
rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of BD data for each
geological period. Blue dots represent mean values, and bars indicate standard deviation.
Letters denote statistically significant differences between periods. The number of data
samples (n) for each boxplot is shown above.

Figure S5.5 - Distribution of data for maximum values of BD on the turbinate monodont in
the Eastern Mediterranean. (A) Temporal distribution of records, including major climatic
events (purple rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of BD
data for each geological period. Blue dots represent mean values, and bars indicate standard
deviation. Letters denote statistically significant differences between periods. The number of
data samples (n) for each boxplot is shown above.

S6. Sampling site and references

Table S6 – Table with information on the coordinates of each sampling site, with corresponding variable from which data were extracted, and the literary reference.

Latitude	Longitude	Reference
40,86	8,62	(Andrews *et al.* 2023)
37,65	12,58	(Andrews *et al.* 2023)
41,01	28,95	(Andrews *et al.* 2023)
38,11	13,37	(Andrews *et al.* 2023)
36,68	15,13	(Andrews *et al.* 2023)
36,118	-5,342	(Andrews *et al.* 2022)
36,762	-3,844	(Andrews *et al.* 2022)
37,422	23,131	(Andrews *et al.* 2022)
37,471	24,43	(Andrews *et al.* 2022)
42,969	16,718	(Andrews *et al.* 2022)
35,694	34,584	(Andrews *et al.* 2022)
43,345	5,271	(Andrews *et al.* 2022)
37,551	22,717	(Andrews *et al.* 2022)
39,347	22,939	(Andrews *et al.* 2022)
39,915	26,151	(Andrews *et al.* 2022)
38,03	22,949	(Andrews *et al.* 2022)
39,957	26,239	(Andrews *et al.* 2022)
35,018	24,76	(Andrews *et al.* 2022)
36,359	25,397	(Andrews *et al.* 2022)
37,126	25,209	(Andrews *et al.* 2022)
40,455	-0,131	(Andrews *et al.* 2022)
37,052	10,058	(Andrews *et al.* 2022)
35,889	-5,321	(Andrews *et al.* 2022)
35,874	8,785	(Andrews *et al.* 2022)
44,612	33,49	(Andrews *et al.* 2022)
36,851	10,327	(Andrews *et al.* 2022)
36,418	25,428	(Andrews *et al.* 2022)
37,175	21,92	(Andrews *et al.* 2022)
45,351	36,469	(Andrews *et al.* 2022)
39,068	8,455	(Andrews *et al.* 2022)
37,52	23,455	(Andrews *et al.* 2022)
38,222	15,241	(Andrews *et al.* 2022)
36,737	-4,109	(Andrews *et al.* 2022)
42,989	10,492	(Andrews *et al.* 2022)
40,171	22,484	(Andrews *et al.* 2022)
36,722	-4,417	(Andrews *et al.* 2022)
40,872	8,629	(Andrews *et al.* 2022)
40,623	8,197	(Andrews *et al.* 2022)
43,085	3,064	(Andrews *et al.* 2022)
41,006	28,952	(Andrews *et al.* 2022)
41,032	28,95	(Andrews *et al.* 2022)
41,033	28,734	(Andrews *et al.* 2022)

37,669	13,648	(Andrews *et al.* 2022)
38,11	13,365	(Andrews *et al.* 2022)
38,109	13,37	(Andrews *et al.* 2022)
37,654	12,591	(Andrews *et al.* 2022)
40,472	8,563	(Andrews *et al.* 2022)
35,889	5,305	(Andrews *et al.* 2022)
40,863	8,624	(Andrews *et al.* 2022)
40,729	8,557	(Andrews *et al.* 2022)
40,72	8,55	(Andrews *et al.* 2022)
43,302	5,367	(Andrews *et al.* 2022)
38,322	14,749	(Andrews *et al.* 2022)
38,243	13,414	(Andrews *et al.* 2022)
34,328	35,531	(Bosch *et al.* 2018)
43,111	16,727	(Branscombe, Bosch and Miracle 2021)
40,12	-1,049	(Fernández-López De Pablo and Gabriel 2016)
35,372	35,937	(Fuller *et al.* 2020)
33,558	35,375	(Fuller *et al.* 2020)
33,053	35,609	(Guy *et al.* 2018)
31,669	34,575	(Guy *et al.* 2018)
31,326	34,649	(Guy *et al.* 2018)
31,69	34,576	(Guy *et al.* 2018)
31,561	34,844	(Guy *et al.* 2018)
32,457	35,498	(Guy *et al.* 2018)
32,661	35,11	(Guy *et al.* 2018)
31,779	34,851	(Guy *et al.* 2018)
31,766	35,221	(Guy *et al.* 2018)
32,543	34,911	(Guy *et al.* 2018)
31,14	34,458	(Guy *et al.* 2018)
30,882	34,634	(Guy *et al.* 2018)
32,852	35,202	(Guy *et al.* 2018)
37,935	12,348	(Lo Presti *et al.* 2019)
38,009	12,329	(Lo Presti *et al.* 2019)
38,006	12,322	(Lo Presti *et al.* 2019)
38,002	12,322	(Lo Presti *et al.* 2019)
38,164	13,305	(Mannino *et al.* 2011)
43,567	3,914	(Mercier *et al.* 2012)
43,438	4,947	(Mion *et al.* 2022)
39,853	3,131	(Oliver 2015)
32,9	22,053	(Prendergast *et al.* 2016)
35,215	3,254	(Sanchez *et al.* 2022)
33,82	10,843	(Trixl *et al.* 2020)
31,776	35,238	(Tütken *et al.* 2020)
31,326	34,649	(Tütken *et al.* 2020)
35,372	35,937	(Linseele *et al.* 2019)
42,97	16,717	(Rainsford, O'Connor and

		Miracle 2014)
		(Chakroun and Zaghbib-Turki 2017)
36,639	10,852	(Hunt *et al.* 2011)
32,9	22,052	(Hunt *et al.* 2011)
38,19	13,355	(Mannino *et al.* 2011)
38,167	13,348	(Mannino *et al.* 2011)
38,935	1,507	(Giner and Ribas 2008)
39,233	8,367	(Addis *et al.* 2008)
39,183	8,3	(Addis *et al.* 2008)
37,5	12,2	(Longo and Clark 2012)
37,93	12,328	(Longo and Clark 2012)
41	8	(Ganzedo *et al.* 2016)
38,7	8,4	(Ganzedo *et al.* 2016)
39,33	8,1	(Ganzedo *et al.* 2016)
39,75	8,1	(Ganzedo *et al.* 2016)
38,5	12,6	(Ganzedo *et al.* 2016)
38	12,1	(Ganzedo *et al.* 2016)
36,684	15,137	(Andrews *et al.* 2022)

S7. Species' number of samples per variable

Table S7 – Information on the number of samples (n) per variable for each species. NISP – Number of Identified Specimens; MNI – Minimum Number of Species; TL – Total Length; BD – Basal Diameter (including the number of samples for mean and maximum measurements).

Species	Variable		n
Atlantic bluefin tuna	NISP		130
	Catch		1225
	TL		68
Gilthead sea bream	NISP		193
	TL		154
Banded-dye murex	MNI		24
Turbinate monodont	NISP		44
	MNI		41
	BD	Mean	14
		Maximum	57

References

Andreou S, Fotiadis M, Kotsakis K. Review of Aegean prehistory V: The Neolithic and Bronze Age of northern Greece. *American Journal of Archaeology* 1996;**100(3)**:537–97.

Athanassopoulos EF. Historical archaeology of medieval Mediterranean landscapes. In: University of Pennsylvania Museum of Archaeology and Anthropology (ed.), *Mediterranean Archaeological Landscapes: Current Issues*. University of Pennsylvania Museum of Archaeology and Anthropology, 2004, 81–98.

Addis P, Dean JM, Pesci P *et al.* Effects of local scale perturbations in the Atlantic bluefin tuna (Thunnus thynnus L.) trap fishery of Sardinia (W. Mediterranean). *Fish Res* 2008;**92**:242–54.

Andrews AJ, Di Natale A, Addis P *et al.* Vertebrae reveal industrial-era increases in Atlantic bluefin tuna catch-at-size and juvenile growth. Andrews AH (ed.). *ICES J Mar Sci* 2023;**80**:836–47.

Andrews AJ, Di Natale A, Bernal-Casasola D *et al.* Exploitation history of Atlantic bluefin tuna in the eastern Atlantic and Mediterranean—insights from ancient bones. Poulsen B (ed.). *ICES J Mar Sci* 2022;**79**:247–62.

Benjamin J, Rovere A, Fontana A *et al.* Late Quaternary sea-level changes and early human societies in the central and eastern Mediterranean Basin: An interdisciplinary review. *Quat Int* 2017;**449**:29–57.

Bosch MD, Mannino MA, Prendergast AL *et al.* Year-round shellfish exploitation in the Levant and implications for Upper Palaeolithic hunter-gatherer subsistence. *J Archaeol Sci Rep* 2018;**21**:1198–214.

Branscombe TL, Bosch MarjoleinD, Miracle PT. Seasonal Shellfishing across the East Adriatic

Mesolithic-Neolithic Transition: Oxygen Isotope Analysis of *Phorcus turbinatus* from

Vela Spila (Croatia). *Environ Archaeol* 2021;**26**:497–510.

Chakroun A, Zaghbib-Turki D. Facies and fauna proxies used to reconstruct the MIS 5 and

MIS 7 coastal environments in eastern Tunisia. *Geol Q* 2017;**61**, DOI:

10.7306/gq.1312.

Cisneros M, Cacho I, Frigola J *et al.* Sea surface temperature variability in the central-western

Mediterranean Sea during the last 2700 years: a multi-proxy and multi-record

approach. *Clim Past* 2016;**12**:849–69.

Fernández-López De Pablo J, Gabriel S. El Collado shell midden and the exploitation patterns

of littoral resources during the Mesolithic in the Eastern Iberian Peninsula. *Quat Int*

2016;**407**:106–17.

Fuller BT, Van Neer W, Linseele V *et al.* Fish $\delta 13C$ and $\delta 15N$ results from two Bronze/Iron

Age sites (Tell Tweini & Sidon) along the Levantine coast. *J Archaeol Sci Rep*

2020;**29**:102066.

Ganzedo U, Polanco-Martínez JM, Caballero-Alfonso ÁM *et al.* Climate effects on historic

bluefin tuna captures in the Gibraltar Strait and Western Mediterranean. *J Mar Syst*

2016;**158**:84–92.

Giner CA, Ribas BC. Methodological aspects of purple dye production on Ibiza: The new site

of Cala Olivera. *Purpureae Vestes*. 2008, 195–208.

Guy S-V, Thomas T, Irit Z *et al.* Tooth oxygen isotopes reveal Late Bronze Age origin of

Mediterranean fish aquaculture and trade. *Sci Rep* 2018;**8**:14086.

Hunt CO, Reynolds TG, El-Rishi HA *et al.* Resource pressure and environmental change on the

North African littoral: Epipalaeolithic to Roman gastropods from Cyrenaica, Libya.

Quat Int 2011;**244**:15–26.

Krauß R, Marinova E, De Brue H *et al.* The rapid spread of early farming from the Aegean into

the Balkans via the Sub-Mediterranean-Aegean Vegetation Zone. *Quat Int*

2018;**496**:24–41.

Linseele V, Marinova E, De Cupere B *et al.* Bronze and Iron Age palaeo-economy in a

changing environment. The bioarchaeology of Tell Tweini, on the northern Levantine

coast. *About Tell Tweini (Syria): Artefacts, Ecofacts and Landscape*. Leuven: Peeters,

2019, 417–617.

Lionello P, Giorgi F, Rohling E *et al.* Mediterranean climate. *Oceanography of the*

Mediterranean Sea. Elsevier, 2023, 41–91.

Lo Presti V, Antonioli F, Palombo MR *et al.* Palaeogeographical evolution of the Egadi Islands

(western Sicily, Italy). Implications for late Pleistocene and early Holocene sea

crossings by humans and other mammals in the western Mediterranean. *Earth-Sci*

Rev 2019;**194**:160–81.

Longo SB, Clark B. The Commodification of Bluefin Tuna: The Historical Transformation of

the Mediterranean Fishery. *J Agrar Change* 2012;**12**:204–26.

Mannino MA, Thomas KD, Leng MJ *et al.* Stuck to the shore? Investigating prehistoric

hunter-gatherer subsistence, mobility and territoriality in a Mediterranean coastal

landscape through isotope analyses on marine mollusc shell carbonates and human

bone collagen. *Quat Int* 2011;**244**:88–104.

Mercier L, Mouillot D, Bruguier O *et al.* Multi-element otolith fingerprints unravel sea–lagoon lifetime migrations of gilthead sea bream Sparus aurata. *Mar Ecol Prog Ser* 2012;**444**:175–94.

Mion L, André T, Mailloux A *et al.* Contribution to Mediterranean medieval dietary studies: Stable carbon and nitrogen isotope data of marine and catadromous fish from Provence (9th–14th CE). *Data Brief* 2022;**41**:108016.

Morand N. The exploitation of molluscs and other invertebrates in Alexandria (Egypt) from the Hellenistic period to Late Antiquity: food, usage, and trade. *Anthropozoologica* 2020;**55**:1.

Oliver AV. An ancient fishery of Banded dye-murex (Hexaplex trunculus): zooarchaeological evidence from the Roman city of Pollentia (Mallorca, Western Mediterranean). *J Archaeol Sci* 2015;**54**:1–7.

Prendergast AL, Stevens RE, O'Connell TC *et al.* Changing patterns of eastern Mediterranean shellfish exploitation in the Late Glacial and Early Holocene: Oxygen isotope evidence from gastropod in Epipaleolithic to Neolithic human occupation layers at the Haua Fteah cave, Libya. *Quat Int* 2016;**407**:80–93.

Rainsford C, O'Connor T, Miracle P. Fishing in the Adriatic at the Mesolithic–Neolithic transition: Evidence from Vela Spila, Croatia. *Environ Archaeol* 2014;**19**:311–20.

Roberts N, Brayshaw D, Kuzucuoğlu C *et al.* The mid-Holocene climatic transition in the Mediterranean: Causes and consequences. *The Holocene* 2011;**21**:3–13.

Sanchez W, Yanes Y, Linstädter J *et al.* Chronology, time averaging, and oxygen isotope composition of harvested marine mollusk assemblages from Ifri Oudadane, northeast Morocco. *Quat Res* 2022;**106**:147–61.

Stiner MC, Kuhn SL. Changes in the 'Connectedness' and Resilience of Paleolithic Societies in

Mediterranean Ecosystems. *Hum Ecol* 2006;**34**:693–712.

Stuiver M, Reimer PJ. A Computer Program for Radiocarbon Age Calibration. *Radiocarbon*

1986;**28**:1022–30.

Trixl S, Ben Tahar S, Ritter S *et al.* Wool sheep and purple snails—Long-term continuity of

animal exploitation in ancient Meninx (Jerba/Tunisia). *Int J Osteoarchaeol*

2020;**30**:811–23.

Tütken T, Weber M, Zohar I *et al.* Strontium and Oxygen Isotope Analyses Reveal Late

Cretaceous Shark Teeth in Iron Age Strata in the Southern Levant. *Front Ecol Evol*

2020;**8**:570032.

Tortosa JEA. El Magdaleniense Mediterráneo: La Cova del Parpalló (Gandia, Valencia). 1995.

1006

1007