Climate and human impacts on key marine fauna of the Mediterranean Sea

during the pre-industrial period

Daniela Leal[1,2*], Konstantina Agiadi[1,3], Marta Coll[1,4], Maria Bas[1]

[1]Institut de Ciències del Mar (ICM-CSIC), Passeig Marítim de la Barceloneta 37-49, 08003

Barcelona, Spain

[2]University of Algarve, Campus de Gambelas, 8005-139 Faro, Portugal

[3]Department of Geology, University of Vienna, Josef-Holaubek-Platz 2, 1090 Vienna, Austria

[4]Ecopath International Initiative (EII), Barcelona, Spain

*Corresponding author: dani.leal2000@gmail.com; +351 96 048 52 94

Keywords: Atlantic bluefin tuna, Banded-dye murex, Climate change, Gilthead sea bream,

Holocene, Overexploitation.

ABSTRACT

Climate changes and human activities have impacted the Mediterranean ecosystem for

thousands of years, but the available quantitative evidence is incomplete, hindering a

detailed assessment of the marine ecosystem's baseline state and resources. To disentangle

these impacts, we investigated changes in the distribution, abundance and body size of commercially important species of marine mammals, fishes, and molluscs over the last 130,000 (130 ka) years until the Industrial Revolution (1850) in three Mediterranean sub-regions: Western, Eastern and Adriatic. We identified and integrated quantitative data available from the literature on two marine mammals, thirteen fish and three molluscs, revealing consistent and significant climatic and anthropogenic impacts on three species: the banded-dye murex, the Atlantic bluefin tuna and the gilthead sea bream. Climate change, particularly temperature changes, was the most important factor influencing the past abundance and body size of Atlantic bluefin tuna, while sub-regional differences in the stable isotopic composition of archaeological skeletal remains reflected shifts in the system's productivity. Human activities, particularly overexploitation, impacted the abundance of both banded-dye murex and Atlantic bluefin tuna, as well as the body size of gilthead sea bream and Atlantic bluefin tuna. Our results highlight the significance of studying the relationships between climate, human activities, and marine resources in this large marine ecosystem before the Industrial Revolution and showcases the relevance of integrating past quantitative data with contemporary data to reconstruct ecological baselines and obtain longer and more comprehensive time series of ecosystem change.

1. INTRODUCTION

Marine ecosystems are among the world's most diverse and socio-economically valuable systems, providing invaluable ecosystem services (Millennium Ecosystem Assessment 2005; Halpern *et al.* 2008). The Mediterranean Sea is a marine ecosystem that has provided resources to human populations for millennia (Madariaga 1964; Mehvar *et al.* 2018) and

sheltered significant biodiversity (Coll *et al.* 2010). Evidence shows shellfish exploitation may

have already started in Mediterranean coastal areas as early as the Lower Paleolithic (Lotze

et al. 2006). For example, processed shells of marine molluscs have been recovered from

Terra Amata (southern France), dating back to ~300 ka (De Lumley 1966). The exploitation of

marine resources expanded over time, eventually evolving into extensive fisheries and trade

networks for marine products as early as the Roman period (Lotze, Coll and Dunne 2011;

Marzano 2013). Today, the Mediterranean Sea harbours 7.5% of the world's marine

diversity, including several emblematic species such as turtles, marine mammals, and fish,

among others (Coll *et al.* 2010), and it is considered the second-largest biodiversity hotspot

globally (IUCN-MED 2009).

Climate change and human activities significantly impact the Mediterranean Sea, leading to

important changes in marine biota (Lotze *et al.* 2006; Coll *et al.* 2010, 2012; Moullec *et al.*

2019). In recent decades, the Mediterranean Sea has experienced warming well exceeding

the global average (Salat *et al.* 2019), which has profoundly affected marine ecosystems

(Smale *et al.* 2019; Trisos, Merow and Pigot 2020), altering species distributions and

abundances (Garrabou *et al.* 2009; Ouled-Cheikh *et al.* 2022). Key human-related threats to

marine ecosystems include habitat loss, degradation, and overexploitation of resources

(Halpern *et al.* 2008; Coll *et al.* 2010, 2012; Micheli *et al.* 2013), resulting in changes in

species productivity, fitness, and distribution (Pauly *et al.* 1998; Howarth *et al.* 2014).

Most ecological research focuses on short-term data from the past century (Jackson 1997;

Dayton *et al.* 1998), whereas geological, archaeological, and historical data provide long-

term information that is essential for fully understanding how marine ecosystems and

species responded to changes over time (Jackson *et al.* 2001; Lotze *et al.* 2006; Fortibuoni et

al. 2010; Agiadi *et al.* 2024). Marine organisms and their remains are used as proxies for

reconstructing paleoclimatic and paleoenvironmental conditions (e.g., Richardson 2001;

Wanamaker *et al.* 2008; Sisma-Ventura *et al.* 2019) and past human subsistence strategies

(e.g., Bosch *et al.* 2018; Branscombe, Bosch and Miracle 2021) in decadal to million-year

timescales. Additionally, recent studies have revealed the high potential of geohistorical data

for understanding changes in marine ecosystem structure and processes in the past, and

how human-induced and environmental changes impacted marine ecosystems over the long

term (e.g., Calkin, Wiles and Barclay 2001; Lotze, Coll and Dunne 2011; Agiadi *et al.* 2024).

The present study aims to understand how climatic changes and human activities impacted

commercially important species in the Mediterranean Sea over the last 130 ka. We use the

information provided by Leal, Agiadi and Bas (2025), who conducted a systematic review of

marine mammals, fishes and molluscs in this region. We hypothesise that climatic changes,

in particular temperature oscillations associated with warm and cold periods of the Earth's

history, affected the targeted marine species across the Mediterranean Sea throughout

time. In addition, human activities affected this marine fauna through periods of extensive

exploitation across the Mediterranean over the last 130ka. Our ultimate goal is to offer

insights to aid in the management and conservation efforts of Mediterranean marine

communities and ecosystems, learning from past evidence.

2. MATERIALS AND METHODS

2.1 Study area and species data collection

The Mediterranean Sea is a semi-enclosed basin extending over 3 million km^2 with a

maximum depth of approximately 5,300 m (Robinson and Golnaraghi 1994). We divided the

study area into three different sub-regions bounded by the straits of Sicily and Otranto: the

Western and Eastern Mediterranean, and the Adriatic Sea (Millot and Taupier-Letage 2005;

Rio *et al.* 2007; Robinson *et al.* 2009). The Marmara Sea and the Black Sea were included in

the Eastern Mediterranean analyses. We used data on eighteen commercially relevant

species of marine mammals, fish and molluscs gathered by Leal, Agiadi and Bas (2025) (Table

1).

Table 1 – Species analysed in this work.

Scientific Name	Common Name	Functional Group
Delphinus delphis Linnaeus, 1758	Common Dolphin	Marine Mammal
Monachus monachus (Hermann, 1779)	Mediterranean monk seal	Marine Mammal
Engraulis encrasicolus (Linnaeus, 1758)	European anchovy	Fish
Sardina pilchardus (Walbaum, 1792)	European pilchard	Fish
Sardinella aurita Valenciennes, 1847	Round sardinella	Fish
Sardinella maderensis (Lowe, 1838)	Madeiran sardinella	Fish
Sparus aurata Linnaeus, 1758	Gilthead sea bream	Fish
Diplodus annularis (Linnaeus, 1758)	Annular sea bream	Fish
Diplodus bellottii (Steindachner, 1882)	Senegal sea bream	Fish
Diplodus cervinus (Lowe, 1838)	Zebra sea bream	Fish
Diplodus levantinus Fricke, Golani & Appelbaum-Golani, 2016	-	Fish
Diplodus puntazzo (Walbaum, 1792)	Sharpsnout sea bream	Fish
Diplodus sargus (Linnaeus, 1758)	White sea bream	Fish
Diplodus vulgaris (Geoffroy Saint-Hilaire, 1817)	Common two-banded sea bream	Fish
Thunnus thynnus (Linnaeus, 1758)	Atlantic bluefin tuna	Fish
Ostrea edulis Linnaeus, 1758	European flat oyster	Mollusc
Hexaplex trunculus (Linnaeus, 1758)	Banded dye-murex	Mollusc
Phorcus turbinatus (Born, 1778)	Turbinate monodont	Mollusc

From the available records, we analysed data on species: 1) presence and abundance, as

reflected in Number of Identified Specimens – NISP (Grayson 1984) and Minimum Number

of Individuals – MNI (Lyman 1994); 2) catches; 3) body size (as reflected in fish length and

molluscs basal diameter), and 4) stable isotopic composition of fish bones (δ^{13}C and δ^{15}N).

The study conducted by Leal, Agiadi and Bas (2025) revealed important available data on

Atlantic bluefin tuna catches from Italian tuna traps spanning between 1700 and 1936 (Addis

et al. 2008; Longo and Clark 2012; Polanco-Martínez *et al.* 2018). Using long-term

fluctuations in catches as a proxy for abundance (Ravier 2001), we analysed these data to

detect temporal and spatial patterns in Atlantic bluefin tuna abundance in the

Mediterranean during the study period. To place these patterns in the paleoclimatic context,

we obtained Sea Surface Temperature (SST) data for the same period (Jalali *et al.* 2016).

Finally, we investigated the relationships between these catch patterns, major human

civilisation events, and advancements in fisheries.

Furthermore, we analysed δ^{13}C and δ^{15}N data extracted from bones of Atlantic bluefin tuna

(Andrews *et al.* 2023b, 2023c) and the gilthead sea bream (Fuller *et al.* 2020). No corrections

were applied to stable isotope values, as data for each species were produced by the same

methodology.

2.2 Data Analyses

Species with the most data over time in each sub-region were selected, and data were

categorised into four geological periods: Late Pleistocene (131,949 - 11,650 years BP; Head

et al. 2021), and Early (11,650 – 8,186 years BP; Stuiver and Reimer 1986)), Middle (8,186 –

4,200 years BP; Stuiver and Reimer 1986)) and Late Holocene (4,200 – 0 years BP; Stuiver

and Reimer 1986)). The temporal coverage of each climatic and cultural event is displayed in

Table S1. Normality (Shapiro-Wilk test) and homoscedasticity (Levene's test, *car* package;

Fox and Weisberg 2019) were assessed for all variables. For NISP, MNI, body size, and stable

isotopic compositions, non-parametric tests (data non-normally distributed) were used:

Mann-Whitney *U* for two-group comparisons and Kruskal-Wallis for more than two-group

comparisons. The post-hoc pairwise Wilcox test was subsequently used. For the gilthead sea

bream (body size and isotopic information) and turbinate monodont's body size, parametric

tests (ANOVA or Welch ANOVA) were conducted using the post-hoc Tukey's HSD tests. For

catch data, monotonic trends were identified using the Mann-Kendall test (*trend* package;

Pohlert 2023). Data were divided into intervals using the Jenks natural breaks classification

method (Jenks 1977) (*classInt* package in R; Bivand 2023), and the detected differences

between periods were assessed using Kruskal-Wallis and the pairwise Wilcox test. A *p-value*

threshold of ≤0.05 was applied when comparing two groups, with the threshold adjusted

using the Bonferroni correction. Maps were created using QGIS 3.34.3 (QGIS.org) with

bathymetry data from GEBCO (GEBCO Compilation Group 2022). All statistical analyses were

performed in R version 4.4.1 (R Core Team 2023).

137   3. RESULTS

138   The literature review did not provide results for the Senegal seabream, Zebra seabream,

139   *Diplodus levantinus*, and the Sharpsnout seabream. Most data came from the Western and

140   Eastern Mediterranean, particularly during the Holocene period (11,650-0 years BP; Stuiver

141   and Reimer 1986). In the present work, we presented the statistically significant results of

142   species analyses, for more detailed information, see supplementary material S2 to S6.

3.1 Presence and abundance of targeted species

The NISP data was found for most species, except the annular sea bream, European pilchard, round sardinella, and Madeiran sardinella. Fish presented the most data, followed by molluscs and marine mammals. The data among functional groups were distributed between 100,000–500 years BP, with most samples coming from the Western and Eastern Mediterranean, and few samples from the Adriatic Sea. The analyses performed revealed no significant differences between periods (supplementary material S3).

The MNI data, primarily distributed between 100,000–17,000 years BP, were available for *Diplodus spp.,* banded-dye murex, Mediterranean monk seal, European flat oyster, turbinate monodont and the gilthead sea bream (Figure S3.4). The mollusc turbinate monodont and the banded-dye murex had the most data across periods. Data for the turbinate monodont were available for both the Western and the Eastern Mediterranean, and no significant differences were detected between periods (Figures S3.4.1B and S3.4.2B) in both areas. For the banded-dye murex, significant differences were observed between periods in the Western Mediterranean ($H_2 = 15.792$; $p = 0.0004$, Figure 1A), particularly between the Early and Late Holocene ($p = 0.0013$; Figure 1B, Table S2.1).

Figure 1 - Distribution of MNI (Minimum Number of Individuals) data for banded-dye murex in the
Western Mediterranean. (A) Temporal distribution of records, major climatic events (purple
rectangles) and human cultural periods (delimited using black dashed lines); (B) Boxplots of MNI data
for each geological period: EH (Early Holocene), MH (Middle Holocene), and LH (Late Holocene). Blue
dots represent mean values and bars indicate standard deviation. Letters denote statistically
significant differences between periods. The number of data samples (n) for each boxplot is shown
above.

3.2 Catch records

Catch data were available only for the Atlantic bluefin tuna, ranging from zero to
approximately 12,500 individuals caught between 250 and 14 years BP (1700-1936 AD;
Figure 2). Two peaks were observed: the first (~10,000 individuals) occurred around 200
years BP (~1750 AD) during the Little Ice Age, and the second (~12,000 individuals) between
75–45 years BP (1875–1905 AD), after the onset of industrial fisheries. Declines in catches

were evident ($S = -3.98 \, e^5$; Kendall's tau coefficient = $-2.39 \, e^{-1}$; $p < 2.2 \, e^{-16}$), with significant

differences between some of the periods ($H_5 = 407.55$; $p < 2.2e^{-16}$; Table S2.2 - S2.2.1, Figure

S4).

Figure 2 – Atlantic Bluefin Tuna catch records between 250 and 14 years BP (1700–1936 AD), from

tuna traps in the Western Mediterranean Sea. The black points refer to data collected between 250

and 100 years BP (1700–1850 AD), and the light grey points indicate the catch records between 99

and 14 years BP (1851–1936 AD). The red points indicate reconstructed SST records from the Gulf of

Lions (France) (data from Jalali *et al.* 2016). The light blue rectangle refers to the LIA (Little Ice Age)

period and the light grey rectangle refers to the onset of Industrial fisheries, with minor

modifications made to tuna traps until the 1960s (Doumenge 1999), including WWI (World War I).

3. 3 Body size measures

We analysed Total Length (TL) data due to the limited availability of Standard Length (SL)

measurements (Figures S5.1 and S5.2). The TL data were available for Atlantic bluefin tuna

and gilthead sea bream. For the Atlantic bluefin tuna in the Western Mediterranean, we

found significant differences between the Medieval Warm Period and the Little Ice Age

($t_{15.141}$ = 2.967; p = 0.01; Figure 3A, Table S2.3 and Figure S5.3). In the Eastern

Mediterranean, significant differences in Atlantic bluefin tuna TL were observed between

the period before (2,500–1,050 years BP) and during the Medieval Warm Period (1,050–650

years BP; W = 8; p = 0.01035) despite differences in sample sizes (Figure 3B, Table S2.3 and

Figure S5.4).

Figure 3 - Distribution of data for the TL (Total Length) of Atlantic bluefin tuna in the Western and

Eastern Mediterranean. Temporal distribution of records, including major climatic events (purple

rectangles) and anthropogenic events (black dashed lines); (A) Temporal distribution of records for

the Western Mediterranean; (B) Temporal distribution of records for the Eastern Mediterranean.

The TL data for gilthead sea bream were limited to the Eastern Mediterranean (Figure 4).

Pairwise comparisons revealed significant differences in sea bream TL between the Early and

Late Holocene (p = 0.0061), the Middle and Late Holocene (p = 0.001; Figure 4B), and

between the Bronze and Iron Ages and the Byzantine Period (H_2 = 8.501; p = 0.014, Table

S2.3 and Figure S5.5).

Figure 4 - Distribution of TL (Total Length) data for gilthead sea bream in the Eastern Mediterranean.

(A) Temporal distribution of records, including major climatic events (purple rectangles) and

anthropogenic events (black dashed lines); (B) Boxplots of TL data for each geological period: EH

(Early Holocene), MH (Middle Holocene), and LH (Late Holocene). Blue dots represent mean values,

and bars indicate standard deviation. Letters denote statistically significant differences between

periods. The number of data samples (n) for each boxplot is shown above.

The Basal diameter (BD) data for the turbinate monodont were available only for the Eastern

Mediterranean, specifically mean and maximum values (Table S2.4 and Figures S5.6 and

S5.7). No significant differences were found between periods for both mean and maximum

BD (F_{mean} = 2.021; p_{mean} = 0.195; F_{max} = 0.3701; p_{max} = 0.7767; Figures S5.6B and S5.7B,

respectively).

3.4 Stable isotopic data

For the Atlantic bluefin tuna significant differences were observed between the Western and

Eastern Mediterranean between 1,050-850 years BP (Medieval Warm Period) for both δ^{13}C

(W = 1, p = 3.927 e^{-9}; Figure 5A, Table S2.5) and δ^{15}N (W = 16, p = 1.2 e^{-7}; Figure 5B, Table

S2.5).

227
Figure 5 - Distribution of δ^{13}C and δ^{15}N data for Atlantic bluefin tuna in the Eastern and Western

Mediterranean for 1,050–850 years BP. (A) Boxplots showing δ^{13}C values for the Eastern and

Western Mediterranean; (B) Boxplots showing δ^{15}N values for the same sub-regions. Blue dots

represent mean values, and bars indicate standard deviation. The number of data samples (n) for

each boxplot is shown above. See Figure S6.1 for complete isotopic data dispersion.

For the gilthead sea bream, a few δ^{13}C and δ^{15}N records were available covering the Bronze

and Iron Ages (Figures S6.2A and S6.3A). Statistical analysis revealed no significant

differences between geological periods for either stable isotope ($H_{\delta^{13}C}$ = 0.70028, $p_{\delta^{13}C}$ =

0.705; $F_{\delta^{15}N}$ = 2.344, $p_{\delta^{15}N}$ = 0.177, Table S2.5). However, δ^{15}N increased sharply from 4,199 to

3,625 years BP, followed by a decrease towards 2,675 years BP (Figure S5.3B).

4. DISCUSSION

Climate change and human activities have significantly impacted the Mediterranean Sea

(Lotze *et al.* 2006; Coll *et al.* 2010; Moullec *et al.* 2019), highlighting the need to understand

species' responses to these pressures in the past (Jackson *et al.* 2001). In this study, we

compiled new information pointing to the importance of studying the impact of pre-

industrial climate change and human activities on key marine fauna. Available data was

found for banded-dye murex, Atlantic bluefin tuna, and gilthead sea bream, of the

Mediterranean Sea. These species have been important for the Mediterranean ecosystem

structure and functioning and crucial for sustaining the socioeconomic systems of the peri-

Mediterranean countries for thousands of years.

Seawater temperature influences Atlantic bluefin tuna's growth, reproduction, mortality,

and migration patterns (Pepin 1991; Graham and Dickson 2004). This highly migratory

species moves between cold feeding waters like the North Atlantic Ocean (Mather *et al.*

1995; Block *et al.* 2001, 2005) and warm spawning areas such as the Mediterranean Sea

(Mather *et al.* 1995; Block *et al.* 2005; Rooker *et al.* 2006). The high number of catches in the

Western Mediterranean during the latter stages of the Little Ice Age (650–100 years BP;

Briffa 2000) may be linked to a decline in the SST. During this period, there may have been

changes in the species' migration patterns due to cooler and unstable meteorological

conditions (strong storm activity and cold, windy conditions, etc.), reducing the efficiency of

tuna fisheries (Ravier and Fromentin 2004; Ganzedo *et al.* 2016). Temperature also affects

plankton production and distribution (e.g., McGowan, Cayan and Dorman 1998; Beaugrand

et al. 2002) and thus food availability for both juvenile and adult tuna (Mather *et al.* 1995),

driving the spatiotemporal dynamics of tuna movements between the spawning and feeding

grounds (Ravier and Fromentin 2004).

Pleistocene-Holocene climatic variability may have also impacted the body size of tuna, as

seen for other species (Agiadi *et al.* 2023; Porz *et al.* 2024). Recent studies suggest stronger

consumer-driven control of food webs with warming (Allen, Gillooly and Brown 2005; López-

Urrutia *et al.* 2006), which enhances heterotroph abundance, strengthening consumer-

resource interactions (O'Connor 2009). This can benefit high trophic-level species (Archibald

et al. 2022), such as Atlantic bluefin tuna. Species from temperate ecosystems generally

have broader niche breadths, making them more adaptive (MacArthur 1972; Olesen and

Jordano 2002). The TL increase observed in this study may have been caused by this dietary

flexibility, combined with stronger consumer-resource interactions, allowing tuna to thrive

during past climatic fluctuations. This hypothesis is consistent with the study by Andrews *et*

al. (2023a), who observed increased juvenile growth between the 16[th] and 18[th] centuries in

the Mediterranean Sea, likely linked to warmer sea surface temperature and a positive

North Atlantic Oscillation phase.

The Mediterranean Sea is broadly oligotrophic with a persistent east-to-west gradient in

productivity (Coll *et al.* 2010). Historical data suggest reduced primary productivity in the

Eastern Mediterranean (e.g., Kuhnt *et al.* 2008) and higher productivity in the Western

Mediterranean (e.g., Incarbona *et al.* 2010). These productivity differences likely explain the

δ^{13}C and δ^{15}N variations of Atlantic bluefin tuna's bones from 1,050 to 850 years BP. The

δ^{13}C values are good indicators of primary production sources (Hobson, Piatt and Pitocchelli

1994; France 1995). Tissues presenting low-δ^{13}C values can be associated with species

inhabiting environments dominated by pelagic primary producers or enriched by terrestrial

organic matter over benthic carbon sources (Barnes and Jennings, Jon. T. Barry 2009;

Magozzi *et al.* 2017). The higher δ^{13}C values in the Western Mediterranean suggest that this

species may have used coastal waters in this area, while it might have used offshore areas in

the Eastern Mediterranean. This aligns with the use of Western waters for spawning (Oray

and Karakulak 2005; Alemany *et al.* 2010) and Eastern waters for offshore foraging,

potentially linked to opportunities for deep diving (Wilson and Block 2009; Olafsdottir *et al.*

2016) and/or reduced benthic-pelagic coupling (Magozzi *et al.* 2017). Similarly, δ^{15}N values,

which are used to estimate the trophic position in the food web (Sigman *et al.* 2009), were

higher in the Western Mediterranean, likely reflecting greater secondary productivity.

Despite its oligotrophy (Coll *et al.* 2010), Mediterranean coastal areas experience nutrient

enrichment through local factors such as wind conditions, currents, and fluvial discharges

(Zavatarelli *et al.* 1998; Bosc, Bricaud and Antoine 2004). These localised nutrient inputs,

combined with the generalist feeding strategy of Atlantic bluefin tuna (MacArthur 1972;

Olesen and Jordano 2002), may have facilitated its success in the West, while in the East the

more oligotrophic conditions may have led to more specialised diets. Nevertheless, a

previous study using long-term records from eastern Atlantic, Mediterranean and Black Sea

locations found no significant shifts in the trophic position of this species over the past

millennia despite extensive modifications in marine ecosystems (Andrews *et al.* 2023c).

This study also highlights the impact of sustained exploitation on the abundance and body

sizes of different species. The banded-dye murex was extensively harvested during the Late

Holocene (4,200 years BP to the present day; Stuiver and Reimer 1986), particularly in the

Roman Period (2,249-1,400 years BP; Roberts *et al.* 2011). This gastropod inhabits the

intertidal zone (Bonanno *et al.* 2016), making it accessible to hunter-gatherers (e.g.,

Fernández-López De Pablo and Gabriel 2016) and later to civilisations that harvested it at an

industrial scale (Michel and McGovern 1987; Oliver 2015). The Romans used this species,

along with *Bolinus brandaris* and *Stramonita haemastoma* to produce Purple Dye,

highlighting its economic significance (Oliver 2015).

The Atlantic bluefin tuna abundance seems to have also been influenced by industrial

fisheries (Polanco-Martínez *et al.* 2018; this study). The most common method of capture

was tuna traps (*tonnare* in Italian), which caught the fish near the shore where they

spawned (Longo 2012), making it a sustainable fishery (De la Serna *et al.* 2012). With

industrial fishing, *tonnare* were slowly modified with more robust materials (e.g., iron

anchors, steel cables, etc.; Lentini 2001) and later replaced by modernised fishing gears, like

purse-seiners and longlines (Joseph 2003). The 19[th] century, which roughly coincides with

the increasing use of modernised gears, was marked by the starting decades of intense

exploitation of tuna (Andrews *et al.* 2022). The high number of catches after the 19[th] century

might reflect this slow transformation from a low-scale, sustainable fishery into a large-scale,

unsustainable fishery. Generally, removing top predators may lead to changes in the food

web structure through top-down control (Pauly *et al.* 1998; Scheffer, Carpenter and Young

2005), although the ecosystem impacts of these removals remain understudied. Similar

processes are documented for sea otters, heavily hunted by Aboriginal Aleuts 2,500 years

ago, that led to their population decline and an increase in the sea urchin population,

resulting in deep changes to kelp forest ecosystems (Simenstad, Estes and Kenyon 1978;

Jackson *et al.* 2001).

Our findings indicate that the overexploitation of Atlantic bluefin tuna and gilthead sea

bream may have resulted in body size changes. Historical records show tuna catch declined

in Spain between mid-16th and 18th centuries (Sarmiento 1757). Overexploitation in previous

centuries of juveniles and seabed disturbances near tuna traps likely contributed to changes

in tuna migration routes (García 2016). Andrews *et al.* (2023a) proposed that the increased

growth in juveniles between the 16th and 21st centuries in the Mediterranean Sea was linked

to fisheries-induced evolution. This process selects species traits (early maturation,

increased reproductive output, and reduced growth post-maturation) that favour faster life

histories through enhanced survival and the number of offspring of species subjugated to

intensive fishing (Law 2000; Enberg *et al.* 2012). In contrast, the gilthead sea bream

experienced overexploitation in the Late Holocene in the Eastern Mediterranean with a body

size decrease (Guy *et al.* 2018; this study). This pattern of body size reduction also occurred

with *Archosargus probatocephalus* due to long-term fishing activities (Guiry *et al.* 2021).

These size reductions, driven by overfishing and size-selective gear, can have profound

effects on trophic interactions within marine ecosystems (Shackell *et al.* 2010; Garcia *et al.*

2012).

It is necessary to be cautious when analysing our results due to some assumptions. The

study is constrained towards economically relevant species, potentially overlooking other

key species. The lack of comprehensive data for the Adriatic Sea suggests gaps in earlier

studies or unpublished data. Variations in data quality and standardisation (dating systems

and body size measures) presented challenges in analysing data. The resolution needed to

reconstruct past population sizes limited the assessment of the magnitude of change.

Despite these limitations, our study provides important insights into how past environmental

and human impacts have shaped species abundance, body size, and trophic ecology of key

species in the Mediterranean Sea.

Conclusions

This work underscores the importance of studying the impact of pre-industrial climate

change and human activities on Mediterranean species, focusing on abundance, body size,

and trophic ecology. We show that the Atlantic bluefin tuna was likely affected by both

factors, while the banded-dye murex and gilthead sea bream were mainly impacted by

human exploitation. Understanding the history of marine ecosystems and species-specific

responses to environmental and anthropogenic disturbances is essential to ensure the

development of effective present and future environmental management strategies while

supporting the implementation of targeted adaptive conservation measures.

ACKNOWLEDGEMENTS

This work was supported by the Erasmus+ Program Grant no. 2022-1-PT01-KA131-HED-

000061155. We acknowledge the institutional support of the Severo Ochoa Centre of

Excellence accreditation (CEX2019-000928-S). This research contributes to the objectives of

Q-MARE (a PAGES working group). This research is part of the Integrated Marine Ecosystem

Assessments (iMARES) research group funded by Agència de Gestió d'Ajuts Universitaris i de

Recerca (Generalitat de Catalunya) Grant no. 2021 SGR 00435.

AUTHOR CONTRIBUTIONS

374 **Daniela Leal:** Conceptualisation, Data Curation, Methodology, Formal analysis, Writing -

375 original draft, Funding acquisition. **Konstantina Agiadi:** Conceptualisation, Methodology,

376 Writing - Review & Editing, Supervision. **Marta Coll:** Writing - Review & Editing. **Maria Bas:**

377 Conceptualisation, Methodology, Writing - Review & Editing, Supervision.

DECLARATION OF INTERESTS

380 The authors declare that they have no known competing financial interests or personal

381 relationships that could have appeared to influence the work reported in this paper.

FUNDING

384 This work was supported by the Erasmus+ Program Grant no. 2022-1-PT01-KA131-HED-

385 000061155.

DATA AVAILABILITY

All data produced for this work have been made available within the main manuscript and in

the supplementary material.

REFERENCES

Addis P, Dean JM, Pesci P *et al.* Effects of local scale perturbations in the Atlantic bluefin

tuna (Thunnus thynnus L.) trap fishery of Sardinia (W. Mediterranean). *Fish Res*

2008;**92**:242–54.

Agiadi K, Caswell BA, Almeida R *et al.* Geohistorical insights into marine functional

connectivity. Selden R (ed.). *ICES J Mar Sci* 2024;**81**:1884–911.

Agiadi K, Quillévéré F, Nawrot R *et al.* Palaeontological evidence for community-level

decrease in mesopelagic fish size during Pleistocene climate warming in the eastern

Mediterranean. *Proc R Soc B Biol Sci* 2023;**290**:20221994.

Alemany F, Quintanilla L, Velez-Belchí P *et al.* Characterisation of the spawning habitat of

Atlantic bluefin tuna and related species in the Balearic Sea (western Mediterranean). *Prog*

Oceanogr 2010;**86**:21–38.

Allen AP, Gillooly JF, Brown JH. Linking the global carbon cycle to individual metabolism.

Funct Ecol 2005;**19**:202–13.

Andrews AJ, Di Natale A, Addis P *et al.* Vertebrae reveal industrial-era increases in Atlantic

bluefin tuna catch-at-size and juvenile growth. Andrews AH (ed.). *ICES J Mar Sci*

2023a;**80**:836–47.

Andrews AJ, Di Natale A, Bernal-Casasola D *et al.* Exploitation history of Atlantic bluefin tuna

in the eastern Atlantic and Mediterranean—insights from ancient bones. Poulsen B (ed.).

ICES J Mar Sci 2022;**79**:247–62.

Andrews AJ, Orton D, Onar V *et al.* Isotopic life-history signatures are retained in modern

and ancient Atlantic bluefin tuna vertebrae. *J Fish Biol* 2023b;**103**:118–29.

Andrews AJ, Pampoulie C, Di Natale A *et al.* Exploitation shifted trophic ecology and habitat

preferences of Mediterranean and Black Sea bluefin tuna over centuries. *Fish Fish*

2023c;**24**:1067–83.

Archibald KM, Dutkiewicz S, Laufkötter C *et al.* Thermal Responses in Global Marine

Planktonic Food Webs Are Mediated by Temperature Effects on Metabolism. *J Geophys Res*

Oceans 2022;**127**:e2022JC018932.

Barnes C, Jennings, Jon. T. Barry S. Environmental correlates of large-scale spatial variation in

the δ13C of marine animals. *Estuar Coast Shelf Sci* 2009;**81**:368–74.

Beaugrand G, Reid PC, Ibañez F *et al.* Reorganization of North Atlantic Marine Copepod

Biodiversity and Climate. *Science* 2002;**296**:1692–4.

Bivand R. *classInt: Choose Univariate Class Intervals*. R package version 0.4-10, 2023.

https://CRAN.R-project.org/package=classInt.

Block BA, Dewar H, Blackwell SB *et al.* Migratory Movements, Depth Preferences, and

Thermal Biology of Atlantic Bluefin Tuna. *Science* 2001;**293**:1310–4.

Block BA, Teo SLH, Walli A *et al.* Electronic tagging and population structure of Atlantic

bluefin tuna. *Nature* 2005;**434**:1121–7.

Bonanno A, Barra M, Basilone G *et al.* Environmental processes driving anchovy and sardine

distribution in a highly variable environment: the role of the coastal structure and riverine

input. *Fish Oceanogr* 2016;**25**:471–90.

Bosc E, Bricaud A, Antoine D. Seasonal and interannual variability in algal biomass and

primary production in the Mediterranean Sea, as derived from 4 years of SeaWiFS

observations. *Glob Biogeochem Cycles* 2004;**18**:2003GB002034.

Bosch MD, Mannino MA, Prendergast AL *et al.* Year-round shellfish exploitation in the Levant

and implications for Upper Palaeolithic hunter-gatherer subsistence. *J Archaeol Sci Rep*

2018;**21**:1198–214.

Branscombe TL, Bosch MarjoleinD, Miracle PT. Seasonal Shellfishing across the East Adriatic

Mesolithic-Neolithic Transition: Oxygen Isotope Analysis of *Phorcus turbinatus* from Vela

Spila (Croatia). *Environ Archaeol* 2021;**26**:497–510.

Briffa KR. Annual climate variability in the Holocene: interpreting the message of ancient

trees. *Quat Sci Rev* 2000;**19**:87–105.

Calkin PE, Wiles GC, Barclay DJ. Holocene coastal glaciation of Alaska. *Quat Sci Rev*

2001;**20**:449–61.

Coll M, Piroddi C, Steenbeek J *et al.* The Biodiversity of the Mediterranean Sea: Estimates,

Patterns, and Threats. Bograd SJ (ed.). *PLoS ONE* 2010;**5**:e11842.

Coll M, Piroddi C, Albouy C *et al.* The Mediterranean Sea under siege: spatial overlap

between marine biodiversity, cumulative threats and marine reserves. *Glob Ecol Biogeogr*

2012;**21**:465–80.

Dayton PK, Tegner MJ, Edwards PB, Riser KL. Sliding baselines, ghosts, and reduced

expectations in kelp forest communities. *Ecological Applications* 1998;**8(2)**:309–22.

De la Serna JM, Macías D, de Urbina JO, Rodríguez-Marin E, Abascal F. Study on the eastern

Atlantic and Mediterranean bluefin tuna stock using the Spanish traps as scientific

observatories. *Collect Vol Sci Pap ICCAT* 2012;**67(1)**:331–43.

De Lumley H. Les fouilles de Terra Amata à Nice. Premiers résultats. *Bulletin du Musée*

d'Anthropologie Préhistorique de Monaco 1966;**13**:29–51.

Doumenge F. La storia delle pesche tonniere. *Biol Mar Medit* 1999;**6(2)**:107–48.

Enberg K, Jørgensen C, Dunlop ES, Varpe Ø, Boukal DS, et al. Fishing-induced evolution of

growth: Concepts, mechanisms, and the empirical evidence. *Marine Ecology* 2012;**33(1)**:1–

25.

Fox J, Weisberg S. *An R Companion to Applied Regression*, 3rd ed. Sage, Thousand Oaks, CA,

2019. https://socialsciences.mcmaster.ca/jfox/Books/Companion/.

Fernández-López De Pablo J, Gabriel S. El Collado shell midden and the exploitation patterns

of littoral resources during the Mesolithic in the Eastern Iberian Peninsula. *Quat Int*

2016;**407**:106–17.

Fortibuoni T, Libralato S, Raicevich S *et al.* Coding Early Naturalists' Accounts into Long-Term

Fish Community Changes in the Adriatic Sea (1800–2000). Thrush S (ed.). *PLoS ONE*

2010;**5**:e15502.

France RL. Differentiation between littoral and pelagic food webs in lakes using stable

carbon isotopes. *Limnol Oceanogr* 1995;**40**:1310–3.

Fuller BT, Van Neer W, Linseele V *et al.* Fish δ13C and δ15N results from two Bronze/Iron

Age sites (Tell Tweini & Sidon) along the Levantine coast. *J Archaeol Sci Rep* 2020;**29**:102066.

Ganzedo U, Polanco-Martínez JM, Caballero-Alfonso ÁM *et al.* Climate effects on historic

bluefin tuna captures in the Gibraltar Strait and Western Mediterranean. *J Mar Syst*

2016;**158**:84–92.

García FDLCG. *Doñana en su historia: Cuatro siglos entre la explotación y la conservación*

bajo la posesión de la Casa de los Guzmanes. Organismo Autónomo Parques Nacionales,

2016.

Garcia SM, Kolding J, Rice J *et al.* Reconsidering the Consequences of Selective Fisheries.

Science 2012;**335**:1045–7.

Garrabou J, Coma R, Bensoussan N *et al.* Mass mortality in Northwestern Mediterranean

rocky benthic communities: effects of the 2003 heat wave. *Glob Change Biol* 2009;**15**:1090–

103.

GEBCO Compilation Group. *GEBCO 2022 Grid* [Data set]. 2022.

https://doi.org/10.5285/a29c5465-b138-234d-e053-6c86abc0df45.

Graham JB, Dickson KA. Tuna comparative physiology. *J Exp Biol* 2004;**207**:4015–24.

Grayson DK. *Quantitative Zooarchaeology: Topics in the Quantification of Archaeofaunas*.

Academic Press, Orlando, 1984.

Guiry EJ, Kennedy JR, O'Connell MT *et al.* Early evidence for historical overfishing in the Gulf

of Mexico. *Sci Adv* 2021;**7**:eabh2525.

Guy S-V, Thomas T, Irit Z *et al.* Tooth oxygen isotopes reveal Late Bronze Age origin of

Mediterranean fish aquaculture and trade. *Sci Rep* 2018;**8**:14086.

Halpern BS, Walbridge S, Selkoe KA *et al.* A Global Map of Human Impact on Marine

Ecosystems. *Science* 2008;**319**:948–52.

Head MJ, Pillans B, Zalasiewicz JA *et al.* Formal ratification of subseries for the Pleistocene

Series of the Quaternary System. *Episodes* 2021;**44**:241–7.

Hobson KA, Piatt JF, Pitocchelli J. Using Stable Isotopes to Determine Seabird Trophic

Relationships. *J Anim Ecol* 1994;**63**:786.

Howarth LM, Roberts CM, Thurstan RH *et al.* The unintended consequences of simplifying

the sea: making the case for complexity. *Fish Fish* 2014;**15**:690–711.

Incarbona A, Ziveri P, Di Stefano E *et al.* The Impact of the Little Ice Age on Coccolithophores

in the Central Mediterranea Sea. *Clim Past* 2010;**6**:795–805.

Jackson JBC. Reefs since Columbus. *Coral Reefs* 1997;**16**:S23–32.

Jackson JBC, Kirby MX, Berger WH *et al.* Historical Overfishing and the Recent Collapse of

Coastal Ecosystems. *Science* 2001;**293**:629–37.

Jalali B, Sicre M-A, Bassetti M-A *et al.* NOAA/WDS Paleoclimatology - Gulf of Lion,

Mediterranean Sea 10,000 Year Biomarker Data and SST Reconstruction. 2016, DOI:

10.25921/6S87-Y952.

Jenks GF. Optimal data classification for choropleth maps. *Department of Geography,*

University of Kansas Occasional Paper, 1977.

Joseph J. *Managing Fishing Capacity of the World Tuna Fleet*. Food and Agriculture Organization, Rome, Italy, 2003.

Kuhnt T, Schmiedl G, Ehrmann W *et al.* Stable isotopic composition of Holocene benthic foraminifers from the Eastern Mediterranean Sea: Past changes in productivity and deep water oxygenation. *Palaeogeogr Palaeoclimatol Palaeoecol* 2008;**268**:106–15.

Law R. Fishing, selection, and phenotypic evolution. *ICES Journal of Marine Science* 2000;**57(3)**:659–68.

Leal D, Agiadi K, Bas M. (2025) 'Commercially Relevant Species in the Mediterranean Sea: A Perspective from Late Pleistocene to the Industrial Revolution', bioRxiv, DOI: 10.1101/2025.02.04.636444, 07-02-2025.

Lentini R. Favignana nella seconda metà dell'800: Innovazioni e mercato. In: Doneddu G, Fiori A (eds.), *La pesca in Italia tra età moderna e contemporanea: Produzione, mercato, consumo*. Editrice Democratica Sarda, Sassari, Italy, 2001, 507–20.

Longo SB. Mediterranean Rift: Socio-Ecological Transformations in the Sicilian Bluefin Tuna Fishery. *Crit Sociol* 2012;**38**:417–36.

Longo SB, Clark B. The Commodification of Bluefin Tuna: The Historical Transformation of the Mediterranean Fishery. *J Agrar Change* 2012;**12**:204–26.

López-Urrutia Á, San Martin E, Harris RP *et al.* Scaling the metabolic balance of the oceans. *Proc Natl Acad Sci* 2006;**103**:8739–44.

Lotze HK, Coll M, Dunne JA. Historical Changes in Marine Resources, Food-web Structure and Ecosystem Functioning in the Adriatic Sea, Mediterranean. *Ecosystems* 2011;**14**:198–222.

Lotze HK, Lenihan HS, Bourque BJ *et al.* Depletion, Degradation, and Recovery Potential of

Estuaries and Coastal Seas. *Science* 2006;**312**:1806–9.

Lyman RL. *Vertebrate Taphonomy*. Cambridge University Press, 1994.

MacArthur RH. *Geographical Ecology: Patterns in the Distribution of Species*. Princeton

University Press, 1972.

Madariaga B. *El mar y el hombre prehistórico*. Universidad De Salamanca, 1964.

http://hdl.handle.net/10366/70977.

Magozzi S, Yool A, Vander Zanden HB *et al.* Using ocean models to predict spatial and

temporal variation in marine carbon isotopes. *Ecosphere* 2017;**8**:e01763.

Marzano A. *Harvesting the Sea: The Exploitation of Marine Resources in the Roman*

Mediterranean. Oxford University Press, 2013.

Mather FJ, Mason JM Jr, Jones A. Historical document: Life history and fisheries of Atlantic

bluefin tuna. *NOAA Technical Memorandum NMFS-SEFSC-370*, Miami, FL, 1995, 165 pp.

McGowan JA, Cayan DR, Dorman LM. Climate-Ocean Variability and Ecosystem Response in

the Northeast Pacific. *Science* 1998;**281**:210–7.

Mehvar S, Filatova T, Dastgheib A *et al.* Quantifying Economic Value of Coastal Ecosystem

Services: A Review. *J Mar Sci Eng* 2018;**6**:5.

Michel RH, McGovern PE. The chemical processing of Royal Purple dye: Ancient descriptions

as elucidated by modern science. *Archeomaterials* 1987;**1(2)**:135–43.

Micheli F, Halpern BS, Walbridge S *et al.* Cumulative Human Impacts on Mediterranean and

Black Sea Marine Ecosystems: Assessing Current Pressures and Opportunities. Meador JP

(ed.). *PLoS ONE* 2013;**8**:e79889.

Millennium Ecosystem Assessment. *Ecosystems and Human Well-Being: Current State and*

Trends. Island Press, 2005.

Millot C, Taupier-Letage I. Circulation in the Mediterranean Sea. In: Saliot A (ed.), *The*

Mediterranean Sea. Vol 5K. Berlin, Heidelberg: Springer Berlin Heidelberg, 2005, 29–66.

Moullec F, Velez L, Verley P *et al.* Capturing the big picture of Mediterranean marine

biodiversity with an end-to-end model of climate and fishing impacts. *Prog Oceanogr*

2019;**178**:102179.

O'Connor MI. Warming strengthens an herbivore–plant interaction. *Ecology* 2009;**90**:388–

98.

Olafsdottir D, MacKenzie BR, Chosson-P V *et al.* Dietary Evidence of Mesopelagic and Pelagic

Foraging by Atlantic Bluefin Tuna (Thunnus thynnus L.) during Autumn Migrations to the

Iceland Basin. *Front Mar Sci* 2016;**3**, DOI: 10.3389/fmars.2016.00108.

Olesen JM, Jordano P. GEOGRAPHIC PATTERNS IN PLANT–POLLINATOR MUTUALISTIC

NETWORKS. *Ecology* 2002;**83**:2416–24.

Oliver AV. An ancient fishery of Banded dye-murex (Hexaplex trunculus): zooarchaeological

evidence from the Roman city of Pollentia (Mallorca, Western Mediterranean). *J Archaeol Sci*

2015;**54**:1–7.

Oray IK, Karakulak FS. Further evidence of spawning of bluefin tuna (Thunnus thynnus L.,

1758) and the tuna species (Auxis rochei Ris., 1810, Euthynnus alletteratus Raf., 1810) in the

eastern Mediterranean Sea: preliminary results of TUNALEV larval survey in 2004. *J Appl*

Ichthyol 2005;**21**:236–40.

Ouled-Cheikh J, Coll M, Cardona L *et al.* Fisheries-enhanced pressure on Mediterranean

regions and pelagic species already impacted by climate change. *Elem Sci Anthr*

2022;**10**:00028.

Pauly D, Christensen V, Dalsgaard J *et al.* Fishing Down Marine Food Webs. *Science*

1998;**279**:860–3.

Pepin P. Effect of Temperature and Size on Development, Mortality, and Survival Rates of

the Pelagic Early Life History Stages of Marine Fish. *Can J Fish Aquat Sci* 1991;**48**:503–18.

Pohlert T. *trend: Non-Parametric Trend Tests and Change-Point Detection*. R package version

1.1.6, 2023. https://CRAN.R-project.org/package=trend.

Polanco-Martínez JM, Caballero-Alfonso ÁM, Ganzedo U *et al.* A reconstructed database of

historic bluefin tuna captures in the Gibraltar Strait and Western Mediterranean. *Data Brief*

2018;**16**:206–10.

Porz A, Zuschin M, Strotz L *et al.* Controls on long-term changes in bathyal bivalve biomass:

The Pleistocene glacial–interglacial record in the eastern Mediterranean. *Deep Sea Res Part*

Oceanogr Res Pap 2024;**203**:104224.

QGIS.org. QGIS Geographic Information System. *QGIS Association*. http://www.qgis.org

(accessed April 2024).

R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for

Statistical Computing, Vienna, Austria, 2023. https://www.R-project.org/.

Ravier C. Long-term fluctuations in the eastern Atlantic and Mediterranean bluefin tuna

population. *ICES J Mar Sci* 2001;**58**:1299–317.

Ravier C, Fromentin J. Are the long-term fluctuations in Atlantic bluefin tuna (*Thunnus*

thynnus) population related to environmental changes? *Fish Oceanogr* 2004;**13**:145–60.

Richardson CA. Molluscs as archives of environmental change. In: *Oceanography and Marine*

Biology. Taylor & Francis Ltd, 2001, 103–64.

Rio M-H, Poulain P-M, Pascual A *et al.* A Mean Dynamic Topography of the Mediterranean

Sea computed from altimetric data, in-situ measurements and a general circulation model. *J*

Mar Syst 2007;**65**:484–508.

Roberts N, Brayshaw D, Kuzucuoğlu C *et al.* The mid-Holocene climatic transition in the

Mediterranean: Causes and consequences. *The Holocene* 2011;**21**:3–13.

Robinson AR, Golnaraghi M. The Physical and Dynamical Oceanography of the

Mediterranean Sea. In: Malanotte-Rizzoli P, Robinson AR (eds.), *Ocean Processes in Climate*

Dynamics: Global and Mediterranean Examples. Dordrecht: Springer Netherlands, 1994,

255–306.

Robinson AR, Leslie WG, Theocharis A, Lascaratos A. Mediterranean Sea circulation. *Ocean*

Currents, 2009, 283–98.

Rooker JR, Secor DH, De Metrio G, Rodríquez-Marín E, Fenech Farrugia A. Evaluation of

population structure and mixing rates of Atlantic bluefin tuna from chemical signatures in

otoliths. *Collect Vol Sci Pap ICCAT* 2006;**59**:813–18.

Salat J, Pascual J, Flexas M *et al.* Forty-five years of oceanographic and meteorological

observations at a coastal station in the NW Mediterranean: a ground truth for satellite

observations. *Ocean Dyn* 2019;**69**:1067–84.

Sarmiento FM. *De los atunes y sus transmigraciones y conjeturas sobre la decadencia de las*

almadrabas y sobre los medios para restituirlas. Caixa de Pontevedra, 1757.

Scheffer M, Carpenter S, Young B. Cascading effects of overfishing marine systems. *Trends*

Ecol Evol 2005;**20**:579–81.

Shackell NL, Frank KT, Fisher JAD *et al.* Decline in top predator body size and changing

climate alter trophic structure in an oceanic ecosystem. *Proc R Soc B Biol Sci* 2010;**277**:1353–

60.

Sigman DM, Karsh KL, Casciotti KL. Nitrogen isotopes in the ocean. In: Steele JH (ed.),

Encyclopedia of Ocean Sciences. Academic Press, 2009, 40–54.

Simenstad CA, Estes JA, Kenyon KW. Aleuts, Sea Otters, and Alternate Stable-State

Communities. *Science* 1978;**200**:403–11.

Sisma-Ventura G, Tütken T, Peters STM *et al.* Past aquatic environments in the Levant

inferred from stable isotope compositions of carbonate and phosphate in fish teeth. Puceat

E (ed.). *PLOS ONE* 2019;**14**:e0220390.

Smale DA, Wernberg T, Oliver ECJ *et al.* Marine heatwaves threaten global biodiversity and

the provision of ecosystem services. *Nat Clim Change* 2019;**9**:306–12.

Stuiver M, Reimer PJ. A Computer Program for Radiocarbon Age Calibration. *Radiocarbon*

1986;**28**:1022–30.

Trisos CH, Merow C, Pigot AL. The projected timing of abrupt ecological disruption from

climate change. *Nature* 2020;**580**:496–501.

Wanamaker AD, Kreutz KJ, Schöne BR *et al.* Coupled North Atlantic slope water forcing on

Gulf of Maine temperatures over the past millennium. *Clim Dyn* 2008;**31**:183–94.

Wilson S, Block B. Habitat use in Atlantic bluefin tuna Thunnus thynnus inferred from diving

behavior. *Endanger Species Res* 2009;**10**:355–67.

Zavatarelli M, Raicich F, Bregant D *et al.* Climatological biogeochemical characteristics of the

Adriatic Sea. *J Mar Syst* 1998;**18**:227–63.

  Supplementary material

  S1. List of climatic and cultural events

  Table S1 – List of climatic and cultural events, with corresponding names, time intervals and
  references.

Type of Event	Name	Interval (years BP)	Reference
Climatic	Last Glacial Maximum (LGM)	32,000 - 21,000	Lionello *et al.* (2023)
	Younger Dryas	14,749 - 13,646	Benjamin *et al.* (2017)
	Holocene 1st cooling event (Event 1)	10,500 - 9,800	Lionello *et al.* (2023)
	Rapid Climate Change (RCC)	8,499 - 7,999	Krauß *et al.* (2018)
	Holocene 2nd cooling event (Event 2)	8,000 - 7,000	Lionello *et al.* (2023)
	Holocene 3rd cooling event (Event 3)	5,000 - 4,800	Lionello *et al.* (2023)
	Medieval Warm Period (MWP)	1,050 - 650	Briffa (2000)
	Little Ice Age (LIA)	650 - 100	Briffa (2000)
Cultural	Late Paleolithic	51,949 - 21,949	Stiner and Kuhn (2006)
	Mesolithic	21,949 - 11,949	Stiner and Kuhn (2006)
	Epipaleolithic	11,949 - 8,949	Tortosa (1995); Sánchez (2007)
	Neolithic (Eastern Mediterranean)	8,649 - 5,049	Andreou *et al.* (1996)
	Neolithic (Western Mediterranean)	7,000 - 5,000	Stuiver and Reimer (1986)
	Bronze Age (Eastern Mediterranean)	5,049 - 2,999	Andreou *et al.* (1996)
	Chalcolithic-Bronze (Western Mediterranean)	5,000 - 3,000	Stuiver and Reimer (1986)

Iron Age	3,000 - 2,200	Stuiver and Reimer (1986)
Hellenistic Period	2,349 - 1,950	Morand (2020)
Roman Period (Western Mediterranean)	2,249 - 1,400	Roberts *et al.* (2011)
Roman Period (Eastern Mediterranean)	1,950 - 1,350	Morand (2020)
Byzantine Period	1,626 - 497	Athanassopoulos (2004)
Dark Age	1,400 - 1,090	Roberts *et al.* (2011)

654 **S2. Results from statistical analyses**

655 **Table S2.1 –** Results from the statistical analyses performed regarding MNI data for the
656 banded-dye murex for the Western Mediterranean. Numbers in bold represent statistical
657 differences.

Species/Tests	Shapiro-Wilk test	Levene's test	Kruskal-Wallis test	Pairwise Wilcox test
Banded-dye murex	$W = 0.480$ $p = 1.349e^{-7}$	$F = 4.270$ $p = 0.03$	$H_2 = 15.792$ **$p = 0.0004$**	**$p = 0.0013$**

659 **Table S2.2 –** Results from the statistical analyses performed regarding catch data for the
660 Atlantic bluefin tuna for the Western Mediterranean, for periods divided using the Jenks
661 natural breaks classification method (*classInt* package; Bivand 2023).

Tests	Shapiro-Wilk test	Levene's test	Kruskal-Wallis test
Atlantic bluefin tuna	$W = 0.91514$ $p = 3.12e^{-12}$	$F = 33.624$ $p < 2.2e^{-16}$	$H_5 = 407.55$ **$p < 2.2e^{-16}$**

663 **Table S2.2.1 –** Results from the pairwise comparisons between periods for the Atlantic
664 bluefin tuna, adjusted by the Bonferroni correction method. Numbers in bold represent the
665 periods that differ significantly from each other.

	[14,95]	(54, 95]	(95,136]	(136,175]	(175,212]
(54, 95]	**$< 2e^{-16}$**				
(95,136]	1	**$< 2e^{-16}$**			
(136,175]	1	**$< 2e^{-16}$**	1		
(175,212]	**$< 2e^{-16}$**	1	**$< 2e^{-16}$**	**$< 2e^{-16}$**	
(212,250]	**$6e^{-04}$**	**$< 2e^{-16}$**	**$3e^{-04}$**	**$1.1e^{-05}$**	**$< 2e^{-16}$**

667 **Table S2.3 –** Results from the statistical analyses performed regarding total length (TL) data
668 for the Atlantic bluefin tuna and the gilthead sea bream by subregions of the Mediterranean
669 Sea.

Tests	Shapiro-Wilk test		Levene's test		Two samples t-test	Mann-Whitney U test	one-way ANOVA
Species/ Subregion	Western	Eastern	Western	Eastern	Western	Eastern	Eastern
Atlantic bluefin tuna	$W = 0.9593$ $p = 0.1686$	$W = 0.9091$ $p = 0.0163$	$F = 2.103$ $p = 0.1554$	$F = 0.9167$ $p = 0.3468$	$t = 2.967$ **$p = 0.01$**	$W = 8$ $p =$ **0.01035**	
Gilthead sea bream		$W = 0.9898$ $p = 0.4921$		$F = 1.5013$ $p = 0.227$			$F = 10.28$ **$p = 7.54e^{-05}$**

671 **Table S2.4 –** Results from the statistical analyses performed regarding basal diameter (BD)
672 data (mean and maximum values) for the Turbinate monodont for the Eastern Mediterranean.

Tests	Shapiro-Wilk test		Levene's test		one-way ANOVA	Welch ANOVA
Variable/ Subregion	Mean	Maximum	Mean	Maximum	Mean	Maximum
Turbinate monodont	$W = 0.9102$ $p = 0.245$	$W = 0.9794$ $p = 0.4367$	$F = 1.2519$ $p = 0.3364$	$F = 4.4836$ $p = 0.0071$	$F = 2.021$ $p = 0.195$	$F = 0.3701$ $p = 0.7767$

674 **Table S2.5 –** Results from the statistical analyses performed regarding stable isotopic data
675 ($\delta^{13}C$ and $\delta^{15}N$ values) for the Atlantic bluefin tuna between the Western and Eastern
676 Mediterranean and the gilthead sea bream for the Eastern Mediterranean.

Tests	Shapiro-Wilk test		Levene's test		Mann-Whitney U test		Kruskal-Wallis	one-way ANOVA
Species/ Stable Isotopes	$\delta^{13}C$	$\delta^{15}N$	$\delta^{13}C$	$\delta^{15}N$	$\delta^{13}C$	$\delta^{15}N$	$\delta^{13}C$	$\delta^{15}N$
Atlantic bluefin tuna	$W = 0.9335$ $p = 0.0091$	$W = 0.8727$ $p = 0.0002$	$F = 1.141$ $p = 0.291$	$F = 41.743$ $p = 8.626e^{-8}$	$W = 1$ **p = 3.927e^{-9}**	$W = 16$ **p = 1.2e^{-7}**		
Gilthead sea bream	$W = 08855$ $p = 0.1794$	$W = 0.9172$ $p = 0.3695$	$F = 45.719$ $p = 0.0002$	$F = 3.7329$ $p = 0.0885$			$H = 0.70028$ $p = 0.7046$	$F = 2.344$ $p = 0.177$

S3. Presence and abundance of the targeted species

The temporal distribution of data varies across these groups: fish data are concentrated

between 100,000–12,500 years BP, mollusc data between 100,000–50,000 years BP, and

marine mammal data between 9,000–500 years BP (Figures S3.1, S3.2, and S3.3).

Geographically, the majority of data samples are from the Western (n = 194) and Eastern (n

= 326) Mediterranean, with only a few data samples from the Adriatic Sea (n = 11). Among

marine mammals, the monk seal shows a peak in NISP (~70) around 12,500 years BP in the

Western Mediterranean. Within the Western and Eastern Mediterranean, the gilthead sea

bream, Atlantic bluefin tuna, and turbinate monodont have the highest amounts of data. For

fish, data for the gilthead sea bream and Atlantic bluefin tuna span the entire time interval.

However, comparisons between periods for these two species within each region yielded no

significant differences (Figures S3.1.1B–S3.1.4B). Regarding molluscs, the turbinate

monodont is the only species with sufficient data to analyse differences between periods

across the Mediterranean. This species shows two distinct NISP peaks: one between 45,000–

30,000 years BP in the Eastern Mediterranean and another between 15,000–10,000 years

BP, primarily in the Western Mediterranean. Despite these patterns, comparisons across

periods revealed no significant outcomes (Figures S3.2.1B and S3.2.2B).

Figure S3.1 - Distribution of data for NISP for fishes. Colours separate the data for each
Mediterranean of the Mediterranean Sea.

Figure S3.1.1 - Distribution of data for NISP on the Atlantic bluefin tuna in the Western Mediterranean. (A) Temporal distribution of records, including major climatic events (purple rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each geological period. Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences between periods. The number of data samples (n) for each boxplot is shown above. The plot inside the second figure is a zoom-in of the data, for better visualisation of the increase in NISP values towards the Late Holocene (no outliers were excluded).

Figure S3.1.2 - Distribution of data for NISP on the Atlantic bluefin tuna in the Eastern
Mediterranean. (A) Temporal distribution of records, including major climatic events (purple
rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each
geological period. Blue dots represent mean values, and bars indicate standard deviation.
Letters denote statistically significant differences between periods. The number of data
samples (n) for each boxplot is shown above.

Figure S3.1.3 - Distribution of data for NISP on the gilthead sea bream in the Western
Mediterranean. (A) Temporal distribution of records, including major climatic events (purple
rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each
geological period. Blue dots represent mean values, and bars indicate standard deviation.
Letters denote statistically significant differences between periods. The number of data
samples (n) for each boxplot is shown above.

Figure S3.1.4 - Distribution of data for NISP on the gilthead sea bream in the Eastern Mediterranean. (A) Temporal distribution of records, including major climatic events (purple rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each geological period. Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences between periods. The number of data samples (n) for each boxplot is shown above.

Figure S3.2 - Distribution of data for NISP for molluscs. Colours separate the data for each Mediterranean of the Mediterranean Sea.

Figure S3.2.1 - Distribution of data for NISP on the turbinate monodont in the Western
Mediterranean. (A) Temporal distribution of records, including major climatic events (purple
rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each
geological period. Blue dots represent mean values, and bars indicate standard deviation.
Letters denote statistically significant differences between periods. The number of data
samples (n) for each boxplot is shown above.

Figure S3.2.2 - Distribution of data for NISP on the turbinate monodont in the Eastern
Mediterranean. (A) Temporal distribution of records, including major climatic events (purple
rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of NISP data for each
geological period. Blue dots represent mean values, and bars indicate standard deviation.
Letters denote statistically significant differences between periods. The number of data
samples (n) for each boxplot is shown above.

764
765 **Figure S3.3 -** Distribution of data for NISP for marine mammals. Colours separate the data
766 for each Mediterranean of the Mediterranean Sea.

768
769 **Figure S3.4 –** MNI species distribution data. Colours separate the data for each
770 Mediterranean of the Mediterranean Sea.

774 **Figure S3.4.1 -** Distribution of data for MNI on the turbinate monodont in the Western
775 Mediterranean. (A) Temporal distribution of records, including major climatic events (purple
776 rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of MNI data for each

geological period. Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences between periods. The number of data samples (n) for each boxplot is shown above.

Figure S3.4.2 - Distribution of data for MNI on the turbinate monodont in the Eastern Mediterranean. (A) Temporal distribution of records, including major climatic events (purple rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of MNI data for each geological period. Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences between periods. The number of data samples (n) for each boxplot is shown above.

S4. Catch records

Figure S4 – Catches records of the Atlantic bluefin tuna divided into six periods, using the
Jenks Natural Breaks Method, between 250–14 years BP. The blue dots, connected by the
blue line, are the mean values for each period. The number of samples (n) included in each
box plot is indicated above. The black dots are outliers.

S5. Body size measures

803
804   **Figure S5.1 –** Fishes distribution of data for Standard Length (SL). The colour indicates that
805   the data comes from the Eastern Mediterranean Sea.

808   **Figure S5.2 –** Fishes distribution of data for Total Length (TL). The colour indicates from
809   which sub-basin of the Mediterranean Sea the data comes.

810
811
812

813
814
815
816

Figure S5.3 – Distribution of data for TL on the Atlantic bluefin tuna in the Western
Mediterranean. (A) Boxplots of TL data for each geological period. Blue dots represent mean
values, and bars indicate standard deviation. Letters denote statistically significant
differences between periods. The number of data samples (n) for each boxplot is shown
above. (B) Total length differences between the MWP (Medieval Warm Period) and LIA
(Little Ice Age).

Figure S5.4 – Distribution of data for TL on the Atlantic bluefin tuna in the Eastern
Mediterranean. (A) Boxplots of TL data for each geological period. Blue dots represent mean

values, and bars indicate standard deviation. Letters denote statistically significant
differences between periods. The number of data samples (n) for each boxplot is shown
above. (B) differences between the before and during the MWP (Medieval Warm Period).

Figure S5.5 – Distribution of data for the TL on the Gilthead sea bream in the Eastern
Mediterranean. Differences between the Bronze and Iron Ages and the Byzantine period.

Figure S5.6 - Distribution of data for mean BD on the turbinate monodont in the Eastern
Mediterranean. (A) Temporal distribution of records, including major climatic events (purple
rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of BD data for each
geological period. Blue dots represent mean values, and bars indicate standard deviation.
Letters denote statistically significant differences between periods. The number of data
samples (n) for each boxplot is shown above.

Figure S5.7 - Distribution of data for maximum values of BD on the turbinate monodont in
the Eastern Mediterranean. (A) Temporal distribution of records, including major climatic
events (purple rectangles) and anthropogenic events (black dashed lines); (B) Boxplots of BD
data for each geological period. Blue dots represent mean values, and bars indicate standard
deviation. Letters denote statistically significant differences between periods. The number of
data samples (n) for each boxplot is shown above.

S6. Stable isotopic data

890   **Figure S6.1 –** Distribution of data for the δ^{13}C and δ^{15}N of the Atlantic bluefin tuna from the
891   Western and Eastern Mediterranean. (A) The δ^{13}C values fall within the Medieval Warm
892   Period (MWP); (B) The δ^{15}N values fall within the Medieval Warm Period (MWP).

894
895

Figure S6.2 – Distribution of data for the δ^{13}C of the gilthead sea bream from the Eastern
Mediterranean. (A) The δ^{13}C values fall between the Bronze and Iron Ages; (B) Boxplots of
the differences between data with three mean ages. Blue dots represent mean values, and
bars indicate standard deviation. Letters denote statistically significant differences between
periods. The number of data samples (n) for each boxplot is shown above.

Figure S6.3 – Distribution of data for the δ^{15}N of the gilthead sea bream from the Eastern
Mediterranean. (A) The δ^{15}N values fall between the Bronze and Iron Ages; (B) Boxplots of

the differences between data with three mean ages. Blue dots represent mean values, and bars indicate standard deviation. Letters denote statistically significant differences between periods. The number of data samples (n) for each boxplot is shown above.

References

Andreou S, Fotiadis M, Kotsakis K. Review of Aegean prehistory V: The Neolithic and Bronze Age of northern Greece. *American Journal of Archaeology* 1996;**100(3)**:537–97.

Athanassopoulos EF. Historical archaeology of medieval Mediterranean landscapes. In: University of Pennsylvania Museum of Archaeology and Anthropology (ed.), *Mediterranean Archaeological Landscapes: Current Issues*. University of Pennsylvania Museum of Archaeology and Anthropology, 2004, 81–98.

Benjamin J, Rovere A, Fontana A *et al.* Late Quaternary sea-level changes and early human societies in the central and eastern Mediterranean Basin: An interdisciplinary review. *Quat Int* 2017;**449**:29–57.

Bivand R. *classInt: Choose Univariate Class Intervals*. R package version 0.4-10, 2023. https://CRAN.R-project.org/package=classInt.

Briffa KR. Annual climate variability in the Holocene: interpreting the message of ancient trees. *Quat Sci Rev* 2000;**19**:87–105.

Krauß R, Marinova E, De Brue H *et al.* The rapid spread of early farming from the Aegean into the Balkans via the Sub-Mediterranean-Aegean Vegetation Zone. *Quat Int* 2018;**496**:24–41.

Lionello P, Giorgi F, Rohling E *et al.* Mediterranean climate. *Oceanography of the Mediterranean Sea*. Elsevier, 2023, 41–91.

Morand N. The exploitation of molluscs and other invertebrates in Alexandria (Egypt) from

the Hellenistic period to Late Antiquity: food, usage, and trade. *Anthropozoologica*

2020;**55**:1.

Sánchez MC. *Cueva Bajondillo (Torremolinos): Secuencia cronocultural y paleoambiental del*

cuaternario reciente en la bahía de Málaga. Centro de Ediciones de La Diputación de

Málaga, 2007.

Roberts N, Brayshaw D, Kuzucuoğlu C *et al.* The mid-Holocene climatic transition in the

Mediterranean: Causes and consequences. *The Holocene* 2011;**21**:3–13.

Stiner MC, Kuhn SL. Changes in the 'Connectedness' and Resilience of Paleolithic Societies in

Mediterranean Ecosystems. *Hum Ecol* 2006;**34**:693–712.

Stuiver M, Reimer PJ. A Computer Program for Radiocarbon Age Calibration. *Radiocarbon*

1986;**28**:1022–30.

Tortosa JEA. El Magdaleniense Mediterráneo: La Cova del Parpalló (Gandia, Valencia). 1995.