Life history evolution of insects in response to climate variation: seasonal timing versus thermal physiology.

Karl Gotthard[1]*, David Berger[2]* & Patrick Rohner[3]*

*All authors contributed equally

Affiliations:

1. *KG: Department of Zoology, Stockholm University, 106 91 Stockholm, Sweden.* ORCID: https://orcid.org/0000-0002-4560-6271

2. *DB: Department of Ecology and Genetics, Uppsala University, Uppsala, 752 36 Sweden. ORCID: https://orcid.org/0000-0003-0196-6109*

3. *PTR: Department of Ecology, Behavior, and Evolution, University of California San Diego, La Jolla, CA 92093, USA. ORCID: https://orcid.org/0000-0002-9840-1050*

Abstract

Climate adaptation in insects can proceed via responses in life history traits and their thermal plasticity, and through phenological shifts mediated by responses to photoperiodic cues ("photoperiodism"). While, experimental studies demonstrate evolutionary potential for both modes of adaptation, it remains unclear how evolution will unfold in natural populations, limiting our ability to predict how insects will respond to climate change. Here we review the literature and perform an analysis of published studies revealing that photoperiodism for diapause induction evolves predictably along latitude, with high-latitude populations entering diapause earlier. In contrast, although a few species showed clinal variation in life history and thermal plasticity, the direction of these clines were not consistent across taxa. This suggests that while life history traits sometime evolve rapidly, photoperiodism is a more predictable response to climate, and thus, climate adaptation is more likely to proceed via phenological shifts than physiological adaptations to changing temperatures.

Introduction

Ecological and evolutionary processes influenced by climate variation are widely studied due to the urgent need to understand the impact of global climate change on biodiversity (3, 49, 102, 105, 133, 134, 143). It is well established that global warming has already had strong ecological effects on biological communities by affecting the distribution, abundance, and phenology (seasonal timing) of many organisms, including insects (27, 38, 45, 48, 84, 103, 104, 133, 139). However, the mechanisms by which insects respond and adapt to climate change is less well understood (3, 23, 52, 70). This presents a challenge not only for assessing future extinction risk of focal species, but also for predicting stability of species interactions and ecosystems as functions of alternative evolutionary strategies in response to a warming world (6, 24, 77). For instance, while changes in climate can lead to the evolution of physiological responses that mitigate environmental stress (52), the evolution of phenology might offer an alternate route to adaptation (17, 20, 68) (Figure 1). While physiological and phenological responses may both facilitate adaptation, their effects on insect growth rates and distribution ranges are likely to differ. Here, we review the literature on insect life history evolution in response to variation in climate, with a particular focus on work that has explored genetic responses in both thermal plasticity and phenology via the use of photoperiodic cues (from hereon: "*photoperiodism*").

Physiological temperature adaptation in life-history traits

Life history theory sees the scheduling of life cycle events such as growth, maturation and reproduction as the result of strategic decisions made over an organism's life (88, 99, 115, 126). Spatiotemporal variation in climate variables such as temperature, precipitation, and humidity are major sources of natural selection on life history strategies in general, and maybe in particular for ectotherms like insects (3, 23, 52, 59, 99, 131). Virtually all physiological processes of insects are at least somewhat dependent on temperature, leading to abundant phenotypic plasticity in life history responses to ambient temperatures (3, 23, 99). For instance, decreased temperature often leads to plastic increases in adult body size in arthropods (the so-called temperature-size rule) (5, 53) and prolongs egg-to-adult development time (3, 23, 99). Nevertheless, theory predicts that an organism's range of tolerated and preferred temperatures should evolve to match those experienced in its environment (3, 57), and indeed, there are often large evolved species differences in the degree to which species are affected by temperature (3, 23, 99). Moreover, there is often abundant genetic variation within species, indicating that thermal plasticity can evolve, sometimes even on ecological timescales (12, 46, 82). This suggests that natural selection may alter temperature dependence of important physiological and life history traits (3, 12, 36, 46, 86, 141), offering an avenue to adapt to climate change (Figure 1C-E).

Seasonal adaptation in life-history via photoperiodism

Most organisms experience some type of seasonal variation in climate that also drive associated variation in biotic resources and interactions (22, 34, 131). Seasonality is consequently a highly predictable form of temporal variation in climate and other ecological conditions (59, 131). Adaptation to seasonality requires that organisms can exploit the favorable seasons, avoid or mitigate the effects of the unfavorable seasons, and adequately time the transitions between respective stages in their life cycle (22). This is especially important for insects because most species use some kind of dormancy to withstand harsh conditions (34, 131) (although there are notable examples of insects that migrate long distances to evade hazardous conditions (28, 131, 148)). In temperate insects, the typical adaptation to mitigate the effects of the unfavorable winter season is to enter diapause before

the onset of cold conditions (i.e., a state of dormancy that arrests or greatly prolongs
development). The individual then remains in diapause until there is no longer a risk of
developing prematurely and finally restarts development and reproduction when the thermal
conditions again become permissive (34, 74, 131, 142). Although variation in thermal
conditions and biotic resources are likely to be the major sources of selection on life cycle
timing, temperate insects have commonly evolved to use the photoperiod as a cue of seasonal
change (34, 59, 131). Photoperiod provides almost noise-free information on seasonal
timing, allowing insects to appropriately schedule diapause. The sensitivity to photoperiod
typically follows a threshold function where in certain photoperiods (typically long summer
daylengths) individuals follow non-diapause (or 'direct') development, while in other
photoperiods (typically shorter late summer/autumn daylengths) they instead enter diapause
(68, 131). The inflection-point of this threshold function is referred to as the critical
daylength (CDL) and is often used as a metric for assessing genetic variation in
photoperiodism between populations or species (16, 22, 34, 68, 124, 131). This type of
photoperiodic plasticity allows a given genotype to develop without a diapause in the warm
summer months while still inducing diapause in individuals that are active later in the season.
The facultative induction of diapause allows insect populations to produce multiple
generations within a given year (multivoltinism) and genetic variation for CDL is often
strongly associated with variation in voltinism (81, 124), and the length of the permissive
season for growth and reproduction (68).
111
112   **Figure 1**

113
114

Figure 1: Insects have several mechanisms to respond to changing climates. A & B) Populations may track their preferred environment by shifting their geographic distribution range (option 1). Alternatively, they may remain in place and adapt to the new climatic conditions (option 2). C) This adaptation can occur through the evolution of thermal tolerance (option 2a) and/or changes in seasonal timing (option 2b). D) An organism's thermal tolerance evolves through changes in the minimal (CT_{min}), optimal (T_{opt}), and/or maximal (CT_{max}) temperature at which it can operate. As thermal tolerance is governed by trade-offs between performance at different temperatures, climate warming can result in evolutionary shifts in all these parameters (blue to red reaction norm). Note, however, that climate adaptation of growth and development also can occur through vertical shifts of the entire reaction norm (between blue and broken red reaction norm) so that growth is faster in northern populations that have less time to complete development, or slows down in future warming climates (so called "countergradient" variation). E) An organism can change its seasonal timing via the evolution of photoperiodism; by modification of how it uses and interprets daylength. Adaptation to climate warming by shifts or prolongations of the timing for growth and reproduction towards colder parts of the year (option 2b) is predicted to be associated with changes in the critical daylength (CDL) at which diapause is induced (blue to red reaction norm). Additionally, if species respond to warming climates by migrating closer to the poles (option 1), they also need to evolve changes in how they interpret daylength in order to time their life cycle. Importantly, the evolution of seasonal timing can relax selection for changes in thermal tolerance (and vice versa), and changes in seasonal timing may also require life history adjustments to accommodate changes in season length, such as overall increases or decreases in development rate and size at maturation, predicting correlated evolution of photoperiodism and life history.

Two modes of adaptation

Temperature variation during the favorable season is likely to exert strong selection on the thermal plasticity of life history traits, whereas temperature variation at the end of this season will exert selection on the photoperiodic regulation of diapause induction. It is important to note that these two types of selection pressures interact with each other. On one hand, evolved responses in photoperiodism can shield the organism from unfavorable temperatures. On the other hand, evolved responses in thermal plasticity can in turn relax selection for shifts in phenology. Hence, we expect these two interacting modes of adaptation to exert correlational selection on the optimal life history strategy to mitigate climate variability (Figure 1C-E). However, the relative significance of selection on thermal physiology versus selection on photoperiodism is not well understood. Here we review how insect life histories have evolved in response to climate variation across both time and space by comparing cases where climate variation is expected to exert natural selection on life history traits and their thermal plasticity, as well as on photoperiodism for life cycle regulation.

Evolution of thermal performance and photoperiodism in selection experiments
Experiments using artificial selection or experimental evolution to study the evolution of thermal reaction norms (i.e. thermal plasticity) have been performed in many species (3, 85). These experiments provide a proof-of-principle of how natural selection can shape thermal plasticity in life history. A recent meta-analysis of such experiments on ectotherms compared thermal reaction norms of evolved populations after thermal selection, to control (or ancestral) populations, allowing quantification of thermal adaptation (85). About a quarter of the studies included in the meta-analysis were done on insects. The results show a general positive response to thermal selection, where estimates of fitness in the assay temperature increased after selection. Moreover, thermal selection in the laboratory also had the capacity to change the shape of reaction norms so that adaptation to warmer temperatures generally led to lower fitness at colder temperatures (85). Although these effects were generally strongest in unicellular organisms, they still suggest that selection over relatively short timespans has capacity to alter the thermal sensitivity of life history traits and that there may be genetic trade-offs between performing at different thermal conditions.

Different types of artificial selection experiments that explore the evolutionary potential of
photoperiodism for diapause induction and duration have been performed on species from all
large insect orders (131). Typically, they show that these seasonal adaptations can change
relatively rapidly in laboratory experiments (131). For instance, artificial selection on the
photoperiodic induction of reproductive diapause in the fly *Drosophila montana*
demonstrated a rapid change in critical daylength (69). Using a baseline population derived
from a northern location in Finland, selection for eight generations where only non-
diapausing individuals were allowed to breed under continuously shorter daylengths resulted
in shorter CDL in all replicate lines compared to the control. These selected lines evolved a
response to photoperiod that is typical for more southern populations of *D. montana*.
Similarly, bidirectional artificial selection for non-diapause and diapause development led to
the evolution of photoperiodic plasticity for diapause induction in the North American
mosquito *Wyeomyia smithii* after 12 generations (18), and in the chrysomelid beetle
Ophraella communa after 5-8 generations (129).
182
183 In summary these examples demonstrate that both thermal reaction norms and photoperiodic
184 reaction norms have the potential to evolve relatively rapidly in response to directional
185 selection in the laboratory, suggesting that there is standing genetic variation present within
186 natural populations that can fuel rapid evolutionary change in the wild.
187
Life history evolution in response to recent climate change
Rising temperatures are altering patterns of seasonality and phenology — a trend expected to
be further exacerbated in the foreseeable future (3). Against the backdrop of strong declines
in insect abundances in recent years (118, 139), there is increasing interest in the effects
of climate on insect populations and their life histories (49, 143). However, to detect and
estimate phenotypic evolution in natural populations it is necessary to measure heritable (i.e.,
'genetic') changes in traits in the same populations repeatedly (at least twice) over several
years or even decades.
196
In insects these types of studies are relatively rare as they require controlled common garden
experiments to be replicated over extended time periods. Still, there are some notable
examples, spearheaded by work on the North American Pitcher plant mosquito (*Wyeomyia*
smithii) by Bradshaw and Holtzapfel (16). The authors demonstrated that the photoperiodic
regulation of the life cycle, measured by the CDL for diapause induction in a range of
latitudinally separated populations, showed a general genetic shift towards shorter CDLs, that
are typical in more southern populations. This shift likely represents adaptive evolution in
response to a warming climate as shorter CDLs result in a longer growth season and a later
date for diapause induction. Surprisingly, this evolutionary change was detectable over only
five years of evolution in the field (16). Similar results have since been found in other insects
(41, 96, 97), indicating that changes in photoperiodism for life cycle regulation may
constitute a common evolutionary response to climate change.
209
However, there have also been documented evolutionary changes in life history traits over
time. For instance, the Yellow dung fly (*Scathophaga stercoraria*) shows genetically based
reductions in body size (13), and a study on the Winter moth (*Operophtera brumata*) found a
reduction in egg development rate (137). The latter example presumably represents a rare
demonstration of how climate change has led to the evolution of thermal plasticity of
development rate. Interestingly, this response also seems related to selection for a change in
phenology (137) as several populations in the Netherlands have evolved a reduction in egg
development rate in a range of relevant spring temperatures, most likely due to selection for

maintaining synchronization of egg hatching with the budburst of its host plant, the Oak
(Quercus robur), in the now warmer spring.
220
Life history evolution in response to climate variation in space
Climate and seasonality vary greatly in space, and this variation is particularly predictable
along latitudinal gradients (59). Studies of life history evolution in insects have a long
tradition of exploring this type of latitudinal variation in climate (3, 14, 114, 123, 131). In
many cases researchers have exploited this spatial variation to explore the likely effects of
current climate change by the so-called "space-for-time" substitution (59).
227
There is a large body of literature that explores the thermal biology of life history evolution
across insect species (3). In particular variation in thermal tolerance limits (i.e., the critical
minimum and maximum temperatures, CT_{min} and CT_{max}, respectively) across species and
geographic locations are well-studied (2, 11, 51, 67, 127). A general pattern across insects
(2, 51) and in ectotherms in general (11, 127) is that lower thermal limits show greater
adaptive variation across climate regions at the species-level, compared to upper thermal
limits. As a consequence, many researchers have converged on the conclusion that there may
be limited scope for rapid evolution of CT_{max} in response to a warming world (11, 51, 67).
Still, artificial selection experiments show that populations of the Fruit fly (*Drosophila*
melanogaster) harbors additive genetic variation that allows successful laboratory evolution
of increased heat tolerance, although the level detected may still be insufficient to keep up
with the predicted increase in future temperatures (46). Moreover, as temperatures during the
hottest parts of the year vary less across the typical geographic regions studied than do
thermal minima, it is also possible that the relatively modest differentiation in CT_{max} can in
part be explained by similar selection pressure across geographic regions (65). Studies
exploring latitudinal variation in thermal tolerance within species provide a similar, albeit
less strong, pattern as the between species comparisons (7, 9, 71, 72, 92, 112, 120, 122, 140,
147, 153). For example, within the limited data we reviewed for our quantitative analysis
(see below), we found estimates of latitudinal variation in heat and cold tolerance for 4 and 8
dipterans, respectively. The observed clinal variation was not very consistent among species
but perhaps slightly more so for cold tolerance, which tended to be higher in more northern
populations. Interestingly, insects that have colonized urban environments show parallel
evolution of both CT_{max} and CT_{min} (36, 86, 90), due to local adaptation to the generally
warmer urban environment. Together, this suggests that adaptive responses to natural
selection on survival at thermal extremes do occur as insects colonize new climate
zones.
254
There is also a large number of insect studies that have explored the evolution of
development rates, body size, and their thermal plasticity across latitude (3, 14, 58, 73, 87,
100, 123, 125). However, predictions of how natural selection should affect
development rate, size and their temperature dependencies are not always
straightforward (15, 29, 53, 123). A common expectation is that because both mean
temperatures and the length of insect growth seasons decrease with distance from the equator,
natural selection should favor faster development rates (shorter development times) and
thermal plasticity that maximizes growth in colder conditions in populations living at higher
latitudes (73, 114). This pattern has been called "counter gradient variation" or "genetic
compensation", as genetic adaptation across latitude opposes the direct effect of temperature
on plasticity in growth and development rates (30, 31, 44, 79). As a consequence of the
shortened time for growth and development, it can be expected that adult size of insects
should become smaller when moving away from the equator (114). However,

physiological models assuming that rates of growth and development have different thermal
plasticity predict that there are situations when insects should grow to larger sizes at colder
latitudes (138, 144). Studies of latitudinal clines in body size within insect species have
indeed found evidence for both these patterns (15, 73, 123). For instance, Shelomi (2012)
reviewed a large set of both inter- and intraspecific latitudinal and altitudinal clines in insect
body sizes, and while there were approximately equal numbers of positive and negative size
clines, the most common pattern was the absence of a size cline altogether (123). Similar
mixed patterns were found in a metanalysis of within-species size clines in arthropods
that was mainly based on insects (53). Interestingly, this study found strong evidence that
terrestrial taxa were more likely to show reductions in size with latitude, whereas aquatic taxa
typically showed the opposite pattern. This result may be explained by the difference between
water and air in how oxygen limitation affects both the plasticity and evolution of growth and
development. Moreover, in the terrestrial taxa the pattern of voltinism influenced the
latitudinal size clines as species with one or two generations per year decreased in size with
latitude, while multivoltine species instead showed a weak positive size cline with latitude.
This suggests that when growth season length increases towards the equator, species showing
strong latitudinal clines in voltinism (multivoltine) are under selection to produce additional
generations, while in univoltine (and bivoltine) species selection instead favors an increase in
adult size via prolongation of development (53).
287
Because photoperiods change very predictably across latitudes, it is expected that natural
selection should favor latitudinal variation in adaptations for seasonal timing in response to
daylength. Indeed, there is a large number of examples demonstrating local adaptation in
photoperiodism of insects, in particular in plasticity for diapause induction (33, 59, 68, 131).
The most common pattern is that the critical daylength for diapause induction (CDL)
increases with latitude across populations, which shows that more northern populations with
a shorter growth season typically enter diapause earlier in year (68, 131). Latitudinal clines
in CDL can evolve relatively rapidly in introduced species (10, 40, 131, 136), species
colonizing urban areas (89), and in species expanding their ranges due to climate change
(60).
298
The available evidence consequently suggests that life history traits such as development
rates, body size and their thermal dependencies, as well as the photoperiodic plasticity for
seasonal life cycle regulation do evolve across space and to some degree also with ongoing
climate change. However, it is still unclear whether natural populations are more likely to
adapt through changes in thermal plasticity, the adjustment of seasonal activity (through the
evolution of photoperiodism), or both. Previous attempts to assess whether thermal- or
photoperiodic adaptations are more likely to evolve in response to rapid climate change
suggested that genetic changes in photoperiodism for seasonal life cycle regulation was a
considerably more likely "first response" compared to thermal adaptation (17, 20). As these
early reviews were based on the relatively limited set of then known examples and that
selection due to climate change is likely to be multifaceted, other researchers have argued
that this conclusion is premature (52). Unfortunately, well established examples of
evolutionary change due to recent climate change are still rare, and for insects we have
not found more than the handful of examples reviewed above. However, because of the
large number of studies exploring evolution of life history traits and their thermal
dependencies, and photoperiodism in response to climate variation in across latitude, it is
possible to explore this question in the spatial dimension. Below, we summarize and integrate
the results of previously published studies to more formally quantify the amount of

population differentiation in life history traits and their plasticity across latitude and compare
this to the corresponding amount found for photoperiodism using the same species.
319
Analysis of latitudinal clines in photoperiodism, life history, and thermal
plasticity
Full details of the literature search and analyses can be found in the supplementary material.
In brief, we looked for published common garden datasets that quantified latitudinal clines in
critical daylength as well as means and thermal plasticity of the life history traits body size
and development rate. The dataset on critical daylength was based on an initial data set
generated by Joschinski and Bonte (2021) (68). We added additional data that has since been
published and had at least 3 populations and 3 photoperiod treatments. Data on life history
was collected based on an independent literature search targeted species with photoperiodism
data, again focusing on common garden studies that reported latitudinal clines. We included
all studies that had at least two temperature treatments to calculate an index of thermal
plasticity (i.e. how temperature affected a linear response in trait means). Our final dataset
included 24 insect species that had estimates for critical daylength and development rate
and/or body size. This included representatives of the orders Coleoptera (n = 5), Diptera (n =
5), Hemiptera (n = 3), Lepidoptera (n = 10), as well as one species of Neuroptera (1, 4, 8, 14,
18, 26, 32, 39, 43, 50, 54–56, 60, 62, 63, 66, 76, 78, 81, 83, 93–95, 98, 101, 106, 107, 110,
113, 116, 117, 119, 121, 125, 128, 132, 135, 136, 145, 146, 149–152, 154).
337
We found that the vast majority of species showed an increase in CDL with latitude,
indicating that populations closer to the poles enter diapause earlier (Figs. 2,3). This is
consistent with the findings of other studies (68) and indicates a strong and consistent
evolutionary pattern found across orders. Based on critical daylength estimates, we also
calculated the mean ordinal day of the year at diapause onset. If CDL remains constant across
a species' range, one would expect populations at high latitudes to enter diapause earlier due
to the faster decline in daylengths towards the end of the season. However, we did not find a
consistent latitudinal pattern, putatively because populations evolve longer CDLs towards
high latitudes.
347
In contrast, in the same set of species, we find no consistent latitudinal effect for any of the
investigated life history traits (development rate, body size and the thermal plasticity of these
traits). This finding contrasts with the notion that counter gradient adaptation in development
rate is one of the main hypotheses for the evolution of latitudinal clines in life histories of
insects. Importantly, however, there are many instances of life history clines in the data. In
fact, latitude explained almost as much of the observed population differences within species
for mean development rate and body size (43% and 47%, respectively) as it did for
photoperiodism (54%) (Fig. 2B). For instance, the speckled wood butterfly (*Pararge aegeria*)
and southwestern corn borer (*Diatraea grandiosella*) showed rather strong genetic decreases
in development rate with latitude (43, 81, 128). However, patterns were inconsistent, and
sometimes in opposing directions, across species (Fig. 3) and we even found an instance of
inconsistent patterns within species. In the Colorado potato beetle (*Leptinotarsa*
decemlineata), development rate increases with latitude in European populations but
decreased in towards the pole in North America (62, 78). As latitudinal clines in body size
often are considered to be results of a trade-off with development time (potentially also
affected by shifts in voltinism), which showed no consistent latitudinal pattern here, the
idiosyncratic responses in body size clines are perhaps less surprising. We also found no
general pattern in thermal plasticity of body size or developmental rate. Although sample
sizes were limited for those comparisons, most estimates of the latitudinal effect on thermal

plasticity for individual species hoover around zero (Figs. 2B and 3C, F). Taken together,
these findings suggest that the mechanisms underpinning seasonal timing evolve predictably
across orders. In contrast, while life history traits often do evolve, the patterns are not
consistent across insects and remain hard to predict.
371
372
373 **Figure 2**
374

A) Geographic origin of studied populations

B) Standardized latitudinal effect across species

375
376
Figure 2: A) The geographical locations of all populations that were included in the analysis of life history
and photoperiodism. B) shows the global effects of latitude on photoperiodism, mean life history, and thermal
plasticity in life history (means and corresponding 95% confidence limits). Only critical daylength shows a
strong and consistent clinal pattern across species. The standardized latitudinal effect refers to the global slope
estimated in a Bayesian mixed multilevel model divided by the standard deviation in the slope observed across
species. The horizontal bars to the right show the average proportion of population differentiation within each
study that was explained by latitude. Note that although life history traits did not show consistent clinal variation
across species, latitude still explains a large fraction of the observed population-level variance, indicating
evolution of latitudinal clines in opposing directions across species.
386

387
An important caveat is that there are still a relatively small number of species for which we
were able to find both photoperiodism and life history data and we were not able to generate
replicated measurements within species. This limits our ability to differentiate between
evolved differences among species and spurious experimental effects, despite researchers'
best efforts to implement common garden rearings. However, by including only species
where latitudinal variation in life history and CDL for diapause induction have been
estimated, we were able to assess the relative likelihood of the two modes of adaptation in
life history strategies. Although this reduced the number of species in our analysis, the results
nevertheless revealed a very strong pattern in CDL and no consistent patterns in the other
traits. In combination with the limited set of examples that have documented evolutionary
changes due to recent climate change, it seems reasonable to assume that this higher
predictability of CDL evolution pertains also to the future expected changes in climate.
400
The difference in predictability of evolutionary change may have several explanations.
Selection on CDL may simply be stronger and less dependent on other ecological
circumstances that differ among species and geographic locations. For instance, variation in
behavioral thermoregulation may affect the exposure to temperature variation, and moderate
selection on thermal plasticity of life history traits (23). Along the same lines, and perhaps
more pertinent to our analysis, it has been suggested that evolutionary changes in seasonal
life cycle regulation may mitigate selection on thermal sensitivity (23) (Fig. 1). This type of
correlational selection would arise if selection typically favors a seasonal regulation that
consistently exposes the active part of the life cycle to the time of the year that is most
thermally similar at different locations, thereby reducing the difference in thermal selection
across space; after all this is one of the main adaptive functions of insect diapause in general.
412
It is also possible that, compared to CDL, the amount of standing genetic variation is
typically lower for life history traits and their thermal plasticity, which would limit the
genetic response to selection. However, given that many studies indicate abundant genetic
variation for life history (25, 47), this hypothesis seems unlikely to be the only explanation
for the observed patterns. It is also possible that, even though life history traits harbor
significant genetic variation, general differences in the genetic architecture of these different
traits affect their evolvability. For instance, selection on life history traits and their thermal
plasticity are likely to affect many genes with strong pleiotropic effects, which could lead to
correlated evolution in other traits with maladaptive effects in the natural environment. For
example, artificial selection causing adaptation to higher temperatures in a range of
ectotherms led to correlated evolution of maladaptation at colder temperatures (85). As
temperature has such a multifaceted influence on virtually all aspects of insect biology it is
perhaps not surprising if a genetic change in one aspect of this temperature dependence also
affects the temperature dependence of other important physiological processes. In a natural
environment with highly variable thermal conditions such correlated effects are likely to
lower the rate of evolution in response to a higher (or lower) mean temperature. In contrast,
genes affecting photoperiodism may have less systemic effects and therefore be subject to
less pleiotropic constraints. Indeed, several studies have found major effect loci that underpin
intraspecific variation in diapause induction (42, 64, 75, 108, 109, 111, 130). However,
other studies suggest that photoperiodism is genetically highly complex (19, 21, 80, 111,
124) and some suggest that diapause induction and life history might be genetically
correlated through linkage or pleiotropy (91).
435

Poleward range expansions are one of the most common effects of ongoing climate change
(27, 103). Considering the general process that (re-)distributes insects across latitudes may
provide yet another explanation for why genetic clines in CDL are more predictable than any
of the other traits examined here. When insects colonize new locations across latitudes it is
likely that their existing thermal adaptations are strong determinants of where they can
establish successfully (61). This means that they may colonize new locations when thermal
conditions have changed enough to allow positive population growth. If so, selection for
rapid thermal adaptation may initially be comparatively weak in these newly colonized
locations. In contrast, all expansions across latitudes will inevitably mean that they are
exposed to a novel regime of photoperiodic variation and if a non-optimal CDL does not
preclude establishment, the novel environment will invariably exert selection on
photoperiodic plasticity and CDL (60). The latitudinal snapshots of this process that are
represented in our examples may be a reflection of this process.
449
450
451 **Figure 3**
452

453
454
Figure 3: Population differentiation in photoperiodism, life history traits, and their thermal plasticity for each
species. A) Critical day length increases consistently with latitude. In contrast, average life history (B, E) and
life history plasticity (C, F) do not show consistent latitudinal patterns across species. D) The average day of the
year at which the critical daylength for diapause is reached at each population's location ("diapause timing")
tends to decrease with latitude but shows large inconsistency across species. Development rate and body size
were standardized relative to the average trait value within each study and are expressed as percent change
relative to this average. Thermal plasticity is expressed as the percent change in trait value (relative to the global
average) per degree Celsius . Error bars indicate standard errors.
463
464
465

Conclusions

Several studies indicate that insects can relatively rapidly evolve in response to climate change, either through the evolution of thermal physiology or shifts in seasonal activity. Our review suggests that changes in photoperiodism, that is, the timing of life history transitions via use of photoperiod cues, is a predictable way in which various insect species adapt to climate variability. In contrast, while life history traits and their thermal plasticity often evolve, there is little support for a general pattern that holds broadly across species. In the context of climate adaptation, evolution of seasonal timing thus emerges as a more predictable factor than physiological adaptations to thermal conditions. As such, our updated review and more formal analysis of life history evolution in response to climate variation gives credence to the statement of Bradshaw and Holzapfel (2008): it's seasonal timing that matters.

Because the evolution of photoperiodism (or lack thereof) can lead to phenological mismatches and affect species interactions and ecosystem functioning, these evolutionary changes may have major impacts on the structure and resilience of natural systems. Our review thus offers some hope regarding our ability to predict shifts in insect phenology and its associated consequences in future climates. However, it should be noted that obligate univoltine as well as tropical insects are less likely to rely on photoperiodic plasticity for seasonal life cycle regulation and diapause induction. These insects may instead be more dependent on thermal variation also for life cycle regulation and potentially be more likely to show predictable clines in life history and thermal plasticity in relation to spatial variation in climate. Moreover, despite relatively rapid evolution of seasonal life cycle regulation, many insect populations will ultimately face significantly hotter conditions during important parts of their life cycles. The comparatively slow evolution of upper thermal limits shown in many insects is therefore clearly worrying (11, 51, 67).

Predicting how climate change will impact insect life history is crucial to mitigating both biodiversity loss and pest outbreaks (35, 48, 77). Current statistical projection models indicate that climate warming will lead to deterministic changes in insect life history via temperature's effects on biological rates (37), leading to range expansions and increases in abundance in temperate regions, but population decline in the tropics (35). However, these projections may have poor accuracy as they do not factor in evolutionary responses. For example, a recent study on the pest insect *Callosobruchus maculatus* showed that a decade of life history adaptation at hot temperature almost doubled its predicted costs to agriculture compared to simple statistical projections (24). The unpredictability of the observed life history evolution in the species analyzed here suggests that forecasting responses in insect growth rates and distribution ranges will be a significant challenge. Our review highlights four critical areas for future work that may help tackle this problem. First, the number of species in which clines in diapause induction, life history, and various physiological responses have been tested in a common garden experiment remain limited, taxonomically biased, and rarely replicated within species. Future research should aim at closing this gap. Second, there is a need for more studies that investigate how natural populations adapt in situ to changing climates (13, 16, 41, 96, 97, 137). Although these longitudinal studies suffer from a variety of experimental challenges, they provide important insights into climate-induced life history evolution in the wild. Third, studying the genetic architecture of photoperiodism, life history, and thermal physiology will provide critical insights into the evolvability of natural populations - at the moment it remains unclear if the number of loci and their pleiotropic relationships differ qualitatively between life history and seasonal timing traits, and how important genetic constraints are in explaining the observed patterns. Finally,

studies that characterize genetic (co)variation in photoperiodism and life history traits within and between natural populations should ideally combine these estimates with measures of their fitness consequences, as this will offer insights into the role of correlational selection in driving the observed clinal patterns. Important work lies ahead.

References

1. Aalberg Haugen, IM, Gotthard, K 2015. Diapause induction and relaxed selection on alternative developmental pathways in a butterfly. *J Anim Ecol* 84:464-472
2. Addo-Bediako, A, Chown, SL, Gaston, KJ 2000. Thermal tolerance, climatic variability and latitude. *Proc. R. Soc. Lond. B* 267:739-745
3. Angilletta, MJ 2009. *Thermal Adaptation: A theoretical and empirical synthesis* Oxford:Oxford University Press
4. Armbruster, P, Conn, JE 2006. Geographic Variation of Larval Growth in North American Aedes albopictus (Diptera: Culicidae). *Ann Entomol Soc Am* 99:1234-1243
5. Atkinson, D 1994. Temperature and organism size - a biological law for ectotherms? *Adv Eco Res* 25:1-58
6. Bale, JS, Masters, GJ, Hodkinson, ID, Awmack, C, Bezemer, TM et al. 2002. Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Glob Chan Biol* 8:1-16
7. Baskauf, SJ, McCauley, DE 2001. Evaluation of Low Temperature Mortality as a Range-limiting Factor for the Southwestern Corn Borer (Lepidoptera: Crambidae). *Environ Entomol* 30:181-188
8. Bauerfeind, SS, Schäfer, MA, Berger, D, Blanckenhorn, WU, Fox, CW 2018. Replicated latitudinal clines in reproductive traits of European and North American yellow dung flies. *Oikos* 127:1619-1632
9. Bauerfeind, SS, Sørensen, JG, Loeschcke, V, Berger, D, Broder, ED et al. 2018. Geographic variation in responses of European yellow dung flies to thermal stress. *J Therm Biolo* 73:41-49
10. Bean, DW, Dalin, P, Dudley, TL 2012. Evolution of critical day length for diapause induction enables range expansion of Diorhabda carinulata, a biological control agent against tamarisk (Tamarix spp.). *Evol Appl* 5:511-523
11. Bennett, JM, Sunday, J, Calosi, P, Villalobos, F, Martínez, B et al. 2021. The evolution of critical thermal limits of life on Earth. *Nat Comm* 12:1198
12. Berger, D, Postma, E, Blanckenhorn, WU, Walters, RJ 2013. Quantitative Genetic Divergence And Standing Genetic (Co)Variance In Thermal Reaction Norms Along Latitude. *Evolution* 67:2385-2399
13. Blanckenhorn, WU 2015. Investigating yellow dung fly body size evolution in the field: Response to climate change. *Evolution* 69:2227-2234
14. Blanckenhorn, WU, Bauerfeind, SS, Berger, D, Davidowitz, G, Fox, CW et al. 2018. Life history traits, but not body size, vary systematically along latitudinal gradients on three continents in the widespread yellow dung fly. *Ecography* 41:2080-2091
15. Blanckenhorn, WU, Demont, M 2004. Bergmann and converse Bergmann latitudinal clines in arthropods: Two ends of a continuum? *Integr Comp Biol* 44:413-424
16. Bradshaw, WE, Holzapfel, CM 2001. Genetic shift in photoperiodic response correlated with global warming. *Proc Natl Acad Sc USA* 98:14509-14511
17. Bradshaw, WE, Holzapfel, CM 2008. Genetic response to rapid climate change: it's seasonal timing that matters. *Mol Ecol* 17:157-166
18. Bradshaw, W, Quebodeaux, M, Holzapfel, C 2003. Circadian Rhythmicity and Photoperiodism in the Pitcher Plant Mosquito: Adaptive Response to the Photic Environment or Correlated Response to the Seasonal Environment. *Am Nat* 161:735-748
19. Bradshaw, WE, Emerson, KJ, Catchen, JM, Cresko, WA, Holzapfel, CM 2012. Footprints in time: comparative quantitative trait loci mapping of the pitcher-plant mosquito, Wyeomyia smithii. *Proc Roy Soc B* 279:4551-4558

569   20.   Bradshaw, WE, Holzapfel, CM 2006. Evolutionary Response to Rapid Climate Change. *Science*
570   312:1477-1478

571   21.   Bradshaw, WE, Holzapfel, CM 2017. Natural Variation and Genetics of Photoperiodism in
572   Wyeomyia smithii. In *99*, ed. Goodwin, SF, pp. 39-71. Academic Press

573   22.   Bradshaw, WE, Zani, PA, Holzapfel, CM 2004. ADAPTATION TO TEMPERATE
574   CLIMATES. *Evolution* 58:1748-1762

575   23.   Buckley, LB, Kingsolver, JG 2021. Evolution of Thermal Sensitivity in Changing and Variable
576   Climates. *Annu. Rev. Ecol. Evol. Syst.* 52:563-586

577   24.   Burc, E, Girard-Tercieux, C, Metz, M, Cazaux, E, Baur, J et al. 2025. Life-history adaptation
578   under climate warming magnifies the agricultural footprint of a cosmopolitan insect pest. *Nat*
579   *Comm* 16:827

580   25.   Carbonell, JA, Wang, Y-J, Stoks, R 2021. Evolution of cold tolerance and thermal plasticity in
581   life history, behaviour and physiology during a poleward range expansion. *J Anim Ecol*
582   90:1666-1677

583   26.   Chanthy, P, Martin, RJ, Gunning, RV, Andrew, NR 2015. Influence of temperature and
584   humidity regimes on the developmental stages of green vegetable bug, "Nezara viridula" (l.)
585   (Hemiptera: Pentatomidae) from inland and coastal populations in Australia. *Gen Appl Ent: J*
586   *Ent Soc New South Wales* 43:37-55

587   27.   Chen, I-C, Hill, JK, Ohlemüller, R, Roy, DB, Thomas, CD 2011. Rapid Range Shifts of Species
588   Associated with High Levels of Climate Warming. *Science* 333:1024

589   28.   Chowdhury, S, Zalucki, MP, Amano, T, Woodworth, BK, Venegas-Li, R, Fuller, RA 2021.
590   Seasonal spatial dynamics of butterfly migration. *Ecol Lett* 24:1814-1823

591   29.   Chown, SL, Gaston, KJ 2010. Body size variation in insects: a macroecological perspective.
592   *Biol Rev* 85:139-169

593   30.   Conover, DO, Present, TMC 1990. Countergradient variation in growth rate: compensation for
594   length of the growing season among Atlantic silversides from different latitudes. *Oecologia*
595   83:316-324

596   31.   Conover, DO, Duffy, TA, Hice, LA 2009. The Covariance between Genetic and Environmental
597   Influences across Ecological Gradients. *Annals N Y Acad Sci* 1168:100-129

598   32.   Couper, LI, Farner, JE, Lyberger, KP, Lee, AS, Mordecai, EA 2024. Mosquito thermal
599   tolerance is remarkably constrained across a large climatic range. *Proc Roy Soc B*
600   291:20232457

601   33.   Danilevskii, AS 1965. *Photoperiodism and Seasonal Development of Insects.* London:Oliver &
602   Boyd

603   34.   Denlinger, DL 2022. *Insect diapause* Cambridge University Press

604   35.   Deutsch, CA, Tewksbury, JJ, Tigchelaar, M, Battisti, DS, Merrill, SC et al. 2018. Increase in
605   crop losses to insect pests in a warming climate. *Science* 361:916-919

606   36.   Diamond, SE, Chick, LD, Perez, A, Strickler, SA, Martin, RA 2018. Evolution of thermal
607   tolerance and its fitness consequences: parallel and non-parallel responses to urban heat islands
608   across three cities. *Proc Roy Soc B* 285:20180036

609   37.   Dillon, ME, Wang, G, Huey, RB 2010. Global metabolic impacts of recent climate warming.
610   *Nature* 467:704-706

611   38.   Forrest, JRK 2016. Complex responses of insect phenology to climate change. *Curr Opp Insect*
612   *Sci* 17:49-54

613   39.   Gilbert, N 1988. Control of Fecundity in Pieris rapae. V. Comparisons Between Populations. *J*
614   *Anim Ecol* 57:395-410

615   40.   Gomi, T 2007. Seasonal adaptations of the fall webworm Hyphantria cunea (Drury)
616   (Lepidoptera: Arctiidae) following its invasion of Japan. *Ecol Res* 22:855-861

617   41.   Gomi, T, Nagasaka, M, Fukuda, T, Hagihara, H 2007. Shifting of the life cycle and life-history
618   traits of the fall webworm in relation to climate change. *Entomol Exp Appl* 125:179-184

619   42.   Goto, SG 2013. Roles of circadian clock genes in insect photoperiodism. *Ent Sci* 16:1-16

620   43.   Gotthard, K, Nylin, S, Wiklund, C 1994. Adaptive variation in growth rate - life history costs
621   and consequences in the speckled wood butterfly, pararge aegeria. *Oecologia* 99:281-289

622   44.   Grether, G 2005. Environmental Change, Phenotypic Plasticity, and Genetic Compensation. *Am*
623   *Nat* 166:E115-E123

45.   Halsch, CA, Shapiro, AM, Fordyce, JA, Nice, CC, Thorne, JH et al. 2021. Insects and recent
climate change. *P Natl Acad Sci USA* 118:e2002543117
46.   Hangartner, S, Hoffmann, AA 2016. Evolutionary potential of multiple measures of upper
thermal tolerance in Drosophila melanogaster. *Funct Ecol* 30:442-452
47.   Hansen, TF, Pélabon, C, Houle, D 2011. Heritability is not Evolvability. *Evol Biol* 38:258-277
48.   Harvey, JA, Heinen, R, Gols, R, Thakur, MP 2020. Climate change-mediated temperature
extremes and insects: From outbreaks to breakdowns. *Glob. Change Biol.* 26:6685-6701
49.   Harvey, JA, Tougeron, K, Gols, R, Heinen, R, Abarca, M et al. 2023. Scientists' warning on
climate change and insects. *Ecol Mono* 93:e1553
50.   Hashimoto, K-i, Iijima, K, Ogawa, K-i 2008. Geographic Variation in Photoperiodic Response
for the Induction of Pupal Diapause in the White Cabbage Butterfly, *Pieris rapae crucivora*
Boisuduval (Lepidoptera: Pieridae). *Jap J Appl Ent Zool* 52:201-206
51.   Hoffmann, AA, Chown, SL, Clusella-Trullas, S 2013. Upper thermal limits in terrestrial
ectotherms: how constrained are they. *Funct Ecol* 27:934-949
52.   Hoffmann, AA, Sgrò, CM 2011. Climate change and evolutionary adaptation. *Nature* 470:479-
485
53.   Horne, CR, Hirst, AG, Atkinson, D 2015. Temperature-size responses match latitudinal-size
clines in arthropods, revealing critical differences between aquatic and terrestrial species. *Ecol
Lett* 18:327-335
54.   Hou, Y-Y, Xu, L-Z, Wu, Y, Wang, P, Shi, J-J, Zhai, B-P 2016. Geographic Variation of
Diapause and Sensitive Stages of Photoperiodic Response in Laodelphax striatellus Fallén
(Hemiptera: Delphacidae). *J Insect Sci* 16:13
55.   Hsiao, TH 1978. Host plant adaptation among geographic populations of the Colorado potato
beetle. *Entomol Exp Appl* 24:437-447
56.   Huang, L, Tang, J, Chen, C, He, H, Gao, Y, Xue, F 2020. Diapause incidence and critical day
length of Asian corn borer (Ostrinia furnacalis) populations exhibit a latitudinal cline in both
pure and hybrid strains. *J Pest Sci* 93:559-568
57.   Huey, RB, Kingsolver, J 1989. Evolution of thermal sensitivity of ectoterm performance. *Trend
Ecol Evolut* 4:131-135
58.   Huey, RB, Gilchrist, GW, Carlson, ML, Berrigan, D, Serra, §L 2000. Rapid Evolution of a
Geographic Cline in Size in an Introduced Fly. *Science* 287:308-309
59.   Hut, RA, Paolucci, S, Dor, R, Kyriacou, CP, Daan, S 2013. Latitudinal clines: an evolutionary
view on biological rhythms. *Proc Roy Soc B* 280:20130433
60.   Ittonen, M, Hagelin, A, Wiklund, C, Gotthard, K 2022. Local adaptation to seasonal cues at the
fronts of two parallel, climate-induced butterfly range expansions. *Ecol Lett* 25:2022-2033
61.   Ittonen, M, Roberts, KT, Lehmann, P, Gotthard, K 2023. A range-expanding butterfly is
susceptible to cold and long winters but shows no signs of local adaptation to winter conditions.
Funct Ecol 37:3064-3078
62.   Izzo, VM, Hawthorne, DJ, Chen, YH 2014. Geographic variation in winter hardiness of a
common agricultural pest, Leptinotarsa decemlineata, the Colorado potato beetle. *Evol Ecol*
28:505-520
63.   James, AC, Azevedo, RB, Partridge, L 1995. Cellular basis and developmental timing in a size
cline of Drosophila melanogaster. *Genetics* 140:659-666
64.   Jin, M, Peng, Y, Peng, J, Yu, S, Wu, C et al. 2024. A supergene controls facultative diapause in
the crop pest Helicoverpa armigera. *Cell Rep* 43
65.   Johansson, F, Orizaola, G, Nilsson-Örtman, V 2020. Temperate insects with narrow seasonal
activity periods can be as vulnerable to climate change as tropical insect species. *Sci Rep*
10:8822
66.   Jordan, RG, Bradshaw, WE 1978. Geographic Variation in the Photoperiodic Response of the
Western Tree-Hole Mosquito, Aedes sierrensis. *Ann Entomol Soc Am* 71:487-490
67.   Jørgensen, LB, Ørsted, M, Malte, H, Wang, T, Overgaard, J 2022. Extreme escalation of heat
failure rates in ectotherms with global warming. *Nature* 611:93-98
68.   Joschinski, J, Bonte, D 2021. Diapause and bet-hedging strategies in insects: a meta-analysis of
reaction norm shapes. *Oikos* 130:1240-1250

69.   Kauranen, H, Kinnunen, J, Hiillos, A-L, Lankinen, P, Hopkins, D et al. 2019. Selection for
679   reproduction under short photoperiods changes diapause-associated traits and induces
680   widespread genomic divergence. *J Exp Biol* 222:jeb205831

681   70.   Kellermann, V, van Heerwaarden, B 2019. Terrestrial insects and climate change: adaptive
682   responses in key traits. *Physiol. Entomol.* 44:99-115

683   71.   Kimura, MT 1988. Adaptations to Temperate Climates and Evolution of Overwintering
684   Strategies in the Drosophila melanogaster Species Group. *Evolution* 42:1288-1297

685   72.   Kimura, MT 2004. Cold and heat tolerance of drosophilid flies with reference to their latitudinal
686   distributions. *Oecologia* 140:442-449

687   73.   Kivelä, SM, Välimäki, P, Carrasco, D, Mäenpää, MI, Oksanen, J 2011. Latitudinal insect body
688   size clines revisited: a critical evaluation of the saw-tooth model. *J Anim Ecol* 80:1184-1195

689   74.   Kostal, V 2006. Eco-physiological phases of insect diapause. *J Insect Physiol* 52:113-127

690   75.   Kozak, GM, Wadsworth, CB, Kahne, SC, Bogdanowicz, SM, Harrison, RG et al. 2019.
691   Genomic Basis of Circannual Rhythm in the European Corn Borer Moth. *Curr Biol* 29:3501-
692   3509.e5

693   76.   Kurota, H, Shimada, M 2003. Geographical variation in photoperiodic induction of larval
694   diapause in the bruchid beetle, Bruchidius dorsalis: polymorphism in overwintering stages.
695   *Entomol Exp Appl* 107:11-18

696   77.   Lehmann, P, Ammunét, T, Barton, M, Battisti, A, Eigenbrode, SD et al. 2020. Complex
697   responses of global insect pests to climate warming. *Front Ecol Environ* 18:141-150

698   78.   Lehmann, P, Piiroinen, S, Lyytinen, A, Lindström, L 2015. Responses in metabolic rate to
699   changes in temperature in diapausing Colorado potato beetle eptinotarsa decemlineata from
700   three European populations. *Physiol. Entomol.* 40:123-130

701   79.   Levins, R 1969. Thermal Acclimation and Heat Resistance in Drosophila Species. *Am Nat*
702   103:483-499

703   80.   Lindestad, O, Nylin, S, Wheat, CW, Gotthard, K 2024. Testing for variation in photoperiodic
704   plasticity in a butterfly: Inconsistent effects of circadian genes between geographic scales. *Ecol*
705   *Evol* 14:e11713

706   81.   Lindestad, O, Wheat, CW, Nylin, S, Gotthard, K 2019. Local adaptation of photoperiodic
707   plasticity maintains life cycle variation within latitudes in a butterfly. *Ecology* 100:e02550

708   82.   Logan, ML, Cox, CL 2020. Genetic Constraints, Transcriptome Plasticity, and the Evolutionary
709   Response to Climate Change. *Front Gene* 11

710   83.   Lushai, G, Hardie, J, Harrington, R 1996. Diapause termination and egg hatch in the bird cherry
711   aphid, Rhopalosiphum padi. *Entomol Exp Appl* 81:113-115

712   84.   Macgregor, CJ, Thomas, CD, Roy, DB, Beaumont, MA, Bell, JR et al. 2019. Climate-induced
713   phenology shifts linked to range expansions in species with multiple reproductive cycles per
714   year. *Nat Comm* 10:4455

715   85.   Malusare, SP, Zilio, G, Fronhofer, EA 2023. Evolution of thermal performance curves: A meta-
716   analysis of selection experiments. *J Evol Biol* 36:15-28

717   86.   Martin, RA, Chick, LD, Garvin, ML, Diamond, SE 2021. In a nutshell, a reciprocal transplant
718   experiment reveals local adaptation and fitness trade-offs in response to urban evolution in an
719   acorn-dwelling ant. *Evolution* 75:876-887

720   87.   Masaki, S 1967. Geographic Variation and Climatic Adaptation in a Field Cricket (Orthoptera:
721   Gryllidae). *Evolution* 21:725-741

722   88.   McNamara, JM, Houston, AI 1996. State-dependent life histories. *Nature* 380:215-221

723   89.   Merckx, T, Nielsen, ME, Heliölä, J, Kuussaari, M, Pettersson, LB et al. 2021. Urbanization
724   extends flight phenology and leads to local adaptation of seasonal plasticity in Lepidoptera.
725   *Proc Natl Acad Sci USA* 118:e2106006118

726   90.   Merckx, T, Nielsen, ME, Kankaanpää, T, Kadlec, T, Yazdanian, M, Kivelä, SM 2024.
727   Continent-wide parallel urban evolution of increased heat tolerance in a common moth. *Evol*
728   *Appl* 17:e13636

729   91.   Mori, A, Romero-Severson, J, Severson, DW 2007. Genetic basis for reproductive diapause is
730   correlated with life history traits within the Culex pipiens complex. *Insect Mol Biol* 16:515-524

92. Murata, Y, Ideo, S, Watada, M, Mitsui, H, Kimura, MT 2009. Genetic and physiological
variation among sexual and parthenogenetic populations of Asobara japonica (Hymenoptera:
Braconidae), a larval parasitoid of drosophilid flies. *Eur. J. Entomol.* 106:171-178
93. Mushegian, AA, Neupane, N, Batz, Z, Mogi, M, Tuno, N et al. 2021. Ecological mechanism of
climate-mediated selection in a rapidly evolving invasive species. *Ecol Lett* 24:698-707
94. Musolin, DL, Tougou, D, Fujisaki, K 2011. Photoperiodic response in the subtropical and
warm-temperate zone populations of the southern green stink bug Nezara viridula: why does it
not fit the common latitudinal trend. *Physiol Entomol* 36:379-384
95. Nechols, JR, Tauber, MJ, Tauber, CA 1987. Geographical variability in ecophysiological traits
controlling dormancy in Chrysopa oculata (Neuroptera: Chrysopidae). *J Insect Physiol* 33:627-
633
96. Nielsen, ME, Kingsolver, JG 2020. Compensating for climate change–induced cue-environment
mismatches: evidence for contemporary evolution of a photoperiodic reaction norm in Colias
butterflies. *Ecol Lett* 23:1129-1136
97. Nielsen, ME, Nylin, S, Wiklund, C, Gotthard, K 2023. Evolution of butterfly seasonal plasticity
driven by climate change varies across life stages. *Ecol Lett* 26:1548-1558
98. Noda, H 1992. Geographic-Variation of Nymphal Diapause in the Small Brown Planthopper in
Japan. *Jap Agri Res Q* 26:124-129
99. Nylin, S, Gotthard, K 1998. Plasticity in life history traits. *Annu Rev Entomol* 43:63-83
100. Nylin, S, Svärd, L 1990. Latitudinal patterns in the size of European butterflies. *Holarct Ecol*
101. Parker, AL, Kingsolver, JG 2023. Population divergence in nutrient-temperature interactions in
Pieris rapae. *Front Insect Sci* 3
102. Parmesan, C 2006. Ecological and evolutionary responses to recent climate change. *Annu Rev*
Ecol Syst 37:637-669
103. Parmesan, C, Ryrholm, N, Stefanescu, C, Hill, JK, Thomas, CD et al. 1999. Poleward shifts in
geographical ranges of butterfly species associated with regional warming. *Nature* 399:579-583
104. Parmesan, C, Morecroft, MD, Trisurat, Y, Adrian, R, Anshari, GZ et al. 2022. Terrestrial and
Freshwater Ecosystems and Their Services. In *Climate Change 2022: Impacts, Adaptation and*
Vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the
Intergovernmental Panel on Climate Change, eds. Pörtner, H-O, Roberts, DC, Tignor, M,
Poloczanska, ES, Mintenbeck, K et al., pp. 197-377. Cambridge, UK and New York, NY, USA:
Cambridge University Press,
105. Parmesan, C, Yohe, G 2003. A globally coherent fingerprint of climate change impacts across
natural systems. *Nature* 421:37-42
106. Pegoraro, M, Zonato, V, Tyler, ER, Fedele, G, Kyriacou, CP, Tauber, E 2017. Geographical
analysis of diapause inducibility in European Drosophila melanogaster populations. *J Insect*
Physiol 98:238-244
107. Peng, X, Song, C-M, Wang, K, Chen, M-H 2017. Geographical Variations in the Life Histories
of Rhopalosiphum padi (Hemiptera: Aphididae) in China. *J Econ Entomol* 110:961-970
108. Pruisscher, P, Nylin, S, Wheat, CW, Gotthard, K 2021. A region of the sex chromosome
associated with population differences in diapause induction contains highly divergent alleles at
clock genes. *Evolution* 75:490-500
109. Pruisscher, P, Nylin, S, Gotthard, K, Wheat, CW 2018. Genetic variation underlying local
adaptation of diapause induction along a cline in a butterfly. *Mol Ecol* 27:3613-3626
110. Ragland, GJ, Kingsolver, JG 2008. The effect of fluctuating temperatures on ectotherm life-
history traits: comparisons among geographic populations of Wyeomyia smithii. *Evol Ecol Res*
10:29-44
111. Ragland, GJ, Armbruster, PA, Meuti, ME 2019. Evolutionary and functional genetics of insect
diapause: a call for greater integration. *Curr Opin Insect Sci* 36:74-81
112. Ragland, GJ, Kingsolver, JG 2008. EVOLUTION OF THERMOTOLERANCE IN
SEASONAL ENVIRONMENTS: THE EFFECTS OF ANNUAL TEMPERATURE
VARIATION AND LIFE-HISTORY TIMING IN WYEOMYIA SMITHII. *Evolution* 62:1345-
1357

113. Reznik, SY, Dolgovskaya, MY, Ovchinnikov, AN, Belyakova, NA 2015. Weak photoperiodic
response facilitates the biological invasion of the harlequin ladybird armonia axyridis (Pallas)
(Coleoptera: Coccinellidae). *J. Appl. Entomol.* 139:241-249
114. Roff, DA 1980. Optimizing development time in a seasonal environment: the 'ups and downs'
of clinal variation. *Oecologia* 45:202-208
115. Roff, DA 1992. *The Evolution of Life Histories* 535 New York:Chapman & Hall
116. Sadakiyo, S, Ishihara, M 2011. Rapid seasonal adaptation of an alien bruchid after introduction:
geographic variation in life cycle synchronization and critical photoperiod for diapause
induction. *Entomol Exp Appl* 140:69-76
117. Sadakiyo, S, Ishihara, M 2012. The role of host seed size in mediating a latitudinal body size
cline in an introduced bruchid beetle in Japan. *Oikos* 121:1231-1238
118. Sánchez-Bayo, F, Wyckhuys, KAG 2019. Worldwide decline of the entomofauna: A review of
its drivers. *Biol Conserv* 232:8-27
119. Scharf, I, Bauerfeind, SS, Blanckenhorn, WU, Sch 2010. Effects of maternal and offspring
environmental conditions on growth, development and diapause in latitudinal yellow dung fly
populations. *Clim Res* 43:115-125
120. Schmidt, PS, Paaby, AB 2008. Reproductive Diapause and Life-History Clines in North
American Populations of Drosophila melanogaster. *Evolution* 62:1204-1215
121. Seiter, S, Kingsolver, J 2013. Environmental determinants of population divergence in
life☐history traits for an invasive species: climate, seasonality and natural enemies. *J Evol Biol*
26:1634-1645
122. SgrÒ, CM, Overgaard, J, Kristensen, TN, Mitchell, KA, Cockerell, FE, Hoffmann, AA 2010. A
comprehensive assessment of geographic variation in heat tolerance and hardening capacity in
populations of Drosophila melanogaster from eastern Australia. *J Evol Biol* 23:2484-2493
123. Shelomi, M, Associate, ETD, Editor: MA 2012. Where Are We Now? Bergmann's Rule Sensu
Lato in Insects. *Am Nat* 180:511-519
124. Shoshan, AB, Pindeler, U, Wheat, CW, Gotthard, K 2025. Repeated evolution of photoperiodic
plasticity by different genetic architectures during recurrent colonizations in a butterfly. *Proc*
Roy Soc B 292:20242195
125. Siemers, I, von Schmalensee, L, Ziesemer, S, Gotthard, K, Lehmann, P 2024. Predictable local
adaptation in butterfly photoperiodism but not thermal performance along a latitudinal cline.
Ecosphere 15:e4842
126. Stearns, SC 1992. *The Evolution of Life Histories* 249 Oxford:Oxford University Press
127. Sunday, J, Bennett, JM, Calosi, P, Clusella-Trullas, S, Gravel, S et al. 2019. Thermal tolerance
patterns across latitude and elevation. *Philos Trans R Soc Lond B* 374:20190036
128. Takeda, M, Chippendale, GM 1982. Environmental and genetic control of the larval diapause of
the Southwestern corn borer, Diatraea grandiosella. *Physiol Entomol* 7:99-110
129. Tanaka, K, Murata, K 2017. Genetic Basis Underlying Rapid Evolution of an Introduced Insect
Ophraella communa (Coleoptera: Chrysomelidae): Heritability of Photoperiodic Response.
Environ Entomol 46:167-173
130. Tauber, E, Zordan, M, Sandrelli, F, Pegoraro, M, Osterwalder, N et al. 2007. Natural Selection
Favors a Newly Derived *timeless* Allele in *Drosophila melanogaster*. *Science* 316:1895
131. Tauber, MJ, Tauber, CA, Masaki, S 1986. *Seasonal Adaptations of Insects.* Oxford:Oxford
University Press
132. Terao, M, Tokuda, M, Shintani, Y 2021. Geographic Variation in Photoperiodic Response for
Induction of Pseudopupal Diapause in Epicauta gorhami (Coleoptera: Meloidae). *Environ*
Entomol 50:1145-1150
133. Thackeray, SJ, Henrys, PA, Hemming, D, Bell, JR, Botham, MS et al. 2016. Phenological
sensitivity to climate across taxa and trophic levels. *Nature* 535:241-245
134. Thomas, CD, Cameron, A, Green, RE, Bakkenes, M, Beaumont, LJ et al. 2004. Extinction risk
from climate change. *Natue* 427:145-148
135. Trotta, V, Calboli, FCF, Ziosi, M, Guerra, D, Pezzoli, MC et al. 2006. Thermal plasticity in
Drosophila melanogaster: A comparison of geographic populations. *BMC Evol Biol* 6:67

136. Urbanski, J, Mogi, M, O'Donnell, D, DeCotiis, M, Toma, T, Armbruster, P 2012. Rapid
Adaptive Evolution of Photoperiodic Response during Invasion and Range Expansion across a
Climatic Gradient. *Am Nat* 179:490-500
137. van Asch, M, Salis, L, Holleman, LJM, van Lith, B, Visser, ME 2013. Evolutionary response of
the egg hatching date of a herbivorous insect under climate change. *Nat Clim Chan* 3:244-248
138. van der Have, TM, deJong, G 1996. Adult size in ectotherms: Temperature effects on growth
and differentiation. *J Theo Biol* 183:329-340
139. van Klink, R, Bowler, DE, Gongalsky, KB, Swengel, AB, Gentile, A, Chase, JM 2020. Meta-
analysis reveals declines in terrestrial but increases in freshwater insect abundances. *Science*
368:417-420
140. Vesala, L, Hoikkala, A 2011. Effects of photoperiodically induced reproductive diapause and
cold hardening on the cold tolerance of Drosophila montana. *J Insect Physiol* 57:46-51
141. von Schmalensee, L, Caillault, P, Gunnarsdóttir, KH, Gotthard, K, Lehmann, P 2023. Seasonal
specialization drives divergent population dynamics in two closely related butterflies. *Nat*
Comm 14:3663
142. von Schmalensee, L, Süess, P, Roberts, KT, Gotthard, K, Lehmann, P 2024. A quantitative
model of temperature-dependent diapause progression. *P Natl Acad Sci USA* 121:e2407057121
143. Wagner, DL, Grames, EM, Forister, ML, Berenbaum, MR, Stopak, D 2021. Insect decline in
the Anthropocene: Death by a thousand cuts. *P Natl Acad Sci USA* 118:e2023989118
144. Walters, R, Hassal, M 2006. The temeprature-size rule in ecotherms: may a general explanation
exist after all? *Am Nat* 167:510-523
145. Wang, X-P, Yang, Q-S, Dalin, P, Zhou, X-M, Luo, Z-W, Lei, C-L 2012. Geographic variation
in photoperiodic diapause induction and diapause intensity in Sericinus montelus
(Lepidoptera: Papilionidae). *Insect Sci* 19:295-302
146. Wang, X-M, Yang, X, Zang, L-S, Wang, Z, Ruan, C-C, Liu, X-J 2017. Effect of geographic
variation on biology and cold tolerance of Harmonia axyridis in China. *Entomol Gener* 36:239-
250
147. Wiberg, RAW, Tyukmaeva, V, Hoikkala, A, Ritchie, MG, Kankare, M 2021. Cold adaptation
drives population genomic divergence in the ecological specialist, Drosophila montana. *Mol*
Ecol 30:3783-3796
148. Williams, CB 1957. Insect Migration. *Annu Rev Entomol* 2:163-180
149. Wipking, W 1988. Repeated larval diapause and diapause-free development in geographic
strains of the burnet moth Zygaena trifolii Esp. (Insecta, Lepidoptera). *Oecologia* 77:557-564
150. Xiao, L, He, H-M, Huang, L-L, Geng, T, Fu, S, Xue, F-S 2016. Variation of life-history traits of
the Asian corn borer, Ostrinia furnacalis in relation to temperature and geographical latitude.
Ecol Evol 6:5129-5143
151. Yoshio, M, Ishii, M 1998. Geographical variation of pupal diapause in the great mormon
butterfly, Papilio memnon L. (Lepidoptera : Papilionidae), in Western Japan. *Appl Entomol*
Zool 33:281-288
152. Yoshio, M, Ishii, M 2001. Relationship between cold hardiness and northward invasion in the
great mormon butterfly, *Papilio memnon* L. (Lepidoptera: Papilionidae) in Japan. *Appl Entomol*
Zool 36:329-335
153. Zani, PA, Swanson, SET, Corbin, D, Cohnstaedt, LW, Agotsch, MD et al. 2005.
GEOGRAPHIC VARIATION IN TOLERANCE OF TRANSIENT THERMAL STRESS IN
THE MOSQUITO WYEOMYIA SMITHII. *Ecology* 86:1206-1211
154. Zheng, X-L, Yang, Q-S, Hu, Y-W, Lei, C-L, Wang, X-P 2015. Latitudinal variation of
morphological characteristics in the swallowtail ericinus montelus Gray, 1798 (Lepidoptera:
Papilionidae). *Acta Zool* 96:242-252
885
886