Life history evolution of insects in response to climate variation:

seasonal timing versus thermal physiology.

Karl Gotthard[1,2]* (ORCID: https://orcid.org/0000-0002-4560-6271)

David Berger[3]* (ORCID: https://orcid.org/0000-0003-0196-6109)

Patrick Rohner[4]* (ORCID: https://orcid.org/0000-0002-9840-1050)

Affiliations:

1. *Department of Zoology, Stockholm University, 106 91 Stockholm, Sweden.*

2. *The Bolin Centre for Climate Research, Stockholm University, 106 91 Stockholm, Sweden.*

3. *Department of Ecology and Genetics, Uppsala University, Uppsala, 752 36 Sweden.*

4. *Department of Ecology, Behavior, and Evolution, University of California San Diego, La Jolla, CA 92093, USA.*

*All authors contributed equally

Abstract

Climate adaptation in insects can proceed via responses in life history traits and their thermal
plasticity, and through phenological shifts mediated by responses to photoperiodic cues
("photoperiodism"). While, experimental studies demonstrate evolutionary potential for both
modes of adaptation, it remains unclear how evolution will unfold in natural populations,
limiting our ability to predict how insects will respond to climate change. Here we review the
literature and perform an analysis of published studies revealing that photoperiodism for
diapause induction evolves predictably along latitude, with high-latitude populations entering
diapause earlier. In contrast, although a few species showed clinal variation in life history and
thermal plasticity, the direction of these clines were not consistent across taxa. This suggests
that while insect life history and physiological adaptation to temperature can evolve,
phenological shifts via evolution of photoperiodism is likely to be a more common and
predictable response to future climate change.

Introduction

Ecological and evolutionary processes influenced by climate variation are widely studied due to the urgent need to understand the impact of global climate change on biodiversity (3, 54, 114, 117, 148, 149, 153, 161). It is well established that global warming has already had strong ecological effects on biological communities by affecting the distribution, abundance, and phenology (seasonal timing) of many organisms, including insects (28, 41, 51, 53, 95, 115, 116, 148, 157). However, the mechanisms by which insects respond and adapt to climate change is less well understood (3, 23, 45, 58, 78). This presents a challenge not only for assessing future extinction risk of focal species, but also for predicting stability of species interactions and ecosystems as functions of alternative evolutionary strategies in response to a warming world (6, 24, 88). For instance, while changes in climate can lead to the evolution of physiological responses that mitigate environmental stress (45, 58), the evolution of phenology might offer an alternate route to adaptation (17, 20, 76) (Figure 1). While physiological and phenological responses may both facilitate adaptation, their effects on insect growth rates and distribution ranges are likely to differ. Here, we review the literature on insect life history evolution in response to variation in climate, with a particular focus on work that has explored genetic responses in both thermal plasticity and phenology via the use of photoperiodic cues (from hereon: "*photoperiodism*").

Physiological temperature adaptation in life-history traits
Life history theory sees the scheduling of life cycle events such as growth, maturation and reproduction as the result of strategic decisions made over an organism's life (99, 111, 128, 140). Spatiotemporal variation in climate variables such as temperature, precipitation, and humidity are major sources of natural selection on life history strategies in general, and maybe in particular for ectotherms like insects (3, 23, 45, 58, 66, 111, 146). Virtually all physiological processes of insects are at least somewhat dependent on temperature, leading to abundant phenotypic plasticity in life history responses to ambient temperatures (3, 23, 111). For instance, decreased temperature often leads to plastic increases in adult body size in arthropods (the so-called temperature-size rule) (5, 59) and prolongs egg-to-adult development time (3, 23, 111). Nevertheless, theory predicts that an organism's range of tolerated and preferred temperatures should evolve to match those experienced in its environment (3, 63), and indeed, there are often evolved species differences in the degree to which species are affected by temperature (3, 23, 111). Moreover, there is often noticeable genetic variation within species, indicating that thermal plasticity can evolve, sometimes even on ecological timescales (12, 52, 93). This suggests that natural selection may alter temperature dependence of important physiological and life history traits (3, 12, 38, 52, 97, 159), offering an avenue to adapt to climate change (Figure 1C-E).

Seasonal adaptation in life-history via photoperiodism
Most organisms experience some type of seasonal variation in climate that also drive associated variation in biotic resources and interactions (22, 36, 146, 150). Seasonality is consequently a highly predictable form of temporal variation in climate and other ecological conditions (66, 146). Adaptation to seasonality requires that organisms can exploit the favorable seasons, avoid or mitigate the effects of the unfavorable seasons, and adequately time the transitions between respective stages in their life cycle (22). This is especially important for insects because most species use some kind of dormancy to withstand harsh conditions (36, 146, 150) (although there are notable examples of insects that migrate long distances to evade hazardous conditions (29, 146, 169)). In temperate insects, the typical adaptation to mitigate the effects of the unfavorable winter season is to enter diapause before the onset of cold conditions (i.e., a state of dormancy that arrests or greatly prolongs

development). The individual then remains in diapause until there is no longer a risk of
developing prematurely and finally restarts development and reproduction when the thermal
conditions again become permissive (36, 83, 146, 160). Although variation in thermal
conditions and biotic resources are likely to be the major sources of selection on life cycle
timing, temperate insects have commonly evolved to use the photoperiod as a cue of seasonal
change (36, 66, 146, 150). Photoperiod provides almost noise-free information on seasonal
timing, allowing insects to appropriately schedule diapause. The sensitivity to photoperiod
typically follows a threshold function where in certain photoperiods (typically long summer
daylengths) individuals follow non-diapause (or 'direct') development, while in other
photoperiods (typically shorter late summer/autumn daylengths) they instead enter diapause
(76, 146). The inflection-point of this threshold function is referred to as the critical
daylength (CDL) and is often used as a metric for assessing genetic variation in
photoperiodism between populations or species (16, 22, 36, 76, 138, 146). This type of
photoperiodic plasticity allows a given genotype to develop without a diapause in the warm
summer months while still inducing diapause in individuals that are active later in the season.
The facultative induction of diapause allows insect populations to produce multiple
generations within a given year (multivoltinism) and genetic variation for CDL is often
strongly associated with variation in voltinism (92, 138), and the length of the permissive
season for growth and reproduction (76).
106
107

Two modes of adaptation
Temperature variation during the favorable season is likely to exert strong selection on
thermal plasticity of life history traits, whereas temperature variation at the end of the active
season will exert selection on photoperiodic regulation of diapause induction. It is important
to note that these two types of selection pressures interact with each other. On one hand,
evolved responses in photoperiodism can shield the organism from unfavorable temperatures.
On the other hand, evolved responses in thermal plasticity can in turn relax selection for
shifts in phenology. Hence, we expect these two interacting modes of adaptation to exert
correlational selection on the optimal life history strategy to mitigate climate variability
(Figure 1C-E) (23, 45, 86, 168). However, the relative significance of selection on thermal
physiology versus selection on photoperiodism is not well understood. Here we review how
insect life histories have evolved in response to climate variation across both time and space
by comparing cases where climate variation is expected to exert natural selection on life
history traits and their thermal plasticity, as well as on photoperiodism for life cycle
regulation.
123

Evolution of thermal performance and photoperiodism in selection experiments
Experiments using artificial selection or experimental evolution to study the evolution of
thermal reaction norms (i.e. thermal plasticity) have been performed in many species (3, 96).
These experiments provide a proof-of-principle of how natural selection can shape thermal
plasticity in life history. A recent meta-analysis of such experiments on ectotherms compared
thermal reaction norms of evolved populations after thermal selection, to control (or
ancestral) populations, allowing quantification of thermal adaptation (96). About a quarter of
the studies included in the meta-analysis were done on insects. The results show a general
positive response to thermal selection, where estimates of fitness in the assay temperature
increased after selection. Moreover, thermal selection in the laboratory also had the capacity
to change the shape of reaction norms so that adaptation to warmer temperatures generally
led to lower fitness at colder temperatures (96). Although these effects were generally
strongest in unicellular organisms, they still suggest that selection over relatively short

timespans has capacity to alter the thermal sensitivity of life history traits and that there may
be genetic trade-offs between performing at different thermal conditions.
139
Different types of artificial selection experiments that explore the evolutionary potential of
photoperiodism for diapause induction and duration have been performed on species from all
large insect orders (146). Typically, they show that these seasonal adaptations can change
relatively rapidly in laboratory experiments (146). For instance, artificial selection on the
photoperiodic induction of reproductive diapause in the fly *Drosophila montana*
demonstrated a rapid change in critical daylength (77). Using a baseline population derived
from a northern location in Finland, selection for eight generations where only non-
diapausing individuals were allowed to breed under continuously shorter daylengths resulted
in shorter CDL in all replicate lines compared to the control. These selected lines evolved a
response to photoperiod that is typical for more southern populations of *D. montana*.
Similarly, bidirectional artificial selection for non-diapause and diapause development led to
the evolution of photoperiodic plasticity for diapause induction in the North American
mosquito *Wyeomyia smithii* after 12 generations (18), and in the chrysomelid beetle
Ophraella communa after 5-8 generations (144).
154
155 In summary these examples demonstrate that both thermal reaction norms and photoperiodic
156 reaction norms have the potential to evolve relatively rapidly in response to directional
157 selection in the laboratory, suggesting that there is standing genetic variation present within
158 natural populations that can fuel rapid evolutionary change in the wild.
159
Life history evolution in response to recent climate change
Rising temperatures are altering patterns of seasonality and phenology — a trend expected to
be further exacerbated in the foreseeable future (3). Against the backdrop of strong declines
in insect abundances in recent decades (132, 157), there is increasing interest in the effects of
climate on insect populations and their life histories (54, 161). However, to detect and
estimate phenotypic evolution in natural populations it is necessary to measure heritable (i.e.,
'genetic') changes in traits in the same populations repeatedly (at least twice) over several
years or even decades.
168
In insects these types of studies are relatively rare as they require controlled common garden
experiments to be replicated over extended time periods. Still, there are some notable
examples, spearheaded by work on the North American Pitcher plant mosquito (*Wyeomyia*
smithii) by Bradshaw and Holtzapfel (16). The authors demonstrated that the photoperiodic
regulation of the life cycle, measured by the CDL for diapause induction in a range of
latitudinally separated populations, showed a general genetic shift towards shorter CDLs, that
are typical in more southern populations. This shift likely represents adaptive evolution in
response to a warming climate as shorter CDLs result in a longer growth season and a later
date for diapause induction. Surprisingly, this evolutionary change was detectable over only
five years of evolution in the field (16). Similar results have since been found in other insects
(44, 107, 108), indicating that changes in photoperiodism for life cycle regulation may
constitute a common evolutionary response to climate change.
181
However, there have also been documented evolutionary changes in life history traits over
time. For instance, the Yellow dung fly (*Scathophaga stercoraria*) shows genetically based
reductions in body size (13), and a study on the Winter moth (*Operophtera brumata*) found a
reduction in egg development rate (155). The latter example presumably represents a rare
demonstration of how climate change has led to the evolution of thermal plasticity of

development rate. Interestingly, this response also seems related to selection for a change in
phenology (155) as several populations in the Netherlands have evolved a reduction in egg
development rate in a range of relevant spring temperatures, most likely due to selection for
maintaining synchronization of egg hatching with the budburst of its host plant, the Oak
(Quercus robur), in the now warmer spring. Perhaps the most extreme example comes from a
recent study on fruit flies, showing evidence for rapid life history adaptation to climate
fluctuations over a single season (spring to fall), corresponding to approximately 10
overlapping fly generations (129).
195
Life history evolution in response to climate variation in space
Climate and seasonality vary greatly in space, and this variation is particularly predictable
along latitudinal gradients (66). Studies of life history evolution in insects have a long
tradition of exploring this type of latitudinal variation in climate (3, 14, 127, 137, 146). In
many cases researchers have exploited this spatial variation to explore the likely effects of
current climate change by the so-called "space-for-time" substitution (66).
202
There is a large body of literature that explores the thermal biology of life history evolution
across insect species (3). In particular variation in thermal tolerance limits (i.e., the critical
minimum and maximum temperatures, CT_{min} and CT_{max}, respectively) across species and
geographic locations are well-studied (2, 11, 57, 75, 142). Recent research has also
highlighted the importance of studying corresponding measures of thermal tolerance in
reproductive traits (40, 162), which may shed further light on adaptive potential and genetic
constraints on climate adaptation in insect life history. A general pattern across insects (2, 57)
and in ectotherms in general (11, 142) is that lower thermal limits show greater adaptive
variation across climate regions at the species-level, compared to upper thermal limits. As a
consequence, many researchers have converged on the conclusion that there may be limited
scope for rapid evolution of CT_{max} in response to a warming world (11, 57, 75). Still,
artificial selection experiments show that populations of the Fruit fly *(Drosophila*
melanogaster) harbors additive genetic variation that allows successful laboratory evolution
of increased heat tolerance, even if it may still be insufficient to keep up with the predicted
increase in future temperatures (52). Moreover, as temperatures during the hottest parts of the
year vary less across the typical geographic regions studied than do thermal minima, it is also
possible that the relatively modest differentiation in CT_{max} can in part be explained by similar
selection pressure across geographic regions (72). Studies exploring latitudinal variation in
thermal tolerance within species provide a similar, albeit less strong, pattern as the between
species comparisons (7, 9, 79, 80, 103, 125, 134, 136, 158, 167, 175). For example, within
the limited data we reviewed for our quantitative analysis (see below), we found estimates of
latitudinal variation in heat and cold tolerance for 4 and 8 dipterans, respectively. The
observed clinal variation was not very consistent among species but perhaps slightly more so
for cold tolerance, which tended to be higher in more northern populations. Interestingly,
insects that have colonized urban environments show parallel evolution of both CT_{max} and
CT_{min} (38, 97, 101), presumably due to local adaptation to the generally warmer urban
environment. Together, this suggests that even though adaptive responses to natural selection
on survival at thermal extremes do occur as insects colonize new climate zones, the rate of
evolution may often be relatively slow.
232
233 There are also a large number of insect studies that have explored the evolution of
234 development rates, body size, and their thermal plasticity across latitude (3, 14, 64, 73, 81,
235 98, 109, 112, 137, 139). However, predictions of how natural selection should affect
236 development rate, size and their temperature dependencies are not always straightforward

(15, 30, 59, 137). A common expectation is that because both mean temperatures and the length of insect growth seasons decrease with distance from the equator, natural selection should favor faster development rates (shorter development times) and thermal plasticity that maximizes growth in colder conditions in populations living at higher latitudes (81, 82, 127). This quite frequently observed pattern has been called "counter gradient variation" or "genetic compensation", as genetic adaptation across latitude opposes the direct effect of temperature on plasticity in growth and development rates (32, 33, 48, 81, 82, 90). As a consequence of the shortened time for growth and development, it can be expected that adult size of insects should become smaller when moving away from the equator (127). However, physiological models assuming that rates of growth and development have different thermal plasticity predict that there are situations when insects should grow to larger sizes at colder latitudes (156, 163). Studies of latitudinal clines in body size within insect species have indeed found evidence for both these patterns (15, 81, 137). For instance, Shelomi (2012) reviewed a large set of both inter- and intraspecific latitudinal and altitudinal clines in insect body sizes, and while there were approximately equal numbers of positive and negative size clines, the most common pattern was the absence of a size cline altogether (137). Similar mixed patterns were found in a metanalysis of within-species size clines in arthropods that was mainly based on insects (59). Interestingly, this study found strong evidence that terrestrial taxa were more likely to show reductions in size with latitude, whereas aquatic taxa typically showed the opposite pattern. This result may be explained by the difference between water and air in how oxygen limitation affects both the plasticity and evolution of growth and development. Moreover, in the terrestrial taxa the pattern of voltinism influenced the latitudinal size clines as species with one or two generations per year decreased in size with latitude, while multivoltine species instead showed a weak positive size cline with latitude. This suggests that as the length of the growth season increases towards the equator, species with strong latitudinal clines in voltinism (multivoltine) are under selection to produce additional generations, while in uni- and bivoltine species selection instead favors an increase in adult size via prolongation of development (59).

Because photoperiods change very predictably across latitudes, it is expected that natural selection should favor latitudinal variation in adaptations for seasonal timing in response to daylength. Indeed, there is a large number of examples demonstrating local adaptation in photoperiodism of insects, in particular in plasticity for diapause induction (35, 66, 76, 146). The most common pattern is that the critical daylength for diapause induction (CDL) increases with latitude across populations, which shows that more northern populations with a shorter growth season typically enter diapause earlier in year (76, 146). Latitudinal clines in CDL can evolve relatively rapidly in introduced species (10, 43, 146, 154), species colonizing urban areas (100), and in species expanding their ranges due to climate change (67).

The available evidence consequently suggests that life history traits such as development rates, body size and their thermal dependencies, as well as the photoperiodic plasticity for seasonal life cycle regulation do evolve across space and to some degree also with ongoing climate change. However, it is still unclear whether natural populations are more likely to adapt through changes in thermal plasticity, the adjustment of seasonal activity (through the evolution of photoperiodism), or both. Previous attempts to assess whether thermal- or photoperiodic adaptations are more likely to evolve in response to rapid climate change suggested that genetic changes in photoperiodism for seasonal life cycle regulation was a considerably more likely "first response" compared to thermal adaptation (17, 20). As these early reviews were based on the relatively limited set of then known examples and that

selection due to climate change is likely to be multifaceted, other researchers have argued
that this conclusion is premature (58). Unfortunately, well established examples of
evolutionary change due to recent climate change are still rare, and for insects we have not
found more than the handful of examples reviewed above. However, because of the large
number of studies exploring evolution of photoperiodism, as well as life history traits and
their thermal dependencies in response to climate variation across latitudes, it is possible to
explore this question in the spatial dimension. Below, we first summarize and integrate the
results of previously published studies to more formally quantify the amount of population
differentiation in photoperiodism across latitude. We subsequently compare this to the
corresponding latitudinal variation in life history traits and their thermal plasticity in the same
set of species.
298
299

Analysis of latitudinal clines in photoperiodism, life history, and thermal plasticity

Full details of the literature search and analyses can be found in the supplementary material.
In brief, we looked for published common garden datasets that quantified latitudinal clines in
critical daylength as well as means and thermal plasticity of the life history traits body size
and development rate. The dataset on critical daylength was based on an initial data set
generated by Joschinski and Bonte (2021) (76). We added additional data that has since been
published and had at least 2 populations and 3 photoperiod treatments. Data on life history
was collected based on an independent literature search that targeted species with
photoperiodism data, again focusing on common garden studies that reported latitudinal
clines. We calculated an index of thermal plasticity (i.e. how temperature affected a linear
response in trait means) for those studies that included at least two temperature treatments to.
Our final dataset included 308 populations from 24 insect species that had estimates of
latitudinal clines in both critical daylength and development rate and/or body size. This
included representatives of the orders Coleoptera (n = 5), Diptera (n = 5), Hemiptera (n = 3),
Lepidoptera (n = 10), as well as one species of Neuroptera, from the original publications: (1,
4, 8, 14, 18, 27, 34, 42, 47, 55, 60–62, 67, 69, 70, 74, 85, 89, 92, 94, 104–106, 110, 113, 118,
119, 123, 126, 130, 131, 133, 135, 139, 143, 147, 151, 154, 164, 165, 171–174, 176).
318
We found that the vast majority of species showed an increase in CDL with latitude,
indicating that populations closer to the poles have adapted to the generally longer summer
photoperiods at these locations (Figures 2,3). Based on critical daylength estimates, we also
calculated the mean ordinal day of the year at diapause onset at each population's location.
Average diapause onset showed a significant decrease with latitude, indicating that
populations closer to the poles generally enter diapause earlier in the year. These findings are
consistent with other studies (76) and indicate a strong and consistent evolutionary pattern of
photoperiodism evolution across insect orders.
327
In contrast, in the same set of species, we found no consistent latitudinal effect for any of the
investigated life history traits (development rate, body size and the thermal plasticity of these
traits). This finding contrasts with the notion that counter gradient adaptation in development
rate is one of the main hypotheses for the evolution of latitudinal clines in life histories of
insects. Importantly, however, there are many instances of life history clines in the data. In
fact, latitude explained a considerable fraction of the observed population differentiation
within species for mean development rate and body size (45% and 48%, respectively) (Figure
2B). For instance, the speckled wood butterfly (*Pararge aegeria*) and southwestern corn
borer (*Diatraea grandiosella*) showed rather strong genetic decreases in development rate

with latitude (47, 92, 143). However, patterns were inconsistent, and sometimes in opposing directions, across species (Figure 3) and we even found a couple of instances of inconsistent patterns within species. For example, in the Colorado potato beetle (*Leptinotarsa decemlineata*), development rate increases with latitude in European populations but decreased towards the pole in North America (69, 89). As latitudinal clines in body size often are considered to be results of a trade-off with development time (potentially also affected by shifts in voltinism), which showed no consistent latitudinal pattern here, the idiosyncratic responses in body size clines are perhaps less surprising. We also found no general pattern in thermal plasticity of body size or developmental rate. Although sample sizes were limited for those comparisons, most estimates of the latitudinal effect on thermal plasticity for individual species hoover around zero (Figures 2B and 3C, F). Taken together, these findings suggest that the mechanisms underpinning seasonal timing evolve predictably across orders. In contrast, while life history traits often do evolve, the patterns are not consistent across insects and remain hard to predict.

An important caveat is that there is still a relatively small number of species for which we were able to find both photoperiodism and life history data and we were not able to generate replicated measurements within species (except for a few species measured from more than one continent). This limits our ability to differentiate between evolved differences among species and spurious experimental effects, despite researchers' best efforts to implement common garden rearings. However, by including only species where latitudinal variation in life history and CDL for diapause induction have been estimated, we were able to assess the relative likelihood of the two modes of adaptation in life history strategies. Although this reduced the number of species in our analysis, the results nevertheless revealed a very strong pattern in CDL and mean diapause onset and no consistent patterns in the other traits. In combination with the limited set of examples that have documented evolutionary changes due to recent climate change, it seems reasonable to assume that this higher predictability of the evolution of photoperiodism pertains also to the expected future changes in climate.

The difference in predictability of evolutionary change may have several explanations. Compared to life history traits, selection on CDL may simply be stronger and less dependent on other ecological circumstances that differ among species and geographic locations. For instance, variation in behavioral thermoregulation may affect the exposure to temperature variation, and moderate selection on thermal plasticity of life history traits (23). Along the same lines, and perhaps more pertinent to our analysis, it has been suggested that evolutionary changes in seasonal life cycle regulation may mitigate selection on thermal sensitivity (23) (Figure 1). This type of correlational selection would arise if selection typically favors a seasonal regulation that consistently exposes the active part of the life cycle to the time of the year that is most thermally similar at different locations, thereby reducing the difference in thermal selection across space; after all this is one of the main adaptive functions of insect diapause in general.

It is also possible that, compared to CDL, the amount of standing genetic variation is typically lower for thermal plasticity in life history traits, which would limit the response to selection. However, some studies indicate noticeable genetic variation for thermal plasticity of life history traits (26, 93), suggesting that it is unlikely to be the only explanation for the observed patterns. It is also possible that, even though thermal plasticity of life history traits harbor significant genetic variation, general differences in the genetic architecture of these different traits affect their evolvability. For instance, selection on life history traits and their thermal plasticity are likely to affect many genes with strong pleiotropic effects, which could

lead to correlated evolution in other traits with maladaptive effects in the natural
environment. For example, artificial selection causing adaptation to higher temperatures in a
range of ectotherms led to correlated evolution of maladaptation at colder temperatures (96).
As temperature has such a multifaceted influence on virtually all aspects of insect biology it
is perhaps not surprising if a genetic change in one aspect of this temperature dependence
also affects the temperature dependence of other important physiological processes. In a
natural environment with highly variable thermal conditions such correlated effects are likely
to lower the rate of evolution in response to a higher (or lower) mean temperature. In
contrast, genes affecting photoperiodism may have less systemic effects and therefore be
subject to less pleiotropic constraints. Indeed, several studies have found major effect loci
that underpin intraspecific variation in diapause induction (46, 71, 84, 121, 122, 124, 145).
However, other studies suggest that photoperiodism is genetically highly complex (19, 21,
91, 124, 138) and some suggest that diapause induction and life history might be genetically
correlated through linkage or pleiotropy (102).
401
Poleward range expansions are one of the most common effects of ongoing climate change
(28, 115). Considering the general process that (re-)distributes insects across latitudes may
provide yet another explanation for why genetic clines in CDL are more predictable than any
of the other traits examined here. When insects colonize new locations across latitudes it is
likely that their existing thermal adaptations are strong determinants of where they can
establish successfully (68). This means that they may colonize new locations when thermal
conditions have changed enough to allow positive population growth. If so, selection for
rapid thermal adaptation may initially be comparatively weak in these newly colonized
locations. In contrast, all expansions across latitudes will inevitably mean that they are
exposed to a novel regime of photoperiodic variation and if a non-optimal CDL does not
preclude establishment, the novel environment will consistently exert selection on
photoperiodic plasticity and CDL (67). The latitudinal snapshots represented in our collated
data may be a reflection of this process.
415

Conclusions

416
Several studies indicate that insects can relatively rapidly evolve in response to climate
change, either through the evolution of thermal physiology or shifts in seasonal activity. Our
review suggests that changes in photoperiodism, that is, the timing of life history transitions
via use of photoperiod cues, is a predictable way in which various insect species adapt to
climate variability. In contrast, while changes in life history traits often evolve, there is little
support for a general pattern that holds broadly across species. In the context of climate
adaptation, evolution of seasonal timing consequently emerges as a more predictable factor
than physiological adaptations to thermal conditions. As such, our updated review and more
formal analysis of life history evolution in response to climate variation gives credence to the
statement of Bradshaw and Holzapfel (2008): it's seasonal timing that matters.
427
Because the evolution of photoperiodism (or lack thereof) can lead to phenological
mismatches and affect species interactions and ecosystem functioning, these evolutionary
changes may have major impacts on the structure and resilience of natural systems. Our
review thus offers some hope regarding our ability to predict shifts in insect phenology and
its associated consequences in future climates. However, it should be noted that obligate
univoltine as well as tropical insects are less likely to rely on photoperiodic plasticity for
seasonal life cycle regulation and diapause induction. These insects may instead be more
dependent on thermal variation also for life cycle regulation and potentially be more likely to
show predictable clines in life history and thermal plasticity in relation to spatial variation in

climate (49, 50). Moreover, despite relatively rapid evolution of seasonal life cycle regulation, many insect populations will ultimately face significantly hotter conditions during important parts of their life cycles. Notable in this regard, our analysis is heavily biased towards terrestrial species with good capacity to disperse, whereas aquatic taxa and those with generally more stationary lifestyles are not represented. Such taxa may be particularly vulnerable to climate change (56). The comparatively slow evolution and limited scope for adaptive acclimation of upper thermal limits shown in many insects is therefore clearly worrying (11, 57, 75, 120, 166). It should also be noted that the analyzed data suffers from the general sampling bias in studies of biodiversity, with a strong underrepresentation of not only the tropics but also the southern hemisphere (25, 65) (Fig. 2A). Because the distribution of biodiversity as well as current and future climate niches are geographically heterogenous, future forecasting will need to account for the fact that climate change itself is expected to progress at different pace across the globe (31, 39, 170).

Predicting how climate change will impact insect life history is crucial to mitigating both biodiversity loss and pest outbreaks (37, 53, 88). Current statistical projection models indicate that climate warming will lead to deterministic changes in insect life history via temperature's effects on biological rates (39) leading to range expansions and increases in abundance in temperate regions, but population decline in the tropics (37). However, these projections may have poor accuracy as they do not factor in evolutionary responses. For example, a recent study on the beetle pest *Callosobruchus maculatus* showed that less than a decade of life history adaptation at hot temperature in the laboratory almost doubled its predicted costs to agriculture compared to simple statistical projections (24). The unpredictability of observed life history evolution in the species analyzed here adds to the notion that forecasting responses in insect growth rates and distribution ranges will be a significant challenge (87). Our review highlights four critical areas for future work that may help tackle this problem. First, the number of species in which clines in diapause induction, life history, and various physiological responses have been tested in a common garden experiment remains small, taxonomically biased, and rarely replicated within species. Future research should aim at closing this gap. One possibility here is also to study variation along altitude, which typically replicates climate variation at a smaller spatial scale compared to latitudes. For instance, a recent study of altitudinal variation among populations of burying beetles found that while photoperiodism showed systematic population differentiation, there was no similar effect on thermal plasticity in life history (152), much in agreement with the general pattern in our multispecies analysis across latitudes. Second, there is a need for more studies that investigate how natural populations adapt in situ to changing climates (13, 16, 44, 107, 108, 129, 155). Although these longitudinal studies suffer from a variety of experimental challenges, they provide important insights into climate-induced life history evolution in the wild (141). Third, studying the genetic architecture of photoperiodism, life history, and thermal physiology will provide critical insights into the evolvability of natural populations - at the moment it remains unclear if the number of loci and their pleiotropic relationships differ qualitatively between thermal plasticity in life history and seasonal timing traits, and how important genetic constraints are in explaining the observed patterns. Finally, studies that characterize genetic (co)variation in photoperiodism and life history traits within and between natural populations should ideally combine these estimates with measures of their fitness consequences, as this will offer insights into the role of correlational selection in driving the observed clinal patterns. It is clear that these suggested research strategies will require major effort and skill in combination with significant funding, but given the magnitude of expected effects of climate change for both natural environments and human well-being, it is likely to be a valuable investment.

Acknowledgements

This research was supported by funding from the Swedish research council (VR 2021-04258 to KG), the Swedish research council FORMAS (2022-01117 to DB). We would like to thank David Holway for helpful comments on earlier versions of this manuscript.

References

1. Aalberg Haugen, IM, Gotthard, K 2015. Diapause induction and relaxed selection on alternative developmental pathways in a butterfly. *J Anim Ecol* 84:464-472
2. Addo-Bediako, A, Chown, SL, Gaston, KJ 2000. Thermal tolerance, climatic variability and latitude. *Proc. R. Soc. Lond. B* 267:739-745
3. Angilletta, MJ 2009. *Thermal Adaptation: A theoretical and empirical synthesis* Oxford:Oxford University Press
4. Armbruster, P, Conn, JE 2006. Geographic Variation of Larval Growth in North American Aedes albopictus (Diptera: Culicidae). *Ann Entomol Soc Am* 99:1234-1243
5. Atkinson, D 1994. Temperature and organism size - a biological law for ectotherms? *Adv Eco Res* 25:1-58
6. Bale, JS, Masters, GJ, Hodkinson, ID, Awmack, C, Bezemer, TM et al. 2002. Herbivory in global climate change research: direct effects of rising temperature on insect herbivores. *Glob Chan Biol* 8:1-16
7. Baskauf, SJ, McCauley, DE 2001. Evaluation of Low Temperature Mortality as a Range-limiting Factor for the Southwestern Corn Borer (Lepidoptera: Crambidae). *Environ Entomol* 30:181-188
8. Bauerfeind, SS, Schäfer, MA, Berger, D, Blanckenhorn, WU, Fox, CW 2018. Replicated latitudinal clines in reproductive traits of European and North American yellow dung flies. *Oikos* 127:1619-1632
9. Bauerfeind, SS, Sørensen, JG, Loeschcke, V, Berger, D, Broder, ED et al. 2018. Geographic variation in responses of European yellow dung flies to thermal stress. *J Therm Biolo* 73:41-49
10. Bean, DW, Dalin, P, Dudley, TL 2012. Evolution of critical day length for diapause induction enables range expansion of Diorhabda carinulata, a biological control agent against tamarisk (Tamarix spp.). *Evol Appl* 5:511-523
11. Bennett, JM, Sunday, J, Calosi, P, Villalobos, F, Martínez, B et al. 2021. The evolution of critical thermal limits of life on Earth. *Nat Comm* 12:1198
12. Berger, D, Postma, E, Blanckenhorn, WU, Walters, RJ 2013. Quantitative Genetic Divergence And Standing Genetic (Co)Variance In Thermal Reaction Norms Along Latitude. *Evolution* 67:2385-2399
13. Blanckenhorn, WU 2015. Investigating yellow dung fly body size evolution in the field: Response to climate change. *Evolution* 69:2227-2234
14. Blanckenhorn, WU, Bauerfeind, SS, Berger, D, Davidowitz, G, Fox, CW et al. 2018. Life history traits, but not body size, vary systematically along latitudinal gradients on three continents in the widespread yellow dung fly. *Ecography* 41:2080-2091
15. Blanckenhorn, WU, Demont, M 2004. Bergmann and converse Bergmann latitudinal clines in arthropods: Two ends of a continuum? *Integr Comp Biol* 44:413-424
16. Bradshaw, WE, Holzapfel, CM 2001. Genetic shift in photoperiodic response correlated with global warming. *Proc Natl Acad Sc USA* 98:14509-14511
17. Bradshaw, WE, Holzapfel, CM 2008. Genetic response to rapid climate change: it's seasonal timing that matters. *Mol Ecol* 17:157-166
18. Bradshaw, W, Quebodeaux, M, Holzapfel, C 2003. Circadian Rhythmicity and Photoperiodism in the Pitcher-Plant Mosquito: Adaptive Response to the Photic

Environment or Correlated Response to the Seasonal Environment. *Am Nat* 161:735-748

19. Bradshaw, WE, Emerson, KJ, Catchen, JM, Cresko, WA, Holzapfel, CM 2012. Footprints in time: comparative quantitative trait loci mapping of the pitcher-plant mosquito, Wyeomyia smithii. *Proc Roy Soc B* 279:4551-4558

20. Bradshaw, WE, Holzapfel, CM 2006. Evolutionary Response to Rapid Climate Change. *Science* 312:1477-1478

21. Bradshaw, WE, Holzapfel, CM 2017. Natural Variation and Genetics of Photoperiodism in Wyeomyia smithii. *Advances in Genetics* 99: Natural Variation and Clocks:39-71

22. Bradshaw, WE, Zani, PA, Holzapfel, CM 2004. Adaptation to temperate climates. *Evolution* 58:1748-1762

23. Buckley, LB, Kingsolver, JG 2021. Evolution of Thermal Sensitivity in Changing and Variable Climates. *Annu. Rev. Ecol. Evol. Syst.* 52:563-586

24. Burc, E, Girard-Tercieux, C, Metz, M, Cazaux, E, Baur, J et al. 2025. Life-history adaptation under climate warming magnifies the agricultural footprint of a cosmopolitan insect pest. *Nat Comm* 16:827

25. Caldwell, IR, Hobbs, J-PA, Bowen, BW, Cowman, PF, DiBattista, JD et al. 2024. Global trends and biases in biodiversity conservation research. *Cell Reports Sustainability* 1

26. Carbonell, JA, Wang, Y-J, Stoks, R 2021. Evolution of cold tolerance and thermal plasticity in life history, behaviour and physiology during a poleward range expansion. *J Anim Ecol* 90:1666-1677

27. Chanthy, P, Martin, RJ, Gunning, RV, Andrew, NR 2015. Influence of temperature and humidity regimes on the developmental stages of green vegetable bug, "Nezara viridula" (l.) (Hemiptera: Pentatomidae) from inland and coastal populations in Australia. *Gen Appl Ent: J Ent Soc New South Wales* 43:37-55

28. Chen, I-C, Hill, JK, Ohlemüller, R, Roy, DB, Thomas, CD 2011. Rapid Range Shifts of Species Associated with High Levels of Climate Warming. *Science* 333:1024

29. Chowdhury, S, Zalucki, MP, Amano, T, Woodworth, BK, Venegas-Li, R, Fuller, RA 2021. Seasonal spatial dynamics of butterfly migration. *Ecol Lett* 24:1814-1823

30. Chown, SL, Gaston, KJ 2010. Body size variation in insects: a macroecological perspective. *Biol Rev* 85:139-169

31. Coelho, MTP, Barreto, E, Rangel, TF, Diniz-Filho, JAF, Wüest, RO et al. 2023. The geography of climate and the global patterns of species diversity. *Nature* 622:537-544

32. Conover, DO, Present, TMC 1990. Countergradient variation in growth rate: compensation for length of the growing season among Atlantic silversides from different latitudes. *Oecologia* 83:316-324

33. Conover, DO, Duffy, TA, Hice, LA 2009. The Covariance between Genetic and Environmental Influences across Ecological Gradients. *Annals N Y Acad Sci* 1168:100-129

34. Couper, LI, Farner, JE, Lyberger, KP, Lee, AS, Mordecai, EA 2024. Mosquito thermal tolerance is remarkably constrained across a large climatic range. *Proc Roy Soc B* 291:20232457

35. Danilevskii, AS 1965. *Photoperiodism and Seasonal Development of Insects.* London:Oliver & Boyd

36. Denlinger, DL 2022. *Insect diapause* Cambridge University Press

37. Deutsch, CA, Tewksbury, JJ, Tigchelaar, M, Battisti, DS, Merrill, SC et al. 2018. Increase in crop losses to insect pests in a warming climate. *Science* 361:916-919

38. Diamond, SE, Chick, LD, Perez, A, Strickler, SA, Martin, RA 2018. Evolution of
thermal tolerance and its fitness consequences: parallel and non-parallel responses to
urban heat islands across three cities. *Proc Roy Soc B* 285:20180036
39. Dillon, ME, Wang, G, Huey, RB 2010. Global metabolic impacts of recent climate
warming. *Nature* 467:704-706
40. Dougherty, LR, Frost, F, Maenpaa, MI, Rowe, M, Cole, BJ et al. 2024. A systematic
map of studies testing the relationship between temperature and animal reproduction.
Ecol Solut Evid 5:e12303
41. Forrest, JRK 2016. Complex responses of insect phenology to climate change. *Curr
Opp Insect Sci* 17:49-54
42. Gilbert, N 1988. Control of Fecundity in Pieris rapae. V. Comparisons Between
Populations. *J Anim Ecol* 57:395-410
43. Gomi, T 2007. Seasonal adaptations of the fall webworm Hyphantria cunea (Drury)
(Lepidoptera: Arctiidae) following its invasion of Japan. *Ecol Res* 22:855-861
44. Gomi, T, Nagasaka, M, Fukuda, T, Hagihara, H 2007. Shifting of the life cycle and life-
history traits of the fall webworm in relation to climate change. *Entomol Exp Appl*
125:179-184
45. González-Tokman, D, Córdoba-Aguilar, A, Dáttilo, W, Lira-Noriega, A, Sánchez-
Guillén, RA, Villalobos, F 2020. Insect responses to heat: physiological mechanisms,
evolution and ecological implications in a warming world. *Biol Rev* 95:802-821
46. Goto, SG 2013. Roles of circadian clock genes in insect photoperiodism. *Ent Sci* 16:1-
16
47. Gotthard, K, Nylin, S, Wiklund, C 1994. Adaptive variation in growth rate - life history
costs and consequences in the speckled wood butterfly, pararge aegeria. *Oecologia*
99:281-289
48. Grether, G 2005. Environmental Change, Phenotypic Plasticity, and Genetic
Compensation. *Am Nat* 166:E115-E123
49. Halali, S, Brakefield, PM, Brattström, O 2024. Phenotypic plasticity in tropical
butterflies is linked to climatic seasonality on a macroevolutionary scale. *Evolution*
78:1302-1316
50. Halali, S, Halali, D, Barlow, HS, Molleman, F, Kodandaramaiah, U et al. 2021.
Predictability of temporal variation in climate and the evolution of seasonal
polyphenism in tropical butterfly communities. *J. evol. Biol.* 34:1362-1375
51. Halsch, CA, Shapiro, AM, Fordyce, JA, Nice, CC, Thorne, JH et al. 2021. Insects and
recent climate change. *P Natl Acad Sci USA* 118:e2002543117
52. Hangartner, S, Hoffmann, AA 2016. Evolutionary potential of multiple measures of
upper thermal tolerance in Drosophila melanogaster. *Funct Ecol* 30:442-452
53. Harvey, JA, Heinen, R, Gols, R, Thakur, MP 2020. Climate change-mediated
temperature extremes and insects: From outbreaks to breakdowns. *Glob. Change Biol.*
26:6685-6701
54. Harvey, JA, Tougeron, K, Gols, R, Heinen, R, Abarca, M et al. 2023. Scientists'
warning on climate change and insects. *Ecol Mono* 93:e1553
55. Hashimoto, K-i, Iijima, K, Ogawa, K-i 2008. Geographic Variation in Photoperiodic
Response for the Induction of Pupal Diapause in the White Cabbage Butterfly, Pieris
rapae crucivora Boisuduval (Lepidoptera: Pieridae). *Jap J Appl Ent Zool* 52:201-206
56. Heino, J, Virkkala, R, Toivonen, H 2009. Climate change and freshwater biodiversity:
detected patterns, future trends and adaptations in northern regions. *Biol Rev* 84:39-54
57. Hoffmann, AA, Chown, SL, Clusella-Trullas, S 2013. Upper thermal limits in terrestrial
ectotherms: how constrained are they. *Funct Ecol* 27:934-949

58. Hoffmann, AA, Sgrò, CM 2011. Climate change and evolutionary adaptation. *Nature*
470:479-485
59. Horne, CR, Hirst, AG, Atkinson, D 2015. Temperature-size responses match
latitudinal-size clines in arthropods, revealing critical differences between aquatic and
terrestrial species. *Ecol Lett* 18:327-335
60. Hou, Y-Y, Xu, L-Z, Wu, Y, Wang, P, Shi, J-J, Zhai, B-P 2016. Geographic Variation of
Diapause and Sensitive Stages of Photoperiodic Response in Laodelphax striatellus
Fallén (Hemiptera: Delphacidae). *J Insect Sci* 16:13
61. Hsiao, TH 1978. Host plant adaptation among geographic populations of the Colorado
potato beetle. *Entomol Exp Appl* 24:437-447
62. Huang, L, Tang, J, Chen, C, He, H, Gao, Y, Xue, F 2020. Diapause incidence and
critical day length of Asian corn borer (Ostrinia furnacalis) populations exhibit a
latitudinal cline in both pure and hybrid strains. *J Pest Sci* 93:559-568
63. Huey, RB, Kingsolver, J 1989. Evolution of thermal sensitivity of ectoterm
performance. *Trend Ecol Evolut* 4:131-135
64. Huey, RB, Gilchrist, GW, Carlson, ML, Berrigan, D, Serra, §L 2000. Rapid Evolution
of a Geographic Cline in Size in an Introduced Fly. *Science* 287:308-309
65. Hughes, AC, Orr, MC, Ma, K, Costello, MJ, Waller, J et al. 2021. Sampling biases
shape our view of the natural world. *Ecography* 44:1259-1269
66. Hut, RA, Paolucci, S, Dor, R, Kyriacou, CP, Daan, S 2013. Latitudinal clines: an
evolutionary view on biological rhythms. *Proc Roy Soc B* 280:20130433
67. Ittonen, M, Hagelin, A, Wiklund, C, Gotthard, K 2022. Local adaptation to seasonal
cues at the fronts of two parallel, climate-induced butterfly range expansions. *Ecol Lett*
25:2022-2033
68. Ittonen, M, Roberts, KT, Lehmann, P, Gotthard, K 2023. A range-expanding butterfly is
susceptible to cold and long winters but shows no signs of local adaptation to winter
conditions. *Funct Ecol* 37:3064-3078
69. Izzo, VM, Hawthorne, DJ, Chen, YH 2014. Geographic variation in winter hardiness of
a common agricultural pest, Leptinotarsa decemlineata, the Colorado potato beetle. *Evol*
Ecol 28:505-520
70. James, AC, Azevedo, RB, Partridge, L 1995. Cellular basis and developmental timing
in a size cline of Drosophila melanogaster. *Genetics* 140:659-666
71. Jin, M, Peng, Y, Peng, J, Yu, S, Wu, C et al. 2024. A supergene controls facultative
diapause in the crop pest Helicoverpa armigera. *Cell Rep* 43
72. Johansson, F, Orizaola, G, Nilsson-Örtman, V 2020. Temperate insects with narrow
seasonal activity periods can be as vulnerable to climate change as tropical insect
species. *Sci Rep* 10:8822
73. Johansson, F, Stoks, R, Rowe, L, De Block, M 2001. Life history plasticity in a
damselfly: effects of combined time and biotic constraints. *Ecology* 82:1857-1869
74. Jordan, RG, Bradshaw, WE 1978. Geographic Variation in the Photoperiodic Response
of the Western Tree-Hole Mosquito, Aedes sierrensis. *Ann Entomol Soc Am* 71:487-490
75. Jørgensen, LB, Ørsted, M, Malte, H, Wang, T, Overgaard, J 2022. Extreme escalation
of heat failure rates in ectotherms with global warming. *Nature* 611:93-98
76. Joschinski, J, Bonte, D 2021. Diapause and bet-hedging strategies in insects: a meta-
analysis of reaction norm shapes. *Oikos* 130:1240-1250
77. Kauranen, H, Kinnunen, J, Hiillos, A-L, Lankinen, P, Hopkins, D et al. 2019. Selection
for reproduction under short photoperiods changes diapause-associated traits and
induces widespread genomic divergence. *J Exp Biol* 222:jeb205831
78. Kellermann, V, van Heerwaarden, B 2019. Terrestrial insects and climate change:
adaptive responses in key traits. *Physiol. Entomol.* 44:99-115

79. Kimura, MT 1988. Adaptations to Temperate Climates and Evolution of Overwintering
Strategies in the Drosophila melanogaster Species Group. *Evolution* 42:1288-1297
80. Kimura, MT 2004. Cold and heat tolerance of drosophilid flies with reference to their
latitudinal distributions. *Oecologia* 140:442-449
81. Kivelä, SM, Välimäki, P, Carrasco, D, Mäenpää, MI, Oksanen, J 2011. Latitudinal
insect body size clines revisited: a critical evaluation of the saw-tooth model. *J Anim*
Ecol 80:1184-1195
82. Kojima, W, Nakakura, T, Fukuda, A, Lin, C-P, Harada, M et al. 2020. Latitudinal cline
of larval growth rate and its proximate mechanisms in a rhinoceros beetle. *Funct Ecol*
34:1577-1588
83. Kostal, V 2006. Eco-physiological phases of insect diapause. *J Insect Physiol* 52:113-
127
84. Kozak, GM, Wadsworth, CB, Kahne, SC, Bogdanowicz, SM, Harrison, RG et al. 2019.
Genomic Basis of Circannual Rhythm in the European Corn Borer Moth. *Curr Biol*
29:3501-3509.e5
85. Kurota, H, Shimada, M 2003. Geographical variation in photoperiodic induction of
larval diapause in the bruchid beetle, Bruchidius dorsalis: polymorphism in
overwintering stages. *Entomol Exp Appl* 107:11-18
86. Lancaster, LT 2016. Widespread range expansions shape latitudinal variation in insect
thermal limits. *Nat Clim Chang* 6:618-621
87. Lawlor, JA, Comte, L, Grenouillet, G, Lenoir, J, Baecher, JA et al. 2024. Mechanisms,
detection and impacts of species redistributions under climate change. *Nature Reviews*
Earth & Environment 5:351-368
88. Lehmann, P, Ammunét, T, Barton, M, Battisti, A, Eigenbrode, SD et al. 2020. Complex
responses of global insect pests to climate warming. *Front Ecol Environ* 18:141-150
89. Lehmann, P, Piiroinen, S, Lyytinen, A, Lindström, L 2015. Responses in metabolic rate
to changes in temperature in diapausing Colorado potato beetle Leptinotarsa
decemlineata from three European populations. *Physiol. Entomol.* 40:123-130
90. Levins, R 1969. Thermal Acclimation and Heat Resistance in Drosophila Species. *Am*
Nat 103:483-499
91. Lindestad, O, Nylin, S, Wheat, CW, Gotthard, K 2024. Testing for variation in
photoperiodic plasticity in a butterfly: Inconsistent effects of circadian genes between
geographic scales. *Ecol Evol* 14:e11713
92. Lindestad, O, Wheat, CW, Nylin, S, Gotthard, K 2019. Local adaptation of
photoperiodic plasticity maintains life cycle variation within latitudes in a butterfly.
Ecology 100:e02550
93. Logan, ML, Cox, CL 2020. Genetic Constraints, Transcriptome Plasticity, and the
Evolutionary Response to Climate Change. *Front Gene* 11
94. Lushai, G, Hardie, J, Harrington, R 1996. Diapause termination and egg hatch in the
bird cherry aphid, Rhopalosiphum padi. *Entomol Exp Appl* 81:113-115
95. Macgregor, CJ, Thomas, CD, Roy, DB, Beaumont, MA, Bell, JR et al. 2019. Climate-
induced phenology shifts linked to range expansions in species with multiple
reproductive cycles per year. *Nat Comm* 10:4455
96. Malusare, SP, Zilio, G, Fronhofer, EA 2023. Evolution of thermal performance curves:
A meta-analysis of selection experiments. *J Evol Biol* 36:15-28
97. Martin, RA, Chick, LD, Garvin, ML, Diamond, SE 2021. In a nutshell, a reciprocal
transplant experiment reveals local adaptation and fitness trade-offs in response to
urban evolution in an acorn-dwelling ant. *Evolution* 75:876-887
98. Masaki, S 1967. Geographic Variation and Climatic Adaptation in a Field Cricket
(Orthoptera: Gryllidae). *Evolution* 21:725-741

99. McNamara, JM, Houston, AI 1996. State-dependent life histories. *Nature* 380:215-221

100. Merckx, T, Nielsen, ME, Heliölä, J, Kuussaari, M, Pettersson, LB et al. 2021.
Urbanization extends flight phenology and leads to local adaptation of seasonal
plasticity in Lepidoptera. *Proc Natl Acad Sci USA* 118:e2106006118

101. Merckx, T, Nielsen, ME, Kankaanpää, T, Kadlec, T, Yazdanian, M, Kivelä, SM 2024.
Continent-wide parallel urban evolution of increased heat tolerance in a common moth.
Evol Appl 17:e13636

102. Mori, A, Romero-Severson, J, Severson, DW 2007. Genetic basis for reproductive
diapause is correlated with life history traits within the Culex pipiens complex. *Insect*
Mol Biol 16:515-524

103. Murata, Y, Ideo, S, Watada, M, Mitsui, H, Kimura, MT 2009. Genetic and
physiological variation among sexual and parthenogenetic populations of Asobara
japonica (Hymenoptera: Braconidae), a larval parasitoid of drosophilid flies. *Eur. J.*
Entomol. 106:171-178

104. Mushegian, AA, Neupane, N, Batz, Z, Mogi, M, Tuno, N et al. 2021. Ecological
mechanism of climate-mediated selection in a rapidly evolving invasive species. *Ecol*
Lett 24:698-707

105. Musolin, DL, Tougou, D, Fujisaki, K 2011. Photoperiodic response in the subtropical
and warm-temperate zone populations of the southern green stink bug Nezara viridula:
why does it not fit the common latitudinal trend. *Physiol Entomol* 36:379-384

106. Nechols, JR, Tauber, MJ, Tauber, CA 1987. Geographical variability in
ecophysiological traits controlling dormancy in Chrysopa oculata (Neuroptera:
Chrysopidae). *J Insect Physiol* 33:627-633

107. Nielsen, ME, Kingsolver, JG 2020. Compensating for climate change–induced cue-
environment mismatches: evidence for contemporary evolution of a photoperiodic
reaction norm in Colias butterflies. *Ecol Lett* 23:1129-1136

108. Nielsen, ME, Nylin, S, Wiklund, C, Gotthard, K 2023. Evolution of butterfly seasonal
plasticity driven by climate change varies across life stages. *Ecol Lett* 26:1548-1558

109. Nilsson-Örtman, V, Stoks, R, De Block, M, Johansson, F 2012. Generalists and
specialists along a latitudinal transect: patterns of thermal adaptation in six species of
damselflies. *Ecology* 93:1340-1352

110. Noda, H 1992. Geographic-Variation of Nymphal Diapause in the Small Brown
Planthopper in Japan. *Jap Agri Res Q* 26:124-129

111. Nylin, S, Gotthard, K 1998. Plasticity in life history traits. *Annu Rev Entomol* 43:63-83

112. Nylin, S, Svärd, L 1991. Latitudinal patterns in the size of European butterflies. *Holarct*
Ecol 14:192-202

113. Parker, AL, Kingsolver, JG 2023. Population divergence in nutrient-temperature
interactions in Pieris rapae. *Front Insect Sci* 3

114. Parmesan, C 2006. Ecological and evolutionary responses to recent climate change.
Annu Rev Ecol Syst 37:637-669

115. Parmesan, C, Ryrholm, N, Stefanescu, C, Hill, JK, Thomas, CD et al. 1999. Poleward
shifts in geographical ranges of butterfly species associated with regional warming.
Nature 399:579-583

116. Parmesan, C, Morecroft, MD, Trisurat, Y, Adrian, R, Anshari, GZ et al. 2022.
Terrestrial and Freshwater Ecosystems and Their Services. In *Climate Change 2022:*
Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Sixth
Assessment Report of the Intergovernmental Panel on Climate Change, eds. Pörtner, H-
O, Roberts, DC, Tignor, M, Poloczanska, ES, Mintenbeck, K et al., pp. 197-377.
Cambridge, UK and New York, NY, USA: Cambridge University Press,

117. Parmesan, C, Yohe, G 2003. A globally coherent fingerprint of climate change impacts
across natural systems. *Nature* 421:37-42
118. Pegoraro, M, Zonato, V, Tyler, ER, Fedele, G, Kyriacou, CP, Tauber, E 2017.
Geographical analysis of diapause inducibility in European Drosophila melanogaster
populations. *J Insect Physiol* 98:238-244
119. Peng, X, Song, C-M, Wang, K, Chen, M-H 2017. Geographical Variations in the Life
Histories of Rhopalosiphum padi (Hemiptera: Aphididae) in China. *J Econ Entomol*
110:961-970
120. Pigot, AL, Merow, C, Wilson, A, Trisos, CH 2023. Abrupt expansion of climate change
risks for species globally. *Nat Ecol Evol* 7:1060-1071
121. Pruisscher, P, Nylin, S, Wheat, CW, Gotthard, K 2021. A region of the sex
chromosome associated with population differences in diapause induction contains
highly divergent alleles at clock genes. *Evolution* 75:490-500
122. Pruisscher, P, Nylin, S, Gotthard, K, Wheat, CW 2018. Genetic variation underlying
local adaptation of diapause induction along a cline in a butterfly. *Mol Ecol* 27:3613-
3626
123. Ragland, GJ, Kingsolver, JG 2008. The effect of fluctuating temperatures on ectotherm
life-history traits: comparisons among geographic populations of Wyeomyia smithii.
Evol Ecol Res 10:29-44
124. Ragland, GJ, Armbruster, PA, Meuti, ME 2019. Evolutionary and functional genetics of
insect diapause: a call for greater integration. *Curr Opin Insect Sci* 36:74-81
125. Ragland, GJ, Kingsolver, JG 2008. Evolution of thermotolerance in seasonal
environments: the effects of annual temperature variation and life-history timing in
wyeomyia smithii. *Evolution* 62:1345-1357
126. Reznik, SY, Dolgovskaya, MY, Ovchinnikov, AN, Belyakova, NA 2015. Weak
photoperiodic response facilitates the biological invasion of the harlequin ladybird
armonia axyridis (Pallas) (Coleoptera: Coccinellidae). *J. Appl. Entomol.* 139:241-249
127. Roff, DA 1980. Optimizing development time in a seasonal environment: the 'ups and
downs' of clinal variation. *Oecologia* 45:202-208
128. Roff, DA 1992. *The Evolution of Life Histories* 535 New York:Chapman & Hall
129. Rudman, SM, Greenblum, SI, Rajpurohit, S, Betancourt, NJ, Hanna, J et al. 2022.
Direct observation of adaptive tracking on ecological time scales in Drosophila. *Science*
375:eabj7484
130. Sadakiyo, S, Ishihara, M 2011. Rapid seasonal adaptation of an alien bruchid after
introduction: geographic variation in life cycle synchronization and critical photoperiod
for diapause induction. *Entomol Exp Appl* 140:69-76
131. Sadakiyo, S, Ishihara, M 2012. The role of host seed size in mediating a latitudinal body
size cline in an introduced bruchid beetle in Japan. *Oikos* 121:1231-1238
132. Sánchez-Bayo, F, Wyckhuys, KAG 2019. Worldwide decline of the entomofauna: A
review of its drivers. *Biol Conserv* 232:8-27
133. Scharf, I, Bauerfeind, SS, Blanckenhorn, WU, Sch 2010. Effects of maternal and
offspring environmental conditions on growth, development and diapause in latitudinal
yellow dung fly populations. *Clim Res* 43:115-125
134. Schmidt, PS, Paaby, AB 2008. Reproductive Diapause and Life-History Clines in North
American Populations of Drosophila melanogaster. *Evolution* 62:1204-1215
135. Seiter, S, Kingsolver, J 2013. Environmental determinants of population divergence in
life-history traits for an invasive species: climate, seasonality and natural enemies. *J*
Evol Biol 26:1634-1645
136. SgrÒ, CM, Overgaard, J, Kristensen, TN, Mitchell, KA, Cockerell, FE, Hoffmann, AA
2010. A comprehensive assessment of geographic variation in heat tolerance and

hardening capacity in populations of Drosophila melanogaster from eastern Australia. *J*
Evol Biol 23:2484-2493

137. Shelomi, M, Associate, ETD, Editor: MA 2012. Where Are We Now? Bergmann's
Rule Sensu Lato in Insects. *Am Nat* 180:511-519

138. Shoshan, AB, Pindeler, U, Wheat, CW, Gotthard, K 2025. Repeated evolution of
photoperiodic plasticity by different genetic architectures during recurrent colonizations
in a butterfly. *Proc Roy Soc B* 292:20242195

139. Siemers, I, von Schmalensee, L, Ziesemer, S, Gotthard, K, Lehmann, P 2024.
Predictable local adaptation in butterfly photoperiodism but not thermal performance
along a latitudinal cline. *Ecosphere* 15:e4842

140. Stearns, SC 1992. *The Evolution of Life Histories* 249 Oxford:Oxford University Press

141. Stroud, JT, Ratcliff, WC 2025. Long-term studies provide unique insights into
evolution. *Nature* 639:589-601

142. Sunday, J, Bennett, JM, Calosi, P, Clusella-Trullas, S, Gravel, S et al. 2019. Thermal
tolerance patterns across latitude and elevation. *Philos Trans R Soc Lond B*
374:20190036

143. Takeda, M, Chippendale, GM 1982. Environmental and genetic control of the larval
diapause of the Southwestern corn borer, Diatraea grandiosella. *Physiol Entomol* 7:99-
110

144. Tanaka, K, Murata, K 2017. Genetic Basis Underlying Rapid Evolution of an
Introduced Insect Ophraella communa (Coleoptera: Chrysomelidae): Heritability of
Photoperiodic Response. *Environ Entomol* 46:167-173

145. Tauber, E, Zordan, M, Sandrelli, F, Pegoraro, M, Osterwalder, N et al. 2007. Natural
Selection Favors a Newly Derived *timeless* Allele in *Drosophila melanogaster*. *Science*
316:1895

146. Tauber, MJ, Tauber, CA, Masaki, S 1986. *Seasonal Adaptations of Insects.*
Oxford:Oxford University Press

147. Terao, M, Tokuda, M, Shintani, Y 2021. Geographic Variation in Photoperiodic
Response for Induction of Pseudopupal Diapause in Epicauta gorhami (Coleoptera:
Meloidae). *Environ Entomol* 50:1145-1150

148. Thackeray, SJ, Henrys, PA, Hemming, D, Bell, JR, Botham, MS et al. 2016.
Phenological sensitivity to climate across taxa and trophic levels. *Nature* 535:241-245

149. Thomas, CD, Cameron, A, Green, RE, Bakkenes, M, Beaumont, LJ et al. 2004.
Extinction risk from climate change. *Nature* 427:145-148

150. Tougeron, K 2019. Diapause research in insects: historical review and recent work
perspectives. *Entomol Exp Appl* 167:27-36

151. Trotta, V, Calboli, FCF, Ziosi, M, Guerra, D, Pezzoli, MC et al. 2006. Thermal
plasticity in Drosophila melanogaster: A comparison of geographic populations. *BMC*
Evol Biol 6:67

152. Tsai, H-Y, Rubenstein, DR, Fan, Y-M, Yuan, T-N, Chen, B-F et al. 2020. Locally-
adapted reproductive photoperiodism determines population vulnerability to climate
change in burying beetles. *Nat Comm* 11:1398

153. Urban, MC, Swaegers, J, Stoks, R, Snook, RR, Otto, SP et al. 2024. When and how can
we predict adaptive responses to climate change. *Evol Lett* 8:172-187

154. Urbanski, J, Mogi, M, O'Donnell, D, DeCotiis, M, Toma, T, Armbruster, P 2012. Rapid
Adaptive Evolution of Photoperiodic Response during Invasion and Range Expansion
across a Climatic Gradient. *Am Nat* 179:490-500

155. van Asch, M, Salis, L, Holleman, LJM, van Lith, B, Visser, ME 2013. Evolutionary
response of the egg hatching date of a herbivorous insect under climate change. *Nat*
Clim Chan 3:244-248

156. van der Have, TM, deJong, G 1996. Adult size in ectotherms: Temperature effects on
growth and differentiation. *J Theo Biol* 183:329-340
157. van Klink, R, Bowler, DE, Gongalsky, KB, Swengel, AB, Gentile, A, Chase, JM 2020.
Meta-analysis reveals declines in terrestrial but increases in freshwater insect
abundances. *Science* 368:417-420
158. Vesala, L, Hoikkala, A 2011. Effects of photoperiodically induced reproductive
diapause and cold hardening on the cold tolerance of Drosophila montana. *J Insect*
Physiol 57:46-51
159. von Schmalensee, L, Caillault, P, Gunnarsdóttir, KH, Gotthard, K, Lehmann, P 2023.
Seasonal specialization drives divergent population dynamics in two closely related
butterflies. *Nat Comm* 14:3663
160. von Schmalensee, L, Süess, P, Roberts, KT, Gotthard, K, Lehmann, P 2024. A
quantitative model of temperature-dependent diapause progression. *P Natl Acad Sci*
USA 121:e2407057121
161. Wagner, DL, Grames, EM, Forister, ML, Berenbaum, MR, Stopak, D 2021. Insect
decline in the Anthropocene: Death by a thousand cuts. *P Natl Acad Sci USA*
118:e2023989118
162. Walsh, BS, Parratt, SR, Hoffmann, AA, Atkinson, D, Snook, RR et al. 2019. The
Impact of Climate Change on Fertility. *Trends in Ecology & Evolution* 34:249-259
163. Walters, R, Hassal, M 2006. The temperature-size rule in ecotherms: may a general
explanation exist after all? *Am Nat* 167:510-523
164. Wang, X-P, Yang, Q-S, Dalin, P, Zhou, X-M, Luo, Z-W, Lei, C-L 2012. Geographic
variation in photoperiodic diapause induction and diapause intensity in Sericinus
montelus (Lepidoptera: Papilionidae). *Insect Sci* 19:295-302
165. Wang, X-M, Yang, X, Zang, L-S, Wang, Z, Ruan, C-C, Liu, X-J 2017. Effect of
geographic variation on biology and cold tolerance of Harmonia axyridis in China.
Entomol Gener 36:239-250
166. Weaving, H, Terblanche, JS, Pottier, P, English, S 2022. Meta-analysis reveals weak
but pervasive plasticity in insect thermal limits. *Nat Comm* 13:5292
167. Wiberg, RAW, Tyukmaeva, V, Hoikkala, A, Ritchie, MG, Kankare, M 2021. Cold
adaptation drives population genomic divergence in the ecological specialist,
Drosophila montana. *Mol Ecol* 30:3783-3796
168. Willi, Y, Van Buskirk, J 2022. A review on trade-offs at the warm and cold ends of
geographical distributions. *Philosophical Transactions of the Royal Society B:*
Biological Sciences 377:20210022
169. Williams, CB 1957. Insect Migration. *Annu Rev Entomol* 2:163-180
170. Williamson, J, Lu, M, Camus, MF, Gregory, RD, Maclean, IMD et al. 2025. Clustered
warming tolerances and the nonlinear risks of biodiversity loss on a warming planet.
Philosophical Transactions of the Royal Society B: Biological Sciences 380:20230321
171. Wipking, W 1988. Repeated larval diapause and diapause-free development in
geographic strains of the burnet moth Zygaena trifolii Esp. (Insecta, Lepidoptera).
Oecologia 77:557-564
172. Xiao, L, He, H-M, Huang, L-L, Geng, T, Fu, S, Xue, F-S 2016. Variation of life-history
traits of the Asian corn borer, Ostrinia furnacalis in relation to temperature and
geographical latitude. *Ecol Evol* 6:5129-5143
173. Yoshio, M, Ishii, M 1998. Geographical variation of pupal diapause in the great
mormon butterfly, Papilio memnon L. (Lepidoptera : Papilionidae), in Western Japan.
Appl Entomol Zool 33:281-288

174. Yoshio, M, Ishii, M 2001. Relationship between cold hardiness and northward invasion
in the great mormon butterfly, Papilio memnon L. (Lepidoptera: Papilionidae) in Japan.
Appl Entomol Zool 36:329-335
175. Zani, PA, Swanson, SET, Corbin, D, Cohnstaedt, LW, Agotsch, MD et al. 2005.
Geographic variation in tolerance of transient thermal stress in the mosquito wyeomyia
smithii. *Ecology* 86:1206-1211
176. Zheng, X-L, Yang, Q-S, Hu, Y-W, Lei, C-L, Wang, X-P 2015. Latitudinal variation of
morphological characteristics in the swallowtail ericinus montelus Gray, 1798
(Lepidoptera: Papilionidae). *Acta Zool* 96:242-252
941
942

943 **FIGURE 1**

944
945

Figure 1: Insects have several mechanisms to respond to changing climates.
A & B) Populations may track their preferred environment by shifting their geographic
distribution range (option 1). Alternatively, they may remain in place and adapt to the new
climatic conditions (option 2). C) This adaptation can occur through the evolution of thermal
tolerance (option 2a) and/or changes in seasonal timing (option 2b). D) An organism's
thermal tolerance evolves through changes in the minimal (CT_{min}), optimal (T_{opt}), and/or
maximal (CT_{max}) temperature at which it can operate. As thermal tolerance is governed by
trade-offs between performance at different temperatures, climate warming can result in
evolutionary shifts in all these parameters (blue to red reaction norm). Note, however, that
climate adaptation of growth and development also can occur through vertical shifts of the
entire reaction norm (between blue and broken red reaction norm) so that growth is faster in
northern populations that have less time and permissible thermal conditions to complete
development, or so that growth slows down in future warming climates to counteract the
direct effect of temperature on biological rates (so called "countergradient" variation). E) An
organism can change its seasonal timing via the evolution of photoperiodism; by modification
of how it uses and interprets daylength. Adaptation to climate warming by shifts or
prolongations of the timing for growth and reproduction towards colder parts of the year
(option 2b) is predicted to be associated with changes in the critical daylength (CDL) at
which diapause is induced (blue to red reaction norm). Additionally, if species respond to
warming climates by migrating closer to the poles (option 1), they also need to evolve
changes in how they interpret daylength in order to time their life cycle. Importantly, the
evolution of seasonal timing can relax selection for changes in thermal tolerance (and vice
versa), and changes in seasonal timing may also require life history adjustments to
accommodate changes in season length, such as overall increases or decreases in
development rate and size at maturation, predicting correlated evolution of photoperiodism
and life history.

Figure 2: A) The geographical locations of all 308 populations from the 24 species that were included in the analysis of life history and photoperiodism. B) shows the global effects of latitude on photoperiodism, mean life history, and thermal plasticity in life history (means and corresponding 95% confidence limits). Only photoperiodism shows a strong and consistent clinal pattern across species. The standardized latitudinal effect refers to the global slope estimated in a Bayesian multilevel model divided by the standard deviation in the slope observed across species. Thus, this effect size estimate corresponds to a measure of repeatability across species. The horizontal bars to the right show the average proportion of population differentiation within each species that was explained by latitude (as measured by partial R^2). Note that although life history traits did not show consistent clinal variation across species, latitude still explains a considerable fraction of the observed population-level variance in mean body size and development rate indicating evolution of latitudinal clines in opposing directions across species. Details are provided in the Supplementary material.

989
990
Figure 3: Population differentiation in photoperiodism, life history traits, and their thermal
plasticity in relation to absolute latitude of sampled populations for each species (indicated by
separate slopes). A) Critical day length increases consistently with latitude. In contrast,
average life history (B, E) and life history plasticity (C, F) do not show consistent latitudinal
patterns across species. D) The average day of the year at which the critical daylength for
diapause is reached at each population's location ("diapause timing") decreases with latitude,
indicating that diapause is induced at earlier dates at higher latitudes. The life stage in which
development rates were estimated is specified in B and C. To facilitate comparison
development rate and body size were standardized relative to the average trait value within
each study and are expressed as percent change relative to this average. Thermal plasticity is
expressed as the percent change in trait value (relative to the global average) per degree
Celsius. Error bars indicate standard errors. Slopes are drawn separately per study in panel F;
corresponding to 9 species with data from 12 studies.
1004
1005

1006