Cardiac-specific GCN5L1 deficiency promotes MASLD in HFpEF

Paramesha Bugga[1,2], Bellina A.S. Mushala[1,2], Michael W. Stoner[1,2], Janet R. Manning[1,2], Nisha Bhattarai[1,2], Maryam Sharifi-Sanjani[1,2], Amber Vandevender[3], Raja G.R. Mooli[3], Sadeesh K. Ramakrishnan[3], Brett A. Kaufman[1,2], Sruti S. Shiva[1,3], Cassandra L. Happe[4], Steven J. Mullet[4], Stacy L. Gelhaus[4], Michael J. Jurczak[2], and Iain Scott[1,2]*

[1]Vascular Medicine Institute, [2]Divisions of Cardiology, [2]Endocrinology and Metabolism, [3]Classical Hematology, Department of Medicine, [4]Health Sciences Mass Spectrometry Core, University of Pittsburgh

*For correspondence:

Iain Scott, PhD

University of Pittsburgh

200 Lothrop Street

Pittsburgh, PA 15261

iain.scott@pitt.edu

Keywords: GCN5L1, HFpEF, BCAAs, BCKAs, Heart, Liver, MASLD

Running Title: Heart-to-Liver Crosstalk in HFpEF

Word Count: 1199

Abstract

The prevalence of cardiometabolic heart failure with preserved ejection fraction (HFpEF) continues to grow, representing over half of heart failure cases in the United States. As no specific medication for HFpEF exists, treatment guidelines focus on the management of comorbidities related to metabolic syndrome (e.g. obesity, diabetes, hypertension) that promote the disease[1]. These same comorbidities also drive pathology in non-cardiac tissues, and the links between cardiometabolic disease presentations in different organs are increasingly being recognized. Preclinical studies on the potential crosstalk between HFpEF and metabolic disease in the liver (e.g. metabolic dysfunction-associated liver disease; MASLD) have focused on how liver dysfunction may affect the heart, particularly through the release of secreted liver proteins. This may reflect the situation in the clinic, where incident MASLD is a risk factor for future HFpEF development. Here, in contrast to this developing paradigm of liver-initiated cardiac disease, we report for the first time a defect in cardiac metabolism related to the mitochondrial metabolic protein GCN5L1 that drives hepatic steatosis and MASLD in HFpEF.

Treatment guidelines for cardiometabolic heart failure with preserved ejection fraction (HFpEF) focus on the management of comorbidities related to metabolic syndrome (*e.g.* obesity, hypertension) that promote disease progression[1]. These comorbidities also drive dysfunction in non-cardiac tissues, and the links between cardiometabolic disease presentations in different organs are increasingly being recognized. Preclinical studies on the potential crosstalk between HFpEF and metabolic disease in the liver (*e.g.* metabolic dysfunction-associated steatotic liver disease; MASLD) have largely focused on how liver dysfunction may affect the heart[2]. This may reflect clinical situations, where incident MASLD is a risk factor for future HFpEF development[3], and the general rarity of cardiac metabolic defects driving pathophysiology in other organs[4]. Here, in contrast to this paradigm, we report for the first time a cardiac-specific gene deletion that promotes MASLD in mice with HFpEF.

GCN5L1, a ubiquitously expressed protein, modulates cellular metabolism via interactions with mitochondrial and endo-lysosomal protein regulatory complexes[5]. To examine its potential function as a metabolic regulator in HFpEF, young adult C57BL/6NJ male wildtype (WT) and inducible, cardiomyocyte-specific GCN5L1 knockout (cKO) mice were placed on either a chow or HFpEF diet (60% HFD + 0.5 g/L L-NAME)[2] for 15 weeks (**Figures 1A-B**). All studies were carried out with approval from the University of Pittsburgh IACUC. HFpEF cKO mice displayed significantly worse diastolic dysfunction relative to WT mice, along with elevated body weight and fasting blood glucose levels (**Figures 1C-F**). Necropsy analysis following these unexpected findings demonstrated that liver steatosis (including histology-quantified MASLD) was a contributor to the increased body weight observed (**Figures 1G-I, S1**). Biochemical and chromatography analysis confirmed that significantly elevated liver triglycerides and cholesterol levels were the basis of the observed steatosis in cKO HFpEF livers (**Figures 1J-L**).

Bulk RNA-seq of HFpEF mouse livers demonstrated that cholesterol metabolism pathways were the most highly dysregulated in cKO mice. This manifested as a downregulation of genes involved in

lysosomal cholesterol trafficking (*e.g. Npc1*), with the subsequent accumulation of unprocessed

cholesterol likely suppressing hepatic cholesterol synthesis pathway genes (*e.g. Hmgcr*) (**Figures**

1M-N). Combined with an increased abundance of unprocessed cholesterol esters (**Figure 1J**), these

findings suggest that cKO HFpEF mouse livers develop steatosis partly via the abnormal

accumulation of cholesterol in lysosomes.

We next attempted to identify the cardiac-derived factor driving liver disease in cKO mice. Unbiased

proteomic detection of pooled WT and cKO plasma samples did not detect an obvious candidate. Of

note, serum amyloid proteins (*e.g.* SAA1, SAA2), previously identified as potential liver-secreted

factors driving heart fibrosis in HFpEF[2], were unexpectedly decreased in cKO HFpEF mice (**Figure**

1O, Table S1). Untargeted metabolomic analysis showed that the branched-chain amino acids

(BCAAs) leucine and isoleucine, and their branched-chain keto acid (BCKA) metabolites, were

significantly upregulated in the plasma of cKO HFpEF mice (**Figure 1P, Table S2**). Concordantly,

genes involved in BCAA/BCKA metabolism (*e.g. Klf15, Bckdha*) were significantly downregulated in

the hearts, but not livers, of cKO HFpEF mice (**Figures 1Q-R**). Furthermore, there was decreased

expression of cardiac genes involved in the metabolism of leucine/isoleucine BCKA derivatives (*Ivd*

and *Acadsb*, respectively), but not valine (*Acad8*), which matched the plasma accumulation of their

respective BCAAs and BCKAs (KIC and KMV) (**Figures 1Q-S**). In the liver, WT mice showed a

significant decline in leucine in HFpEF, while cKO mice maintained hepatic levels of this BCAA under

these conditions (**Figure 1U**). Combined, these data suggest that reduced BCAA/BCKA metabolism

in cKO mouse hearts, and the subsequent release of cardiac BCAAs/BCKAs into the circulation, may

be the heart-derived factor driving MASLD in cKO HFpEF mice.

Finally, we examined how BCAAs/BCKAs may negatively impact liver metabolism and drive MASLD.

Impaired cholesterol trafficking in endothelial cells inhibits the activation of mTOR[6], a master regulator

of autophagy and lysosomal function. We hypothesized that excess circulating BCAAs/BCKAs may

impede hepatic cholesterol trafficking, leading to mTOR-mediated lysosomal dysfunction and liver

cholesterol accumulation. In agreement with this hypothesis, mTOR activation (measured by p-mTOR

and p-P70S6K immunoblotting) was reduced in cKO livers relative to WT mice (**Figures 1V, S2A**).

Concordantly, treatment of liver HepG2 cells with the leucine-derived BCKA, KIC, resulted in a similar

reduction in mTOR activity (**Figures 1W, S2B**). Combined, these data suggest that elevated cardiac-

derived BCAA/BCKAs in the circulation negatively impact liver cholesterol metabolism via the mTOR

pathway.

In summary, we provide the first example of a cardiac-specific gene deletion driving fatty liver disease

in HFpEF. Loss of GCN5L1 in the heart during HFpEF led to disrupted cardiac BCAA/BCKA

metabolism, which resulted in hepatic cholesterol accumulation. These findings suggest that the

heart-liver signaling axis in HFpEF is bidirectional, and should not be viewed solely as a liver-initiated

process in this disease.

 FIGURE 1 – Cardiac-specific GCN5L1 deficiency promotes MASLD in HFpEF.

 A. Experimental protocol. **B.** Confirmation of cardiac-specific GCN5L1 deletion in cKO hearts. **C.** Left

 ventricle ejection fraction. **D.** Left ventricle E/e' ratio. **E.** Body weight. **F.** Fasting blood glucose. **G.**

 Gross anatomy and histological staining (H&E and Masson's Trichrome; MT) of livers. **H.** Liver weight

 I. NAFLD activity score (NAS). **J-L.** Liver triglycerides (TAG) and cholesterol (Chol). **M.** ShinyGO

 transcriptome analysis. **N.** Liver cholesterol metabolism genes. **O-P.** Plasma proteomics (pooled

 samples) and untargeted plasma metabolomics (BioCyc pathway analysis). **Q-R.** BCAA and BCKA

 metabolism genes in heart and liver **S.** Plasma BCAAs **T.** Plasma KIC and KMV. **U.** Liver leucine

 levels. **V.** mTOR and p70S6K phosphorylation in HFpEF livers. **W.** mTOR and p70S6K

 phosphorylation after KIC treatment in liver HepG2 cells. **X.** Proposed model. All graphs represent

 mean $\pm$ SD. Mann-Whitney U-test or Student's T-test was used for comparisons between two groups,

 one-way ANOVA was used for comparisons between multiple groups.

BIBLIOGRAPHY

1. Kittleson MM, et al. 2023 ACC Expert Consensus Decision Pathway on Management of Heart Failure With Preserved Ejection Fraction: A Report of the American College of Cardiology Solution Set Oversight Committee. J Am Coll Cardiol. (2023) 81:1835-1878.

2. Strocchi S, et al. Systems Biology Approach Uncovers Candidates for Liver-Heart Interorgan Crosstalk in HFpEF. Circ Res. (2024) 135: 873-876.

3. Chang KC, et al. Metabolic dysfunction-associated steatotic liver disease is associated with increased risks of heart failure. Eur J Heart Fail. (2025) doi: 10.1002/ejhf.3567. Epub ahead of print.

4. Sen MG, et al. Heart-derived factors and organ cross-talk in settings of health and disease: new knowledge and clinical opportunities for multimorbidity. J Physiol. (2025) doi: 10.1113/JP287400. Epub ahead of print.

5. Wu K, et al. The emerging roles of GCN5L1 in mitochondrial and vacuolar organelle biology. Biochim Biophys Acta Gene Regul Mech. (2021) 1864:194598.

6. Xu J, et al. Cholesterol trafficking is required for mTOR activation in endothelial cells. Proc Natl Acad Sci U S A. (2010) 107: 4764-9.