Cardiac-specific deletion of GCN5L1 promotes fatty liver disease in HFpEF

Paramesha Bugga[1,2], Bellina A.S. Mushala[1,2], Michael W. Stoner[1,2], Janet R. Manning[1,2], Nisha Bhattarai[1,2], Maryam Sharifi-Sanjani[1,2], Amber Vandevender[3], Raja G.R. Mooli[3], Sadeesh K. Ramakrishnan[3], Brett A. Kaufman[1,2], Sruti S. Shiva[1,3], Cassandra L. Happe[4], Steven J. Mullet[4], Stacy L. Gelhaus[4], Michael J. Jurczak[2], and Iain Scott[1,2]*

[1]Vascular Medicine Institute, [2]Division of Cardiology, [2]Division of Endocrinology and Metabolism, [3]Division of Classical Hematology, Department of Medicine, [4]Health Sciences Mass Spectrometry Core, University of Pittsburgh, Pittsburgh, PA 15261

*For correspondence:

Iain Scott, PhD, FAHA, FCVS

University of Pittsburgh

Pittsburgh, PA 15261

iain.scott@pitt.edu

Word Count: 800

Keywords: GCN5L1, HFpEF, BCAAs, Heart, Liver, Steatosis, MAFLD

Running Title: Heart-to-Liver Crosstalk in HFpEF

The authors have declared that no conflict of interest exists.

ABSTRACT

The prevalence of cardiometabolic heart failure with preserved ejection fraction (HFpEF) continues to grow, representing over half of heart failure cases in the United States. As no specific medication for HFpEF exists, treatment guidelines focus on the management of comorbidities related to metabolic syndrome (e.g. obesity, diabetes, hypertension) that promote the disease. These same comorbidities also drive pathology in non-cardiac tissues, and the links between cardiometabolic disease presentations in different organs are increasingly being recognized. Preclinical studies on the potential crosstalk between HFpEF and metabolic disease in the liver (e.g. metabolic-associated fatty liver disease; MAFLD) have focused on how liver dysfunction may affect the heart, particularly through the release of secreted liver proteins. This may reflect the situation in the clinic, where incident MAFLD is a risk factor for future HFpEF development. Here, in contrast to this developing paradigm of liver-initiated cardiac disease, we report for the first time a defect in cardiac metabolism related to the mitochondrial metabolic protein GCN5L1 that drives hepatic steatosis and MAFLD in HFpEF.

The prevalence of cardiometabolic heart failure with preserved ejection fraction (HFpEF) continues to grow, representing over half of heart failure cases in the United States[1]. As no specific medication for HFpEF exists, treatment guidelines focus on the management of comorbidities related to metabolic syndrome (*e.g.* obesity, diabetes, hypertension) that promote the disease[1]. These same comorbidities also drive pathology in non-cardiac tissues, and the links between cardiometabolic disease presentations in different organs are increasingly being recognized[2].

Preclinical studies on the potential crosstalk between HFpEF and metabolic disease in the liver (*e.g.* metabolic-associated fatty liver disease; MAFLD) have focused on how liver dysfunction may affect the heart, particularly through the release of secreted liver proteins[2]. This may reflect the situation in the clinic, where incident MAFLD is a risk factor for future HFpEF development[3]. Here, in contrast to this developing paradigm of liver-initiated cardiac disease, we report for the first time a defect in cardiac metabolism that drives hepatic steatosis and MAFLD in HFpEF.

Young adult C57BL/6NJ male wildtype (WT) and cardiomyocyte-specific knockout mice of the mitochondrial metabolic regulatory protein GCN5L1 (GCN5L1 cKO)[4] were placed on either a chow or HFpEF diet (60% high fat diet + 0.5 g/L L-NAME)[2] for 15 weeks (**Figure [A]**). All studies were carried out with approval from the University of Pittsburgh IACUC and followed NIH guidelines. While both WT and GCN5L1 cKO HFpEF mice developed diastolic dysfunction in the absence of systolic impairment as expected, GCN5L1 cKO animals displayed significantly elevated body weight and fasting blood glucose levels relative to WT mice (**Figure [A]**). Necropsy analysis following this unexpected finding demonstrated that GCN5L1 cKO mice on a HFpEF diet had significantly increased liver weights and histology-quantified MAFLD (NAS; **Figure [B]**). Biochemical and chromatography analysis showed that significantly elevated liver triglycerides and cholesterol levels were the basis for the observed hepatic steatosis in GCN5L1 cKO HFpEF mice (**Figure [B]**).

Bulk RNA-seq on HFpEF mouse livers demonstrated that transcriptional pathways related to cholesterol were highly dysregulated in GCN5L1 cKO mice. This manifested as a downregulation of genes involved in lysosomal cholesterol processing (*e.g. Npc1, Lamp2*), along with the suppression of hepatic cholesterol synthesis pathways, in GCN5L1 cKO mice (**Figure [C]**). Combined with the increased abundance of unprocessed cholesterol esters in chromatography analysis (**Figure [B]**), these findings suggest that the livers of GCN5L1 cKO HFpEF mice develop steatosis partly from an abnormal accumulation of cholesterol in lysosomes.

We next attempted to determine the cardiac-derived factor that was driving liver disease in GCN5L1 cKO mice. Unbiased proteomic detection of pooled WT and GCN5L1 cKO plasma samples did not detect an obvious protein signal. Of note, serum amyloid proteins (*e.g.* SAA1, SAA2, SAMP), previously identified as potential liver-secreted factors driving heart fibrosis in HFpEF[2], were significantly decreased in GCN5L1 cKO HFpEF mice (**Figure [D]**). Untargeted metabolomic analysis of plasma samples showed that branched-chain amino acids (BCAAs) and their metabolites were significantly upregulated in GCN5L1 cKO HFpEF mice (**Figure [D]**). We examined BCAA metabolic gene expression in WT and GCN5L1 cKO HFpEF mice by bulk RNA-seq, and found that genes involved in BCAA and branched-chain ketoacid (BCKA) metabolism (*e.g. Klf15, Bckdha, Bckdhb*) were significantly downregulated in the hearts, but not livers, of GCN5L1 cKO HFpEF mice (**Figure [E]**). Furthermore, there was a downregulation of cardiac genes involved in the metabolism of leucine and isoleucine BCKA derivatives (*Ivd* and *Acadsb*, respectively), but not valine (*Acad8*), which matched the plasma accumulation of their BCAA parent metabolites (**Figure [E]**). Lastly, while the livers of WT mice showed a significant decline in leucine after HFpEF diet exposure, GCN5L1 cKO mice maintained their tissue levels of this BCAA under HFpEF conditions (**Figure [E]**), potentially via uptake from the circulation. Combined, these data suggest that reduced cardiac BCAA/BCKA metabolism in GCN5L1 cKO mice, and the subsequent release of cardiac BCAAs/BCKAs into the circulation, may be the factor driving the observed hepatic steatosis.

Finally, as lysosomal processes are involved in cholesterol handling, and BCAAs are known to inhibit hepatic autophagy[5], we hypothesized there may be a defect in these pathways in GCN5L1 cKO mouse livers. As predicted, immunoblotting for the key autophagosome regulatory protein LC3 demonstrated a reduction in both total and active (LC3-II) forms of this protein (**Figure [F]**), suggesting an overall suppression of autophagy/lysosomal processes in the livers of cardiac-specific GCN5L1 cKO HFpEF mice.

In summary, we provide the first example of a cardiac-specific gene deletion driving metabolic dysfunction and liver disease progression in HFpEF. Loss of GCN5L1 in the heart during HFpEF led to disrupted cardiac BCAA metabolism and subsequent elevation of plasma BCAA/BCKA metabolites, which promoted lipid accumulation and autophagy dysfunction in the liver. These findings suggest that the heart-liver signaling axis in HFpEF is bidirectional, and that future studies of interorgan crosstalk will require sophisticated mechanisms to determine the tissue source of the signal.

FIGURE – Cardiac-specific deletion of GCN5L1 promotes the development of liver steatosis.

A. Wildtype (WT) and cardiomyocyte-specific GCN5L1 knockout (cKO) mice displayed normal systolic function (ICVT) and impaired diastolic function (IVRT) after 15 weeks of HFpEF diet (60% HFD + 0.5 g/L L-NAME in drinking water). GCN5L1 cKO mice displayed significantly increased body weight and fasting blood glucose. N = 8-12, One-way ANOVA. **B.** GCN5L1 cKO mice displayed significantly increased liver steatosis, liver weights, and NAFLD activity score (NAS). N = 7-12, One-way ANOVA. Liver triglycerides (TAG) and cholesterol (Chol) were increased in GCN5L1 cKO mice by thin-layer chromatography (TLC) and biochemical quantification. N = 5, Mann-Whitney. **C.** KEGG pathway analysis using ShinyGO demonstrated that cholesterol and lipid pathways were dysregulated in GCN5L1 cKO mice. N = 4, Adjusted *P* values. **D.** Plasma proteomics (pooled samples, N = 5 per group) and untargeted plasma metabolomics (N = 6, KEGG pathway analysis using Metaboanalyst) showed that branched-chain amino acids (BCAAs) were among the top modified pathways in GCN5L1 cKO mice. **E.** BCAA metabolism genes were downregulated in GCN5L1 cKO hearts, with minimal changes observed in liver. N = 4, Adjusted *P* values. Targeted metabolomics demonstrated ~25% higher leucine and isoleucine in plasma of GCN5L1 cKO mice. N = 6, Fold change). WT HFpEF mice displayed a significant decrease in liver leucine levels that was absent in GCN5L1 cKO HFpEF mice. N = 6, One-way ANOVA. **F.** Autophagosome LC3 levels are significantly reduced in GCN5L1 cKO mouse livers. N = 6, Two-tailed T-test.

BIBLIOGRAPHY

1. Kittleson MM, Panjrath GS, Amancherla K, Davis LL, Deswal A, Dixon DL, Januzzi JL Jr, Yancy CW. 2023 ACC Expert Consensus Decision Pathway on Management of Heart Failure With Preserved Ejection Fraction: A Report of the American College of Cardiology Solution Set Oversight Committee. J Am Coll Cardiol. 2023 May 9;81(18):1835-1878. doi: 10.1016/j.jacc.2023.03.393. Epub 2023 Apr 19. PMID: 37137593.

2. Strocchi S, Liu L, Wang R, Häseli SP, Capone F, Bode D, Nambiar N, Eroglu T, Santiago Padilla L, Farrelly C, Vacca A, Mascagni M, Oeing CU, Kintscher U, Jung S, Diezel SA, Liévano Contreras SV, Zhou M, Seldin M, Schiattarella GG. Systems Biology Approach Uncovers Candidates for Liver-Heart Interorgan Crosstalk in HFpEF. Circ Res. 2024 Sep 27;135(8):873-876. doi: 10.1161/CIRCRESAHA.124.324829. Epub 2024 Aug 29. PMID: 39206552; PMCID: PMC11427132.

3. Chang KC, Su TH, Wu CK, Huang SC, Tseng TC, Hong CM, Hsu SJ, Liu CH, Yang HC, Liu CJ, Kao JH. Metabolic dysfunction-associated steatotic liver disease is associated with increased risks of heart failure. Eur J Heart Fail. 2025 Jan 8. doi: 10.1002/ejhf.3567. Epub ahead of print. PMID: 39777761.

4. Manning JR, Thapa D, Zhang M, Stoner MW, Traba J, McTiernan CF, Corey C, Shiva S, Sack MN, Scott I. Cardiac-specific deletion of GCN5L1 restricts recovery from ischemia-reperfusion injury. J Mol Cell Cardiol. 2019 Apr;129:69-78. doi: 10.1016/j.yjmcc.2019.02.009. Epub 2019 Feb 15. PMID: 30776374; PMCID: PMC6486843.

5. Zhang F, Zhao S, Yan W, Xia Y, Chen X, Wang W, Zhang J, Gao C, Peng C, Yan F, Zhao H, Lian K, Lee Y, Zhang L, Lau WB, Ma X, Tao L. Branched Chain Amino Acids Cause Liver Injury in Obese/Diabetic Mice by Promoting Adipocyte Lipolysis and Inhibiting Hepatic Autophagy.

ACKNOWLEDGEMENTS

This work was supported by: National Institutes of Health Fellowships (F31DK134089 and T32HL110849) to B.A.S.M; National Institutes of Health Shared Instrument Grants (S10OD023402, S10OD032141) to S.L.G.; National Institutes of Health Research Grants (R01HL147861, R0HL156874) and American Heart Association Established Investigator Award (23EIA1037834) to I.S. The University of Pittsburgh Center for Metabolism and Mitochondrial Medicine is supported by the Pittsburgh Foundation (MR2020 109502) grant to M.J.J. This project used the University of Pittsburgh Medical Center (UPMC) Hillman Cancer Center and Tissue and Research Pathology/Pitt Biospecimen Core shared resource, which is supported in part by National Institutes of Health Center Granty P30CA047904. Echocardiography was carried out by the University of Pittsburgh Rodent Ultrasonography Core, which received funding from the NIH Shared Instrumentation Grant Program (S10OD023684).

AUTHOR CONTRIBUTIONS

P.B. and I.S. conceived the study and designed experiments. P.B., B.A.S.M., M.W.S., J.R.M., N.B., M.S-S., A.M.V., R.G.R.M, C.L.H, and S.J.M. performed experiments. P.B., C.L.H., S.J.M., R.G.R.M., and I.S. analyzed data. P.B. and I.S. prepared figures. S.S.S., B.A.K., S.K.R., S.L.G, and M.J.J. provided critical input and expertise. I.S. drafted the manuscript. P.B. and I.S. edited and revised the manuscript.