Binding Affinity Ranking at the Molecular Initiating Event (BARMIE): An open-source computational pipeline for ecological hazard ranking of endocrine disrupting chemicals.

Fernando Calahorro [1*], Parsa Fouladi[2*], Alessandro Pandini[2], Matloob Khushi[2], Yogendra Gaihre[1], Nic Bury[1#]

1. University of Southampton, School of Ocean and Earth Science, National Oceanography Centre, European Way, Southampton, SO14 2ZH, United Kingdom.
2. Brunel University of London, College of Engineering, Design and Physical Sciences, Department of Computer Science, Wilfred Brown Building, Uxbridge, UB8 3PH, United Kingdom.

* Contributed equally

Corresponding author

Abstract

One of the key challenges in ecological risk assessment lies in identifying the chemicals that pose the greatest threat and determining the species that are most vulnerable to their effects. Computational prediction of protein binding affinity can help in assessing the risk of chemicals to species. In this study we developed and validated an open-source tool called BARMIE (Binding Affinity Ranking at the Molecular Initiating Event) to rank chemical hazards and identify species that are most susceptible based on the binding affinity of the chemical to steroid receptor proteins. As an exemplar of BARMIE's output we focus on 163 teleost fish glucocorticoid receptors (GRs) and the natural ligand cortisol and 10 synthetic glucocorticoid (GCs) drugs and five other potential chemical GR agonists. The hazard ranking is based on the likelihood that the chemicals with the highest binding affinity are likely to outcompete cortisol at the receptor binding site. In this analysis, halcinonide, a GC, was predicted to be the most hazardous based on its binding affinity and the superorder Protacanthopterygii species, including the Esociformes and Salmoniformes, were identified as the most vulnerable. This computational pipeline can be expanded to evaluate more chemicals, species, and proteins as part of an in silico chemical hazard assessment tool.

Introduction

Approximately 350,000 synthetic chemicals are produced globally, yet most have not

undergone any human or environmental risk assessment[1]. Consequently, the impact of

these novel substances on human and wildlife health is likely underestimated. To protect

wildlife, hazard assessors, who evaluate the potential of a chemical to cause harm, and risk

assessors, who determine the likelihood of that harm occurring, face the challenge of

identifying which of these chemicals are of concern and the species that are most vulnerable

to them.

All vertebrates have a similar steroid hormone/receptor based endocrine system that

regulates a plethora of physiological and developmental processes. These processes are

controlled via the binding of the steroid ligand to their steroid receptor to initiate ligand

inducible transactivation or transrepression, or other cellular signalling pathways[2]. Due to

their significance the steroid receptors (glucocorticoid, mineralocorticoid, androgen,

oestrogen, and progesterone) are highly conserved within the vertebrate subphylum[3]. The

importance of this system for health is also reflected in concerns over endocrine active

substances that may perturb hormonal actions via these receptors leading to reproductive

and metabolic disorders as well as neural development.

In response to this concern national and international governments implemented testing

programs over 10 years ago to elucidate the endocrine disrupting potential of synthetic and

natural compounds[4,5]. These initially focused on the EATS (estrogen, androgen, thyroid and

steroidogenesis) modalities, but in more recent years several New Approach Methodologies

(NAMs) utilizing in silico techniques and novel in vitro methodologies have expanded the

screening of chemical hazards for non-EATS modalities[6] (e.g. glucocorticoid and

progesterone receptor and non-steroidal receptors). NAMs provide valuable insights into a

chemical's mode of action (MOA) and its potential effects[7-11], and it is expected that NAMs

will be integrated into regulatory frameworks in the future[12]. The majority of NAMs research

efforts have concentrated on human risk assessment, which aims to protect the individual. In

contrast, environmental and ecological risk assessments are far more ambitious, seeking to

protect numerous species and maintain ecosystem function.

Models such as EcoDrug[3] and Sequence Alignment to Predict Across Species Susceptibility

(SeqAPASS)[13] make use of protein conservation across phyla to identify non-target species

susceptible to drugs based on the presence of human or veterinary drug targets. Two recent

studies have combined the sequence information with empirical toxicity information to

provide greater information on the potential environmental impacts of chemicals. SeqAPASS

in combination with Genes to Pathway – Species Conservation Analysis (G2P-SCAN) uses

network analysis, based on Adverse Outcome Pathway (AOP) information, to identify

conserved Reactome[14] pathways and points of departure in toxic outcomes[15], and

RASRTox (Rapidly Acquire, Score, and Rank Toxicology data) links the sequence

information with toxicological databases (ECOTOX, ToxCast21) and QSAR models to

develop tool for ranking chemicals for hazard assessment[16]. However, one issue that is

difficult to avoid if the aim is to protect species and ecosystem function, is the lack of

species-specific chemical impact data and the models are restricted to data-rich resources

for only a few species (e.g. humans, mice, zebrafish). It is known that there are huge

differences in the concentrations of a chemical that induce a toxic response between species

within a taxon[17] likely due to species specific toxicokinetic and/or toxicodynamic

parametres[18]. Within the AOP framework[15] a potential reason for enhanced or decreased

chemical sensitivity are differences in binding affinity at the molecular initiating protein. If the

mutations in the protein that confer different binding affinity characteristics[19] are conserved

within orders or families of species or are species specific, then it is possible to develop a

more granular chemical hazard assessment based on interaction at the MIE for groups of

species, species or individuals.

This study focuses on glucocorticoid receptors (GRs) in teleost fish as an exemplar of this

approach, however examples for other EATS and non-EATS modalities are provided in

Supplementary Information (SI 4). The reason for focusing on the GR is two-fold. Firstly,

there are species differences in the binding affinity (K_d), transactivation (EC50) and trans

repression activity for both natural and synthetic glucocorticoids (GCs)[20-24], and previous

research has indicated a correlation between GR hormone binding affinity and hormone

EC50 for ligand-inducible gene transactivation in rainbow trout GR mutants[25]. Secondly,

synthetic glucocorticoids, which are commonly used to treat various health conditions, pose

growing environmental concerns with 17% of the GCs assessed are predicted to exist at

concentrations that pose a risk to fish[16]. The computational pipeline [Binding Affinity Ranking

at the Molecular Initiating Event (BARMIE)] ranks chemical binding affinities to GR across a

wide range of fish species to determine the binding affinity of synthetic glucocorticoids and

other chemicals to identify chemicals and species of concern. This is based on the

hypothesis that species with GRs exhibiting the highest binding affinity are more likely to

respond to lower concentrations of natural GCs or GR agonists or antagonists in the

bloodstream.

Materials and Methods

The novel computational pipeline Binding Affinity Ranking at the Molecular Initiating Event

(BARMIE) uses open-source database APIs and software (UniProt[27], Chembl[28],

OpenBabel[29], PyMol[30] and AutoDock Vina[31]). The code to estimate receptor binding

affinities, summary of the procedural steps and user instructions are available at

https://github.com/ParsaFouladi/Barmie. The pipeline was run on the University of

Southampton HPC Iridis 6, and the example provide is specific to teleost fish GRs. The

pipeline can be adapted for use with other HPC architecture as well as other species and

proteins.

Several steps were necessary in developing BARMIE. Firstly, genome annotation is based

on sequence homology to proteins of known function, and the level of validation may vary

depending on the database. There are over 250 genomes available from Ensembl and

UniProt provides verified and non-verified protein structures, and it is recommended to use

only those verified. Secondly, several receptor isoforms (e.g., splice variants) are predicted

in Uniport. Because very few isoforms have been characterized in fish[32] we did not remove

these. Thirdly, we only used the Uniport proteins with confirmed structures. However,

bespoke structures predicted from amino acid sequence information[33] can be integrated in

the pipeline. Fourthly, the absolute orientation of the receptor structure when imported into

AutoDock Vina generally differs between structures, making it difficult to automate the

definition of the box coordinates to correctly encompass the ligand binding domain. To

overcome this, structures are automatically aligned using PyMol scripting, so that the LBD

location is aligned for all the proteins. The only manual step in the pipeline is the definition of

the box coordinates for docking exploration of the ligand binding pocket. This is set to a

reference protein, in this example *Oncorhynchus mykis*s GR AF-P49843, and is identical for

all receptor due to the three- dimensional alignment in PyMol. For the GR used in this study,

the box size was set to 20, 20, 20 Å, and the coordinates X=5, Y=2, Z=-15.

Docking binding affinity were estimated for 163 teleost GR proteins (SI 1 contains the

Uniprot ID codes) in complex with the natural ligand cortisol (CHEMBL389621), the synthetic

glucocorticoids beclomethasone (CHEMBL1586), clobetasol (CHEMBL1201362),

dexamethasone (CHEMBL 384467), flumetasone (CHEMBL1201392), halcinonide

(CHEMBL1200845), mapracorat (CHEMBL2103876), mometasone (CHEMBL1201404),

prednicarbate (CHEMBL1200386), prednisolone (CHEMBL131), and triamcinolone

(CHEMBL1451) as well as the herbicide atrazine [CHEMBL15063. Unlikely to interact with

the GR][34], insecticide glyphosate [CHEMBL95764. Reported interactions with GR] [35],

anesthetic sevoflurane [CHEMBL1200694. Reported to affect the immune system but

interaction with the GC pathway unknown] [36], cortisol synthesis inhibitor osilodrosat

[CHEMBL3099695][37] and antibiotic triclocarban [CHEMBL1076347. A reported GR

antagonist][38]. AutoDock Vina has a stochastic algorithm to explore ligand binding poses, and

thus, docking searches were run 5 times and an average binding affinity are reported.

Docking simulations were run with exhaustiveness of 8, 32, and 128 to assess consistency

of results. Amino acid interaction fingerprint was conducted with the protein/ligand pose in

LigPlot$^+$ v2.2 (https://www.ebi.ac.uk/thornton-srv/software/LigPlus/).

Results and Discussion

The described pipeline referred to as BARMIE (Binding Affinity Ranking at the molecular

initiating event) predicts the binding affinity of 163 fish GRs to the natural ligand cortisol

(Figure 1a, for an example of cortisol docking with *Pygocentrus nattereri* AOA3B4D0J0 GR

and the amino acid interaction fingerprint), various synthetic GCs, and potential GR agonists.

We observed no difference between results with AutoDock Vina at exhaustiveness 8, 32,

and 128 (SI 3) and we have reported an average at exhaustiveness of 32. This produces a

ranking of binding affinity highlighting those species whose GRs bind with the highest affinity

to these chemicals and provides an *in-silico* hazard screening protocol based on binding to

the receptor, or in AOP parlance the MIE[39]. As an example of the potential outputs in the top

five out of 2822 combinations, we find the GC halcinonide is predicted to bind to 4 GRs with

the greatest affinity, and of these 4 GRs belong to the superorder Protacanthopterygii

containing the Esociformes and Salmoniformes (Figure 1b) and suggesting in this exercise

that the synthetic GC halcinonide may be considered the most significant chemical hazard

and the Protacanthopterygii as a group that may be vulnerable. However, the data generated

can be used in more nuanced ways by focusing on specific chemicals (Figure 2) and species

(Table 1) and for all GCs, we see that 55% of the five most sensitive species belong to the

Protacanthopterygii, with the species Northern pike (*Esox lucius*) ranked 1st for 7 of the 11

GCs tested (Table 1). There is limited empirical data on fish GR cortisol binding and

transactivation activity, but where this is available, there is a correlation between measured

GR hormone binding affinity and hormone EC50 for ligand-inducible gene transactivation for

rainbow trout GR[25]. The variance in binding in rainbow trout is due to mutations within the

receptors[25], similarly human receptor mutants that cause disease show differences in

hormone binding and transactivation characteristics [e.g., 40,41].

Our analysis shows that the Protacanthopterygii, which include the commercially important

salmon and trout, are an order of fish particularly susceptible to synthetic glucocorticoids.

However, a limitation should be noted that only 86 fish genomes have been annotated out of

the potentially 30,000 teleost species[42], and of these genomes, a high proportion, 8%, are

Protacanthopterygii. Thus, this order is overrepresented. The number of species genomes

sequenced is rapidly expanding for example the Earth BioGenome Project Network[43] has 63

global partners with an ambitious target of "[…] characteriz(ing) the genomes of all of Earth's

eukaryotic biodiversity over a period of ten years". This will provide the necessary genetic

information for BARMIE to provide a more granular order, family or species chemical risk

assessment.

The range of binding affinities is reduced for the non-GC chemicals atrazine, glyphosate,

sevoflurane, and osilodrosat (~-5 to ~-6 Kcal/mol,) compared to natural and synthetic GCs

(Figure 2 and SI 1). This would suggest that these chemicals would be classed low in an

MIE binding affinity-based hazard assessment because they are unlikely to out-compete

cortisol at the binding site. However, for risk assessment, the potential to cause harm is

based on the exposure concentration as well as the hazard. Thus, if fish accumulate these

chemicals to such an extent that this level far exceeds those of the natural ligand, it poses a

risk.

Environmental Implications

The challenge for ecological risk assessors is identifying those chemicals of concern

amongst the thousands in production and those species most vulnerable. The current

pipeline that has been developed enables us to rank chemicals and species based on the

receptor's binding affinity. This can be used for hazard assessment because the chemical

with a higher binding affinity is likely to out-compete the natural ligand and thus be a more

potent agonist or antagonist. The results are an exemplar of what the pipeline can offer, it

can be applied to any protein where the binding affinity of a ligand is crucial for its function.

This is particularly relevant for all steroid and non-steroid receptors enabling the

identification of potent endocrine-disrupting chemicals. BARMIE is a screening tool that

allows a toxicity testing program for regulatory purposes focusing on those chemicals and

species of concern, reducing costs and the number of animals used for testing.

Acknowledgements: This study was funded by the Natural Environment Research Council

in the United Kingdom, grant nos: NE/X000192/1 awarded to NB and MK.

References

1. Wang, Z., Walker, G.W., Muir, D.C.G., Nagatani-Yoshida, K et al., 2020. Toward a global understanding of chemical pollution: A first comprehensive analysis of national and regional chemical inventories. Environ. Sci. Technol. 54, 2575–2584. DOI: 10.1021/acs.est.9b06379.

2. Oakley, R.H., Cidlowski, J.A. 2013. The biology of the glucocorticoid receptor: New signaling mechanisms in health and disease. J. Allergy Clin. Immunol. 132, 1033–1044. doi: 10.1016/j.jaci.2013.09.007.

3. Verbruggen, B., Gunnarsson, L., Kristiansson, E., Österlund, T., Owen, S.F., Snape, J.R., Tyler, C.R. 2018. ECOdrug: a database connecting drugs and conservation of their targets across species. Nucleic Acids Res. 46, D930-D936. DOI:10.1093/nar/gkx1024.

4. Rotroff, D.M., Dix, D.J., Houck, K.A., Knudsen, T.B., Martin, M.T., McLaurin, K.W., Reif, D.M., Crofton, K.M., Singh, A.V., Xia, M., Huang, R., Judson, R.S. 2013. Using in vitro high throughput screening assays to identify potential endocrine-disrupting chemicals Environ. Health Perspect. 121, 7–14. http://dx.doi.org/10.1289/ehp.1205065

5. Hecker, M., Hollert, H. 2011. Endocrine disruptor screening: regulatory perspectives and needs. Environ. Sci. Eur. 23, 15 https://doi.org/10.1186/2190-4715-23-15.

6. Martyniuk, C.J., Martínez, R., Navarro-Martin, L., Kamstra, J.H., Schwednt, A., Reynaud, S., Chalifour, L. 2022. Emerging concepts and opportunities for endocrine disruptor screening of the non-EATS modalities. Environ. Res. 204, 111904. https://doi.org/10.1016/j.envres.2021.111904

7. Mondou, M., Hickey, G.M., Rahman, H.M.T., Maguire, s., Pain, G., Crump, D., Hecker, M., Basu, N. 2020. Factors affecting the perception of New Approach Methodologies (NAMs) in the ecotoxicology community. Integr. Environ. Asses. 16, 269–28. DOI:10.1002/ieam.4244.

8. Di Nicola, M.R., Cattaneo, I., Nathanail, A.V., Carnesecchi, E., Astuto, M.C., Steinbach, M., Williams, A.J., Charles, S., Gestin, O., Lopes, C., Lamonica, D., Tarazona, J.V., Dorne, J.M.C.M. 2023. The use of new approach methodologies for the environmental risk assessment of food and feed chemicals. Curr. Opin. Environ. Sci. Health. 31, 1-8. DOI: 10.1016/j.coesh.2022.100416.

9. van der Zalm, A.J., Barroso, J., Browne, P., Casey, W., Gordon, J., Henry, T.R., Kleinstreuer, N.C., Lowit, A.B., Perron, M., Clippinger, A.J. 2022. A framework for establishing scientific confidence in new approach methodologies. Arch. Toxicol. 96, 2865-2879. DOI:10.1007/s00204-022-03365-4.

10. Parish, S.Y., Aschner, S.T., Casey, W., Corvaro, M., Embry, M.R., Fitzpatrick, S., Kidd, D., Kleinstreuer, N.C., Lima, B.S., Settivari, R.S., Wolf, D.C., Yamazaki, D., Boobis, A. 2020. An evaluation framework for new approach methodologies (NAMs) for human health safety assessment. Regul. Toxicol. Pharmacol. 112, 104592. DOI: 10.1016/j.yrtph.2020.104592.

11. Lee, K-C., Guiney, P.D., Menzie, C.A., Belanger, S.E. 2023. Advancing the weight of evidence approach to enable chemical environmental risk assessment for decision-making and achieving protection goals. Integr. Environ. Asses. 19, 1188-1191. DOI: 10.1002/ieam.4803.

12. Schmeisser, S., Miccoli, A., von Bergen, M., Berggren, E., Braeuning, A., Busch, W., Desaintes, C., Gourmelon, A., Grafström, R., Harrill, J., Hartung, T., Herzler, M., Kass, J.E.N., Kleinstreuer, N., Leist, M., Luijten, M., Marx-Stoelting, P., Poetz O., van Ravenzwaay, B., Roggeband, R., Rogiers, V., Roth, A., Sanders, P., Thomas, R.S., Vinggaard, A. M., Vinken, M., van de Water, B., Luch, A., Tralau, T. 2023. New approach methodologies in human regulatory toxicology - Not if, but how and when! Environ Int. 178:108082. doi: 10.1016/j.envint.2023.108082.

13. Lalone, C.A., Villeneuve, D.L., Lyons, D., Helgen, H.W., Robinson, S.L., Swintek, J.A., Saari, T.W., Ankley, G.T. 2016. Sequence Alignment to Predict Across Species Susceptibility (SeqAPASS): A web-based tool for addressing the challenges of cross-species extrapolation of chemical toxicity. Toxicol. Sci. 153, 228-45. doi: 10.1093/toxsci/kfw119.

14. Milacic, M., Beavers, D., Conley, P., Gong, C., Gillespie, M., Griss, J., Haw, R., Jassal, B., Matthews, L., May, B., Petryszak, R., Ragueneau, W., Rothfels, K., Sevilla, G., Shamovsky, V., Stephan, R., Tiwari, K., Varusai, T., Weiser, J., Wright, A., Wu, G., Stein, L., Hermjakob, H., D'Eustachio, P. 2024 The Reactome Pathway Knowledgebase 2024. Nucleic Acids Res. 52, D672–D678. https://doi.org/10.1093/nar/gkad1025.

15. Schumann, P., Rivetti, C., Houghton, J., Campos, B., Hodges, G., LaLone, C. 2024. Combination of computational new approach methodologies for enhancing evidence of biological pathway conservation across species. Sci Total Environ 912:168573. doi: 10.1016/j.scitotenv.2023.168573

16. Schaupp, C.M., Byrne, G., Chan, M., Haggard, D.E., Hazemi, M., Jankowski, M.D., LaLone, C.A., LaTier, A., Mattingly, K.Z., Olker, J.H., Renner, J., Sharma, B., Villeneuve, D.L.. 2024. An automated computational data pipeline to rapidly acquire, score, and rank toxicological data for ecological hazard assessment. Integr Environ Assess Manag 20:2203-2217. doi: 10.1002/ieam.4945.

17. Posthuma, L., van Gils, J., Zijp, M.C., van de Meent, D., de Zwart, D. 2019. Species sensitivity distributions for use in environmental protection, assessment, and management of aquatic ecosystems for 12 386 chemicals. Environ. Toxicol. Chem. 38, 905-917. DOI:10.1002/etc.4373.

18. Nyman, A.M., Schirmer, K., Ashauer, R. 2014. Importance of toxicokinetics for interspecies variation in sensitivity to chemicals. Environ. Sci. Tecnolo. 48, 5946-5954. https://doi.org/10.1021/es5005126

19. Gerasimavicius, L., Livesey, B.J. Marsh, J.A. 2022. Loss-of-function, gain-of-function and dominant-negative mutations have profoundly different effects on protein structure. Nat Commun 13, 3895. https://doi.org/10.1038/s41467-022-31686-6

20. Bury, N.R., Sturm, A., Le Rouzic, P., Lethimonier, C., Ducouret, B., Guiguen, Y., Robinson-Rechavi, M., Laudet, V., Rafestin-Oblin, M.E., Prunet, P. 2013. Evidence for two distinct functional glucocorticoid receptors in teleost fish. J. Mol. Endocrinol. 31. 141-56. doi: 10.1677/jme.0.0310141.

21. Bury, N.R., Sturm, A. 2007. Evolution of the corticosteroid receptor signalling pathway in fish. Gen. Comp. Endocrinol. 153, 47-56. doi: 10.1016/j.ygcen.2007.03.009.

22. Stolte, E.H., de Mazon, A.F., Leon-Koosterziel, K.M., Jesiak, M., Bury, N.R., Sturm, A., Savelkoul, H.F., van Kemenade, B.M., Flik, G. 2008. Corticosteroid receptors involved in stress regulation in common carp, *Cyprinus carpio*. J. Endocrinol. 198, 403-17. doi: 10.1677/JOE-08-0100.

23. Bury, N.R. 2017. The evolution, structure and function of the ray finned fish (Actinopterygii) glucocorticoid receptors. Gen. Comp. Endocrinol. 251, 4-11. doi: 10.1016/j.ygcen.2016.06.030.

24. Kugathas, S., Sumpter, J.P. 2011. Synthetic glucocorticoids in the environment: first results on their potential impacts on fish. Environ Sci Technol. 45, 2377-83. doi: 10.1021/es104105e.

25. Sturm, A., Colliar, L., Leaver, M.J., Bury, N.R. 2011. Molecular determinants of hormone sensitivity in rainbow trout glucocorticoid receptors 1 and 2. Mol. Cell. Endocrinol. 333, 181-9. doi: 10.1016/j.mce.2010.12.033.

26. Hamilton, C.M., Winter, M.J., Margiotta-Casaluci, L., Owen, S.F., Tyler, C.R. 2022. Are synthetic glucocorticoids in the aquatic environment a risk to fish? Environ. Int. 162, 107163. doi: 10.1016/j.envint.2022.107163.

27. UniProt Consortium T. 2018. UniProt: the universal protein knowledgebase. Nucleic Acids Res. 46, 2699. doi: 10.1093/nar/gky092.

28. Zdrazil, B., Felix, E., Hunter, F., Manners, E.J., Blackshaw, J., Corbett, S., de Veij, M., Ioannidis, H., Lopez, D.M., Mosquera, J.F., Magarinos, M.P., Bosc, N., Arcila, R., Kizilören, T., Gaulton, A., Bento, A.P., Adasme, M.F., Monecke, P., Landrum, G.A., Leach, A.R. 2024. The ChEMBL Database in 2023: a drug discovery platform spanning multiple bioactivity data types and time periods. Nucleic Acids Res. 52, D1180-D1192. doi: 10.1093/nar/gkad1004.

29. OpenBabbel. https://openbabel.org/index.html

30. Schrödinger, L., DeLano, W. PyMOL [Internet]. 2020. http://www.pymol.org/pymol

31. Eberhardt, J., Santos-Martins, D., Tillack, A.F., Forli, S., 2021. AutoDock Vina 1.2.0: New docking methods, expanded force field, and python bindings. J. Chem. Inf. Model. 61, 3891-3898. doi: 10.1021/acs.jcim.1c00203.

32. Romero, A., Vega. M., Santibáñez, N., Spies, J., Pérez, T., Enríquez R., Kausel G., Oliver C., Oyarzún R., Tort L., Vargas-Chacoff L. 2020. *Salmo salar* glucocorticoid receptors analyses of alternative splicing variants under stress conditions. Gen. Comp. Endocrinol. 293, 113466. doi: 10.1016/j.ygcen.2020.113466.

33. Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O., Tunyasuvunakool, K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S.A.A., Ballard, A.J., Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Petersen, S., Reiman, D., Clancy, E., Zielinski, M., Steinegger, M., Pacholska, M., Berghammer, T., Bodenstein, S., Silver, D., Vinyals, O., Senior, A.W., Kavukcuoglu, K., Kohli, P., Hassabis, D. 2021. Highly accurate protein structure prediction with AlphaFold. Nature 596, 583-589. doi: 10.1038/s41586-021-03819-2.

34. Devos, S., De Bosscher, K., Staels, B., Bauer, E., Roels, F., Vanden Berghe, W., Haegeman, G., Hooghe, R., Hooghe-Peters, E.L. 2016. Inhibition of cytokine production by the herbicide atrazine - Search for nuclear receptor targets. Bioche, Pharmacol. 65, 303-308. DOI10.1016/S0006-2952(02)01507-1

35. Spulber, S., Reis, L., Alexe, P., Ceccatelli, S. 2024. Decreased activity in zebrafish larvae exposed to glyphosate-based herbicides during development-potential mediation by glucocorticoid receptor. Front. Toxicol. 6 DOI10.3389/ftox.2024.1397477

36. Hajjo, R., Sabbah, D.A., Bardaweel, S.K. 2020. Chemocentric informatics analysis: Dexamethasone versus combination therapy for COVID-19. ACS OMEGA 5, 29765-29779. DOI10.1021/acsomega.0c03597

37. Fleseriu, M., Pivonello, R., Young, J., Hamrahian, A.H., Molitch, M.E., Shimizu, C., Tanaka, T., Shimatsu, A., White, T., Hilliard,A., Tian, C., Sauter, N., Biller, B.M.K., Bertagna, X. 2015. Osilodrostat, a potent oral 11β-hydroxylase inhibitor: 22-week, prospective, Phase II study in Cushing's disease. Pituitary, 19, 138-148. doi: 10.1007/s11102-015-0692-z.

38. Kenda, M., Kuzelicki, N.K., Iida, M., Kojima, H., Dolenc, M.S. 2020. Triclocarban, Triclosan, Bromochlorophene, Chlorophene, and Climbazole Effects on Nuclear Receptors: An in Silico and in Vitro Study. Environ. Health Perspec. 128 (10). DOI10.1289/EHP6596.

39. Villeneuve, D.L., Crump, D., Garcia-Reyero, N., Hecker, M., Hutchinson, T.H., LaLone, C.A., Landesmann, B., Lettieri, T., Munn, S., Nepelska, M., Ottinger, M.A., Vergauwen, L., Whelan M., 2014. Adverse Outcome Pathway (AOP) Development I: Strategies and Principles. Toxicol. Sci. 142, 312-320. https://doi.org/10.1093/toxsci/kfu199

40. Malchoff, D.M., Brufsky, A., Reardon, G., McDermott, P., Javier, E.C., Bergh, C.H., Rowe, D. 1993. A mutation of the glucocorticoid receptor in primary cortisol resistance. J. Clin. Invest. 91, 1918–1925. doi: 10.1172/JCI116410

41. Kino, T. 2018. Single nucleotide variations of the human GR gene manifested as pathologic mutations or polymorphisms. Endocrinology 159, 2506-2519. doi: 10.1172/JCI116410.

363   42. Ravi, V., Venkatesh, B. 2018. The Divergent Genomes of Teleosts. Annu Rev Anim
364   Biosci. 6, 47-68. doi: 10.1146/annurev-animal-030117-014821.
365   43. https://www.earthbiogenome.org/

Figure 1. A. Image of cortisol docking into the *Pygocentrus nattereri* AOA3B4D0J0
glucocrticoid receptor with a binding affinity -8.43 Kcal/mol and the corresponding amino
acid interaction fingerprint. B. The predicted binding affinity for all chemical and species
combinations. Inset, the top 5 species + chemical binding affinities (see supplementary
information for values).

 Figure 2. Binding affinities for the 163 teleost fish glucocorticoid receptors per chemical (see
 supplementary information for values)

Table 1 The five species with the highest binding affinity to each chemical.

Chemical	Class	Species Rank (*Latin name* and UniProt ID) of 5 most sensitive to each chemical [ΔG (kcal/mol)]				
		1	2	3	4	5
Cortisol	Natural steroid	*Esox lucius* A0A6Q2WV38 (-10.6856)	*Coregonus maraena* A0A0D6E1Q9 (-10.456)	*Anabas testudineus* A0A7N6BVF0 -10.4278	*Salmo trutta* A0A673WVI2 -10.2504	*Scophthalmus maximus* A0A2U9C6P4 -10.1376
Atrazine	Herbicide	*Anabas testudineus* A0A7N6B9C8 -6.216	*Mastacembulus armateus* A0A7N9ALU0 -6.0674	*Scophthalmus maximus* A0A2U9C6P4 -6.0612	*Poecilia lapipinna* A0A3B3V5Z8 -6.0596	*Xiphororous maculatus* M4AXU0 -6.0576
Beclomethasone	GC	*Esox lucius* A0A6Q2WV38 -10.6038	*Coregonus maraena* A0A0D6E1Q9 -10.3024	*Anabas testudineus* A0A7N6BVF0 -10.1674	*Mastacembulus armateus* A0A7N8X2D0 -9.9236	*Salmo trutta* A0A673WVI2 -9.7614
Clobetasol	GC	*Esox lucius* A0A6Q2WV38 -11.2956	*Coregonus maraena* A0A0D6E1Q9 -11.1776	*Scophthalmus maximus* A0A2U9C6P4 -10.7898	*Anabas testudineus* A0A7N6BVF0 -10.746	*Salmo trutta* A0A673WVI2 -10.742
Clofenotane,	Insecticide	*Oncorhynchus mykiss* Q6RKQ3 -8.1072	*Nothobranchius pienaari* A0A1A8M2D1 -8.0478	*Xiphororous maculatus* M4AXU0 -8.0214	*Scophthalmus maximus* A0A2U9C6P4 -7.9704	*Sinocyclocheilus rhinocerous* A0A673HKV4 -7.91
Dexamethasone	GC	*Esox lucius* A0A6Q2WV38 -11.144	*Coregonus maraena* A0A0D6E1Q9 -11.0824	*Xiphororous maculatus* M4AXU0 -10.9068	*Scophthalmus maximus* A0A2U9C6P4 -10.8816	*Salmo trutta* A0A673WVI2 -10.7178
Flumetasone	GC	*Esox lucius* A0A6Q2WV38 -11.4008	*Coregonus maraena* A0A0D6E1Q9 -11.2524	*Xiphororous maculatus* M4AXU0 -11.1978	*Scophthalmus maximus* A0A2U9C6P4 -11.1266	*Oncorhynchus mykiss* Q6RKQ3 -11.0974
Glyphosate	Herbicide	*Amphilophus citrinellus* A0A3Q0S1C3 -4.7726	*Anabas testudineus* A0A7N6BVF0 -4.7142	*Hucho hucho* A0A4W5NMM7 -4.6994	*Ictalurus punctatus* A0A2D0SEE8 -4.6884	*Electrophorus electricus* A0A4W4EHL7 -4.6826
Halcinonide	GC	*Coregonus maraena* A0A0D6E1Q9 -12.1068	*Esox lucius* A0A6Q2WV38 -11.7716	*Xiphororous maculatus* M4AXU0 -11.4424	*Salmo trutta* A0A673WVI2 -11.3964	*Astatotilapia calliptera* A0A3P8PIE4 -11.3412
Mapracorat	GC	*Electrophorus electricus* A0A4W4EII3 -10.0314	*Stegastes partitus* A0A3B5AC37 -9.8402	*Salarias fasciatus* A0A672FRL6 -9.3478	*Stegastes partitus* A0A3B4ZYN2 -9.2704	*Anabas testudineus* A0A7N6BVF0 -9.1282
Momentasone	GC	*Salarias fasciatus* A0A672FRL6 -10.371	*Coregonus maraena* A0A0D6E1Q9 -10.138	*Esox lucius* A0A6Q2WV38 -10.1016	*Scophthalmus maximus* A0A2U9C6P4 -9.8704	*Anabas testudineus* A0A7N6BVF0 -9.6366
Osilodrosat	Synthesis Inhibitor	*Nothobranchius pienaari* A0A1A8MV28 -7.8638	*Nothobranchius kadleci* A0A1A8C987 -7.8162	*Nothobranchius kuhntae* A0A1A8HYB3 -7.7868	*Scophthalmus maximus* A0A2U9C6P4 -7.7864	*Iconisemion striatum* A0A1A7WDX0 -7.7602
Prednicarbate	GC	*Paralichthys olivaceus* O73673 -7.7606	*Coregonus maraena* A0A0D6E1Q9 -7.7018	*Esox lucius* A0A6Q2WV38 -7.6118	*Electrophorus electricus* A0A4W4EII3 -7.6044	*Salmo trutta* A0A673WVI2 -7.5676
Prednisolone	GC	*Esox lucius* A0A6Q2WV38 -10.5004	*Anabas testudineus* A0A7N6BVF0 -10.2672	*Coregonus maraena* A0A0D6E1Q9 -10.184	*Salmo trutta* A0A673WVI2 -9.9726	*Scophthalmus maximus* A0A2U9C6P4 -9.9648
Sevoflurane	Anaesthetic	*Pundamilia nyererei* A0A3B4GAN3 -6.0414	*Electrophorus electricus* A0A4W4EIA3 -5.9786	*Hucho hucho* A0A4W5NM33 -5.8264	*Hucho hucho* A0A4W5NMM7 -5.7428	*Seriola lalandi dorsalis* A0A3B4WZ32 -5.5394
Triamcinolone	GC	*Esox lucius* A0A6Q2WV38 -11.198	*Coregonus maraena* A0A0D6E1Q9 -10.8506	*Anabas testudineus* A0A7N6BVF0 -10.7634	*Salmo trutta* A0A673WVI2 -10.5826	*Scophthalmus maximus* A0A2U9C6P4 -10.4684
Triclocarban	Antimicrobial	*Hucho hucho* A0A4W5NM33 -8.4626	*Hucho hucho* A0A4W5NMM7 -8.1526	*Electrophorus electricus* A0A4W4EII3 -8.1472	*Fundulus heteroclitus* A0A3Q2Q9J2 -7.9158	*Gasterosteus aculeatus* G3QBY4 -7.9082

Species order colour code: Escociformes & Salmoniformes; Pleuronectiformes; Anabantiformes; Synbranchiformes; Cyprinodontiformes; Cichliformes; Cypriniformes;

Siluriformes; Blenniiformes; Gymnotiformes; Carangiformes; Scorpaeniformes.