Proteomics-based models of gene expression and cellular control of cotton fiber

development

Running title:

Linking proteome dynamics with fiber development

Keywords: G*ossypium hirsutum*, trichoblast, proteomics, machine learning, cellulose synthase,

multi-omics

Youngwoo Lee[1,2], Pengcheng Yang[3], Heena Rani[4], Gideon Miller[3], Corrinne E Grover[5],

Sivakumar Swaminathan[6], Olga A Zabotina[6], Jonathan F Wendel[5], Jun Xie[3], and Daniel B

Szymanski[1,*]

1 Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN, 47907

2 Center for Plant Biology, Purdue University, West Lafayette, IN, 47907

3 Department of Statistics, Purdue University, West Lafayette, IN, 47907

4 Cereal Crops Research Unit, USDA-ARS, 502 Walnut Street, Madison, WI, 53762

5 Department of Ecology, Evolution, and Organismal Biology, Iowa State University, Ames, IA,

50011

6 Roy J Carver Department of Biochemistry, Biophysics and Molecular Biology, Iowa State

University, Ames, IA, 50011

*Correspondence to: szymandb@purdue.edu

Abstract

The cotton textile industry is based on the material properties of terminally differentiated

trichoblasts. Tens of thousands emerge from the seed coat epidermis and march through a

reproducible developmental program that includes interconversion between cell-autonomous and

tissue-based morphogenesis. Proteomic analyses of purified fibers provide a way to associate

molecules with cellular processes that directly affect morphological transitions and material

properties of the harvested cells. However, insufficient temporal sampling and poor protein

coverage have limited the extent to which proteomic data can predict control mechanisms. Here

we quantified the subcellular proteomes of purified fibers daily over a 20-day interval that

includes diverse modes of elongation and cell wall remodeling. Thousands of reliable protein

abundance profiles were assigned to expression groups and tested for associations with a broad

array of phenotypes. These analyses were validated by the presence of proteins with known

localization and stage-specific functions and also include novel functional predictions for

hundreds of genes with no known function. Integration of proteomic and RNA-sequencing data

revealed distinct modes of transcriptional and post-transcriptional control. Collectively

proteomics data resource provides a valuable foundation for large-scale gene function predictions

and provides a path to engineer cotton fiber material properties.

Introduction

Cotton (*Gossypium spp*.) is the world's most important fiber crop and the foundation of a multi-billion dollar renewable textile industry (Huang et al., 2021; Wilkins et al., 2010). Tens of thousands of unicellular seed coat trichoblasts synchronously differentiate and transform their lenth-to-width aspect ratios to ~3000:1 during development (Stewart, 1975). At maturity, each fiber contains a helically patterned cellulose-rich wall that desiccates after programmed cell death to generate harvestable thin, strong, spinnable fibers (Butterworth et al., 2009; Kim and Triplett, 2001; Seagull, 1992). Fiber size, spinnability, and strength are the output of a morphogenetic program of anisotropic diffuse growth (Ryser, 1977; Yanagisawa et al., 2022) with an often underestimated degree of complexity (Figure 1). Phenotyping daily reveals an early tapering phase that is completed by 2 days post anthesis (DPA), during which the hemispherical fiber initial is converted to a thin elongated trichoblast (Graham and Haigler, 2021; Yanagisawa *et al.*, 2022). After tapering the cell wall (CW) thickens as it enters a protracted phase of decelerating anisotropic diffuse growth (Schubert et al., 1973; Wilson et al., 2024), while simultaneously modulating plasmodesmata functions and turgor pressure (Ruan et al., 2001). Given progress in understanding the molecular and biomechanical control of trichoblast morphogenesis (Yanagisawa et al., 2015; Yanagisawa *et al.*, 2022; Yu et al., 2019), there are emerging opportunities to use biotechnological interventions to engineer fiber traits (Liu et al., 2025; Qin et al., 2022; Sun et al., 2025). The goal of this research is to integrate high-temporal resolution quantitative proteomics, RNA-seq, and phenotyping to broadly predict cellular networks that mediate key developmental transitions.

A complexity of studying fiber cell "omics" is their coordinated development with the organ-scale growth and physiology of the enclosing capsule (colloquially termed a "boll") chamber (Figure 1A. and 1B; Supplemental Video 1). Fiber development also includes reversible transitions between cell-autonomous growth and tissue-morphogenesis during which the cells are tightly adhered (Singh et al., 2009). Cotton has long served as a useful model for CW assembly and morphogenesis (Meinert and Delmer, 1977; Seagull, 1993; Tiwari and Wilkins, 1995), and components of the molecular control continue to emerge (Huang *et al.*, 2021). A highly conserved cellulose patterning module underlies polarized diffuse growth in the plant kingdom (Baskin, 2005), and the details of how interactions among cortical microtubules, patterned cellulose synthase (CESA) activity, and CW stress-sensing operate to control cell shape changes

are beginning to emerge (Coen and Cosgrove, 2023; Li et al., 2023). Strong lateral interactions

between microfibrils (Yanagisawa *et al.*, 2015; Yanagisawa *et al.*, 2022; Zhang et al., 2021) and

interactions between fiber and matrix components generate domains of the CW that deform

asymmetrically in response to turgor-generated mechanical stress in the CW (Yanagisawa *et al.*,

2015; Yanagisawa *et al.*, 2022; Zhang *et al.*, 2021). Spatially patterned cortical microtubules

influence microfibril deposition throughout development to maintain cell diameter and pattern a

helical secondary CW (SCW) (Seagull, 1992; Tiwari and Wilkins, 1995; Wilson *et al.*, 2024;

Yanagisawa *et al.*, 2022). Important cotton microtubule-associated proteins are being discovered

using forward and reverse genetic approaches (Chen et al., 2024; Rajangam et al., 2008; Zang et

al., 2021). The microtubule-CESA module does not function in isolation; actin-based

intracellular transport, vesicle trafficking, metabolism and redox control, and hormone signaling

function in concert during fiber development (Huang *et al.*, 2021).

Proteomic profiling provides a way to obtain direct measurements of the protein

machineries and post-translational controls that enable effective morphogenetic work and

coordinate diverse functions, some of which reside in different subcellular compartments.

Historically, proteomic analyses of cotton fibers have identified proteins with altered abundances

as a function of domestication (Bao et al., 2011; Hu et al., 2013; Qin et al., 2017), hormone or

inhibitor responses (Wang et al., 2020a; Yang et al., 2023a), and developmental stages (Jiang et

al., 2022; Mujahid et al., 2016; Zhou et al., 2019). In general, the number of discovered proteins

is low, and temporal sampling of fiber development is limited to 5-day intervals at best. Gel-free

workflows and increasingly sensitive instrumentation enable the quantification of thousands,

rather than dozens, of proteins in a typical experiment (Hu et al., 2015; Lee and Szymanski,

2021; Yang *et al.*, 2023a). This presents an opportunity to gain broad insights into the protein

dynamics that underlie the numerous developmental transitions during fiber development that are

not known.

Here we combined daily sampling and clean cell fractionation procedures to analyze how

protein abundances change in distinct subcellular compartments during fiber development. Our

analytical pipeline allowed us to generate thousands of reliable protein abundance profiles

spanning a broad period of fiber development that encompasses primary cell wall synthesis and

the transition into secondary wall synthesis. Clustering methods identified unique protein

expression groups that reflected specialized functions of the apoplast, cytosol, and membrane-

associated proteins. Multiple groups had clear temporal transitions and broad memberships
across the cell fractions. Correlation analysis with a wide range of phenotypes quantified across
the same timeline provided a way to predict suites of genes that coordinate the unidirectional
progression of fiber development. Integration of RNA-seq and proteomic data can provide large-
scale predictions about general modes of gene expression control (Manzoni et al., 2018; Nie et
al., 2006). These results were validated by the presence of proteins with known localizations and
stage-specific functions in secondary cell wall synthesis. We used combinations of cross-
correlation analysis and machine learning methods to globally predict modes of transcriptional
and post-transcriptional control. Collectively, our resource article provides valuable data and
analytical methods that provide an entry point to broadly analyze the types of proteins and
cellular processes that may control morphology in cottons. Because many of these processes and
machineries are conserved across the plant kingdom, the data have broad utility in the analysis of
cellular growth control.
118
119

Results
Protein profiling of developing cotton fiber by proteomics
Fiber development is reproducible across a wide range of environmental conditions (Jan
et al., 2022; Schubert *et al.*, 1973), including fibers that are cultured in vitro (Meinert and
Delmer, 1977). A central goal was to generate reliable methods to collect time-series data of
protein abundance in multiple subcellular fractions from purified fiber tissue (Figure 1). Methods
to isolate highly purified fiber samples and generate high-quality cell fractions with limited
cross-contamination with broken cells were developed (Lee et al., 2025). The method enables
precursor ion-based quantification of a diverse population of proteins within and between cell
fractions. Here we scaled up the procedure to increase coverage of the crude cytosolic (S200)
fraction. Each replicate included multiple locules from a single boll and included three replicates
daily from 5-24 DPA (Figure 1C). The entire developmental progression, from anthesis at 0 DPA
to boll opening at 48 DPA, is shown in Figure 1A and Supplemental Video 1. Protein yields in
the different cell fractions are included in Data S1. The earliest timepoint was chosen based on
the minimum requirements for adequate protein precipitation from the soluble fractions, and the
final time point was selected in order to capture a complete transition to SCW synthesis.

A crude apoplast (APO_T) fraction was generated after careful dissection and brief

immersion of the intact locules in the isolation buffer. This protocol minimizes contamination

from broken cells and has been shown to generate a distinct apoplast fraction with unique protein

multimerization behaviors and composition compared to cytosol and microsome fractions (Lee *et*

al., 2025). After immersion, fibers were manually dissected, homogenized, and subjected to

differential centrifugation to generate a crude microsomal (P200) and a soluble (S200) fraction

(Figure 1C). Across the 5 to 24 DPA window, 4316 proteins were reproducibly identified and

quantified in 2 of 3 biological replicates of at least one cell fraction, providing the most

comprehensive cotton fiber proteome coverage to date (Figure 1D; Data S3).

Cotton (*G. hirsutum*) is an allopolyploid with two subgenomes (A, D), and as such the

duplicated genes (homoeologs) are encoded by the A- and D-subgenomes, such that they

potentially can be distinguished based on diverged peptide sequences (Hu *et al.*, 2015; Lee and

Szymanski, 2021). Diagnostic homoeolog-specific peptides were detected for ~55% of the cases

and these proteins are annotated with "A" and "D" suffixes following the Locus ID (Data S3).

The mRNA and protein search databases (Supplemental Files S1 & S2) were generated based on

aggregated "A" and "D" subgenome SNPs relative to a base diploid assembly (Grover et al.,

2025). These databases enabled high coverage matches with the LC/MS spectra and accurate

comparisons of transcript and protein abundance profiles for specific homoeologs (see below).

The search databases and locus ID conventions used here were cross-referenced locus IDs across

several commonly used cotton genome assemblies to facilitate gene matching (call out table

exactly here??). The P200 fraction consistently had the most diverse protein populations over the

time course (Data 3C), and, as observed previously (Lee *et al.*, 2025), was more similar to the

APO_T compared to S200. The APO_T fraction primarily consists of membrane-enclosed particles

with a highly heterogeneous composition (Lee *et al.*, 2025). Widespread overlap among the

different cell fractions can occur because many proteins are dually localized (McBride et al.,

2017). Our pipeline generated protein quantification across over 6 orders of magnitude (Figure

1E). The protein types amenable to proteomic analyses are extremely diverse (Data S3). The

coverage of metabolism in this dataset is deep (Supplemental Figure 1) and provides broad

experimental evidence for the specific gene products that operate in central metabolism,

phenylpropanoid biosynthesis (Tuttle et al., 2015; Yang *et al.*, 2023a), fatty acid synthesis (Qin et

al., 2007), and cell wall assembly (Pettolino et al., 2022).

A protein was retained for abundance profiling across the time course if it had non-zero

values in three or more fractions with at least two of them on 2 consecutive days. The protein

population composed of those detected in at least 2 of 3 consecutive DPAs and the subset

detected in 10 or more DPAs had similar intensity distributions (Figure 1E; Supplemental Figure

5E), indicating that signal intensity is not the primary determinant of expression profile

variability. The raw and normalized intensity data and plotted distributions are provided in Data

S3 and Supplemental Figures 2–4. Our deep proteome profiling provided a broad depiction of

compartmentalized protein behaviors during development, identifying diverse groups of proteins

that coordinate fiber development. However, the widespread occurrence of missing values

generated between biological replicates is a widely recognized shortcoming of label-free

quantitative proteomics (Hamid et al., 2022; Jörnsten et al., 2005; Wang et al., 2020b). We

developed a highly stringent data filtering procedure to minimize the effects of artifactual

missing values between DPA and eliminate potentially unreliable profiles and generate a

restricted dataset containing informative expression profiles from three subcellular locales (see

below and methods).

To identify expression groups, the informative profiles were analyzed using a machine

learning-based clustering method termed Self-Organizing Map (SOM) analysis (Kohonen, 1990;

Wehrens and Kruisselbrink, 2018). In an SOM, the input data are mapped onto a discrete

topological grid formed by a set of groups. The groups are typically organized in a 2-dimensional

grid, with rows and columns. As the model is trained on input data, the groups gradually adjust

their weights to become more similar to the pattern of the input data in their neighborhood,

resulting in clusters of the input profile data. The number of SOM groups was determined by

evaluating cluster stability, ensuring that proteins consistently grouped with the same or

neighboring clusters as the number of groups increased. This strategy minimized empty or

single-member clusters, providing meaningful resolution and interpretability. Each SOM group

has a "code" that reflects a central expression pattern of the members (Figure 2A; Data S4). A

group was scored as unreliable if: 1) the code profile was punctuated by missing values across

the timeline; 2) the profile pattern reflected variability in protein coverage in specific DPAs

(Data S3D); or 3) the pattern was defined by strings of 3 adjacent zeros that could simply reflect

random gaps in the data that were not filled by interpolation. Following this filtering, 39 groups

were classified as unreliable, and the remaining 44 potentially informative groups were subjected

to further filtering to remove unreliable members with discordant features in their expression. A

multi-step quality control procedure was then applied to each SOM cluster to exclude outliers,

unreliable expression patterns, and low-abundance proteins, yielding high-confidence protein

expression profiles (Supplemental Figures 2–5). For each group, member intensities were

normalized to a 0–1 range based on their maximum values and averaged to generate

representative expression patterns on a consistent scale (Figure 2B). This process resulted in 44

distinct cell fraction protein expression group patterns (12 in APO_T, 23 in P200, and 9 in S200),

where a 10% change in the normalized scale corresponds to an about 4-fold difference in

abundance (or 2-fold on the log2 scale) due to the high signal intensity of the included profiles.

In the end, these 44 expression groups provide large-scale functional data on gene sets and

cellular activities (Figure 2B; Data S5), and provide clues about gene functions that may operate

during specific developmental windows in particular subcellular locales.

Many of the profiles were non-uniform, with sharp abundance transitions often reflecting

a shift to or from consecutive days of non-detection. Those classified as "increasing" had a

clearly defined timepoint at which the protein transitioned from below to above the detection

limit (Figure 2B). The inverse was true for the profiles classified as "decreasing". We

hypothesize that these clearly varying expression patterns reflect rearrangements of suites of

cellular machineries as the fiber population undergoes developmental transitions. Some non-

uniform expression groups were defined by a specific subcellular compartment. Gene ontology

(GO) enrichment analysis shows that P200 protein expression groups with an increasing pattern

have hydrolase activity, guanyl nucleotide binding, and catalytic activity terms, whereas S200

was enriched in response to stress term (Data S6). Constitutively expressed proteins classified as

non-uniform were flagged if there was at least a 4-fold increase or decrease in the mean protein

abundance of the group members during development (Data S5). These "constitutive

increase/decrease" groups may reflect broad reprogramming of gene expression relative to the

transition to SCW synthesis (Grover *et al.*, 2025) and/or proteins with unequal activities in the

elongation and SCW phases. A large number of proteins had flat relative abundances and were

classified as uniform. These proteins may define part of the "constitutivome" that includes

essential functions that operate across the developmental phases.

Integration of protein dynamics modules with fiber phenotypes

We next wanted to correlate protein expression profiles with phenotypic transitions. To

generate a merged set of profiles aggregating protein abundance dynamics across the cell

fractions, we turned again to SOM. After evaluating SOM configurations with 9, 20, 30, and 40

groups, we selected the 20-groups SOM as it minimized empty or single-member clusters while

maintaining stable group memberships (Data S7A; Supplemental Figures 6–9). This

configuration yielded 15 distinct merged protein expression modules (Figure 3A). 11groups

contained clearly distinct and non-uniform profiles (Figure 3A; Data S7). The mean normalized

intensities of the merged group members were used to estimate the behavior of the group and test

for correlations with phenotypes. Aggregation of 7 developmentally variable fiber phenotypes

from the same or similar developmental windows (Swaminathan et al., 2024; Wilson *et al.*, 2024)

provided a way to test for significant associations between protein expression groups and

phenotypic transitions using correlation analyses (Figure 3C; Data S8).

As an example, the M16 group was most strongly positively correlated with the transition

to SCW synthesis, which is characterized by transitions in cellulose microfibril angle, increases

in cell wall thickness, and an increasing proportion of cellulose in the cell wall (Figure 3C; Data

S8B). M16 contained proteins present in all three subcellular compartments and included many

known proteins involved in CW biosynthesis, as well as many previously uncharacterized

proteins (Figure 3D; Data S8 and S9). M16 also contained *G. hirsutum* orthologs of each of the

SCW-associated CESAs (Grover *et al.*, 2025; Kim et al., 2019) that were induced at ~16 DPA to

similar levels (Figure 4D; Data S9A). M19 was negatively correlated with SCW transition and

may contain proteins that are specifically involved in the fiber elongation and remodeling of the

primary CW (Figure 3C; Data 7C and 8B). M14 was also negatively correlated with SCW

transition, but this group was composed solely of APO_T members. By comparing these data with

those in a purified cotton EV dataset (Lee *et al.*, 2025), we found that M14 members were either

specific to the particulate fraction or dual-localized in the soluble and particulate fractions of the

apoplast (Data 7D). M14 contained more than half of the total APO_T proteome. The particulate

fraction was enriched in spherical stain-excluding compartments that could be visualized using

negative staining and transmission electron microscopy (Figure 3B). To determine if the

apoplastic particles (APO_{p200}) had a unique polysaccharide composition, their glycome profiles

were compared to those of the P200 and APO_{s200} fractions. Several epitopes related to xylan and

xyloglucan were specifically enriched in the APO_{p200} particles compared to the other cell

fractions and many other epitopes were selectively depleted (Data S10). These data provide

further evidence for their unique composition and may reflect their mode of generation. M9 and

M18 were also APO$_T$-specific expression groups with similar profiles (Figure 3C). These profiles

may reflect the developmentally regulated biogenesis of extracellular vesicles.

M20 declined from maximal levels at 5 DPA and was most strongly correlated with the

decelerating relative growth rate and a reduced proportion of pectin (Figure 3C). M20 may

contain groups of proteins associated with a fiber state with a faster relative growth rate. M11

and M17 were anti-correlated with relative growth rate and auxin-response factors, ABA

receptors, and vesicle trafficking machineries (Data S7C, column T, M11) that enable persistent

axial growth and/or early components that prime the cell for the transition to tissue-based

morphogenesis (Singh *et al.*, 2009) or SCW synthesis (Pettolino *et al.*, 2022). Many of the

merged protein groups correlated with phenotypic profiles, and these analyses provide models of

how diverse types of gene products function in multiple cellular locations to coordinate

developmental transitions.

Integration of mRNA and protein dynamics: distinct modes of gene expression control

To globally analyze potential modes of gene expression control, we integrated the

proteomic profile data with fiber transcript data obtained using the same conditions and

developmental time points (Grover *et al.*, 2025). Pearson correlation coefficients (PCC) were

calculated for 6809 genes in which protein profiles could be matched to a single homoeolog (see

methods, case 1) or a pair of homoeologs (Figure 4A; Supplemental Figures 10-12). We initially

focused on proteins with a non-uniform expression pattern, i.e., the mRNA/protein correlations

in the "increasing" or "decreasing" categories that were stronger in the P200 and S200 fractions

than APO$_T$ fraction (Figure 4B; Data S12). About ~ 48 % of mRNA/protein pairs in P200 and

S200 had PCC values > 0.75, and protein abundance could be explained by transcriptional

control. In general, the APO$_T$ proteins in both expression groups were weakly correlated with

their mRNAs because this cell fraction was dominated by putative extracellular vesicle proteins

that may have reduced abundance after ~20 DPA (Figure 3A). In addition to the APO$_T$ groups,

those showing a negative correlation could be explained by the spatial and functional

compartmentalization of gene expression regulation, highlighting the importance of subcellular

localization information to understand mRNA-protein correlations. However, there was evidence

for transcriptionally regulated proteins in the apoplast. A cotton ortholog of secreted signaling
molecule STOMAGEN, usually associated with non-cell-autonomous cell proliferation (Cui et
al., 2014; Sugano et al., 2010), was downregulated at ~14 DPA and correlated with transcript
levels (Data S7).

To conduct a more open-ended test for commonly observed or non-linear patterns of gene
expression control, a weighted SOM clustering was conducted by combining the mRNA and
protein data for 1228 different P200 proteins with informative profiles (Figure 4C; Data S13).
The full clustering result is shown graphically in Supplemental Figure 13. The analysis produced
three groups (V8-V10) with a total of 48 members with coherently increasing protein and mRNA
expression, and three groups (V81,91,92) with a total of 50 members with coherently decreasing
expression (Figure 4C), of which 48 of 98 were captured in the PCC analysis described in Figure
4B. Among the large population of proteins with a more uniform expression, there were 79 genes
in which mRNA levels fell by at least 8-fold over the time course (Figure 4C). These examples
could reflect post-transcriptional controls in which the mRNA translation efficiency increases or
protein stability increases during development. In V30, transcript levels increased 8-fold, but
protein levels were stable (Figure 3C), indicating that translation of the mRNA may be regulated
or the steady state of the protein may be the result of physical interactions with a stabilizing
factor. The SOM-weighted cross-correlation analyses provide quantitative information about
how diverse types and timings of gene expression control operate during fiber development.

To partially validate our analyses with knowns, we queried our list of genes predicted to
be subject to temporal transcriptional control with those known to be involved in CW assembly
or the transition to SCW synthesis (Data S14). For example, among the 67 proteins present in the
M16 module, 44 were predicted to be transcriptionally regulated. Several orthologs of
documented players in SCW synthesis (e.g., cell wall CESAs, COBRAs, and WAVE-
DAMPENED-LIKE2 proteins) were reliably detected in the P200 fraction from fiber cell
extracts (Figure 4D) (Chen *et al.*, 2024; He et al., 2024; Wen et al., 2022). GhCESA4-B, 7-A/B,
and 8-B proteins were first detected with a ~2-day lag compared to when the corresponding
mRNAs first increased. A database of cotton orthologs to predicted Arabidopsis cell wall genes
(Cell Wall Genomics database; https://cellwall.genomics.purdue.edu/) was created, and a subset
of those were present in the set of predicted transcriptionally controlled genes (Data S14). These
data summarized in this resource article provide a glimpse into the regulatory complexities of

fiber development and a source of promoters and gene products that appear to be subject to

transcriptional control and more easily modulated in transformed lines.

Discussion

In this resource article, we combine sensitive and accurate quantitative proteomics with

the synchronized process of cotton fiber morphogenesis to broadly predict protein networks that

orchestrate fiber development. The data are derived from a single cell type that develops in the

insulated environment of the developing capsule. Daily sampling revealed an unexpected

complexity: an extended phase of growth deceleration in early development, followed by waves

of gene activation and suppression that precede the transition to SCW synthesis and multiple

phases of cellulose microfibril and cell wall matrix remodeling (Grover *et al.*, 2025; Lee *et al.*,

2025; Swaminathan *et al.*, 2024; Wilson *et al.*, 2024). These transitions are not simply anchored

to a general transition to SCW synthesis. Our proteomic dataset provides a conceptual model of

how networks of proteins in multiple subcellular locations coordinate cellular functions and fiber

phenotypes during development.

The proteomic resources generated here provide experimental support for over 4,300

fiber proteins, and reliable abundance profiles over a wide window of development for more than

2,900. This dataset provides useful time-series data on the localization and abundance of a

functionally diverse population of enzymes, hormone signaling proteins like auxin response

factors and ABA-receptors, as well as proteins that span the full breadth of molecular functions

in the cell. There are many enzymes that are specifically associated with fiber elongation,

including those involved in flavonoid, anthocyanin, and wogonin metabolism (Supplemental

Figure 1; Data S5 and S6). The roles of these small molecules in fiber morphogenesis merit

further analyses. All of the proteomic and associated RNA-seq data are publicly available in a

simple search interface (link to be provided).

The SOM clustering method and cross-correlation with time-resolved phenotypes were

used to generate models of how protein groups in multiple cellular compartments are linked to

specific developmental transitions (Figure 3). The merged profiles captured constitutively

expressed proteins with uniform (M2,M3,M4,M5), increasing (M6,M7), and decreasing

(M9,M10) abundances across the timeline. These proteins likely provide essential functions that

may be more likely to be subject to post-translational controls. Many proteins were assigned to
groups with clear time points of abundance change (M11,M14,M16-M20). This approach
enabled us to identify groups of up- and down-trending proteins that transition at a similar time
point. There were expression group pairs that were most strongly associated with growth rate
variability (M20,M17) and others with the transition to SCW synthesis (M16,M19). Gene
knockout experiments will be needed to determine how accurately these gene products are linked
to these specific developmental transitions.

In the case of the transition to SCW synthesis, many key players are conserved across
species and cell types with a thickened SCW (Kumar et al., 2016). Our dataset contains orthologs
of each of the three CESA clades (Wen *et al.*, 2022) and COBRA-like (He *et al.*, 2024) proteins
that are known to influence SCW synthesis in cotton, and many of them appear to be
transcriptionally regulated (Figures 3D and 4D). SCW patterning is a complex process that
includes a radical reorganization of the cytoskeletal, vesicle trafficking, and cell wall assembly
systems (Hoffmann et al., 2021). This proteomics analysis provides new insights into the
particular CESAs and systems of proteins that operate prior to (M19) and after (M16) the
transition to SCW synthesis. Some of expression groups may reflect reversible transitions in cell-
cell adhesion as fibers interconvert between cell-autonomous and tissue-like morphogenesis at
~5 and ~20 DPA (Singh *et al.*, 2009). These different modes of growth may partially explain
relative growth rate deceleration and the functional roles of protein expression groups correlated
with growth rate control (Figure 3C). A prior publication on the apoplast proteome (Lee *et al.*,
2025) and the time-series analyses here point to the existence of heterogeneously sized
membrane-enclosed compartments in the extracellular space (Figure 3B). The merged expression
group profiles M18 and M14 reflect time points of transition to and from tissue-based
morphogenesis (Figure 3A). We speculate that these vesicle-enclosed compartments cannot
simply diffuse from the plasma membrane to the extracellular space but are generated when cell-
to-cell contact structures that include cytoplasmic components experience nanoscale mechanical
failure as bundled fibers grow at different rates or directions.

Integration of proteomic and RNA-seq data can reveal distinct modes of gene expression
control. These data also define novel temporal relationships between transcriptional
upregulation, protein detection, and associated phenotypes (e.g., increases in cellulose content at
17 DPA). We identified a large set of transcriptionally activated and repressed genes at different

stages of development (Figure 4B and 4C). These lists are likely reliable as they identified

specific *G. hirsutum* loci encoding orthologs of known players (Chen *et al.*, 2024; He *et al.*,

2024; Wen *et al.*, 2022) as well as a broader set of unknown genes that have been annotated to

have CW-related functions (Figure 4D). These gene sets provide a potential toolkit of promoters

and genes that may be effectively used to modulate fiber traits. Other potentially useful gene sets

with constitutive expression or discordant transcript and protein profiles likely reflect unknown

post-translational control mechanisms and will require further research to develop methods to

alter their activities. The validation of these findings is an important next step, but it is not

addressed in this resource article.

Conclusions and future challenges

Fiber development is an experimentally tractable system that has enormous potential in

the context of architectural engineering for enhanced fiber traits. This potentially is likely to be

increased by future improvements to proteomic pipelines. Only ~10% of the expressed genes

were detected here (Figure 4A), and there are too many missing values in the daily data matrix.

In this respect, improved instrumentation and organelle separation methods would facilitate new

insights into the overall proteomic composition and transformation states during cotton fiber

development. Finite element biomechanical models of growing trichoblasts predict key CW

material properties and CW geometry/force-sensing mechanisms underlie morphogenesis (Li *et*

al., 2023; Yanagisawa *et al.*, 2022). Discovery of key controlling genes can be accelerated by

finding better ways to integrate time-series mRNA, proteome, and glycome data with genetic

maps and gene function predictions. The barriers to cotton fiber transformation are diminished,

so it is now possible to validate these predictions at a reasonable scale (Kangben et al., 2024).

These improvements increase the prospects for crop enhancement via cotton fiber engineering.

We anticipate growing proteomic datasets and additional high-dimensional transcriptomic and

metabolomic data (Qin *et al.*, 2022; Sun *et al.*, 2025; Tuttle *et al.*, 2015) will be fully aggregated

in public databases (Yang et al., 2023b; Yu et al., 2021) to fuel machine learning and AI-driven

approaches that broadly predict relationships among genetic maps, molecular features, cellular

functions, and important fiber traits.

Methods

Cotton growth conditions

Gossypium hirsutum cv. TM1 plants were cultivated in a Conviron® E15 growth chamber

(Conviron, Pembina, ND, USA) at the Plant Growth Center in Purdue University as previously

described (Lee *et al.*, 2025). Seeds were sown in a 3-gallon pot with a soil mixture composition

prepared as 4:2:2:1 soil:perlite:bark:chicken grit. The growth chamber was controlled to retain

50-70 % of relative humidity and remain with a day/night setting of 28/23 °C and 16/8 h at a

light intensity of 500 µmol m^{-2} s^{-1} (fluorescent lamps of 28 Sylvania F72T12/CW/VHO 100W

and 4 Sylvania F24T12/CW/HO 35W; incandescent bulbs of GE 60W light 130V A19). The 30

mins two-step ramp-up and 30 mins two-step ramp-down periods (15 minutes at 166 µmol

photons + 15 minutes at 336 µmol photons) were programmed at the daytime beginning and the

ending, respectively. At anthesis, flowers were marked as 0 day post anthesis (0 DPA). Three

bolls were collected daily from 5 to 24 DPA for a total of 60 samples (20 days × 3 replicates).

The harvested cotton bolls were immediately maintained on ice and carefully dissected to get

four intact ovules from each boll.

Cell fractionation into apoplast (APO$_T$), microsome (P200), and cytosolic (S200) fractions

The APO$_T$, P200, and S200 fractions were prepared as previously described (Lee *et al.*,

2025). One intact ovule (700 mg) in each locule was dipped in 5 mL of pre-chilled microsome

isolation buffer (MIB) [50 mM Hepes/KOH (pH 7.5), 250 mM sorbitol, 50 mM KOAc, 2 mM

Mg(OAc)$_2$, 1 mM EDTA, 1 mM EGTA, 1 mM dithiothreitol (DTT), 2 mM PMSF and 1% (v/v)

protein inhibitor cocktail (160 mg/mL benzamidine-HCl, 100 mg/mL leupeptin, 12 mg/mL

phenanthroline, 0.1 mg/mL aprotinin, and 0.1 mg/mL pepstatin A)]. After 10 minutes of

incubation under gentle shaking, the resulting buffer solution was recovered through 2 layers of

cheesecloth. Then, the entire APO$_T$ fraction was used for protein precipitation using an acetone

precipitation method.

For P200 and S200 fractions, fibers were directly isolated from the ovule material used

for the above APO$_T$ collection (Lee *et al.*, 2025). Purified fresh fiber tissues were homogenized

under the cold MIB (sample to MIB ratio was 1:4) using Polytron homogenizer (Brinkmann

Instruments, New York, NY, USA) with chilled blade tip with 10 seconds grinding, 1 minute rest

on ice, and another 10 seconds grinding. The homogenate was filtered with 4 layers of

cheesecloth pre-soaked in cold MIB, and the cheesecloth was further squeezed to get a residual

sample in the cheesecloth. Then the filtered homogenate was further centrifuged on an Allegra

X-30R centrifuge (Beckman Coulter Life Sciences, Indianapolis, IN, USA) at 1,000 x g for 10

min to remove debris. The supernatant was fractionated at 200 k x g for 20 minutes at 4°C using

a Beckman Optima Ultracentrifuge with TLA110 rotor (Beckman Coulter Life Sciences,

Indianapolis, IN, USA) into a cytosolic (S200) fraction and pellet. The resulting pellet was

washed with cold MIB, incubated on ice for 10 minutes, and ultracentrifuged at the same

condition as shown above, and the washing step was repeated. The pellet was defined as a

microsome fraction (P200) and directly subjected to protein precipitation using an acetone

precipitation method. Protein yield information for the three cell fractions is provided in Data S1.

Determination of protein concentration

For the APO_T and S200 cellular fractions, dried pellets in the acetone precipitation were

dissolved and denatured in 100 μL of 8 M Urea for 1 hour at room temperature. For the P200

fraction, 200 μL of 8 Urea was directly added over the final pellet and incubated for 1 hour at

room temperature to denature proteins from membranes. Undissolved debris was removed by

centrifugation at 12 k x g for 15 min using an Allegra X-30R centrifuge (Beckman Coulter Life

Sciences, Indianapolis, IN, USA). Protein assay was performed using a BCA assay kit, according

to the manufacturer's protocol (Thermo Fisher Scientific, Waltham, MA, USA).

Electron and fluorescence microscopy of isolated extracellular vesicles (APO_{p200})

A freshly prepared APO_{p200} pellet isolated from a 9 DPA apoplastic wash solution was

isolated as described above and diluted 100 times in phosphate buffered saline (pH 7). One

microliter of the resuspended vesicles representing APO_{p200} was negatively stained with 1%

uranyl acetate on plasma-treated copper grids. Imaging was performed using an accelerating

voltage of 40 to 100kV on a Phillips CM-100 Transmission Electron Microscope (Philips/FEI

Corporation, Eindhoven, Holland).

Glycome profiling

APO_T was fractionated into APO_{S200} and APO_{P200} fractions by ultracentrifugation as

described previously (Lee et $al.$, 2025). The supernatant (APO_{S200} fraction) after

ultracentrifugation was lyophilized completely. The pellet containing apoplastic vesicles

(APO$_{P200}$ fraction) were lysed by sonication and ultracentrifuged to collect the soluble fraction

with the released polysaccharides (Okekeogbu et al., 2019). The lyophilized APO$_{S200}$ and

APO$_{P200}$ fractions were redissolved in water, the sugar contents were measured, and 2 µg of

polysaccharides were spotted in each well of a 96-well plate and probed with 71 different

antibodies as described previously (Swaminathan *et al.*, 2024). The crude microsomes obtained

from lysed fiber cells (P200) were processed identically and used as an additional control.

LC-MS/MS sample preparation

For LC-MSMS analysis, proteins were digested using trypsin as described previously

(Lee *et al.*, 2025; McBride et al., 2019). We used all protein samples we gained from the APO$_T$,

and S200 preparations, while 50 µg of proteins in the P200 cell fractions. Denatured protein

samples were reduced in 10 mM DTT for 45 min at 60°C and then alkylated with 20 mM

iodoacetamide for 45 min at room temperature under the dark. The urea concentration of the

solution was brought down to below 1.5 M by adding 50 mM NH$_4$HCO$_3$ before trypsin T6567

(Sigma-Aldrich, St. Louis, MO, USA)) digestion. The digestion was proceeded at an enzyme-to-

protein ratio of 1 to 50 at 37 °C. After overnight digestion, trifluoroacetic acid (TFA) was added

to end digestion. The digested peptides were purified using C18 Micro Spin Columns 74-4601

(Harvard Apparatus, Holliston, MA, USA). One µg of each peptide sample was injected onto the

LC-MS/MS system.

LC-MS/MS data acquisition

LC-MS/MS analysis was carried out as described previously (Barabas et al., 2019).

Peptide samples were analyzed by a reverse-phase LC-ESI-MS/MS system using a Dionex

UltiMate 3000 RSLCnano System coupled with the Orbitrap Fusion Lumos Tribrid Mass

Spectrometer (Thermo Fisher Scientific, Waltham, MA, USA). Peptides were loaded onto a trap

column (300 µm ID × 5 mm) packed with 5 µm 100 Å PepMap C18 medium and then separated

on a reverse phase column (75 µm ID × 50 cm) packed with 2 µm 100-Å PepMap C18 medium

(Thermo Fisher Scientific, Waltham, MA, USA). Peptides were resolved at a flow rate of 200

nL/min over a 160-min LC gradient. All the data were acquired in the positive ion mode in the

Orbitrap mass analyzer using a higher energy collisional dissociation (HCD) fragmentation

scheme. The MS scan range was from 375 to 1,500 m/z at a resolution of 120,000. The automatic

gain control target was set as standard, maximum injection time (50 ms), dynamic exclusion 60

s, and intensity threshold 5.0e3. MS data were acquired in data-dependent mode with a cycle

time of 3 s/scan.

Peptide Identification and Quantification

Andromeda search engine on MaxQuant (version 1.6.14.0) was used for relative protein

abundance quantification and protein identification (Tyanova et al., 2016). The search was

conducted as described (Lee et al., 2021). To minimize locus identity (Locus ID) assignment

errors between cell fractions LC/MS raw files were searched in a single MaxQuant job, and

signal intensities were normalized (Cox et al., 2014; Tyanova *et al.*, 2016) across all time points

(Data S2). The three different parameter groups for the three cellular fractions were defined to

specify different parameters for each cell fraction. The search parameters were as follows: Match

between runs was set with the default setting; one percent of the Protein and Peptide false

discovery rate (FDR) was chosen; 1 was chosen for Min. unique peptides; Label-free

quantification was selected within each parameter group; all other parameters were set as default.

The species-specific and homoeolog-diagnostic cotton reference was built (Grover *et al.*, 2025)

using *G. raimondii* genome annotation to facilitate transcriptome (Grover *et al.*, 2025) and

proteome data integration. The reference (Supplemental File S2) includes species/homoeolog-

specific SNP and had been validated as performing well in *G. hirsutum* (Hu et al., 2021). To

promote consistent locus ID assignments between annotated *Gossypium hirsutum* genome

annotations, a locus ID conversion table is provided. Identified peptides, proteins, and a locus ID

conversion table can be found in Data S2.

Data filtering for protein identification in each cellular fraction (APO$_T$, P200, S200)

Two layers of filtering strategies were applied for each cellular fraction dataset to identify

proteins in each cellular fraction as described before (Lee *et al.*, 2025). First, those proteins

identified with MS/MS count(s) $\geq$ 1 in at least 2 out of 3 replicates were chosen as identified

proteins in each cellular fraction. Secondly, quantified proteins with valid LFQ values in at least

2 out of 3 replicates in each cellular fraction were selected from the above count-based datasets.

These reproducible proteins defined by MS/MS count(s) and LFQ criteria are provided in Data

S3A and S3B, respectively.

Profile interpolation

Up to two missing values, which are denoted as zeros, at the two ends or in the middle of

a profile were interpolated. In the middle of a profile, a single zero was interpolated by averaging

its neighboring non-zero values. In cases where two consecutive zeros occur within an

abundance profile, a linear interpolation method was applied. At the two ends of a profile, a

single zero was replaced by its neighboring non-zero value, while two consecutive zeros were

directly replaced by the nearest non-zero value. The interpolated profiles were used in

downstream analyses and are available in Data S3C.

Self-organizing maps (SOM) for protein expression patterns

Self-organizing maps (SOM), a robust unsupervised machine learning method (Kohonen,

1990), were used for analyzing protein expression patterns. We applied a SOM algorithm

(Wehrens and Kruisselbrink, 2018), based on an inner product distance metric, to cluster the

interpolated protein profiles of each cell fraction. Choosing the number of groups in the SOM

analysis involves a trade-off. A large number of groups can lead to empty clusters or clusters

with only a single member, while too few groups may fail to capture the underlying clustering

structure in the data. We selected a balanced number of groups that preserves meaningful cluster

structure while avoiding empty or single-member clusters. The SOM clusters defined the

corresponding protein expression profile patterns (Figure 2A). The SOM clustering results for

the protein profiles in each group are available in Data S4 and Supplemental Figures 2–4. Cluster

profiles were flagged as unreliable if they reflected the overall distribution pattern of protein

coverage (Data S3D) or were dominated by patterns that can be explained by the interpolation,

e.g., a gap of three consecutive zeros. After this inspection, 12 groups in APO_T, 23 groups in

P200, and 9 groups in S200 fractions were selected as containing informative protein profiles

(Data S4).

Quality control for reliable protein expression patterns

To ensure reliable protein expression patterns, a multi-step quality control procedure was

applied to each SOM cluster to remove outliers, unreliable profiles, and low abundance profiles

(Supplemental Figures 2–4).

Step 1: Outlier detection

Outliers were identified by calculating all pairwise similarities defined by the inner

product of two profiles in a cluster:

$$\frac{x_i^T x_j}{\sqrt{x_i^T x_i}\sqrt{x_j^T x_j}}$$

If the similarity between a protein profile within a group and the group centroid was less than

either the group's median similarity or 0.8, the protein was treated as an outlier and removed

(Supplemental Figures 2–4, Step 1).

Step 2: Removal of unreliable profiles

Unreliable profiles were defined as those with isolated non-zeros, including profiles with

a single non-zero value flanked by zeros or three consecutive zeros in the middle of the

expression pattern. These profiles were removed from the clusters (Supplemental Figures 2–4,

Step 2).

Step 3: Filtering based on abundance

To further validate the clustering and filtering steps, we conducted a detailed analysis to

assess the relationship between protein expression classes–including non-reproducible proteins,

outliers (Step 1), unreliable profiles (Step 2), and reliable profiles–and signal intensity

(Supplemental Figure 5). Proteins with below median abundance values across DPAs were more

likely to exhibit missing values and contain more zeros in their profiles. When the median

abundance dropped below 20 (log2-transformed) for APO_T, 19 for P200, and 20 for S200, these

proteins were considered low abundance and removed. These cutoffs were derived from the

median abundance values of proteins excluded during the earlier steps (Supplemental Figures 2–

4, Step 3).

Informative cell fraction protein expression group patterns

To generate representative expression patterns for each protein groups, member

intensities were normalized to a 0–1 range based on their own maximum values within the group

(Figure 2B). The normalized profiles were then averaged to produce a consistent scale across all

patterns. This procedure resulted in 44 distinct cell fraction protein expression group patterns,

including 12 in APO_T, 23 in P200, and 9 in S200, as detailed in Data S5. In this dataset, a 10 %

increase or decrease in a normalized (0–1) profile corresponds approximately to a 4-fold changed

(equivalent to a 2-fold change in the log2 scale), because the reliable profiles used in these

protein expression groups have averaged protein abundances of at least 20 in the log2-

transformed values.

Merging protein expression patterns for data integration.

Similar protein expression patterns were merged into modules through additional SOM

clustering analysis, using the 44 previously generated protein expression group patterns. We

tested different SOM methods with 9, 20, 30, and 40 groups to optimize group number in the

SOM analysis (Data 7A; Supplemental Figures 6–9). As the number of groups increased, the

SOM produced more empty clusters or clusters containing only a single group pattern, typically

with flat profile patterns, while cluster memberships remained unchanged. Therefore, we

selected the 20-cluster SOM method, which defined 15 distinct clusters as merged protein

expression modules (Figure 3A). The profiles of these merged modules, along with the proteins

in each module, were used as protein phenotypes in data integration analyses with phenotypic

and transcriptomic data, which are available in Data S7B and S7C.

Correlation analysis and classification of mRNA and protein pair

In this study, we used a published cotton fiber transcriptome dataset, which profiled

transcripts present in developing fiber tissues obtained in the same time window (Grover *et al.*,

2025). We downloaded Supplementary Table 1 which includes transcript profiles of 74,776 genes

and further filtered out low abundance transcripts with less than one transcript per kilobase

million (TPM) to generate high-quality, reliable transcriptome information. Data S11 provides

the filtered 50,512 protein-coding genes used in this comparison analysis.

MaxQuant assigns proteins into a single protein group if they share all their peptides with

other proteins (Tyanova *et al.*, 2016), so ambiguously identified homoeologous or orthologous

proteins within a protein group can limit the ability to link proteins to their corresponding

mRNAs. To address this, we used unique peptide information to assign representative proteins to

each protein group. In this assignment, if proteins were detected by at least a quarter of the

maximum number of unique peptides found in the leading protein of a protein group, those

proteins were considered identified for that group. Isoform identifications were then condensed

to representative gene model IDs used in the transcriptome dataset.

631   Based on the homoeolog/ortholog assignment results, protein and mRNA pairs were

classified into three cases for Pearson correlation coefficient (PCC) analyses. Case 1 includes

homoeologs that were identified unambiguously, where unique peptides map to only one

homoeolog. Case 2 involves cases in which unique peptides map to both homoeologs; in this

case, we retained the homoeolog with the higher PCC. Case 3 includes homoeologs and

orthologs identified by unique peptides that map to multiple homoeologs and more than one

ortholog within a protein group. Using this protein assignment (annotated in Data S4), we

integrated the protein profiles (in protein expression groups shown in Figure 2B) with their

mRNA profiles and calculated PCCs between mRNA-protein pairs in each of the cell fractions

(Supplemental Figures 10–12; Data S12).

RNA-protein relationship patterns via SOM

643   To identify relationships between mRNAs and proteins that exhibit constitutively

expressed profile patterns, P200 proteins in Case 1 with flat profiles were paired with their

corresponding transcripts. We applied SOM again to cluster these pairs of protein and mRNA

profiles. In this analysis, 100 groups (10 rows and 10 columns) were used (Supplemental Figure

13). The SOM clusters defined the relationship patterns between mRNAs and proteins over

cotton development. Details of the clustering results and the profiles of paired mRNAs and

proteins are provided in Data S13.

Statistic tests and data analysis

652   Statistical analysis was performed using R version 4.2.0 (R Core Team, 2018) on RStudio

653   2022.07.1 (RStudio Team, 2018). The Flexible Self-Organizing Maps in Kohonen 3.0 package

654   for R (Wehrens and Kruisselbrink, 2018). Microsoft Excel on Office 365 for Mac was used to

655   organize and display the analyzed data.

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository (Perez-Riverol et al., 2021) with the dataset identifier PXD051704.

Code availability

The source codes for the protein SOM clustering analyses and the weighted cross-correlation SOM clustering analysis are publicly available on GitHub (https://github.com/yangpengchengstat/Proteomics-based-models-of-cotton-fiber-protein-dynamics-and-gene-expression-control).

Acknowledgements

This material is based upon work supported by the National Science Foundation under Grant No. 1951819 to D.B.S. We would like to thank Dr. Uma Aryal at the Purdue Proteomics Facility for running the LC/MS samples. We are grateful to Eileen L. Mallery for growing cotton plants and to Dr. Pragya Barua for helping with sample preparation.

Contributions

Y.L.: Methodology, Formal analysis, Investigation, Data Curation, Writing - Original Draft, Visualization

P.Y.: Methodology, Data Curation, Software, Formal analysis, Writing - Review & Editing

H.R.: Investigation

C.E.G.: Investigation, Writing - Review & Editing

S.S.: Investigation, Writing - Review & Editing

G.M.: Data Curation

J.X.: Methodology, Formal analysis, Writing - Review & Editing, Supervision, Funding acquisition

O.A.Z.: Writing - Review & Editing, Supervision, Funding acquisition

J.F.W.: Writing - Review & Editing, Supervision, Funding acquisition

D.B.S.: Conceptualization, Methodology, Formal analysis, Investigation, Writing - Original

Draft, Writing - Review & Editing, Supervision, Project administration, Funding acquisition

Competing interests

The authors declare no competing interests.

Figures

Figure 1. Deep proteome profiling across fiber development in cotton (*G. hirsutum*) plants:
cell fractionation and quantitative proteomics.

(A) Key developmental stages of cotton fibers adapted and re-illustrated based on published
descriptions (Grover *et al.*, 2025; Swaminathan *et al.*, 2024; Wilson *et al.*, 2024). Flower
opening day is defined as 0 days post anthesis (DPA).

(B) Phenotypes of flowers, dissected bolls, and combed fibers at selected time points for
proteomics analysis (5 to 24 DPAs). The white arrow indicates the opening of the boll at
48 DPA. One sepal was removed from the flower for visualization. A movie of cotton
flower and boll development is available in Supplemental Video 1.

(C) Cell fractionation of triplicate samples for quantitative proteomics analysis (Lee *et al.*,

2025).

(D) Proteome coverages across the three different subcellular compartments.

(E) Distributions of average protein abundances across DPAs showing missing values.

APO_T: apoplast; P200: crude microsome; and S200: soluble fractions. Parentheses show

numbers of proteins in each cellular proteome.

Figure 2. Generating a core set of protein expression modules during developing cotton fibers.

(A) Grouping of similar protein profiles in the three different subcellular proteomes using a machine learning-based self-organizing map (SOM) approach. The number of SOM groups was selected based on group density and distribution to minimize empty groups or multiple groups with similar patterns.

728   (B) A set of informative protein expression groups across fiber development in the three

729   different cellular fractions.

731   APO_T: apoplast; P200: crude microsome; and S200: soluble fractions. Parentheses show

732   numbers of proteins in each cellular proteome or each group.

Figure 3. Integration of protein dynamics modules with fiber phenotypes.

(A) Protein expression modules merged from the APO$_T$, P200, and S200 cellular fraction protein expression groups. The number of SOM groups was selected based on group

density and distribution to minimize empty groups or multiple groups with similar

patterns. Empty SOM groups were not visualized.

(B) M14 protein expression group reflects developmentally regulated extracellular vesicles.

The proteomics pipeline used to identify APO_{P200} and APO_{S200} proteins (Lee *et al.*, 2025).

TEM images of vesicles present in APO_{P200}. Overlap of APO_T proteins in M14 with

APO_{P200} proteins (Data S7D).

(C) Integration analyses of fiber phenotypes and protein network modules. Pearson

correlation coefficient (PCC) between protein patterns and phenotypes (Swaminathan *et*

al., 2024; Wilson *et al.*, 2024) was calculated. Module profiles show the merged protein

expression profiles.

(D) M16 protein expression module shows similar and unique compartment specific profiles

that are associated with cellulose content (proportional) increase across development. A

full list of proteins in the M16 can be found in Data S9A.

APO_T: apoplast; P200: crude microsome; and S200: soluble fractions. Parentheses show

numbers of proteins in each cellular proteome or each group.

Figure 4. Global analysis transcript and protein profiles across fiber development.

(A) Coverages of proteins present in the cell fraction protein expression groups over the cotton fiber transcriptome data (Grover *et al.*, 2025). Maximum mRNA and protein abundances across fiber development were visualized. The filtered transcriptome dataset is available in Data S11.

(B) Pearson correlation coefficient (PCC) between mRNAs and proteins (Case 1 proteins; please see the methods) present in the protein expression groups with up or down profiles in Figure 2B.

(C) SOM-based weighted cross-correlation of paired mRNA and proteins that have flat profiles across development. Protein patterns were defined in Figure 2B, and mRNA patterns were defined using fold change (8-times abundance change). The entire SOM result can be found in Data S13 and Supplemental Figure 13.

(D) Transcriptionally controlled known protein markers that function in secondary cell wall synthesis.

Example mRNA and protein abundance profiles of transcriptionally regulated genes present both in the M16 module and assigned to the V8, V9, V10, and V92 groups of the weighted cross-correlation SOM in Figure 4C. A broader list of the predicted cell-wall transcriptionally regulated dataset of the abundance profiles of proteins across fiber development and PCC values are provided in Data S14. At and Dt indicate homoeologs for the A and D genomes.

APO$_T$: apoplast; P200: crude microsome; and S200: soluble fractions.

Supplemental information

Supplemental Figure 1. Enzyme coverages of metabolic pathways.

Supplemental Figure 2. SOM grouping and profile filtering (APO$_T$).

Supplemental Figure 3. SOM grouping and profile filtering (P200).

Supplemental Figure 4. SOM grouping and profile filtering (S200).

Supplemental Figure 5. Analysis of proteins removed at each data filtering step while identifying

the subset of reliable members.

Supplemental Figure 6. Merged SOM protein expression modules (9 cluster model).

Supplemental Figure 7. Merged SOM protein expression modules (20 cluster model).

Supplemental Figure 8. Merged SOM protein expression modules (30 cluster model).

Supplemental Figure 9. Merged SOM protein expression modules (40 cluster model).

Supplemental Figure 10. Profiles of paired mRNA and proteins (APO$_T$).

Supplemental Figure 11. Profiles of paired mRNA and proteins (P200).

Supplemental Figure 12. Profiles of paired mRNA and proteins (S200).

Supplemental Figure 13. Weighted cross-correlation SOM of paired mRNA and proteins (P200).

Supplemental Video 1. A movie of the cotton flower and boll development (-5 to 53 DPAs).

Supplemental File 1. The mRNA search databases (AD1_AtDt.transcripts.fa).

Supplemental File 2. The protein search databases (AD1_AtDt.pep.fasta).

Data S1. Fiber protein yields in the APO$_T$, P200, and S200 fractions.

• The protein amount yielded in each DPA from three different cellular fractions was measured

by BCA assay.

Data S2. Raw abundances of peptides and proteins (APO$_T$, P200, and S200).

• Raw peptide and protein abundance information obtained from a MaxQuant search were
reported.
• Locus ID converter from Gorai to Gohir ID system
818
Data S3. Raw and interpolated protein profiles across fiber development (APO$_T$, P200, and
S200).
• The averaged abundances of proteins identified by MS/MS count(s) > 0 in at least 2 out of
the 3 replicates and quantified by LFQ > 0 in at least 2 out of the 3 replicates
• **Interpolated protein profiles**
• Numbers of proteins at each filtering step, i.e. raw (unfiltered), MS/MS, and LFQ.
825
826 Data S4. SOM clustering results and filtering for generating protein expression group profiles
827 (APO$_T$, P200, and S200).
828 • SOM group profiles in each cell fraction
829 • **Profiles of members in each SOM group and data filtering were summarized.**
830
831 Data S5. SOM protein expression group profiles (APO$_T$, P200, and S200).
832 • **The SOM groups with their profiles in the three cell fractions were reported and**
833 **visualized in Figure 2B.**
834
835 Data S6. Gene ontology enrichment analysis of P200 and S200 protein expression groups with
836 increasing patterns.
837
Data S7. Merged SOM protein expression modules.
• Optimization of the number of groups defined for SOM-based clustering
• **Informative merged SOM groups with their profiles and profiles of members in each**
merged SOM group were reported. The reliable profiles reported in Data S4 were pooled,
so localization and cell fraction group information can be found in column O.
• M14 APO$_T$ analysis
844
Data S8. Integration analyses of fiber phenotypes and merged protein expression modules.

•   Phenotype estimation data (profiles of each phenotype from 6 to 24 DPA)

•   PCC and p-values of correlation result between phenotypes and merged modules. This was

visualized in Figure 3C.

Data S9. Gene ontology enrichment analysis of the M16 protein expression module.

•   This was visualized in Figure 3D.

Data S10. Glycome profiling of polysaccharide epitopes (APO$_{p200}$ vs. APO$_{s200}$ vs. P200)

.

Data S11. Abundance profiles of transcripts in cotton fiber across development (Grover *et al.*,

2025).

•   Filtered mRNA dataset downloaded from Grover et al, 2025 paper.

Data S12. Analyses of Pearson correlation coefficient (PCC) between mRNAs and proteins

present in cell fraction protein expression groups with trend-up or -down patterns.

Data S13. Weighted cross-correlation SOM of paired mRNA and proteins (P200).

•   SOM group profiles with fold changes in their expression profiles. This was used for data

visualization in Figure 4C.

•   mRNA and protein profiles were assigned in each SOM group

Data S14. Integrative analysis of mRNAs and proteins annotated as CW-related gene families.

•   The list of Cell Wall Related Gene Families was downloaded at

https://cellwall.genomics.purdue.edu/

•   Data integration and correlation analysis results of CW-related gene families with their

mRNA and protein profiles were reported.

References

Bao, Y., Hu, G., Flagel, L.E., Salmon, A., Bezanilla, M., Paterson, A.H., Wang, Z., and Wendel, J.F. (2011). Parallel up-regulation of the profilin gene family following independent domestication of diploid and allopolyploid cotton (Gossypium). Proceedings of the National Academy of Sciences **108**:21152-21157.

Barabas, A.J., Aryal, U.K., and Gaskill, B.N. (2019). Proteome characterization of used nesting material and potential protein sources from group housed male mice, Mus musculus. Scientific Reports **9**:17524. 10.1038/s41598-019-53903-x.

Baskin, T.I. (2005). Anisotropic expansion of the plant cell wall. Annu. Rev. Cell Dev. Biol. **21**:203-222. 10.1146/annurev.cellbio.20.082503.103053.

Butterworth, K.M., Adams, D.C., Horner, H.T., and Wendel, J.F. (2009). Initiation and early development of fiber in wild and cultivated cotton. Int. J. Plant Sci. **170**:561-574.

Chen, B., Tian, Z., Fu, G., Zhang, A., Sun, Y., Wang, J., Pan, Z., Li, H., Hu, D., Xia, Y., et al. (2024). GhWDL3 is involved in the formation and development of fiber cell morphology in upland cotton (Gossypium hirsutum L.). Journal of Cotton Research **7**:6. 10.1186/s42397-024-00167-0.

Coen, E., and Cosgrove, D.J. (2023). The mechanics of plant morphogenesis. Science **379**:eade8055. 10.1126/science.ade8055.

Cox, J., Hein, M.Y., Luber, C.A., Paron, I., Nagaraj, N., and Mann, M. (2014). Accurate proteome-wide label-free quantification by delayed normalization and maximal peptide ratio extraction, termed MaxLFQ. Mol Cell Proteomics **13**:2513-2526. 10.1074/mcp.M113.031591.

Cui, H.C., Kong, D.Y., Wei, P.C., Hao, Y.L., Torii, K.U., Lee, J.S., and Li, J. (2014). SPINDLY, ERECTA, and Its Ligand STOMAGEN Have a Role in Redox-Mediated Cortex Proliferation in the Root. Mol Plant **7**:1727-1739. 10.1093/mp/ssu106.

Graham, B.P., and Haigler, C.H. (2021). Microtubules exert early, partial, and variable control of cotton fiber diameter. Planta **253**:47. 10.1007/s00425-020-03557-1.

Grover, C.E., Jareczek, J.J., Swaminathan, S., Lee, Y., Howell, A.H., Rani, H., Arick, M.A., Leach, A.G., Miller, E.R., Yang, P., et al. (2025). A high-resolution model of gene expression during Gossypium hirsutum (cotton) fiber development. BMC Genomics **26**:221. 10.1186/s12864-025-11360-z.

Hamid, Z., Zimmerman, K.D., Guillen-Ahlers, H., Li, C., Nathanielsz, P., Cox, L.A., and Olivier, M. (2022). Assessment of label-free quantification and missing value imputation for proteomics in non-human primates. BMC Genomics **23**:496. 10.1186/s12864-022-08723-1.

He, Q., Yu, Y., Qin, Z., Duan, Y., Liu, H., Li, W., Song, X., Zhu, G., Shang, X., and Guo, W. (2024). COBRA-LIKE 9 modulates cotton cell wall development via regulating cellulose deposition. Plant Physiol **197**10.1093/plphys/kiae675.

Hoffmann, N., King, S., Samuels, A.L., and McFarlane, H.E. (2021). Subcellular coordination of plant cell wall synthesis. Dev Cell **56**:933-948. 10.1016/j.devcel.2021.03.004.

Hu, G., Koh, J., Yoo, M.J., Chen, S., and Wendel, J.F. (2015). Gene-Expression Novelty in Allopolyploid Cotton: A Proteomic Perspective. Genetics 10.1534/genetics.115.174367.

Hu, G., Koh, J., Yoo, M.J., Grupp, K., Chen, S., and Wendel, J.F. (2013). Proteomic profiling of developing cotton fibers from wild and domesticated *Gossypium barbadense*. New Phytol **200**:570-582. 10.1111/nph.12381.

Hu, G., Grover, C.E., Arick, M.A., II, Liu, M., Peterson, D.G., and Wendel, J.F. (2021). Homoeologous gene expression and co-expression network analyses and evolutionary inference in allopolyploids. Briefings in Bioinformatics **22**:1819-1835. 10.1093/bib/bbaa035.

Huang, G., Huang, J.Q., Chen, X.Y., and Zhu, Y.X. (2021). Recent Advances and Future Perspectives in Cotton Research. Annu Rev Plant Biol **72**:437-462. 10.1146/annurev-arplant-080720-113241.

Jan, M., Liu, Z., Guo, C., and Sun, X. (2022). Molecular Regulation of Cotton Fiber Development: A Review. International Journal of Molecular Sciences.

Jiang, J., Shi, Z., Ma, F., and Liu, K. (2022). Identification of key proteins related to high-quality fiber
in Upland cotton via proteomics analysis. Plant Cell Rep **41**:893-904. 10.1007/s00299-021-02825-y.
Jörnsten, R., Wang, H.-Y., Welsh, W.J., and Ouyang, M. (2005). DNA microarray data imputation and
significance analysis of differential expression. Bioinformatics **21**:4155-4161.
10.1093/bioinformatics/bti638.
Kangben, F., Kumar, S., Li, Z.G., Sreedasyam, A., Dardick, C., Jones, D., and Saski, C.A. (2024).
Phylogenetic and functional analysis of tiller angle control homeologs in allotetraploid cotton. Frontiers
in Plant Science **14**ARTN 1320638
10.3389/fpls.2023.1320638.
Kim, H.J., and Triplett, B.A. (2001). Cotton fiber growth in planta and in vitro. Models for plant cell
elongation and cell wall biogenesis. Plant Physiol **127**:1361-1366.
Kim, H.J., Thyssen, G.N., Song, X., Delhom, C.D., and Liu, Y. (2019). Functional divergence of
cellulose synthase orthologs in between wild Gossypium raimondii and domesticated G. arboreum diploid
cotton species. Cellulose **26**:9483-9501. 10.1007/s10570-019-02744-y.
Kohonen, T. (1990). The self-organizing map. Proceedings of the IEEE **78**:1464-1480. 10.1109/5.58325.
Kumar, M., Campbell, L., and Turner, S. (2016). Secondary cell walls: biosynthesis and manipulation.
Journal of Experimental Botany **67**:515-531. 10.1093/jxb/erv533.
Lee, Y., and Szymanski, D.B. (2021). Multimerization variants as potential drivers of
neofunctionalization. Sci Adv **7**10.1126/sciadv.abf0984.
Lee, Y., Okita, T.W., and Szymanski, D.B. (2021). A co-fractionation mass spectrometry-based
prediction of protein complex assemblies in the developing rice aleurone-subaleurone. The Plant Cell
33:2965-2980. 10.1093/plcell/koab182.
Lee, Y., Rani, H., Mallery, E.L., and Szymanski, D.B. (2025). A cell fractionation and quantitative
proteomics pipeline to enable functional analyses of cotton fiber development. The Plant Journal
121:e17246. https://doi.org/10.1111/tpj.17246.
Li, J., Szymanski, D.B., and Kim, T. (2023). Probing stress-regulated ordering of the plant cortical
microtubule array via a computational approach. BMC Plant Biol **23**:308. 10.1186/s12870-023-04252-5.
Liu, Z., Fan, L., Shu, S., Qanmber, G., Chen, E., Huang, J., Li, F., and Yang, Z. (2025). Cotton
metabolism regulatory network: Unraveling key genes and pathways in fiber development and growth
regulation. Plant Communications **6**10.1016/j.xplc.2024.101221.
Manzoni, C., Kia, D.A., Vandrovcova, J., Hardy, J., Wood, N.W., Lewis, P.A., and Ferrari, R.
(2018). Genome, transcriptome and proteome: the rise of omics data and their integration in biomedical
sciences. Brief Bioinform **19**:286-302. 10.1093/bib/bbw114.
McBride, Z., Chen, D., Reick, C., Xie, J., and Szymanski, D.B. (2017). Global analysis of membrane-
associated protein oligomerization using protein correlation profiling. Molecular & Cellular
Proteomics:mcp.000276.002017. 10.1074/mcp.RA117.000276.
McBride, Z., Chen, D., Lee, Y., Aryal, U.K., Xie, J., and Szymanski, D.B. (2019). A label-free mass
spectrometry method to predict endogenous protein complex composition. Molecular & Cellular
Proteomics **18**:1588. 10.1074/mcp.RA119.001400.
Meinert, M.C., and Delmer, D.P. (1977). Changes in biochemical composition of the cell wall of the
cotton fiber during development. Plant Physiol **59**:1088-1097.
Mujahid, H., Pendarvis, K., Reddy, J.S., Nallamilli, B.R., Reddy, K.R., Nanduri, B., and Peng, Z.
(2016). Comparative Proteomic Analysis of Cotton Fiber Development and Protein Extraction Method
Comparison in Late Stage Fibers. Proteomes **4**10.3390/proteomes4010007.
Nie, L., Wu, G., and Zhang, W. (2006). Correlation of mRNA expression and protein abundance
affected by multiple sequence features related to translational efficiency in Desulfovibrio vulgaris: a
quantitative analysis. Genetics **174**:2229-2243. 10.1534/genetics.106.065862.
Okekeogbu, I.O., Pattathil, S., González Fernández-Niño, S.M., Aryal, U.K., Penning, B.W., Lao, J.,
Heazlewood, J.L., Hahn, M.G., McCann, M.C., and Carpita, N. (2019). Glycome and Proteome
Components of Golgi Membranes Are Common between Two Angiosperms with Distinct Cell-Wall
Structures. The Plant Cell **31**:1094-1112. 10.1105/tpc.18.00755.

Perez-Riverol, Y., Bai, J., Bandla, C., García-Seisdedos, D., Hewapathirana, S., Kamatchinathan, S.,
Kundu, Deepti J., Prakash, A., Frericks-Zipper, A., Eisenacher, M., et al. (2021). The PRIDE
database resources in 2022: a hub for mass spectrometry-based proteomics evidences. Nucleic Acids
Research **50**:D543-D552. 10.1093/nar/gkab1038.
Pettolino, F.A., Yulia, D., Bacic, A., and Llewellyn, D.J. (2022). Polysaccharide composition during
cotton seed fibre development: temporal differences between species and in different seasons. Journal of
Cotton Research **5**ARTN 27
10.1186/s42397-022-00136-5.
Qin, Y., Wei, H.L., Sun, H.R., Hao, P.B., Wang, H.T., Su, J.J., and Yu, S.X. (2017). Proteomic
Analysis of Differences in Fiber Development between Wild and Cultivated Gossypium hirsutum L.
Journal of Proteome Research **16**:2811-2824. 10.1021/acs.jproteome.7b00122.
Qin, Y., Sun, M., Li, W., Xu, M., Shao, L., Liu, Y., Zhao, G., Liu, Z., Xu, Z., You, J., et al. (2022).
Single-cell RNA-seq reveals fate determination control of an individual fibre cell initiation in cotton
(Gossypium hirsutum). Plant Biotechnology Journal **20**:2372-2388. https://doi.org/10.1111/pbi.13918.
Qin, Y.M., Hu, C.Y., Pang, Y., Kastaniotis, A.J., Hiltunen, J.K., and Zhu, Y.X. (2007). Saturated very-
long-chain fatty acids promote cotton fiber and Arabidopsis cell elongation by activating ethylene
biosynthesis. Plant Cell **19**:3692-3704. 10.1105/tpc.107.054437.
R Core Team (2018). R: A language and environment for statistical computing. Version 3.5.1.
http://www.R-project.org/.
Rajangam, A.S., Kumar, M., Aspeborg, H., Guerriero, G., Arvestad, L., Pansri, P., Brown, C.J.,
Hober, S., Blomqvist, K., Divne, C., et al. (2008). MAP20, a microtubule-associated protein in the
secondary cell walls of hybrid aspen, is a target of the cellulose synthesis inhibitor 2,6-
dichlorobenzonitrile. Plant Physiol. **148**:1283-1294. pp.108.121913 [pii]
10.1104/pp.108.121913 [doi].
RStudio Team (2018). RStudio: Integrated Development Environment for R. Version 1.1.463.
https://rstudio.com/.
Ruan, Y., Llewellyn, D., and Furbank, R. (2001). The control of single-celled cotton fiber elongation by
developmentally reversible gating of plasmodesmata and coordinated expression of sucrose and k(+)
transporters and expansin. Plant Cell **13**:47-60.
Ryser, U. (1977). Cell-Wall Growth in Elongating Cotton Fibers - Autoradiographic Study. Cytobiologie
15:78-84.
Schubert, A.M., Benedict, C.R., Berlin, J.D., and Kohel, R.J. (1973). Cotton Fiber Development -
Kinetics of Cell Elongation and Secondary Wall Thickening. Crop Science **13**:704-709. DOI
10.2135/cropsci1973.0011183X001300060035x.
Seagull, R.W. (1992). A Quantitative Electron-Microscopic Study of Changes in Microtubule Arrays and
Wall Microfibril Orientation during Invitro Cotton Fiber Development. Journal of Cell Science **101**:561-
577.
Seagull, R.W. (1993). Cytoskeletal Involvement in Cotton Fiber Growth and Development. Micron
24:643-660. Doi 10.1016/0968-4328(93)90042-Y.
Singh, B., Avci, U., Inwood, S.E., Grimson, M.J., Landgraf, J., Mohnen, D., Sorensen, I., Wilkerson,
C.G., Willats, W.G., and Haigler, C.H. (2009). A Specialized Outer Layer of the Primary Cell Wall Joins
Elongating Cotton Fibers Into Tissue-like Bundles. Plant Physiol.
Stewart, J.M. (1975). Fiber initiation on the cotton ovule. Am J Bot **62**:723-730.
Sugano, S.S., Shimada, T., Imai, Y., Okawa, K., Tamai, A., Mori, M., and Hara-Nishimura, I. (2010).
Stomagen positively regulates stomatal density in Arabidopsis. Nature **463**:241-244.
10.1038/nature08682.
Sun, X., Qin, A., Wang, X., Ge, X., Liu, Z., Guo, C., Yu, X., Zhang, X., Lu, Y., Yang, J., et al. (2025).
Spatiotemporal transcriptome and metabolome landscapes of cotton fiber during initiation and early
development. Nature Communications **16**:858. 10.1038/s41467-025-55869-z.
Swaminathan, S., Grover, C.E., Mugisha, A.S., Sichterman, L.E., Lee, Y., Yang, P., Mallery, E.L.,
Jareczek, J.J., Leach, A.G., Xie, J., et al. (2024). Daily glycome and transcriptome profiling reveals

polysaccharide structures and correlated glycosyltransferases critical for cotton fiber growth. The Plant
Journal **120**:1857-1879. https://doi.org/10.1111/tpj.17084.
Tiwari, S.C., and Wilkins, T.A. (1995). Cotton (*Gossypium hirsutum*) seed trichomes expand via diffuse
growing mechanism. Can. J. Bot. **73**:746-757.
Tuttle, J.R., Nah, G., Duke, M.V., Alexander, D.C., Guan, X.Y., Song, Q.X., Chen, Z.J., Scheffler,
B.E., and Haigler, C.H. (2015). Metabolomic and transcriptomic insights into how cotton fiber
transitions to secondary wall synthesis, represses lignification, and prolongs elongation. Bmc Genomics
16ARTN 477
10.1186/s12864-015-1708-9.
Tyanova, S., Temu, T., and Cox, J. (2016). The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. Nature Protocols **11**:2301-2319. 10.1038/nprot.2016.136.
Wang, L., Liu, C., Liu, Y., and Luo, M. (2020a). Fumonisin B1-Induced Changes in Cotton Fiber
Elongation Revealed by Sphingolipidomics and Proteomics. Biomolecules **10**10.3390/biom10091258.
Wang, S., Li, W., Hu, L., Cheng, J., Yang, H., and Liu, Y. (2020b). NAguideR: performing and
prioritizing missing value imputations for consistent bottom-up proteomic analyses. Nucleic Acids
Research **48**:e83-e83. 10.1093/nar/gkaa498.
Wehrens, R., and Kruisselbrink, J. (2018). Flexible Self-Organizing Maps in kohonen 3.0. Journal of
Statistical Software **87**:1 - 18. 10.18637/jss.v087.i07.
Wen, X., Zhai, Y., Zhang, L., Chen, Y., Zhu, Z., Chen, G., Wang, K., and Zhu, Y. (2022). Molecular
studies of cellulose synthase supercomplex from cotton fiber reveal its unique biochemical properties. Sci
China Life Sci **65**:1776-1793. 10.1007/s11427-022-2083-9.
Wilkins, O., Brautigam, K., and Campbell, M.M. (2010). Time of day shapes Arabidopsis drought
transcriptomes. Plant J **63**:715-727. 10.1111/j.1365-313X.2010.04274.x.
Wilson, M.C., Howell, A.H., Sood, A., Lee, Y., Yang, P., Yu, E., Rani, H., Mallery, E.L.,
Swaminathan, S., Grover, C.E., et al. (2024). Developmental variability in cotton fiber cell wall
properties linked to important agronomic traits. bioRxiv:2024.2008.2019.607249.
10.1101/2024.08.19.607249.
Yanagisawa, M., Keynia, S., Belteton, S., Turner, J.A., and Szymanski, D.B. (2022). A conserved
cellular mechanism for cotton fibre diameter and length control. . *in silico* Plants **4**10.1093.
Yanagisawa, M., Desyatova, A.S., Belteton, S., Mallery, E.M., Turner, J.A., and Szymanski, D.B.
(2015). Patterning mechanisms of cytoskeletal and cell wall systems during leaf trichome morphogenesis.
Nature Plant **1**:15014. doi:10.1038/nplants.2015.14.
Yang, Y., Lai, W., Long, L., Gao, W., Xu, F., Li, P., Zhou, S., Ding, Y., and Hu, H. (2023a).
Comparative proteomic analysis identified proteins and the phenylpropanoid biosynthesis pathway
involved in the response to ABA treatment in cotton fiber development. Sci Rep **13**:1488.
10.1038/s41598-023-28084-3.
Yang, Z., Wang, J., Huang, Y., Wang, S., Wei, L., Liu, D., Weng, Y., Xiang, J., Zhu, Q., Yang, Z., et
al. (2023b). CottonMD: a multi-omics database for cotton biological study. Nucleic Acids Res **51**:D1446-
D1456. 10.1093/nar/gkac863.
Yu, J., Jung, S., Cheng, C.H., Lee, T., Zheng, P., Buble, K., Crabb, J., Humann, J., Hough, H.,
Jones, D., et al. (2021). CottonGen: The Community Database for Cotton Genomics, Genetics, and
Breeding Research. Plants (Basel) **10**10.3390/plants10122805.
Yu, Y., Wu, S., Nowak, J., Wang, G., Han, L., Feng, Z., Mendrinna, A., Ma, Y., Wang, H., Zhang, X.,
et al. (2019). Live-cell imaging of the cytoskeleton in elongating cotton fibres. Nat Plants **5**:498-504.
10.1038/s41477-019-0418-8.
Zang, Y., Hu, Y., Xu, C., Wu, S., Wang, Y., Ning, Z., Han, Z., Si, Z., Shen, W., Zhang, Y., et al.
(2021). GhUBX controlling helical growth results in production of stronger cotton fiber. iScience
24:102930. 10.1016/j.isci.2021.102930.
Zhang, Y., Yu, J., Wang, X., Durachko, D.M., Zhang, S., and Cosgrove, D.J. (2021). Molecular
insights into the complex mechanics of plant epidermal cell walls. Science **372**:706-711.
10.1126/science.abf2824.

Zhou, X., Hu, W., Li, B., Yang, Y., Zhang, Y., Thow, K., Fan, L., and Qu, Y. (2019). Proteomic
profiling of cotton fiber developmental transition from cell elongation to secondary wall deposition. Acta
Biochim Biophys Sin (Shanghai) **51**:1168-1177. 10.1093/abbs/gmz111.
1078

A

APO₁ (1846) V26 → V30

P200 (3299) V36 → V40

S200 (763) V16 → V20

SOM Group
Member (Protein)
Group profile
5 — 24
DPA

B

Cell fraction protein expression group

APO₁ (939)   P200 (1780)   S200 (254)

Increase (159)

V23 (6) V28 (4)
V32 (7) V33 (12)
V36 (60) V38 (27)
V1 (25) V2 (10) V6 (8)

Decrease (683)

V2 (8) V3 (84)
V4 (5) V5 (476) V8 (12)
V1 (31) V6 (36) V11 (25)
V18 (6)

Constitutive increase (396)

V22 (37) V27 (54)
V23 (96) V25 (135)
V28 (4) V29 (3)
V7 (2) V11 (65)

Constitutive decrease (183)

V21 (137)
V12 (1)
V17 (45)

Transient increase (71)

V25 (34)
V12 (37)

Uniform (1481)

V26 (82) V4 (96) V8 (19) V9 (130) V10 (156) V13 (138) V14 (120)
V15 (145) V18 (147) V19 (120) V20 (140) V24 (132) V16 (56)

Group profiles

Normalized abundance
Group ID
Group profile
No. of members
V2 (8)
5 10 15 20
DPA