Elevated lysosomal mass and enzyme activity in fibroblasts of the Mediterranean mouse *Mus spretus*

Melissa Sui[1]*, Joanne Teh[1]*, Kayleigh Fort[1], Daniel E. Shaw[2], Peter Sudmant[3], Tsuyoshi Koide[4], Jeffrey M. Good[2], Juan M. Vazquez[#3], Rachel B. Brem[#1]

Departments of [1]Plant and Microbial Biology and [3]Integrative Biology, University of California, Berkeley, Berkeley, CA 94720, USA

[2]Division of Biological Sciences, University of Montana, Missoula, MT 59812, USA

[4]National Institute of Genetics, Mishima, Shizuoka 411-8540, Japan

*Authors contributed equally to this work.

[#]To whom correspondence should be addressed: juan@vazquez.bio, rbrem@berkeley.edu.

Abstract

Failures of the lysosome-autophagy system are a hallmark of aging and many disease states. As a consequence, interventions that enhance lysosome function are of keen interest in the context of drug development. Throughout the biomedical literature, evolutionary biologists have found cases in which challenges faced by humans in clinical settings have been resolved by non-model organisms adapting to wild environments. Here, we used a primary cell culture approach to survey lysosomal characteristics in species of the genus *Mus*. We found that fibroblasts from *M. spretus*, a wild Mediterranean mouse, exhibited elevated lysosomal mass and enzyme activity along with reduced activity of β-galactosidase, a classical marker of cellular senescence, compared to those from *M. musculus*, a related species adapted to human-associated environments. We propose that classic laboratory models of lysosome function and senescence may reflect characters that diverge from the phenotypes of wild mice. The *M. spretus* phenotype may ultimately serve as a blueprint for interventions that ameliorate lysosomal dysfunction under conditions of stress and disease.

Introduction

Many aspects of metazoan health hinge on the ability of the lysosome-autophagy pathway to recycle damaged macromolecules and to direct cell growth decisions (Shin & Zoncu 2020). Indeed, interventions that act broadly to promote health and longevity often require the lysosome-autophagy system (Aman *et al*. 2021; Hansen *et al*. 2018; Bareja *et al*. 2019). More specific mechanisms to boost autophagy are also of potential clinical interest, particularly for treatment of proteinopathies and aging etiologies (Bonam *et al*. 2019; Hansen *et al*. 2018). In practice, whether and how to stimulate proteostasis machinery to advance organismal health remains an open question, and the literature describing such manipulations is in its infancy (Pyo *et al*. 2013; Leiva-Rodríguez *et al*. 2018; Shin *et al*. 2013; Liu *et al*. 2023; Bhuiyan *et al*. 2013; Sami et al. 2002; Bonam et al. 2019; Grabowski & Mistry 2022).

Against a backdrop of decades of work in laboratory systems, ecologists have cataloged stress- and disease-resistance traits in wild genotypes from unusual niches, perhaps most famously in long-lived animal species (Oka *et al*. 2023; Chusyd *et al*. 2021; Finch 2009). Cell-based surveys represent a powerful approach to find and dissect these natural resilience programs. One particularly fruitful discipline has profiled the variation in chemical stress resistance across primary cells from panels of non-model animals, including, in landmark cases, discovery of the underlying mechanisms (Tian *et al*. 2019; Harper *et al*. 2007; Attaallah *et al*. 2020; Sulak *et al.* 2016). Cellular senescence has also been shown to vary across animal species in *in vitro* models (Attaallah *et al*. 2020; Kang *et al*. 2023; Zhao *et al*. 2018; Gomes *et al.* 2011).

In this study, we aimed to leverage species-specific variation in lysosomal markers within the *Mus* genus, to investigate how evolutionary processes have shaped the function of this organelle. Mouse species in this genus shared a common ancestor 7-8 million years ago, and as they radiated across Eurasia, *M. musculus* subspecies came to occupy human-associated

niches, whereas other taxa are still found exclusively in the wild (Pagès *et al.* 2015; Smissen &
Rowe 2018). Our previous case study (Kang *et al.* 2023) found differences in senescence
behaviors, including lysosome markers, across fibroblasts from *M. musculus* subspecies and *M.*
spretus, a wild relative that diverged 1-3 million years ago (Morgan *et al.* 2022; Dejager *et al.*
2009). Here, we aimed to build on these observations to gain a deeper understanding of
lysosomal programs in non-model mice, focusing on the contrasts between the genotypes of
human commensal and laboratory mice and those of their wild relatives.
63
64   ## Results
65
66   ### *M. spretus* fibroblasts exhibit reduced lysosomal β-galactosidase activity
67
We previously reported divergence among fibroblasts from several *Mus* genotypes (Kang *et al.*
2023) in characteristics of cellular senescence, a stress-induced program of cell cycle arrest,
resistance to apoptosis, and activation of immune signaling (Campisi & d'Adda di Fagagna
2007; Campisi 2005; Hornsby 2002). Most salient from our initial findings were differences
between *Mus* fibroblasts in activity of lysosomal β-galactosidase, which is used as a marker of
senescence by virtue of its accumulation to detectable levels at suboptimal pH, due to the
expanded lysosomal content of senescent cells (Robbins *et al.* 1970; Magalhães & Passos
2018; Dodig *et al.* 2019; Curnock *et al.* 2023).To explore this variation further, we established a
panel of primary tail skin fibroblasts from four wild-derived strains of *M. m. musculus* (PWK/PhJ,
BLG2/Ms, CHD/Ms, MSM/Ms); two wild-derived strains of *M. m. domesticus* (ManB/NachJ,
TUCA/NachJ); an admixed laboratory strain of mostly *M. m. domesticus* origin (C57BL/6J); a
wild-derived strain of the steppe mouse *M. spicilegus* (ZRU); and two wild-derived strains of the
Mediterranean mouse *M. spretus* (STF/Pas, SFM/Pas). We subjected each culture to 15 Gy of
ionizing radiation and incubated for 10 days, which is sufficient to induce DNA damage
response, arrested cell growth, and senescence expression programs in fibroblasts from across
the genus (Kang *et al.* 2023, Hernandez-Segura *et al.* 2018). In assays of β-galactosidase,
fibroblasts of each genotype displayed the anticipated increase in activity following irradiation,
compared to their respective control counterparts (Figure 1A-B and Figure S1). Whereas cells
from strains of each species behaved similarly, we observed significant differences between
species: irradiated *M. spretus* fibroblasts exhibited β-galactosidase signal ~2-fold lower than did
cells from *M. m. musculus* and *M. m. domesticus,* and staining of *M. spicilegus* cells was
between the two extremes (Figure 1A-B and Figure S1). To follow up on this difference, we
used an alternative approach for senescence induction via the radiomimetic drug
neocarzinostatin, which induces senescence in mouse cells (Ito *et al.* 2018; Correia□Melo *et al.*
2016, Hernandez-Segura *et al.* 2018). Fibroblasts treated with neocarzinostatin exhibited a
pattern mimicking that we had seen under radiation, with the treatment inducing elevated β-
galactosidase activity in all cultures, and cells from *M. spretus* staining lower than those of sister
taxa (Figure 1C and Figure S1). Furthermore, the same trend was detectable in cells cultured in
the absence of DNA damage: fibroblasts from *M. musculus* subspecies exhibited higher β-
galactosidase staining than those of *M. spicilegus*, which stained higher than *M. spretus*. The
latter species differences from untreated cells were of smaller magnitude than those manifesting
after stress, and this dependence on treatment was statistically significant (Figures 1, S1, and

S2). Thus, *M. spretus* fibroblasts exhibited dampened β-galactosidase activity relative to other
genotypes, with amplification of the effect under stress. Controls ruled out culture passage
number as a driver of the variation (Figure S3).
103
In principle, differences in the senescence-apoptosis fate choice after DNA damage (Childs *et*
al. 2014; Zhao *et al.* 2018; Attaallah *et al.* 2020) could contribute to the divergence between
mouse genotypes that we had seen in fibroblast β-galactosidase staining. To explore this, we
measured caspase 3/5 activity in fibroblasts in response to irradiation and neocarzinostatin
treatment (Figure S4). The results showed no consistent patterns of apoptosis either within or
between species' genotypes, arguing against a role for apoptosis activation in species
differences in fibroblast β-galactosidase activity.
111
To establish further the robustness of *Mus* species differences in fibroblast β-galactosidase
activity, we considered the dose-response relationship with stress. We compared primary
fibroblasts from *M. m. musculus* PWK and *M. spretus* STF as representatives of their respective
species, in each case assaying β-galactosidase upon irradiation at increasing doses. The
results revealed a consistent ~2-fold difference between the genotypes at each dose (Figure
S5), a magnitude slightly exceeding the species divergence in the untreated control, consistent
with our survey across genotypes at fixed stress doses (Figure 1). These data ruled out a switch
by *M. spretus* fibroblasts into a high-amplitude, *M. musculus*-like program above a certain
threshold of stress exposure. We conclude instead that *M. spretus* cells are hard-wired for lower
β-galactosidase activity in all tested conditions—including the resting state.
122
To advance our search to understand *Mus* species differences in β-galactosidase activity in the
fibroblast model, we next focused on regulation of the enzyme itself. Transcriptional profiling
data from fibroblasts revealed 1.5 to 2-fold higher expression of the lysosomal β-galactosidase
GLB1 in *M. musculus* fibroblasts than in those of *M. spretus*, regardless of treatment (Figure
S6A). Allele-specific expression measurements in stressed and control fibroblasts from an
interspecies F1 hybrid made clear that this change was regulated in *cis*: that is, the *M. musculus*
allele of *GLB1* drove higher expression of its own encoding locus than the *M. spretus* allele,
when both were in the same nucleus (Figure S6B). Thus, the species changes in β-
galactosidase activity we had noted in terms of cell biology in fibroblasts (Figure 1) were
mirrored by regulation of gene expression, including robust differences between genotypes in
unstressed control conditions.
134
135 **No consistent divergence in senescence signaling across *Mus* species fibroblasts**
136
We reasoned that the differences in β-galactosidase activity in fibroblasts among the *Mus*
species we tested were likely mediated by mechanisms unrelated to senescence itself. As a first
investigation of this idea, we explored senescence signaling, using as a readout p21, a key
regulator of cell cycle arrest in senescence (Gu *et al.* 2013; Yew *et al.* 2011; Kreis *et al.* 2016).
We quantified levels of p21 protein in response to either irradiation or neocarzinostatin, with
fibroblasts from *M. m. musculus* PWK and *M. spretus* STF as a testbed. Results showed the
expected induction of p21 in fibroblasts from both species in response to irradiation or

neocarzinostatin (Figure S7). In investigating quantitative patterns among these p21 induction
behaviors, we noted some species-specific differences, though they were not of consistent
direction: *M. spretus* cells induced p21 protein abundance more than did *M. musculus* under
neocarzinostatin treatment but less following irradiation and in untreated controls (Figure S7).
These changes could reflect the differential activity of alternate senescence signaling pathways
(Saul *et al.* 2023) between species in some treatments. However, given our focus on lysosomal
β-galactosidase differences between *M. musculus* and *M. spretus* across all tested conditions
(Figure 1), we concluded that regulation of p21 abundance was not a consistent correlate of this
phenotype (although our results do not rule out p21 localization differences between the
genotypes). Indeed, in analysis of mRNA expression profiles (Kang *et al.* 2023), we detected no
significant difference in induction of the p21 gene (*Cdkn1A*) between *M. musculus* and *M.*
spretus fibroblasts in response to stress (Figure S8).
156
157 **Increased lysosomal enzyme activity and lysosomal mass in *M. spretus* fibroblasts**
158
We hypothesized that the higher β-galactosidase expression and activity we had seen in *M.*
musculus fibroblasts could be a consequence of heightened need owing to failures elsewhere in
the proteostasis system, relative to cells of *M. spretus*. To explore this, we first assayed acidic
compartments with the LysoTracker stain, again making use of *M. m. musculus* PWK and *M.*
spretus STF as a test system. Conforming to our prediction, quantitative microscopy revealed a
~2.5-fold increase in the cellular area stained by LysoTracker in untreated *M. spretus* fibroblasts
compared with *M. musculus* (Figure 2A), and the species difference persisted in cultures
induced to senesce with irradiation and neocarzinostatin (Figure 2A). A separate assay design
using flow cytometry to quantify LysoTracker staining yielded similar results, with ~3-fold higher
levels in *M. spretus* cells in all conditions (Figure S9). LysoTracker staining was absent from
both *M. musculus* and *M. spretus* fibroblasts following treatment with the lysosomal H+-ATPase
poison bafilomycin A1, confirming the dependence of this readout on lysosomal acidification
(Figure S10).
172
To investigate further the potential for enhanced lysosomal enzyme function in *M. spretus* cells
compared to *M. musculus*, we developed an assay targeting the lysosomal enzyme cathepsin
D. This assay employed a BODIPY-labeled pepstatin A conjugate, a fluorescent probe with
established use in studies of lysosomal enzyme activity (Boland *et al.* 2008; Lee *et al.* 2010).
Measurements from experiments of this design made clear that *M. spretus* fibroblasts exhibited
~2-fold higher area staining for cathepsin D activity than did *M. musculus* cells under untreated
conditions, and similar though slightly dampened differences between the genotypes after
irradiation or neocarzinostatin treatment (Figure 2B). Taken together, our results define a trait
syndrome in *M. spretus* fibroblasts characterized by elevated lysosomal mass with acidic pH
and enzymatic activity, relative to other *Mus* species, under both basal and senescent
conditions.
184
185 **Lysosomal gene expression divergence between *Mus* fibroblasts**
186

To unearth clues to mechanisms of the divergence between *Mus* species fibroblasts in
lysosomal characters, we revisited the expression programs of these cells, this time with a
broad survey of lysosome-associated genes. The results revealed marked expression
differences between purebred *M. spretus* and *M. musculus* fibroblasts (Figure S11), including
high expression in *M. spretus* cells of lysosomal biogenesis factors (namely *Dtnbp1* and
members of the *Bloc1s* family); the lysosomal regulator *Lamtor1*; and subunits of the lysosomal
H+-ATPase domain *ATP6v1*. Measurements of allele-specific expression in interspecies F1
hybrid fibroblasts established major contributions of *cis*-regulatory variation at many such genes
(Figure S11), meaning that the underlying genetic basis of a given gene's expression change
between purebreds could often be ascribed to heritable variants at the locus itself. We conclude
that lysosomal gene regulatory changes between the fibroblasts of *Mus* species, controlled in
part by *cis*-regulatory variants, parallel the divergence in lysosomal mass and enzyme activity—
as expected if *M. spretus* cells made more lysosomes, and populated them to a greater extent
with more functional enzymes, as a direct consequence of their distinct regulatory program.
201

Discussion

For decades, biomedical researchers have relied on strains from the *M. musculus* clade as the
standard for studying lysosomal and senescence biology. In this work, we have demonstrated
that lysosomal phenotypes vary quantitatively across *Mus* in fibroblast culture models, and that
in contrast to *M. musculus*, cells from the non-commensal mouse *M. spretus* exhibit a program
of lysosomal function of the kind that, in the experimental literature, has been associated with
healthy aging and disease resistance.

When experimentally induced in laboratory cell models, a backup in the lysosomal-autophagy
system triggers compensatory increases in lysosomal mass and number, for which high β-
galactosidase activity is a robust marker (Curnock *et al.* 2023; Delfarah *et al.* 2021; Lee *et al.*
2006, Dimri *et al.* 1995, Rovira *et al.* 2022). We propose a similar causal relationship between
the phenotypes we have observed as they vary across *Mus* fibroblasts. Under this model, the
elevated β-galactosidase activity we have seen in *M. musculus* cells would represent
compensation for their limited lysosomal mass and enzyme activity in comparison to *M. spretus*
genotypes, even in the absence of stress. The *M. musculus cis*-regulatory allele upregulating β-
galactosidase that we have noted in fibroblasts could well represent a genetically encoded
component of such a program, a constitutive boost in protein degradation capacity by a
regulatory mechanism, in the face of lysosomal defects. Indeed, we consider the broader set of
lysosomal genes at which we have detected regulatory variation between the species as
compelling candidate components of the mechanism underlying increased lysosomal mass and
function in *M. spretus* cells.

Our findings do leave open whether the compromised levels of these phenotypes in *M.
musculus* cells represent an ancestral state in *Mus* that was resolved in the *M. spicilegus*–*M.
spretus* lineage, or a derived trait that emerged alongside commensalism in *M. musculus*. In
either case, the evidence that we have seen for conservation across *M. musculus* strains and
subspecies strongly suggests a history of constraint in this lineage. If the lysosomal behaviors

we have studied here prove to have evolved under selection, they could conceivably relate to
body size, response to immune challenges, or association with human ecology as they differ
between *M. musculus* and *M. spretus* (Dejager *et al.* 2009; Mahieu *et al.* 2006; Kawakami &
Yamamura 2008; Staelens *et al.* 2002; Blanchet *et al.* 2011; Pérez del Villar *et al.* 2013; Harr *et*
al. 2016).

Ultimately, as its mechanism is revealed, the elevated mass and enzyme activity of lysosomes
that we have seen in *M. spretus* fibroblasts are of interest in the search for mimetics that would
boost lysosomal function in a human clinical setting. More broadly, our results provide a way
forward for the use of wild mouse species as models for lysosomal function and senescence,
including future comparisons to the lysosomes of human cells and their phenotypes in disease.

Methods

Primary tail fibroblasts were extracted as described by Khan and Gasser (2016). Subsequent
culture and experiments used complete medium (DMEM, 10% FBS, 1% pen-strep) in T25 flasks
at 37˚C, 3% O_2, and 10% CO_2. Cells were passaged based on confluence using trypsin.
Experimental treatments included 15 GY of ionizing irradiation or 3.6 µM neocarzinostatin.
Analysis kits used were the Abcam Ltd. Senescence Detection Kit (Cat. #ab65351), ApoTox-
Glo™ Triplex Assay (Promega Cat. #G6321), LysoTracker™ Green DND-26 (Thermo Fisher
Cat. #L7526), and Pepstatin A BODIPY™ FL Conjugate (Thermo Fisher Cat. #P12271).
Primary and secondary antibodies used in western blots were the rabbit monoclonal Anti-p21
antibody (ab188224, Abcam, 1:1000), mouse monoclonal anti-β-tubulin (T9026, Sigma-Aldrich,
1:1000), Goat Anti-Mouse IgG(H+L) Human ads-HRP (Cat#1031-05, Southern Biotech,
1:5000), and Goat Anti-Rabbit IgG(H+L) Human ads-HRP (Cat#4050-05, Southern Biotech,
1:5000). Additional details of methods available in Supplementary Materials.

Acknowledgements

The authors thank Vera Gorbunova, Emilie Tu, Helen Bateup, Samantha Jackson, Linda
Wilbrecht, Polly Campbell, and Michael Nachman for animal material; Diana Bautista, Britt
Glaunsinger, José Pablo Vázquez-Medina, and Mary West for their generosity with space and
resources; Juliana Valencia Lesmes, Mara Baylis, Sam Rider, Harriet Song and William
D'Angelo for technical contributions; and members of the Brem lab for helpful discussions. This
work was supported by National Institutes of Health R01NS116992 to RBB and R01GM120430
to RBB and JMG. JMV was supported by the National Science Foundation Postdoctoral
Research Fellowship in Biology (#2109915) and by the National Institute for Health
T32AG000266 and 1K99AG088361.

Data Accessibility Statement

272 Data used in this manuscript can be found in an external data repository on Figshare at
273 https://doi.org/10.6084/m9.figshare.c.8086870.v1. RNA-sequencing data used in this

274   manuscript can be found in the NCBI Gene Expression Omnibus (GEO;
275   https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE201217.
276
277   Figure captions
278
Figure 1. Low β-galactosidase activity in _M. spretus_ fibroblasts relative to _M. musculus_.
In a given panel, in the plot at the top, each bar length displays the mean percentage of primary
fibroblasts of the indicated strain and species that stained positive after administration of the
colorimetric β-galactosidase substrate X-Gal. From top to bottom, complete names of the strains
are as follows: _M. m. musculus_ (PWK/PhJ, BLG2/Ms, CHD/Ms, MSM/Ms), laboratory strain of
mostly _M. m. domesticus_ origin (C57BL/6J), _M. m. domesticus_ (TUCA/NachJ, ManB/NachJ), _M._
spicilegus, and _M. spretus_ (SFM, STF/Pas). Representative images of primary fibroblasts from
M. musculus PWK and _M. spretus_ STF are shown at the bottom, with black arrows indicating
dead cells. Panels report results from (A) untreated cells, (B) cells treated with ionizing
irradiation followed by a 10-day incubation, or (C) cells after 1 hour of neocarzinostatin
treatment followed by 24 hours of incubation. Points report biological and technical replicates
collected over at least two separate days. Error bars report one standard error above and below
the mean. ***, two-tailed Wilcoxon $p < 10^{-7}$ comparing _M. spretus_ with all other genotypes; _M._
spretus and _M. spicilegus_ also differed in each panel (Wilcoxon $p < 0.05$). For (B) and (C), in a
comparison between the respective treatment and the untreated control in (A), a two-factor
ANOVA with treatment and genotype as factors yielded $p < 2e^{-16}$ for the interaction between the
two factors. Replicate numbers are provided in Figure S12.
296
Figure 2. Elevated lysosomal mass and enzyme activity in _M. spretus_ fibroblasts relative
to _M. musculus_. (A) Shown are results of quantitative microscopy analyses of staining by the
lysosomal acidity reporter LysoTracker in primary fibroblasts of the indicated genotype (_M._
spretus, strain STF; _M. m. musculus_, strain PWK). The _y_-axis reports the number of fluorescent
pixels per cell, normalized to the control _M. musculus_ sample for each experiment. Pairs of bars
report results from untreated cells (left), or cells treated with irradiation followed by a 10-day
incubation (middle) or after 1 hour of neocarzinostatin (NCS) treatment followed by 24 hours of
incubation (right). At the bottom are shown representative fluorescence microscopy images of
LysoTracker staining in primary fibroblasts from _M. musculus_ PWK and _M. spretus_ STF in
untreated control, neocarzinostatin (NCS) treatment, and irradiation treatment. (B) Shown are
results from assays of cathepsin D activity via BODIPY-pepstatin A fluorescence staining in
fibroblasts of the indicated genotype (_M. spretus,_ strain STF; _M. m. musculus_, strain PWK). In
the plot at the top, the _y_-axis reports the number of fluorescent pixels per cell, normalized to the
control _M. musculus_ sample for each experiment. Pairs of bars report results from untreated
cells (left), or cells treated with irradiation followed by a 10-day incubation (middle) or after 1
hour of neocarzinostatin (NCS) treatment followed by 24 hours of incubation (right). At the
bottom are shown representative fluorescence microscopy images of BODIPY-pepstatin A
staining in primary fibroblasts from _M. musculus_ PWK and _M. spretus_ STF in untreated control,
neocarzinostatin (NCS) treatment, and irradiation treatment.. Data points correspond to
biological and technical replicates collected over at least two different days, and the bar height

reports their mean. Error bars report one standard error above and below the mean. *, Wilcoxon
$p < 0.05$, **, Wilcoxon $p < 0.01$, ***, Wilcoxon $p < 0.001$. Replicate numbers are provided in
Figure S12.
320

Supplementary figure captions

Figure S1. Impact of genotype and treatment on β-galactosidase activity in *Mus*
fibroblasts. Data are those of Figure 1 of the main text, grouped by genotype to emphasize
treatment effects. A given set of three bars represents results from primary fibroblasts of the
indicated strain from three conditions: untreated cells (darkest color), cells exposed to ionizing
irradiation followed by a 10-day incubation (medium color), and cells treated with
neocarzinostatin for 1 hour followed by 24 hours of incubation (lightest color).

Figure S2. Low normalized β-galactosidase activity in *M. spretus* fibroblasts. Data are as
in Figure 1 of the main text, except that measurements from the culture of a given genotype and
treatment were normalized to the average of all measurements from untreated controls of that
genotype. ***, two-tailed Wilcoxon $p < 0.001$ comparing *M. spretus* with all other genotypes for
both treatments.

Figure S3. Fibroblast passage number has no detectable effect on X-Gal staining. In a
given panel, each row reports results from assays of the β-galactosidase substrate X-Gal on
fibroblasts of the indicated genotype shown in Figure 1 of the main text, and each point reports
one replicate culture. The x-axis reports the passage number of the respective culture and the
y-axis reports X-Gal staining. (A) Cells induced to senesce with irradiation (right) and their
paired controls (left). (B) Cells induced to senesce with neocarzinostatin (right) and their paired
controls (left). Replicate numbers are provided in Figure S12.

Figure S4. No consistent variation between *Mus* species fibroblasts in apoptosis activity.
Data are as in Figure 1 of the main text, except that for a given genotype and treatment, the *x*-
axis reports caspase activity of the culture measured with the Apotox-Glo assay, normalized to
the average from untreated cultures of the respective genotype for each experiment. (A)
Outliers trimmed for visualization. (B) Full data set view. Replicate numbers are provided in
Figure S12.

Figure S5. Dose dependence of β-galactosidase staining in *Mus* fibroblasts. Each trace
shows results from assays of the β-galactosidase substrate X-Gal on primary fibroblasts of the
indicated genotype (*M. spretus*, strain STF; *M. m. musculus*, strain PWK). For a given trace, the
x-axis reports irradiation dosage (in centigray, cGy) at 0 (Control), 1000, 1500, 2000, and 2500
cGy, and the y-axis reports the percentage of X-Gal positive cells in untreated controls or in
irradiated cells after 10 days of incubation. Smaller points represent biological and technical
replicates, and larger points indicate their mean values. ***, two-factor ANOVA with dosage and
species as factors yielded $p < 2e^{-16}$ for each factor. Replicate numbers are provided in Figure
S12.

Figure S6. Variation between *Mus* fibroblasts in Glb1 mRNA expression. (A) Shown are
measurements of mRNA expression of the β-galactosidase gene from profiling of fibroblasts of
the indicated genotype (*M. spretus*, strain STF; *M. m. musculus*, strain PWK), untreated or
treated with irradiation followed by a 10-day incubation (Kang *et al.* 2023). Data points within
each column represent biological and technical replicates and the bar height reports their mean.
Only the effect of genotype was significant in a two-factor ANOVA with condition and species as
factors (*, $p < 0.05$). Wilcoxon test yielded no significant difference between the species for
each condition. (B) Data are as in (A), except that allele-specific mRNA expression of the β-
galactosidase gene was measured from profiling of fibroblasts of an F1 hybrid between *M.*
spretus, strain STF and *M. m. musculus*, strain PWK (Kang *et al.* 2023). In a two-factor ANOVA
with condition and genotype as factors, both had significant effects (species $p < 0.0005$, ***;
condition $p < 0.05$, *). Replicate numbers are provided in Figure S12.

Figure S7. No consistent variation between *Mus* fibroblasts in p21 abundance. Shown are
results of Western blot assays of abundance of the senescence regulator p21 in primary
fibroblasts of the indicated genotype (*M. spretus*, strain STF; *M. m. musculus*, strain PWK). (A)
Top, representative blot showing tubulin (55 kDa) and p21 (21 kDa) levels in 25 µg of whole cell
lysate (WCL) from untreated control cells or those irradiated (Xray) and incubated for 10 days to
establish senescence. Bottom, blot as at top except that cells were treated with 1 hour of
neocarzinostatin (NCS) and incubated for 24 hours to establish senescence. (B) Each column
reports quantification of p21 protein abundance normalized to tubulin for the indicated genotype
and treatment. Data points correspond to biological and technical replicates collected over at
least two different days, and the bar height reports their mean. Error bars report one standard
error above and below the mean. No comparisons were significant in pairwise Wilcoxon tests. *,
two-factor ANOVA with condition and species as factors yielded p < 0.05 for species. Replicate
numbers are provided in Figure S12.

Figure S8. No significant variation between *Mus* fibroblasts in Cdkn1a mRNA expression.
Shown are measurements of mRNA expression of the p21 gene Cdkn1a, from profiling of
fibroblasts of the indicated genotype (*M. spretus*, strain STF; *M. m. musculus*, strain PWK),
untreated or treated with irradiation followed by a 10-day incubation (Kang *et al.* 2023). Data
points within each column represent biological and technical replicates and the bar height
reports their mean. Error bars report one standard error above and below the mean. Only
condition had a significant impact on expression in a two-factor ANOVA with treatment and
genotype as factors (p < 0.005, **). Replicate numbers are provided in Figure S12.

**Figure S9. Fluorescent signal quantification of LysoTracker staining indicates enhanced
intracellular acidity in *M. spretus* fibroblasts.** Shown are results from assays of the
intracellular acidity reporter LysoTracker on primary fibroblasts of the indicated genotype (*M.*
spretus, strain STF; *M. m. musculus*, strain PWK). The *y*-axis reports mean fluorescence of the
indicated culture normalized to the control *M. musculus* sample for each experiment. Data
points correspond to biological and technical replicates collected over at least two different
days, and the bar height reports their mean. Error bars report one standard error above and

below the mean. *, Wilcoxon $p < 0.05$, **, Wilcoxon $p < 0.01$. Replicate numbers are provided in
Figure S12.

Figure S10. Bafilomycin A1 treatment abrogates LysoTracker fluorescent staining. (A)
Shown are representative microscopy images of intracellular acidity reporter LysoTracker in
untreated primary fibroblasts of the indicated genotype (*M. spretus,* strain STF; *M. m. musculus*,
strain PWK) after bafilomycin A1 treatment. (B) Alamar Blue cell viability stain results are shown
for primary fibroblasts of the indicated genotypes (*M. spretus*, strain STF; *M. m. musculus*,
strain PWK) either with or without bafilomycin A1 treatment, or a no-cell control (Blank). The y-
axis indicates absorbance of cell cultures at 570 nm, corresponding to Alamar Blue after
reduction by active cellular metabolism. Data points within each column represent technical
replicates and the bar height reports their mean.

Figure S11. Variation between species fibroblasts in lysosomal gene expression. Each
point reports the effect of genetic variation between *M. spretus* STF and *M. m. musculus* PWK
on the expression of a lysosomal gene in primary skin fibroblasts from (Kang *et al.* 2023). The *x*-
axis shows the ratio of expression of the *M. spretus* allele relative to that of the *M. musculus*
allele in an F1 hybrid background, indicating the contribution of *cis*-acting regulatory variants.
The *y*-axis displays gene expression in purebred *M. spretus* fibroblasts normalized to that of *M.*
musculus. Genes positioned near the diagonal line exhibit expression differences primarily
driven by *cis*-regulatory variation. Red-labeled genes are highlighted in the main text.

Figure S12. Sample sizes. Shown are sample sizes for each experiment presented in the main
and supplementary figures. (n) denotes the number of independent animals used per
experiment, (c) denotes the number of independent culture replicates used per experiment, and
(q) denotes the number of cells quantified in the experiment.

References

Aman Y, Schmauck-Medina T, Hansen M, Morimoto RI, Simon AK, Bjedov I, Palikaras K, Simonsen A, Johansen T, Tavernarakis N, Rubinsztein DC, Partridge L, Kroemer G, Labbadia J & Fang EF (2021) Autophagy in healthy aging and disease. *Nat Aging* 1, 634–650.

Attaallah A, Lenzi M, Marchionni S, Bincoletto G, Cocchi V, Croco E, Hrelia P, Hrelia S, Sell C & Lorenzini A (2020) A pro longevity role for cellular senescence. *Geroscience* 42, 867–879.

Bareja A, Lee DE & White JP (2019) Maximizing Longevity and Healthspan: Multiple Approaches All Converging on Autophagy. *Front Cell Dev Biol* 7, 183.

Bhuiyan MS, Pattison JS, Osinska H, James J, Gulick J, McLendon PM, Hill JA, Sadoshima J & Robbins J (2013) Enhanced autophagy ameliorates cardiac proteinopathy. *J Clin Invest* 123, 5284–5297.

Blanchet C, Jaubert J, Carniel E, Fayolle C, Milon G, Szatanik M, Panthier J-J & X Montagutelli (2011) Mus spretus SEG/Pas mice resist virulent Yersinia pestis, under multigenic control. *Genes Immun* 12, 23–30.

Boland B, Kumar A, Lee S, Platt FM, Wegiel J, Yu WH & Nixon RA (2008) Autophagy Induction and Autophagosome Clearance in Neurons: Relationship to Autophagic Pathology in Alzheimer's Disease. *J Neurosci* 28, 6926–6937.

Bonam SR, Wang F & Muller S (2019) Lysosomes as a therapeutic target. *Nat Rev Drug Discov* 18, 923–948.

Campisi J (2005) Senescent cells, tumor suppression, and organismal aging: good citizens, bad neighbors. *Cell* 120, 513–522.

Campisi J & d'Adda di Fagagna F (2007) Cellular senescence: when bad things happen to good cells. *Nat Rev Mol Cell Biol* 8, 729–740.

Childs BG, Baker DJ, Kirkland JL, Campisi J & van Deursen JM (2014) Senescence and apoptosis: dueling or complementary cell fates? *EMBO Rep* 15, 1139–1153.

Chusyd DE, Ackermans NL, Austad SN, Hof PR, Mielke MM, Sherwood CC & Allison DB (2021) Aging: What We Can Learn From Elephants. *Front Aging* 2, 726714.

Correia‑Melo C, Marques FD, Anderson R, Hewitt G, Hewitt R, Cole J, Carroll BM, Miwa S, Birch J, Merz A, Rushton MD, Charles M, Jurk D, Tait SW, Czapiewski R, Greaves L, Nelson G, Bohlooly-Y M, Rodriguez-Cuenca S, Vidal-Puig A, Mann D, Saretzki G, Quarato G, Green DR, Adams PD, Von Zglinicki T, Korolchuk VI & Passos JF (2016) Mitochondria are required for pro-ageing features of the senescent phenotype. *The EMBO Journal* 35, 724–742.

Curnock R, Yalci K, Palmfeldt J, Jäättelä M, Liu B & Carroll B (2023) TFEB‑dependent lysosome biogenesis is required for senescence. *EMBO J* 42, e111241.

De Magalhães JP & Passos JF (2018) Stress, cell senescence and organismal ageing. *Mechanisms of Ageing and Development* 170, 2–9.

Del Villar LP, Vicente B, Galindo-Villardón P, Castellanos A, Pérez-Losada J & Muro A (2013) Schistosoma mansoni experimental infection in Mus spretus (SPRET/EiJ strain) mice. *Parasite*, 20, 27.

Dejager L, Libert C & Montagutelli X (2009) Thirty years of Mus spretus: a promising future. *Trends Genet* 25, 234–241.

Delfarah A, Hartel NG, Zheng D, Yang J & Graham NA (2021) Identification of a Proteomic Signature of Senescence in Primary Human Mammary Epithelial Cells. *J Proteome Res* 20, 5169–5179.

Dimri GP, Lee X, Basile G, Acosta M, Scott G, Roskelley C, Medrano EE, Linskens M, Rubelj I & Pereira-Smith O (1995) A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proceedings of the National Academy of Sciences* 92, 9363–9367.

Dodig S, Čepelak I & Pavić I (2019) Hallmarks of senescence and aging. *Biochem Med (Zagreb)* 29, 030501.

Finch CE (2009) Update on slow aging and negligible senescence--a mini-review. *Gerontology* 55, 307–313.

Gomes NMV, Ryder OA, Houck ML, Charter SJ, Walker W, Forsyth NR, Austad SN, Venditti C, Pagel M, Shay JW & Wright WE (2011) Comparative biology of mammalian telomeres: hypotheses on ancestral states and the roles of telomeres in longevity determination. *Aging Cell* 10, 761–768.

Grabowski GA & Mistry PK (2022) Therapies for lysosomal storage diseases: Principles,
practice, and prospects for refinements based on evolving science. *Mol Genet Metab*
137, 81–91.

Green DR, Galluzzi L & Kroemer G (2011) Mitochondria and the autophagy-inflammation-
cell death axis in organismal aging. *Science* 333, 1109–1112.

Gu Z, Jiang J, Tan W, Xia Y, Cao H, Meng Y, Da Z, Liu H & Cheng C (2013) p53/p21
Pathway involved in mediating cellular senescence of bone marrow-derived
mesenchymal stem cells from systemic lupus erythematosus patients. *Clin Dev*
Immunol 2013, 134243.

Gupta P, Soyombo AA, Shelton JM, Wilkofsky IG, Wisniewski KE, Richardson JA &
Hofmann SL (2003) Disruption of PPT2 in mice causes an unusual lysosomal storage
disorder with neurovisceral features. *Proceedings of the National Academy of Sciences*
100, 12325–12330.

Hansen M, Rubinsztein DC & Walker DW (2018) Autophagy as a promoter of longevity:
insights from model organisms. *Nat Rev Mol Cell Biol* 19, 579–593.

Harper JM, Salmon AB, Leiser SF, Galecki AT & Miller RA (2007) Skin-derived fibroblasts
from long-lived species are resistant to some, but not all, lethal stresses and to the
mitochondrial inhibitor rotenone. *Aging Cell* 6, 1–13.

Harr B, Karakoc E, Neme R, Teschke M, Pfeifle C, Pezer Ž, Babiker H, Linnenbrink M,
Montero I, Scavetta R, Abai MR, Molins MP, Schlegel M, Ulrich RG, Altmüller J,
Franitza M, Büntge A, Künzel S & Tautz D (2016) Genomic resources for wild
populations of the house mouse, Mus musculus and its close relative Mus spretus. *Sci*
Data 3, 160075.

Hernandez-Segura A, Brandenburg S & Demaria M (2018) Induction and Validation of
Cellular Senescence in Primary Human Cells. *J Vis Exp*, 57782.

Hornsby PJ (2002) Cellular senescence and tissue aging in vivo. *J Gerontol A Biol Sci Med*
Sci 57, B251-256.

Imai Y, Tanave A, Matsuyama M & Koide T (2022) Efficient genome editing in wild strains
of mice using the i-GONAD method. *Sci Rep* 12, 13821.

Ito T, Teo YV, Evans SA, Neretti N & Sedivy JM (2018) Regulation of Cellular Senescence
538   by Polycomb Chromatin Modifiers through Distinct DNA Damage-and Histone
539   Methylation-Dependent Pathways. *Cell Rep* 22, 3480–3492.

540   Kang T, Moore EC, Kopania EEK, King CD, Schilling B, Campisi J, Good JM & Brem RB
541   (2023) A natural variation-based screen in mouse cells reveals USF2 as a regulator of
542   the DNA damage response and cellular senescence. *G3 (Bethesda)* 13, jkad091.

543   Kawakami M & Yamamura K-I (2008) Cranial bone morphometric study among mouse
544   strains. *BMC Evolutionary Biology*, 8 (1), 73.

545   Khan M & Gasser S (2016) Generating Primary Fibroblast Cultures from Mouse Ear and
546   Tail Tissues. *J Vis Exp*, 53565.

547   Kreis N-N, Friemel A, Zimmer B, Roth S, Rieger MA, Rolle U, Louwen F & Yuan J (2016)
548   Mitotic p21Cip1/CDKN1A is regulated by cyclin-dependent kinase 1 phosphorylation.
549   *Oncotarget* 7, 50215–50228.

550   Lee BY, Han JA, Im JS, Morrone A, Johung K, Goodwin EC, Kleijer WJ, DiMaio D & Hwang
551   ES (2006) Senescence-associated beta-galactosidase is lysosomal beta-
552   galactosidase. *Aging Cell* 5, 187–195.

553   Lee J-H, Yu WH, Kumar A, Lee S, Mohan PS, Peterhoff CM, Wolfe DM, Martinez-Vicente
554   M, Massey AC, Sovak G, Uchiyama Y, Westaway D, Cuervo AM & Nixon RA (2010)
555   Lysosomal Proteolysis and Autophagy Require Presenilin 1 and Are Disrupted by
556   Alzheimer-Related PS1 Mutations. *Cell* 141, 1146–1158.

557   Leiva-Rodríguez T, Romeo-Guitart D, Marmolejo-Martínez-Artesero S, Herrando-Grabulosa
558   M, Bosch A, Forés J & Casas C (2018) ATG5 overexpression is neuroprotective and
559   attenuates cytoskeletal and vesicle-trafficking alterations in axotomized motoneurons.
560   *Cell Death Dis* 9, 626.

561   Liu X, Ye Q, Huang Z, Li X, Zhang L, Liu X, Wu Y-C, Brockmeier U, Hermann DM, Wang Y-
562   C & Ren L (2023) BAG3 Overexpression Attenuates Ischemic Stroke Injury by
563   Activating Autophagy and Inhibiting Apoptosis. *Stroke* 54, 2114–2125.

564   Mahieu T, Park JM, Revets H, Pasche B, Lengeling A, Staelens J, Wullaert A, Vanlaere I,
565   Hochepied T, Van Roy F, Karin M, Libert C (2006) The wild-derived inbred mouse
566   strain SPRET/Ei is resistant to LPS and defective in IFN-β production. *Proceedings of*
567   *the National Academy of Sciences* 103 (7), 2292–2297.

Morgan AP, Hughes JJ, Didion JP, Jolley WJ, Campbell KJ, Threadgill DW, Bonhomme F,
Searle JB & de Villena FP-M (2022) Population structure and inbreeding in wild house
mice (Mus musculus) at different geographic scales. *Heredity (Edinb)* 129, 183–194.

Oka K, Yamakawa M, Kawamura Y, Kutsukake N & Miura K (2023) The Naked Mole-Rat
as a Model for Healthy Aging. *Annu Rev Anim Biosci* 11, 207–226.

Pagès M, Fabre P, Chaval Y, Mortelliti A, Nicolas V, Wells K, Michaux JR & Lazzari V
(2015) Molecular phylogeny of South-East Asian arboreal murine rodents. *Zoologica*
Scripta 45 (4), 349–364.

Pyo J-O, Yoo S-M, Ahn H-H, Nah J, Hong S-H, Kam T-I, Jung S & Jung Y-K (2013)
Overexpression of Atg5 in mice activates autophagy and extends lifespan. *Nat*
Commun 4, 2300.

Robbins E, Levine EM & Eagle H (1970) Morphologic changes accompanying senescence
of cultured human diploid cells. *J Exp Med* 131, 1211–1222.

Rovira M, Sereda R, Pladevall-Morera D, Ramponi V, Marin I, Maus M, Madrigal-Matute J,
Díaz A, García F, Muñoz J, Cuervo AM & Serrano M (2022) The lysosomal proteome
of senescent cells contributes to the senescence secretome. *Aging Cell* 21, e13707.

Sami S, Liu G, Hornicek F, Cates JM & Mankin HJ (2002) Membranous lipodystrophy. A
case report. *J Bone Joint Surg Am* 84, 630–633.

Saul D, Jurk D, Doolittle ML, Kosinsky RL, Monroe DG, LeBrasseur NK, Robbins PD,
Niedernhofer LJ, Khosla S & Passos JF (2023) Distinct secretomes in p16- and p21-
positive senescent cells across tissues. *bioRxiv*, 2023.12.05.569858.

Shin HR & Zoncu R (2020) The Lysosome at the Intersection of Cellular Growth and
Destruction. *Dev Cell* 54, 226–238.

Shin JY, Hong S-H, Kang B, Minai-Tehrani A & Cho M-H (2013) Overexpression of beclin1
induced autophagy and apoptosis in lungs of K-rasLA1 mice. *Lung Cancer* 81, 362–
370.

Smissen PJ & Rowe KC (2018) Repeated biome transitions in the evolution of Australian
rodents. *Molecular Phylogenetics and Evolution* 128, 182–191.

Staelens J, Wielockx B, Puimège L, Van Roy F, Guénet J-L & Libert C (2002)
Hyporesponsiveness of SPRET/Ei mice to lethal shock induced by tumor necrosis

factor and implications for a TNF-based antitumor therapy. *Proceedings of the National*
Academy of Sciences, 99 (14), 9340–9345.

Sulak M, Fong L, Mika K, Chigurupati S, Yon L, Mongan NP, Emes RD & Lynch VJ (2016)
TP53 copy number expansion is associated with the evolution of increased body size
and an enhanced DNA damage response in elephants. *eLife* 5, e11994.

Tian X, Firsanov D, Zhang Z, Cheng Y, Luo L, Tombline G, Tan R, Simon M, Henderson S,
Steffan J, Goldfarb A, Tam J, Zheng K, Cornwell A, Johnson A, Yang J-N, Mao Z,
Manta B, Dang W, Zhang Z, Vijg J, Wolfe A, Moody K, Kennedy BK, Bohmann D,
Gladyshev VN, Seluanov A & Gorbunova V (2019) SIRT6 Is Responsible for More
Efficient DNA Double-Strand Break Repair in Long-Lived Species. *Cell* 177, 622-
638.e22.

Wannemacher R, Stegmann F, Eikelberg D, Bühler M, Li D, Kohale SK, Asawapattanakul
T, Ebbecke T, Raulf M-K, Baumgärtner W, Lepenies B & Gerhauser I (2025) Infection
of a β-galactosidase-deficient mouse strain with Theiler's murine encephalomyelitis
virus reveals limited immunological dysregulations in this lysosomal storage disease.
Front Immunol 16, 1467207.

Yew T-L, Chiu F-Y, Tsai C-C, Chen H-L, Lee W-P, Chen Y-J, Chang M-C & Hung S-C
(2011) Knockdown of p21(Cip1/Waf1) enhances proliferation, the expression of
stemness markers, and osteogenic potential in human mesenchymal stem cells. *Aging*
Cell 10, 349–361.

Yuan Y, Kilpatrick BS, Gerndt S, Bracher F, Grimm C, Schapira AH & Patel S (2021) The
lysosomotrope GPN mobilises Ca2+ from acidic organelles. *Journal of Cell Science*
134, jcs256578.

Zaidi N, Maurer A, Nieke S & Kalbacher H (2008) Cathepsin D: A cellular roadmap.
Biochemical and Biophysical Research Communications 376, 5–9.

Zhao Y, Tyshkovskiy A, Muñoz-Espín D, Tian X, Serrano M, de Magalhaes JP, Nevo E,
Gladyshev VN, Seluanov A & Gorbunova V (2018) Naked mole rats can undergo
developmental, oncogene-induced and DNA damage-induced cellular senescence.
Proc Natl Acad Sci U S A 115, 1801–1806.

A

| Strain | Irradiated | NCS |

B

| Strain | Irradiated | NCS |

A

	M. musculus	*M. spretus*
Control		
Bafilomycin A1		

B

Species/Strain	M. musculus PWK	M. musculus BLG2	M. musculus CHD	M. musculus MSM	Black 6	M. domesticus TUCA	M. domesticus ManB	M. spicilegus ZRU	M. spretus SFM	M. spretus STF	M. spretus (STF)/ M. musculus (PWK) hybrid
Figure 1A.	c=45, n=3	c=6, n=2	c=8, n=2	c=8, n=2	c=8, n=2	c=8, n=3	c=10, n=3	c=9, n=2	c=12, n=2	c=25, n=3	
Figure 1B.	c=24, n=3	c=6, n=2	c=4, n=2	c=4, n=1	c=4, n=1	c=4, n=1	c=6, n=2	c=4, n=1	c=6, n=2	c=14, n=2	
Figure 1C.	c=30, n=3	c=4, n=1	c=4, n=1	c=4, n=1	c=4, n=1	c=4, n=1	c=4, n=1	c=12, n=2	c=6, n=1	c=18, n=3	
Figure 2A.	Control c=19, n=2, q=139 Irradiation c=10, n=1, q=64 Neocarzinostatin c=7, n=1, q=111									Control c=10, n=1, q=85 Irradiation c=7, n=1, q=45 Neocarzinostatin c=8, n=1, q=53	
Figure 2B.	Control c=25, n=2, q=49 Irradiation c=16, n=1, q=33 Neocarzinostatin c=17, n=1, q=66									Control c=24, n=1, q=44 Irradiation c=17, n=1, q=32 Neocarzinostatin c=19, n=1, q=62	
Figure S3A.	Control c=22, n=2 Irradiation c=24, n=3	Control c=6, n=2 Irradiation c=6, n=2	Control c=4, n=1 Irradiation c=4, n=2	Control c=4, n=1 Irradiation c=4, n=1	Control c=4, n=1 Irradiation c=4, n=1	Control c=4, n=1 Irradiation c=4, n=1	Control c=6, n=2 Irradiation c=6, n=2	Control c=3, n=1 Irradiation c=4, n=1	Control c=6, n=2 Irradiation c=6, n=2	Control c=13, n=2 Irradiation c=14, n=2	
Figure S3B.	Control c=23, n=2 Neocarzinostatin c=30, n=3	Control c=3, n=1 Neocarzinostatin c=4, n=1	Control c=4, n=1 Neocarzinostatin c=4, n=1	Control c=4, n=1 Neocarzinostatin c=4, n=1	Control c=4, n=1 Neocarzinostatin c=4, n=1	Control c=4, n=1 Neocarzinostatin c=4, n=1	Control c=4, n=1 Neocarzinostatin c=4, n=1	Control c=6, n=2 Neocarzinostatin c=12, n=2	Control c=6, n=1 Neocarzinostatin c=6, n=1	Control c=12, n=2 Neocarzinostatin c=18, n=3	
Figure S4A. & Figure S4B.	Control c=131, n=2 Irradiation c=150, n=2 Neocarzinostatin c=96, n=2	Control c=71, n=2 Irradiation c=48, n=1 Neocarzinostatin c=28, n=1	Control c=52, n=1 Irradiation c=80, n=2 Neocarzinostatin c=32, n=1	Control c=71, n=2 Irradiation c=100, n=1 Neocarzinostatin c=48, n=1	Control c=24, n=1 Irradiation c=24, n=1 Neocarzinostatin c=24, n=1	Control c=60, n=1 Irradiation c=80, n=1 Neocarzinostatin c=32, n=1	Control c=52, n=1 Irradiation c=80, n=1 Neocarzinostatin c=40, n=1	Control c=12, n=1 Irradiation c=16, n=1 Neocarzinostatin c=8, n=1	Control c=48, n=1 Irradiation c=48, n=1 Neocarzinostatin c=32, n=1	Control c=56, n=2 Irradiation c=56, n=2 Neocarzinostatin c=48, n=1	
Figure S5.	Control c=8, n=1 Irrad Dose 1000 c=10, n=1 Irrad Dose 1500 c=10, n=1 Irrad Dose 2000 c=10, n=1 Irrad Dose 2500 c=10, n=1									Control c=8, n=1 Irrad Dose 1000 c=10, n=1 Irrad Dose 1500 c=10, n=1 Irrad Dose 2000 c=10, n=1 Irrad Dose 2500 c=10, n=1	
Figure S6A.	Control c=3, n=1 Irradiation c=3, n=1									Control c=3, n=1 Irradiation c=3, n=1	
Figure S6B.											Control c=3, n=1 Irradiation c=3, n=1
Figure S7A. & Figure S7B.	Control c=6, n=1 Irradiation c=3, n=1 Neocarzinostatin c=3, n=1										Control c=6, n=1 Irradiation c=3, n=1 Neocarzinostatin c=3, n=1
Figure S8.	Control c=3, n=1 Irradiation c=3, n=1										Control c=3, n=1 Irradiation c=3, n=1
Figure S9.	Control c=6, n=1 Irradiation c=7, n=1 Neocarzinostatin c=7, n=1									Control c=7, n=1 Irradiation c=7, n=1 Neocarzinostatin c=7, n=1	
Figure S10A. & Figure S10B.	Blank c=9, n=1 Control c=20, n=1 Bafilomycin A1 c=16, n=1									Blank c=6, n=1 Control c=20, n=1 Bafilomycin A1 c=16, n=1	