1
2

Human differentiated adipocytes can serve as surrogate mature adipocytes for adipocyte-derived extracellular vesicle analysis.

Bradley L. Butsch[1#], Mangesh Dattu Hade[1#], Paola Loreto Palacio[1], Kim Truc Nguyen[2], Dharti Shantaram[3], Sabrena Noria[4], Stacy A. Brethauer[4], Bradley J. Needleman[4], Willa Hsueh[3], Eduardo Reátegui[2], Setty M. Magaña[1]*

[1]Department of Pediatrics, Center for Clinical and Translational Research, Nationwide Children's Hospital, Columbus, Ohio.

[2]Department of Chemical and Biomedical Engineering, The Ohio State University, Columbus, Ohio.

[3]Diabetes and Metabolism Research Center, The Ohio State University, Columbus, Ohio.

[4]Center for Minimally Invasive Surgery, Department of Surgery, The Ohio State University College of Medicine, Columbus, Ohio.

# Authors contributed equally.

Running Title: Human differentiated adipocyte extracellular vesicles

*Corresponding Author: Setty Magaña, MD, PhD

Address: Nationwide Children's Hospital

                  Department of Pediatrics, Division of Neurology

                  700 Children's Drive

                  Columbus, OH 43205

Telephone: 614-722-4625

Fax: 614-722-4633

Email: Setty.Magana@nationwidechildrens.org

Abstract

Obesity is a growing global health concern, contributing to diseases such as cancer, autoimmune disorders, and neurodegenerative conditions. Adipose tissue dysfunction, characterized by abnormal adipokine secretion and chronic inflammation, plays a key role in these conditions. Adipose-derived extracellular vesicles (ADEVs) have emerged as critical mediators in obesity-related diseases. However, the study of mature adipocyte-derived EVs (mAdipo-EVs) is limited due to the short lifespan of mature adipocytes in culture, low EV yields, and the low abundance of these EV subpopulations in the circulation. Additionally, most studies rely on rodent models, which have differences in adipose tissue biology compared to humans. To overcome these challenges, we developed a standardized approach for differentiating human preadipocytes (preAdipos) into mature differentiated adipocytes (difAdipos), which produce high-yield, human adipocyte EVs (Adipo-EVs). Using visceral adipose tissue from bariatric surgical patients, we isolated the stromal vascular fraction (SVF) and differentiated preAdipos into difAdipos. Brightfield microscopy revealed that difAdipos exhibited morphological characteristics comparable to mature adipocytes (mAdipos) directly isolated from visceral adipose tissue, confirming their structural similarity. Additionally, qPCR analysis demonstrated decreased preadipocyte markers and increased mature adipocyte markers, further validating successful differentiation. Functionally, difAdipos exhibited lipolytic activity comparable to mAdipos, supporting their functional resemblance to native adipocytes. We then isolated preAdipo-EVs and difAdipo-EVs using tangential flow filtration and characterized them using bulk and single EV analysis. DifAdipo-EVs displayed classical EV and adipocyte-specific markers, with significant differences in biomarker expression compared to preAdipo-EVs. These findings demonstrate that difAdipos serve as a reliable surrogate for mature adipocytes, offering a consistent and scalable source of adipocyte-derived EVs for studying obesity and its associated disorders.

Keywords: extracellular vesicles, adipocyte, adipose, adipocyte-derived extracellular vesicles, obesity

Introduction

   The widespread prevalence of obesity has escalated into a formidable global
health emergency, placing significant burdens on public health systems worldwide [1,2]. In
2019, an estimated 5 million obesity-related deaths occurred worldwide [3]. Obesity is
intricately linked to various metabolic disorders, including type 2 diabetes mellitus,
cardiovascular diseases and non-alcoholic fatty liver disease [4-6]. Moreover, obesity is
also implicated in the pathogenesis of several cancers, autoimmune diseases, and
neurodegenerative disorders [7,8]. The complex interplay between obesity and these
diseases underscores the multifaceted role of adipose tissue in maintaining metabolic
homeostasis and overall health [9,10]. Adipose tissue, a dynamic endocrine organ,
significantly influences energy balance, metabolic regulation, and immune responses [11].
Dysfunctional adipose tissue, characterized by altered adipokine secretion, chronic
inflammation, and impaired adipocyte differentiation, is a hallmark of obesity and a
crucial factor in the emergence of obesity-related comorbidities [12,13]. Recent research
advances have illuminated the pivotal role of extracellular vesicles (EVs) in mediating
intercellular communication and regulating metabolic processes [14,15]. EVs are small,
membrane-bound particles secreted by cells that transport bioactive molecules (e.g.
proteins, lipids, and nucleic acids) to recipient cells, thereby altering recipient cell
genotypic and phenotypic function [14-16].

   Mounting evidence indicates that obesity is associated with enhanced production
of adipose-derived EVs (ADEVs), which are instrumental in the pathogenesis of obesity
and its related metabolic complications [17-19]. ADEVs are implicated in various
physiological and pathological processes, including inflammation, insulin resistance,
and lipid metabolism [19,20].

   Despite the growing interest in studying ADEVs, there is a dearth of human ADEV
studies, partly due to the technical and biological limitations of obtaining EVs from
mature human adipocytes (mAdipo-EVs). mAdipos exhibit short viability in culture,
produce low yields of EVs, and are difficult to handle due to their nonadherent
properties [21-23]. To address these limitations, and with the end goal of developing a

robust human translational paradigm for isolating mature adipocyte-derived EVs, we

first developed a surrogate mAdipo model by differentiating human preAdipos into

mature-like difAdipos with functional, genotypical, and phenotypical similarities to

mature adipocytes. Secondly, we used the validated difApos as a reliable and

surrogate source of human adipocyte EVs, demonstrating classic EV and adipose

markers.

In summary, our study provides a reproducible and reliable source of human

mature-like adipose EVs. By leveraging human-based models, we aim to provide a

more relevant understanding of the role of adipocyte-derived EVs in obesity-related

disorders, thus addressing the critical need for human-based obesity EV paradigms [24].

Results

Multiparametric isolation and characterization of ADEVs

We have developed a comprehensive pipeline for isolating and characterizing EVs from

various human adipose tissue biospecimens (**Figure 1, Table S2)[25]**. While most human

EV biomarker studies have focused on biofluids or in vitro EV sources, isolating tissue-

derived EVs presents technical challenges that must be addressed to maintain the

integrity of both the cells and the EVs[29]. To ensure our isolation method produced intact

adipocyte-derived EVs, we performed extensive bulk and single EV characterization, as

detailed below in **Figure 1.**

Figure 1. Workflow for the isolation, differentiation, and characterization of human

preadipocytes, differentiated adipocytes, and their EVs. (A): Visceral adipose tissue

obtained from bariatric patients was enzymatically digested using Collagenase A at

37°C for 30–40 minutes, followed by filtration through 300 μm and 100 μm cell strainers.

Subsequent centrifugation (500 × g, RT, 10 min) yielded three distinct fractions: stromal

vascular fraction (SVF), SVF supernatant (SVF-SN), and mature adipocytes (mAdipos).

The SVF-derived preadipocytes (preAdipos) were cultured and induced to differentiate

into differentiated mature adipocytes (difAdipos) over 28 days. Conditioned media from

adipocyte cultures were collected and processed using tangential flow filtration (TFF) for

EV isolation. (B): Characterization of preAdipos, difAdipos and mAdipos included

functional assays assessing lipolysis, morphological analysis via brightfield microscopy,

and gene expression analysis using quantitative PCR. (C): Isolated EVs from preAdipos

(preAdipo-EVs) and difAdipos (difAdipo-EVs) underwent extensive characterization,

including immunoblot analysis of EV and adipocyte markers (CD9, CD81, CD63, and

ADIPOQ), size distribution and concentration analysis via microfluidic resistive pulse

sensing (MRPS), ultrastructural evaluation using transmission electron microscopy

(TEM), and single-EV multiplexing analysis. extracellular vesicles (EVs); stromal

vascular fraction (SVF), SVF supernatant (SVF-SN), mature adipocytes (mAdipos);

preadipocytes (preAdipos); differentiated mature adipocytes (difAdipos); microfluidic

resistive pulse sensing (MRPS); transmission electron microscopy (TEM)

Visceral adipose-derived preadipocytes serve as a surrogate source for mature

adipocytes

Our robust and systematic human-based protocol for isolating, differentiating, and

characterizing SVF-derived preadipocytes yielded a reliable surrogate source of

differentiated mature adipocytes (difAdipos) (see material and methods). Preadipocytes

exhibited significant morphological changes during the differentiation process,

accumulating larger lipid droplets and developing a mature phenotype, as confirmed by

brightfield imaging (Figure 2B).

Figure 2. Human preadipocytes undergo differentiation and acquire a mature adipocyte phenotype. (A) Differentiation timeline: At Day -1, cells reached ~80% confluency in growth medium (GM). Differentiation was initiated on Day 0 following a 24-hour incubation in GM by replacing the medium with adipogenic induction medium 1 (AIM1). The transition to adipogenic induction medium 2 (AIM2) was performed to facilitate further maturation, leading to the acquisition of a differentiated mature adipocyte (difAdipo) phenotype by Day 28. (B) Morphological changes during adipogenesis: (a) Representative brightfield microscopy images illustrate the progressive morphological changes occurring during preadipocyte differentiation into difAdipos. Lipid droplet accumulation becomes progressively evident throughout the differentiation process. (b) The morphology of difAdipos at Day 28 closely resembles that of primary mature adipocytes (mAdipos), which were directly isolated from patient-derived visceral adipose tissue and maintained in culture for 24 hours. Images are representative of three independent biological replicates (n = 3). Dashed boxes highlight selected regions, which are shown as enlarged high-magnification images in the subsequent panels to provide detailed visualization of cellular morphology. Scale bar = 50 μm for main images; 10 μm for enlarged high-magnification images. GM, growth medium; AIM1, adipogenic induction medium 1; AIM2, adipogenic induction medium 2. differentiated mature adipocyte (difAdipo); mature adipocytes (mAdipos)

Differential expression of preadipocyte and mature adipocyte markers during adipocyte differentiation

We next investigated gene transcripts of known preAdipo markers (OCT4, PREF-1, and GATA3) [30-33] [30] and mature adipocyte markers (ADIPOQ, PLIN1, and PPARG) [34] at different time points during adipocyte differentiation (**Figure 3**). We hypothesized that difAdipos would acquire a genetic program similar to mAdipos and lose the undifferentiated transcript profile that characterizes preAdipos. As expected [35,36], OCT4 expression was low at Day 0 (D0), peaked at D7 (P< 0.001), and decreased as differentiation progressed and cells committed adipogenesis (**Figure 3A**).

We saw a similar pattern in PREF-1 expression levels. PREF-1 is an inhibitor of

adipogenesis and it is downregulated as cells undergo differentiation [36]. Consistent with

this observation, preAdipo PREF-1 levels were relatively low at D0, peaked at D7(P<

0.0001), and were decreased by D28 and in mAdipos (**Figure 3A**). In contrast to OCT4

and PREF-1, GATA3 levels persisted in both the early and intermediate differentiation

stages. GATA3 expression peaked at D7 (P< 0.01) and D14, decreased through D21

and D28, but remained elevated in mAdipo compared to D28 (**Figure 3A**). This

expression pattern is consistent with studies showing GATA3's involvement in regulating

adipogenesis [37,38].

Concurrently, the expression of the mature adipocyte markers ADIPOQ, PLIN1,

and PPARG, showed a clear increasing trend during differentiation, as previously

reported [38-40]. ADIPOQ expression was low at D0 and D7, increased steadily from D14

onwards, and peaked at D28 (P< 0.0001), with levels comparable to those seen in our

mAdipos (**Figure 3B**). PLIN1 plays a crucial role in regulating lipid storage and

mobilization by protecting lipid droplets from lipolysis in adipocytes [41]. We observed

gradually increasing levels of PLIN1 expression from D0 onward, significantly increasing

at D21 and D28 (P< 0.0001), signifying lipid accumulation and adipocyte maturation.

Similarly, PPARG expression progressively increased from D0 to D28, with the highest

expression observed at D21 and D28 (P< 0.0001), and slightly lower expression in

mAdipo, highlighting its crucial role as a regulator of adipogenesis (**Figure 3B**). In

summary, the dynamic transcriptional program we observed reflects the successful

differentiation of preadipocytes into difAdipos exhibiting mature adipocyte transcriptional

profiles.

A

B

Figure 3. Transcript profile in preadipocytes, differentiated adipocytes and

mature adipocyte (A) Relative mRNA expression levels of preadipocyte markers

(OCT4, PREF1, GATA3) and **(B)** mature adipocyte markers (ADIPOQ, PLIN1, PPARG)

were measured at Day 0 (D0), D7, D14, D21, D28, and in terminally differentiated

mAdipo. OCT4 expression peaked at D7 and decreased thereafter, indicating early

differentiation. PREF-1 expression increased to a peak at D7 and decreased towards

D28, consistent with its inhibitory role in adipogenesis. GATA3 expression peaked at D7

and D14 and remained elevated in mAdipo, indicating its involvement in early

differentiation. Adipocyte markers ADIPOQ, PLIN1, and PPARG exhibited a progressive

increase, peaking at D28 and in mAdipo, reflecting mature adipocyte function (n=3). The

statistical analysis was performed using ordinary one-way ANOVA with Bartlett's test.

OCT4, octamer-binding transcription factor 4; PREF-1, preadipocyte factor 1; GATA3,

GATA-binding protein 3; ADIPOQ, adiponectin; PLIN, perilipin; PPARG, peroxisome

proliferator-activated receptor gamma. A value of P < 0.05 was deemed statistically

significant (*< 0.05; **< 0.01; ***< 0.001; ****< 0.0001).

Differentiated adipocytes exhibit lipolytic function comparable to mature

adipocytes

Preadipocytes differentiated in vitro may not retain the same biophysical properties of

mature adipocytes that are differentiated in vivo[42]. Therefore, we next sought to

functionally validate the mAdipo phenotype of our difAdipos by investigating their ability

to perform the prototypic mature adipocyte function of lipolysis [43-45]. Cells were

stimulated with the lipolytic agent, isoproterenol, and lipolysis was assessed by

quantifying the stimulated release of glycerol into the culture medium (**Figure 4**). We

hypothesized that the lipolytic activity of difAdipos would not differ from that of mAdipos,

suggesting that our difAdipos could serve as functional surrogates of mAdipos.

Compared to undifferentiated D0 preadipocytes, D28 difAdipos achieved a comparable

degree of lipolytic activity as mAdipos. These findings demonstrate our ability to

generate human difAdipos that are not only morphologically and genotypically

comparable to mAdipos, but also functionally resemble mAdipos.

Figure 4. Differentiated adipocytes (difAdipos) exhibit lipolytic function comparable to mature adipocytes (mAdipos). Lipolytic activity was assessed in preAdipos, difAdipos, and mAdipos by stimulating cells with isoproterenol and measuring glycerol release. D0 difAdipos reflect undifferentiated adipocytes and exhibited low-level lipolytic activity similar to HEK293T cells (negative control). D28 difAdipos exhibited lipolytic activity comparable to mAdipos and glycerol (positive control), validating the differentiation protocol and confirming that difAdipos functionally resemble mature adipocytes in their lipolytic activity. Statistical analysis was performed using one-way ANOVA with Sidaks multiple comparison test. A value of $P < 0.05$ was deemed statistically significant (*< 0.05; **< 0.01; ***< 0.001; ****< 0.0001).

Multiparametric characterization of EVs from differentiated adipocytes

After validating our difAdipos as robust surrogates for mAdipos, we next sought to characterize EVs from paired preAdipos, difAdipos, and mAdipos obtained from the

same individual (**Figure 5A**). Microfluidic resistive pulse sensing analysis revealed a

relatively narrow particle size distribution among the EVs from preAdipo, difAdipo, and

mAdipo. Median particle diameters were similar across the three Adipo populations

(88.5 nm for preAdipo, 91.8 nm for difAdipo, and 85.1 nm for mAdipo). All three ADEV

sources had similar mean particle concentrations: 6.26E+10 particles/mL for mAdipos,

3.47E+10 particles/mL for preAdipos and 2.33E+10 particles/mL for difAdipos **(Figure**

5A). These results suggest that our differentiation protocol did not alter the ability of

difAdipos to generate small EVs that were similar in size and concentration to native

populations (ie. preAdipos and mAdipos), which did not undergo in vitro maturation.

TEM revealed a polydispersed and prominent population of small EVs with the

characteristic artefactual 'cup-shaped' morphology **(Figure 5B).** Immunoblotting of

difAdipo-EVs demonstrated a heterogeneous expression of the typical EV markers

CD9, CD81, and CD63 and the absence of the non-EV marker calnexin (CANX) **(Figure**

5C). The adipocyte marker, ADIPOQ, was also detected (**Figure 5C**). Collectively, these

findings highlight the diversity of tissue-derived EV populations from varied adipocyte

sources within the same individual and at different stages of adipocyte differentiation.

Figure 5. Multiparametric characterization of matched adipose tissue-derived EVs from the same bariatric patients. (A) MRPS analysis of EVs from preAdipo, difAdipo, and mAdipo from the same individuals (n = 3 biological replicated). Median particle diameters: 88.5 nm (preAdipo), 91.8 nm (difAdipo), 85.1 nm (mAdipo). Mean particle concentrations: 3.47E+10 particles/mL (preAdipo), 2.33E+10 particles/mL (difAdipo), 6.26E+10 particles/mL (mAdipo). (B) TEM imaging showed a polydispersed population of particles, including small EVs with a characteristic 'cup-shaped' morphology. (C) Immunoblot analysis of EVs from difAdipo reveals heterogeneous expression of typical EV (CD9, CD81, CD63), non-EV (CANX), and adipose-specific (ADIPOQ) markers. MRPS, microfluidic resistive pulse sensing; TEM, transmission electron microscopy.

High-resolution, simultaneous detection of human adipose-derived EV proteins and RNA using total internal reflection fluorescence (TIRF) microscopy

TIRF microscopy was utilized to achieve simultaneous detection of human adipose-derived EV proteins and RNA and colocalization of the same (**Figure 6 A, B).**

First, we investigated the colocalization of EV surface proteins and EV mRNA cargo in

preAdipo-EVs and difAdipo-EVs to confirm the specificity of adipocyte-EVs. DifAdipo-

EVs revealed high colocalization between OCT4 protein and ADIPOQ mRNA,

confirming the adipocyte-specificity of the EVs. The tetraspanin marker CD63 showed

less to no colocalization with other biomarkers (Figure 6A), underscoring the

heterogeneous expression profile of tetraspanins. Similar results were observed for

preAdipo-EVs (results not shown).

Next, we quantified the relative fluorescence intensity of EV surface markers and mRNA

cargo (**Figure 6B**). DifAdipo-EVs collected on day 28, compared to preAdipo-EVs

collected on day 2, expressed a significantly higher level of the canonical EV marker

CD63 (***< 0.001). Interestingly, the protein marker for undifferentiated adipocytes,

OCT4, was also significantly higher on difAdipo-EVs than preAdipo-EVs (*< 0.05). The

expression levels of adipose tissue-specific protein and gene markers, such as

adipocyte plasma membrane-associated protein (APMAP) and ADIPOQ, were similar

between the difAdipo-EVs and preAdipo-EVs. In summary, APMAP and ADIPOQ

exhibited stable expression levels across the maturation process, while CD63 and

OCT4 were dynamically regulated during the same differentiation process.

Figure 6. Single EV multiparametric characterization comparing preAdipo-EVs (day 2) and difAdipo-EVs (day 28). (A) Representative TIRFM images illustrating (day 28) EVs were captured by anti-CD9 and anti-CD63 antibodies and immobilized on a biochip surface. Detection antibodies against CD63, OCT, and APMAP were used to identify EV surface proteins. A fluorescence-labeled molecular beacon detected the mRNA ADIPOQ within those EVs. The merged image reveals difAdipo-EVs defined by colocalization of OCT4 and mRNA ADIPOQ, CD63 and Oct4, CD63 and mRNA ADIPOQ, CD63 and APMAP Scale 10 µm (B) The relative fluorescence intensity of EVs collected at day 2 and day 28 was analyzed for the tetraspanin marker CD63, the undifferentiated cell marker OCT4, the adipocyte plasma membrane-associated protein APMAP, and the adipose tissue gene marker mRNA ADIPOQ. OCT4, octamer-binding transcription factor 4; ADIPOQ, adiponectin; APMAP, astrocyte membrane-associated protein. Statistical analysis was performed using students test. A value of P < 0.05 was deemed statistically significant (*< 0.05; **< 0.01; ***< 0.001; ****< 0.0001).

Discussion

Adipose-derived EVs are increasingly recognized as critical orchestrators in obesity-associated diseases[46]. Limited studies have characterized human adipocyte EVs[47-51], however, no study has systematically studied paired difAdipo-EVs and mAdipo-EVs at bulk and single EV resolutions. Herein, we present a reproducible and scalable surrogate source of human mature-like adipocytes (difAdipos) that exhibit phenotypic, genotypic, and functional properties paralleling those of paired mature adipocytes (mAdipos). Furthermore, we demonstrate that in vitro difAdipos can release ADEVs comparable to ADEVs released by mature adipocytes. Finally, we provide the first high-resolution, single EV analysis of human difAdipo-EVs, demonstrating simultaneous adipocyte-enriched gene transcript and surface protein markers.

Due to recognized cross-species differences[52,53] between mouse and human adipose biology, adipocyte EVs from rodent models have limited direct human translational utility. Several alternate approaches to circumvent the cross-species differences include

studying in vitro differentiated human adipocytes, adipose tissue explants, and

circulating adipocyte-enriched EV subpopulations in human biofluids[54]. Kranendonk and

colleagues were amongst the earliest groups to characterize human adipocyte EVs

(defined as adiponectin and fatty-acid binding protein 4 positive EVs), isolated from the

Human Simpson Golabi Behmel Syndrome (SGBS) cell line[55] . More recently, a study

by Clement and collaborators isolated primary human adipocytes from dermolipectomy-

derived subcutaneous adipose tissue and demonstrated the functional ability of isolated

human adipocyte EVs to metabolically remodel melanoma cells into a more invasive

phenotype[56]. Several human ADEV studies have utilized tissue explants for ADEVs

rather than mature adipocytes[51,57] . Human adipose tissue explants have the advantage

of overcoming the cross-species limitations of mouse models but confound the ability to

investigate the potential pathobiological role of EVs shed solely from mature

adipocytes[58]which are the most abundant and primary functional metabolic cells within

visceral adipose tissue[49,59].

A less invasive and more feasible approach for studying human ADEVs involves

interrogating the circulating pool of all EVs and attempting to enrich for EV

subpopulations that contain 'adipocyte-specific' markers. Connolly and colleagues were

the first to confirm human EVs containing adipose-enriched markers within the

circulating pool of plasma-derived EVs from obese individuals[60] . In this study, the

authors applied contemporaneous ISEV guidelines[61] for EV research and used

complementary EV isolation and enrichment methods to attempt to account for non-

vesicular co-isolates and non-adipose EV populations.

One of the formidable challenges in studying cell and tissue-specific EVs, such as

ADEVs, arises from their low abundance and high signal-to-noise ratio[58,62] . Adipose

EVs are estimated to comprise >80% of tissue-derived EVs in human plasma and can

be the primary source of circulating miRNAs; however, tissue-derived EVs comprise

<0.2 % of the circulating human plasma pool [63-65]. This limitation underscores the

critical need for more advanced methods to study low-abundant, rare EV

subpopulations. To address this limitation, we first produced a surrogate source of

functional mature adipocytes in vitro and demonstrated their ability to shed EVs

comparable in adipocyte-specific markers to EVs shed by adipocytes that matured in
vivo.

To our knowledge, no studies have characterized the biophysical properties of human
adipose-derived EVs in the pre- and post-adipogenic phases from paired human
biospecimens. One study conducted a systematic phenotypic comparison between
immature and mature mouse ADEVs[66]. In this study, the authors characterized the
biophysical properties of pre- and post-adipogenic adipocyte-derived EVs from 3T3-L1
cells. Similar to the Connolly study, we found the highest expression of the preAdipo
marker PREF-1 in the earliest stages of differentiation and the highest expression of the
mAdipo marker ADIPOQ in the later stages of maturation. They also found increased
secretion of small EVs in the early stages of differentiation, slightly increased CD9 and
CD63 protein via time-resolved fluorescence, and a higher proportion of phospholipids
associated with cell signaling. Interestingly, the authors found EV-unique
characteristics that differed from the 3T3-L1 parental cell, suggesting modifications that
could impact EV-mediated intercellular communication[66].

Finally, we applied single EV multiplexing analytics to validate the adipocyte-specificity
of our difAdipo-EVs. Colocalization of EV surface proteins and internal cargo can
increase sensitivity for detecting low-abundance EV subpopulations and can also
enhance the biomarker utility of EVs to detect earlier stages of disease[70,71] . Therefore,
our single ADEV multiplexing platform holds great promise for investigating novel and
robust, adipocyte-specific EVs in obesity-associated diseases.

Conclusion

Our study presents a novel paradigm for advancing the translational application of
human ADEVs by using human difAdipos as a surrogate for mature adipocytes, thereby
augmenting the harvesting of viable human-derived, mature adipocyte EVs. The model
established provides a high-yield, functionally equivalent source of EVs, facilitating more
consistent and relevant research into the role of ADEVs in obesity and metabolic
diseases. Our findings underscore the importance of adipocyte differentiation on EV

content and function, revealing significant differences in biomarker profiles between

preAdipo-EVs and difAdipo-EVs. This differentiation may impact the bioactive cargo

within the EVs, influencing recipient cell behavior and contributing to metabolic

complications associated with obesity.

Future research should explore the specific mechanisms by which difAdipo-EVs

influence metabolic processes and their potential as biomarkers or therapeutic targets in

obesity-related conditions. This study herein lays the groundwork for developing

therapeutic strategies that modulate ADEVs' effects, offering promising avenues for

mitigating the adverse consequences of obesity on metabolic health and reducing the

risk of associated diseases.

Materials and methods

Study participants

This study complies with all relevant ethical guidelines. The Ohio State University

Institutional Review Board (IRB# 2014H0471) approved the study, and all participants

provided written informed consent. Visceral adipose tissue (VAT) samples were

collected from patients (n = 11, mean age: 41.6 years, SD: 12.98; mean BMI: 40.97, SD:

6.00) undergoing elective bariatric surgery at The Ohio State University (OSU) Center

for Minimally Invasive Surgery. Clinical data are summarized in Table S1.

Adipose tissue dissociation and mature adipocyte culture for EV isolation

Adipose tissue samples were collected in the operating room during the bariatric

surgery and processed within 1 hr using collagenase digestion, as previously described

[25]. The cell suspension was centrifuged at 500 x g for 10 min to generate three distinct

phases (mature adipocytes, the stromal vascular fraction (SVF), and the SVF

secretome). Floating mature adipocytes were carefully removed from the top layer and

kept for culture. The supernatant was then carefully aspirated to avoid disturbing the

SVF pellet, which was resuspended in sorting media for the pre-adipocyte isolation and

differentiation protocol (see below). Mature adipocytes were cultured for 24 hr in

Dulbecco's Modified Eagle Medium/Ham's Nutrient Mixture F-12 (DMEM/F12),

supplemented with 40% M199 basal medium with Earle's salts and L-glutamine

(MCDB201), 2% fetal bovine serum (FBS), 1x penicillin/streptomycin, 1 nM

dexamethasone, 0.1 mM L-ascorbic acid 2-phosphate (LAAP), 1x Insulin-Transferrin-

Selenium (ITS) Mix, 1x linoleic acid-albumin, 5 μg/μL insulin, and 1 nM triiodothyronine

(T3).

402   Preadipocyte (preAdipo) isolation and differentiation into differentiated mature

403   adipocytes (difAdipos)

The resuspended SVF pellet was filtered through a 100 μm cell strainer into fresh tubes,

repeating the washing step to ensure thorough isolation of SVF cells. Following another

round of centrifugation (at 1200 rpm for 10 min at RT or 4 °C), and aspiration of the

supernatant, the SVF pellet was treated with ACK lysis buffer on ice for three minutes to

lyse any remaining red blood cells, followed by dilution and filtration through a 40 μm

cell strainer. Finally, the SVF pellet containing cells were re-suspended in sorting media

and seeded into culture plates until the cells reached approximately 80% confluency.

The differentiation process commenced with a 24 hr incubation in growth media,

designated as day 0. Throughout the differentiation period, media was replaced every

48 hr. Cells were maintained for up to 28 days in adipogenic induction media 1 and 2

(AIM1 and AIM 2). For media composition, see Table S3.

416   ADEV isolation via tangential flow filtration (TFF)

PreAdipo, difAdipo and mAdipo culture media was filtered through 0.2 μm filters and

then concentrated to 5 mL. The concentrated media was diafiltered using TFF with a

500 kDa filter for purification at a flow rate of 35 mL/min, as previously described [25].

Following TFF, the retentates were further concentrated using centrifugal units (30 kDa

MWCO, MilliporeSigma Amicon Ultra Centrifugal Filter Unit, Fisher Scientific) at 4000 ×

g for 30 min to a final volume of 100 μL.

423   Microfluidic resistive pulse sensing (MRPS)

424   Particle concentration and size distribution were measured using microfluidic resistive

425   pulse sensing with the Spectradyne nCS1 instrument (Spectradyne, Torrance, CA),

426 equipped with C-400 polydimethylsiloxane cartridges to cover a size range of

427 approximately 65-400 nm [25]. Data analysis was performed using the nCS1 Data

428 Analyzer (Spectradyne, Torrance, CA).

Western blotting

EV samples were lysed in radioimmunoprecipitation assay (RIPA) buffer (Thermo

Scientific) and treated with protease and phosphatase inhibitor (Thermo Scientific) for

15 min on ice. Protein concentration was quantified using a Micro BCA™ Protein Assay

Kit (Thermo Scientific). Equivalent amounts of protein in Laemmli buffer (with reducing

agent 2-mercaptoethanol, Sigma-Aldrich) were electrophoresed on 4-20% Mini-

PROTEAN® TGX Stain-Free gels (Bio-Rad) and then transferred onto a polyvinylidene

fluoride (PVDF) membrane (Bio-Rad). Membranes were blocked and then probed with

primary antibody diluted in TBS-T overnight at 4 °C, followed by incubation with

horseradish peroxidase-conjugated secondary antibody for 1 hr at room temperature.

Immunoreactivity was detected using enhanced chemiluminescence (ECL) solutions

(Bio-Rad) and visualized using a Bio-Rad ChemiDoc™ MP imaging system.

Quantification of real-time polymerase chain reaction (qRT-PCR)

Total RNA was isolated using the RNeasy Mini Kit (Qiagen) according to the

manufacturer's instructions and quantified using a NanoDrop spectrophotometer. The

isolated RNA was reverse-transcribed into cDNA using the High-Capacity cDNA

Reverse Transcription Kit (Applied Biosystems). Real-time PCR was performed with the

PowerUp™ SYBR™ Green Master Mix (Thermo Fisher) on an Applied Biosystems real-

time PCR system. The sequences of the forward and reverse primers are listed in Table

S4. β-actin was used as the internal normalization control.

Lipolysis assay

To investigate mature adipocyte functional activity, a lipolysis assay was performed

using the Lipolysis Assay Kit (Abcam) per the manufacturer's instructions. Day 0

(preAdipo), day 28 (difAdipo), and cultured HEK293 (negative control) cells were

washed twice with Lipolysis Wash Buffer, then stimulated with 100 nM isoproterenol for

4 hr. Absorbance at 570 nm was measured using the SpectraMax iD3 (Molecular
Devices), and glycerol concentration was quantified using a standard curve.

Transmission electron microscopy (TEM)

TEM was performed as previously described[26]. Briefly, TEM grids were first treated with
plasma for high surface EVs absorption. Subsequently, 10 µL of EVs samples were
drop-casted onto the plasma-treated surface. The samples were incubated for 1 minute
and then blotted with filter paper to remove excess liquid. The TEM grids were washed
twice in DI water and stained with UranyLess EM contrast stain (Electron Microscopy
Science) for 22 sec. The TEM grids were dried overnight prior to imaging. TEM images
were obtained using Tecnai TF-20 microscope under 200kV, bright-field imaging mode.

Single EV analysis using total internal reflection fluorescence microscopy (TIRF)

The isolated EVs were captured and analyzed on a gold functionalized biochip following
our previous protocol [25]. Briefly, the EVs collected on day 2 (preAdipo-EVs) and day 28
(difAdipo-EVs) were permeabilized and hybridized with molecular beacons (Table S5) in
0.5 × Tris EDTA (TE) buffer at 0.2 µM concentration for 2 hr at 37 °C in a dark
environment to facilitate molecular beacon hybridization to the target RNA. The EV
samples were incubated on the biochip surface and captured by functionalized anti-
CD63 and anti-CD9 antibodies. Proteins were detected using fluorescent-dye-
conjugated antibodies. Finally, a 10 × 10 array of images was acquired via total internal
reflection fluorescence microscope (TIRFM; Nikon, Melville, NY) for each well. Relative
and total fluorescence intensities of the sample were obtained from custom-built
algorithms that were previously reported [27,28]
476

Data availability

478 All data generated or analyzed during this study are included in this manuscript and its
479 supplementary information file. Further inquiries can be directed to the corresponding
480 author.
481

Author contributions

Conceptualization: BLB., M.D.H., P.L.P., W.H., E.R., S.M.M,

Study design and funding acquisition: E.R., S.M.M.

Investigation and performed research: M.D.H., B.L.B., P.L.P., D.S., K.T.N., S.N., S.A.B.,

B.J.N.

Organization and data analysis: M.D.H., B.L.B, K.T.N

Completed the original draft of the manuscript: M.D.H., B.L.B., S.M.M

Data interpretation, revision, and manuscript correction: M.D.H., B.L.B., K.T.N., E.R.,

S.M.M.

All authors approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Correspondence and requests for materials should be addressed to Senior

Corresponding author.

Acknowledgments

We would like to thank William Ray for his illustrative media expertise in designing the

figures for this project. SMM was supported by intramural funding from the Abigail

Wexner Research Institute at Nationwide Children's Hospital and NINDS

(5K12NS098482-05). ER was supported by National Institutes of Health (NIH) grants

UG3/UH3TR002884 and U18TR003807. The content is solely the responsibility of the

authors and does not necessarily represent the official views of the National Institutes of

Health.

References

1   Blüher, M. Obesity: global epidemiology and pathogenesis. *Nature Reviews*
Endocrinology **15**, 288-298 (2019). https://doi.org:10.1038/s41574-019-0176-8
2   Seidell, J. C. & Halberstadt, J. The global burden of obesity and the challenges of
prevention. *Ann Nutr Metab* **66 Suppl 2**, 7-12 (2015). https://doi.org:10.1159/000375143
3   Chong, B. *et al.* Trends and predictions of malnutrition and obesity in 204 countries and
territories: an analysis of the Global Burden of Disease Study 2019. *eClinicalMedicine*
57, 101850 (2023). https://doi.org:https://doi.org/10.1016/j.eclinm.2023.101850

4   Powell-Wiley, T. M. *et al.* Obesity and Cardiovascular Disease: A Scientific Statement
From the American Heart Association. *Circulation* **143**, e984-e1010 (2021).
https://doi.org:10.1161/CIR.0000000000000973
5   Avgerinos, K. I., Spyrou, N., Mantzoros, C. S. & Dalamaga, M. Obesity and cancer risk:
Emerging biological mechanisms and perspectives. *Metabolism* **92**, 121-135 (2019).
https://doi.org:10.1016/j.metabol.2018.11.001
6   Sarwar, R., Pierce, N. & Koppe, S. Obesity and nonalcoholic fatty liver disease: current
perspectives. *Diabetes Metab Syndr Obes* **11**, 533-542 (2018).
https://doi.org:10.2147/dmso.S146339
7   Picone, P., Di Carlo, M. & Nuzzo, D. Obesity and Alzheimer's disease: Molecular bases.
Eur J Neurosci **52**, 3944-3950 (2020). https://doi.org:10.1111/ejn.14758
8   Shaikh, S. R., Beck, M. A., Alwarawrah, Y. & MacIver, N. J. Emerging mechanisms of
obesity-associated immune dysfunction. *Nature Reviews Endocrinology* **20**, 136-148
(2024). https://doi.org:10.1038/s41574-023-00932-2
9   Kawai, T., Autieri, M. V. & Scalia, R. Adipose tissue inflammation and metabolic
dysfunction in obesity. *Am J Physiol Cell Physiol* **320**, C375-c391 (2021).
https://doi.org:10.1152/ajpcell.00379.2020
10 Kershaw, E. E. & Flier, J. S. Adipose tissue as an endocrine organ. *J Clin Endocrinol*
Metab **89**, 2548-2556 (2004). https://doi.org:10.1210/jc.2004-0395
11 Booth, A., Magnuson, A., Fouts, J. & Foster, M. T. Adipose tissue: an endocrine organ
playing a role in metabolic regulation. *Horm Mol Biol Clin Investig* **26**, 25-42 (2016).
https://doi.org:10.1515/hmbci-2015-0073
12 Ouchi, N., Parker, J. L., Lugus, J. J. & Walsh, K. Adipokines in inflammation and
metabolic disease. *Nature Reviews Immunology* **11**, 85-97 (2011).
https://doi.org:10.1038/nri2921
13 Longo, M. *et al.* Adipose Tissue Dysfunction as Determinant of Obesity-Associated
Metabolic Complications. *Int J Mol Sci* **20** (2019). https://doi.org:10.3390/ijms20092358
14 Kalluri, R. & LeBleu, V. S. The biology, function, and biomedical applications of
exosomes. *Science* **367** (2020). https://doi.org:10.1126/science.aau6977
15 Hade, M. D., Suire, C. N. & Suo, Z. Mesenchymal Stem Cell-Derived Exosomes:
Applications in Regenerative Medicine. *Cells* **10**, 1959 (2021).
16 Tkach, M. & Thery, C. Communication by Extracellular Vesicles: Where We Are and
Where We Need to Go. *Cell* **164**, 1226-1232 (2016).
https://doi.org:10.1016/j.cell.2016.01.043
17 Connolly, K. D., Rees, D. A. & James, P. E. Role of adipocyte-derived extracellular
vesicles in vascular inflammation. *Free Radic Biol Med* **172**, 58-64 (2021).
https://doi.org:10.1016/j.freeradbiomed.2021.04.031
18 Isaac, R., Reis, F. C. G., Ying, W. & Olefsky, J. M. Exosomes as mediators of
intercellular crosstalk in metabolism. *Cell Metab* **33**, 1744-1762 (2021).
https://doi.org:10.1016/j.cmet.2021.08.006
19 Zhou, C., Huang, Y.-Q., Da, M.-X., Jin, W.-L. & Zhou, F.-H. Adipocyte-derived
extracellular vesicles: bridging the communications between obesity and tumor
microenvironment. *Discover Oncology* **14**, 92 (2023). https://doi.org:10.1007/s12672-
023-00704-4
20 Kumar, V., Kiran, S., Kumar, S. & Singh, U. P. Extracellular vesicles in obesity and its
associated inflammation. *Int Rev Immunol* **41**, 30-44 (2022).
https://doi.org:10.1080/08830185.2021.1964497
21 Crewe, C. The challenges of interrogating adipose tissue extracellular vesicle functions
in physiology. *Commun Biol* **5**, 581 (2022). https://doi.org:10.1038/s42003-022-03511-9

Crewe, C. *et al.* An Endothelial-to-Adipocyte Extracellular Vesicle Axis Governed by Metabolic State. *Cell* **175**, 695-708 e613 (2018). https://doi.org:10.1016/j.cell.2018.09.005

Akbar, N., Pinnick, K. E., Paget, D. & Choudhury, R. P. Isolation and Characterization of Human Adipocyte-Derived Extracellular Vesicles using Filtration and Ultracentrifugation. *J Vis Exp* (2021). https://doi.org:10.3791/61979

Borgeson, E., Boucher, J. & Hagberg, C. E. Of mice and men: Pinpointing species differences in adipose tissue biology. *Front Cell Dev Biol* **10**, 1003118 (2022). https://doi.org:10.3389/fcell.2022.1003118

Palacio, P. L. *et al.* Novel multiparametric bulk and single EV pipeline for adipose cell-specific biomarker discovery in paired human biospecimens. *bioRxiv*, 2024.2004.2018.590172 (2024). https://doi.org:10.1101/2024.04.18.590172

Nguyen, K. T. *et al.* Integrated Techniques for Extracellular Particle Separation and Single-Particle Multiparametric Characterization to Track Cancer Biomarkers from Tissue to Biofluids. *bioRxiv*, 2025.2001.2009.632270 (2025). https://doi.org:10.1101/2025.01.09.632270

Rima, X. Y. *et al.* Microfluidic harvesting of breast cancer tumor spheroid-derived extracellular vesicles from immobilized microgels for single-vesicle analysis. *Lab Chip* **22**, 2502-2518 (2022). https://doi.org:10.1039/d1lc01053k

Nguyen, L. T. H. *et al.* An immunogold single extracellular vesicular RNA and protein ((Au) SERP) biochip to predict responses to immunotherapy in non-small cell lung cancer patients. *J Extracell Vesicles* **11**, e12258 (2022). https://doi.org:10.1002/jev2.12258

Vella, L. J. *et al.* A rigorous method to enrich for exosomes from brain tissue. *J Extracell Vesicles* **6**, 1348885 (2017). https://doi.org:10.1080/20013078.2017.1348885

Tong, Q. *et al.* Function of GATA Transcription Factors in Preadipocyte-Adipocyte Transition. *Science* **290**, 134-138 (2000). https://doi.org:doi:10.1126/science.290.5489.134

da Silva, C., Durandt, C., Kallmeyer, K., Ambele, M. A. & Pepper, M. S. The Role of Pref-1 during Adipogenic Differentiation: An Overview of Suggested Mechanisms. *Int J Mol Sci* **21** (2020). https://doi.org:10.3390/ijms21114104

Tseng, Y.-H. *et al.* Prediction of preadipocyte differentiation by gene expression reveals role of insulin receptor substrates and necdin. *Nature Cell Biology* **7**, 601-611 (2005). https://doi.org:10.1038/ncb1259

Rim, J. S. *et al.* Direct Conversion of Human Preadipocytes into Hematopoietic, Neuronal, and Pancreatic α Cells by Oct4 and Klf4 Overexpression. *BioResearch Open Access* **4**, 389-397 (2015). https://doi.org:10.1089/biores.2015.0028

Dang, T. N. *et al.* Expression of the preadipocyte marker ZFP423 is dysregulated between well-differentiated and dedifferentiated liposarcoma. *BMC Cancer* **22**, 300 (2022). https://doi.org:10.1186/s12885-022-09379-6

Noguchi, Y., Murakami, M., Murata, M. & Kano, F. Microscopic image-based classification of adipocyte differentiation by machine learning. *Histochemistry and Cell Biology* **159**, 313-327 (2023). https://doi.org:10.1007/s00418-022-02168-z

Audano, M. *et al.* Regulatory mechanisms of the early phase of white adipocyte differentiation: an overview. *Cellular and Molecular Life Sciences* **79**, 139 (2022). https://doi.org:10.1007/s00018-022-04169-6

Tong, Q. *et al.* Function of GATA transcription factors in preadipocyte-adipocyte transition. *Science* **290**, 134-138 (2000). https://doi.org:10.1126/science.290.5489.134

Al-Mansoori, L. *et al.* Suppression of GATA-3 increases adipogenesis, reduces inflammation and improves insulin sensitivity in 3T3L-1 preadipocytes. *Cellular Signalling* **75**, 109735 (2020). https://doi.org:https://doi.org/10.1016/j.cellsig.2020.109735

Fu, Y., Luo, N., Klein, R. L. & Garvey, W. T. Adiponectin promotes adipocyte differentiation, insulin sensitivity, and lipid accumulation. *Journal of Lipid Research* **46**, 1369-1379 (2005). https://doi.org:https://doi.org/10.1194/jlr.M400373-JLR200

Itabe, H., Yamaguchi, T., Nimura, S. & Sasabe, N. Perilipins: a diversity of intracellular lipid droplet proteins. *Lipids in Health and Disease* **16**, 83 (2017). https://doi.org:10.1186/s12944-017-0473-y

Sztalryd, C. & Brasaemle, D. L. The perilipin family of lipid droplet proteins: Gatekeepers of intracellular lipolysis. *Biochim Biophys Acta Mol Cell Biol Lipids* **1862**, 1221-1232 (2017). https://doi.org:10.1016/j.bbalip.2017.07.009

Daya, S., Loughlin, A. J. & MacQueen, H. A. Culture and differentiation of preadipocytes in two-dimensional and three-dimensional in vitro systems. *Differentiation* **75**, 360-370 (2007). https://doi.org:https://doi.org/10.1111/j.1432-0436.2006.00146.x

Nielsen, T. S., Jessen, N., Jørgensen, J. O. L., Møller, N. & Lund, S. Dissecting adipose tissue lipolysis: molecular regulation and implications for metabolic disease. *Journal of Molecular Endocrinology* **52**, R199-R222 (2014). https://doi.org:10.1530/jme-13-0277

Yang, A. & Mottillo, E. P. Adipocyte lipolysis: from molecular mechanisms of regulation to disease and therapeutics. *Biochem J* **477**, 985-1008 (2020). https://doi.org:10.1042/bcj20190468

Cho, C. H., Patel, S. & Rajbhandari, P. Adipose tissue lipid metabolism: lipolysis. *Current Opinion in Genetics & Development* **83**, 102114 (2023). https://doi.org:https://doi.org/10.1016/j.gde.2023.102114

Crewe, C. & Scherer, P. E. Intercellular and interorgan crosstalk through adipocyte extracellular vesicles. *Rev Endocr Metab Disord* **23**, 61-69 (2022). https://doi.org:10.1007/s11154-020-09625-x

Kranendonk, M. E. *et al.* Human adipocyte extracellular vesicles in reciprocal signaling between adipocytes and macrophages. *Obesity (Silver Spring)* **22**, 1296-1308 (2014). https://doi.org:10.1002/oby.20679

Lazar, I. *et al.* Adipocyte Exosomes Promote Melanoma Aggressiveness through Fatty Acid Oxidation: A Novel Mechanism Linking Obesity and Cancer. *Cancer Research* **76**, 4051-4057 (2016). https://doi.org:10.1158/0008-5472.CAN-16-0651

Clement, E. *et al.* Adipocyte extracellular vesicles carry enzymes and fatty acids that stimulate mitochondrial metabolism and remodeling in tumor cells. *Embo j* **39**, e102525 (2020). https://doi.org:10.15252/embj.2019102525

Matilainen, J. *et al.* Increased secretion of adipocyte-derived extracellular vesicles is associated with adipose tissue inflammation and the mobilization of excess lipid in human obesity. *J Transl Med* **22**, 623 (2024). https://doi.org:10.1186/s12967-024-05249-w

Kulaj, K. *et al.* Adipocyte-derived extracellular vesicles increase insulin secretion through transport of insulinotropic protein cargo. *Nat Commun* **14**, 709 (2023). https://doi.org:10.1038/s41467-023-36148-1

Börgeson, E., Boucher, J. & Hagberg, C. E. Of mice and men: Pinpointing species differences in adipose tissue biology. *Front Cell Dev Biol* **10**, 1003118 (2022). https://doi.org:10.3389/fcell.2022.1003118

Muniyappa, R. *et al.* Comparison between surrogate indexes of insulin sensitivity/resistance and hyperinsulinemic euglycemic clamp estimates in rats. *Am J Physiol Endocrinol Metab* **297**, E1023-1029 (2009). https://doi.org:10.1152/ajpendo.00397.2009

Gómez-Serrano, M., Ponath, V., Preußer, C. & Pogge von Strandmann, E. Beyond the Extracellular Vesicles: Technical Hurdles, Achieved Goals and Current Challenges When Working on Adipose Cells. *Int J Mol Sci* **22** (2021). https://doi.org:10.3390/ijms22073362

55   Kranendonk, M. E. G. *et al.* Human adipocyte extracellular vesicles in reciprocal
signaling between adipocytes and macrophages. *Obesity* **22**, 1296-1308 (2014).
https://doi.org:https://doi.org/10.1002/oby.20679
56   Sarango, G. *et al.* The Autophagy Receptor TAX1BP1 (T6BP) improves antigen
presentation by MHC-II molecules. *EMBO Rep* **23**, e55470 (2022).
https://doi.org:10.15252/embr.202255470
57   Chatterjee, E. *et al.* The extracellular vesicle transcriptome provides tissue-specific
functional genomic annotation relevant to disease susceptibility in obesity. *medRxiv*
(2024). https://doi.org:10.1101/2024.11.18.24317277
58   Gómez-Serrano, M., Ponath, V., Preußer, C. & Pogge von Strandmann, E. Beyond the
Extracellular Vesicles: Technical Hurdles, Achieved Goals and Current Challenges
When Working on Adipose Cells. *International Journal of Molecular Sciences* **22**, 3362
(2021).
59   Sethi, J. K. & Vidal-Puig, A. J. Thematic review series: Adipocyte Biology. Adipose tissue
function and plasticity orchestrate nutritional adaptation. *Journal of Lipid Research* **48**,
1253-1262 (2007). https://doi.org:https://doi.org/10.1194/jlr.R700005-JLR200
60   Connolly, K. D. *et al.* Evidence for Adipocyte-Derived Extracellular Vesicles in the
Human Circulation. *Endocrinology* **159**, 3259-3267 (2018).
https://doi.org:10.1210/en.2018-00266
61   Lötvall, J. *et al.* Minimal experimental requirements for definition of extracellular vesicles
and their functions: a position statement from the International Society for Extracellular
Vesicles. *J Extracell Vesicles* **3**, 26913 (2014). https://doi.org:10.3402/jev.v3.26913
62   Adnan, S.-s., Benjamin, G. T. & David, R. W. Advances in extracellular vesicle isolation
methods: a path towards cell-type specific EV isolation. *Extracellular Vesicles and
Circulating Nucleic Acids* **4**, 447-460 (2023). https://doi.org:10.20517/evcna.2023.14
63   Beck, B. R., Shin, B., Choi, Y., Park, S. & Kang, K. Predicting commercially available
antiviral drugs that may act on the novel coronavirus (SARS-CoV-2) through a drug-
target interaction deep learning model. *Comput Struct Biotechnol J* **18**, 784-790 (2020).
https://doi.org:10.1016/j.csbj.2020.03.025
64   Thomou, T. *et al.* Adipose-derived circulating miRNAs regulate gene expression in other
tissues. *Nature* **542**, 450-455 (2017). https://doi.org:10.1038/nature21365
65   Li, Y. *et al.* EV-origin: Enumerating the tissue-cellular origin of circulating extracellular
vesicles using exLR profile. *Computational and Structural Biotechnology Journal* **18**,
2851-2859 (2020). https://doi.org:https://doi.org/10.1016/j.csbj.2020.10.002
66   Connolly, K. D. *et al.* Characterisation of adipocyte-derived extracellular vesicles
released pre- and post-adipogenesis. *J Extracell Vesicles* **4**, 29159 (2015).
https://doi.org:10.3402/jev.v4.29159
67   Liangsupree, T., Multia, E. & Riekkola, M.-L. Modern isolation and separation techniques
for extracellular vesicles. *Journal of Chromatography A* **1636**, 461773 (2021).
https://doi.org:https://doi.org/10.1016/j.chroma.2020.461773
68   Ter-Ovanesyan, D. *et al.* Framework for rapid comparison of extracellular vesicle
isolation methods. *Elife* **10** (2021). https://doi.org:10.7554/eLife.70725
69   Di, K. *et al.* Highly efficient and automated isolation technology for extracellular vesicles
microRNA. *Front Bioeng Biotechnol* **10**, 948757 (2022).
https://doi.org:10.3389/fbioe.2022.948757
70   Ferguson, S. *et al.* Single-EV analysis (sEVA) of mutated proteins allows detection of
stage 1 pancreatic cancer. *Sci Adv* **8**, eabm3453 (2022).
https://doi.org:10.1126/sciadv.abm3453
71   Li, Z. *et al.* Colocalization of protein and microRNA markers reveals unique extracellular
vesicle subpopulations for early cancer detection. *Sci Adv* **10**, eadh8689 (2024).
https://doi.org:10.1126/sciadv.adh8689
719