Single-nucleus analysis reveals oxidative stress in Down syndrome basal forebrain neurons at birth

Nicole R. West[1,2], Kalpana Hanthanan Arachchilage[1], Sara Knaack[1], Shawn MacGregor[1,] Masoumeh Hosseini[1], Ryan D. Risgaard[1,3], Pubudu Kumarage[1], Jose L. Martinez[1], Su-Chun Zhang[1,4], Daifeng Wang[1,5], Andre M.M. Sousa[1,6], Anita Bhattacharyya[1,7]*

[1]Waisman Center, University of Wisconsin-Madison, 1500 Highland Ave., Madison, WI 53705

[2]Cellular and Molecular Biology Graduate Program, University of Wisconsin-Madison, 1525 Linden Dr., Madison, WI 53706

[3]Medical Scientist Training Program, University of Wisconsin-Madison School of Medicine and Public Health, 750 Highland Ave., Madison, WI 53705

[4]Director and Professor, Center for Neurologic Diseases, Sanford Burnham Prebys Medical Discovery Institute, 10901 N Torrey Pines Rd La Jolla, CA 92037

[5]Department of Biostatistics and Medical Informatics, University of Wisconsin-Madison School of Medicine and Public Health, 610 Walnut St., Madison, WI 53726

[6]Department of Neuroscience, University of Wisconsin-Madison School of Medicine and Public Health, 1111 Highland Ave., Madison, WI 53705

[7]Department of Cell and Regenerative Biology, University of Wisconsin-Madison School of Medicine and Public Health, 1111 Highland Ave., Madison, WI 53705

*Corresponding author: bhattacharyy@waisman.wisc.edu,

Waisman Center, University of Wisconsin-Madison, 1500 Highland Ave., Madison, WI 53705

Abstract:

INTRODUCTION: Basal forebrain cholinergic neurons (BFCNs) are integral to learning, attention, and memory, and are prone to degeneration in Down syndrome (DS), Alzheimer's disease, and other neurodegenerative diseases. However, the mechanisms that lead to the degeneration of these neurons are not known.

METHODS: Single-nucleus gene expression and ATAC sequencing were performed on postmortem human basal forebrain from unaffected control and DS tissue samples at 0-2 years of age (n=4 each).

RESULTS: Sequencing analysis of postmortem human basal forebrain identifies gene expression differences in DS early in life. Genes encoding proteins associated with energy metabolism pathways, specifically oxidative phosphorylation and glycolysis, and genes encoding antioxidant enzymes are upregulated in DS BFCNs.

DISCUSSION: Multiomic analyses reveal that energy metabolism may be disrupted in DS BFCNs by birth. Increased oxidative phosphorylation and the accumulation of reactive oxygen species byproducts may be early contributors to DS BFCN neurodegeneration.

1. Background

Cholinergic projection neurons of the basal forebrain (BFCNs), are the primary cholinergic input to the cortex, hippocampus, and amygdala, regulating cognitive functions including learning, attention, and memory[1]. The cholinergic hypothesis, proposed nearly 50 years ago, posits that the dysfunction or loss of cholinergic neurons is an early driver of cognitive decline associated with age and Alzheimer's disease (AD)[2-4]. BFCNs are some of the first neurons to degenerate in the progression of AD[5-7]. Tau tangles accumulate in the basal forebrain in AD before the entorhinal cortex[5, 6, 8]. Studies suggest that BFCNs seed the cortex with pathology through the trans-synaptic spread of misfolded Tau[6, 7]. Consequently, BFCNs have been a target of therapeutics to reduce degeneration, slow the spread of AD pathology, and ultimately to slow cognitive decline in AD[9, 10]. Although the link between BFCN degeneration and memory decline is well understood, early molecular events that occur in BFCNs that increase susceptibility to degeneration later are unknown.

BFCN dysfunction and degeneration also occur in Down syndrome (DS, trisomy 21, T21)[11-15] and other neurodegenerative diseases, including Parkinson's disease (PD)[12, 16-20] and Dementia with Lewy bodies (DLB)[21-24]. Nearly all individuals with DS develop AD (DS-AD), making DS the leading genetic cause of AD[25]. The progression of DS-AD pathology and the onset of dementia occurs in a consistent and predictable manner[26, 27], enabling examination of successive stages of disease progression in DS. BFCN degeneration in the anteromedial and posterior basal forebrain in individuals with DS begins about 30 years prior to the median onset of prodromal AD[14]. Post-mortem studies validate fewer neurons are present in the basal forebrain in

71   DS relative to unaffected controls(11). Further, DS is a neurodevelopmental disorder, and

72   so we hypothesized that early molecular changes in BFCNs increase their susceptibility

73   to degeneration in DS. Identifying these molecular changes may provide insight into

74   drivers of BFCN degeneration in the progression of DS-AD.

75   We performed unbiased single-nucleus gene expression and ATAC multiomic

analysis of early human postnatal DS and unaffected control basal forebrain (BF) to

identify molecular changes in DS that precede BFCN dysfunction or degeneration(14, 28-

30). We identified cell type-specific differential gene expression across all eight cell types

in the DS basal forebrain. The results suggest that energy metabolism, specifically the

upregulation of glycolysis and oxidative phosphorylation (OXPHOS) genes, is

dysregulated in DS BFCNs. Consequently, DS BFCNs have increased gene expression

of antioxidant enzymes, possibly to regulate the reactive oxygen species (ROS) that

accumulate as a byproduct of OXPHOS. The increased ROS over a sustained time may

increase the vulnerability of DS BFCNs, causing their early degeneration in DS. These

results identify potential novel targets and a timeframe for therapeutic intervention to

delay BFCN dysfunction and mitigate disease progression.

2. Methods

2.1 Tissue Samples

90   Human basal forebrain post-mortem samples from Down syndrome and

unaffected control individuals were obtained from the University of Maryland Brain and

Tissue Bank, as part of the National Institutes of Health NeuroBioBank. Acquisition of the

de-identified samples was approved by the Health Sciences Institutional Review Board

at the University of Wisconsin-Madison (Protocol #2016-0979) and certified exempt from

IRB oversight. Sample information is provided in **Figure 1A** and **Supplement Table 1**.

2.2 Nuclei Isolation

Frozen post-mortem human basal forebrain (BF) tissue sections were pulverized

into a powder in liquid nitrogen over dry ice, using a mortar and pestle (Fisherscientific

#FB961A and #FB961K). 25-35 mg of pulverized BF tissue was used for nuclei isolation.

All buffers were prepared fresh and maintained on ice. 3 mL of ice-cold buffer B (Iodixanol

buffer) [50% (v/v) Optiprep (Iodixanol) solution (Sigma # D1556); 25 mM KCl (Sigma

#60142); 5 mM MgCl2 (Sigma #M1028); 20 mM Tris-HCl (pH 7.5) (Invitrogen #15567-

027); 1% cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail (Roche

#11836170001); 1% BSA (GeminiBio #700-100p); RNase inhibitor (80U/mL) (Roche

#03335402001); 1 mM DTT (Sigma #43186)] was transferred to a 15 mL tube.

1 mL buffer A (lysis buffer) [250 mM Sucrose (Sigma #S0389); 25 mM KCl2 (Sigma

#21115); 5 mM MgCl2 (Sigma #63052); 20 mM Tris-HCl (pH 7.5) (Invitrogen #15567-

027); 1% cOmplete™, Mini, EDTA-free Protease Inhibitor Cocktail (Roche

#11836170001); RNase inhibitor (40U/mL) (Roche #03335402001); 1 mM DTT (Sigma

#43186); 0.1 % (v/v) IGEPAL CA-630 (Sigma #I8896)] was added to the pulverized BF

tissue in a tube. 1 mL of ice-cold lysis buffer was added to the Dounce tissue grinder (15

mL volume, Wheaton #357544; autoclaved, RNase free, ice-cold). 1 mL of lysis buffer

was added to the pulverized tissue tube to rinse and collect all tissue. The suspension

was transferred to Dounce tissue grinder and homogenized with loose and tight pestles,

30 cycles each, with constant pressure and without the introduction of air. The solution

was transferred to the 15 mL tube containing buffer B and mixed by inverting the tube 10

times. The homogenate was filtered through a 40-um cell strainer (Falcon #352340) which

was pre-washed with lysis buffer. Samples were centrifuged at 1000 g for 30 min at 4°C

(Eppendorf #5910 Ri). Following centrifugation, the supernatant was carefully and

completely removed and the pellet was resuspended in 1 mL of wash buffer [10 mM NaCl

(Sigma #60142); 3 mM MgCl2 (Sigma #M1028); 10 mM Tris-HCl (pH 7.5) (Invitrogen

#15567-027); 1% BSA (GeminiBio #700-100p); RNase inhibitor (1000U/mL) (Roche

#03335402001); 1mM DTT (Sigma #43186)]. The homogenate was filtered through a 40-

um cell strainer (Falcon #352340) pre-washed with wash buffer to eliminate large clumps

and cell debris. Samples were then centrifuged at 500 g for 5 min at 4°C.

Supernatants were carefully and completely removed. Pellets were gently

dissolved by adding 200 mL and 300 mL of buffer C [10 mM NaCl (Sigma #60142); 3 mM

MgCl2 (Sigma #M1028); 10 mM Tris-HCl (pH 7.5) (Invitrogen #15567-027); 0.01%

Tween-20 (Bio-Rad #1662404); 0.001% Digitonin (Thermo Fisher #BN2006); 0.01% (v/v)

IGEPAL CA630 (Sigma#I8896); 1% cOmplete™, Mini, EDTA-free Protease Inhibitor

Cocktail (Roche#11836170001); 1% BSA (GeminiBio #700-100p); RNase inhibitor

(1000U/mL) (Roche #03335402001); 1mM DTT (Sigma #43186)].The solution was

incubated on ice for 5 minutes. After incubation 500 mL of buffer D [10 mM NaCl (Sigma

#60142); 3 mM MgCl2 (Sigma #M1028); 10 mM Tris-HCl (pH 7.5) (Invitrogen #15567-

027); 0.1% Tween-20 (Bio-Rad #1662404); 1% BSA (GeminiBio #700-100p); RNase

inhibitor (1000U/mL) (Roche #03335402001); 1mM DTT (Sigma #43186)) was added to

the solution. After resuspension, nuclei quality was assessed at 40X magnification and

were manually counted using a hemocytometer. The sample was centrifuged at 500 g for

5 min at 4°C. The pellet was resuspended in buffer E [1X nuclei buffer (10x Genomics

#2000207); RNase inhibitor (1000U/mL) (Roche #03335402001); 1mM DTT (Sigma

#43186)] at a final concentration of 5 million nuclei/mL.

2.3 snMultiomic Library Generation and Sequencing

For each sample, 10,000 nuclei were targeted. Nuclei suspension was first

incubated in a transposition mix. Thereafter, along with the oligo-coated gel beads and

partitioning oil (10x Genomics #PN-2000190), the single nuclei master mixture containing

tagmented single nuclei suspension was transferred onto a Next GEM Chip J (10x

Genomic #PN-2000264), and the chip was loaded to the Chromium Controller for GEM

generation. After post GEM incubation clean up, preamplification of samples was

performed and ATAC libraries were generated utilizing the Single Index Kit N Set A (10x

Genomics #PN-1000212). The snRNA-seq libraries were generated using the Library

Construction Kit (10x Genomics #PN-1000190) and Dual Index Kit TT Set A (10x

Genomics #PN-1000215), following the manufacturer's recommended protocol. At each

step, the concentration and quality of cDNA, ATAC library and GEX libraries were

assessed by 4200 TapeStation (Agilent). Sequencing was carried out with Illumina

NovaSeq X Plus for a targeted depth of 50,000 raw reads per nucleus.

2.4 snMultiomic Alignment

The 10x multiomic data was processed according to BICAN default methods. The

snRNA-seq data were aligned with STAR v2.7.11a(31) and aggregated into count

matrices with STARSolo. The alignments were performed locally using defaults from the

WARP Multiome (v5.9.0) pipeline(32). Initial quality control checks included assessment

of the percentages of uniquely- and multiply-mapped reads, and statistics of corrected

barcodes and UMI. The snRNA-seq data alignment was done utilizing the

GeneFull_Ex50pAS argument for STAR to ensure inclusion of alignments that overlap

exonic ends within genic regions, in consideration of nuclear RNA processing biology.

The snATAC-seq data was likewise processed according to the WARP Multiome

(v5.9.0) pipeline. This includes ibarcode correction with the fastqProcessing processing

tool from WarpTools (32), and subsequent trimming with Cutadapt (v4.4)(33). Alignment

was performed with BWA-MEM2(34). Finally, generation of fragment files and initial QC

was performed with SnapATAC (v2.3.1)(35). This was facilitated with an updated version

of the snATAC component of the WARP Multiome (v5.9.0) pipeline to facilitate this

workflow on Amazon AWS EC2 instances. Specifically, docker images for Cutadapt (v4.4)

and for the BWA alignment were prepared to leverage AVX2 processing for improved

performance. In this stage of the analysis all samples for either unaffected control or DS

donors were treated identically.

2.5 snRNA-seq Processing and QC

Data processing and downstream analysis were conducted in R (4.4.0). Packages

and versions are listed in **Supplement Table 2**. Since DS samples carry an extra

chromosome, we modified the standard preprocessing pipeline to incorporate genotype-

specific differences in the transcript abundance. For instance, it is standard practice to

impose upper and lower thresholds on the number of UMIs (nCount_RNA). The upper

threshold is primarily used to remove potential homotypic doublets. Applying a single

upper UMI threshold would have either led to a loss of many high-quality DS cells or the

retention of potential doublets in control samples. To address this, we set separate upper

UMI thresholds for control and DS samples based on their respective distributions

(**Supplement 1A**). Identifying highly variable genes separately per genotype enhances

the transcriptomic signal related to batch effects and cell types, ultimately improving

clustering accuracy.

The snRNA-seq datasets were preprocessed and analyzed using the Seurat

(v5.1.0)(36, 37) R package. mRNA contamination caused by cell-free ambient RNA in the

gene expression data was corrected using the SoupX (v1.6.2)(38) package. Low-quality

nuclei were then identified and removed based on stringent quality control thresholds:

fewer than 200 expressed genes, ribosomal gene content exceeding 40%, mitochondrial

gene content exceeding 5%, and a UMI count lower than 800 or higher than $Q_3 + 3(Q_3 -$

$Q_1)$, where Q_1 and Q_3 are the lower and upper quartiles(39, 40) (**Supplement 1A-B**). The

upper UMI thresholds used were 29,284 in control and 34,160 in DS (**Supplement 1A**).

After quality control, the datasets were subjected to doublet removal. Given the

uncertainties inherent in doublet detection methods, an ensemble approach was

employed, incorporating three techniques: DoubletFinder (v2.0.4)(41), Scrublet(42), and

scDBIFinder v(1.18.0)(43). A cell identified as a doublet by at least two of the three

methods was classified as a doublet and removed from the dataset.

SoupX-corrected UMI counts were log-normalized using Seurat. The top 3,000

highly variable genes were identified using the default variance-stabilizing process. Gene

expression data for these highly variable genes were then scaled, and dimensionality

reduction was performed using principal component analysis (PCA). Batch effects were

subsequently removed using Harmony (v1.2.3)(44) implemented within Seurat.

Clustering and cell type annotations were carried out in two steps. In the first step,

Louvain clustering(37) was applied using the first 30 Harmony components with a cluster

resolution of 0.5. Cell classes (e.g., neurons and non-neuronal cells) were assigned to

the clusters based on the relative expression of a curated list of marker genes. In the

second step, cells from each cell class were isolated and sub-clustered to identify more

granular clusters representing specific cell subclasses and subtypes based on a list of

known marker genes (**Supplement Table 3**). This led to the identification of six non-

neuronal subtypes: astrocytes, endothelial cells, microglia, oligodendrocytes,

oligodendrocyte precursor cells (OPCs), and glial precursor cells (GPCs). Inhibitory

neurons were the only subclass of neurons identified, and BFCNs were identified as a

subtype within the inhibitory neuron population. Additional 'low quality cells' that could not

confidently be identified with known marker genes were removed (**Supplement 1C**). After

all the quality control steps and cell type annotations, the dataset retained a total of 37,467

cells and all donors were represented (**Supplement 1D-E**). There is a positive correlation

(0.92) between the number of unique molecular identifiers (RNA Count) and the number

of genes (RNA Features) (**Supplement 1F**).

Following separate QC and cell annotations, control and DS data were merged

and batch corrected using Harmony integration in Seurat. The FindMarkers() function in

Seurat was used with the MAST model and sample age as a latent variable to calculate

differentially expressed genes (DEGs). For this analysis, we required that at least 25% of

cells in a cluster expressed the gene for it to be considered a DEG. DEGs for each cell

type are listed in **Supplement Table 4**. Hsa21 DEGs for each cell type are listed in

Supplement Table 5. DEGs were mapped to chromosomes using the online tool,

MG2C(45). Gene set enrichment analysis (GSEA) was performed using the FGSEA

method(46). The scProgram and clusterProfiler(47) packages were used for KEGG

pathway analysis.

2.6 snATAC-seq Processing and QC

238   The snATAC fragment data files were merged into a single master fragment file by

1) mapping the snATAC barcodes in the prepared per-sample fragment files to the

corresponding snRNA seq barcodes (for direct integration), and 2) subsequently

coordinate-sorting, compressing and tabulating these data for analysis in Seurat

(v5.1.0)(36, 37, 48) and Signac (v1.14)(49). The multiomic integration analysis proceeded

by assessing and filtering the snATAC data for those barcodes meeting the snRNA-seq

quality control criteria (**Supplement G-H**).

245   Barcodes meeting the following criteria were selected for the integrated analysis:

246   1) number of fragments between 1k and 100k, 2) A TSS enrichment score >=2 and 3)

247   nucleosome fraction score < 4 (**Supplement 1I**). These criteria resulted in 31,411 total

248   barcodes for multiomic analysis (**Supplement 1J**).

250   **2.7 Peak Calling**

251   Peak calling was facilitated with the Signac CallPeaks() function, using MACS2

(v2.2.9.1)(50, 51). This was done for the integrated barcode data set partitioned by both

annotated cell type (per the snRNA-seq analysis) and disease condition of donors

(unaffected control and DS, respectively). For each cell type and condition, peaks were

called, and a merged set of peak regions were compiled as implemented in CallPeaks().

A chromatin assay object representing the merged regions was aggregated by

barcode and prepared in a merged Seurat object. The chief quality control measure

assessed for the peak calling was the fraction of reads in peaks (FRiP) per barcode

(**Supplement 1K**).

2.8 Peak-to-Gene Links

We estimated peak-to-gene links (i.e., nearby peaks correlated with gene expression of

a given gene) for a few selected cell types to examine characteristics of regulatory

elements across control and DS samples. These calculations were done separately

across cell type and genotype. Peak-to-gene links within 500 kb of the corresponding

transcription start sites (TSS) were obtained using LinkPeaks() function. The loci of genes

of interest were then visualized using the coveragePlot() function in Signac, highlighting

the differences in the regulatory landscape across control and DS samples.

2.9 Protein Carbonyl Enzyme-linked Immunosorbent Assay

15mg of frozen pulverized tissue was homogenized in 100uL of 1x PBS with 1% Protease

and Phosphatase Inhibitor Cocktail (Sigma-Aldrich #PPC1010). Protein concentration

was quantified with a *DC* Protein Assay following the manufacturer's protocol (Bio-Rad

#5000116). Protein oxidation was measured with a Protein Carbonyl ELISA Kit following

the manufacturer's protocol (abcam #ab238536). Samples and standards were run in

duplicate and absorbance was read on a Molecular Devices VersaMax plate reader.

2. Malondialdehyde Lipid Peroxidation Enzyme-linked Immunosorbent Assay

15mg of frozen pulverized tissue was homogenized in 100uL of 1x PBS with 1% Protease and Phosphatase Inhibitor Cocktail (Sigma-Aldrich #PPC1010). Protein concentration was quantified with a *DC* Protein Assay following the manufacturer's protocol (Bio-Rad #5000116). Lipid peroxidation was measured with an MDA Assay Kit (competitive ELISA) (abcam #ab238537). Protein samples were diluted to 1mg/mL and were assayed according to the manufacture's protocol. Samples were assayed twice using fresh preparations of protein extracts for each assay run. Samples and standards were run in duplicate and absorbance was read on a Molecular Devices VersaMax plate reader. The average concentration from both assay runs were averaged.

3. Results

3.1 Overview of Human Basal Forebrain Tissue

Our data represent the first single-nucleus sequencing of the human basal forebrain. Single-nucleus multiomic analysis of gene expression and chromatin accessibility (**Figure 1B**) was performed on four unaffected control and four DS basal forebrain samples matched for age, sex, and PMI (**Figure 1A; Supplement Table 1**). After quality control (**Supplement 1A-C**), 34,467 cells were used for downstream analysis (**Figure 1C**). Cells from all donors were represented (**Supplement 1D-E**). The cell types identified in the basal forebrain were astrocytes, BFCNs, endothelial cells, inhibitory neurons, glial progenitor cells (GPCs), microglia, oligodendrocytes, and oligodendrocyte precursor cells (OPCs) (**Figure 1C**), and all cell types are present in both control and DS BF tissue samples.

Cell annotations of control and DS samples were performed separately. Initially,

cells were annotated by major cell class- excitatory neurons (ExN), inhibitory neurons

(InN), and non-neuronal cells (NNC). Only InNs and NNCs were identified in the initial

classification (**Supplement 2A**). NNCs were sub-clustered to subclasses annotated as

glial or endothelial cells (**Supplement 2B**). Astrocytes, GPCs, microglia,

oligodendrocytes, and OPCs were annotated in the glial subclass (**Figure 1C;**

Supplement 2C). BFCNs were identified as a subtype within inhibitory neurons (**Figure**

1C; Supplement 2C). BFCNs, which are capable of co-transmitting ACh and GABA(52,

53), were initially annotated as InNs based on their expression of *GAD1, GABBR1,* and

SLC6A1 (**Supplement 2A-B**).

After separate cluster annotation, control and DS samples were integrated and

batch corrected with Harmony integration. Following integration, cell types from control

and DS cluster together (**Figure 1D).** While all cell types are present in both control and

DS, the proportion of each cell type differs (**Figure 1D**). Proportional to total cell number,

there are fewer astrocytes and more inhibitory neurons and microglia in the DS basal

forebrain (**Figure 1D**). Principal component analysis (PCA) of pseudobulk data,

categorized by cell type and genotype, reveals that cell type, not genotype, has the

greatest influence on gene expression between cell types (**Figure 1E**). Non-neuronal glial

cells (astrocytes, GPCs, microglia, oligodendrocytes and OPCs) cluster closely together.

The non-neuronal endothelial cells of the vasculature cluster separate from the other

subtypes. Neurons, inhibitory and BFCNs, form a cluster (**Figure 1E**). The differences in

gene expression that exist due to genotype are not enough to cause control and DS to

form distinct clusters in the PCA plot. In all cell types in the basal forebrain, differentially

expressed genes (DEGs) in DS compared to control are encoded across the genome

(**Figure 1F; Supplement 2D**), consistent with other gene expression data from DS

samples(54-61).

Next, we assessed differences in chromatin accessibility and peak-to-gene

linkages of Hsa21-encoded genes that could account for differential gene expression in

the DS basal forebrain. Following QC of the ATAC-seq data (**Supplement 1G-I**), 16,771

control cells and 14,640 DS cells were used for downstream analysis (**Supplement 1J**).

The integrated multiomic data was clustered, revealing that all cell types passed ATAC

QC and were present in both control and DS samples (**Figure 1G**).

3.2 Hsa21-Encoded Genes

We investigated whether and how Hsa21-encoded genes are dysregulated in DS.

Of the 221 predicted Hsa21 protein-coding genes annotated in the GRCh38.p14

reference assembly, 85 Hsa21 genes are dysregulated between astrocytes, BFCNs,

inhibitory neurons, microglia, and oligodendrocytes, with most genes dysregulated in

more than one cell type (**Figure 2A; Supplement Table 5**). Hsa21 genes represent 1.7%

of astrocyte DEGs, 10.6% of BFCN DEGs, 2.0% of inhibitory neuron DEGs, 3.1% of

microglia DEGs, and 2.0% of oligodendrocyte DEGs (**Figure 2B; Supplement 3A-E**).

These dysregulated Hsa21 genes account for 26.7%, 4.1%, 33.0%, 10.9%, and 21.3% of

total Hsa21 protein-coding genes, respectively (**Figure 2C**). 56 Hsa21 genes are

upregulated and 3 are downregulated in DS astrocytes (**Figure 2A, 2D**); 9 are

upregulated in DS BFCNs (**Figure 2A, 2E**); 72 are upregulated and 1 is downregulated

in DS inhibitory neurons (**Figure 2A, 2F**); 21 are upregulated and 3 are downregulated in

DS microglia (**Figure 2A, 2G**); 42 are upregulated and 5 are downregulated in DS

oligodendrocytes (**Figure 2A, 2H**). 79 Hsa21-encoded genes are dysregulated between

endothelial cells, GPCs, and OPCs (**Supplement 3F–I**), accounting for 2.4%, 3.1%, and

1.8% of total DEGs in these cells (**Supplement 3J**). Hsa21-encoded genes account for

22.2%, 24.9%, and 22.6% of total Hsa21 protein-coding genes, respectively

(**Supplement 3K-N**). Gene set enrichment analysis (GSEA) of biological processes

reveals that upregulated Hsa21 genes in all cell types are enriched in regulation of

developmental process, neuron differentiation, and neurogenesis (**Figure 2I**). These

results reveal dysregulated Hsa21 genes in the basal forebrain that may contribute to the

heterochronic development reported in DS(62-64).

To further investigate the impact of the additional copy of Hsa21 in basal forebrain

cells, we analyzed the chromatin accessibility and peak-to-gene linkages of differentially

expressed Hsa21 genes in astrocytes, BFCNs, inhibitory neurons, microglia, and

oligodendrocytes. However, the number of BFCNs that passed ATAC QC was too low to

calculate statistically significant peak-to-gene linkages in these cells. Two genes per cell

subtype were selected from the Hsa21 DEGs to calculate peak-to-gene linkages for the

other subtypes (**Supplement 4**). In DS astrocytes and microglia, larger peaks are present

at the transcription start site (TSS) of *APP* and *S100B* (**Supplement 4A, 4C**). DS

inhibitory neurons have larger peaks at the TSS of *APP* and *SOD1* (**Supplement 4B**).

TSS peaks in *APP* and *OLIG1* are increased in DS oligodendrocytes (**Supplement 4E**).

Since peaks mark regions of accessible chromatin, these results suggest increased

chromatin accessibility, facilitating the recruitment of transcription factors and enhancers,

which in turn promote active gene transcription. The peak-to-gene links observed in *APP*

and *S100B* (DS astrocytes), *APP* (DS inhibitory neurons), *APP* and *S100B* (DS

microglia), and *OLIG1* (DS oligodendrocytes) may provide insight into potential gene

regulatory elements that control expression of these genes in DS (**Supplement 4A-D**).

Future work identifying these potential regulatory motifs and determining their necessity

for gene regulation will provide insight into the mechanisms controlling the expression of

these genes in DS.

3.3 Astrocytes

Recently, the important role of glia in neurodegenerative diseases, including

DS(65, 66), has emerged. The proportion of astrocytes is decreased and the proportion

of microglia is increased in DS relative to total glial cells (**Supplement 5A**) and total cells

(**Figure 1D**). The differential proportions of these cells may have functional consequences

on both development and degeneration in the BF.

We analyzed the differential gene expression in astrocytes, microglia, and

oligodendrocytes to identify signatures of early deficits in glial cells of the DS basal

forebrain. Identification of DEGs in DS astrocytes (adjusted *P*<.05) revealed dysregulated

genes across the genome(**Figure 3A; Supplement Table 4**). Hsa21 genes constitute

only 1.7% of the dysregulated genes in astrocytes (**Figure 2B**). More Hsa21 genes are

represented than all other chromosomes except Hsa13 and Hsa18 in DS astrocytes when

normalizing the DEGs per chromosome to the number of protein-coding genes of the

chromosome (**Figure 3B**). The top 30 DEGs (from the largest absolute values of the

LFCs) in DS astrocytes are all upregulated genes (**Figure 3C**) GSEA of biological

processes reveals that upregulated genes in DS astrocytes are enriched in ribosome

biogenesis, translation, and ensheathment of neurons and axons (**Figure 3D**).

Meanwhile, downregulated genes are enriched in protein folding and protein maturation

(**Figure 3D**). The five most dysregulated pathways in DS astrocytes- ribosome

biogenesis, cytoplasmic translation, translation, ensheathment of neurons, and axon

ensheathment- consist of genes that are upregulated are shown in the gene-concept

network plot (cnet plot) (**Figure 3E; Supplement Table 6**). The dysregulation of ribosome

and translation genes and genes involved in neuron ensheathment suggests that DS

astrocytes are increasing the production of proteins that mediate interactions with

neurons in the BF.

3.4 Microglia

DEGs (adjusted P<.05) in DS microglia are encoded across the genome(**Figure

4A; Supplement Table 4**). Hsa21-encoded genes are represented at a higher

percentage in DS microglia when taking into account the number of protein-coding genes

per chromosome (**Figure 4B**). The top 30 DEGs in DS microglia comprised of 20

upregulated genes and 10 downregulated genes (**Figure 4C**). GSEA reveals that

upregulated genes in DS microglia of the basal forebrain are enriched in complement

activation and translation whereas downregulated genes are enriched in transcription,

protein folding, and protein maturation (**Figure 4D**). The five most dysregulated pathways

in DS basal forebrain microglia are upregulated complement activation and cytoplasmic

translation processes. Nucleic acid metabolic process, protein maturation processes are

downregulated in DS basal forebrain microglia (**Figure 4E; Supplement Table 6**). These

results reveal that, at birth, there is evidence of an immune response via activation of the

complement system and dysregulation of microglial function in the DS BF.

3.5 Oligodendrocytes

In oligodendrocytes, DEGs (adjusted *P*<.05) are encoded across the

genome(**Figure 5A; Supplement Table 4**). Similar to the other glial cells, Hsa21 genes

are represented at a higher percentage in DS oligodendrocytes when normalizing DEGs

to the chromosome's number of protein-coding genes (**Figure 5B**). The top 30 DEGs

include 28 downregulated and 2 upregulated genes in DS oligodendrocytes (**Figure 5C**).

Upregulated genes in DS oligodendrocytes are enriched in metabolic and secretion

processes (**Figure 5D)**. Downregulated genes are enriched in protein folding and

response to stimuli processes (**Figure 5D**). The top five dysregulated processes-

response to temperature stimulus, response to heat, chaperone-mediated protein folding,

'de novo' protein folding, and protein folding- are all downregulated in in DS

oligodendrocytes of the basal forebrain (**Figure 5E; Supplement Table 6**). These results

suggest that DS oligodendrocytes may have an increase in protein misfolding and may

not be as responsive to environmental cues, potentially contributing to the reduced

myelination reported in DS (56, 67, 68).

3.6 Inhibitory Neurons

Inhibitory neurons account for 9.5% of control cells and 34.1% of DS cells in our

data (**Figure 1D; Supplement 2A-B**). DS inhibitory neurons have DEGs (adjusted

P<.05) distributed across the genome (**Figure 6A; Supplement Table 4**). Hsa21 genes

make up a small percentage of DS inhibitory DEGs (**Figure 2B**) but are overrepresented

when normalizing the DEGs per chromosome to the number of chromosome protein-

coding genes (**Figure 6B**). The top 30 DEGs in DS inhibitory neurons based on the

absolute value of the LFC are all upregulated (**Figure 6C**). GSEA reveals that upregulated

genes in DS Inhibitory neurons of the basal forebrain are enriched in processes of

translation, ATP synthesis, and energy metabolism (**Figure 6D**). Downregulated genes

are enriched in cell adhesion and synaptic transmission pathways (**Figure 6D**). The top

five dysregulated pathways- ribosomal small subunit biogenesis, cytoplasmic translation,

proton transmembrane transport, ATP synthesis coupled electron transport, and oxidative

phosphorylation- are upregulated processes in DS Inhibitory neurons of the basal

forebrain (**Figure 6E; Supplement Table 6**). DS Inhibitory neurons display dysregulation

of genes associated with ATP production, cell adhesion, and synapse formation,

suggesting mitochondrial defects and cell-cell communication deficits may already be

present at birth in the DS basal forebrain.

3.7 Two Distinct BFCN Populations

To identify the BFCNs for downstream analysis, we sub-clustered the inhibitory

neurons and annotated BFCN clusters based on known markers (**Figure 7A;**

Supplement Table 3). Of the total inhibitory neurons in control and DS, BFCNs comprise

5.2% and 2.4%, respectively (**Figure 7A**). snRNA-seq provides an unbiased approach to

identify potential novel markers for BFCNs. However, relative to inhibitory neurons, many

of the top BFCN marker genes have already been characterized in cholinergic neurons

(**Supplement 5B**).

Interestingly, BFCNs formed 2 separate cluster (**Figure 7A**). We then subset the

BFCNs and sub-clustered them, again resulting in the separation of two distinct clusters

(**Figure 7B**). The two clusters were annotated as BFCNs 1 and BFCNs 2 (**Figure 7B**). In

controls these two populations of BFCNs are roughly of equal proportion, whereas in DS

there are more BFCNs 1 (**Figure 7C-D**). Differentially expressed genes calculated

between BFCNs 1 and BFCNs 2 reveal that several tubulin genes and several genes

involved in energy metabolism are upregulated in BFCNs 1 (**Figure 7E**). PCA of the

BFCN subpopulations shows that there is greater variability in the BFCNs 1 population

(**Figure 7F**). These results suggest that BFCNs 1 are more impacted by DS than BFCNs

2.

We compared the expression of known BFCN markers to identify differences in

these two BFCN populations. Both BFCN populations express the established marker

genes of BFCNs, including genes encoding enzymes and transporters in the

acetylcholine (ACh) pathway (*CHAT, ACHE, SLC18A3, SLC5A7)*, neurotrophic receptors

required for the maintenance and survival of BFCNs (*NTRK1, NGFR*), and transcription

factors that regulate these genes (*ISL1, LHX8*). However, BFCNs 2 express most of these

genes at lower levels compared to BFCNs 1 (**Figure 7G**). BFCNs 2 have a significant

reduction in the expression of *SLC18A3,* a transmembrane protein responsible for

transporting ACh into secretory vesicles for release, and *ACHE*, which hydrolyzes ACh

into choline that is recycled for continued ACh synthesis (**Figure 7G**). These results

suggest that BFCNs 2, with reduced expression of essential components for ACh

neurotransmission, are likely not fully functional cholinergic neurons.

Given the developmental stage of the samples and the reduced expression of genes in

the ACh pathway, we suspected that BFCNs 2 are immature compared to BFCNs 1. We

assessed the expression of mature and immature neuron markers in these two

populations (**Figure 7H**). Compared to BFCNs 1, BFCNs 2 have decreased expression

of *ENO2*, a mature neuron marker, and increased expression of *DCX*, an immature

neuron marker (**Figure 7H**). Additionally, BFCNs 2 exhibit decreased expression of the

GABA receptor, *GABBR1,* along with decreased *GAD1* expression, which encodes

GAD65, the enzyme that catalyzes the conversion of glutamate into GABA (**Figure 7H**).

GSEA reveals that energy metabolism pathways are activated in BFCNs 1 whereas cell

adhesion and processes related to synaptic formation are suppressed (**Supplement 5C;**

Supplement Table 6). KEGG pathway analysis reveals upregulated genes in BFCNs 1

are enriched in oxidative phosphorylation and glycolysis/gluconeogenesis pathways

(**Supplement 5D**). GSEA and KEGG analysis suggests that BFCNs 1 are likely more

metabolically active, characteristic of more mature neurons. This increased metabolic

activity coupled with the increased expression of genes in the ACh and GABA pathways,

the increased expression of *ENO2*, and the decreased expression of *DCX* suggest

BFCNs 1 are a more mature population of cells relative to BFCNs 2. The increased

proportion of mature BFCNs 1 in DS supports precocious development of DS BFCNs

(**Figure 7D**). The limited number of DEGs in DS BFCNs 2 (**Supplement 5E; Supplement**

Table 4) precludes pathway analysis and so further analysis was performed only on the

BFCNs 1 population.

3.8 Analysis of Mature BFCNs

We analyzed the DEGs (adjusted $P<.05$) of BFCNs 1 to understand the cellular mechanisms that may contribute to DS BFCN degeneration so early in life. Differentially expressed genes in BFCNs are distributed across the genome (**Figure 8A**). DS BFCNs 1 have 89 upregulated genes and six downregulated genes relative to control (**Figure 8A; Supplement Table 4**). Hsa21 genes make up 10.6% of DS BFCNs 1 DEGs (**Figure 2B**) but are overrepresented when normalizing the DEGs per chromosome to the number of chromosome protein-coding genes (**Figure 8B**). Of the top 30 DEGs in DS BFCNs 1, all are upregulated (**Figure 8C; Supplement Table 4**). Genes upregulated in DS include those encoding subunits of the oxidative phosphorylation pathway (*NDUFS2, COX5A, ATP5PO, ATP5PF,* and *UQCRC2),* antioxidant enzymes (*SOD1, PRDX1,* and *PRDX2)*, and subunits of the vacuolar-type ATPase (V-ATPase) (*ATP6V0B, ATP6V0D1,* and *ATP6V0C)* (**Figure 8C; Supplement Table 4**). Overexpression of V-ATPase, an ATP-driven proton pump that regulates cellular pH and plays a role in overall cell homeostasis, has been linked to several human diseases(69). Additionally, genes associated with glycolysis (*PGAM1, LDHB, LDHA, PDHA1, TPI1, GAPDH,* and *PGK1*) are also upregulated in DS BFCNs (**Figure 8C; Supplement Table 4**). The increased expression of glycolysis-associated genes in neurons that should be relying primarily on the oxidative phosphorylation pathway for ATP production suggests DS BFCNs 1 may be starting to shift toward glycolysis as the primary source for energy production, a shift that is linked to mitochondrial dysfunction and several neurodegenerative diseases(70). Although several genes related to energy metabolism are dysregulated in DS BFCNs, potential compensatory mechanisms may also be at play, evidenced by the increased expression

and increased chromatin accessibility of antioxidant enzymes (**Supplement 5F**) as well

as the increased expression of V-ATPase subunits, all of which help maintain cellular

homeostasis.

GSEA of biological processes reveals that metabolic and regulation of pH

processes are activated in DS BFCNs 1 of the basal forebrain (**Figure 8D**). Response to

stimuli and synapse formation processes are suppressed in DS BFCNs 1 (**Figure 8D**).

Gene-concept network analysis of KEGG pathway enrichment performed with

clusterProfiler reveals that, in addition to oxidative phosphorylation and

glycolysis/gluconeogenesis pathways, genes upregulated in DS BFCNs 1 are enriched in

pathways associated with several neurodegenerative diseases, including AD, PD,

Huntington's disease, prion disease, and amyotrophic lateral sclerosis (ALS) (**Figure 8E;**

Supplement Table 6). KEGG enrichment analysis performed with scProgram reveals

that DS BFCNs 1 DEGs are enriched in the oxidative phosphorylation pathway (**Figure**

8F), further suggesting that there is metabolic dysregulation in DS BFCNs 1.

Gene expression and chromatin accessibility of antioxidant genes indicates there

may be an increase in OXPHOS and resulting ROS in DS BFCNs 1. To validate this

finding, we probed for lipid peroxidation and protein oxidation, downstream consequences

of ROS, to determine if ROS is accumulating in the DS basal forebrain at birth. Protein

carbonylation, the oxidation of protein side chains induced by ROS, is not altered in the

DS basal forebrain (**Figure 8G**). Though variable and not statistically significant, there is

an increase in lipid peroxidation in the DS basal forebrain (**Figure 8H**). Changes in lipid

peroxidation but not in protein oxidation in DS samples suggests that ROS are just

beginning to accumulate and induce damage to lipids, which are more susceptible to

ROS(71-73), in the DS basal forebrain. Altogether, these results suggest that metabolic

dysregulation and ROS are present in the DS basal forebrain at birth, potentially

contributing to the early degeneration of DS BFCNs(14).

4. Discussion

4.1 Summary of findings

Our results provide the first gene expression and chromatin accessibility data of the

human basal forebrain from either healthy or diseased individuals. Cholinergic neurons

in the basal forebrain are vulnerable in age-associated degeneration and implicated in

the progression of multiple neurodegenerative diseases, including AD, PD, and

DLB(74). BFCN degeneration is particularly early in the progression of DS-AD,

detectable when individuals are in their twenties(14), so these data establish a rich

resource for further investigation of neurodegeneration. Thus, our findings of early

neuropathology in both neurons and glia of the basal forebrain in DS may provide

insight into shared mechanisms between several neurodegenerative disorders.

4.2 Limitations of tissue analysis

All studies analyzing post-mortem human tissue suffer from specific features of the

donor and tissue including cause of death, post-mortem interval, and sample integrity

that can affect the results. Samples used in this study were from donors with varied

causes of death (**Supplement Table 1**) and so transcriptional changes due to specific

causes of death are unlikely. Analyzed samples are from early postnatal donors, but we

lack information of gestational duration. We cannot rigorously match samples for stages

of development and analyzed samples that correspond to neonatal and very early

infancy, a developmental phase that includes rapid growth, synaptic remodeling and

myelination(75).

The organization of cholinergic neurons into four nuclei (Ch1-4) whose

anatomical boundaries are not discrete(76, 77) limits our ability to determine in which

nucleus or nuclei the cholinergic neurons we analyzed reside. However, degeneration

of the anteromedial basal forebrain (Ch1-3) and posterior basal forebrain (Ch4) occurs

concomitantly in DS(14), so any changes likely apply to BFCNs from all nuclei.

4.3 Single cell transcriptomic signature of Early Postnatal Basal Forebrain

All expected cell types are present in both control and DS early postnatal DS

basal forebrain, but cell type proportions are altered in DS. Interestingly, the proportions

of neurons and glia are contrary to reports in the DS cortex, where neurons are

reduced, and astrocytes are increased compared to controls(62, 64, 78). These results

suggest that prenatal development of the basal forebrain may be altered in DS, leading

to the generation of different numbers of progenitors and/or neurons or, alternatively,

that neuron degeneration begins prenatally. Analysis of prenatal DS tissue and induced

pluripotent stem cell models are needed to interrogate earlier developmental time

periods to test these possibilities.

We identified molecular events that occur in DS prior to cholinergic

dysfunction(28-30) that provide clues to the vulnerability of BFCNs. We uncovered

dysregulation of genes in all cell types in the early postnatal DS basal forebrain. Few

dysregulated genes and molecular pathways were shared across cell types, suggesting

that the gene expression differences in DS are largely cell type-specific. Although

Hsa21 genes were overrepresented when normalizing to the protein-coding genes,

Hsa21-encoded genes were a small proportion of total dysregulated genes in all cell

types. Functional validation of these gene expression differences will define how each

cell type is affected in DS.

We uncovered two populations of BFCNs (BFCNs 1, BFCNs 2) in the early

postnatal forebrain that are present in control and DS samples. Both populations

express established cholinergic marker genes, although BFCNs 2 express these genes

at much lower levels. Additionally, BFCNs 2 have increased expression of the immature

neuron marker *DCX*, suggesting that these cells are less mature and not fully functional.

It is likely, given the early age of the samples, that BFCNs are still developing, resulting

in immature and mature populations. However, the DS samples have a higher

proportion of the BFCNs 1 population. If the BFCNs 1 population indeed represent

mature neurons, then these data align with the idea of developmental heterochrony that

has been proposed in DS(62-64), in which development progresses precociously in DS.

4.4 BFCN Degeneration

Despite the fact that BFCNs degenerate earlier in DS than in other

neurodegenerative disorders (14), there is no analysis of the human DS basal forebrain

in early life. Previous studies have examined the basal forebrain cholinergic system in

DS and DS-AD, but the youngest individuals in these studies are adolescents and

young adults(11, 14) where AD pathology has already begun (26, 27). We sought to

identify molecular signatures defining vulnerability that occur in DS prior to cholinergic

dysfunction(28-30). Our results reveal that basal forebrain pathology is present as early

as birth in DS, indicating pathological processes begin prenatally. These results suggest

that the perinatal period is a potential window of therapeutic opportunity to mitigate

emerging neuropathology in the DS basal forebrain.

Mitochondrial dysfunction and dysregulated energy metabolism are emerging as

hallmarks of many neurodegenerative diseases, including those that include BFCN

degeneration(70, 79). Relative to control, the more mature population of DS BFCNs

(BFCNs 1) upregulate several genes that encode components of the OXPHOS

pathway, along with antioxidant enzymes, *PRDX1, PRDX2,* and *SOD1*, that detoxify

reactive oxygen species (ROS) byproducts generated from OXPHOS. In response to

ROS accumulation, NRF2 is activated which regulates the expression of antioxidant

enzymes, including superoxide dismutases and peroxiredoxins(80). The upregulation of

these antioxidant enzymes is potentially a compensatory mechanism to detoxify

excessive ROS levels in the DS basal forebrain. We hypothesize that this early

dysregulation of the OXPHOS pathway leads to an accumulation of ROS and the

resulting oxidative stress increases the vulnerability of DS BFCNs. Our gene expression

and biochemical assay results suggest that dysregulated energy metabolism and the

early accumulation of ROS in the DS basal forebrain underlies the susceptibility of

BFCNs to degeneration later in life.

While our results indicate that dysregulated genes in DS BFCNs are associated

with several neurodegenerative diseases, OXPHOS genes are typically downregulated

with age and in neurodegenerative diseases as cells shift toward glycolysis as the

primary source for ATP production, a shift known as the Warburg effect(70, 81). Several

nuclear-encoded OXPHOS subunits are upregulated in DS BFCNs, suggesting that the

OXPHOS pathway is still utilized at birth in DS. In the Ts65Dn mouse model of DS and

AD, OXPHOS genes are downregulated in BFCNs at 6 months of age(82, 83),

approximately when BFCN dysfunction and degeneration begins in this model(84, 85).

The upregulation of OXPHOS genes in early postnatal human BFCNs and the

downregulation in Ts65Dn 6-month BFCNs supports this potential shift in energy

metabolism as these neurons begin to degenerate. The increase in genes encoding

glycolytic enzymes suggests DS BFCNs may be in the early stages of shifting toward

glycolysis as the primary source for energy production. A prolonged shift from OXPHOS

to glycolysis can create an energy deficit that makes cells more susceptible to oxidative

stress and cell death(70). The accumulation of ROS coupled with a shift from OXPHOS

to glycolysis by birth may be an early driver of BFCN vulnerability in DS.

Alternatively, the upregulation of genes encoding antioxidant enzymes and

components of the glycolysis pathway may be attributed to the predominance of female

samples in our study. Biological sex influences the progression of DS-AD

pathogenesis(86), and sex differences have been reported in the basal forebrain

cholinergic system of the Ts65Dn mouse model of DS and AD(87). Recent spatial

transcriptomic analyses of DS-AD samples reveal that genes involved in oxidative

stress and glucose metabolism are upregulated in females compared to males(88). The

upregulation of genes encoding antioxidant enzymes, the primary defense against

oxidative stress, and components of the glycolysis pathway in our study may result from

the fact that three out of four control and DS samples are from female donors. However,

our study is not sufficiently powered to assess the impact of sex differences on the

transcriptome of DS BFCNs.

4.5 Non-neuronal cell dysregulation

669   Differential gene expression in astrocytes, microglia, and oligodendrocytes

indicates neuroinflammation and myelination deficits are present in the DS basal

forebrain as early as birth. DS astrocytes and microglia upregulate genes involved in

translation and are potentially altering their proteomic profiles as they transition from a

homeostatic state to a reactive or activated state, respectively. Activation of the

complement system further suggests that DS microglia of the basal forebrain are

activated. These results are consistent with reports of neuroinflammation, primarily from

the cortex, as DS-AD progresses (89). Protein folding and protein maturation processes

are downregulated in DS basal forebrain oligodendrocytes, suggesting that DS

oligodendrocytes have impaired production of myelin components, possibly contributing

to the reduced myelination of neurons in DS(56, 67, 68). Future work will need to

validate these cell specific transcriptomic changes with biochemical assays and

elucidate how the non-neuronal cells contribute to basal forebrain pathology and early

degeneration of DS BFCNs.

4.6 Future Directions

Taken as a whole, our results reveal that metabolic dysfunction and oxidative stress are

present in the DS basal forebrain by birth. Sustained metabolic dysregulation and

exposure to oxidative stress from birth likely contributes to the susceptibility of BFCNs

so early in the progression of DS-AD in individuals with DS. Elucidation of the shift in

energy metabolism across the DS lifespan with human basal forebrain tissue or in stem

cell models of BFCNs (90-95) will validate the transcriptomic changes. If these

metabolic changes hold true, regulation of the OXPHOS pathway and ROS

accumulation could provide targets for early therapeutic intervention in DS BFCNs prior

to degeneration.

5. References

1.   Ballinger EC, Ananth M, Talmage DA, Role LW. Basal Forebrain Cholinergic Circuits
and Signaling in Cognition and Cognitive Decline. Neuron. 2016;91(6):1199-218.
2.   Davies P, Maloney AJ. Selective loss of central cholinergic neurons in Alzheimer's
disease. Lancet. 1976;2(8000):1403.
3.   Bartus RT, Dean RL, 3rd, Beer B, Lippa AS. The cholinergic hypothesis of geriatric
memory dysfunction. Science. 1982;217(4558):408-14.
4.   Bowen DM, Smith CB, White P, Davison AN. Neurotransmitter-related enzymes and
indices of hypoxia in senile dementia and other abiotrophies. Brain. 1976;99(3):459-96.
5.   Schmitz TW, Nathan Spreng R. Basal forebrain degeneration precedes and predicts the
cortical spread of Alzheimer's pathology. Nat Commun. 2016;7:13249.
6.   Fernández-Cabello S, Kronbichler M, Van Dijk KRA, Goodman JA, Spreng RN, Schmitz
TW. Basal forebrain volume reliably predicts the cortical spread of Alzheimer's degeneration.
Brain. 2020;143(3):993-1009.
7.   Schmitz TW, Mur M, Aghourian M, Bedard M-A, Spreng RN. Longitudinal Alzheimer's
Degeneration Reflects the Spatial Topography of Cholinergic Basal Forebrain Projections. Cell
Reports. 2018;24(1):38-46.
8.   Geula C, Nagykery N, Nicholas A, Wu CK. Cholinergic neuronal and axonal
abnormalities are present early in aging and in Alzheimer disease. Journal of neuropathology
and experimental neurology. 2008;67(4):309-18.
9.   Moss DE, Perez RG. The phospho-tau cascade, basal forebrain neurodegeneration, and
dementia in Alzheimer's disease: Anti-neurodegenerative benefits of acetylcholinesterase
inhibitors. J Alzheimers Dis. 2024;102(3):617-26.
10.   Giacobini E, Cuello AC, Fisher A. Reimagining cholinergic therapy for Alzheimer's
disease. Brain. 2022;145(7):2250-75.
11.   Casanova MF, Walker LC, Whitehouse PJ, Price DL. Abnormalities of the nucleus
basalis in Down's syndrome. Annals of neurology. 1985;18(3):310-3.
12.   Rogers JD, Brogan D, Mirra SS. The nucleus basalis of Meynert in neurological disease:
a quantitative morphological study. Annals of neurology. 1985;17(2):163-70.
13.   Price DL, Whitehouse PJ, Struble RG, Coyle JT, Clark AW, Delong MR, et al.
Alzheimer's Disease and Down's Syndrome. Annals of the New York Academy of Sciences.
1982;396(1):145-64.
14.   Rozalem Aranha M, Iulita MF, Montal V, Pegueroles J, Bejanin A, Vaqué-Alcázar L, et
al. Basal forebrain atrophy along the Alzheimer's disease continuum in adults with Down

syndrome. Alzheimer's & dementia : the journal of the Alzheimer's Association.
2023;19(11):4817-27.
15. Russell JK, Conley AC, Boyd BD, Begnoche JP, Schlossberg R, Stranick A, et al. Age-
Related Changes in the Cholinergic System in Adults with Down Syndrome Assessed Using
[(18)F]-Fluoroethoxybenzovesamicol Positron Emission Tomography Imaging. medRxiv. 2024.
16. Candy JM, Perry RH, Perry EK, Irving D, Blessed G, Fairbairn AF, et al. Pathological
changes in the nucleus of meynert in Alzheimer's and Parkinson's diseases. Journal of the
Neurological Sciences. 1983;59(2):277-89.
17. Nakano I, Hirano A. Parkinson's disease: neuron loss in the nucleus basalis without
concomitant Alzheimer's disease. Annals of neurology. 1984;15(5):415-8.
18. Tagliavini F, Pilleri G, Bouras C, Constantinidis J. The basal nucleus of Meynert in
idiopathic Parkinson's disease. Acta Neurol Scand. 1984;70(1):20-8.
19. Whitehouse PJ, Hedreen JC, White CL, 3rd, Price DL. Basal forebrain neurons in the
dementia of Parkinson disease. Annals of neurology. 1983;13(3):243-8.
20. Bohnen NI, Albin RL. The cholinergic system and Parkinson disease. Behav Brain Res.
2011;221(2):564-73.
21. Grothe MJ, Schuster C, Bauer F, Heinsen H, Prudlo J, Teipel SJ. Atrophy of the
cholinergic basal forebrain in dementia with Lewy bodies and Alzheimer's disease dementia.
Journal of neurology. 2014;261(10):1939-48.
22. Lippa CF, Smith TW, Perry E. Dementia with Lewy bodies: choline acetyltransferase
parallels nucleus basalis pathology. J Neural Transm (Vienna). 1999;106(5-6):525-35.
23. Tiraboschi P, Hansen LA, Alford M, Merdes A, Masliah E, Thal LJ, et al. Early and
Widespread Cholinergic Losses Differentiate Dementia With Lewy Bodies From Alzheimer
Disease. Archives of General Psychiatry. 2002;59(10):946-51.
24. Schumacher J, Ray NJ, Hamilton CA, Donaghy PC, Firbank M, Roberts G, et al.
Cholinergic white matter pathways in dementia with Lewy bodies and Alzheimer's disease.
Brain. 2022;145(5):1773-84.
25. Fortea J, Zaman SH, Hartley S, Rafii MS, Head E, Carmona-Iragui M. Alzheimer's
disease associated with Down syndrome: a genetic form of dementia. The Lancet Neurology.
2021;20(11):930-42.
26. Lott IT, Head E. Dementia in Down syndrome: unique insights for Alzheimer
disease research. Nat Rev Neurol. 2019;15(3):135-47.
27. Wiseman FK, Al-Janabi T, Hardy J, Karmiloff-Smith A, Nizetic D, Tybulewicz VL, et al. A
genetic cause of Alzheimer disease: mechanistic insights from Down syndrome. Nature reviews
Neuroscience. 2015.
28. Kish S, Karlinsky H, Becker L, Gilbert J, Rebbetoy M, Chang LJ, et al. Down's syndrome
individuals begin life with normal levels of brain cholinergic markers. Journal of neurochemistry.
1989;52(4):1183-7.
29. Brooksbank BW, Walker D, Balázs R, Jørgensen OS. Neuronal maturation in the foetal
brain in Down's syndrome. Early Hum Dev. 1989;18(4):237-46.
30. Lubec B, Yoo BC, Dierssen M, Balic N, Lubec G. Down syndrome patients start early
prenatal life with normal cholinergic, monoaminergic and serotoninergic innervation. J Neural
Transm Suppl. 2001(61):303-10.
31. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21.
32. Degatano K, Awdeh A, Dingman W, Grant G, Khajouei F, Kiernan E, et al. WDL Analysis
Research Pipelines: Cloud-Optimized Workflows for Biological Data Processing and
Reproducible Analysis. Preprints. 2024.
33. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. 2011. 2011;17(1):3.

34. Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 2009;25(14):1754-60.
35. Zhang K, Zemke NR, Armand EJ, Ren B. A fast, scalable and versatile tool for analysis
of single-cell omics data. Nature Methods. 2024;21(2):217-27.
36. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, et al. Integrated
analysis of multimodal single-cell data. Cell. 2021;184(13):3573-87.e29.
37. Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, 3rd, et al.
Comprehensive Integration of Single-Cell Data. Cell. 2019;177(7):1888-902.e21.
38. Young MD, Behjati S. SoupX removes ambient RNA contamination from droplet-based
single-cell RNA sequencing data. Gigascience. 2020;9(12).
39. Mathys H, Peng Z, Boix CA, Victor MB, Leary N, Babu S, et al. Single-cell atlas reveals
correlates of high cognitive function, dementia, and resilience to Alzheimer's disease pathology.
Cell. 2023;186(20):4365-85.e27.
40. Gao Y, Dong Q, Arachchilage KH, Risgaard R, Sheng J, Syed M, et al. Multimodal
analyses reveal genes driving electrophysiological maturation of neurons in the primate
prefrontal cortex. bioRxiv. 2024.
41. McGinnis CS, Murrow LM, Gartner ZJ. DoubletFinder: Doublet Detection in Single-Cell
RNA Sequencing Data Using Artificial Nearest Neighbors. Cell Syst. 2019;8(4):329-37.e4.
42. Wolock SL, Lopez R, Klein AM. Scrublet: Computational Identification of Cell Doublets in
Single-Cell Transcriptomic Data. Cell Syst. 2019;8(4):281-91.e9.
43. Germain PL, Lun A, Garcia Meixide C, Macnair W, Robinson MD. Doublet identification
in single-cell sequencing data using scDblFinder. F1000Res. 2021;10:979.
44. Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, et al. Fast, sensitive and
accurate integration of single-cell data with Harmony. Nat Methods. 2019;16(12):1289-96.
45. Chao J, Li Z, Sun Y, Aluko OO, Wu X, Wang Q, et al. MG2C: a user-friendly online tool
for drawing genetic maps. Molecular Horticulture. 2021;1(1):16.
46. Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. Fast gene
set enrichment analysis. bioRxiv. 2021:060012.
47. Yu G, Wang L-G, Han Y, He Q-Y. clusterProfiler: an R Package for Comparing
Biological Themes Among Gene Clusters. OMICS: A Journal of Integrative Biology.
2012;16(5):284-7.
48. Hao Y, Stuart T, Kowalski MH, Choudhary S, Hoffman P, Hartman A, et al. Dictionary
learning for integrative, multimodal and scalable single-cell analysis. Nature Biotechnology.
2024;42(2):293-304.
49. Stuart T, Srivastava A, Madad S, Lareau CA, Satija R. Single-cell chromatin state
analysis with Signac. Nat Methods. 2021;18(11):1333-41.
50. Liu T. Use model-based Analysis of ChIP-Seq (MACS) to analyze short reads generated
by sequencing protein-DNA interactions in embryonic stem cells. Methods in molecular biology
(Clifton, NJ). 2014;1150:81-95.
51. Gaspar JM. Improved peak-calling with MACS2. bioRxiv. 2018:496521.
52. Takács VT, Cserép C, Schlingloff D, Pósfai B, Szőnyi A, Sos KE, et al. Co-transmission
of acetylcholine and GABA regulates hippocampal states. Nature Communications.
2018;9(1):2848.
53. Saunders A, Granger AJ, Sabatini BL. Corelease of acetylcholine and GABA from
cholinergic forebrain neurons. eLife. 2015;4:e06412.
54. Lockstone HE, Harris LW, Swatton JE, Wayland MT, Holland AJ, Bahn S. Gene
expression profiling in the adult Down syndrome brain. Genomics. 2007;90(6):647-60.
55. Mao R, Zielke CL, Zielke HR, Pevsner J. Global up-regulation of chromosome 21 gene
expression in the developing Down syndrome brain. Genomics. 2003;81(5):457-67.

56.   Olmos-Serrano JL, Kang HJ, Tyler WA, Silbereis JC, Cheng F, Zhu Y, et al. Down
Syndrome Developmental Brain Transcriptome Reveals Defective Oligodendrocyte
Differentiation and Myelination. Neuron. 2016;89(6):1208-22.
57.   Aït Yahya-Graison E, Aubert J, Dauphinot L, Rivals I, Prieur M, Golfier G, et al.
Classification of human chromosome 21 gene-expression variations in Down syndrome: impact
on disease phenotypes. Am J Hum Genet. 2007;81(3):475-91.
58.   Mao R, Wang X, Spitznagel EL, Jr., Frelin LP, Ting JC, Ding H, et al. Primary and
secondary transcriptional effects in the developing human Down syndrome brain and heart.
Genome Biol. 2005;6(13):R107.
59.   Pelleri MC, Cattani C, Vitale L, Antonaros F, Strippoli P, Locatelli C, et al. Integrated
Quantitative Transcriptome Maps of Human Trisomy 21 Tissues and Cells. Front Genet.
2018;9:125-.
60.   FitzPatrick DR, Ramsay J, McGill NI, Shade M, Carothers AD, Hastie ND. Transcriptome
analysis of human autosomal trisomy. Human molecular genetics. 2002;11(26):3249-56.
61.   Letourneau A, Santoni FA, Bonilla X, Sailani MR, Gonzalez D, Kind J, et al. Domains of
genome-wide gene expression dysregulation in Down's syndrome. Nature.
2014;508(7496):345-50.
62.   Russo ML, Sousa AMM, Bhattacharyya A. Consequences of trisomy 21 for brain
development in Down syndrome. Nature reviews Neuroscience. 2024;25(11):740-55.
63.   Zhou Y, Tao L, Zhu Y. TempShift Reveals the Sequential Development of Human
Neocortex and Skewed Developmental Timing of Down Syndrome Brains. Brain Sciences.
2023;13(7):1070.
64.   Stagni F, Giacomini A, Emili M, Guidi S, Bartesaghi R. Neurogenesis impairment: An
early developmental defect in Down syndrome. Free Radical Biology and Medicine.
2018;114:15-32.
65.   Stevenson R, Samokhina E, Rossetti I, Morley JW, Buskila Y. Neuromodulation of Glial
Function During Neurodegeneration. Frontiers in Cellular Neuroscience. 2020;14.
66.   García O, Flores-Aguilar L. Astroglial and microglial pathology in Down syndrome: Focus
on Alzheimer's disease. Front Cell Neurosci. 2022;16:987212.
67.   Wisniewski KE, Schmidt-Sidor B. Postnatal delay of myelin formation in brains from
Down syndrome infants and children. Clinical neuropathology. 1989;8(2):55-62.
68.   Ábrahám H, Vincze A, Veszprémi B, Kravják A, Gömöri É, Kovács GG, et al. Impaired
myelination of the human hippocampal formation in Down syndrome. International Journal of
Developmental Neuroscience. 2012;30(2):147-58.
69.   Santos-Pereira C, Rodrigues LR, Côrte-Real M. Emerging insights on the role of V-
ATPase in human diseases: Therapeutic challenges and opportunities. Med Res Rev.
2021;41(4):1927-64.
70.   Jadiya P, Garbincius JF, Elrod JW. Reappraisal of metabolic dysfunction in
neurodegeneration: Focus on mitochondrial function and calcium signaling. Acta
neuropathologica communications. 2021;9(1):124.
71.   Endale HT, Tesfaye W, Mengstie TA. ROS induced lipid peroxidation and their role in
ferroptosis. Frontiers in Cell and Developmental Biology. 2023;Volume 11 - 2023.
72.   Yang WS, Kim KJ, Gaschler MM, Patel M, Shchepinov MS, Stockwell BR. Peroxidation
of polyunsaturated fatty acids by lipoxygenases drives ferroptosis. Proceedings of the National
Academy of Sciences. 2016;113(34):E4966-E75.
73.   Su L-J, Zhang J-H, Gomez H, Murugan R, Hong X, Xu D, et al. Reactive Oxygen
Species-Induced Lipid Peroxidation in Apoptosis, Autophagy, and Ferroptosis. Oxidative
Medicine and Cellular Longevity. 2019;2019(1):5080843.
74.   Perry EK. Nerve growth factor and the basal forebrain cholinergic system: a link in the
etiopathology of neurodegenerative dementias? Alzheimer Dis Assoc Disord. 1990;4(1):1-13.

75.   Kang HJ, Kawasawa YI, Cheng F, Zhu Y, Xu X, Li M, et al. Spatio-temporal
transcriptome of the human brain. Nature. 2011;478(7370):483-9.
76.   Mesulam MM, Mufson EJ, Levey AI, Wainer BH. Cholinergic innervation of cortex by the
basal forebrain: cytochemistry and cortical connections of the septal area, diagonal band nuclei,
nucleus basalis (substantia innominata), and hypothalamus in the rhesus monkey. J Comp
Neurol. 1983;214(2):170-97.
77.   Mesulam MM, Geula C. Nucleus basalis (Ch4) and cortical cholinergic innervation in the
human brain: observations based on the distribution of acetylcholinesterase and choline
acetyltransferase. J Comp Neurol. 1988;275(2):216-40.
78.   Zdaniuk G, Wierzba-Bobrowicz T, Szpak GM, Stępień T. Astroglia disturbances during
development of the central nervous system in fetuses with Down's syndrome. Folia
Neuropathol. 2011;49(2):109-14.
79.   Hroudová J, Singh N, Fišar Z. Mitochondrial dysfunctions in neurodegenerative
diseases: relevance to Alzheimer's disease. Biomed Res Int. 2014;2014:175062.
80.   Kasai S, Shimizu S, Tatara Y, Mimura J, Itoh K. Regulation of Nrf2 by Mitochondrial
Reactive Oxygen Species in Physiology and Pathology. Biomolecules. 2020;10(2).
81.   Emelyanova L, Preston C, Gupta A, Viqar M, Negmadjanov U, Edwards S, et al. Effect
of Aging on Mitochondrial Energetics in the Human Atria. J Gerontol A Biol Sci Med Sci.
2018;73(5):608-16.
82.   Alldred MJ, Lee SH, Stutzmann GE, Ginsberg SD. Oxidative Phosphorylation Is
Dysregulated Within the Basocortical Circuit in a 6-month old Mouse Model of Down Syndrome
and Alzheimer's Disease. Frontiers in Aging Neuroscience. 2021;13.
83.   Alldred MJ, Penikalapati SC, Lee SH, Heguy A, Roussos P, Ginsberg SD. Profiling
Basal Forebrain Cholinergic Neurons Reveals a Molecular Basis for Vulnerability Within the
Ts65Dn Model of Down Syndrome and Alzheimer's Disease. Mol Neurobiol. 2021;58(10):5141-
62.
84.   Holtzman DM, Li YW, DeArmond SJ, McKinley MP, Gage FH, Epstein CJ, et al. Mouse
model of neurodegeneration: atrophy of basal forebrain cholinergic neurons in trisomy 16
transplants. Proceedings of the National Academy of Sciences of the United States of America.
1992;89(4):1383-7.
85.   Granholm A-CE, Sanders LA, Crnic LS. Loss of Cholinergic Phenotype in Basal
Forebrain Coincides with Cognitive Decline in a Mouse Model of Down's Syndrome.
Experimental neurology. 2000;161(2):647-63.
86.   Andrews EJ, Martini AC, Head E. Exploring the role of sex differences in Alzheimer's
disease pathogenesis in Down syndrome. Frontiers in Neuroscience. 2022;16.
87.   Kelley CM, Powers BE, Velazquez R, Ash JA, Ginsberg SD, Strupp BJ, et al. Sex
differences in the cholinergic basal forebrain in the Ts65Dn mouse model of Down syndrome
and Alzheimer's disease. Brain Pathol. 2014;24(1):33-44.
88.   Miyoshi E, Morabito S, Henningfield CM, Das S, Rahimzadeh N, Shabestari SK, et al.
Spatial and single-nucleus transcriptomic analysis of genetic and sporadic forms of Alzheimer's
disease. Nat Genet. 2024;56(12):2704-17.
89.   Flores-Aguilar L, Iulita MF, Kovecses O, Torres MD, Levi SM, Zhang Y, et al. Evolution
of neuroinflammation across the lifespan of individuals with Down syndrome. Brain.
2020;143(12):3653-71.
90.   Wu Y, West NR, Bhattacharyya A, Wiseman FK. Cell models for Down syndrome-
Alzheimer's disease research. Neuronal Signaling. 2022;6(1):NS20210054.
91.   Watson LA, Meharena HS. From neurodevelopment to neurodegeneration: utilizing
human stem cell models to gain insight into Down syndrome. Front Genet. 2023;14.
92.   Muñoz SS, Engel M, Balez R, Do-Ha D, Cabral-da-Silva MC, Hernández D, et al. A
Simple Differentiation Protocol for Generation of Induced Pluripotent Stem Cell-Derived Basal

Forebrain-Like Cholinergic Neurons for Alzheimer's Disease and Frontotemporal Dementia
Disease Modeling. Cells. 2020;9(9):2018.
93.   Tao R, Yue C, Guo Z, Guo W, Yao Y, Yang X, et al. Subtype-specific neurons from
patient iPSCs display distinct neuropathological features of Alzheimer's disease. Cell
Regeneration. 2024;13(1):21.
94.   Hu Y, Qu Z-y, Cao S-y, Li Q, Ma L, Krencik R, et al. Directed differentiation of basal
forebrain cholinergic neurons from human pluripotent stem cells. Journal of Neuroscience
Methods. 2016;266:42-9.
95.   Duan L, Bhattacharyya BJ, Belmadani A, Pan L, Miller RJ, Kessler JA. Stem cell derived
basal forebrain cholinergic neurons from Alzheimer's disease patients are more susceptible to
cell death. Molecular Neurodegeneration. 2014;9(1):3.
940

6. Acknowledgements

Human tissue was obtained from the NIH NeuroBioBank at the University of

Maryland, Baltimore, MD. N.R.W. and A.B. would like to acknowledge the contributions

of Bhattacharyya Lab members Dr. Matthew Russo, Grace Branger, Kamryn Witkowiak,

Rachel Lichte, and Sahith Puthireddy for their preliminary stereological analysis of the

basal forebrain that played a key role in the conceptualization of this project. We would

like the acknowledge the feedback on this study provided by members of Dr. Su-Chun

Zhang's lab, members of Dr. Andre Sousa's lab, Dr. Xinyu Zhao, Dr. Luigi Puglielli, Dr.

Elizabeth Head, and Dr. Carissa Sirois. N.R.W. received training in bioinformatic analysis

from the INCLUDE Data Science for Developing Scholars in Down Syndrome (DS3)

which is supported by the National Institutes of Health INCLUDE Project under

R25HD114950 from the Eunice Kennedy Shriver National Institute of Child Health and

Human Development.

7. Conflicts of Interest

The authors have no conflicts of interest to disclose.

8. Funding Sources

Research reported in this publication was supported by the National Institutes of Health under F99AG086925 from the Office of the Director, National Institutes of Health and the National Institute on Aging awarded to N.R.W.; GM140935 Medical Scientist Training Program T32 and F30MH140382 Ruth L. Kirschstein Individual Predoctoral NRSA from the National Institutes of Health awarded to R.D.R.; R36AG072108 from the National Institute on Aging awarded to J.L.M; R01HD106197 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development awarded to S-C.Z., D.F., A.M.M.S., and A.B; and P50HD105353 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development awarded as a core grant to the Waisman Center. Additional funding came from the University of Wisconsin-Madison Sophomore Research Fellowship awarded to S.M.; BRFSG-2023-11 from the Brain Research Foundation awarded to A.M.M.S; and the William F. Vilas Trust Estate Award, Wisconsin Alzheimer's Disease Research Center Research Education Component Award, and the Wisconsin Partnership Program New Investigator Award from the University of Wisconsin-Madison to A.B. The content of this study is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

9. Consent Statement

Acquisition of the de-identified human tissue samples was approved by the Health Sciences Institutional Review Board at the University of Wisconsin-Madison (Protocol #2016-0979) and certified exempt from human subjects IRB oversight.

 10. Key Words:

Down syndrome; Basal Forebrain Cholinergic Neurons (BFCNs); snMultiomic Analysis;

Oxidative Phosphorylation (OXPHOS); Reactive Oxygen Species (ROS); Glycolysis

11. Figure Legends

Figure 1. A) BF samples from four control and four DS donors matched for age, sex, and

PMI were used in this study. B) Schematic of BF snMultiomic analysis (Created with

BioRender.com). C) Cell types and the numbers identified in control and DS samples. D)

Gene expression UMAP of cell clusters by subtype. Cell subtypes identified in the human

basal forebrain were astrocytes, BFCNs, endothelial cells, GPCs, inhibitory neurons,

microglia, oligodendrocytes, and OPCs. Percentages of cell types in control (N=20,826)

and DS (N=16,641). There are fewer astrocytes and more inhibitory neurons and

microglia in the DS basal forebrain. E) PCA analysis of cell type and genotype. Cell types

from control and DS are largely the same, with control and DS clustering together on the

plot. Cell subtypes cluster by subclass. F) DEGs per chromosome by cell type. DEGs are

distributed across the genome in all cell types. G) Gene expression and ATAC integrated

UMAP of cell clusters by subtype.

Figure 2. A) Hsa21-encoded genes differentially expressed in astrocytes, BFCNs,

inhibitory neurons, microglia, and oligodendrocytes. Boxes next to each gene represent

the cell types that gene is dysregulated in. B) Percent of Hsa21 DEGs relative to total

DEGs per cell type. C) Percent of Hsa21 DEGs normalized to total Hsa21 protein-coding

genes per cell type. D) Volcano plot of dysregulated genes in DS astrocytes with Hsa21

genes of interest labeled. E) Volcano plot of dysregulated genes in DS BFCNs with Hsa21

genes of interest labeled. F) Volcano plot of dysregulated genes in DS inhibitory neurons

with Hsa21 genes of interest labeled. G) Volcano plot of dysregulated genes in DS

microglia with Hsa21 genes of interest labeled. H) Volcano plot of dysregulated genes in

DS oligodendrocytes with Hsa21 genes of interest labeled. Hsa21 genes of interest that

are labeled on the volcano plots are *APP, S100B, DYRK1A, DSCAM, SOD1, RCAN1,*

IFNAR1, IFNAR2, IFNGR2, SYNJ1, BACE2, and *BACH1*. I) Gene set enrichment

analysis of Hsa21 genes dysregulated across all subtypes. Dysregulated Hsa21 genes

are enriched in regulation of developmental process, neuron differentiation, and

neurogenesis processes.

1014

Figure 3. A) DEGs are distributed across the genome in DS astrocytes. B) Percent of

DEGs normalized to protein-coding genes per chromosome in DS astrocytes with Hsa21

highlighted in red. C) Top 30 DEGs in DS astrocytes. D) DS astrocyte upregulated genes

are enriched in ribosome biogenesis, translation, and ensheathment of neurons and

axons while downregulated genes are enriched in protein folding and protein maturation.

E) CNET plot of the top 5 dysregulated biological processes in DS astrocytes. Genes for

each category node are listed in Supplement Table 6.

1022

1023   **Figure 4.** A) DEGs are distributed across the genome in DS microglia. B) Percent of

1024   DEGs normalized to protein-coding genes per chromosome in DS microglia with Hsa21

1025   highlighted in red. C) Top 30 DEGs in DS microglia. D) DS microglia upregulated genes

1026   are enriched in complement activation and translation whereas downregulated genes are

enriched in transcription, protein folding, and protein maturation. E) CNET plot of the top

5 dysregulated biological processes in DS microglia. Genes for each category node are

listed in Supplement Table 6.

Figure 5. DEGs are distributed across the genome in DS oligodendrocytes. B) Percent

of DEGs normalized to protein-coding genes per chromosome in DS oligodendrocytes

with Hsa21 highlighted in red. C) Top 30 DEGs in DS oligodendrocytes. D) DS

oligodendrocyte upregulated genes are enriched in metabolic and secretion processes,

and downregulated genes are enriched in protein folding and response to stimuli

processes. E) CNET plot of the top 5 dysregulated biological processes in DS

oligodendrocytes. Genes for each category node are listed in Supplement Table 6.

Figure 6. A) DEGs are distributed across the genome in DS inhibitory neurons. B) Percent

of DEGs normalized to protein-coding genes per chromosome in DS inhibitory neurons

with Hsa21 highlighted in red. C) Top 30 DEGs in DS inhibitory neurons. D) DS inhibitory

neuron upregulated genes are enriched in processes of translation, ATP synthesis, and

energy metabolism. Downregulated genes are enriched in cell adhesion and synaptic

transmission pathways. E) CNET plot of the top 5 dysregulated biological processes in

DS inhibitory neurons. Genes for each category node are listed in Supplement Table 6.

Figure 7. A) UMAP of all inhibitory neurons. BFCNs cluster in two distinct clusters.

Proportion of BFCN subpopulations in control and DS. B) BFCNs were subset from the

data and reclustered, again clustering into two distinct populations that were labeled

BFCNs 1 and BFCNs 2. C) Counts of BFCNs1 and BFCNs per condition. D) Proportion

of BFCN populations in control and DS. E) BFCNs 1 DEGs relative to BFCNs 2. F) PCA

of BFCN populations. There is more variability between control and DS BFCNs 1. G)

Expression of known cholinergic marker genes in BFCN subpopulations. BFCNs 2 have

lower expression of most of these genes. H) Expression of immature and mature neuron

marker genes in BFCN subpopulations. BFCNs 1 have higher expression of *ENO2*, a

mature neuron marker, while BFCNs 2 have increased expression of DCX, an immature

neuron marker. BFCNs 1 also have higher expression of *GAD1*, encoding the enzyme

that catalyzes the conversion of glutamate into GABA, and *GABBR1*, a GABA receptor

1059

Figure 8. A) DEGs are distributed across the genome in DS BFCNs 1. B) Percent of

DEGs normalized to protein-coding genes per chromosome in DS BFCNs 1 with Hsa21

highlighted in red. C) Top 30 DEGs in DS BFCNs 1. D) Metabolic and regulation of pH

processes are activated in DS BFCNs 1. Response to stimuli and synapse formation

processes are suppressed in DS BFCNs 1. E) CNET plot of 5 dysregulated KEGG

pathways in DS BFCNs 1 calculated with clusterProfiler. Dysregulated BFCNs 1 genes

are enriched oxidative phosphorylation and glycolysis/gluconeogenesis pathways and

are associated with several neurodegenerative diseases. Genes for each category node

are listed in Supplement Table 6. F) KEGG pathway analysis of enriched genes in control

and DS BFCNs 1 calculated with scProgram. In DS BFCNs 1, DEGs are enriched in the

oxidative phosphorylation pathway and these DEGs are associated with Parkinson's and

Huntington's diseases. G) Protein oxidation assay from the bulk basal forebrain tissue.

There is no difference in the protein carbonyl content in control and DS, indicating no

change in protein oxidation in the DS basal forebrain. H) Lipid peroxidation assay from

the bulk basal forebrain tissue. There is an increase in the malondialdehyde (MDA)

content in DS, indicating increased lipid peroxidation in the DS basal forebrain. The large

variability between DS samples suggests that ROS are just beginning to accumulate in

the DS basal forebrain during this period of development.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8