DEEP ORIGINS AND DISTINCT ADAPTATIONS INDICATED FOR A GLACIAL RELICT SEAL

Ari Löytynoja[1,*], Jaakko Pohjoismäki[2,*], Mia Valtonen[3], Juha Laakkonen[4], Wataru Morita[5], Mervi Kunnasranta[2,6], Risto Väinölä[7], Morten Tange Olsen[8], Petri Auvinen[9], Jukka Jernvall[9,10,*]

[1]Organismal and Evolutionary Biology Research Programme, Faculty of Biological and Environmental Sciences, University of Helsinki, P.O. Box 65, 00014 Helsinki, Finland
[2]Department of Environmental and Biological Sciences, University of Eastern Finland, P.O. Box 111, 80101 Joensuu, Finland
[3]Natural Resources Institute Finland (Luke), P.O. Box 2, 00791 Helsinki, Finland
[4]Division of Veterinary Anatomy and Developmental Biology, Department of Veterinary Biosciences, Faculty of Veterinary Medicine, P.O. Box 66, University of Helsinki, 00014 Helsinki, Finland
[5]Department of Anthropology, National Museum of Nature and Science, Tsukuba, Ibaraki, Japan
[6]Natural Resources Institute Finland (Luke), Yliopistokatu 6, 80100 Joensuu, Finland
[7]Finnish Museum of Natural History, P.O. Box 17, University of Helsinki, 00014 Helsinki, Finland
[8]Section of Molecular Ecology and Evolution, Globe Institute, University of Copenhagen, Copenhagen, Denmark
[9]Institute of Biotechnology, HiLIFE, P.O. Box 56, University of Helsinki, 00014 Helsinki, Finland
[10]Department of Geosciences and Geography, P.O. Box 64, University of Helsinki, 00014 Helsinki, Finland

Email: ari.loytynoja@helsinki.fi, jaakko.pohjoismaki@uef.fi. jernvall@fastmail.fm

Abstract

Isolated populations of postglacial relics are known from many regions and are typically found on mountains for terrestrial species and in lakes for aquatic species. Among the few aquatic mammalian relics, the Saimaa ringed seal (*Pusa hispida saimensis*) has been land-locked in Lake Saimaa, Finland, for the last 10,000 years. Saimaa ringed seals show genetic, behavioral, and morphological differences to the other ringed seal subspecies, but the extent these differences stem from the end of the last glacial period remains unclear. Here, we demonstrate with comprehensive sampling and state-of-the-art genomic methods that the Saimaa ringed seals are much older than the lake they inhabit, having formed a separate evolutionary branch for at least 60,000 years ago. This deep evolutionary origin of the Saimaa ringed seals is further underscored by our ecomorphological analyses revealing adaptively distinct features in their dentition and tongue. Overall, glacial relics, many threatened by extinction, may harbor a richer selection of evolutionary history than might be expected from their recent isolation history alone.

Keywords: mammal, subspecies, genomic analysis, phenotypic, freshwater pinniped, Lake Saimaa

Introduction

Ringed seals *Pusa hispida* (Schreber, 1775) are the most common Arctic pinnipeds, found throughout all seasonally ice-covered northern seas and two freshwater lakes (Figure 1A). They are highly ice-associated, adapted to maintaining breathing holes and constructing subnivean lairs for parturition, nursing, and resting (McLaren 1958, Kelly 2022). The nominate su bspecies, *P. hispida hispida* (Schreber, 1775) (hereafter the "Arctic ringed seal"), is the most widespread, inhabiting the circumpolar Arctic Ocean. Three subspecies are present in the Fennoscandia (Figure 1B), the Baltic ringed seal *P. h. botnica* (Gmelin, 1788), the Ladoga ringed seal *P. h. ladogensis* (Nordqvist, 1899), and the Saimaa ringed seal *P. h. saimensis* (Nordqvist, 1899). The presence of these three Fennoscandian subspecies has been explained by the isolation of the ringed seals into the Baltic Sea basin after the last glacial period with further entrapment of seals into Lake Ladoga and Lake Saimaa (Ukkonen et al. 2014). Lake Saimaa is a freshwater lake with a surface area of 4400 km^2 and houses a population of close to 500 endemic ringed seals, making these one of the most endangered pinnipeds in the world (Kovacs et al. 2012, https://www.metsa.fi/en/nature-and-heritage/species/saimaa-ringed-seal/). Currently, the population is slowly growing due to active conservation efforts, but the Saimaa ringed seal is still threatened by bycatch mortality and climate change driven habitat change (Kunnasranta et al. 2021).

The morphological and ecological differences between the Saimaa ringed seal and the other subspecies have been long known (Nordqvist 1899, Sipilä et al. 1990, , Hyvärinen and Nieminen 1990, Sipilä and Hyvärinen 1998, Kunnasranta 2001, Amano et al. 2002), historically assumed to derive from the postglacial isolation of Saimaa population from Baltic Sea population. However, genetic analyses of mitochondrial DNA (mtDNA) haplotypes (Valtonen et al. 2012, 2014, Heino et al. 2023), microsatellite loci (Palo et al. 2001, 2003, Nyman et al. 2014), and genome-wide variation (Löytynoja et al. 2023, Sundell et al. 2023) have revealed greater genetic differences between Saimaa and other populations than can be explained by postglacial drift and recent adaptation. Limited sampling in many of these studies has left open the possible contribution of other ringed seal populations, especially in the eastern hemisphere, whose distribution might have reached Fennoscandia during the last glacial period.

In this study, we further explore the genomic differences between the Saimaa ringed seal and other ringed seals, including the previously unsampled Arctic ringed seals in northern Eurasia, as well as the Okhotsk ringed seal *P. h. ochotensis* (Pallas, 1811), an eastern hemisphere representative for the species (Figure 1). In addition to mitochondrial and nuclear genomes, we examine feeding morphology differences to address the magnitude of ecological differentiation among the subspecies. We show that even with the circumpolar sampling of the Arctic ringed seal populations, the Saimaa ringed seal retains its genetic uniqueness. Based on the genomic divergence from the other ringed seals, together with specialized morphological features linked

81 to feeding ecology, we argue that the Saimaa ringed seal is evolutionary more unique than
82 previously thought.

Results

Genetic origins and differentiation of the Saimaa ringed seal

Our phylogenetic analysis of mtDNA data, representing the first truly global sampling of
mitochondrial diversity in ringed seals, confirms earlier findings and shows Saimaa forming a
tight cluster with closely-related haplotypes present but rare among the Baltic and Arctic ringed
seals (Figure S1) (Valtonen et al. 2012, Nyman et al. 2014, Heino et al. 2023). The wider
sampling from the eastern hemisphere fails to establish a recent connection between Saimaa and
other populations, including Okhotsk seals. Similarly to Artic and Baltic ringed seals, the
haplotypes of the sampled Okhotsk individuals are spread around the phylogenetic tree with the
exception of the Saimaa clade (Figure S1). Despite being embedded among the global mtDNA
variation, the Saimaa population is highly unique among the ringed seals and, as the only
distinguished subspecies, forms a monophyletic lineage of mtDNA haplotypes.

While the nonrecombining mtDNA haplotypes reveal deep relationships, the nuclear genome is
expected to reflect the more recent events. Still, the Saimaa sample forms a tight cluster in the
principal component analysis (PCA) based on 1.808 million SNPs across the whole nuclear
genomes from five individuals per population. The Saimaa sample is separated from all the
others, including the Okhotsk ringed seal, by PC1 whereas the other populations are separated by
PC2 (Figure 2A). The full 46 individual dataset produces largely similar division between the
populations but places the Arctic and Okhotsk samples closer to each other (Figure 2B), possibly
as an artifact of uneven sampling (Elhaik 2022). Furthermore, the derived allele statistics show
that the Saimaa ringed seals retain a high representation of population-specific variants, a pattern
preserved when multiple individuals are compared (Figure 2C). The relative magnitude of the
private variants in Saimaa makes it highly implausible that this variation has resulted from drift
during the 10 kya, or roughly 1000-generation entrapment in the lake. This raises the question
how deep are the roots of the Saimaa population.

To better understand the population histories, we focused on the divergence times among the
three Fennoscandian populations (Baltic, Ladoga, Saimaa) in reference to the Arctic population
using MSMC-IM (Wang et al. 2020), a demographic inference method providing time-dependent
estimates of population-specific N_e as well as gene flow. Even with this improved modeling
compared to a previous analysis (Löytynoja et al. 2023), the inferred N_e trajectories for the
Saimaa population are clearly different from the other comparisons and show the dramatic
decrease in effective population size over the past several thousand years (Figure 3A). The
estimates of migration rate over time (Figure 3B) demonstrate the recent strict isolation of the
Saimaa population but also show the Saimaa lineage to have been in contact with the Baltic and
Ladoga around 8–10 kya and the migration been at its greatest around 50–80 kya (Figure 3B).

The gene flow within the three Fennoscandian populations at 8–10 kya provides an explanation
for the Saimaa-like mtDNA haplotypes and microsatellite alleles being present but rare in the
Baltic population (Palo et al. 2001, Palo et al. 2003, Valtonen et al. 2012, Nyman et al. 2014). On
the other hand, compared to the Baltic, the slightly earlier isolation of Ladoga ringed seals from
the Arctic (Figure 3B) can explain the reduced mtDNA haplotype variation observed in that
population (Figure S1).

To further date the roots of the Saimaa lineage, we computed M, the cumulative migration
probability over time, between each population pair (Figure 3C). These confirm the delayed
accumulation of migration between Saimaa and the three other populations, while the separate
trajectories for the Arctic-Ladoga pairs support multiple waves of gene flow from the Atlantic to
the Baltic region, some of that not reaching Ladoga. As a proxy for the population split times
(Wang et al. 2020), we estimated the times for M reaching 50% (M_{50}) as well as the more
stringent thresholds of 95% (M_{95}) and 99% (M_{99}) (Table 2). With the applied parameters of μ
and g used, M_{50} for Saimaa varies between 60.2 kya (with Baltic) and 65.8 kya (with Arctic).
These are much deeper in the past than among the three non-Saimaa populations, varying from
29.2 kya to 41.7 kya (Table 1). In comparisons including Saimaa individuals, the average M_{95}
are over 261 kya and M_{99} are over 372 kya, indicating very deep roots for the Saimaa lineage
(Table 1). It is also notable that the M cut-off values for the Saimaa-Ladoga pairs are deeper in
the past than they are for the Saimaa-Baltic pairs, highlighting the distinct origins of the two lake
populations.

The early isolation of the Saimaa lineage between 10–60 kya could be explained by a refugium
in the glacial lakes either on the eastern or southern side of the Fennoscandian ice sheet. During
90–50 kya, large glacial lakes existed in the West Siberian Plains and White Sea Basin with
water routes to the eastern edge of the Fennoscandian ice sheet (Mangerud et al. 2001,
Mangerud, et al. 2004, Krinner et al. 2004). Similarly, a chain of ice lakes existed at the
southeastern border of the glacier, in current-day western Russia and Belarus, with connections
to the Caspian Sea and Black Sea (Gorlach et al. 2017). After the latest glacial maximum, around
11.6 kya, the Baltic Ice Lake drained to the Atlantic, creating a brief phase of brackish-water
Yoldia Sea (Patton et al. 2017) followed by the re-enclosed Ancylus Lake (Bjorck 1995).
Migration of seals between the Atlantic and the Baltic Sea basin has been permanently possible
since the opening of the connection through the Danish strait 8.5 kya (Ukkonen et al. 2014), and
also detectable in the genetic data (Martinez-Bakker et al. 2013).

Taken together, the genomic evidence demonstrates that Saimaa ringed seals represent an
evolutionary lineage with deep roots, clearly distinct from the contemporary Baltic and Ladoga
ringed seals. The M_{50} estimates date this split latest during the Middle Weichselian (Table 1), but
the divergence of the lineages appears to have started significantly earlier (Figure 3C, Table 1).
The split dates provide a plausible scenario where the ancestral population was formed from ice-

dammed marine ringed seals in an eastern or southeastern refugium or possibly in multiple
landlocked refugia. Regardless of the timing and geographical origin, the results support the
hypothesis that the Saimaa ringed seal is the sole remaining representative of proglacial ringed
seals that were mostly replaced elsewhere by the contemporary Arctic ringed seals at the end of
the ice age (Heino et al. 2023). Along with this unique ancestry, the Saimaa lineage also carries
ancestry derived from a postglacial admixture with Atlantic ringed seals. Currently, the land-
locked Saimaa ringed seals are effectively allopatric and thus continuing their distinct
evolutionary trajectory into the foreseeable future.
168

Phenotypic differentiation of the Saimaa ringed seal relating to feeding ecology
Given the deep ancestry of the Saimaa lineage, next we examined phenotypic features that may
relate to feeding ecology differences between marine and freshwater habitats. Previous analyses
have demonstrated morphological features, especially related to the skull that can be used to
differentiate Saimaa ringed seals from other ringed seals (Nordqvist 1899, Hyvärinen and
Nieminen 1990, Sipilä and Hyvärinen 1998, Amano et al. 2002). A short postcanine tooth row
has been reported to characterize the Saimaa ringed seal (Amano et al. 2002) and here we
analyzed the dentition in greater detail. In the seals, the spacing of teeth, hence the length of the
tooth row increases as the individual grows. Thus, as with other cranial and skeletal features,
age-controlled sampling must be used for tooth row length. In contrast, individual tooth
measurements are affected only by wear, rarely an issue with ringed seals as they lack exact
occlusion and do not chew their food. Instead, the laterally compressed postcanines are used for
biting and piercing of food, or retention of food in the oral cavity when the water is expelled
(Loughlin 1982, Adam and Berta 2002, Hocking et al. 2017, Watanabe et al. 2020).
183

First, we compared the sizes of the largest four lower postcanines P_2, P_3, P_4, and P_5. These teeth
are characteristic to species and in ringed seals have typically three to five sharp cusps. The
postcanine lengths of a 326 specimen sample covering Arctic, Okhotsk, Baltic, Ladoga, and
Saimaa show that the P_2, to P_4, fit between the longer Baltic, Ladoga, and Arctic, and the shorter
Okhotsk specimens (Figure 4A, Table 2). However, the last postacanine P_5 (that is actually the
first molar) is distinctly short in Saimaa, even shorter than the P_5 of Okhotsk seals, the smallest
of the studied ringed seals ($P = 0.0000$ to 0.0011, Figure 4A, Table 2). Examining Saimaa tooth
heights reveals that especially their anterior P_2 and P_3 are tall, and their overall height/length
ratios, measuring relative height of the crown profile, are by far the highest among the
populations (Figure 4B). Furthermore, the top-cusp angles, measuring relative height of the
lateral cusps, are the smallest in Saimaa (Figure 4C), further indicating a sharper crown profiles
for the Saimaa ringed seal. This morphological pattern contrast with that of the characteristic
ringed seal morphology observed in the other population samples, irrespective of their tooth size.
197

To examine the diagnostic value of tooth measures in identifying Saimaa ringed seals from the
whole sample, we computed the receiver operating characteristic (ROC) curves for each tooth

(Figure 4D). For the tooth length the areas under the ROC curves, which measure the
performance, are 0.64, 0.56, 0.53, and 0.88 for P_2, P_3, P_4, and P_5, respectively (Figure 4D).
Because the P_4 appears to perform closest to random classification (area is closest to 0.5 and the
line follows the diagonal in Figure 4D), we tested whether it could be used to remove overall size
variation from P_5 by calculating the P_5/P_4 ratio for each specimen. The P_5/P_4 ROC curve shows
even better performance than the P_5 length alone, enclosing 0.92 of the area (Figure 4D). A cut-
off of approximately 0.88 for the P_5/P_4 ratio is indicated by the ROC curve as a diagnostic
threshold value in the identification of Saimaa ringed seal material. Importantly, this size ratio is
not affected by the overall small size of Okhotsk seals, and can also be used irrespective of age
and sex, although females show better separation due to smaller teeth (Figure S2). In addition to
tooth length, both height/length ratios and top-cusp angles of P_5 provide diagnostic value,
enclosing 0.87 and 0.79 of the ROC area, respectively (Figure 4D).
212
Because seals swallow food with little or no processing, their tongue plays an important role in
feeding, especially in species with a suction feeding strategy (Adam and Berta 2002). Although
the availability of soft tissue anatomy is generally limited, next we examined the overall tongue
morphology as it may help to explain the distinctive dental morphology of Saimaa ringed seals
(Table S7, Figure 5A). We observed that the shape of the tip of the tongue differs between the
Saimaa and Baltic ringed seals, and only the former shows intermolar elevations on the lateral
sides of the tongue (asterisks in Figure 5A). The width of the tongue is already over 70% of the
final width at 20% from the tongue tip in the Saimaa specimens, whereas the Baltic specimen has
much pointier shape, reaching 70% of the width closer to 40% from the tongue tip (Figure 5B,
Table S7). The tongue anatomy was recently examined in Antarctic seals and the crabeater (or
the krill-eater seal, *Lobodon carcinophaga*) and the leopard seal (*Hydrurga leptonyx*) of that
study are of special interest in comparison to ringed seals (Loza et al. 2023). The former is a
suction and filter-feeding specialist that uses its complex teeth to retain the krill in the oral
cavity. Its overall tongue shape appears to be an intermediate between the Saimaa and Baltic
ringed seals (Figure 5B). In contrast, the leopard seal tongue is quite similar in overall shape to
that of the Baltic ringed seal (Figure 5B). Whereas leopard seals consume krill much the same
fashion as crabeater seals, they diversify their diet to vertebrate prey as they grow larger,
switching from mainly suction-filter feeding to a grip-and-tear feeding mode.
231
The known diets of ringed seals appear to incipiently align within the extremes of larger bodied
crabeater and leopard seal diets. Ringed seals in the Arctic feed on small fish, typically around
10 cm, and rarely over 20 cm in length (Holst et al. 2001). However, Arctic ringed seals also
consume zooplankton as seals in some regions show a seasonal switch to feeding on amphipod
crustaceans, such as *Parathemisto libellula* (Lowry et al. 1980; Labansen et al. 2011, see also
Geptner et al. 1988). In addition to seasonal feeding, amphipods are also regionally important
dietary source for young seals (Holst et al. 2001). Although data is more limited for the Okhotsk
population, also these seals have been reported to eat planktonic crustaceans in addition to fish

(Fedoseev 1964, Geptner et al. 1988). Apart from the species composition, the Baltic ringed seals
have comparable diets with the Arctic (Suuronen and Lehtonen 2012). In contrast, the Lake
Saimaa is depauperate of large planktonic crustaceans and its seals feed almost entirely on small
fish, such as vendace (*Coregonus albula*), smelt (*Osmerus eperlanus*) and perch (*Perca*
fluviatilis) (Kunnasranta et al. 1999, Auttila et al. 2015).
245
Taken together, the addition of zooplankton to fish diet and the postcanines with well-developed
cuspal comb (Ishihara et al. 2024) in marine ringed seals indicate adaptations to the filter feeding
of zooplankton reminiscent of *Lobodon* and *Hydrurga* (Figure 5C, Table S8). In comparison,
cusp relief needed for filtering small prey is reduced in Saimaa ringed seals (Figure 5C, Figure 4,
Table S8). Rather, Saimaa ringed seals have sharper crown profiles emphasizing the central cusp
at the expense of the lateral cusps, especially in the anterior postcanines (Figure 4). These
morphological features suggest increased specialization to catching, biting and piercing of fish
prey. This dental change is unlikely to be the result of a recent drift due to small population size
because the broadened tongue implies integrated changes in the feeding apparatus. Kinematic
analysis of feeding of marine ringed seals has shown them to be able to suction feed despite the
lack of apparent specializations to suction in the skull morphology (Kienle et al. 2018).
Considering the oral anatomy, we postulate that Saimaa ringed seals should have relatively
strong suction feeding performance. Additional changes linked to feeding are suggested by
subtle differences between jaw musculature between Saimaa and Baltic ringed seals (Laakkonen
and Jernvall 2020), and the length of the intestinal tract of the Saimaa ringed seal being relatively
shorter than those of marine ringed seals (Kunnasranta et al. 1999). Interestingly, the Ladoga
ringed seals have not been documented to feed on zooplankton (Tormosov and Filatov 1979, but
see Geptner et al. 1988), yet their teeth retain the typical ringed seal morphology (Figure 4). This
suggests that the derived oral anatomy of the Saimaa ringed seal reflects a long regime of natural
selection. To this end, the deep evolutionary roots of the Saimaa lineage have led to an opposite
outcome compared to another long-isolated land-locked seal, the Baikal seal (*Pusa sibirica*).
These seals have highly complex teeth, well suited for complementing their fish diet with
suction-filter feeding of Lake Baikal's abundant zooplankton (Watanabe et al. 2004, 2020,
Ishihara et al. 2024).
270

Discussion

Genome wide analyses of species relationships are providing more detailed, and increasingly
more complex view into the history of many mammalian lineages (e.g., Cahill et al. 2018,
Ferreira et al. 2021, Meneganzin and Bernardi 2023, Bertola et al. 2024). This can complicate
taxonomic certainty in delineating species, and approaches integrating multiple lines of evidence
have been advocated (Groves et al. 2016, Kitchener et al. 2022). For example, the traffic-light
system proposed by Kitchener et al. (2022) uses three types of independent evidence:
morphological, genetic and biogeographical. Accordingly, a simple genetic distance alone, even

if large, does not provide taxonomic certainty. Integrative analyses combining multiple lines of
evidence have shown that they can result in a reduction of the number of recognized species
(e.g., van Elst et al. 2024), but also in identification of new taxa (e.g. Nater et al. 2017, Carroll et
al. 2021). Given that the Lake Saimaa has formed a geographic barrier for dispersal during the
last 10 kya, here we focused on genetic and morphological evidence to examine whether the
ancestry of the Saimaa ringed seals could be older than the Lake Saimaa.
285
Our genetic analyses confirm that the seals in Lake Saimaa are not only clearly different from
other ringed seals, but importantly, also have an independent and deep evolutionary history. The
Saimaa lineage has twice as distant split times to the three Atlantic ringed seal subspecies as
those have amongst them. Similarly, the Saimaa population has the greatest fraction of unique
SNPs among the ringed seal populations, and the PCA component separating the Saimaa
individuals from all the ringed seals explains nearly three times the amount of variance that the
second component separating the four remaining ringed seal subspecies. Additionally, our
analyses reveal a brief period of more recent contact between the Saimaa lineage and the
lineages forming the current Ladoga and Baltic populations around 8–10 kya. The existence of
this kind of intermittent gene flow is a recurrent observation of genetic admixture and
introgression among many *bona fide* species in the wild (Huerta-Sanchez et al. 2014, Cahill et al.
2018, Ferreira et al. 2021, Bertola et al. 2024).
298
The deep genetic origin of the Saimaa lineage is further underscored by the morphological data.
The distinct skull morphology of Saimaa ringed seals has been long recognized (e.g. Amano et al
2002), and here we demonstrate that their dental and tongue morphologies are indicative of
specialization on feeding exclusively on fish. This specialized feeding niche differs from the
other ringed seals that have more complex dental morphologies reflecting the ability to include
zooplankton in their diet. Taken together, given the combination of genomic and phenotypic
evidence, we propose that the Saimaa ringed seal is evolutionary more unique than previously
thought.
307
The genetic differences and the timing of the population splits (Table 1, Figure 3C) are
consistent with the scenario where the ancestral Saimaa population was formed from ice-
dammed marine ringed seals in an eastern or southeastern refugium or possibly in multiple
refugia latest during the Middle Weichselian, 90–50 kya ago, when large glacial lakes existed in
the West Siberian Plains and White Sea Basin with water routes to the eastern edge of the
Fennoscandian ice sheet (Mangerud et al. 2001, 2004, Krinner et al. 2004). We note that
although ancient, as a landlocked marine species derived from eastern proglacial refugia, Saimaa
ringed seal is not unique. Relict marine fish and invertebrate species are found in modern Lake
Saimaa and eastern lakes around the Baltic Sea basin (Savolainen 1975, Segerstrale 1976,
Väinölä et al. 1994, 2001, Audzijonyte and Väinölä 2005, Säisä et al. 2005). It is conceivable

that many of the other relicts may also have more ancient origins than previously thought,
harboring unique evolutionary history (e.g. Rosing-Asvid et al. 2023).
320
Although Nordqvist (1899) originally described the Saimaa and Ladoga ringed seals, he also
advocated for the extermination of both populations, due to their claimed nuisance to local
fisheries (Nordqvist 1892). His goal of eradicating the Saimaa population was nearly realized,
with fewer than 100 individuals remaining when the species finally gained legal protection in
1955 (Kunnasranta et al. 2021). The population has gradually recovered since then, reaching
almost 500 individuals in 2024 (https://www.metsa.fi/en/nature-and-heritage/species/saimaa-
ringed-seal/). As ringed seals depend on snow lairs for birthing and nursing, ongoing
anthropogenic climate change presents a significant long-term threat (Jakkila et al. 2024).
However, the Saimaa ringed seal has survived the previous post-glacial warm period and is
likely able to do so in the future, if given the chance.
331

Materials and methods

Material examined

DNA samples from ringed seals were obtained from 12 unique sampling localities in the
northern hemisphere, greatly improving the coverage in the eastern regions compared to the
previous analyses [Figure 1, distribution ranges drawn after Kelly et al. (2010)]. The sources of
the new samples for Baltic and Ladoga are the same as in (Löytynoja et al. 2023). Alaska
samples are from Prudhoe Bay (001B98, AF7110), Peard Bay (G003), and Bering Sea TV13-
15); Chukchi Sea samples are from Lorino; Okhotsk Sea samples are mainly from Chkalov
Island and Odyan Bay (TV16-18); Pechora Sea samples are from Vaigach Island; and White Sea
samples from Belomorsk archipelago. Samples are from bycaught seals or dead beachcast
animals collected for research. Based on earlier analyses (Löytynoja et al. 2023, Rosing-Asvid et
al. 2023), the whole-genome-sequenced individuals were ranked by quality (maximizing the
sequencing coverage and data completeness) and at most five individuals from each population
were included for full genome analyses (in total 46 samples). Four spotted seals (*Phoca largha*;
accession numbers SAMN08238620, SAMN16895771, SAMN31577600 and SAMN31577601;
Park et al. 2018, Dudchenko et al. 2017, Xu et al. 2024) were included for SNP polarization. The
mitochondrial DNA (mtDNA) was analyzed separately and 162 unique ringed seal haplotypes
and the four spotted seal genomes were included. For the dental data, Saimaa material are from
the University of Eastern Finland (UEF) and Metsähallitus (the Finnish Forest Administration),
Ladoga material from Finnish Museum of Natural History (FMNH), University of Helsinki,
University of Eastern Finland and Metsähallitus, Baltic material from Finnish Museum of
Natural History (FMNH), Arctic Greenland material (East and West Greenland, including
Canadian Arctic Archipelago) from National History Museum of Denmark (NHMD), Arctic
Alaska material (Beaufort, Chukchi, and Bering Sea) from University of Alaska Museum of the
North (UAM), and Arctic Kara Sea and Sea of Okhotsk material from the National Museum of

Nature and Science of Tokyo (NSMT). The four distal postcanines show the largest range of
variation among phocid species, and their intra- and interspecies variation have been studied
extensively previously (Jernvall 2000, Savriama et al. 2018, Ishihara et al. 2024). Additionally,
we investigated soft tissue anatomy of tongues from freshly defrozen seal cadavers (*N*=8, from
Metsähallitus (the Finnish Forest Administration). The type specimens are presented in a
separate chapter. Comparative dental specimens to calculate cusp relief (Table S8) originated
from the collections of the Finnish Museum of Natural History (FMNH), the Swedish Museum
of Natural History (NRM), and the National Museum of Victoria, Melbourne (NMV).
365

Genomic analyses

Data mapping and nuclear genome variant calling
The DNA sequencing was performed on Illumina NextSeq500 and NovaSeq 6000 platforms
(Löytynoja et al. 2023, Sundell et al. 2023). Read data for ringed seal samples and four spotted
seal samples were aligned as described previously (Olkkonen and Löytynoja 2023). For the
selected 46 ringed seals and four spotted seals, variants were called by applying GATK4
(v.4.2.5) (Poplin et al. 2018) HaplotypeCaller for each sample, joining the data with
CombineGVCFs, and applying GenotypeGVCFs to the joint data. The sequencing depth of the
studied ringed seal samples varied from 5.0x to 29.5x with the mean coverage of 10.6x, and of
spotted seal samples from 34.8x to 115.7x with the mean coverage of 55.7x.
376

Mitogenome analyses
Variants within the mitochondrial genome was called from all sequenced individuals applying
the same GATK4 tools but defining the ploidy 1. Protein-coding regions were inferred by
mapping the peptide sequences of the *Pusa hispida* mitochondrion (NC_008433) against the
reference mtDNA contig with miniprot (Li, 2023). Binary SNP variants, totalling 1,434, within
the CDS regions were extracted and samples with missing data were discarded. The VCF data
were converted to FASTA sequences using the vcf-to-tab tool from the VCFtools package
(Danecek et al. 2011) and the script from https://github.com/JinfengChen/vcf-tab-to-fasta
modified to work on haploid data. Identical haplotypes (all within a subspecies) were removed,
leaving 162 ringed seals and four spotted seals as the outgoup. The phylogenetic tree was
inferred with RAxML v.8.2.12 (Stamatakis, 2014) using the model ASC_GTRGAMMA and
correcting for the invariable sites (totalling 9,945) missing from the VCF file with the option --
asc-corr=felsenstein. For the search, the rapid option combining the ML search (20 replicates)
and the bootstrap analysis (1000 replicates) was selected (-f a -# 1000). The resulting tree was
visualised in R using the ggtree package (Yu et al. 2017).
392

Data filtering and SNP analyses
The binary SNP data for the full nuclear genomes were filtered and thinned using BCFtools
(v.1.9) (Danecek et al. 2021) and Plink (v.1.9) (Chang et al. 2015). Sites falling outside the
positively masked regions or inside the repeat-masked regions (Olkkonen and Löytynoja 2023),

showing no variation among ringed seals or having >10% of missing data were excluded. The
remaining data, totalling 25,697,406 sites, were further thinned by requiring sites to be at least 1
kbp apart, leaving 1,807,907 variant sites.

A principal component analysis was performed on the thinned data (1.8M sites) with smartpca
(v.16000)(Price, et al. 2006) and the results were visualized with R (R-Core-Team 2020) using
the ggplot2 (Wickham 2016) package. In smartpca, the normalization option (usenorm) was
turned off and the analysis was performed for the full 46 individuals set and for an even
sampling of five individuals per subspecies (Elhaik, 2022), selected based on data completeness
and, in the case of the Arctic subspecies, picking individuals from different geographic locations
(Alaska, Svalbard, White Sea, Pechora Sea, Chukchi Sea). Starting from the 1.8M sites data,
allele-sharing among different subspecies was studied using the same five individuals per
subspecies, including the four spotted seals for derived allele inference. With samples varying
from one to five individuals, the presence of a derived allele in each ringed seal subspecies was
recorded, and the proportion of derived allele patterns unique to the specific subspecies was
calculated. The statistic is highly dependent on the MAF limit applied. With full data (MAF=0),
a large proportion of unique alleles are singletons and reflect the recent demographic events and
the current effective population size. We applied MAF=0.05 (requiring at least five derived
alleles among the 92 sampled genomes in the full data) to understand the more distant
demographic events; however, one cannot rule out the founder effect and recent genetic drift
inflating the Saimaa statistics. The proportion of private variants is expected to decrease with the
number of individuals sampled and, in theory, plateau and stop when all individuals are included.
The allele frequencies in each subset were computed with VCFtools and processed with bash and
R scripts.

Demographic analyses
Demographic analyses were performed with MSMC-IM (Wang et al. 2020) using the whole-
genome data from Löytynoja et al. (2023). For within-population analyses, ten two-individual
sets (i.e., four chromosomes) were randomly chosen from each population and their coalescent
rate histories were computed as explained in Löytynoja et al. (2023). For cross-population
coalescence rate analyses, six pairs of individuals for each population combination (i.e., two
chromosomes from each population, two populations per analysis) were randomly chosen and
their coalescent rate histories were computed using the default time segmenting. The combined
rate estimates were analysed using MSMC_IM.py with the recommended parameters and the
mutation rate (μ) 1.826e-8 estimated for polar bear (Liu et al. 2014). We consider this a robust
estimate as it falls within the mutation rates estimated for different seals (Peart et al. 2020), and
using different estimates does not affect the overall pattern of results. The summary results were
computed and visualized with R using the ggplot2 package and assuming a generation time (g)
of 10 years.

Phenotypic analyses

Lower tooth rows were photographed from the lingual side and maximum tooth lengths, maximum heights, top-cusp angles, and cusps numbers were tabulated using Fiji (Schindelin et al. 2012) and additional analyses were done with PAST (Hammer et al. 2001). Although the dentitions of phocids show high degree of intraspecies variation, the seals are born with their permanent tooth crowns fully formed and in the process of erupting into the oral cavity. Therefore, in contrast to other skeletal features, dental measures do not change after birth. We report the analyses for the four, right side postcanines (P_2, P_3, P_4, P_5), and only for individual specimens from which the measures could be obtained for all the four teeth. We excluded specimens with dental anomalies such as supernumerous teeth. Cusp number and tooth length were measured for 326 individuals and crown height and top-cusp angle for 281 individuals due to cracked or slightly worn specimens. Height and angle measurements were estimated for cases where only the enamel cap of the cusp tip was missing. Exclusion of these data did not alter the results. Cusp were tabulated regardless how small they are. Relative cusp height was measured using the top-cusp angle (Jernvall 2000, Salazar-Ciudad and Jernvall 2010) and the cusp relief as the length of the crown perimeter divided by the length of the tooth at the crown base. *P*-values in Figure 3 and Table 2 are one-tailed and obtained between group means using 10 000 permutations. The receiver operating characteristic (ROC) curves were used to obtain correct classification probabilities. For tooth lengths and top-cusp angles, smaller values denote true positive. Because seal tongue plays an important role in feeding, especially in species with a suction feeding strategy (Hocking et al. 2017, Loza, et al. 2023), we compared the overall morphology of Saimaa ringed seal tongue with that of phocids from the literature (Loza, et al. 2023). Tongues from cadavers were photographed in defrosted stage and compared to those in the literature. To obtain a robust measure of shape apart from size and variation caused by differential preservation of soft tissue anatomy, were divided tongue images into ten-percent bins from the tip to the end of the posterior body, and tabulated widths as relative to the maximum width of the tongue.

Data availability

Most of the genome data have been previously used and are available at the European Nucletiode Archive (ENA) under accession numbers PRJEB56329 and PRJEB56317. The newly generated 16 whole-genome sequencing datasets are deposited under PRJEB56317 with accession numbers ERS23016468-ERS23016483. Instructions for replicating the computational analyses are provided at https://github.com/ariloytynoja/pusa_saimensis

Acknowledgements

For tissue samples, we thank the University of Eastern Finland and Metsähallitus (Saimaa); M. Verevkin (Ladoga); P. Timonen, S. Oksanen, and J. Vierimaa (Baltic); D. Tallmon, A. Rosing-Asvid, S. Ferguson, R. Dietz, M. Zasypkin, P. Strelkov, N. Medvedev and particularly O. Shpak

(Arctic). For access to dental material, we thank the curators of Finnish Museum of Natural History, National History Museum of Denmark, the National Museum of Victoria (Melbourne), the Swedish Museum of Natural History, and University of Alaska Museum of the North (Fairbanks). For advice on taxonomy and analyses we thank S. Viranta, K. Leppälä, and X.-Y. Feng. We thank the personnel of the DNA sequencing and genomics laboratory for performing the NGS samples preparation and sequencing, and the CSC – IT Center for Science for the computational resources. The Jane and Aatos Erkko Foundation (JJ and PA), the LIFE Programme of the European Commission (PA, LIFE19NAT/FI/000832), Research Council of Finland (JJ) are acknowledged for their financial support.

References

Adam, P. J. and Berta A. (2002). Evolution of prey capture strategies and diet in the Pinnipedimorpha (Mammalia, Carnivora). *Oryctos 4*, 83-107.

Amano, M., Hayano, A. and Miyazaki, N. (2002). Geographic variation in the skull of the ringed seal. *Journal of Mammalogy 83*(2), 370-380.

Audzijonytė, A. and Väinölä, R. (2005). Diversity and distributions of circumpolar fresh- and brackish-water *Mysis* (Crustacea: Mysida): descriptions of *M. relicta* Lovén, 1862, *M. salemaai* n. sp., *M. segerstralei* n. sp. *and M. diluviana* n. sp., based on molecular and morphological characters. *Hydrobiologia* 544, 89-141.

Auttila, M., Kurkilahti, M., Niemi, M., Levänen, R., Sipilä, T., Isomursu, M., Koskela, J. and Kunnasranta, M. (2016). Morphometrics, body condition, and growth of the ringed seal (Pusa hispida saimensis) in Lake Saimaa: Implications for conservation. *Marine Mammal Science*, *32*(1), 252-267.

Auttila, M., Sinisalo, T., Valtonen, M., Niemi, M., Viljanen, M., Kurkilahti, M., and Kunnasranta, M. (2015). Diet composition and seasonal feeding patterns of a freshwater ringed seal (*Pusa hispida saimensis*). *Marine Mammal Science, 31*(1), 45-65.

Avise, J.C., and Ball-Jr, R.M. (1990). Principles of genealogical concordance in species concepts and biological taxonomy. *Oxford Surveys in Evolutionary Biology 7*, 45–67.

Berta, A. and Churchill M. (2012). Pinniped taxonomy: review of currently recognized species and subspecies, and evidence used for their description. *Mammal Review 42*(3), 207-234.

Bertola, L. D., Quinn, L., Hanghøj, K., Garcia-Erill, G., Rasmussen, M. S., Balboa, R. F., ... and Heller, R. (2024). Giraffe lineages are shaped by major ancient admixture events. *Current Biology*, *34*(7), 1576-1586.

Bjorck, S. (1995). A Review of the History of the Baltic Sea, 13.0-8.0 Ka Bp. *Quaternary International 27*, 19-40.

Burns, J. J. and Fay, F. H. (1970). Comparative morphology of the skull of the ribbon seal, *Histriophoca fasciata*, with remarks on the systematics of Phocidae. *Journal of Zoology 161*, 363-394.

Cahill, J. A., Heintzman, P. D., Harris, K., Teasdale, M. D., Kapp, J., Soares, A. E., ... and Shapiro, B. (2018). Genomic evidence of widespread admixture from polar bears into brown bears during the last ice age. *Molecular Biology and Evolution*, *35*(5), 1120-1129.

Carroll, E. L., McGowen, M. R., McCarthy, M. L., Marx, F. G., Aguilar, N., Dalebout, M. L., ... and Olsen, M. T. (2021). Speciation in the deep: genomics and morphology reveal a new species of beaked whale *Mesoplodon eueu*. *Proceedings of the Royal Society B, 288(1961)*, 20211213.

Chang, C. C., Chow, C. C., Tellier, L. C., Vattikuti, S., Purcell, S. M., and Lee, J. J. (2015). Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience*, *4*(1), s13742-015.

Conway, J. R., Lex, A., and Gehlenborg, N. (2017). UpSetR: an R package for the visualization of intersecting sets and their properties. *Bioinformatics*, *33*(18), 2938-2940.

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., ... and Li, H. (2021). Twelve years of SAMtools and BCFtools. *Gigascience*, *10*(2), giab008.

Dudchenko, O., Batra, S. S., Omer, A. D., Nyquist, S. K., Hoeger, M., Durand, N. C., Shamim, M. S., Machol, I., Lander, E. S., Aiden, A. P., and Aiden, E. L. (2017). *De novo* assembly of the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. *Science 356*, 92–95.

Elhaik, E. (2022). Principal Component Analyses (PCA)-based findings in population genetic studies are highly biased and must be reevaluated. *Scientific Reports, 12*(1), 14683.

Ellenberg, J. R., and Morrison-Scot, T. C. S. (1951). Checklist of Palearctic and Indian mammals. *British Museum (Natural History), London*.

Fedoseev, G. A. (1964). On the embryonic and postembryonic growth and sexual maturation in the Okhotsk Sea Ringed Seal. *Zoologicheskii Zhurnal 43*(8), 1228-1235. [In Russian.]

Van Elst, T., Sgarlata, G. M., Schüßler, D., Tiley, G. P., Poelstra, J. W., Scheumann, M., ... and Salmona, J. (2024). Integrative taxonomy clarifies the evolution of a cryptic primate clade. *Nature Ecology & Evolution*, 1-16.

Ferreira, M. S., Jones, M. R., Callahan, C. M., Farelo, L., Tolesa, Z., Suchentrunk, F., ... and Melo-Ferreira, J. (2021). The legacy of recurrent introgression during the radiation of hares. *Systematic biology*, *70*(3), 593-607.

Geptner, V. G., Nasimovich, A. A., Bannikov, A. G., and Hoffmann, R. S. (1988). Mammals of the Soviet Union. v. 2, pt. 3, Smithsonian Institution Libraries and National Science Foundation.

Gorlach, A., Hang, T., and Kalm, V. (2017). GIS-based reconstruction of Late Weichselian proglacial lakes in northwestern Russia and Belarus. *Boreas*, *46*(3), 486-502.

Groves, C. P., Cotterill, F. P., Gippoliti, S., Robovský, J., Roos, C., Taylor, P. J., and Zinner, D. (2017). Species definitions and conservation: a review and case studies from African mammals. *Conservation Genetics*, *18*, 1247-1256.

Hammer, Ø., Harper, D. A. T. and Ryan, P. D. (2001). PAST: Paleontological statistics software package for education and data analysis. *Palaeontol. Electron*. *4*, 9pp.

Heino, M. T., Nyman, T., Palo, J. U., Harmoinen, J., Valtonen, M., Pilot, M., ... and Aspi, J. (2023). Museum specimens of a landlocked pinniped reveal recent loss of genetic diversity and unexpected population connections. *Ecology and Evolution*, *13*(1), e9720.

Hocking, D. P., Marx, F. G., Park, T., Fitzgerald, E. M., and Evans, A. R. (2017). A behavioural framework for the evolution of feeding in predatory aquatic mammals. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1850), 20162750.

Holst, M., Stirling, I., and Hobson, K. A. (2001). Diet of ringed seals (*Phoca hispida*) on the east and west sides of the North Water Polynya, northern Baffin Bay. *Marine Mammal Science*, *17*(4), 888-908.

Huerta-Sánchez, E., Jin, X., Asan, Bianba, Z., Peter, B. M., Vinckenbosch, N., Liang, Y., Yi, X., He, M., Somel, M., Ni, P., Wang, B., Ou, X., Huasang, Luosang, J., Cuo, Z. X. P., Li, K., Gao, G., Yin, Y., … and Nielsen, R. (2014). Altitude adaptation in Tibetans caused by introgression of Denisovan-like DNA. *Nature*, *512*(7513), 194-197.

Hyvärinen, H. and Nieminen, M. (1990). Differentiation of the ringed seal in the Baltic sea, lake Ladoga USSR and lake saimaa Finland. *Finnish Game Research 47*, 21-27.

Ishihara, U., Miyazaki, N., Yurkowski, D. J., and Watanabe, Y. Y. (2024). Multi-cusped postcanine teeth are associated with zooplankton feeding in phocid seals. *Marine Ecology Progress Series*, *729*, 233-245.

Jakkila, J., Auttila, M., Tuukkanen, T., and Veijalainen, N. (2024). Modelling climate change impacts on lake ice and snow demonstrates breeding habitat decline of the endangered Saimaa ringed seal. *Climatic Change*, *177*(9), 134.

Jernvall, J. (2000). Linking development with generation of novelty in mammalian teeth. *Proceedings of the National Academy of Sciences*, *97*(6), 2641-2645.

Kitchener, A. C., Hoffmann, M., Yamaguchi, N., Breitenmoser-Würsten, C., and Wilting, A. (2022). A system for designating taxonomic certainty in mammals and other taxa. *Mammalian Biology*, *102*(1), 251-261.

Kelly, B. P., Bengtson, J. L., Boveng, P. L., Cameron, M. F., Dahle, S. P., Jansen, J. K., Logerwell, E. A., Overland, J. E., Sabine, C. L., Waring, G. T., and Wilder, J. M. (2010). Status review of the ringed seal (*Phoca hispida*). *NOAA technical memorandum NMFS-AFSC, 212*. https://repository.library.noaa.gov/view/noaa/3762

Kelly, B. P. (2022). The ringed seal: behavioral adaptations to seasonal ice and snow cover. In *Ethology and behavioral ecology of phocids* (pp. 553-597). Cham: Springer International Publishing.

Kienle, S. S., Hermann-Sorensen, H., Costa, D. P., Reichmuth, C., and Mehta, R. S. (2018). Comparative feeding strategies and kinematics in phocid seals: suction without specialized skull morphology. *Journal of Experimental Biology*, *221*(15), jeb179424.

Kovacs, K. M., Aguilar, A., Aurioles, D., Burkanov, V., Campagna, C., Gales, N., ... & Trillmich, F. (2012). Global threats to pinnipeds. *Marine Mammal Science*, *28*(2), 414-436.

Krinner, G., Mangerud, J., Jakobsson, M., Crucifix, M., Ritz, C. and Svendsen, J. I. (2004). Enhanced ice sheet growth in Eurasia owing to adjacent ice-dammed lakes. *Nature* *427*(6973), 429-432.

Kunnasranta, M. (2001). Behavioural biology of two ringed seal (*Phoca hispida*) subspecies in the large European lakes Saimaa and Ladoga. Joensuu, Finland, University of Joensuu.

Kunnasranta, M., Hyvärinen, H., Sipilä, T., and Koskela, J. T. (1999). The diet of the Saimaa ringed seal Phoca hispida saimensis. *Acta Theriologica*, *44*(4), 443-450.

Kunnasranta, M., Niemi, M., Auttila, M., Valtonen, M., Kammonen, J., and Nyman, T. (2021). Sealed in a lake—Biology and conservation of the endangered Saimaa ringed seal: A review. *Biological Conservation*, *253*, 108908.

Laakkonen, J. and Jernvall, J. (2020). Muscles of mastication and the temporo-mandibular joint of the Saimaa (*Pusa hispida saimensis*) and Baltic (*Pusa hispida botnica*) ringed seals. *Annales Zoologici Fennici 57*(1-6), 21-29.

Labansen, A. L., Lydersen, C., Levermann, N., Haug, T., and Kovacs, K. M. (2011). Diet of ringed seals (*Pusa hispida*) from Northeast Greenland. *Polar Biology*, *34*, 227-234.

Li, H. (2023). Protein-to-genome alignment with miniprot. *Bioinformatics*, *39*(1). https://doi.org/10.1093/bioinformatics/btad014

Lipson, M. (2020). Applying f4 statistics and admixture graphs: Theory and examples. *Molecular Ecology Resources*, *20*(6), 1658-1667.

Liu, S., Lorenzen, E. D., Fumagalli, M., Li, B., Harris, K., Xiong, Z., ... and Wang, J. (2014). Population genomics reveal recent speciation and rapid evolutionary adaptation in polar bears. *Cell*, *157*(4), 785-794.

Lowry, L. F., Frost, K. J., and Burns, J. J. (1980). Variability in the diet of ringed seals, *Phoca hispida*, in Alaska. *Canadian Journal of Fisheries and Aquatic Sciences*, *37*(12), 2254-2261.

Loughlin, T. R. (1982). Functional adaptation of eruption sequence of pinniped postcanine teeth. *Journal of Mammalogy*, *63*(3), 523-525.

Löytynoja, A., Rastas, P., Valtonen, M., Kammonen, J., Holm, L., Olsen, M. T., Paulin, L., Jernvall, J. and Auvinen, P. (2023). Fragmented habitat compensates for the adverse effects of genetic bottleneck. *Current Biology*, *33*(6), 1009-1018.

Loza, C. M., Carlini, A. A., Zarza, R., Krmpotic, C. M., Galliari, F. C., Andrés-Laube, F., ... and Barbeito, C. G. (2023). Description and distribution of the mechanical papillae of the lingual surface of Antarctic seals species (Phocidae: Carnivora) and their relationship with diet and dental morphology. *Polar Biology*, *46*(3), 235-251.

Mangerud, J., Astakhov, V., Jakobsson, M., and Svendsen, J. I. (2001). Huge Ice age lakes in Russia. *Journal of Quaternary Science,*, *16*(8), 773-777.

Mangerud, J., Jakobsson, M., Alexanderson, H., Astakhov, V., Clarke, G. K., Henriksen, M., ... and Svendsen, J. I. (2004). Ice-dammed lakes and rerouting of the drainage of northern Eurasia during the Last Glaciation. *Quaternary Science Reviews*, *23*(11-13), 1313-1332.

Martinez-Bakker, M. E., Sell, S. K., Swanson, B. J., Kelly, B. P., and Tallmon, D. A. (2013).
Combined genetic and telemetry data reveal high rates of gene flow, migration, and long-
distance dispersal potential in Arctic ringed seals (*Pusa hispida*). *PLoS One*, *8*(10),
e77125.
McLaren, I. A. (1958). *The biology of the ringed seal (Phoca hispida Schreber) in the eastern*
Canadian Arctic (Vol. 118). Ottawa: Fisheries Research Board of Canada.
Meneganzin, A., and Bernardi, M. (2023). Were Neanderthals and *Homo sapiens* 'good
species'?. *Quaternary Science Reviews*, *303*, 107975.
Nater, A., Mattle-Greminger, M. P., Nurcahyo, A., Nowak, M. G., De Manuel, M., Desai, T., ...
and Krützen, M. (2017). Morphometric, behavioral, and genomic evidence for a new
orangutan species. *Current Biology*, *27*(22), 3487-3498.
Nordqvist, O. (1892). Hylkeen häwittäminen Itämerestä, Laatokasta ja Saimaasta. <u>*Suomen*</u>
<u>*Kalastuslehti*</u> 4, 56-60.
Nordqvist, O. (1899). Beitrag zur kenttniss der isolierten formen der ringelrobbe (*Phoca foetida*
Fabr.) [Contribution to the knowledge of the isolated forms of the ringed seal (*Phoca*
foetida Fabr.)]. *Acta Societatis pro Fauna et Flora Fennica* XV, 1– 47.
Nyman, T., Valtonen, M., Aspi, J., Ruokonen, M., Kunnasranta, M. and Palo, J. U. (2014).
Demographic histories and genetic diversities of F ennoscandian marine and landlocked
ringed seal subspecies. *Ecology and Evolution*, *4*(17), 3420-3434.
Olkkonen, E. and Löytynoja, A. (2023). Analysis of population structure and genetic diversity in
low-variance Saimaa ringed seals using low-coverage whole-genome sequence data. *STAR*
protocols, *4*(4), 102567.
Palo, J. U., Hyvärinen, H., Helle, E., Mäkinen, H. S. and Väinölä, R. (2003). Postglacial loss of
microsatellite variation in the landlocked Lake Saimaa ringed seal. *Conservation Genetics*,
4, 117-128.
Palo, J. U., Mäkinen, H. S., Helle, E., Stenman, O. and Väinölä, R. (2001). Microsatellite
variation in ringed seals (*Phoca hispida*): genetic structure and history of the Baltic Sea
population. *Heredity*, *86*(5), 609-617.
Park, J. Y., Kim, K., Sohn, H., Kim, H. W., An, Y. R., Kang, J. H., ... and Kim, H. (2018).
Deciphering the evolutionary signatures of pinnipeds using novel genome sequences: the
first genomes of *Phoca largha*, *Callorhinus ursinus*, and *Eumetopias jubatus*. *Scientific*
Reports, *8*(1), 16877.
Patton, H., Hubbard, A., Andreassen, K., Auriac, A., Whitehouse, P. L., Stroeven, A. P.,
Shackleton, C., Winsborrow, M., Heyman, J. and Hall, A. M. (2017). Deglaciation of the
Eurasian ice sheet complex. *Quaternary Science Reviews*, *169*, 148-172.
Peart, C. R., Tusso, S., Pophaly, S. D., Botero-Castro, F., Wu, C. C., Aurioles-Gamboa, D., ...
and Wolf, J. B. (2020). Determinants of genetic variation across eco-evolutionary scales in
pinnipeds. *Nature Ecology & Evolution*, *4*(8), 1095-1104.

Poplin, R., Ruano-Rubio, V., DePristo, M. A., Fennell, T. J., Carneiro, M. O., Van der Auwera,
670   G. A., ... and Banks, E. (2017). Scaling accurate genetic variant discovery to tens of
671   thousands of samples. *BioRxiv*, 201178.

672   Price, A. L., Patterson, N. J., Plenge, R. M., Weinblatt, M. E., Shadick, N. A. and Reich, D.
673   (2006). Principal components analysis corrects for stratification in genome-wide
674   association studies. *Nature genetics*, *38*(8), 904-909.

675   R-Core-Team (2020). R: A l,anguage and environment for statistical computing. R Foundation
676   for Statistical Computing. http://www.R-project.org/.

677   Reich, D., Thangaraj, K., Patterson, N., Price, A. L. and Singh, L. (2009). Reconstructing Indian
678   population history. *Nature*, *461*(7263), 489-494.

679   Rosing Asvid, A., Löytynoja, A., Momigliano, P., Hansen, R. G., Scharff Olsen, C. H.,
680   Valtonen, M., ... and Tange Olsen, M. (2023). An evolutionarily distinct ringed seal in the
681   Ilulissat Icefjord. *Molecular ecology*, *32*(22), 5932-5943.

682   Säisä, M., Koljonen, M. L., Gross, R., Nilsson, J., Tähtinen, J., Koskiniemi, J. and Vasemägi, A.
683   (2005). Population genetic structure and postglacial colonization of Atlantic salmon
684   (Salmo salar) in the Baltic Sea area based on microsatellite DNA variation. *Canadian*
685   *Journal of Fisheries and Aquatic Sciences*, *62*(8), 1887-1904.

686   Salazar-Ciudad, I. and Jernvall, J. (2010). A computational model of teeth and the developmental
687   origins of morphological variation. *Nature*, *464*(7288), 583-586.

688   Savolainen, E. (1975). Distribution and food of *Myoxocephalus quadricornis* (L.) (Teleostei,
689   Cottidae) in fresh waters of eastern Finland. *Annales Zoologici Fennici 12*, 271-274.

690   Savriama, Y., Valtonen, M., Kammonen, J. I., Rastas, P., Smolander, O. P., Lyyski, A., ... and
691   Jernvall, J. (2018). Bracketing phenogenotypic limits of mammalian hybridization. *Royal*
692   *society open science*, *5*(11), 180903.

693   Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., ... and
694   Cardona, A. (2012). Fiji: an open-source platform for biological-image analysis. *Nature*
695   *methods*, *9*(7), 676-682.

696   Schreber, J. C. D. (1776). <u>Die Säugthiere in Abbildungen nach der Natur, mit Beschreibungen.</u>
697   Erlangen, Wolfgang Walther.

698   Segerstrale, S. G. (1976). Immigration of glacial relicts into northern Europe. *Boreas*, *5*(1), 1-7.

699   Sipilä, T., Helle, E., and Hyvärinen, H. (1990). Distribution, population size and reproductivity
700   of the Saimaa ringed seal (*Phoca hispida saimensis* Nordq.) in Finland, 1980–84. *Finnish*
701   *Game Research*, *47*, 3-10.

702   Sipilä, T. and Hyvärinen, H. (1998). Status and biology of Saimaa (*Phoca hispida saimensis*) and
703   Ladoga (*Phoca hispida ladogensis*) ringed seals. *NAMMCO Scientific Publications*, *1*, 83-
704   99.

705   Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of
706   large phylogenies. *Bioinformatics*, *30*(9), 1312-1313.

707   Sundell, T., Kammonen, J. I., Mustanoja, E., Biard, V., Kunnasranta, M., Niemi, M., ... and
708   Auvinen, P. (2023). Genomic evidence uncovers inbreeding and supports translocations in

rescuing the genetic diversity of a landlocked seal population. *Conservation Genetics*,
24(2), 155-165.

Suuronen, P. and Lehtonen, E. (2012). The role of salmonids in the diet of grey and ringed seals
in the Bothnian Bay, northern Baltic Sea. *Fisheries Research*, *125*, 283-288.

Tormosov D. D. and Filatov I. J. 1979. Ladozhskaya nerpaob'ekt issledovaniya. Prioda 3, 103-
107.

Ukkonen, P., Aaris-Sørensen, K., Arppe, L., Daugnora, L., Halkka, A., Lõugas, L., Oinonen,
M.J., Pilot, M. and Storå, J. (2014). An Arctic seal in temperate waters: History of the
ringed seal (*Pusa hispida*) in the Baltic Sea and its adaptation to the changing environment.
The Holocene, *24*(12), 1694-1706.

Väinölä, R., Riddoch, B. J., Ward, R. D. and Jones, R. I. (1994). Genetic zoogeography of the
Mysis relicta species group (Crustacea: Mysidacea) in northern Europe and North
America. *Canadian Journal of Fisheries and Aquatic Sciences*, *51*(7), 1490-1505.

Väinölä, R., Vainio, J. K., & Palo, J. U. (2001). Phylogeography of" glacial relict"
Gammaracanthus (Crustacea, Amphipoda) from boreal lakes and the Caspian and White
seas. *Canadian Journal of Fisheries and Aquatic Sciences*, *58*(11), 2247-2257.

Valtonen, M., Palo, J. U., Aspi, J., Ruokonen, M., Kunnasranta, M. and Nyman, T. (2014).
Causes and consequences of fine-scale population structure in a critically endangered
freshwater seal. *BMC ecology*, *14*, 1-15.

Valtonen, M., Palo, J. U., Ruokonen, M., Kunnasranta, M. and Nyman, T. (2012). Spatial and
temporal variation in genetic diversity of an endangered freshwater seal. *Conservation
Genetics*, *13*, 1231-1245.

Wang, K., Mathieson, I., O'Connell, J. and Schiffels, S. (2020). Tracking human population
structure through time from whole genome sequences. *PLoS genetics*, *16*(3), e1008552.

Watanabe, Y., Baranov, E. A., Sato, K., Naito, Y. and Miyazaki, N. (2004). Foraging tactics of
Baikal seals differ between day and night. *Marine Ecology progress series*, *279*, 283-289.

Watanabe, Y. Y., Baranov, E. A. and Miyazaki, N. (2020). Ultrahigh foraging rates of Baikal
seals make tiny endemic amphipods profitable in Lake Baikal. *Proceedings of the National
Academy of Sciences*, *117*(49), 31242-31248.

Wickham, H. (2016). ggplot2: Elegant graphics for data analysis., Springer International
Publishing.

Xu, S., Xie, X., Zhao, R., Tian, J. and Lu, Z. (2024). Genomic comparison of male and female
spotted seals in the Liaodong Gulf, China. *Regional Studies in Marine Science*, 103591.

Figures and legends

Figure 1. The distribution and sampling of the studied ringed seals and their distribution ranges.
(A) The nominal subspecies of the Arctic ringed seal (*Pusa hispida hispida*) has circumpolar
distribution while the four other subspecies are geographically restricted. (B) The ringed seals
living in the Fennoscandia are made of landlocked populations in the Lake Saimaa and Lake
Ladoga, and the Baltic Sea population. Phenotypic sampling from museum collections roughly
matches those of the genetic data, except for Arctic samples from Kara Sea (square) substituting
for Pechora Sea, and Okhotsk samples being from Hokkaido or nearby islands (square).

Figure 2. Genetic uniqueness of the Saimaa ringed seals. (A) Using five individuals from each population, the Saimaa ringed seals form a tightly defined and separate cluster along the PC1 in the PCA analysis, while the genetic differentiation of the non-Saimaa seals creates a geographic continuum along the PC2: the Baltic are placed between the Ladoga and Arctic, and Okhotsk further away. The Arctic individual placed between the rest of the Arctic and Okhotsk is from the Bering Strait. (B) Comparable patterns are obtained using the full 46 individual data, although the uneven sampling hides the separation between Arctic and Okhotsk. The x-axis is reversed to visually match (A). (C) With MAF 0.05 filtering, Saimaa has the highest proportion of population-specific derived alleles in samples of 1-5 individuals per subspecies.

Figure 3. Estimated past effective population sizes and migration rates reveal a deep history for Saimaa ringed seals. (A) Estimates of past effective population sizes (N_e) for the three Fennoscandian and Arctic ringed seals (the distant Okhotsk is not included in these analyses) as computed from the full genome data. (B) Estimates of migration rates over time between pairs of individuals from different populations. The rates are cut at the point where the cumulative migration probability reaches 0.999. Background shadings indicate the average cumulative probability of 0.01, 0.25, 0.75 and 0.99, and dashed lines that of 0.50. (C) Cumulative migration probability between pairs of individuals from different populations. The dotted lines indicate the thresholds M_{50}, M_{95} and M_{99}, the x-coordinates of intersections defining the age of each pair reaching the respective level. Five chromosomes per population were included and each line represents a comparison of individuals from two different populations. Mutation rate (μ) and generation time were 1.826e-8 and 10 years, respectively.

Figure 4. Saimaa ringed seals are distinct in their dental anatomy. (A) Mean lengths of lower
postcanines P_2 to P_4 of Saimaa ringed seals are generally smaller than in Baltic and Ladoga, but
only P_5 is distinctly small compared to all the other populations. In the Saimaa dentition, the
relative tooth heights (B), and the top-cusp angles measuring relative height of the lateral cusps
(C), are the highest and smallest, respectively. These indicate shaper crown profiles in the
Saimaa ringed seal, whereas the other populations retain the more characteristic ringed seal
morphology irrespective of the size (A). Means differing from Saimaa with *P*-values below 0.05
are marked with open symbols in (A) to (C). For the *P* values, see Table 2. (D) ROC curves
showing the performance of different teeth and measures in the classification of Saimaa ringed
seals, P_5 and to a lesser degree P_2 performing the best for all the measures. For the length,
removing absolute size variation by calculating P_5/P_4, improves the ROC curve further (purple).
797

Figure 5. Tongue and dental morphology suggest increased suction but decreased filter-feeding niche for the Saimaa ringed seal. (A) Saimaa ringed seals have a broader tongue with a more rounded bifurcated tip, and intermolar elevations on the lateral side of the tongue (asterisks). (B) Comparison of tongue shapes with crabeater seal (*Lobodon*) and leopard seal (*Hydrurga*) tongues (from Loza et al. 2023) shows that the Saimaa ringed seal have more evenly broad tongue. (C) The cusp relief, measuring the filtering capacity of teeth is lower in Saimaa than in Baltic ringed seals, which in turn have comparable relief to that of the leopard seal. *P*-values between Saimaa sample and the other three are from 0.001 to 0.008 (one-tailed Mann-Whitney *U* test, Table S7). To illustrate the tongue shapes, colors in (A) were adjusted to be similar. Scale bars for (A) and (C), 5 mm.

Tables

814 **Table 1.** Average age (years ago) for cumulative migration probability reaching 50%, 95% and
815 99% for different population pairs. SD=standard deviation.

	M_{50}	SD_{50}	M_{95}	SD_{95}	M_{99}	SD_{99}
Arctic-Saimaa	65,819	1,770	336,303	29,985	490,069	50,029
Baltic-Saimaa	60,220	1,692	261,615	10,421	372,045	14,322
Ladoga-Saimaa	63,152	2,033	294,702	21,567	413,519	32,510
Arctic-Baltic	29,199	2,874	169,737	5,719	272,638	6,849
Baltic-Ladoga	33,502	1,151	148,504	10,678	244,162	12,628
Arctic-Ladoga	41,718	1,369	209,608	8,912	310,000	14,762

816
817
818 **Table 2**. Permutation tests between Saimaa and other samples for tooth length, tooth height,
819 height/length ratio (H/L), and top-cusp angle. *P*-values are one-tailed and except for the ones
820 marked with asterisk, denote lower values for Saimaa lengths and top-cusp angles, and higher
821 values for heights and height/length ratios. Tests are between group means using 10 000
822 permutations, *P*-values below 0.05 are marked in bold.

Length	Ladoga	Baltic	Arctic	Okhotsk	Angle	Ladoga	Baltic	Arctic	Okhotsk
P2	**0.0000**	**0.0000**	**0.0061**	0.0623*	P2	**0.0000**	**0.0000**	**0.0020**	**0.0051**
P3	**0.0202**	**0.0001**	0.2003	**0.0019***	P3	**0.0000**	**0.0001**	**0.0031**	**0.0000**
P4	**0.0095**	**0.0002**	0.3261	**0.0018***	P4	0.2596	0.0641	0.0671*	**0.0260**
P5	**0.0000**	**0.0000**	**0.0000**	**0.0011**	P5	**0.0000**	**0.0000**	**0.0000**	**0.0006**

Height	Ladoga	Baltic	Arctic	Okhotsk	H/L	Ladoga	Baltic	Arctic	Okhotsk
P2	**0.0145**	0.1325	**0.0026**	**0.0000**	P2	**0.0000**	**0.0000**	**0.0000**	**0.0000**
P3	**0.0126**	0.3291	**0.0020**	**0.0000**	P3	**0.0001**	**0.0008**	**0.0000**	**0.0001**
P4	0.1663*	0.1151*	0.2045	**0.0000**	P4	0.0571	0.1181	0.0513	**0.0035**
P5	0.0559*	0.0772*	**0.0012***	**0.0027**	P5	**0.0000**	**0.0000**	**0.0000**	**0.0000**

823
824

Supporting figures

Figure S1. The maximum likelihood tree for the unique mitogenome haplotypes rooted with four
spotted seals. The dots at internal nodes indicate bootstrap support values greater than 50%. In
contrast to the four ringed seal subspecies, all Saimaa samples (in blue) cluster tightly together
and form an evolutionary lineage of their own.

Figure S2. The receiver operating characteristic (ROC) curves for tooth lengths plotted
separately for female and male specimens. Of the 326 specimens measured, 96 (28 from Saimaa)
and 115 (20 from Saimaa) are assigned to female and male, respectively. For both sexes, the
P_5/P_4 performs the best, but female seals show almost perfect classification. Areas below the P_5
and P_5/P_4 curves are 0.90 and 0.98 for females, and 0.81 and 0.87 for males.