Proteomic Landscape of Pattern Triggered Immunity in the Arabidopsis Leaf Apoplast

Running title: PTI proteomics in Arabidopsis leaf apoplast

Hsiao-Chun Chen[1], Carter J. Newton[1], Gustavo Diaz[4], Yaochao Zheng[2], Feng Kong[1], Yao Yao[2],

Li Yang[1, 3], Brian H. Kvitko[1, 3,]*

[1]Department of Plant Pathology, University of Georgia, Athens, Georgia, USA

[2]Department of Animal and Dairy Science, Regenerative Bioscience Center, University of

Georgia, Athens, Georgia, USA

[3]The Plant Center, University of Georgia, Athens, Georgia, USA

[4]Analytical Resources Core: Bioanalysis and Omics, Colorado State University, Fort Collins, CO

*Address correspondence to Brian H. Kvitko, bkvitko@uga.edu.

The author responsible for distribution of materials integral to the findings presented in this

article in accordance with the policy described in the Instructions for Authors

(https://academic.oup.com/plphys/pages/General-Instructions) is Brian H. Kvitko

Abstract

The apoplast is a critical interface in plant-pathogen interactions particularly in the context of

pattern-triggered immunity (PTI), which is initiated by recognition of microbe-associated

molecular patterns (PAMPs). Our study characterizes the proteomic profile of the Arabidopsis

apoplast during PTI induced by flg22, a 22 amino acid bacterial flagellin epitope, to elucidate the

output of PTI. Apoplastic washing fluid (AWF) was extracted with minimal cytoplasmic

contamination for LC-MS/MS analysis. We observed consistent identification of PTI enriched

and depleted peptides across replicates with limited correlation between total protein abundance

and transcript abundance. We observed topological bias in peptide recovery of receptor-like

kinases with peptides predominantly recovered from their ectodomains. Notably, tetraspanin 8,

an exosome marker, was enriched in PTI samples. We additionally confirmed increased

concentrations of exosomes during PTI. This study enhances our understanding of the proteomic

changes in the apoplast during plant immune responses and lays the groundwork for future

investigations into the molecular mechanisms of plant defense under recognition of pathogen

molecular patterns.

Keywords

Pattern-triggered immunity (PTI), Microbe-associated molecular patterns (MAMPs), Apoplast,

Proteomics, Extracellular vesicle, Tetraspanin, flg22

Introduction

Plants are constantly exposed to microbial pathogens, necessitating the evolution of sophisticated

defense mechanisms to ensure survival (Boller and He, 2009; Mott et al., 2014). Pattern-

triggered immunity (PTI) represents the first layer of inducible defense, activated by the

recognition of conserved microbe-associated molecular patterns (MAMPs) through pattern

recognition receptors (PRRs) (Boller and He, 2009). These recognition events trigger a complex

regulatory network that initiates a variety of defense responses, including the production of

reactive oxygen species (ROS), transcriptional reprogramming, and callose deposition (Mott et

al., 2014; DeFalco and Zipfel, 2021). PTI serves as a critical barrier to pathogen invasion and

underscores the importance of studying the molecular mechanisms that underpin this process

(Boller and He, 2009; DeFalco and Zipfel, 2021).

The apoplast is the intercellular space that contains gas and water, situated between cell

membranes and within cell wall matrix (Farvardin et al., 2020). This compartment serves as a

critical battleground for plant-microbe interactions and is the location of colonization by foliar

bacteria and many other pathogens (Roussin-Léveillée et al., 2024). The apoplast of plants is

typically characterized by limited water and nutrient availability, presenting a challenging

environment for microbial proliferation (Freeman and Beattie, 2009; O'Leary et al., 2016; Xin et

al., 2016; Aung et al., 2018; Gentzel et al., 2022; Liu et al., 2022; Lovelace et al., 2022). The

alteration of the apoplastic environment has been proposed to impact pathogen resistance

(Roussin-Léveillée et al., 2024). Studies of apoplastic secreted proteins, peptides, and specialized

metabolites have provided some insights into apoplastic defense (Anderson et al., 2014;

Delaunois et al., 2014; Martínez-González et al., 2018; Gentzel et al., 2022; Serag et al., 2023).

Apoplastic proteins have been found to perform diverse functions, including reinforcing the plant

cell wall, signal transduction, and inhibition of microbial growth (Alexandersson et al., 2013;

Munzert and Engelsdorf, 2025). Secreted proteins include Pathogenesis-related proteins (PRs),

enzymes for cell wall modification, enzymes for generation of ROS and redox regulatory

proteins (van Loon et al., 2006; Camejo et al., 2016; Nishimura, 2016). Secreted proteases,

chitinases, and other hydrolytic enzymes can mediate defense by modification of plant or

pathogen structural targets, or the virulence factor targets such as pathogen effector proteins (van

der Hoorn, 2008). In addition to secreted proteins, plant extracellular vesicles (P-EVs) are

emerging as crucial players in immunity. P-EVs are membrane-bound nanostructures classified

into MVBs (multivesicular bodies), EXPO (exocyst-positive organelle), Penetration 1 (Pen1)-

positive EVs, vacuoles, and autophagosomes based on their biogenesis (Nemati et al., 2022).

They facilitate intercellular and interkingdom communication by transporting bioactive

molecules, including proteins and RNAs (Rutter and Innes, 2017; Cai et al., 2019). EVs are

implicated in enhancing plant defense through the delivery of immune-related cargo, such as

small RNAs and proteins, and contribute to systemic immune signaling (Liu et al., 2021).

Exosomes, a type of MVB-derived EV with a median size of approximately 30-150 nm in

Arabidopsis, have been characterized as a major subtype of P-EVs (Huang et al., 2021). Notably,

markers such as tetraspanin-8 (TET8) have been identified in defense-associated exosomes,

highlighting their potential role in coordinating immune responses (Wang et al., 2023).

Proteomics approaches using suspension cell culture systems have provided important

information of extracellular proteins for understanding the molecular mechanisms involved in

pathogen infections (Kaffarnik et al., 2009; Kim et al., 2009). While suspension cell culture

systems offer the advantage of limiting cytoplasmic contamination, they do not fully replicate the

actual conditions of infection or the specific cellular responses observed in intact tissue apoplast,

which are critical for understanding the spatial and temporal dynamics upon pathogens interact.

Characterization of apoplastic proteomics typically involve isolating apoplastic washing fluid

(AWF), followed by protein identification using techniques such as liquid chromatography-

tandem mass spectrometry (LC-MS/MS) (Jung et al., 2008). Isolation of AWF is typically

conducted using vacuum or pressure infiltration followed by low-speed centrifugation (Agrawal

et al., 2010). This technique is efficient in extracting leaf apoplast contents while minimizing

cytoplasmic contamination. To evaluate potential contamination from cytoplasmic components,

several assessments are employed using enzymatic assay or immunoblotting to ensure the purity

of the AWF (Delaunois et al., 2013; O'Leary et al., 2016). The dynamic nature of the apoplastic

proteome and its response to MAMPs remains an active area of research. The Arabidopsis

apoplastic proteomic profile during PTI is not well described. Further investigation into the

apoplastic proteome under pre-activated PTI will provide crucial insights into the complex

interplay of antimicrobial barrier, ultimately leading to the development of more effective

strategies for disease control.

In this study, we report the apoplastic proteome of *Arabidopsis thaliana* during PTI induced by

flg22 a twenty-two amino acid peptide epitope derived from bacterial flagellin a well-studied

MAMP. AWF was extracted by low-speed centrifugation with minimal cytoplasmic

contamination and analyzed using LC-MS/MS. Our analyses identified proteins significantly

enriched or depleted during PTI. We also compared these proteome profiles with publicly

available transcriptomics time-course data, providing insights into the dynamics of early and late

PTI outputs. Notably, we observed an increase during PTI in the exosome marker tetraspanin 8

and an increase of exosomes based on nanoflow cytometry. This study provides a detailed

snapshot of the proteomic changes in the *A. thaliana* apoplast during PTI elicited by flg22,

offering a foundation for understanding the molecular mechanisms governing plant immune

responses.

Results

Apoplastic washing fluid isolation

A total of 130–150 leaves were collected for apoplastic washing fluid (AWF) isolation from each

sample. 3.5–5.0 mL of AWF was recovered per sample across both mock and flg22 treatments.

Cytoplasmic contamination was assessed by measuring glucose-6-phosphate dehydrogenase

(G6PDH) activity in the AWF, compared to total leaf extracts (Figure S1). The average G6PDH

activity in the mock treatment was 0.55 mU/mL, and in the flg22 treatment was 0.16 mU/mL,

compared to 14.6 mU/mL in the total leaf extract. Both mock and flg22 samples exhibited <5%

G6PDH activity relative to the total leaf extract with no statistic difference between mock and

flg22 treatments, indicating minimal cytoplasmic contamination.

Principal Component Analysis (PCA) of flg22 and mock samples

The overall distribution of log-transformed total protein abundance per sample showed

consistency across replicates. Median $\log$ protein abundances were similar between the

flg22-treated and mock-treated samples, indicating a reliable dataset for further multivariate

analysis (Figure S2). Principal Component Analysis (PCA) was used to examine the proteomic

differences between flg22-treated and mock-treated samples (Figure 1). The first two

components, Dim1 and Dim2, explained 27.9% and 25.9% of the total variance, respectively.

The PCA plot revealed clustering based on sample types.

Comparison of PTI apoplastic proteins enrichment relative to time course transcriptome

profiles

A total of 108 proteins were identified as significantly enriched in samples at 16 hours post flg22

treatment comparing to mock controls (Log$_2$ Fold Change (Log2FC) $\geq$ 1 and p-value $\leq$ 0.05)

(Figure 2). We manually classified these proteins into six groups based on protein annotations in

the open-access information portal Araport: (1) RLKs/RLPs (Receptor-Like Kinases/Receptor-

Like Proteins), (2) redox and redox-associated proteins, (3) hydrolytic enzymes, (4)

antimicrobial peptides, (5) extracellular vesicle-associated proteins, and (6) others. To better

understand to what degree transcriptional regulation relates to protein abundance we integrated

our proteomic data with a published Arabidopsis transcriptomics time course from by Hillmer et

al., 2017. Within each category, proteins were ranked by their log2FC from highest to lowest

abundance. IOS1 and SIF2 were among the most abundant in the RLK/RLP category. Other

immune-associated RLKs like RLK902 and several CRKs (Cysteine-rich receptor-like protein

kinases) were also enriched (Gao et al., 2024). A considerable number of redox-related proteins,

such as peroxidases (PERs) and glutathione-S-transferases (GSTFs) were prominent. Several

hydrolytic enzymes were identified in the apoplast during flg22-induced PTI in *Arabidopsis*.

Enzymes involved in carbohydrate metabolism, including chitinase family proteins (At1g02360),

and beta-hexosaminidase 3 (HEXO3), were enriched. Subtilisin-like protease SBT3.3 and glucan

endo-1,3-beta-glucosidase 14 (At2g27500) were also identified, along with related beta-

glucosidases. Enzymes involved in cell wall modifications, such as pectin methyl-esterases

(PMEs) and xyloglucan endotransglucosylases, were present in the enriched profile. EARLI1

gene in Arabidopsis encodes a lipid transfer protein-like protein that plays a crucial role in

mediating systemic defense mechanisms, specifically systemic acquired resistance (SAR) and

induced systemic resistance (ISR) were also enriched by flg22 (Vlot et al., 2021). Notably, in

flg22-enriched profile, three proteins associated with vesicle trafficking were identified as

significantly enriched under PTI conditions. The enriched proteins include Syntaxin of Plants-

122 (SYP122), a Qa-SNARE which functions in vesicle fusion with the plasma membrane

peptidyl-prolyl cis-trans isomerase FKBP15-1 (FKBP15-1), which may contribute to protein

folding within vesicles, and tetraspanin-8 (TET8), associated with exosome extracellular vesicle

structural integrity (Boavida et al., 2013; Wang et al., 2020; Rubiato et al., 2022).

The heatmap (Figure 2) illustrates how the transcript abundance changes over time, providing

insights into the continuity or discontinuity between transcription and protein accumulation in

response to flg22 treatment. The transcriptomic analysis revealed distinct patterns of gene

expression in response to flg22 treatment, corresponding with the functional roles of the

identified proteins in PTI. There are 22 receptor-like kinases and receptor-like proteins

(RLK/RLP) identified in the enriched profile and 10 of them displayed rapid upregulation at the

first hours then the expression level gradually decreased. Several redox-related proteins and

hydrolytic enzymes, such as peroxidases (PER21, PER52, and PER71) and Glutathione S-

Transferase (GSTF6 and GSTF7) showed sustained transcriptional upregulation at a relatively

later time point. Overall, within each functional group, protein abundance ratios did not

necessarily align with transcriptional trends in a time-course manner.

Recovered peptides of RLKs are heavily biased to ectodomains.

RLKs membrane proteins were enriched in the apoplastic proteome after flg22 treatment.

Considering that these are transmembrane proteins, we would not expect them be found in the

free extracellular fraction. We investigated the topological distribution of detected peptides in 18

identified RLKs in our dataset enriched under PTI conditions (Figure 3). This detection pattern

shows a notable bias toward the extracellular region, suggesting that portions of these

membrane-associated proteins are accessible or released into the apoplast under PTI conditions.

Very few peptides were captured close to the cytoplasmic regions. The extracellular bias aligns

with the apoplast-specific protein collection procedure. The limited peptide coverage in

cytoplasmic regions further emphasize the specificity of the apoplast isolation protocol.

Apoplastic proteins showing lower abundance by flg22 treatment displayed limited

functional clustering and transcript-protein abundance discrepancies.

A total of 255 proteins were identified as significantly decreased following flg22 treatment

compared to mock conditions ((Log$_2$ Fold Change (Log2FC) $\leq$ -1 and p-value $\leq$ 0.05) (Figure 4).

Unlike the distinct functional clustering observed in the enriched protein profile by flg22

treatment, lower abundant proteins did not exhibit clear patterns linked to PTI or disease

response functions. However, based on molecular function GO analysis, these proteins were

categorized into 10 functional groups. We observed some genes displaying high transcript levels

post-flg22 treatment despite their decreased protein abundance in the apoplast, indicating some

discontinuity between transcript and protein abundances.

Gene Ontology (GO) analysis of the enriched protein profile under the flg22 treatment

We performed Gene Ontology (GO) analyses of the 108 significantly enriched proteins in flg22

treatment with TAIR GO Term Enrichment platform. The GO analysis identified significant

enrichment of multiple biological processes in flg22-enriched proteins (Figure 5A). Processes

with the highest gene number include response to stimulus and response to stress, suggesting that

many enriched proteins are involved in both general and specific stress responses. More specific

processes, such as response to biotic stimulus, defense response to Gram-negative bacterium, and

defense response to fungus, highlight these proteins' roles in responding to presence of

pathogens. Specific processes such as response to oxygen levels and response to oxidative stress

are also enriched, suggesting that these proteins may play roles in reactive oxygen species (ROS)

production, which are commonly associated with plant immune responses signaling (Torres et

al., 2006). Notably, highly representative pathway with higher fold enrichment like defense

response to Gram-negative bacterium align with the expected response to the bacterial elicitor

flg22.

The molecular function analysis for flg22-enriched proteins revealed significant enrichment in

various catalytic and kinase activities. Nearly 25% of the proteins identified in the GO analysis

were annotated for catalytic activity. The enrichment of oxidoreductase activity, including

peroxidase and lactoperoxidase functions, suggests roles associated with redox reactions and

potential oxidative stress responses. Proteins with kinase and transferase activities, particularly

those transferring phosphorus-containing groups, were also prominent. Specifically, protein

kinase activity and protein serine/threonine kinase activity were enriched, indicating involvement

in phosphorylation cascades.

Consistent with an apoplastic proteome, the cellular component analysis for flg22-enriched

proteins revealed significant enrichment in regions associated with the cell periphery,

extracellular region, and membrane. Nearly 15% of the proteins identified in the GO analysis

were localized to the cell periphery, suggesting a strong role in cell surface interactions. Notably,

vesicles and components associated with the secretory pathway were also prominent in the

dataset. The presence of proteins localized to vesicles, including secretory and cytoplasmic

vesicles, underscores their potential involvement in protein trafficking and secretion.

In contrast to the flg22-enriched proteins, which exhibited distinct functional categories related

to pattern-triggered immunity (PTI) and defense responses, the proteins with lower abundance

did not display clear patterns linked to these functions (Figure 5B). However, the GO analysis

revealed that these lower abundance proteins were involved in a wide range of cellular processes,

including metabolism and cellular turnover. The cellular component analysis of these proteins

showed their overall localization in the apoplast, with involvement in protein secretion and cell

periphery, albeit with small fold enrichment. The involvement of these lower abundance proteins

in the secretory pathway underscores their role in collecting extracellular proteins in the apoplast,

even if their abundance is reduced.

PTI is associated with increased numbers of extracellular vesicles (EVs) in leaf apoplast

Several studies have reported the enrichment of plant extracellular vesicles (EVs) during

pathogen infection (Rutter and Innes, 2017; Cai et al., 2019). In our dataset, we observed

significantly enriched proteins associated with vesicular export, including SYP122, TET8, and

FKBP15-1. SYP122 and TET8 show increased change of 8.0 and 5.2-fold, respectively, relative

to the mock condition. Notably, TET8 is an exosome marker involved in exosome stability (Cai

et al., 2019; Liu et al., 2024). Exosomes are one of several classes of EVs with a size typically

between 50nm-150nm in dimeter and have been associated with the cross-kingdom delivery of

small RNA for RNAi. We isolated plant EVs using ultracentrifugation and quantified them using

Nano-flow cytometry. The analysis of EV in both flg22-treated and mock samples reveals that

the median size of EVs in flg22-treated samples is 67.8 nm, compared to 70.7 nm in mock

samples (Figure 6A). This similarity in size indicates no significant difference between the two

conditions, suggesting that the majority of these vesicles fall within the size range characteristic

of plant exosomes (Rutter and Innes, 2017; Cai et al., 2019). Exosomes were found at 5.1×10

particles/mL in the flg22-treated samples, nearly three times higher than the $1.7 \times 10\square$

particles/mL observed in the mock samples (Figure 6C). This increased concentration in the

flg22 condition aligns well with the 5-fold higher protein levels identified in the flg22/mock

proteomics analysis. The staining of the AWF-derived EVs with the lipid dye, CFSE, highlighted

the shape and size characteristic of exosomes (Figure 6D).

Discussion

GO analysis of the PTI enriched apoplastic proteome revealed significant enrichment in

predicted biological processes such as defense responses to Gram-negative bacteria and

oxidative stress. The molecular functions enriched in flg22-treated samples included both kinase

and oxidoreductase activities, consistent critical roles of PRRs and redox regulation in plant

defense mechanisms. The cellular component GO terms, as would be expected, emphasized the

extracellular localization of both enriched and depleted proteins, including the cell wall, plasma

membrane, secretory vesicles, and extracellular regions. Unlike the PTI-enriched proteins, which

were prominently associated with defense signaling and oxidative stress management, the PTI-

depleted proteins displayed functions more broadly related to cellular homeostasis and general

secretion mechanisms. This may be due to less cytoplasmic leakage in flg22-treated samples

during the collection process (Figure S1) or may reflect an overall shift in molecular function.

Notably, more than 18% of depleted proteins were associated with functions related to cell

organization and biogenesis, including BRI1 (BRASSINOSTEROID INSENSITIVE 1),

indicating the stabilization of BR signaling under naïve conditions, this can potentially be a

consequence of the growth-defense trade-off (Huot et al., 2014).

We defined five functional categories of apoplastic proteins enriched during flg22-triggered PTI,

including RLK/RLPs, redox-associated proteins, hydrolytic enzymes, antimicrobial peptides,

extracellular vesicle (EV) associated proteins. Among the RLKs, IOS1 (Impaired Oomycete

Susceptibility 1) and SIF2 (Stress Induced Factor 2) were highly enriched. These two malectin-

domain RLKs have both been observed previously to play roles in PTI signaling. IOS1, a

malectin-like leucine-rich repeat RLK, modulates Arabidopsis immunity by forming complexes

with FLS2 and BAK1 to amplify pathogen recognition and defense responses. Loss-of-function

mutants of IOS1 demonstrate increased susceptibility to bacterial pathogens (Yeh et al., 2016).

Similarly, SIF2 also interacts with FLS2 and BAK1 and has been implicated in early flg22-

triggered immune signaling, contributing to downstream defense activation (Chan et al., 2020).

The peptide recovery analysis of RLKs demonstrated a marked bias toward the receptor

extracellular regions, with minimal peptides detected in their cytoplasmic regions. The limited

recovery of cytoplasmic peptides further aligns with the low levels of cytoplasmic contamination

in the apoplastic washing fluid, emphasizing the effectiveness of the apoplast isolation protocol

as described by Lovelace et al., 2022. The recovery of free ectodomains in the apoplast may be

an artifact of the sampling procedure or could indicate that these ectodomains are endogenously

released into the apoplast under the tested conditions through an unknown mechanism.

A comparative analysis of transcriptome and proteome profiles revealed a complex relationship

between transcriptional regulation and endpoint protein abundance during flg22-triggered PTI.

Several high-abundant proteins displayed consistent trends between transcript levels and protein

accumulation over the time course. For example, IOS1 and SIF2, both receptor-like kinases

(RLKs), exhibited rapid transcriptional upregulation within the first few hours of flg22 treatment.

This aligns with their role in early pathogen recognition and activation of downstream signaling

pathways (Bigeard et al., 2015). Similarly, Peroxidases such as PER15 (Peroxidase 15), showed

a strong correlation between transcript and protein levels, consistent with its involvement in

reactive oxygen species (ROS) management and oxidative stress responses during PTI.

However, other high-abundance proteins demonstrated protein enrichment despite declining

transcript levels over time. Cysteine-Rich RLKs, another set of RLKs, have distinct patterns of

expressions despite their role in response to flg22 perceptions (Yadeta et al., 2017). For

instance, CRK11 and CRK12 remained abundant in the apoplast even as their transcript levels

decreased after the initial hours of flg22 exposure. On the other hand, CRK13 exhibited

decreased transcript levels over time but maintained a high expression pattern and protein

abundance during late PTI, suggesting its crucial role in enhanced resistance to the bacterial

pathogen *Pseudomonas syringae* (Acharya et al., 2007). These suggested post-transcriptional

regulatory mechanisms, such as enhanced mRNA stability or protein stabilization, which might

prolong the functional presence of critical immune components. Such discrepancies reflect the

likely importance of post-transcriptional regulation and protein turnover dynamics in fine-tuning

the immune response.

The enrichment of various hydrolytic enzymes in apoplast in response to flg22 treatment

suggested a multifaceted response to PTI. Several enzymes involved in plant cell wall

modification were found to be enriched including pectin methyl-esterases (PME3, PME17),

xyloglucan endotransglucosylase/hydrolase (XTH23), and β-1,3-glucanases (At2g27500 and

At5g56590) (Bethke et al., 2014; Perrot et al., 2022; Zhang et al., 2022). These enzymes

collectively suggest that the plant is actively remodeling its cell wall during PTI, possibly to

enhance its barrier function against potential pathogens. Several hydrolytic enzymes capable of

N-glycosylation of immune receptor and degrading pathogen-associated targets (especially

fungal cell wall) were enriched, such as β-hexosaminidase 3 (HEXO3) and endochitinases

(At2g43620 and At1g02360) (Liebminger et al., 2011; Fiorin et al., 2018). These enzymes

suggest that the plant is actively degrading potential pathogen-derived molecules, which could

both weaken a broad-spectrum of pathogens and generate additional immune-stimulating signals.

Various proteases and protein-modifying enzymes were also enriched. Metalloendoproteinases

(2-MMP and 3-MMP) and subtilisin-like protease (SBT3.3) may be involved in processing

defense-related proteins within the extracellular matrix or degrading pathogen-derived proteins

(Flinn, 2008; Ramírez et al., 2013; Zhao et al., 2017). Plant aspartyl proteases (MMG4.12 and

At3g02740) and cathepsin B-like protease 2 (CATHB2) were also found. These proteases

contribute to plant immunity through various mechanisms, from immune signaling activation and

systemic resistance to direct cleavage of pathogen proteins (McLellan et al., 2009; Figueiredo et

al., 2021). The presence of these proteases suggests active protein processing and turnover,

which could be important for the homeostasis of defense signaling molecules and limiting

pathogen fitness. Other metabolic enzymes, such as NUDT6 and NUDT7, members of the nudix

hydrolase family, were enriched by flg22. They play crucial roles in regulating plant immunity

and stress responses by maintaining the redox balance of NADH and acting as ADP-ribose

pyrophosphatases (Fonseca and Dong, 2014). The coordinated action of these hydrolytic

enzymes with various types of substrates likely contributes to the rapid and effective immune

response against potential pathogens, highlighting the sophisticated defense mechanisms

employed in pathogen recognition and immune activation.

The observation of pathogenesis-related 1 (PR1) protein enrichment in the flg22 treatment

profile, with a notable 4.67-fold change, strongly aligns with key findings in induced pattern-

triggered immunity (PTI). PR1 is a well-established marker for salicylic acid (SA)-regulated

plant immunity, and its secretion is critical for activating systemic acquired resistance (SAR).

Upon flg22 treatment, plants exhibit increased PR1 expression as part of their PTI response

(Djamei et al., 2007). The significant enrichment of PR1 protein abundance observed in this

study confirms its responsiveness to PTI triggered by flg22 treatment and further demonstrates

the effectiveness of the experimental conditions in inducing a robust immune response.

For lower abundant proteins by flg22 treatment, inconsistencies with transcript levels were even

more pronounced. Many of these proteins, such as certain housekeeping enzymes and stress-

related factors, exhibited little to no transcriptional change over the time course. Their reduced

abundance in the apoplast may result from active degradation, inhibited secretion, or preferential

retention in intracellular compartments. Unlike high-abundant proteins, these low-abundant

factors were associated with less PTI-relevant processes, further indicating that their depletion

may reflect cellular reorganization rather than active participation in immune responses. The

integration of transcriptomics and proteomics provides additional insights into the temporal

coordination of immune responses. Further time-resolved multi-omics studies could elucidate the

functional significance of these regulatory discrepancies, revealing additional layers of

complexity in plant immunity.

Several extracellular vesicles (EVs) -associated proteins exhibited a notable increase in flg22-

treated apoplast samples. In particular, tetraspanin-8 (TET8), a well-established exosome marker,

was enriched fourfold compared to mock-treated samples. Cai et al. (2018) previously identified

TET8 as a core component of exosomes involved in stress responses, and pathogen defense as

mediators of interkingdom RNAi. Additionally, the concentration of EVs in flg22-treated

samples was approximately three times higher than in mock-treated samples. This increase in

EVs production suggests enhanced trafficking of defense molecules and signaling factors to the

apoplast. The enrichment of Syntaxin-122 (SYP122), a SNARE protein involved in vesicle

fusion, supports the idea that vesicle-mediated transport is a key component of the immune

response (Waghmare et al., 2018). These results underscore the importance of EVs as critical

components of PTI, facilitating intercellular communication and the targeted delivery of

antimicrobial proteins and signaling molecules. The substantial enrichment of TET8 and the

increased exosome particles suggest that EVs play a pivotal role in coordinating and amplifying

defense responses during pathogen attack.

Overall, our study provides a snapshot of the apoplastic proteome in *Arabidopsis* during the later

phases of flg22-induced pattern-triggered immunity (PTI). A total of 108 significantly enriched

proteins were identified and categorized into key functional groups, including receptor-like

kinases/receptor-like proteins (RLK/RLPs), redox-related proteins, hydrolytic enzymes, small

peptides, and extracellular vesicle (EV)-associated proteins. Our results highlight the complexity

of the apoplastic immune response, where proteins function in distinct stages of PTI activation,

defense reinforcement, and signaling. Notably, exosome-associated proteins such as tetraspanin-

8 (TET8) were enriched fourfold, and exosome particles increased by approximately threefold.

The integration of proteomic and transcriptomic data revealed that transcriptional trends do not

always align with protein abundance. This research advances our understanding of the dynamic

changes occurring in the apoplast during PTI and underscores the multi-faceted strategies plants

employ to defend against pathogen attack. These findings provide a valuable foundation for

future investigations into the molecular mechanisms of PTI.

Materials and Methods

Plant Tissue Preparation

A. thaliana Col-0 seeds suspended in sterile 0.1% agarose were sown in SunGrow 3B

Professional potting mix and stratified in darkness for 2 day at 4°C before being grown in a

growth chamber (Conviron A1000) with 14-h light (70 μmol m^{-2} s^{-1}) at 22°C. After 4 weeks,

plants were transferred to a growth room maintained under a 12-hour light/12-hour dark cycle for

acclimatization. At 4.5 weeks, plants were treated for 16 hours to induce pattern-triggered

immunity (PTI). Treatments were applied using a 1 mL blunt-end syringe to 4-5 adult leaves per

plant. The treatments consisted of 1 μM flg22 peptide to induce PTI and 0.1% dimethyl

sulfoxide (DMSO) as a mock control treatment. The solutions were carefully infiltrated into the

abaxial side of the selected leaves, and plants were maintained under the same growth room

conditions during the 16-hour treatment period.

Extraction of Apoplastic Washing Fluid and Cytoplasmic Contamination Measurement

Apoplastic washing fluid (AWF) was crude extracted using vacuum infiltration as described by

(Lovelace et al., 2022). 130-150 *A. thaliana* leaves were cut and placed into a 500 mL beaker

filled with iced-cold distilled water to top. Repeated cycles of vacuum at 95 kPa for 2 min

followed by slow release of pressure were applied until leaves were fully infiltrated. Excess

water was blotted from plant tissue before rolled into Saran wrap which were placed into 50 mL

conical tubes. Tubes were centrifuged at 1,000 xg for 10 min at 4°C and the fractions were

pooled and stored at -80°C. Cytoplasmic contamination in our AWF samples was examined by

comparisons of cytosolic marker glucose-6-phosphate dehydrogenase- G6PDH activity in

sampled AWF extracts to the total leaf extracts using standard kit (Sigma-Aldrich, St. Louis,

MO, USA). Leaves were homogenized at 4°C. G6PDH activity was assayed

spectrophotometrically every 5 min at 37°C where each reaction contained 50 μL of AWF and

the activity was calculated according to manufacturer's instructions (Sigma-Aldrich, St. Louis,

MO, USA).

Sample Preparation and Mass Spectrometry Analysis - LC-MS/MS

Samples were filtered through a 3 kDa Amicon spin filter to approximately 50 µL. The retentate

was collected via reverse centrifugation, washed with 50 µL of 50 mM Ammonium bicarbonate,

and combined. Total protein content was measured using the Pierce BCA Protein Assay Kit

(ThermoFisher Scientific) and quantified at 550 nm based on a BSA standard curve.

For digestion, 100 µg of protein from each sample was prepared with the EasyPep Mini MS

Sample Prep Kit (ThermoFisher Scientific). After reduction and alkylation at 95°C, samples

were digested with Trypsin/LysC (0.2 µg/µL) at 37°C for 2 hours. Digestion was stopped,

contaminants removed using peptide cleanup columns, and the eluate was dried and resuspended

in 3% acetonitrile/0.1% formic acid. Peptide concentration was measured at 205 nm on a

NanoDrop, calculated using an extinction coefficient (Scopes, 1974). Reverse-phase

chromatography was performed with water + 0.1% formic acid (A) and 80% acetonitrile + 0.1%

formic acid (B). A total of 1 µg of peptides was enriched using a PepMap Neo C18 trap-column,

followed by separation on a Vanquish Neo system (Thermo Scientific) with a C18 nanospray

column at 45°C using a 90 minute gradient at a flow rate of 300 nanoliters/min: 1-6%B over 3

minutes followed by 6-35%B over 70 minutes, 35-45%B over 5 minutes ending in 12 minutes of

washing at 500 nanoliters/minute, 99%B. Peptides were eluted directly into an Orbitrap Eclipse

mass spectrometer with a Nanospray Flex ion source and analyzed in Data Dependent

Acquisition mode. MS spectra were collected over m/z 375–2000 in positive mode, with ions of

charge state +2 or higher selected for MS/MS. Dynamic exclusion was set to 1 MS/MS per m/z

with a 60 s exclusion. MS detection was performed in FT mode at 240,000 resolution, and

MS/MS in ion trap mode with HCD collision energy at 30%.

Data Analysis and Statistics

Data processing was conducted in Proteome Discoverer (PD) 3.0 (Thermo Scientific). A

precursor detector node (S/N=1.5) was used to identify additional precursors within the isolation

window for chimeric spectra. Sequest HT searched spectra with methionine oxidation as a

dynamic modification and cysteine carbamidomethylating as a fixed modification. An intensity-

based rescoring (INFERYS node) using deep learning predicted fragment ion intensities (Zolg et

al., 2021). Data were matched against the *Arabidopsis thaliana* Uniprot proteome

(UP000006548) and cRAP contaminants. Searches used a fragment ion tolerance of 0.60 Da and

parent ion tolerance of 10 PPM. Peptide spectrum matches (PSMs) were validated by Percolator

with FDR ≤1% and protein IDs confirmed by at least one peptide. Normalized abundance was

log2 transformed. Log2 fold changes (FC) identified proteins with higher (Log2 FC ≥ 1) or lower

(Log2 FC ≤ -1) abundance in flg22 vs. mock, with significance tested using t-tests.

Gene Ontology (GO) Analysis

To identify functional categories enriched in the proteomic data, both significantly enriched

proteins and significantly depleted proteins were analyzed using the Gene Ontology (GO)

analysis The Arabidopsis Information Resource (TAIR) GO Term Enrichment tool version 2.

The analysis was conducted using the TAIR10 genome as the reference dataset to determine

over- and under-representation of specific biological processes, molecular functions, and cellular

components. Enriched GO terms were considered significant based on a Bonferroni-corrected p-

value cutoff of <0.05. The results provided insight into functional trends within the identified

apoplastic proteome, including processes associated with plant defense and signal transduction.

Transcriptomics analysis

To investigate the transcriptional profiles of *Arabidopsis thaliana* proteins identified as significantly enriched or significantly lower abundant in response to flg22 treatment, Tag-Seq data for wild-type Col-0 plants were downloaded from Gene Expression Omnibus (GEO, accession number GSE78735) as described by Hillmer et al., 2017. This dataset comprised transcriptome profiles collected at seven time points post flg22 treatment (0, 1, 2, 3, 5, 9, and 18 hours post-infiltration) with three biological replicates per time point. For each gene, read counts from the three biological replicates were averaged at each time point, normalized to the 0-hour time point by calculating fold changes, and log10-transformed. These processed values were visualized as a heatmap using the pheatmap function from the pheatmap package (version 1.0.12) in R.

Topology analysis

To analyze the topological distribution of receptor-like kinases (RLKs) in the high-abundance protein profile in response to flg22/mock treatment, peptides identified via Proteome Discoverer (PD) 3.0 were mapped to specific domains or positions within each RLK. Protein domain annotations for the RLKs were obtained from UniProt. Identified peptides corresponding to RLKs were selected and extracted, and their positions were aligned to the respective RLK sequences. This mapping allowed for the determination of peptide coverage within specific protein domains, including extracellular, transmembrane, and intracellular regions.

Plant Extracellular vesicle isolation and quantification

The plant extracellular vesicles (P-EVs) isolation protocol was modified from previous studies (Huang et al., 2021). Briefly, the crude AWF was centrifuged for 30 min at 4 °C at $2,000 \times g$ to remove large cell debris, and then the supernatant was further centrifuged at $10,000 \times g$ for 30

minutes at 4 °C to remove large insoluble particles. The supernatant (the clean AWF) was then

collected and further processed by ultracentrifugation at 100,000 xg for 1 hour at 4 °C (Optima™

TLX Ultracentrifuge, Beckman Coulter, Indianapolis, IN) to purify and concentrate P-EVs. Each

P-EV pellet was gently resuspended in 50 µL of 0.22nm-filtered PBS, and all samples were

aliquoted and stored at -80 °C until further use. Nano-flow cytometry (NFC, NanoFCM, China)

was used to analyze P-EVs isolated from both flg22 and mock samples, following the

manufacturer's instructions. For each sample, a 0.2 µL aliquot of P-EVs was diluted in 50 µL of

filtered PBS (1:250 dilution) for analysis. Particle concentration and size distribution were

measured in triplicates. To evaluate differences in P-EVs origin between the flg22 and mock

groups, yield was calculated as the number of EVs per gram of leaf tissue used for AWF

collection. Statistical analysis was performed using Student's t-test to determine P-EV yield

differences between groups (*$P<0.05$, **$P<0.01$, NS: not significant).

P-EVs from mock treatment samples were stained with CFSE dye and examined under confocal

microscope. The fluorescence signals were observed by using Zeiss LSM 880 confocal laser-

scanning microscopy. FITC filter (Excitation: 460–500 nm; Emission: 510–560 nm) (Zeiss,

Germany) was used to detect the GFP signal (Excitation: 530–560 nm) (Nikon, Tokyo, Japan).

Images were generated and merged using The Zen 2.3 imaging software.

Data Statement

The mass spectrometry proteomics data is deposited to the ProteomeXchange Consortium via the

PRIDE partner repository (Perez-Riverol et al., 2025) with the dataset identifier PXD060654 and

10.6019/PXD060654. Details regarding experimental design, sample preparation, and LC-

MS/MS parameters are described in the Materials and Methods section of this manuscript. Total

identified proteins, significantly high-abundance proteins by flg22 treatment, and significantly

low abundance by flg22 treatment are listed in the Supplementary Data.

Acknowledgements

This work was supported by the National Science Foundation Grant 1844861 to BK. CJN, LY,

and FK was supported by NIH R35GM143067. YZ and YY were supported by startup funds

from the University of Georgia (to YY). We thank Su-Yun Uhm from Department of Genetics,

University of Georgia for assistance of Arabidopsis planting and AWF collection. We

acknowledge the use of resources provided by the Colorado State University ARC-BIO

(Research Resource ID: SCR_021758). Data was generated using the Orbitrap Eclipse Mass

Spectrometer supported by NSF Grant 2117943. We acknowledge the contribution of ARC-BIO

staff- Dorathea Lee for sample preparation.

Author contributions

HCC, YY, and BHK designed the research; HCC, FK, YZ conducted experiments; HCC, GD,

CJN, YZ analyzed the data; HCC, FK, YZ, and BHK wrote the paper. All listed authors

reviewed and approved draft and final versions of the manuscript.

Conflict of interest

The authors declare no competing interests.

References

Acharya BR, Raina S, Maqbool SB, Jagadeeswaran G, Mosher SL, Appel HM, Schultz JC,

Klessig DF, Raina R (2007) Overexpression of CRK13, an Arabidopsis cysteine-rich

receptor-like kinase, results in enhanced resistance to *Pseudomonas syringa*e. The Plant

Journal **50**: 488–499

Agrawal GK, Jwa N-S, Lebrun M-H, Job D, Rakwal R (2010) Plant secretome: Unlocking

secrets of the secreted proteins. PROTEOMICS **10**: 799–827

Alexandersson E, Ali A, Resjö S, Andreasson E (2013) Plant secretome proteomics. Front

Plant Sci **4**: 9

Anderson JC, Wan Y, Kim Y-M, Pasa-Tolic L, Metz TO, Peck SC (2014) Decreased

abundance of type III secretion system-inducing signals in Arabidopsis mkp1 enhances

resistance against *Pseudomonas syringae*. Proc Natl Acad Sci U S A **111**: 6846–6851

Aung K, Jiang Y, He SY (2018) The role of water in plant–microbe interactions. The Plant

Journal **93**: 771–780

Bethke G, Grundman RE, Sreekanta S, Truman W, Katagiri F, Glazebrook J (2014)

Arabidopsis PECTIN METHYLESTERASEs Contribute to Immunity against *Pseudomonas*

syringae. Plant Physiology **164**: 1093–1107

Bigeard J, Colcombet J, Hirt H (2015) Signaling Mechanisms in Pattern-Triggered Immunity

(PTI). Molecular Plant **8**: 521–539

Boavida LC, Qin P, Broz M, Becker JD, McCormick S (2013) Arabidopsis tetraspanins are

confined to discrete expression domains and cell types in reproductive tissues and form

homo- and heterodimers when expressed in yeast. Plant Physiol **163**: 696–712

Boller T, He SY (2009) Innate immunity in plants: an arms race between pattern recognition

receptors in plants and effectors in microbial pathogens. Science **324**: 742–744

Cai Q, He B, Weiberg A, Buck AH, Jin H (2019) Small RNAs and extracellular vesicles: New

mechanisms of cross-species communication and innovative tools for disease control. PLOS

Pathogens **15**: e1008090

Cai Q, Qiao L, Wang M, He B, Lin F-M, Palmquist J, Huang S-D, Jin H (2018) Plants send

small RNAs in extracellular vesicles to fungal pathogen to silence virulence genes. Science

360: 1126–1129

Camejo D, Guzmán-Cedeño Á, Moreno A (2016) Reactive oxygen species, essential

molecules, during plant-pathogen interactions. Plant Physiol Biochem **103**: 10–23

Chan C, Panzeri D, Okuma E, Tõldsepp K, Wang Y-Y, Louh G-Y, Chin T-C, Yeh Y-H,

Yeh H-L, Yekondi S, Huang Y-H, Huang T-Y, Chiou T-J, Murata Y, Kollisit H,

Zimmerli L (2020) STRESS INDUCED FACTOR 2 Regulates Arabidopsis Stomatal

Immunity through Phosphorylation of the Anion Channel SLAC1. Plant Cell **32**: 2216–

2236

DeFalco TA, Zipfel C (2021) Molecular mechanisms of early plant pattern-triggered immune

signaling. Molecular Cell **81**: 3449–3467

Delaunois B, Colby T, Belloy N, Conreux A, Harzen A, Baillieul F, Clément C, Schmidt J,

Jeandet P, Cordelier S (2013) Large-scale proteomic analysis of the grapevine leaf

apoplastic fluid reveals mainly stress-related proteins and cell wall modifying enzymes.

BMC Plant Biology **13**: 24

Delaunois B, Jeandet P, Clément C, Baillieul F, Dorey S, Cordelier S (2014) Uncovering

plant-pathogen crosstalk through apoplastic proteomic studies. Front Plant Sci. doi:

10.3389/fpls.2014.00249

Djamei A, Pitzschke A, Nakagami H, Rajh I, Hirt H (2007) Trojan Horse Strategy in

Agrobacterium Transformation: Abusing MAPK Defense Signaling. Science **318**: 453–456

Farvardin A, González-Hernández AI, Llorens E, García-Agustín P, Scalschi L, Vicedo B

(2020) The Apoplast: A Key Player in Plant Survival. Antioxidants (Basel) **9**: 604

Figueiredo L, Santos RB, Figueiredo A (2021) Defense and Offense Strategies: The Role of

Aspartic Proteases in Plant–Pathogen Interactions. Biology (Basel) **10**: 75

Fiorin GL, Sanchéz-Vallet A, Thomazella DP de T, Prado PFV do, Nascimento LC do,

Figueira AV de O, Thomma BPHJ, Pereira GAG, Teixeira PJPL (2018) Suppression of

Plant Immunity by Fungal Chitinase-like Effectors. Current Biology **28**: 3023-3030.e5

Flinn BS (2008) Plant extracellular matrix metalloproteinases. Functional Plant Biol **35**: 1183–

1193

Fonseca JP, Dong X (2014) Functional Characterization of a Nudix Hydrolase AtNUDX8 upon

Pathogen Attack Indicates a Positive Role in Plant Immune Responses. PLOS ONE **9**:

e114119

Freeman BC, Beattie GA (2009) Bacterial Growth Restriction During Host Resistance to

Pseudomonas syringae Is Associated with Leaf Water Loss and Localized Cessation of

Vascular Activity in *Arabidopsis thaliana*. MPMI **22**: 857–867

Gao C, Zhao Y, Wang W, Zhang B, Huang X, Wang Y, Tang D (2024)

BRASSINOSTEROID-SIGNALING KINASE 1 modulates OPEN STOMATA 1

phosphorylation and contributes to stomatal closure and plant immunity. The Plant Journal

120: 45–59

Gentzel I, Giese L, Ekanayake G, Mikhail K, Zhao W, Cocuron J-C, Alonso AP, Mackey D

(2022) Dynamic nutrient acquisition from a hydrated apoplast supports biotrophic

proliferation of a bacterial pathogen of maize. Cell Host & Microbe **30**: 502-517.e4

Hillmer RA, Tsuda K, Rallapalli G, Asai S, Truman W, Papke MD, Sakakibara H, Jones

JDG, Myers CL, Katagiri F (2017) The highly buffered Arabidopsis immune signaling

network conceals the functions of its components. PLoS Genet **13**: e1006639

Huang Y, Wang S, Cai Q, Jin H (2021) Effective methods for isolation and purification of

extracellular vesicles from plants. J Integr Plant Biol **63**: 2020–2030

Huot B, Yao J, Montgomery BL, He SY (2014) Growth–Defense Tradeoffs in Plants: A

Balancing Act to Optimize Fitness. Molecular Plant **7**: 1267–1287

Jung Y-H, Jeong S-H, Kim SH, Singh R, Lee J, Cho Y-S, Agrawal GK, Rakwal R, Jwa N-S

(2008) Systematic Secretome Analyses of Rice Leaf and Seed Callus Suspension-Cultured

Cells: Workflow Development and Establishment of High-Density Two-Dimensional Gel

Reference Maps. J Proteome Res **7**: 5187–5210

Kaffarnik FAR, Jones AME, Rathjen JP, Peck SC (2009) Effector Proteins of the Bacterial

Pathogen *Pseudomonas syringae* Alter the Extracellular Proteome of the Host Plant,

Arabidopsis thaliana. Molecular & Cellular Proteomics **8**: 145–156

Kim ST, Kang YH, Wang Y, Wu J, Park ZY, Rakwal R, Agrawal GK, Lee SY, Kang KY

(2009) Secretome analysis of differentially induced proteins in rice suspension-cultured

cells triggered by rice blast fungus and elicitor. Proteomics **9**: 1302–1313

Liebminger E, Veit C, Pabst M, Batoux M, Zipfel C, Altmann F, Mach L, Strasser R (2011)

β-N-Acetylhexosaminidases HEXO1 and HEXO3 Are Responsible for the Formation of

Paucimannosidic N-Glycans in *Arabidopsis thaliana**. Journal of Biological Chemistry **286**:

10793–10802

Liu G, Kang G, Wang S, Huang Y, Cai Q (2021) Extracellular Vesicles: Emerging Players in

Plant Defense Against Pathogens. Front Plant Sci. **12**: 757925

Liu N, Hou L, Chen X, Bao J, Chen F, Cai W, Zhu H, Wang L, Chen X (2024) Arabidopsis

TETRASPANIN8 mediates exosome secretion and glycosyl inositol phosphoceramide

sorting and trafficking. The Plant Cell **36**: 626–641

Liu Z, Hou S, Rodrigues O, Wang P, Luo D, Munemasa S, Lei J, Liu J, Ortiz-Morea FA,

Wang X, et al (2022) Phytocytokine signalling reopens stomata in plant immunity and

water loss. Nature **605**: 332–339

van Loon LC, Rep M, Pieterse CMJ (2006) Significance of inducible defense-related proteins

in infected plants. Annu Rev Phytopathol **44**: 135–162

Lovelace AH, Chen H-C, Lee S, Soufi Z, Bota P, Preston GM, Kvitko BH (2022) RpoS

contributes in a host-dependent manner to *Salmonella* colonization of the leaf apoplast

during plant disease. Front Microbiol **13**: 999183

Martínez-González AP, Ardila HD, Martínez-Peralta ST, Melgarejo-Muñoz LM,

Castillejo-Sánchez MA, Jorrín-Novo JV (2018) What proteomic analysis of the apoplast

tells us about plant–pathogen interactions. Plant Pathology **67**: 1647–1668

McLellan H, Gilroy EM, Yun B-W, Birch PRJ, Loake GJ (2009) Functional redundancy in

the Arabidopsis Cathepsin B gene family contributes to basal defence, the hypersensitive

response and senescence. New Phytologist **183**: 408–418

Mott GA, Middleton MA, Desveaux D, Guttman DS (2014) Peptides and small molecules of

the plant-pathogen apoplastic arena. Front Plant Sci. doi: 10.3389/fpls.2014.00677

Munzert KS, Engelsdorf T (2025) Plant cell wall structure and dynamics in plant–pathogen

interactions and pathogen defence. Journal of Experimental Botany **76**: 228–242

Nemati M, Singh B, Mir RA, Nemati M, Babaei A, Ahmadi M, Rasmi Y, Golezani AG,

Rezaie J (2022) Plant-derived extracellular vesicles: a novel nanomedicine approach with

advantages and challenges. Cell Communication and Signaling **20**: 69

Nishimura M (2016) Cell wall reorganization during infection in fungal plant pathogens.

Physiological and Molecular Plant Pathology **95**: 14–19

O'Leary BM, Neale HC, Geilfus C, Jackson RW, Arnold DL, Preston GM (2016) Early

changes in apoplast composition associated with defence and disease in interactions

between *Phaseolus vulgaris* and the halo blight pathogen *Pseudomonas syringae* pv.

phaseolicola. Plant Cell Environ **39**: 2172–2184

Perez-Riverol Y, Bandla C, Kundu DJ, Kamatchinathan S, Bai J, Hewapathirana S, John

NS, Prakash A, Walzer M, Wang S, Vizcaino JA. (2025) The PRIDE database at 20

years: 2025 update. Nucleic Acids Res 53: D543–D553

Perrot T, Pauly M, Ramírez V (2022) Emerging Roles of β-Glucanases in Plant Development

and Adaptative Responses. Plants (Basel) **11**: 1119

Ramírez V, López A, Mauch-Mani B, Gil MJ, Vera P (2013) An Extracellular Subtilase

Switch for Immune Priming in Arabidopsis. PLOS Pathogens **9**: e1003445

Roussin-Léveillée C, Mackey D, Ekanayake G, Gohmann R, Moffett P (2024) Extracellular

niche establishment by plant pathogens. Nat Rev Microbiol **22**: 360–372

Rubiato HM, Liu M, O'Connell RJ, Nielsen ME (2022) Plant SYP12 syntaxins mediate an

evolutionarily conserved general immunity to filamentous pathogens. eLife **11**: e73487

Rutter BD, Innes RW (2017) Extracellular Vesicles Isolated from the Leaf Apoplast Carry

Stress-Response Proteins. Plant Physiol **173**: 728–741

Scopes RK (1974) Measurement of protein by spectrophotometry at 205 nm. Analytical

Biochemistry **59**: 277–282

Serag A, Salem MA, Gong S, Wu J-L, Farag MA (2023) Decoding Metabolic

Reprogramming in Plants under Pathogen Attacks, a Comprehensive Review of Emerging

Metabolomics Technologies to Maximize Their Applications. Metabolites **13**: 424

Torres MA, Jones JDG, Dangl JL (2006) Reactive oxygen species signaling in response to

pathogens. Plant Physiol **141**: 373–378

Van Der Hoorn RAL (2008) Plant Proteases: From Phenotypes to Molecular Mechanisms.

Annu Rev Plant Biol **59**: 191–223

Vlot AC, Sales JH, Lenk M, Bauer K, Brambilla A, Sommer A, Chen Y, Wenig M, Nayem

S (2021) Systemic propagation of immunity in plants. New Phytologist **229**: 1234–1250

Waghmare S, Lileikyte E, Karnik R, Goodman JK, Blatt MR, Jones AME (2018) SNAREs

SYP121 and SYP122 Mediate the Secretion of Distinct Cargo Subsets1. Plant Physiol **178**:

1679–1688

Wang J, Sun W, Kong X, Zhao C, Li J, Chen Y, Gao Z, Zuo K (2020) The peptidyl-prolyl

isomerases FKBP15-1 and FKBP15-2 negatively affect lateral root development by

repressing the vacuolar invertase VIN2 in Arabidopsis. Planta **252**: 52

Wang Z, Zeng J, Deng J, Hou X, Zhang J, Yan W, Cai Q (2023) Pathogen-Derived

Extracellular Vesicles: Emerging Mediators of Plant-Microbe Interactions. MPMI **36**: 218–

227

Xin X-F, Nomura K, Aung K, Velásquez AC, Yao J, Boutrot F, Chang JH, Zipfel C, He SY

(2016) Bacteria establish an aqueous living space in plants crucial for virulence. Nature

539: 524–529

Yadeta KA, Elmore JM, Creer AY, Feng B, Franco JY, Rufian JS, He P, Phinney B,

Coaker G (2017) A Cysteine-Rich Protein Kinase Associates with a Membrane Immune

Complex and the Cysteine Residues Are Required for Cell Death1. Plant Physiol **173**: 771–

787

Yeh Y-H, Panzeri D, Kadota Y, Huang Y-C, Huang P-Y, Tao C-N, Roux M, Chien H-C,

Chin T-C, Chu P-W, Zipfel C, Zimmerli L (2016) The Arabidopsis Malectin-Like/LRR-

RLK IOS1 Is Critical for BAK1-Dependent and BAK1-Independent Pattern-Triggered

Immunity. The Plant Cell **28**: 1701–1721

Zhang C, He M, Jiang Z, Liu L, Pu J, Zhang W, Wang S, Xu F (2022) The Xyloglucan

Endotransglucosylase/Hydrolase Gene XTH22/TCH4 Regulates Plant Growth by

Disrupting the Cell Wall Homeostasis in Arabidopsis under Boron Deficiency. Int J Mol Sci

23: 1250

Zhao P, Zhang F, Liu D, Imani J, Langen G, Kogel K-H (2017) Matrix metalloproteinases

operate redundantly in Arabidopsis immunity against necrotrophic and biotrophic fungal

pathogens. PLoS One **12**: e0183577

Zolg DP, Gessulat S, Paschke C, Graber M, Rathke-Kuhnert M, Seefried F, Fitzemeier K,

Berg F, Lopez-Ferrer D, Horn D, Henrich C, Huhmer A, Delanghe B, Frejno M (2021)

INFERYS rescoring: Boosting peptide identifications and scoring confidence of database

search results. Rapid Communications in Mass Spectrometry n/a: e9128

Figure legends

Figure 1. Principal Component Analysis (PCA) of flg22 and mock samples. The PCA plot

illustrates the distinct clustering of samples based on their treatment conditions. Flg22-treated

samples and mock-treated samples form separate clusters, indicating clear differences in their

protein abundance profiles. The first principal component (PC1) accounts for 27.9% of the total

variance, while the second principal component (PC2) explains 25.9% of the variance. Each data

point represents an individual biological replicate, with the spatial distribution of points within

each cluster indicating the degree of variability among replicates within the same treatment

group.

Figure 2. PTI-enriched proteins ranked by functional groups and relative to transcriptome

profiles. The heatmap presents flg22-enriched protein abundances on a red scale. The

corresponding log10 transformed transcript abundance is displayed on a blue-yellow scale, with

data are shown at 0, 1, 2, 3, 5, 9, and 18 hours post flg22 treatment relative to the initial time

point (0 hour). Transcriptome data of raw reads were obtained from study of Hillmer et al., 2017.

GEO data series GSE78735.

Figure 3. Peptide distribution analysis of receptor-like kinases (RLKs) in flg22-enriched

proteins. The schematic illustrates the mapping of identified peptides to PTI-enriched receptor-

like kinases (RLKs), represented as short black bars above the protein domain structures. Each

RLK is depicted with its extracellular region (green background) and the cytoplasmic kinase

region (pink background) on the right. The visualization reveals the majority of detected peptides

are localized to the extracellular regions of the RLKs, while the cytoplasmic regions, particularly

the kinase domains, show notably fewer peptide matches.

Figure 4. PTI-lower abundant proteins ranked by functional groups and relative to

transcriptome profiles. The heatmap illustrates proteins showing lower abundance in the flg22-

treated group compared to mock treatment, represented on a green scale. The corresponding

log10 transformed transcript abundance is displayed on a blue-yellow scale, with data are shown

at 0, 1, 2, 3, 5, 9, and 18 hours post flg22 treatment relative to the initial time point (0 hour).

Proteins associated with GO stress responses are highlighted in bold font.

Figure 5. Gene Ontology (GO) analysis of flg22-enriched and flg22-decreased abundance

protein profile. Dot plot represents significant GO terms of (A) 108 significantly enriched

proteins and (B) 255 significantly decreased proteins categorized into cellular component,

molecular function, and biological process. The x-axis represents the gene number of each

category, while the size of the circles indicates fold enrichment. Color intensity of the circles

corresponds to the statistical significance of enrichment ($-$log10 adjusted p-value). GO terms are

arranged on the y-axis, grouped by their respective ontologies (cellular component, molecular

function, and biological process).

Figure 6. Plant extracellular vesicles are enriched in flg22 treatment. (A) Size distribution of

flg22-P-EVs identified by Nano-flow cytometry (NFC). (B) Size distribution of mock-P-EVs

identified by Nano-flow cytometry (NFC). (C) Comparison of P-EV production between the

flg22 and mock groups, showing a significant increase in P-EV production in the flg22 group.

Statistical analysis was performed using two-tailed unpaired Student's t-test to determine P-EV

yield differences between groups with **P <0.01. (D) Confocal images of CFSE-stained P-EV

from mock samples. Scale bar=50μm.

Figures

Figure 1. Principal Component Analysis (PCA) of flg22 and mock samples. The PCA plot
illustrates the distinct clustering of samples based on their treatment conditions. Flg22-treated
samples and mock-treated samples form separate clusters, indicating clear differences in their
protein abundance profiles. The first principal component (PC1) accounts for 27.9% of the total
variance, while the second principal component (PC2) explains 25.9% of the variance. Each data
point represents an individual biological replicate, with the spatial distribution of points within
each cluster indicating the degree of variability among replicates within the same treatment
group.

Figure 2. PTI-enriched proteins ranked by functional groups and relative to transcriptome profiles. The heatmap presents flg22-enriched protein abundances on a red scale. The corresponding log10 transformed transcript abundance is displayed on a blue-yellow scale, with data are shown at 0, 1, 2, 3, 5, 9, and 18 hours post flg22 treatment relative to the initial time point (0 hour). Transcriptome data of raw reads were obtained from study of Hillmer et al., 2017. GEO data series GSE78735.

Figure 3. Peptide distribution analysis of receptor-like kinases (RLKs) in flg22-enriched proteins. The schematic illustrates the mapping of identified peptides to PTI-enriched receptor-like kinases (RLKs), represented as short black bars above the protein domain structures. Each RLK is depicted with its extracellular region (green background) and the cytoplasmic kinase region (pink background) on the right. The visualization reveals the majority of detected peptides are localized to the extracellular regions of the RLKs, while the cytoplasmic regions, particularly the kinase domains, show notably fewer peptide matches.

Figure 4. PTI-lower abundant proteins ranked by functional groups and relative to transcriptome profiles. The heatmap illustrates proteins showing lower abundance in the flg22-treated group compared to mock treatment, represented on a green scale. The corresponding log10 transformed transcript abundance is displayed on a blue-yellow scale, with data are shown at 0, 1, 2, 3, 5, 9, and 18 hours post flg22 treatment relative to the initial time point (0 hour). Proteins associated with GO stress responses are highlighted in bold font.

Figure 5. Gene Ontology (GO) analysis of flg22-enriched and flg22-decreased abundance protein profile. Dot plot represents significant GO terms of (A) 108 significantly enriched proteins and (B) 255 significantly decreased proteins categorized into cellular component, molecular function, and biological process. The x-axis represents the gene number of each category, while the size of the circles indicates fold enrichment. Color intensity of the circles corresponds to the statistical significance of enrichment (–log10 adjusted p-value). GO terms are arranged on the y-axis, grouped by their respective ontologies (cellular component, molecular function, and biological process).

Figure 6. Plant extracellular vesicles are enriched in flg22 treatment. (A) Size distribution of
flg22-P-EVs identified by Nano-flow cytometry (NFC). (B) Size distribution of mock-P-EVs
identified by Nano-flow cytometry (NFC). (C) Comparison of P-EV production between the
flg22 and mock groups, showing a significant increase in P-EV production in the flg22 group.
Statistical analysis was performed using two-tailed unpaired Student's t-test to determine P-EV
yield differences between groups with **P <0.01. (D) Confocal images of CFSE-stained P-EV
from mock samples. Scale bar=50μm.

Supporting figures

Figure S1. Cytoplasmic contamination detection of apoplastic washing fluid. Glucose-6-

phosphate dehydrogenase (G6PDH) enzymatic activity of AWF was tested for cytoplasmic

contamination in mock (blue stripe) or flg22 (red stripe) treatment compared to macerated

sample (solid). Data are presented as mean ± SD (n=3). Statistical analysis was performed using

a two-tailed unpaired Student's t-test. No significant difference was observed between mock and

flg22 treatments (P = 0.256).

Figure S2. Normalized protein abundances (Log10 transformed data) per sample. Mock
treatments are represented in blue boxes (M1-M4) and flg22 treatments are represented in pink
boxes (F1-F4). The y-axis represents the log10-transformed protein abundance. Box plots show
the distribution of protein abundances, with the box representing the interquartile range (IQR),
the horizontal line inside the box indicating the median, and whiskers extending to 1.5 times the
IQR. Outliers beyond the whiskers are depicted as individual points.

820