Moderating effects of cortical thickness, volume, and memory performance on age differences in neural reinstatement of scene information

Joshua M. Olivier[1]

Sabina Srokova[2]

Paul F. Hill[2]

Michael D. Rugg[1]

[1]Center for Vital Longevity and School of Behavioral and Brain Sciences, University of Texas at Dallas

1600 Viceroy Drive, Ste. 800

Dallas, TX 75235

[2]Department of Psychology and Evelyn F. McKnight Brain Institute, University of Arizona

Tucson, AZ 85721

Corresponding Author:

Joshua Olivier

joshua.olivier@utdallas.edu

(972) 883-3753

Center for Vital Longevity and School of Behavioral and Brain Sciences

University of Texas at Dallas

1600 Viceroy Drive, Ste. 800

Dallas, TX 75235

Abstract

The strength of neural reinstatement, a correlate of episodic memory retrieval, reportedly reflects the amount and fidelity of mnemonic content and is weaker in older than younger adults, especially for scene memoranda. Given evidence that age-related declines in cortical thickness and volume contribute to age-related cognitive decline, we analyzed fMRI data acquired from healthy young and older adults to examine relationships between cortical thickness, cortical volume, age, and scene-related reinstatement in the parahippocampal place area (PPA) and medial place area (MPA), two cortical regions implicated in scene processing. A 'reinstatement index' was estimated from fMRI data collected during tests of source memory for scene images, and multiple regression analyses were employed to examine the effects of the variables of interest on scene reinstatement. There were robust age differences in reinstatement, cortical thickness, and cortical volume. In both regions of interest, cortical volume fully mediated the effects of age on reinstatement. Additionally, PPA reinstatement strength predicted source memory performance independently of cortical volume or age. These findings suggest that age differences in scene reinstatement are mediated by cortical volume and that memory performance and cortical volume are associated with unique components of variance in reinstatement strength.

Keywords: episodic memory, fMRI, neural selectivity, neural broadening, structural MRI

Episodic memory, the ability to recollect information about previously experienced events, declines with increasing age (Nilsson 2003; Nyberg et al. 2012). Neuroimaging research has revealed robust evidence for age differences in a variety of functional correlates of episodic memory encoding and retrieval (e.g. Cabeza et al. 2002; Davis et al. 2008; Park et al. 2013; Srokova et al. 2022; Schott et al. 2023). One such correlate is neural reinstatement—retrieval-related neural patterns that partially overlap with those elicited during encoding (for reviews, see Danker and Anderson 2010; Rissman and Wagner 2012; Rugg et al. 2015). It has been reported that stronger reinstatement positively covaries with memory performance (Johnson et al. 2009; Gordon et al. 2014; Thakral et al. 2015; Trelle et al. 2020; Srokova et al. 2021), suggesting that the strength of reinstatement reflects the amount and quality of the retrieved mnemonic content. Additionally, it has sometimes been reported that the strength of neural reinstatement varies with age, with older adults showing weaker effects than young adults (e.g., McDonough et al. 2014; St-Laurent et al. 2014; Johnson et al. 2015; Abdulrahman et al. 2017; Folville et al. 2020; Trelle et al. 2020; for opposing results, see Dulas and Duarte 2012; Wang et al. 2016; Thakral et al. 2017; St-Laurent and Buchsbaum 2019; Thakral et al. 2019). In a study that examined neural reinstatement of face and scene images (Hill et al. 2021), age differences were only evident for scene reinstatement (see also Trelle et al. 2020). In keeping with this finding, based on a critical review of the literature, Rugg and Srokova (2024) concluded that the most secure evidence for age-related attenuation of reinstatement effects comes from studies that employed scene images as the critical memoranda. Here, we address the question of whether these age differences are associated with corresponding differences in brain structural metrics. We also examine whether reinstatement is associated across participants with memory performance and whether any such relationship is moderated by age.

Age differences in structural and functional brain metrics have primarily been studied separately, despite the extensive cross-sectional and longitudinal evidence indicating that structural

metrics such as cortical volume and thickness positively covary with cognitive abilities in older samples (Karama et al. 2014; Salthouse et al. 2015; Sun et al. 2016; Kranz et al. 2018; de Chastelaine et al. 2019; de Chastelaine et al. 2023). In a recent study, it was reported that age differences in gray matter volume mediated corresponding differences in fMRI 'subsequent memory effects,' a functional correlate of episodic memory encoding (Kizilirmak et al. 2024). By contrast, other studies examining relationships between structural and functional measures have reported little evidence for a mediating effect of structure (e.g. Boller et al. 2017; Hou et al. 2020; Hou et al. 2021; Kidwai et al. 2025). Overall, however, the extant literature is limited, and the circumstances under which correlations between brain function and structure can be identified remain unclear.

Motivated by this gap in knowledge, we examined the relationship between neural reinstatement of scene information, cortical thickness, and cortical volume in samples of young and older adults. We combined two independent datasets that comprised a total of 42 young and 40 older adults as they retrieved word-image pairs while undergoing fMRI. The combined dataset provided a substantial boost in statistical power relative to that associated with separate analyses of the two datasets. Our analyses focused on scene reinstatement effects in two regions that have been heavily implicated in visual scene processing—the parahippocampal place area (PPA) and medial place area (MPA) (Epstein and Baker 2019)—with the aim of examining the effects of age on scene reinstatement and the mediating effect, if any, of the structural metrics of cortical thickness and volume.

Materials and Methods

The present report examines the relationship between cortical thickness and volume and scene reinstatement by combining data from two independent, previously reported studies (Hill et al. 2021; Srokova et al. 2021). The experimental procedures and behavioral analyses for each study were detailed in those prior reports and are briefly summarized below for the convenience of the reader. Note that in Hill et al. (2021), neural reinstatement was indexed by multi-voxel pattern analysis rather than the

univariate metric employed in the present study, and the researchers did not differentiate between the PPA and MPA. In Srokova et al. (2021), unlike here, reinstatement was estimated for all correctly recognized test items regardless of the accuracy of the subsequent source memory judgment. Moreover, neither study reported brain structural metrics. Thus, none of the analyses of the functional and structural data reported below have been described previously.

Participants and Neuropsychological Testing

The present report describes analyses performed on 42 younger and 40 older adults across two independent studies. All participants were recruited from the Dallas metropolitan area. All were right-handed, had normal or corrected-to-normal vision, and were fluent English speakers before age 5. Potential participants were excluded if they reported a history of neurological or cardiovascular disease, diabetes, substance abuse, or usage of medication affecting the central nervous system. Demographic data for the combined participant sample can be found in Table 1.

Participants in both studies undertook an extensive neuropsychological test battery, including the Mini-Mental State Examination (MMSE), the California Verbal Learning Test-II (CVLT; Delis et al. 2000), Wechsler Logical Memory (Tests 1 and 2; (Wechsler 2009a), Trail Making (Tests A and B; Reitan and Wolfson 1985), the Symbol Digit Modalities test (SDMT; Smith 1982), the F-A-S subtest of the Neurosensory Center Comprehensive Evaluation for Aphasia (Spreen and Benton 1977), the Wechsler Adult Intelligence Scale-Revised (forward and backward digit span subtests; (Wechsler 1981), Category Fluency test (Benton 1968), Raven's Progressive Matrices (List 1; Raven et al. 2000), and either the Wechsler Test of Adult Reading (WTAR; Wechsler 2001) or the Wechsler Test of Premorbid Functioning (Wechsler 2009b). Participants were excluded if their performance was greater than 1.5 standard deviations below age norms for one or more memory-based tests or two or more non-memory tests.

Participants with scores of less than 27 on the MMSE were also excluded. For a description of the neuropsychological test battery scores in each study, see Hill et al. (2021) and Srokova et al. (2021).

Study 1: 27 young and 33 older adults were recruited to participate in Study 1. Three young and nine older participants were excluded from the initial sample due to inadequate memory performance, incidental MRI findings, technical difficulties, or voluntary withdrawal from the study. Two additional older participants were excluded from the analyses reported here due to the poor quality of their T1-weighted structural images caused by excessive movement. Thus, the final sample for Study 1 included 24 young (age range: 18–28 years) and 22 older adults (age range: 65–75 years).

Study 2: 25 young and 30 older adult participants were recruited in Study 2. Five young and six older participants were excluded from the initial sample due to not completing the scanning session because of claustrophobia or discomfort, technical difficulties, chance source memory performance, or incidental MRI findings. Additionally, two young and two older participants were excluded from the present analyses due to excessive movement during the T1-weighted scan. Four older adults participated in both studies; only their Study 1 data were included in the present analyses. Thus, the final sample for Study 2 included 18 young (age range: 18–30 years) and 18 older adults (age range: 65–75 years).

Table 1: Demographic data for participant sample by study (M ± SD)

	Young adults	Older adults	Group difference (p)
Study 1			
N	24	22	
Sex (F/M)	15/9	13/9	1.00[†]
Age (years)[‡]	22.42 ± 3.24	69.95 ± 3.36	
Years of Education[‡]	15.46 ± 2.65	16.77 ± 2.47	.089
Study 2			
N	18	18	
Sex (F/M)	10/8	7/11	.504[†]
Age (years)[‡]	24.39 ± 3.55	68.17 ± 2.60	
Years of Education[‡]	16.00 ± 1.71	17.61 ± 2.70	**.041***

[†]Result from a χ^2 test with Yates' continuity correction.

[‡]Result from one-way ANOVA revealed no significant differences between studies (Age: p = .830; Years of Education: p = .203).

Experimental materials and procedure

Participants in both studies underwent fMRI during a memory test (Figure 1). In both cases, participants first encoded concrete noun-visual image pairs. The words were paired with images of scenes or faces in Study 1 and with scenes or scrambled images in Study 2. The participants were required to imagine a scenario in which the object depicted by the word interacted with its paired image. They then rated the vividness of the imagined scenario on a 3-point scale ('not vivid' – 'somewhat vivid' – 'very vivid'). In the subsequent retrieval phase, studied and novel words were employed as test items in a two-step source memory task. The requirement was to indicate whether the word was new or old and, if endorsed old, to judge the image category the word had been paired with at study (i.e., a scene vs. a face in Study 1; a scene belonging to a designated 'target' category vs. a non-target scene or scrambled image in Study 2).

Study 1: Participants in Study 1 underwent fMRI during both the encoding and retrieval phases. Experimental items consisted of 288 concrete nouns paired with 96 images of faces and 96 images of scenes. The study phase was undertaken across four scanning runs, each containing 48 trials comprising words paired with an image (50% faces, 50% scenes) and 24 null trials, where a white fixation cross remained on the screen for the entire trial. A red fixation cross was presented for 500ms at the commencement of each trial. The word-image pair remained on the screen for 2000ms, followed by a white fixation cross for another 2000ms.

The test phase was also undertaken across four scanning runs, each comprising 48 studied words, 24 new words, and 24 null trials. Each trial began with a red fixation cross presented for 500ms, the test word for 2000ms, and a white fixation cross for another 2000ms. Participants were instructed to indicate whether the word was "Old" or "New" for each trial. If the endorsement was "Old," participants

were then required to make a source memory judgment, indicating whether the word had been paired

with an image of a face or a scene. A "don't know" option was included to discourage guessing.

Study 2: Participants in Study 2 underwent fMRI during the retrieval phase only. The out-of-

scanner study task employed 240 concrete nouns, 120 color images of scenes (50% urban, 50% rural),

and 60 scrambled (pixelated) scenes. Word-image pairs were presented in one of three randomized

display locations: 20 on the left side of the screen, 20 in the middle, and 20 on the right. Each study

phase trial began with a black fixation cross for 200ms in the location corresponding with the upcoming

word-image pair. The study word was then presented, joined by the background image 200ms later. The

word-image pair remained on the screen for 5500ms.

The test phase was undertaken across five scanner runs, each divided into two mini-blocks. In

the different mini-blocks, memory was tested for either the studied background or the studied location

of the test word. The analyses described below focus only on scene reinstatement associated with

background judgments. Each background task block contained 24 critical trials, comprising 18 studied

word-image pairs and six new trials, intermixed with eight null trials. Each trial began with a red fixation

cross presented for 500ms, followed by the test word, which remained on the screen for 2500ms.

During this time, participants indicated whether they recognized the word from the study phase by

responding "Yes" or "No" to the prompt "Did you see?". For test items endorsed as studied, participants

were then required to respond "Yes," "No," or "Not Sure" to either the prompt "Was it on a rural

scene?" or "Was it on an urban scene?" Task prompts were counterbalanced across participants such

that any given participant received only one of the two prompts for the duration of the study.

Figure 1: Overview of encoding and subsequent memory tasks for **A)** Study 1 and **B)** Study 2.

MRI data acquisition and preprocessing

Structural and functional MRI data were acquired with a Philips Achieva 3T MRI scanner (Philips Medical Systems, Andover, MA) equipped with a 32-channel head coil. Functional images were acquired using a T2*-weighted, blood oxygen level-dependent echoplanar imaging (EPI) sequence (sensitivity

encoding [SENSE] factor 2, flip angle = 70 deg., TE = 30ms, FOV = 24 cm). EPI volumes from Study 1

consisted of 34 slices with a voxel size of 3 × 3 × 3 mm acquired in ascending order (TR = 2s, 1 mm

interslice gap). In Study 2, EPI volumes comprised 44 slices acquired in an interleaved order at a voxel

size of 2.5 x 2.5 x 2.5 mm (TR = 1.6s, 0.5 mm interslice gap) and a multiband factor of 2. T1-weighted

anatomical images in both studies were acquired with a magnetization-prepared rapid gradient echo

(MPRAGE) pulse sequence (FOV = 240 x 240, 1 x 1 x 1 mm isotropic voxels, 160 slices, sagittal

acquisition). In the case of one young and eight older participants, the fMRI and MPRAGE data were

acquired on separate days, with a median of 5.2 months between the two visits.

The fMRI data were preprocessed using Statistical Parametric Mapping (SPM12, Wellcome

Department of Cognitive Neurology, London, UK) and custom MATLAB code. Functional images were

realigned to the mean EPI image, and slice-time was corrected using sinc interpolation to the middle

slice. The images were then reoriented along the anterior and posterior commissures and spatially

normalized to MNI space using a sample-specific EPI template following the procedures described by de

Chastelaine et al. (2011, 2016). During the normalization step, the functional images from Study 2 were

downsampled to a voxel size of 3 × 3 × 3 mm, thus matching the resolution of the functional images

acquired in Study 1.

Structural MRI analysis

Estimation of whole-brain cortical thickness. Estimates of whole-brain cortical thickness were

derived from the T1-weighted image of each participant using Freesurfer's (v6.0) semi-automatic

processing pipeline (https://surfer.nmr.mgh.harvard.edu/fswiki; (Dale et al. 1999; Fischl and Dale 2000;

Fischl et al. 2002). Each 3-dimensional T1 volume underwent skull-stripping and intensity normalization,

followed by reconstruction as an inflated 2-dimensional surface to identify white matter and pial

surfaces. White matter and pial surfaces were visually inspected by one of six trained raters. All raters

were blind to participant age. The raters corrected tissue identification errors by editing the brain masks

returned by the first pass of the automated analysis. 'Control points' were also added to voxels on white

matter paths that the Freesurfer pipeline had not appropriately labeled. Brain regions frequently

requiring manual edits included the boundaries between grey matter and the pial surface in the

temporal cortex, insula, and orbitofrontal cortex and the delineation between the cortex and the

cerebellum. A second rater inspected each scan for quality control purposes. Cortical thickness was

defined as the distance between the pial surface and the grey/white matter boundary on a vertex-by-

vertex basis for the cortical mantle of each hemisphere (de Chastelaine et al. 2023). Each hemisphere's

mean cortical thickness values were averaged to obtain mean whole-brain cortical thickness.

Estimation of whole-brain cortical volume and total intracranial volume. The "CortexVol" and

"estimated Total Intracranial Volume" measures output by Freesurfer were used as estimates of whole-

brain cortical volume and total intracranial volume (ICV), respectively. Before statistical analysis, whole-

brain cortical volume estimates were adjusted for total intracranial volume to account for individual

differences in head size. A linear regression was performed to make this adjustment, with whole-brain

cortical volume as the dependent variable and total intracranial volume as the independent variable.

Residuals were computed by subtracting the predicted cortical volume (based on total ICV) from the

actual cortical volume for each participant. These adjusted cortical volume estimates were used as the

primary volume variable for subsequent analyses. Hence, all references below to cortical volume refer

to these adjusted estimates.

Note that we focus on whole brain measures of cortical thickness and volume because of the

evidence that regional structural measures are highly inter-correlated and, individually, demonstrate

unreliable associations with cognitive metrics (Salthouse et al. 2015; Kranz et al. 2018; Masouleh et al.

2019; Hou et al. 2021; Krogsrud et al. 2021; Masouleh et al. 2022; but, see Lee et al. 2016). Analyses that

used volumetric metrics of the cortical parcels that roughly correspond with the regions occupied by our regions of interest (ROIs) are reported in Supplementary materials.

Functional MRI analysis

ROI selection. We selected *a priori* two scene-selective ROIs for the analyses namely, the parahippocampal place area (PPA) and the medial place area (MPA), as shown in Figure 2. The ROIs were derived from an unpublished independent localizer dataset in which participants viewed images of objects, faces, and scenes. The ROIs were defined using an across-participant scene > object + face contrast at a height threshold of p < 0.05 FWE-corrected. The PPA was defined by restricting the scene-selective cluster to the fusiform and parahippocampal gyrus labels in the Neuromorphometrics atlas available in SPM12. As Neuromorphometrics does not provide an accurate anatomical mask for the MPA, the ROI was defined by inclusively masking the contrast with the outcome of a search in the Neurosynth database using the term "retrosplenial" (search in August 2019; search results were FDR-corrected at p < 0.00001; Yarkoni et al. 2011). The two functional ROIs are illustrated in Figure 2.

Figure 2: Regions of interest: Reinstatement was estimated in the parahippocampal place area (PPA) and the medial place area (MPA).

Reinstatement index

The unsmoothed fMRI data from each participant were concatenated and subjected to a least-squares-all GLM (Rissman et al. 2004; Mumford et al. 2014) in which each trial is modeled with its own regressor in a single design matrix. Every trial was modeled with a delta function time-locked to the onset of the test item and convolved with SPM's canonical hemodynamic response function. Covariates of no interest comprised six motion regressors reflecting rigid body translation and rotation and regressors modeling the session means.

The single trial blood-oxygen-level-dependent (BOLD) parameter estimates obtained from the GLM were employed to compute participant-specific reinstatement indices that quantified the strength of scene reinstatement in each ROI. We extracted the average across-voxel BOLD amplitude for each trial in each ROI, collapsing across the hemispheres. The reinstatement index was computed by subtracting the across-trial mean BOLD responses elicited by test words paired with scrambled images or faces from the mean response elicited by test words paired with the scenes, divided by their pooled standard deviation. Only trials associated with correct source judgments were employed to estimate these indices.

$$Reinstatement\ Index = \frac{\mu_{scene} - \mu_{face/scrambled}}{\sqrt{\dfrac{\sigma^2_{scene} + \sigma^2_{face/scrambled}}{2}}}$$

A positive reinstatement index signifies a greater mean BOLD response for words paired with scenes than for words paired with the alternate image category. Thus, the reinstatement index quantifies retrieval-related neural reinstatement of scene-related information. Of importance, because of the scaling function, the reinstatement index is insensitive to between-participant and between-group differences in the gain of the hemodynamic response function (see, for example, Liu et al. 2013).

Statistical Analysis

The relationships between age, neural reinstatement, whole-brain cortical thickness, and whole-brain cortical volume (adjusted for total intracranial volume) were examined using a series of multiple regression analyses employing the factors of thickness, volume, age group, study, and their interactions as predictors of reinstatement. In cases where a data point deviated by > 2.5 standard deviations from the sample mean, the analyses were re-run after excluding the relevant participant.

Results

Behavioral results

The behavioral results from the two studies have been reported previously (Hill et al. 2021; Srokova et al. 2021) and are only briefly summarized below. Item recognition (*Pr*) was defined as the difference between the probability of a recognition hit and a false alarm. In Study 1, item recognition was lower in older than young adults (p = .003), whereas no age differences were identified in Study 2 (p < .218) (see Table 2).

Source memory performance, operationalized as the probability of source recollection (*pSR*), was defined according to a single high-threshold model (Snodgrass and Corwin 1988) that included a correction for guessing (Park et al. 2008):

$$pSR = \frac{pSource\ Correct - 0.5(1 - pDon't\ Know)}{1 - 0.5(1 - pDon't\ Know)}$$

"*pSource Correct*" and "*pDon't Know*" refer to the proportion of correctly recognized old trials associated with correct source judgments or a don't know response, respectively. *pSR* was significantly lower in older than in young adults in the Study 1 sample (p < .001), but the age group differences in Study 2 did not reach significance (p = .075) (see Table 2).

Table 2: Item and source memory (M ± SD)

	Young adults	Older adults
Study 1		
Source memory (*pSR*)	0.683 ± 0.179	0.502 ± 0.160
Item memory (*Pr*)	0.682 ± 0.173	0.540 ± 0.130
Study 2		
Source memory (*pSR*)	0.489 ± 0.200	0.364 ± 0.211
Item memory (*Pr*)	0.639 ± 0.185	0.568 ± 0.153

Age differences in scene reinstatement

A 2 (age group) x 2 (ROI) x 2 (study) mixed effects ANOVA was conducted on the reinstatement indices derived from the PPA and MPA ROIs, the outcome of which is reported in Table 3 and illustrated in Figure 3. The ANOVA revealed a significant main effect of age group, while the remaining main effects and the 2-way and 3-way interactions failed to reach significance. Thus, scene-related reinstatement was robustly weaker in the older than the young adults in both ROIs and both studies. Reinstatement effects were, however, robustly greater than zero in both age groups (collapsed across ROIs and studies, $p < .001$ in both cases).

Table 3: Outcome of an age group x ROI x study ANOVA on reinstatement

	df	F	η_p^2	p-value
Age group	1,78	12.490	.138	**<.001***
Study	1,78	0.027	.000	.869
ROI	1,78	1.323	.017	.254
Age x study	1,78	0.885	.011	.350
Age x ROI	1,78	0.111	.001	.740
ROI x study	1,78	0.742	.009	.392
Age x ROI x study	1,78	0.145	.002	.705

Figure 3: Distribution of reinstatement indices in Study 1 and Study 2 by age group in (A) the PPA and (B) the MPA. Black dots and bars represent means and ±95% confidence intervals. Removal of outliers (circled) did not change the outcome of the statistical analyses (age group effect p < .001).

Age differences in cortical thickness and volume

We employed 2 (age group) × 2 (study) between-subjects ANOVAs to examine age group differences in cortical thickness and volume estimates. The outcomes of these analyses are reported in Table 4 and illustrated in Figure 4.

As evident from Table 4, the ANOVA of thickness estimates revealed a significant main effect of age group, whereby thickness was lower in the older than the young adult group. Neither the main effect of study nor the age group by study interaction were significant. Thus, neither the thickness estimates nor the moderating effect of age differed significantly between the two studies. Paralleling the thickness analysis, the ANOVA of volume estimates revealed a significant main effect of age group, such that volume was lower in the older than the young adult group. There was also a significant main

effect of study, with mean volume estimates lower in Study 1 than in Study 2; the reason for this

difference is unclear. The age group by study interaction was not significant.

Table 4: Outcome of the Age Group x Study ANOVAs of cortical thickness and volume.

	df	F	η_P^2	p-value
Cortical thickness				
Age group	1,81	84.666	.520	**<.001***
Study	1,81	1.595	.020	.210
Age x study	1,81	2.288	.028	.134
Cortical volume				
Age group	1,81	94.128	.547	**<.001***
Study	1,81	8.679	.100	**.004***
Age x study	1,81	0.412	.005	.523

Figure 4: Distribution of estimates by age group of **(A)** cortical thickness and **(B)** cortical volume, combined across Study 1 and Study 2. The volumetric data are residualized for both ICV and study. Black dots and bars represent means and ±95% confidence intervals. Removal of outliers (circled) did not change the outcome of the statistical analyses (main effects of age p < .001 for both thickness and volume).

Cortical thickness as a predictor of scene reinstatement

Multiple regression was employed to examine the relationships between age group, cortical thickness, and scene reinstatement in each ROI. The initial regression models employed age group, study, thickness, the interactions between age group and thickness, and study and thickness as predictors. The interaction terms were non-significant for both the PPA (age x thickness: p = .391; study x thickness: p = .998) and MPA (age x thickness: p = .939; study x thickness: p = .265) models, and hence the models were re-run after excluding them. The outcomes from these reduced models are presented

in Table 5, where it can be seen that thickness was positively correlated with reinstatement in the PPA

only (see Figure 5).

Table 5: Outcome of linear regression analyses testing whether cortical thickness, age group, and study significantly predicted reinstatement in the PPA and MPA.

	b	SE *b*	β	t	Partial corr.	p-value
PPA reinstatement						
Cortical thickness	.836	.372	.340	2.251	.247	**.027***
Age group	-.051	.095	-.081	-.535	-.060	.594
Study	-.004	.066	-.007	-.065	-.007	.949
MPA reinstatement						
Cortical thickness	.248	.307	.124	.808	.091	.422
Age group	-.136	.078	-.267	-1.748	-.194	.084
Study	-.042	.055	-.081	-.761	-.086	.449

Note: *b,* unstandardized coefficient; SE *b*, standard error of the unstandardized coefficient; β, standardized coefficient.

Figure 5: Scatter plots of the relationships between cortical thickness and scene reinstatement in the **(A)** PPA and **(B)** MPA. The shaded area around each regression line represents the 95% confidence interval. Removal of outliers (circled) did not change the outcome of the statistical analyses (p = .045 for the association between thickness and PPA reinstatement).

Cortical volume as a predictor of scene reinstatement

We used an analogous regression approach to examine the relationships between cortical volume and reinstatement in the PPA and MPA. The age x volume and study x volume interaction terms were again non-significant in the full models for both the PPA (age x volume: p = 0.941; study x volume: p = 0.493) and MPA (age x volume: p = 0.353; study x volume: p = 0.845) and hence were dropped from the final models. The outcomes from these models are described in Table 6 and illustrated in Figure 6. As is evident from the table, cortical volume was a significant predictor of reinstatement in both the PPA and MPA.

Table 6: Outcome of multiple linear regression analyses testing whether cortical volume, age, and study significantly predicted reinstatement in the PPA and MPA.

	b	SE *b*	β	t	Partial corr.	p-value
PPA reinstatement						
Cortical volume	.158	.049	.497	3.238	.344	**.002***
Age group	.022	.094	.035	.233	.026	.816
Study	-.052	.067	-.082	-.779	-.088	.439
MPA reinstatement						
Cortical volume	.113	.040	.439	2.865	.309	**.005***
Age group	-.019	.077	-.037	-.244	-.028	.808
Study	-.084	.054	-.164	-1.554	-.173	.124

Note: *b,* unstandardized coefficient; SE *b*, standard error of the unstandardized coefficient; β, standardized coefficient.

Figure 6: Scatter plots of the relationships between cortical volume and scene reinstatement in the **(A)** PPA and **(B)** MPA. The shaded area around each regression line represents the 95% confidence interval. Removal of outliers (circled) did not change the outcome of the statistical analyses (ps = .015 and .008 for the associations between volume and PPA and MPA reinstatement, respectively).

In a follow-up analysis, we examined whether cortical thickness and volume were independent predictors of reinstatement in the PPA, the ROI where each metric correlated reliably with reinstatement. The outcome of the model is presented in Table 7, where it is evident that only volume accounted for a significant fraction of the variance in reinstatement.

Table 7: Outcome of multiple regression analyses predicting reinstatement in the PPA given the predictors cortical volume, cortical thickness, age group, and study.

	b	SE *b*	β	t	Partial corr.	p-value
PPA reinstatement						
Cortical volume	.134	.054	.422	2.465	.248	**.016***
Cortical thickness	.399	.401	.162	.994	.100	.323
Age group	.061	.102	.098	.600	.060	.550
Study	-.051	.067	-.081	-.769	-.087	.444

Note: *b,* unstandardized coefficient; SE *b*, standard error of the unstandardized coefficient; β, standardized coefficient.

Contributions of broadening and attenuation to age and volume effects on reinstatement

To further elucidate the findings for scene reinstatement, we analyzed the magnitudes of the BOLD responses elicited by the test words studied with scenes vs. the 'unpreferred' image category (i.e., faces in Study 1, scrambled scenes in Study 2; see Figure 7 for plots of the data pooled across experiments and ROIs). We collapsed across ROIs and hemispheres as these factors did not interact with the factors of study or age group in the across-experiment analysis of the reinstatement indices. Thus, the data were subjected to a 2 (study) x 2 (age group) x 2 (image category) mixed-effects ANOVA. The ANOVA gave rise to main effects of study ($F_{1,78} = 36.51$, $p < .001$, $\eta^2 = .319$) and image category ($F_{1,78} = 84.48$, $p < .001$, $\eta^2 = .520$). The latter effect was modified by an age group x category interaction ($F_{1,78} = 16.87$, $p < .001$, $\eta^2 = .178$). As is evident from Figure 7, the interaction reflected an absence of age differences in the responses elicited by the test items paired with scenes ($p = .383$) and a robust difference in favor of the older group in the responses to items paired with the unpreferred image categories ($t_{80} = 3.85$, $p < .001$, $d = .848$). Therefore, these findings indicate that the age-related attenuation of scene reinstatement reported above was a consequence of an age-related enhancement of the BOLD responses elicited by test items paired with the unpreferred images.

Using the same regression approach employed with the reinstatement index, we went on to examine the associations between the BOLD responses elicited by test items paired with scenes and the alternate image categories, age, and cortical volume. In the case of the parameter estimates associated with scene trials, the variables of study, age group, and volume were all significant predictors of the size of the estimates (partial $r = .395$, $p < .001$, .301, $p = .007$, and .314, $p = .005$, respectively). Study and age group were also significant predictors of the size of the responses elicited by the test items paired with the unpreferred categories (partial $r = .573$, $p < .001$, and .384, $p < .001$, respectively), but cortical volume was not significant (partial $r = .087$, $p = .442$). The outcomes of these analyses did not change with the removal of outliers with the exception that age group was no longer a significant predictor of

the estimates associated with scene trials (partial r = .176, p = .127). Thus, the relationship between

scene reinstatement and cortical volume reported above reflects the moderating effect of volume on

BOLD responses associated with scene retrieval.

Figure 7: Plot of the mean beta estimates (collapsed across ROIs) for test words studied with scenes and

unpreferred image categories for the young and older age groups. Black dots and bars represent means

and ±95% confidence intervals. Removal of outliers did not change the outcome of the statistical

analyses (p = .610 for the effect of age on scene trials, p < .001 for the effect of age on unpreferred

trials).

Cortical thickness, cortical volume, and reinstatement as predictors of memory performance

Lastly, we ran a series of multiple regressions to examine whether source memory performance

(pSR) was predicted by reinstatement, cortical thickness, or cortical volume. pSR was residualized by

study to remove study-specific differences in performance. Age group and its interactions with study

and reinstatement were included as predictors in the initial models. In no case were these interactions

significant (ps > .340). As is evident from Table 8 (see also Figure 8), PPA reinstatement and age group

were both significant predictors of memory performance. The relationship remained when cortical

volume and thickness were each added to the model. By contrast, MPA reinstatement did not

significantly predict memory performance (p = .137). Furthermore, after controlling for age group and

study, neither cortical thickness nor volume were, by themselves, significant predictors of memory

performance (r = .170 and .169, respectively, ps > .130).

Figure 8: Scatter plot of the relationship between PPA reinstatement and memory performance (pSR)

after controlling for age group and study. The shaded area around the regression line represents the

95% confidence interval. Removal of the outlier (circled) did not change the outcome of the statistical

analyses (p = .013 for the association between PPA reinstatement and memory performance).

Table 8: Outcome of multiple regression analyses predicting memory performance given the predictors PPA reinstatement, age group, and study. An additional model further includes cortical volume and thickness as predictors.

	b	SE b	β	t	Partial corr.	p-value
Memory performance (pSR)						
PPA reinstatement	.968	.328	.307	2.949	.317	**.004***
Age group	-.593	.206	-.300	-2.879	-.310	**.005***
Study	.002	.196	.001	.009	.001	.993
Memory performance (pSR)						
PPA reinstatement	.869	.354	.276	2.451	.271	**.017***
Cortical volume	.047	.175	.047	.266	.030	.791
Cortical thickness	.879	1.256	.114	.700	.080	.486
Age group	-.383	.319	-.194	-1.203	-.137	.233
Study	-.038	.209	-.019	-.183	-.021	.855

Note: *b,* unstandardized coefficient; SE *b*, standard error of the unstandardized coefficient; β, standardized coefficient.

Discussion

We examined associations between retrieval-related neural reinstatement of scene information in the PPA and MPA, whole brain cortical thickness and volume, and memory performance in samples of young and older adults. As expected, given prior findings (Rugg and Srokova 2024), there were robust age differences in reinstatement strength in both ROIs. Also consistent with the prior literature, cortical volume and thickness were markedly lower in the older age group. Multiple regression revealed associations between each of the structural metrics and scene reinstatement, although the association with thickness was restricted to the PPA. Moreover, only volume accounted for a significant fraction of the variance in PPA reinstatement when the two structural metrics were included in the same regression model. Additionally, the effect of age on scene reinstatement in both ROIs was fully mediated by cortical volume. Lastly, reinstatement strength in the PPA was associated with source memory performance independently of age and cortical volume. Below, we discuss the implications of these findings.

As noted above, the finding that older adults demonstrate weaker scene reinstatement than young participants is unsurprising, given prior reports. The present findings do, however, shed additional light on prior findings, most prominently through the association that was identified between cortical volume and scene reinstatement. The association was highly robust (partial r controlling for age group and study for the combined ROIs = .375, p < .001) and, strikingly, cortical volume fully mediated the association between age group and reinstatement (partial r = .003 after controlling for volume and study). Significantly, the relationship between scene reinstatement and cortical volume remained when the analysis was restricted to the young age group only (partial r = .377, p = .017, after controlling for chronological age and study; note that volume and chronological age also correlated reliably in this age group, partial r = -.327, p = .037).

How does the finding that scene reinstatement covaries not with age but with cortical volume contribute to the understanding of the determinants of the strength of reinstatement? One obvious implication is that prior findings of age-related reductions in scene reinstatement (Johnson et al. 2015; Trelle et al. 2020) may have arisen because of the confounding effects of cortical volume. This leads directly to the question: why should volume be associated with reinstatement? A full answer to this question will require a more comprehensive understanding of the neurobiological determinants of MRI volumetric measures than exists currently. For example, an elucidation of the relative contributions of different microstructural factors to cortical volumetric measures (such as neuron density, size, and arborization, glial and myelin content, etc.) may bring us closer to an answer. The present findings do, however, offer some clues.

First, the finding that the association between reinstatement and cortical volume is age-invariant indicates that while the impact of age on reinstatement is likely attributable to the well-documented finding that cortical volume (and, indeed, total gray matter) declines monotonically across most of the human lifespan (Bethlehem et al. 2022), there is nothing in the present data to suggest that the association strengthens in later life. Thus, the factors responsible for the association are unlikely to include those (such as neuronal loss; Goel et al. 2022) that are responsible for the reductions in cortical volume caused by age-related pathologies like Alzheimer's Disease. Of course, this is not to say that reinstatement strength would not be impacted by such pathologies— this is a question that would be well worth pursuing—but as is evident from the findings from the young age group described above, neuropathological factors do not appear to be the mechanism responsible for variation in reinstatement strength in cognitively healthy adults.

Another clue to the factors responsible for the association between reinstatement and cortical volume comes from the analyses that were conducted on the BOLD responses elicited by the test items paired with the scene and unpreferred image categories. The analyses revealed that volume moderated

reinstatement strength specifically through its impact on functional activity associated with scene retrieval. A speculative explanation for the specificity of this association is that volume acts as a moderator of the dynamic range of BOLD responses elicited in scene-selective neural populations. Whether volume exerts this moderating effect at the neural or the hemodynamic level (or both) remains to be established.

As just alluded to, the analysis of the BOLD responses associated with the scene and unpreferred image categories clearly demonstrated that age-related attenuation of scene reinstatement is a consequence of neural 'broadening' (heightened responses to exemplars of an 'unpreferred' stimulus category) rather than neural 'attenuation' (diminished responses to exemplars belonging to a 'preferred' category). Both patterns of findings have been reported previously in studies describing age-related reductions in the selectivity of BOLD responses to preferred stimulus categories in category-selective cortical regions (e.g. Chee et al. 2006; Park et al. 2012; Koen et al. 2019; Srokova et al. 2020; for review, see Koen and Rugg 2019). However, few studies have examined this issue when category information is retrieved from memory rather than directly perceived (for a recent example, see Hill et al. 2021). The present findings are consistent with evidence (reviewed in Koen and Rugg 2019) that age-related reductions in neural selectivity are a consequence of broader 'tuning' functions in category-selective cortex and suggest that the proposal is equally applicable to the neural selectivity of mnemonic representations.

For the reasons noted in the Introduction, we elected to use whole brain structural metrics when conducting our primary analyses examining possible relationships between brain structure and scene reinstatement. As already discussed, the analyses identified a robust association between whole brain cortical volume and reinstatement in both the PPA and MPA. As reported in the supplementary materials, when these analyses were repeated using as metrics the volumes of cortical regions that roughly corresponded with the regions occupied by our ROIs, no associations with reinstatement were

evident. These null findings are open to at least two interpretations. First, they might signify that reinstatement is sensitive to a generic (trans-region) characteristic of the cortex that is represented most reliably by the whole brain measure. By this argument, the specific microstructural properties of the PPA and MPA are less important determinants of reinstatement strength than a generic factor captured by whole brain volume. Alternately, reinstatement might indeed depend upon specific regional characteristics, but these are sufficiently highly correlated across regions to lead to an association at the whole brain level, while the single region metrics are insufficiently reliable to yield comparable findings (cf. Cox et al. 2019; Masouleh et al. 2019). Resolution of this issue will require research that employs more fine-grained structural analyses than those employed here.

Echoing several prior reports that metrics of neural selectivity derived from the PPA are associated with memory performance (e.g. Koen et al. 2019; Srokova et al. 2020; Srokova et al. 2024), we identified a robust, age- and study-invariant across-participants relationship between PPA scene reinstatement and source memory performance. These findings are consistent with the proposal that the PPA plays a key role in the encoding and retrieval of episodic information, perhaps because of its contribution to the processing of mnemonically relevant contextual information (Eichenbaum et al. 2007; Aminoff et al. 2013). Of importance, this association was unmoderated by cortical volume. Thus, the variance in scene reinstatement shared with memory performance was independent of the variance shared with cortical volume (and, by extension, age group). Identifying the source (or sources) of the first of these variance components could help elucidate the determinants of individual differences in memory function.

Limitations

The present study shares some of the limitations of the two studies that contributed to its dataset. One important limitation comes from the employment of cross-sectional designs.

Consequently, any effects of age group cannot be attributed unambiguously to the effects of aging rather than one or more confounding factors such as selection bias or cohort effects (Rugg 2017). A second limitation is the relatively small sample sizes. Whereas these are, of course, larger than those of the parent studies, our analyses remain underpowered to detect small effects. Lastly, while the reinstatement metric that was employed in our primary analyses should be impervious to age and individual differences in the gain of the hemodynamic transfer function, this is not the case for the analyses of the BOLD responses that contributed to the metric. Therefore, we cannot rule out the possibility that some of the age differences identified in those analyses are attributable to age differences manifest at the neurovascular rather than the neural level.

Conclusions

The present findings advance understanding of the impact of age on retrieval-related neural reinstatement of scene information in two major ways. First, the findings suggest that prior reports of age differences in scene reinstatement strength are likely due to the confounding influence of cortical volume. Here, we found that after controlling for volume, age explained only a trivial and non-significant fraction of the variance in reinstatement. Second, the robust across-participants association we identified between reinstatement strength and memory performance was unmodified by either cortical volume or age group. Therefore, we conclude that there are two independent factors contributing to across-participant variability in reinstatement strength. One of these factors is reflected in brain structural metrics such as cortical volume, which explains the association between reinstatement and age. The other factor, the determinants of which are unknown, is associated with some aspect of cognitive function that contributes to individual differences in performance on the experimental memory test (and, perhaps, performance on other cognitive tests also; cf. Trelle et al. 2020).

It will be of considerable interest to see how well the present findings generalize to other experimental settings and to memoranda that elicit neural reinstatement in other category-selective cortical regions.

Funding: This work was supported by the National Institute on Aging (grant numbers R01AG082680, RF1AG039103), the National Science Foundation (grant number 1633873), and the postdoctoral training program of the Arizona Alzheimer's Consortium (grant number T32AG044402).

Acknowledgments: We gratefully acknowledge the contributions of Marianne de Chastelaine, Mingzhu Hou, Amber Kidwai, Seham Kafafi, and Mel Racenstein to the analysis of the structural MRI data and to the experimental volunteers who contributed their time to these studies.

Conflict of interest statement: None declared.

Address of corresponding author:

Joshua Olivier

1600 Viceroy Dr. #800

Dallas, TX 75325

joshua.olivier@utdallas.edu

(972) 883-3753

References

Abdulrahman H, Fletcher PC, Bullmore E, Morcom AM. 2017. Dopamine and memory dedifferentiation in aging. NeuroImage. 153:211–220. doi:10.1016/j.neuroimage.2015.03.031.

Aminoff EM, Kveraga K, Bar M. 2013. The role of the parahippocampal cortex in cognition. Trends in Cognitive Sciences. 17(8):379–390. doi:10.1016/j.tics.2013.06.009.

Benton AL. 1968. Differential behavioral effects in frontal lobe disease. Neuropsychologia. 6(1):53–60. doi:10.1016/0028-3932(68)90038-9.

Bethlehem R a. I, Seidlitz J, White SR, Vogel JW, Anderson KM, Adamson C, Adler S, Alexopoulos GS, Anagnostou E, Areces-Gonzalez A, et al. 2022. Brain charts for the human lifespan. Nature. 604(7906):525–533. doi:10.1038/s41586-022-04554-y.

Boller B, Mellah S, Ducharme-Laliberté G, Belleville S. 2017. Relationships between years of education, regional grey matter volumes, and working memory-related brain activity in healthy older adults. Brain Imaging and Behavior. 11(2):304–317. doi:10.1007/s11682-016-9621-7.

Cabeza R, Anderson ND, Locantore JK, McIntosh AR. 2002. Aging Gracefully: Compensatory Brain Activity in High-Performing Older Adults. NeuroImage. 17(3):1394–1402. doi:10.1006/nimg.2002.1280.

de Chastelaine M, Donley BE, Kennedy KM, Rugg MD. 2019. Age moderates the relationship between cortical thickness and cognitive performance. Neuropsychologia. 132:107136. doi:10.1016/j.neuropsychologia.2019.107136.

de Chastelaine M, Mattson JT, Wang TH, Donley BE, Rugg MD. 2016. The neural correlates of recollection and retrieval monitoring: Relationships with age and recollection performance. NeuroImage. 138:164–175. doi:10.1016/j.neuroimage.2016.04.071.

de Chastelaine M, Srokova S, Hou M, Kidwai A, Kafafi SS, Racenstein ML, Rugg MD. 2023. Cortical thickness, gray matter volume, and cognitive performance: a crosssectional study of the moderating effects of age on their interrelationships. Cerebral Cortex. 33(10):6474–6485. doi:10.1093/cercor/bhac518.

de Chastelaine M, Wang TH, Minton B, Muftuler LT, Rugg MD. 2011. The Effects of Age, Memory Performance, and Callosal Integrity on the Neural Correlates of Successful Associative Encoding. Cerebral Cortex. 21(9):2166–2176. doi:10.1093/cercor/bhq294.

Chee MWL, Goh JOS, Venkatraman V, Tan JC, Gutchess A, Sutton B, Hebrank A, Leshikar E, Park D. 2006. Age-related Changes in Object Processing and Contextual Binding Revealed Using fMR Adaptation. Journal of Cognitive Neuroscience. 18(4):495–507. doi:10.1162/jocn.2006.18.4.495.

Cox SR, Ritchie SJ, Fawns-Ritchie C, Tucker-Drob EM, Deary IJ. 2019. Structural brain imaging correlates of general intelligence in UK Biobank. Intelligence. 76:101376. doi:10.1016/j.intell.2019.101376.

Dale AM, Fischl B, Sereno MI. 1999. Cortical Surface-Based Analysis: I. Segmentation and Surface Reconstruction. NeuroImage. 9(2):179–194. doi:10.1006/nimg.1998.0395.

Danker JF, Anderson JR. 2010. The ghosts of brain states past: Remembering reactivates the brain regions engaged during encoding. Psychological Bulletin. 136(1):87–102. doi:10.1037/a0017937.

Davis SW, Dennis NA, Daselaar SM, Fleck MS, Cabeza R. 2008. Qué PASA? The Posterior–Anterior Shift in Aging. Cerebral Cortex. 18(5):1201–1209. doi:10.1093/cercor/bhm155.

Delis DC, Kramer JH, Kaplan E, Ober BA. 2000. California verbal learning test, adult version (CVLT-II). Cleveland, Ohio: The Psychological Corporation.

Dulas MR, Duarte A. 2012. The Effects of Aging on Material-Independent and Material-Dependent

Neural Correlates of Source Memory Retrieval. Cerebral Cortex. 22(1):37–50.

doi:10.1093/cercor/bhr056.

Eichenbaum H, Yonelinas AP, Ranganath C. 2007. The Medial Temporal Lobe and Recognition Memory.

Annual Review of Neuroscience. 30(Volume 30, 2007):123–152.

doi:10.1146/annurev.neuro.30.051606.094328.

Epstein RA, Baker CI. 2019. Scene Perception in the Human Brain. Annual Review of Vision Science.

5(1):373–397. doi:10.1146/annurev-vision-091718-014809.

Fischl B, Dale AM. 2000. Measuring the thickness of the human cerebral cortex from magnetic

resonance images. Proceedings of the National Academy of Sciences. 97(20):11050–11055.

doi:10.1073/pnas.200033797.

Fischl B, Salat DH, Busa E, Albert M, Dieterich M, Haselgrove C, van der Kouwe A, Killiany R, Kennedy D,

Klaveness S, et al. 2002. Whole Brain Segmentation: Automated Labeling of Neuroanatomical

Structures in the Human Brain. Neuron. 33(3):341–355. doi:10.1016/S0896-6273(02)00569-X.

Folville A, Bahri MA, Delhaye E, Salmon E, D'Argembeau A, Bastin C. 2020. Age-related differences in the

neural correlates of vivid remembering. NeuroImage. 206:116336.

doi:10.1016/j.neuroimage.2019.116336.

Goel P, Chakrabarti S, Goel K, Bhutani K, Chopra T, Bali S. 2022. Neuronal cell death mechanisms in

Alzheimer's disease: An insight. Front Mol Neurosci. 15. doi:10.3389/fnmol.2022.937133.

Gordon AM, Rissman J, Kiani R, Wagner AD. 2014. Cortical Reinstatement Mediates the Relationship Between Content-Specific Encoding Activity and Subsequent Recollection Decisions. Cerebral Cortex. 24(12):3350–3364. doi:10.1093/cercor/bht194.

Hill PF, King DR, Rugg MD. 2021. Age Differences In Retrieval-Related Reinstatement Reflect Age-Related Dedifferentiation At Encoding. Cerebral Cortex. 31(1):106–122. doi:10.1093/cercor/bhaa210.

Hou M, de Chastelaine M, Donley BE, Rugg MD. 2021. Specific and general relationships between cortical thickness and cognition in older adults: a longitudinal study. Neurobiology of Aging. 102:89–101. doi:10.1016/j.neurobiolaging.2020.11.004.

Hou M, de Chastelaine M, Jayakumar M, Donley BE, Rugg MD. 2020. Recollection-related hippocampal fMRI effects predict longitudinal memory change in healthy older adults. Neuropsychologia. 146:107537. doi:10.1016/j.neuropsychologia.2020.107537.

Johnson JD, McDuff SGR, Rugg MD, Norman KA. 2009. Recollection, Familiarity, and Cortical Reinstatement: A Multivoxel Pattern Analysis. Neuron. 63(5):697–708. doi:10.1016/j.neuron.2009.08.011.

Johnson MK, Kuhl BA, Mitchell KJ, Ankudowich E, Durbin KA. 2015. Age-related differences in the neural basis of the subjective vividness of memories: evidence from multivoxel pattern classification. Cogn Affect Behav Neurosci. 15(3):644–661. doi:10.3758/s13415-015-0352-9.

Karama S, Bastin ME, Murray C, Royle NA, Penke L, Muñoz Maniega S, Gow AJ, Corley J, Valdés Hernández M, Lewis JD, et al. 2014. Childhood cognitive ability accounts for associations between cognitive ability and brain cortical thickness in old age. Mol Psychiatry. 19(5):555–559. doi:10.1038/mp.2013.64.

Kidwai A, Hou M, de Chastelaine M, Rugg MD. 2025. Recollection-related fMRI effects in entorhinal cortex predict longitudinal memory change in healthy older adults. Neurobiology of Aging. 147:150–162. doi:10.1016/j.neurobiolaging.2024.12.011.

Kizilirmak JM, Soch J, Richter A, Schott BH. 2024. Age-related differences in fMRI subsequent memory effects are directly linked to local grey matter volume differences. Neurobiology of Aging. 134:160–164. doi:10.1016/j.neurobiolaging.2023.12.002.

Koen JD, Hauck N, Rugg MD. 2019. The Relationship between Age, Neural Differentiation, and Memory Performance. J Neurosci. 39(1):149–162. doi:10.1523/JNEUROSCI.1498-18.2018.

Koen JD, Rugg MD. 2019. Neural Dedifferentiation in the Aging Brain. Trends in Cognitive Sciences. 23(7):547–559. doi:10.1016/j.tics.2019.04.012.

Kranz MB, Voss MW, Cooke GE, Banducci SE, Burzynska AZ, Kramer AF. 2018. The cortical structure of functional networks associated with age-related cognitive abilities in older adults. PLOS ONE. 13(9):e0204280. doi:10.1371/journal.pone.0204280.

Krogsrud SK, Mowinckel AM, Sederevicius D, Vidal-Piñeiro D, Amlien IK, Wang Y, Sørensen Ø, Walhovd KB, Fjell AM. 2021. Relationships between apparent cortical thickness and working memory across the lifespan - Effects of genetics and socioeconomic status. Developmental Cognitive Neuroscience. 51:100997. doi:10.1016/j.dcn.2021.100997.

Lee S, Habeck C, Razlighi Q, Salthouse T, Stern Y. 2016. Selective association between cortical thickness and reference abilities in normal aging. NeuroImage. 142:293–300. doi:10.1016/j.neuroimage.2016.06.041.

Liu P, Hebrank AC, Rodrigue KM, Kennedy KM, Section J, Park DC, Lu H. 2013. Age-related differences in memory-encoding fMRI responses after accounting for decline in vascular reactivity. NeuroImage. 78:415–425. doi:10.1016/j.neuroimage.2013.04.053.

Masouleh SK, Eickhoff SB, Hoffstaedter F, Genon S, Alzheimer's Disease Neuroimaging Initiative. 2019. Empirical examination of the replicability of associations between brain structure and psychological variables. Nichols TE, Behrens TE, editors. eLife. 8:e43464. doi:10.7554/eLife.43464.

Masouleh SK, Eickhoff SB, Maleki Balajoo S, Nicolaisen-Sobesky E, Thirion B, Genon S. 2022. Empirical facts from search for replicable associations between cortical thickness and psychometric variables in healthy adults. Sci Rep. 12(1):13286. doi:10.1038/s41598-022-17556-7.

McDonough IM, Cervantes SN, Gray SJ, Gallo DA. 2014. Memory's aging echo: Age-related decline in neural reactivation of perceptual details during recollection. NeuroImage. 98:346–358. doi:10.1016/j.neuroimage.2014.05.012.

Mumford JA, Davis T, Poldrack RA. 2014. The impact of study design on pattern estimation for single-trial multivariate pattern analysis. NeuroImage. 103:130–138. doi:10.1016/j.neuroimage.2014.09.026.

Nilsson L-G. 2003. Memory function in normal aging. Acta Neurologica Scandinavica. 107(s179):7–13. doi:10.1034/j.1600-0404.107.s179.5.x.

Nyberg L, Lövdén M, Riklund K, Lindenberger U, Bäckman L. 2012. Memory aging and brain maintenance. Trends in Cognitive Sciences. 16(5):292–305. doi:10.1016/j.tics.2012.04.005.

Park H, Kennedy KM, Rodrigue KM, Hebrank A, Park DC. 2013. An fMRI study of episodic encoding across the lifespan: Changes in subsequent memory effects are evident by middle-age. Neuropsychologia. 51(3):448–456. doi:10.1016/j.neuropsychologia.2012.11.025.

Park H, Uncapher MR, Rugg MD. 2008. Effects of study task on the neural correlates of source encoding. Learn Mem. 15(6):417–425. doi:10.1101/lm.878908.

Park J, Carp J, Kennedy KM, Rodrigue KM, Bischof GN, Huang C-M, Rieck JR, Polk TA, Park DC. 2012. Neural Broadening or Neural Attenuation? Investigating Age-Related Dedifferentiation in the Face Network in a Large Lifespan Sample. J Neurosci. 32(6):2154–2158. doi:10.1523/JNEUROSCI.4494-11.2012.

Raven J, Raven JC, Courth J. 2000. The advanced progressive matrices. In: Manual for Raven's progressive matrices and vocabulary scales, Section 4. San Antonio: Harcourt Assessment.

Reitan R, Wolfson D. 1985. The Halstead-Reitan neuropsychological test battery: therapy and clinical interpretation. Tucson: Neuropsychological.

Rissman J, Gazzaley A, D'Esposito M. 2004. Measuring functional connectivity during distinct stages of a cognitive task. NeuroImage. 23(2):752–763. doi:10.1016/j.neuroimage.2004.06.035.

Rissman J, Wagner AD. 2012. Distributed Representations in Memory: Insights from Functional Brain Imaging. Annual Review of Psychology. 63(1):101–128. doi:10.1146/annurev-psych-120710-100344.

Rugg MD. 2017. Interpreting age-related differences in memory-related neural activity. In: Cognitive neuroscience of aging: Linking cognitive and cerebral aging, 2nd ed. New York, NY, US: Oxford University Press. p. 183–203.

Rugg MD, Johnson JD, Uncapher MR. 2015. Encoding and Retrieval in Episodic Memory. In: The Wiley Handbook on the Cognitive Neuroscience of Memory. John Wiley & Sons, Ltd. p. 84–107.

Rugg MD, Srokova S. 2024. Effects of age on neural reinstatement during memory retrieval. In: Reference Module in Neuroscience and Biobehavioral Psychology. Elsevier.

Salthouse TA, Habeck C, Razlighi Q, Barulli D, Gazes Y, Stern Y. 2015. Breadth and age-dependency of relations between cortical thickness and cognition. Neurobiology of Aging. 36(11):3020–3028. doi:10.1016/j.neurobiolaging.2015.08.011.

Schott BH, Soch J, Kizilirmak JM, Schütze H, Assmann A, Maass A, Ziegler G, Sauvage M, Richter A. 2023. Inhibitory temporo-parietal effective connectivity is associated with explicit memory performance in older adults. iScience. 26(10):107765. doi:10.1016/j.isci.2023.107765.

Smith A. 1982. Symbol digit modalities test (SDMT) manual. Los Angeles: Western Psychological Services.

Snodgrass JG, Corwin J. 1988. Pragmatics of measuring recognition memory: applications to dementia and amnesia. Journal of experimental psychology: General. 117(1):34.

Spreen O, Benton A. 1977. Neurosensory center comprehensive examination for aphasia. Victoria: Neuropsychology Laboratory.

Srokova S, Aktas ANZ, Koen JD, Rugg MD. 2024. Dissociative Effects of Age on Neural Differentiation at the Category and Item Levels. J Neurosci. 44(4):e0959232023. doi:10.1523/JNEUROSCI.0959-23.2023.

Srokova S, Hill PF, Elward RL, Rugg MD. 2021. Effects of age on goal-dependent modulation of episodic

    memory retrieval. Neurobiology of Aging. 102:73–88.

    doi:10.1016/j.neurobiolaging.2021.02.004.

Srokova S, Hill PF, Koen JD, King DR, Rugg MD. 2020. Neural Differentiation is Moderated by Age in

    Scene-Selective, But Not Face-Selective, Cortical Regions. eNeuro. 7(3):ENEURO.0142-20.2020.

    doi:10.1523/ENEURO.0142-20.2020.

Srokova S, Hill PF, Rugg MD. 2022. The Retrieval-Related Anterior Shift Is Moderated by Age and

    Correlates with Memory Performance. J Neurosci. 42(9):1765–1776.

    doi:10.1523/JNEUROSCI.1763-21.2021.

St-Laurent M, Abdi H, Bondad A, Buchsbaum BR. 2014. Memory Reactivation in Healthy Aging: Evidence

    of Stimulus-Specific Dedifferentiation. J Neurosci. 34(12):4175–4186.

    doi:10.1523/JNEUROSCI.3054-13.2014.

St-Laurent M, Buchsbaum BR. 2019. How Multiple Retrievals Affect Neural Reactivation in Young and

    Older Adults. The Journals of Gerontology: Series B. 74(7):1086–1100.

    doi:10.1093/geronb/gbz075.

Sun FW, Stepanovic MR, Andreano J, Barrett LF, Touroutoglou A, Dickerson BC. 2016. Youthful Brains in

    Older Adults: Preserved Neuroanatomy in the Default Mode and Salience Networks Contributes

    to Youthful Memory in Superaging. J Neurosci. 36(37):9659–9668.

    doi:10.1523/JNEUROSCI.1492-16.2016.

Thakral PP, Wang TH, Rugg MD. 2015. Cortical reinstatement and the confidence and accuracy of source

    memory. NeuroImage. 109:118–129. doi:10.1016/j.neuroimage.2015.01.003.

Thakral PP, Wang TH, Rugg MD. 2017. Decoding the content of recollection within the core recollection network and beyond. Cortex. 91:101–113. doi:10.1016/j.cortex.2016.12.011.

Thakral PP, Wang TH, Rugg MD. 2019. Effects of age on across-participant variability of cortical reinstatement effects. NeuroImage. 191:162–175. doi:10.1016/j.neuroimage.2019.02.005.

Trelle AN, Carr VA, Guerin SA, Thieu MK, Jayakumar M, Guo W, Nadiadwala A, Corso NK, Hunt MP, Litovsky CP, et al. 2020. Hippocampal and cortical mechanisms at retrieval explain variability in episodic remembering in older adults. Schlichting ML, Colgin LL, Nyberg L, Koen J, editors. eLife. 9:e55335. doi:10.7554/eLife.55335.

Wang TH, Johnson JD, de Chastelaine M, Donley BE, Rugg MD. 2016. The Effects of Age on the Neural Correlates of Recollection Success, Recollection-Related Cortical Reinstatement, and Post-Retrieval Monitoring. Cerebral Cortex. 26(4):1698–1714. doi:10.1093/cercor/bhu333.

Wechsler D. 1981. Wechsler adult intelligence scale-revised (WAIS-R). Psychological Corporation.

Wechsler D. 2001. Wechsler Test of Adult Reading: WTAR. San Antonio: The Psychological Corporation.

Wechsler D. 2009a. Wechsler memory scale, Ed 4. San Antonio: The Psychological Corporation.

Wechsler D. 2009b. Test of premorbid functioning. San Antonio, TX: The Psychological Corporation.

Yarkoni T, Poldrack RA, Nichols TE, Van Essen DC, Wager TD. 2011. Large-scale automated synthesis of human functional neuroimaging data. Nat Methods. 8(8):665–670. doi:10.1038/nmeth.1635.