Title: Effects of Interleukin-6 dysregulation in a mouse model of Alzheimer's disease:

unraveling the complexity beyond amyloidosis

Authors: Carla Canal [1,2], Kevin Aguilar [1,2,3], Gemma Comes [1,2], Paula Sanchis [1,2,4]

Amalia Molinero [1,2], Mercedes Giralt [1,2], Juan Hidalgo* [1,2]

Authors affiliation:

[1] Department of Cell Biology, Physiology, and Immunology, Animal Physiology Unit,

Faculty of Biosciences, Universitat Autònoma de Barcelona, Bellaterra,

Barcelona, Spain.

[2] Institut de Neurociències, Universitat Autònoma de Barcelona, Bellaterra,

Barcelona, Spain.

[3] Current affiliation: Institute for Research in Biomedicine (IRB Barcelona),

Barcelona, Spain.

[4] Current affiliation: Centre of Inflammation and Metabolism/Centre for Physical

Activity Research (CIM/CFAS), Rigshospitalet, University of Copenhagen,

Copenhagen, Denmark.

Keywords: Alzheimer's disease (AD), Interleukin-6 (IL-6), Tg2576, Amyloid

precursor protein (APP), amyloid-β peptides (Aβ), neuroinflammation, glial

reactivity.

Correspondence: Juan Hidalgo, Department of Cellular Biology, Physiology and

Immunology, Animal Physiology Unit, Faculty of Biosciences, Universitat

Autònoma de Barcelona, Barcelona 08193, Spain. Email: Juan.Hidalgo@uab.es

1. Abstract

Interleukin-6 dysregulation has been implicated in the pathological progression of Alzheimer's disease (AD), a leading cause of dementia in the ageing population. Here we show that chronic neuroinflammation elicited by transgenic expression of IL-6 and systemic IL-6 deficiency influences the phenotype of the Tg2576 mouse model of AD, modulating mortality, body weight, behavioral and cognitive traits, amyloidosis, and neuroinflammation. While the conventional understanding of AD pathophysiology emphasizes the central role of Aβ peptides and amyloid plaques in the development and progression the disease, the absence of amyloidosis in the brain in a specific subset of Tg2576 mice challenges this notion. This suggests that several Tg2576-related traits may manifest independently of Aβ, pointing to a potential contribution of alternative APP-related factors or pathways to the phenotypic alterations observed in Tg2576 mice. Together, our findings underscore the complexity of AD pathogenesis and emphasize the multifaceted nature of IL-6 in both physiological and neurodegenerative processes, particularly in the context of AD.

2. Introduction

Alzheimer's disease (AD) stands as the primary cause of dementia among the elderly, with no significant breakthroughs in the discovery of effective disease-modifying treatments despite extensive research efforts over recent decades (1). The formation of extracellular amyloid deposits of amyloid-β peptides (Aβ) and intracellular neurofibrillary tangles of hyperphosphorylated tau protein, are the two main histopathological hallmarks of AD, predominantly affecting brain regions as the hippocampus and the cortex. Extensive neuronal dysfunction, synaptic loss, vascular damage, and neuroinflammation have also been related to the pathophysiology of the disease (2). Indeed, the inflammatory response to Aβ has been directly associated with the development and pathogenesis of AD. This response involves glial reactivity and the release of inflammatory mediators, likely serving as a defense mechanism to contain Aβ peptides accumulation and regulate the local microenvironment. However, glial reactivity also activates the production and processing of APP leading to sustained elevated levels of Aβ, ultimately contributing to disease progression and severity (3–7).

Interleukin-6 (IL-6) is a pleiotropic cytokine with pro- and anti-inflammatory effects. Ever since its discovery, a significant role for IL-6 in the CNS emerged, and its dysregulation has been associated to the pathological process of AD (8). In fact, IL-6 levels are increased in the serum and brain tissue of AD patients, especially in the area surrounding senile plaques and in the cerebrospinal fluid (7,9). In addition, IL-6 along with other pro-inflammatory cytokines such as IL-1β and TNF-α, are also able to promote amyloid precursor protein (APP) production and enhance BACE1 levels and activity, a critical enzyme in APP cleavage to Aβ, resulting in a self-reinforcing positive feedback loop in AD pathogenesis (7,10). The upregulation of IL-6 is also observed in AD translational mouse models, in which IL-6 is present both around and within β-amyloid plaques (11).

A frequently used and well-characterized transgenic model is the Tg2576 mouse model, which overexpresses human APP (hAPP) harboring the Swedish two-point mutation KM670/671NL (12). This model displays early and aging-associated behavioral and cognitive impairments and pathophysiological features that include Aβ oligomer formation and aggregation into extracellular amyloid deposits, neuritic dystrophy, and neuroinflammation (12–15). However, the mechanisms underlying the effects of hAPP overexpression in Tg2576 are still to be fully elucidated, as recent

evidence highlights that molecular perturbance and cognitive impairments may occur

independently of Aβ formation (16). In mice, the formation of soluble Aβ and other

fragments require transgenic expression of hAPP. Thus, the specific patterns of

transgene expression may contribute to specific pathological and cognitive

disturbances observed in Tg2576 mice before amyloid plaque pathology becomes

apparent, rather resulting from altered APP signaling than of Aβ toxicity (17,18).

Given the potential role of IL-6 in modulating behavior previously demonstrated in

both IL-6 overexpressing mice and IL-6 deficient mice (19–22), we hypothesized that

IL-6 could play a critical role modulating the phenotype of Tg2576 mice. Furthermore,

IL-6 could influence the molecular pathways of Aβ formation and aggregation.

Indeed, evidence suggests a protective role for IL-6 by promoting gliosis, which

reduces Aβ deposition and promotes plaque clearance through phagocytosis (5,23–

25). Here we provide new insights into the impact of IL-6 on the phenotype of Tg2576

mice in two main paradigms: chronic neuroinflammation elicited by transgenic

expression of IL-6 in astrocytes and systemic IL-6 deficiency. Our findings elucidate a

complex interaction between IL-6 and the phenotype of Tg2576 mice, highlighting a

potential role for IL-6 in AD development and progression. This study also challenges

the conventional understanding of the role of Aβ peptides in Tg2576 mice and

underscores the importance of reevaluating existing paradigms in animal models.

3. Materials and methods

3.1. Animals

Tg2576 mice (Taconic Europe) (12), GFAP-IL6 mice (26), and IL-6-DIO-KO mice

(27) were used in this study. Hemizygous Tg2576 male mice were bred with

hemizygous GFAP-IL6 mice to generate WT ($hAPP^{+/+}$, Gfap-Il6$^{+/+}$), GFAP-IL6

($hAPP^{+/+}$, Gfap-Il6$^{Tg/+}$), Tg2576 ($hAPP^{Tg/+}$, Gfap-Il6$^{+/+}$), and Tg2576/GFAP-IL6

($hAPP^{Tg/+}$, Gfap-Il6$^{Tg/+}$). Also, hemizygous Tg2576 mice were bred with homozygous

IL-6-DIO-KO mice following a two-step breeding strategy to generate control

($hAPP^{+/+}$, Il6$^{+/+}$), referred to as WT in the text, IL6KO ($hAPP^{+/+}$, Il6$^{\Delta/+}$), Tg2576

($hAPP^{Tg/+}$, Il6$^{+/+}$), and Tg2576/IL6KO ($hAPP^{Tg/+}$, Il6$^{\Delta/+}$). Genotype was determined by

PCR analysis as previously described (26–28). Mice were group-housed with

littermates in a temperature-controlled environment (22 $\pm$ 2 ºC) under specific

pathogen-free conditions, with a 12-hour dark-light cycle and were maintained with

food and water available *ad libitum*. Experiments were conducted during the light

phase and mice were monitored regularly from weaning to euthanasia at 18 months

of age. Both male and female mice were used unless otherwise noted. All

experimental protocols were approved by the Ethics Committee on Animal and

Human Experimentation (CEEAH) of the Autonomous University of Barcelona (3971

and 9861) and were conducted in accordance with the Spanish legislation and the

EU directive (2010/63/UE) on 'Protection of Animals Used for Experimental and

Other Scientific Purposes'.

3.2.   Behavioral tests

Behavioral evaluation was performed at 5-6 (young) and at 16–17 (old) months of

age, akin to methodology previously described (29,30). Mice were handled for 5 days

before testing for habituation, and a 30-minute acclimation in the home cage within

the testing room preceded each test. Tests were conducted in the following order on

alternate days: open field (OF), hole board (HB), elevated plus maze (EPM), pole

test, Y-maze (YM) and/or Morris water maze (MWM), and nesting. Behavioral data

collection and analyses were performed under blind conditions.

3.2.1. OF, HB, and EPM tests

Locomotor and exploratory activity were assessed in an OF. Mice were placed in a

rectangular apparatus (36,5 x 56 x 31 cm) and allowed to freely explore it for 5 min.

The measured parameters were total horizontal activity (ambulatory distance) and

counts for vertical activity (rearings).

Exploratory behavior was assessed in a HB. Mice were placed in a square

apparatus (40 x 40 x 21,5 cm) with four equally spaced bottom holes (3 cm diameter)

and allowed to freely explore it for 5 min. The measured parameters were the time

and number of head-dipping events.

Anxiety-like behavior was assessed in an EPM. Mice were placed in a cross-

shaped maze consisting of two open arms (20 x 5 cm) and two closed arms (20 x 5 x

15 cm) elevated 47 cm from the floor and were allowed to freely explore the

apparatus for 5 min. The measured parameters were percentage of entries and time

spent in the open arms (OA) of the apparatus (excluding the center).

3.2.2. Pole test

Motor coordination and balance of mice were evaluated using the pole test. Mice

were placed head-up on top of a vertical pole (30 cm height, 1 cm diameter) with a

rough surface for 120 sec at maximum. Mice underwent three trials of habituation

one week before testing. On test day, three consecutive trials were conducted with a

10-min separation, and the time to descend (maximum duration was scored if the

mouse fell down the pole) and the number of failed trials were measured.

3.2.3. YM and MWM tests

A Y-Maze was used to assess working and spatial reference memory. Mice were

placed at the starting arm of a Y-shaped maze consisting of three closed arms (33 x

5 x 15 cm) with external cues placed around the apparatus. To assess working

memory, mice were allowed to freely explore the apparatus in a single 8-min trial.

Arm entries and alternations (three consecutive entries to three different arms) were

calculated as alternations / (total arm entries – 2) * 100. To assess spatial reference

memory, mice were allowed to freely explore the apparatus for 10 min with one arm

closed off (training session), and after 1 and 2 h the blocker was removed, and mice

were allowed to explore the apparatus for 5 min (trial sessions). The measured

parameters were entries and time spent in the novel arm (NA) of the apparatus.

The MWM was used to assess learning and spatial memory. The apparatus

consisted of a round pool (120 cm diameter x 60 cm deep) virtually divided into four

equal quadrants with a removable platform (8 cm diameter). The pool was filled with

opaque water at 22 ± 1 ºC and surrounded by external cues for spatial references.

Animals were tested for 11 days in differentiated phases as in the following protocol

(30). Briefly, animals were challenged to escape from the pool by finding the hidden

platform (submerged 1 cm) in a target quadrant (TQ). Four-trial sessions (1 min

each) during four consecutive days were performed. The platform was removed and

short- (1 hour after the last training session) and long-term memory (24 hours after)

were tested in the probe trials (30 sec each). The evaluation included both spatial

acquisition and spatial reversal (re-learning of a new platform location) phases. The

measured parameters include escape latency (time to find the hidden platform, mean

of four trials) and area under the curve (AUC) calculated for each the training curve,

and latency to reach the platform area (PA) in the probe trials.

3.2.4. Nesting behavior

Natural behavior of mice was assessed with the nesting behavior test. During the

testing period, nesting mice were individually housed, and 3 g of nesting material

(Nestlet) were provided 1 h before the onset of the dark cycle. Nests were assessed

1 h after the next light cycle started and nesting scores were calculated by the extent

of shredded material and the shape of the nest constructed on a scale of 1-5 (from

undisturbed material to near perfect nest) according to published criteria (31). Only

female mice were included in the study.

3.2.5. Analysis of behavior

Tests were recorded by an overhead video camera and videos were analyzed with

either the Ethovision program version 11 (OF, HB, EPM, and YM) or the ANY-maze

version 5.1 (MWM). Data was obtained through both *in situ* manual measurements

and automatic tracking analysis.

3.3.   Neuropathological analysis

3.3.1. Sample collection

Mice were weighed and killed by decapitation at 18 months of age. Whole

blood was collected, allowed to clot, and centrifuged at 6000 x g for 10 min at 4 ºC to

separate serum samples. Brains were extracted and the left hemisphere was fixed in

4% paraformaldehyde (PFA) for 24 h at 4 ºC for further processing. The right

hemisphere was dissected into cortex (anterior and posterior) and hippocampus and

rapidly frozen in liquid nitrogen and stored at -80 ºC for subsequent analysis.

3.3.2. Western Blot

Samples were homogenized by sonication in a lysis buffer (50 mM Tris-HCl pH

7.6, 150 mM NaCl, 2mM EDTA, 0,1% Igepal CA 630, 3% SDS, and 2% sodium

deoxycholate, supplemented with a protease inhibitor cocktail and

phenylmethylsulfonyl fluoride). Total tissue protein quantification was performed with

the Pierce™ BCA Protein Assay Kit (ThermoFisher) according to the manufacturer's

protocol.

Cortical and hippocampal lysates were prepared in NuPAGE™ LDS 4X (Invitrogen

NP0008) and 2-mercaptoethanol (Sigma-Aldrich 805740) and were heat shock for 5

min at 90 ºC. Proteins were separated by electrophoresis on a NuPage™ 4–12% Bis-

Tris Midi protein Gel (Invitrogen) and transferred to 0,2 μm nitrocellulose membranes.

Membranes were blocked with 5% (w/v) skim milk and incubated with primary

antibodies (mouse anti-β-amyloid$_{1-16}$ 6E10 1:2000, 803003 Biolegend; rabbit anti-

GFAP 1:40000, Z0334 Dako; and rabbit anti-Iba-1 1:3000, 019-19741 Wako)

overnight at 4 ºC, and with secondary antibodies (polyclonal rabbit anti-mouse

Ig/HRP 1:5000, P0260 Dako; and goat anti-rabbit IgG (H+L)-HRP conjugate 1:5000

and 1:10000, 1706515 Bio-Rad). Imunoreactive bands were detected using the

Clarity Western ECL Substrate (Bio-Rad), imaged in the Chemidoc Imaging System

(Bio-Rad), and analyzed using Image Lab™ 6.0 Software (Bio-Rad). Monoclonal anti-

β-Actin (1:5000, Sigma-Aldrich) was used as a loading control following membrane

stripping with 2% SDS, Tris HCl, and Tris Base pH 6.7.

3.3.3. Enzyme-Linked ImmunoSorbent assay (ELISA)

ELISA assay was performed to measure $A\beta_{40}$ and $A\beta_{42}$ (Amyloid beta 40 and 42

Human ELISA Kits, ThermoFisher) in cortical and hippocampal homogenates (as

described in 3.3.2) and in serum samples. The protocol was performed according to

the manufacturer's recommendations. The protein concentration of the lysate was

used to normalize the data.

3.3.4. Immunofluorescence (IF)

Fixed tissues were frozen in OCT, and tissue was sagittally cut into serial 20-μm-

thick sections in a cryostat (Leica CM3050 S). All the procedures in this subsection

were performed in free-floating sections. Sections were blocked with 1% BSA in

PBS-Triton 0,2%. For amyloid plaques detection, a 5-minute pre-treatment with 70%

formic acid was performed. Sections were incubated with primary antibodies (mouse

anti-β-amyloid$_{17-24}$ 4G8, Biolegend; rabbit anti-GFAP, Dako; and rabbit anti-IBA-1,

Wako) at 1:500 in blocking solution, and then with secondary antibodies (goat anti-

mouse IgG (H+L) Alexa Fluor™ 488 1:600; Invitrogen A11001; and donkey anti-

rabbit IgG (H+L) Alexa Fluor™ 594 1:600; Invitrogen A21207) in blocking solution.

Slices were mounted and covered with DAPI Fluoromount-G® mounting media

(Southern Biotech 0100-20). For quantification, three non-consecutive sections per

mouse were selected and imaged using an EVOS™ M5000 Imaging System

(Invitrogen) microscope, and images were analyzed using ImageJ software (32). The

sum of intensity of the specific signal (integrated density) was determined for cortex

and hippocampus and normalized by the area of the region of interest (Area ROI).

The integrated density values for GFAP and Iba1 surrounding amyloid plaques were

also quantified and normalized by Area ROI (plaque-shaped areas).

3.4. Statistics

Data are shown as mean ± SEM and plotted using GraphPad Prism version 10.

Statistical analysis was conducted using SPSS version 23 with separate analyses

performed for male and female samples. Survival was evaluated using the Kaplan-

Meier and Log-rank survival tests, with genotype as a four-level factor. Data with two

variables to compare was analyzed either with a Generalized Linear Model (GzLM)

with both genotypes as main factors and appropriate post hoc pairwise comparisons

when needed (sequential Bonferroni's correction), or with Generalized Estimated

Equations (GEE) for repeated measures, including both genotypes as main factors and time as a within-subject factor. Statistical significance was set as $p \leq 0.05$. Sample size is stated in the text or figure as needed.

4. Results

4.1. IL-6 impacts on the mortality and body weight loss observed in AD-like Tg2576 mice in a sex-dependent manner

Survival was significant decreased in both male and female Tg2576 mice **(Fig. 1A, B)**. IL-6 overexpression aggravated Tg2576-associated mortality in female mice, and the same trend was observed in male mice. Conversely, a pro-survival trend was observed in male GFAP-IL6 mice, while minimal mortality was observed in female mice **(Fig. 1A)**. IL-6 deficiency partially rescued the mortality observed in female Tg2576 mice, but not in males. Interestingly, both male and female IL6KO mice also experienced reduced survival **(Fig. 1B)**.

Tg2576 animals also exhibited lower body weight throughout lifetime compared to littermate controls **(Fig. 1C, D)**. Transgenic expression of IL-6 resulted in late-onset decreased body weight in females, especially in GFAP-IL6 mice, but not in males **(Fig. 1C)**. In contrast, IL6KO males developed mature-onset obesity, as evidenced by increased body weight compared to WT littermates, but not in females **(Fig. 1D)**.

4.2. IL-6 modulates AD-like behavior parameters in Tg2576 mice from early stages

Young Tg2576 (5-6 months old) mice exhibited hyperactivity in the OF test, characterized by increased ambulation and rearing behaviors (horizontal and vertical activity, respectively), enhanced exploratory behavior in the HB, as revealed by an increased time and frequency of head-dipping events, and decreased anxiety in the EPM, shown by an increased percentage of entries and time spent in the open arms of the maze. Notably, these changes were predominantly observed in females **(Fig. 2A-F)**. In male GFAP-IL6 mice, particularly Tg2576/GFAP-IL6 mice, increased vertical and horizontal activity was noted compared to littermate controls. In contrast, the opposite trend was observed in female mice **(Fig. 2A)**. Similarly, IL-6 overexpression increased exploratory behavior in male Tg2576 mice **(Fig. 2B)**. Additionally, both male and female IL-6-overexpressing mice exhibited reduced anxiety-like behavior, particularly in Tg2576/GFAP-IL6 mice **(Fig. 2C)**. In contrast, IL-

6 deficiency reversed the hyperactive phenotype in Tg2576/IL6KO mice but did not

affect their exploratory or anxiety behaviors. Notably, no clear-cut effects were

observed in IL6KO mice when compared to WT mice in these behavioral tests **(Fig.**

2D-F).

In aged Tg2576 mice (16-17 months old), behavioral impairments similar to those

observed in young mice persisted **(Fig. 3A-F)**. In contrast to the findings in young

mice, IL-6 overexpression in aged mice reversed hyperactivity in both sexes **(Fig.**

3A) and partially restored exploratory behavior in Tg2576/GFAP-IL6 females while

increasing it in males **(Fig. 3B)**. Additionally, IL-6 overexpression increased

frequency and duration in the OA of the EPM particularly in Tg2576/GFAP-IL6 female

mice, and male mice followed a similar trend **(Fig. 3C)**. Similarly, IL-6 deficiency

reversed hyperactivity in Tg2576 females, reducing both vertical and horizontal

activity **(Fig. 3D)**, but also the increased exploratory behavior in both males in

females **(Fig. 3E)**. Moreover, decreased anxiety in Tg2576 was modulated in a sex-

dependent manner by IL-6 deficiency, as it potentiated it in Tg2576/IL6KO females,

while it reversed it in male mice **(Fig. 3F)**. As in young mice, there were no significant

effects in IL6KO mice compared to WT mice in these behavioral tests, except for

anxiety in male mice, which was increased in IL6KO mice **(Fig. 3D-F)**.

A decline in motor performance was observed in aged Tg2576 mice, especially in

females, as evidenced by a significant increase in the time to descend from the pole

test, although not in the number of failed trials. GFAP-IL6 mice also displayed

impaired motor performance, characterized by an increased time to descend from the

pole, along with a higher number of failed trials. However, the effects of motor

impairment on Tg2576 mice were not compounded by IL-6 overexpression **(Fig. 3G)**.

Impaired nest-building behavior was observed in female Tg2576 mice, as

indicated by a decrease in average nesting scores that persisted through aging.

While IL-6 deficiency did not significantly influence this phenotype in Tg2576/IL6KO

mice, it was enhanced in IL6KO mice compared to WT littermates, especially at aged

stages **(Fig. 3H)**.

4.3.   *IL-6 influences learning and memory impairment of Tg2576 mice*

Working memory was impaired in aged Tg2576 mice, displayed by a decrease in

the percentage of spontaneous alternations in the YM **(Fig. 4A, B)**. IL-6

overexpression improved working memory in both GFAP-IL6 and Tg2576/GFAP-IL6

female mice; in male mice, the same was observed in Tg2576/GFAP-IL6, but the contrary in GFAP-IL6 animals. While a higher number of total arm entries was also observed in Tg2576 mice, no effects were related to IL-6 overexpression **(Fig. 4A)**. In contrast, IL-6 deficiency did not ameliorate working memory of Tg2576/IL6KO mice **(Fig. 4B)**.

Spatial learning and memory were also impaired in aged Tg2576 mice, as evidenced by a decreased percentage of time and entries to the novel arms of the YM test in both sexes **(Fig. 4C)**, as well as an increase in the escape latency curves and in the latency to the platform area of the MWM test, especially in males but not in females **(Fig. 4D, E)**. IL-6 overexpression had opposite effects in females, as it improved memory in GFAP-IL6 mice while worsening it in Tg2576/GFAP-IL6 mice, particularly during the test performed 2 hours after training. Conversely, GFAP-IL6 male mice displayed worsened spatial memory **(Fig. 4C)**. IL-6 deficiency improved learning in females, as seen by a reduction in escape latency and AUC, as well as decreased latency to PA in both spatial acquisition and reversal phases. Similarly, IL-6 deficiency tends to reverse the impaired memory in Tg2576 male mice in the spatial reversal phase **(Fig. 4D, E)**.

4.4.   IL-6 regulates APP-related molecular pathways of Tg2576 mice

The brains of Tg2576 mice are characterized by significant Aβ pathology and progressive development of diffuse and senile plaque resulting from hAPP overexpression and processing. As expected, amyloid plaques were present in Tg2576 animals, as revealed by 4G8 immunostaining performed in the IL-6 overexpression study. Quantitative analysis indicated that amyloid plaques tend to be increased in Tg2576/GFAP-IL6 mice compared to littermate Tg2576 mice **(Fig. 5A)**. Western blot analysis revealed elevated levels of hAPP holoprotein and total Aβ peptides in the cortex and hippocampus of Tg2576 mice, whereas, as expected, they were absent in their wild type littermates. IL-6 overexpression decreased total Aβ peptides levels in the cortex and hippocampus of Tg2576 mice. Likewise, a similar trend was observed for APP levels **(Fig. 5B)**. ELISA analysis of specific Aβ species revealed that IL-6 overexpression increased $A\beta_{42}$ levels, the most pathogenic form, while decreasing the levels of $A\beta_{40}$ in both cortex and hippocampus. Therefore, a dramatic shift in the $A\beta_{42}$ / $A\beta_{40}$ ratio was observed in Tg2576/GFAP-IL6 mice compared to Tg2576 mice **(Fig. 5C)**.

In the IL-6 deficiency study, elevated levels of hAPP holoprotein were observed again by Western Blot in the cortex and hippocampus of Tg2576 mice as compared to wild type littermates. In contrast, we did not detect any Aβ proteolytic fragments in the Tg2576 mice **(Fig. 5D)**. Similar results were found when assessing cortical and hippocampal Aβ$_{40}$ and Aβ$_{42}$ levels by ELISA **(Fig. 5E)**. Control Tg2576 run in parallel showed similar APP levels, which were normally processed rendering elevated Aβ levels **(**blue bar in **Fig. 5D and E)**. In contrast to the results in the brain, Aβ$_{40}$ and Aβ$_{42}$ were detected in Tg2576 serum samples, and these levels were significantly decreased by IL-6 deficiency. Interestingly, Tg2576 mice, heterozygous for IL-6, showed lower levels of Aβ peptides than positive controls, homozygous for IL-6 **(Fig. 5F)**.

4.5.   *IL-6 modulates astrogliosis and microgliosis in Tg2576 mice*

Elevated levels of GFAP and Iba-1 were detected by Western Blot in cortical and hippocampal homogenates of Tg2576 mice compared to littermate controls. Increased levels of both markers were also observed in the cortex and hippocampus of IL-6-overexpressing mice, especially in the hippocampus of IL-6-deficient mice. There was a general sex- and area-dependent trend to an additive effect in the double transgenic mice **(Fig. 6A, B).**

Quantification of GFAP immunostaining also showed increased astrogliosis in the cortex and hippocampus of both Tg2576 and GFAP-IL6 mice compared to WT littermates. In Tg2576/GFAP-IL6 mice, GFAP immunostaining was similar to that of Tg2576 in both brain areas **(Fig. 7A, B)**. Quantification of Iba-1 immunostaining area also revealed increased microgliosis in both cortex and hippocampus of female Tg2576 mice, while no significant effects were observed in males. IL-6 overexpression dramatically increased Iba-1 levels in both brain areas and in both sexes, but interestingly, and in contrast to GFAP, Tg2576/GFAPIL6 mice showed similar Iba-1 immunostaining levels as GFAP-IL6 mice **(Fig. 7D, E)**. We next further analyzed GFAP and Iba-1 immunostaining surrounding amyloid plaques, which revealed distinct patterns in Tg2576 and Tg2576/GFAP-IL6 mice. In the former, astrocytosis was predominantly localized around amyloid plaques, whereas in the latter this was lower in both brain areas **(Fig. 7C)**. Similar results were observed for microgliosis, as Iba-1 immunostaining associated with plaques in Tg2576 mice was higher as compared to Tg2576/GFAP-IL6 mice **(Fig. 7F)**.

5. Discussion

Neuroinflammation plays a key role in the pathogenesis of AD disease, and emerging data suggests that IL-6 is closely related to the development and progression of the disease (9,11,33). Here, we elucidate that IL-6 has a significant impact on various physiological, behavioral, and molecular aspects of the Tg2576 mouse model. Moreover, our study suggests that these effects may be independent of APP processing and could rather be explained by APP overexpression than Aβ formation and deposition.

Premature mortality was observed in both male and female Tg2576 mice compared to their respective controls, as expected. Correlation studies indicate that the underlying mechanisms for Tg2576 mortality may be partly attributed to Aβ toxicity. Hence, preventing Aβ production or enhancing its clearance from the brain can decrease mortality in mice (34). Nevertheless, the impact of APP on premature mortality may extend beyond the well-established amyloid-related pathology associated with AD, including potential factors such as seizures, cerebrovascular dysfunctions, oxidative stress, genetic background, environmental factors, and sex-dependent effects (28,35–38). IL-6 may be involved in several of these factors, as evidenced by its role in maintaining BBB integrity and modulating neuronal responses (39). In our study, IL-6 overexpression in astrocytes increased Tg2576-induced mortality, while IL-6 deficiency rescued it, thus underscoring the crucial role of this cytokine in the pathophysiology of the disease. This aligns with results from studies inhibiting IL-6 trans-signaling, a crucial signaling pathway implicated in the neuroinflammatory response of IL-6 to damage, which partially rescued Tg2576 mortality (30).

Besides mortality, weight loss is also a prominent clinical feature in AD, evident in both human patients and transgenic mouse models (40). As previously reported (30,41), a significant decrease in body weight was observed in Tg2576 mice. IL-6 has long been recognized for its significant role in central regulation of body weight, energy balance, and body fat at the central level (42,43). In our study, IL-6-deficient male mice also displayed late-onset obesity, consistent with these reports. Accordingly, body weight was decreased in IL-6-overexpressing mice, further supporting the role of IL-6 in body weight regulation. Interestingly, while IL-6 affected body weight in wild-type animals, its effects were limited in Tg2576 mice, suggesting

that the specific effects of IL-6 on body weight regulation may be influenced and

altered within the Tg2576 setting.

The behavioral phenotype of Tg2576 mice has been broadly studied at different

stages of the disease, and it has been generally established that these mice display

age-related behavioral alterations at an early onset, including hyperactivity, increased

exploration, and decreased anxiety-like behavior, accompanied by a decline in

cognitive function (12,14,15,44–47). Our study successfully replicated these findings

in both young and aged mice, uncovering age-associated effects and sex-specific

variations. IL-6 signaling appears to modulate a variety of behaviors. Thus, previous

studies on IL-6-deficient mice have reported hypoactivity and decreased exploration

(19,21,29), despite some conflicting results have been reported (48). In our study, the

effects of IL-6 were mostly observed in Tg2576 rather than in wild-type mice,

regulating general activity and exploration. In young Tg2576 mice, IL-6

overexpression tended to potentiate hyperactivity and exploratory behavior, while IL-

6 deficiency tended to blunt activity without clear effects on exploration. Conversely,

in aged mice, both IL-6 overexpression and IL-6 deficiency tended to inhibit activity

and exploratory behavior in Tg2576 mice, and ultimately, in wild-type mice as well.

The anxiolytic-like effect of IL-6 has previously been described (49). IL-6

overexpression further decreased anxiety in both young and aged Tg2576 mice,

while IL-6 deficiency also decreased anxiety in aged Tg2576 female mice but

increased it in both Tg2576 and wild-type males. Accordingly, increased anxiety has

been previously reported in IL-6-deficient mice (19,20,29). These findings underscore

the potential role of IL-6 in modulating behavior and suggest that sex-specific

differences may contribute to IL-6-mediated effects.

In our study, Tg2576 mice exhibited impaired learning and memory compared to

age-matched controls. Remarkably, we identified sex-dependent differences in

cognitive performance, potentially indicating task sensitivity variations, as previously

reported by others (14). The cognitive impairment of Tg2576 mice has been found to

correlate with deficits in long-term potentiation (LTP) in the hippocampus (44,50).

Hippocampal-dependent learning and memory are significantly influenced by IL-6,

and studies have reported a negative association between IL-6 levels and cognitive

function (51,52). Interestingly, blocking endogenous IL-6 in a "healthy" brain has

proven to remarkably prolong LTP and enhance long-term memory in hippocampal-

dependent tasks (25,53). In our study, while IL-6 overexpression improved working

memory in Tg2576 mice, it had a cumulative negative effect on spatial reference

memory. Consistently, IL-6 deficiency enhanced spatial learning and memory. The

age-dependent effect of IL-6 on cognitive impairment had previously been reported

both in IL-6-deficient mice (54,55) and in IL-6-overexpressing mice (56), possibly

affected by the age-dependent neurologic pathology, reduced LTP, and reductions in

cerebellar volume (22,57). Collectively, our results suggest that IL-6 plays a pivotal

role in learning and memory processes, with effects influenced by factors such as

sex, age, cognitive task, and brain regions assessed. Disruptions in IL-6 signaling

may impact brain development and, consequently, brain functions (8,53,54).

Motor impairments are also a common late-onset symptom in AD patients, which

may be related to cognitive decline (58). Motor function abnormalities are observed in

studies performed in APP mouse models of AD, which have been attributed to a

cortical dysfunction in the processing of information induced by Aβ (14,45,59,60). In

line with this, our study revealed decreased motor coordination and balance in

Tg2576 mice. Moreover, our findings align with reports of progressive neurological

disease in GFAP-IL6 mice, characterized by ataxia, tremor, motor impairment,

seizures, and premature mortality (22,26). However, the effect of IL-6 on motor

performance in Tg2576 mice was moderate. Previous studies in various mouse

models of AD have also reported poor nesting performance, which may be influenced

by a combination of sensorimotor deficits, episodic memory issues, and hyperactivity

(46,61,62). Consistently, nesting behavior was significantly decreased in Tg2576

mice in our study. Interestingly, IL-6 deficiency improved the nesting scores in aged

mice, suggesting an age-dependent role for IL-6 in regulating distress.

Mutant hAPP overexpression in Tg2576 mice leads to the secretion of $A\beta_{42}$ and

$A\beta_{40}$ and to the formation of senile plaques with dense amyloid cores (12). In our

study, we confirmed the presence of full-length hAPP in the cortex and hippocampus

of Tg2576 mice. However, the findings regarding amyloid production presented

conflicting results, and are dependent on the specific Tg2576 founder mice

employed, as discussed below. In the IL-6 overexpression study, elevated levels of

Aβ were observed, as expected. It has been previously reported that the expression

of inflammatory mediators, including IL-6, could enhance APP expression and

processing and influence Aβ deposition, potentially establishing a positive feedback

loop that contributes to neurodegeneration (63,64). However, astrocytic IL-6

overexpression decreased APP and total Aβ peptides levels, with opposite effects on

$A\beta_{40}$ and $A\beta_{42}$ isoforms. Interestingly, increased levels of the most pathogenic $A\beta_{42}$

isoform were detected, thereby increasing the $A\beta_{42}/A\beta_{40}$ ratio and suggesting a

potential association with enhanced plaque deposition. Hence, IL-6 overexpression

tended to increase amyloid deposition, contrary to the findings in young TgCRND8

mice with AAV IL-6 overexpression (23).

Glial reactivity is a common feature of Tg2576 mice, occurring alongside amyloid

plaque deposition (3,4), and of GFAP-IL6 mice, which develop a progressive

neuroinflammatory state in the brain parenchyma due to the chronic production of

low levels of IL-6 in the brain (52). While gliosis was predominantly associated with

amyloid plaques in Tg2576 mice, astrocytic IL-6 overexpression dramatically

decreased the glial reactivity responding to amyloid plaques. This strongly suggests

an alteration in the physiological responses of glial cells to amyloid plaques in GFAP-

IL6 mice. Thus, the moderate increase in amyloid plaque load observed in IL-6-

overexpressing mice could be attributed to a potential overload of glial cell

phagocytic capacity, resulting in ineffective clearance. The microenvironment

surrounding amyloid plaques appears to be crucial in modulating glial cell phagocytic

function and influencing Aβ deposit clearance. Indeed, the heterogeneous nature of

microglial responses in the context of plaque formation has already been reported by

other studies (4,65). Finally, it is noteworthy that GFAP-IL6 mice display glial cell

reactivity at early stages mice (52), preceding plaque formation, which could

potentially affect the pattern observed in our mice.

Unexpectedly, in the IL-6 deficiency study, Aβ peptides and amyloid plaques were

notably absent in the neural tissue of Tg2576 and Tg2576/IL6KO mice, despite

normal levels of the precursor protein APP. It must be emphasized that the Tg2576

founder mice varied between the two studies. This discrepancy raised several

hypotheses, including genetic drift in the APP gene, impaired APP processing,

increased Aβ clearance or degradation, or interference with the genetic background

of IL-6-deficient mice. However, the latter seems unlikely, as both GFAP-IL6 and IL-

6KO mice share a similar background. Previous studies had reported that the APP

transgene, although being expressed at the highest level in the brain, is substantially

expressed in other organs as well. Nevertheless, the production of Aβ peptides

seems to be restricted to the brain (66). However, $A\beta_{40}$ and $A\beta_{42}$ were present in

serum samples and significantly decreased by IL-6 deficiency. Remarkably, our

Tg2576 mice, heterozygous for IL-6, exhibited reduced Aβ levels in contrast to

Tg2576 positive controls, homozygous for IL-6, suggesting a potential regulatory role

of IL-6 in modulating circulating Aβ levels. These results point to a potential

independent central or peripheral source of Aβ under the control of IL-6 that deserves

further attention. Interestingly, glial reactivity was also increased regardless of Aβ

production and accumulation in Tg2576 mice. IL-6 deficiency also induced a glial

response, possibly linked to the anti-inflammatory properties of this cytokine.

Therefore, the involvement of IL-6 in mediating responses to various stimuli

highlights the potential impact of IL-6 dysregulation on altered neuroinflammation in

the brain.

Despite the lack of abundant amyloidosis, similar physiological and behavioral

changes were observed, suggesting that these phenotypical traits of Tg2576 mice

may occur independently of significant amounts of Aβ. Prior research has established

that cognitive decline and impaired behavior in Tg2576 mice precede the deposition

of Aβ, emphasizing that plaque formation is not necessarily a prerequisite for

cognitive impairment (12,30,41,67). Consequently, the focus has shifted towards

alternative mechanisms beyond Aβ and its aggregated forms' toxicity, including

neurodevelopmental effects attributed to compensatory processes triggered by APP

overexpression, specific APP metabolites, and/or adjacent background genes

(67,68). In fact, APP and its family members play different roles during brain

development, maturity, and aging, impacting gene regulation and processes like cell

proliferation and neurogenesis (69). APP involvement in synapse formation and

neural networks from early stages suggests its key role in maintaining robust brain

circuits. Disruptions in APP signaling may lead to genetic-dependent dysregulation of

synapse dynamics and cellular signaling, contributing to cognitive alterations

independently of Aβ toxicity. Thus, Aβ alone may be sufficient, but not necessary, for

the development of AD pathology (18,70). Additionally, intracellular APP fragments

including C99 significantly contribute to the early pathological events and to the

initiation of the neurodegenerative process in AD (16).

Taken together, our data provides a novel perspective on the potential

contribution of APP, rather than Aβ, in the pathological manifestations observed in

Tg2576 mice. Previous unsuccessful attempts to develop therapeutic approaches for

AD targeting Aβ dysregulation already suggest that while Aβ may contribute to the

pathogenesis of AD, it is not sufficient to cause cognitive decline or impairments in

global functioning (16,71). Furthermore, our findings highlight the multifaceted nature

of IL-6 in physiological and neurodegenerative disorders as AD. The implication of IL-

6 in the early stages of the disease suggests that addressing IL-6 dyshomeostasis

could be an effective strategy for managing symptoms during the early stages of the

disease.

6. Conclusions

- The study demonstrates that IL-6 plays a complex role in Tg2576 mice, with

varying effects on survival, body weight, behavior, and cognition.

- Contrary to expectations, many observed physiological and behavioral

changes in Tg2576 mice seem to occur independently of significant APP

processing, Aβ peptide production, or amyloid plaque deposition.

- CNS-specific IL-6 overexpression modulates glial reactivity surrounding

amyloid plaques in Tg2576 mice.

7. Declarations

Ethics approval

All experimental protocols were approved by the Ethics Committee on Animal and

Human Experimentation (CEEAH) of the Autonomous University of Barcelona (3971

and 9861) and were conducted in accordance with the Spanish legislation and the

EU directive (2010/63/UE) on 'Protection of Animals Used for Experimental and

Other Scientific Purposes'.

Consent for publication

Not applicable.

Availability of data and materials

The data that support the findings of this study are available from the corresponding

author upon reasonable request.

Competing interests

The authors claim no competing interests.

Funding

This study was supported by Ministerio de Economía y Competitividad y Fondo Europeo de Desarrollo Regional SAF2014-56546-R; Ministerio de Ciencia, Innovación y Universidades y Fondo Europeo de Desarrollo Regional RTI2018-101105-B-I00; and Ministerio de Ciencia e Innovación y Fondo Europeo de Desarrollo Regional PID2021-126602OB-I00.

Authors' contributions

C.C. and J.H. conceived and designed the study. C.C., K.A., G.C., P.S, M.G., and A.M performed research. C.C. performed the formal analysis of the data. C.C. and J.H. wrote the initial draft of the manuscript. All authors edited and reviewed the manuscript.

Acknowledgements

The GFAP-IL6 mice were obtained from The Scripps Research Institute (La Jolla, USA) where they were developed by I. L. Campbell (28). C.C. and K.A. acknowledge the support of PIF UAB and FPU17/02065 fellowships, respectively. P.S. received the Lundbeck R380-2021-1300 fellowship.

8. References

1. Long JM, Holtzman DM. Alzheimer disease: an update on pathobiology and treatment strategies. Cell. 2019 Oct 3;179(2):312–39.

2. Knopman DS, Amieva H, Petersen RC, Chételat G, Holtzman DM, Hyman BT, et al. Alzheimer disease. Nat Rev Dis Primers. 2021 Dec 1;7(1).

3. Heneka MT, Carson MJ, El Khoury J, Landreth GE, Brosseron F, Feinstein DL, et al. Neuroinflammation in Alzheimer's disease. Lancet Neurol. 2015;14(4):388–405.

4. Frautschy SA, Yang F, Irrizarry M, Hyman B, Saido TC, Hsiao K, et al. Microglial response to amyloid plaques in APPsw transgenic mice. Am J Pathol. 1998;152(1):307–17.

5. Glass CK, Saijo K, Winner B, Marchetto MC, Gage FH. Mechanisms underlying inflammation in neurodegeneration. Cell. 2010;140(6):918–34.

6. Saito T, Saido TC. Neuroinflammation in mouse models of Alzheimer's disease. Clin Exp Neuroimmunol. 2018 Nov 1;9(4):211–8.

7. Calsolaro V, Edison P. Neuroinflammation in Alzheimer's disease: current evidence and future directions. Alzheimers Dement. 2016 Jun 1;12(6):719–32.

8. Erta M, Quintana A, Hidalgo J. Interleukin-6, a major cytokine in the central nervous system. Int J Biol Sci. 2012;8(9):1254–66.

9. Hüll M, Strauss S, Berger M, Volk B, Bauer J. The participation of interleukin-6, a stress-inducible cytokine, in the pathogenesis of Alzheimer's disease. Behav Brain Res. 1996;78(1):37–41.

10. Sastre M, Dewachter I, Landreth GE, Willson TM, Klockgether T, Van Leuven F, et al. Nonsteroidal anti-inflammatory drugs and peroxisome proliferator-activated receptor-γ agonists modulate immunostimulated processing of amyloid precursor protein through regulation of β-secretase. J Neurosci. 2003;23(30):9796–804.

11. Apelt J, Schliebs R. β-Amyloid-induced glial expression of both pro- and anti-inflammatory cytokines in cerebral cortex of aged transgenic Tg2576 mice with Alzheimer plaque pathology. Brain Res. 2001;894(1):21–30.

12. Hsiao K, Chapman P, Nilsen S, Eckman C, Harigaya Y, Younkin S, et al. Correlative memory deficits, Aβ elevation, and amyloid plaques in transgenic mice. Science (1979). 1996;274(5284):99–102.

13. Westerman MA, Cooper-Blacketer D, Mariash A, Kotilinek L, Kawarabayashi T, Younkin LH, et al. The relationship between Aβ and memory in the Tg2576 mouse model of Alzheimer's disease. J Neurosci. 2002;22(5):1858–67.

14. King DL, Arendash GW. Behavioral characterization of the Tg2576 transgenic model of Alzheimer's disease through 19 months. Physiol Behav. 2002;75(5):627–42.

15. Ognibene E, Middei S, Daniele S, Adriani W, Ghirardi O, Caprioli A, et al. Aspects of spatial memory and behavioral disinhibition in Tg2576 transgenic mice as a model of Alzheimer's disease. Behav Brain Res. 2005 Jan 30;156(2):225–32.

16. Checler F, Afram E, Pardossi-Piquard R, Lauritzen I. Is γ-secretase a beneficial inactivating enzyme of the toxic APP C-terminal fragment C99? J Biol Chem. 2021 Jan 1;296.

17. Höfling C, Shehabi E, Kuhn PH, Lichtenthaler SF, Hartlage-Rübsamen M, Roßner S. Cell type-specific human APP transgene expression by hippocampal interneurons in the Tg2576 mouse model of Alzheimer's disease. Front Neurosci. 2019;13.

18. Dourlen P, Kilinc D, Malmanche N, Chapuis J, Lambert JC. The new genetic landscape of Alzheimer's disease: from amyloid cascade to genetically driven synaptic failure hypothesis? Acta Neuropathol. 2019 Aug 1;138(2):221–36.

19. Armario A, Hernandez J, Bluethmann H, Hidalgo J. IL-6 deficiency leads to increased emotionality in mice: evidence in transgenic mice carrying a null mutation for IL-6. J Neuroimmunol. 1998;92(1–2):160–9.

680   20.   Butterweck V, Prinz S, Schwaninger M. The role of interleukin-6 in stress-induced
681   hyperthermia and emotional behaviour in mice. Behav Brain Res. 2003 Sep 15;144(1–2):49–
682   56.

683   21.   Baier PC, May U, Scheller J, Rose-John S, Schiffelholz T. Impaired hippocampus-dependent
684   and -independent learning in IL-6 deficient mice. Behav Brain Res. 2009 Jun 8;200(1):192–6.

685   22.   Gyengesi E, Rangel A, Ullah F, Liang H, Niedermayer G, Asgarov R, et al. Chronic microglial
686   activation in the GFAP-IL6 mouse contributes to age-dependent cerebellar volume loss and
687   impairment in motor function. Front Neurosci. 2019;13:1–14.

688   23.   Chakrabarty P, Jansen-West K, Beccard A, Ceballos-Diaz C, Levites Y, Verbeeck C, et al.
689   Massive gliosis induced by interleukin-6 suppresses Aβ deposition in vivo: evidence against
690   inflammation as a driving force for amyloid deposition. FASEB J. 2010 Feb;24(2):548–59.

691   24.   Rothaug M, Becker-Pauly C, Rose-John S. The role of interleukin-6 signaling in nervous tissue.
692   Biochim Biophys Acta Mol Cell Res. 2016 Jun 1;1863(6):1218–27.

693   25.   Lyra e Silva NM, Gonçalves RA, Pascoal TA, Lima-Filho RAS, Resende E de PF, Vieira ELM,
694   et al. Pro-inflammatory interleukin-6 signaling links cognitive impairments and peripheral
695   metabolic alterations in Alzheimer's disease. Transl Psychiatry. 2021 Jun 1;11(2).

696   26.   Campbell IL, Abraham CR, Masliah E, Kemper P, Inglis JD, Oldstone MBA, et al. Neurologic
697   disease induced in transgenic mice by cerebral overexpression of interleukin 6. Proc Natl Acad
698   Sci USA. 1993;90(21):10061–5.

699   27.   Sanchis P, Fernández-Gayol O, Comes G, Aguilar K, Escrig A, Giralt M, et al. A new mouse
700   model to study restoration of interleukin-6 (IL-6) expression in a Cre-dependent manner:
701   microglial IL-6 regulation of experimental autoimmune encephalomyelitis. J Neuroinflammation.
702   2020 Dec 1;17(1).

703   28.   Hsiao KK, Borchelt DR, Olson K, Johannsdottir R, Kitt C, Yunis W, et al. Age-related CNS
704   disorder and early death in transgenic FVB/N mice overexpressing Alzheimer amyloid
705   precursor proteins. Vol. 15, Neuron. 1995. p. 1203–18.

706   29.   Erta M, Giralt M, Esposito FL, Fernandez-Gayol O, Hidalgo J. Astrocytic IL-6 mediates
707   locomotor activity, exploration, anxiety, learning and social behavior. Horm Behav. 2015 Jul
708   1;73:64–74.

709   30.   Escrig A, Canal C, Sanchis P, Fernández-Gayol O, Montilla A, Comes G, et al. IL-6 trans-
710   signaling in the brain influences the behavioral and physio-pathological phenotype of the
711   Tg2576 and 3xTgAD mouse models of Alzheimer's disease. Brain Behav Immun. 2019 Nov
712   1;82:145–59.

713   31.   Deacon RMJ. Assessing nest building in mice. Nat Protoc. 2006 Aug;1(3):1117–9.

714   32.   Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image analysis.
715   Vol. 9, Nature Methods. 2012. p. 671–5.

716   33.   Su F, Bai F, Zhang Z. Inflammatory cytokines and Alzheimer's disease: a review from the
717   perspective of genetic polymorphisms. Neurosci Bull. 2016 Oct 1;32(5):469–80.

718   34.   Leissring MA, Farris W, Chang AY, Walsh DM, Wu X, Sun X, et al. Enhanced proteolysis of β-
719   amyloid in APP transgenic mice prevents plaque formation, secondary pathology, and
720   premature death. Neuron. 2003;40(6):1087–93.

721   35.   Carlson GA, Borchelt DR, Dake A, Turner S, Danielson V, Coffin JD, et al. Genetic modification
722   of the phenotypes produced by amyloid precursor protein overexpression in transgenic mice.
723   Hum Mol Genet. 1997;6(11):1951–9.

724   36.   Krezowski J, Knudson D, Ebeling C, Pitstick R, Giri RK, Schenk D, et al. Identification of loci
725   determining susceptibility to the lethal effects of amyloid precursor protein transgene
726   overexpression. Hum Mol Genet. 2004 Sep 15;13(18):1989–97.

727   37.   Rae EA, Brown RE. The problem of genotype and sex differences in life expectancy in
728   transgenic AD mice. Neurosci Biobehav Rev. 2015 Oct 1;57:238–51.

38. Westmark CJ, Westmark PR, Beard AM, Hildebrandt SM, Malter JS. Seizure susceptibility and
mortality in mice that over-express amyloid precursor protein. Int J Clin Exp Pathol.
2008;1(2):157–68.

39. Abbott NJ, Rönnbäck L, Hansson E. Astrocyte-endothelial interactions at the blood-brain
barrier. Nat Rev Neurosci. 2006 Jan;7(1):41–53.

40. Gillette-Guyonnet S, Nourhashémi F, Andrieu S, de Glisezinski I, Ousset PJ, Rivière D, et al.
Weight loss in Alzheimer disease. Am J Clin Nutr. 2000;71(2):637–42.

41. Manso Y, Carrasco J, Comes G, Meloni G, Adlard PA, Bush AI, et al. Characterization of the
role of metallothionein-3 in an animal model of Alzheimer's disease. Cell Mol Life Sci. 2012
Nov;69(21):3683–700.

42. Wallenius V, Wallenius K, Ahrén B, Rudling M, Carlsten H, Dickson SL, et al. Interleukin-6-
deficient mice develop mature-onset obesity. Nat Med. 2002;8(1):75–9.

43. Señarís RM, Trujillo ML, Navia B, Comes G, Ferrer B, Giralt M, et al. Interleukin-6 regulates
the expression of hypothalamic neuropeptides involved in body weight in a gender-dependent
way. J Neuroendocrinol. 2011 Aug;23(8):675–86.

44. Chapman PF, White GL, Jones MW, Cooper-Blacketer D, Marshall VJ, Irizarry M, et al.
Impaired synaptic plasticity and learning in aged amyloid precursor protein transgenic mice.
Nat Neurosci. 1999;2(3):271–6.

45. Lalonde R, Lewis TL, Strazielle C, Kim H, Fukuchi K. Transgenic mice expressing the
βAPP695SWE mutation: effects on exploratory activity, anxiety, and motor coordination. Brain
Res. 2003 Jul 4;977(1):38–45.

46. Deacon RMJ, Cholerton LL, Talbot K, Nair-Roberts RG, Sanderson DJ, Romberg C, et al. Age-
dependent and -independent behavioral deficits in Tg2576 mice. Behav Brain Res. 2008 May
16;189(1):126–38.

47. Reed MN, Liu P, Kotilinek LA, Ashe KH. Effect size of reference memory deficits in the Morris
water maze in Tg2576 mice. Behav Brain Res. 2010 Sep;212(1):115–20.

48. Swiergiel AH, Dunn AJ. Feeding, exploratory, anxiety-and depression-related behaviors are not
altered in interleukin-6-deficient male mice. Behav Brain Res. 2006;171(1):94–108.

49. Wei H, Chadman KK, McCloskey DP, Sheikh AM, Malik M, Brown WT, et al. Brain IL-6
elevation causes neuronal circuitry imbalances and mediates autism-like behaviors. Biochim
Biophys Acta. 2012 Jun;1822(6):831–42.

50. Huh S, Baek SJ, Lee KH, Whitcomb DJ, Jo J, Choi SM, et al. The reemergence of long-term
potentiation in aged Alzheimer's disease mouse model. Sci Rep. 2016 Jul 5;6.

51. Boulanger LM. Immune proteins in brain development and synaptic plasticity. Neuron. 2009
Oct 15;64(1):93–109.

52. Campbell IL, Hofer MJ, Pagenstecher A. Transgenic models for cytokine-induced neurological
disease. Biochim Biophys Acta. 2010 Oct;1802(10):903–17.

53. Balschun D, Wetzel W, Del Rey A, Pitossi F, Schneider H, Zuschratter W, et al. Interleukin-6: a
cytokine to forget. FASEB J. 2004 Nov;18(14):1788–90.

54. Braida D, Sacerdote P, Panerai AE, Bianchi M, Aloisi AM, Iosuè S, et al. Cognitive function in
young and adult IL (interleukin)-6 deficient mice. Behav Brain Res. 2004 Aug 31;153(2):423–9.

55. Bialuk I, Winnicka MM. Facilitatory effect of IL-6 deficiency on long-term spatial memory in
young adult mice. Behav Genet. 2018 May 1;48(3):236–46.

56. Chesworth R, Gamage R, Ullah F, Sonego S, Millington C, Fernandez A, et al. Spatial memory
and microglia activation in a mouse model of chronic neuroinflammation and the anti-
inflammatory effects of apigenin. Front Neurosci. 2021 Jul 30;15.

57. Steffensen SC, Campbell IL, Henriksen SJ. Site-specific hippocampal pathophysiology due to
cerebral overexpression of interleukin-6 in transgenic mice. Brain Res. 1994;652:149–53.

58. Buchman AS, Bennett DA. Loss of motor function in preclinical Alzheimer's disease. Expert
Rev Neurother. 2011 May;11(5):665–76.

59. Lalonde R, Fukuchi KI, Strazielle C. Neurologic and motor dysfunctions in APP transgenic
mice. Rev Neurosci. 2012 Aug;23(4):363–79.
60. Nyul-Toth A, DelFavero J, Mukli P, Tarantini A, Ungvari A, Yabluchanskiy A, et al. Early
manifestation of gait alterations in the Tg2576 mouse model of Alzheimer's disease.
Geroscience. 2021 Aug 1;43(4):1947–57.
61. Filali M, Lalonde R. Age-related cognitive decline and nesting behavior in an APPswe/PS1
bigenic model of Alzheimer's disease. Brain Res. 2009 Sep 25;1292:93–9.
62. Wesson DW, Wilson DA. Age and gene overexpression interact to abolish nesting behavior in
Tg2576 amyloid precursor protein (APP) mice. Behav Brain Res. 2011 Jan 1;216(1):408–13.
63. Brugg B, Dubreuil YL, Hubert G, Wollmant EE, Delhaye-Bouchaud N, Mariani J. Inflammatory
processes induce β-amyloid precursor protein changes in mouse brain. Proc Natl Acad Sci
USA. 1995;92(7):3032–5.
64. Griffin WST, Sheng JG, Royston MC, Gentleman SM, McKenzie JE, Graham DI, et al. Glial-
neuronal interactions in Alzheimer's disease: the potential role of a 'cytokine cycle' in disease
progression. Brain Pathol. 1998;8(1):65–72.
65. Gyoneva S, Swanger SA, Zhang J, Weinshenker D, Traynelis SF. Altered motility of plaque-
associated microglia in a model of Alzheimer's disease. Neuroscience. 2016 Aug 25;330:410–
20.
66. Kawarabayashi T, Younkin LH, Saido TC, Shoji M, Ashe KH, Younkin SG. Age-dependent
changes in brain, CSF, and plasma amyloid protein in the Tg2576 transgenic mouse model of
Alzheimer's disease. J Neurosci. 2001;21(2):372–81.
67. King DL, Arendash GW, Crawford F, Sterk T, Menendez J, Mullan MJ. Progressive and
gender-dependent cognitive impairment in the APPSW transgenic mouse model for
Alzheimer's disease. Behav Brain Res. 1999;103:145–62.
68. Johnson ECB, Ho K, Yu GQ, Das M, Sanchez PE, Djukic B, et al. Behavioral and neural
network abnormalities in human APP transgenic mice resemble those of App knock-in mice
and are modulated by familial Alzheimer's disease mutations but not by inhibition of BACE1.
Mol Neurodegener. 2020 Dec 1;15(1).
69. Zhang YW, Thompson R, Zhang H, Xu H. APP processing in Alzheimer's disease. Mol Brain.
2011;4(3).
70. Priller C, Bauer T, Mitteregger G, Krebs B, Kretzschmar HA, Herms J. Synapse formation and
function is modulated by the amyloid precursor protein. J Neurosci. 2006;26(27):7212–21.
71. Sun BL, Li WW, Zhu C, Jin WS, Zeng F, Liu YH, et al. Clinical research on Alzheimer's
disease: progress and perspectives. Neurosci Bull. 2018 Dec 1;34(6):1111–8.
813
814

Figure 1. Effect of IL-6 on life span and body weight of Tg2576 mice. Kaplan-

Meier survival curves for the IL-6 overexpression study (A) and for the IL-6 deficiency

study (B). Results are analyzed by log-rank test; ● p ≤ .05 vs WT, △ p ≤ .05 vs

GFAP-IL6 or IL6KO, ○ p ≤ .05 vs Tg2576. Body weight curves and body weight at

euthanasia (inset) for the IL-6 overexpression study (C) and for the IL-6 deficiency

study (D). Results are mean ± SEM and analyzed by GEE and GzLM statistical tests;

▲ p ≤ .05 Tg2576 effect, * p ≤ .05 IL-6 effect. n = 18 - 30 in females, n = 20 - 38 in

males at 3 weeks; n = 11 - 30 in females, n = 14 - 26 in males at 80 weeks for the IL-

6 overexpression study; n = 51 - 72 in females, n = 51 - 67 in males at 3 weeks; n =

20 - 33 in females, n = 13 - 26 in males at 80 weeks for the IL-6 deficiency study.

Figure 2. Impact of IL-6 on behavior of young Tg2576 mice. Number of rearings and total distance travelled in the OF test (A), number and time of head-dipping events in the HB test (B), and percentages of entries and time in the OA in the EPM test (C) of the IL-6 overexpression study, OA open arms. Similar results are represented for the IL-6 deficiency study in the OF, HB and EPM tests (D-F). Results are mean ± SEM and analyzed by GzLM statistical test; ▲ p ≤ .05 Tg2576 effect, * p ≤ .05 IL-6 effect, ♦ p ≤ .05 interaction between both factors. n = 13 - 16 in females, n = 13 - 16 in males (A-C); n = 16 - 27 in females, n = 12 - 19 in males (D-F).

Figure 3. Impact of IL-6 on behavior of aged Tg2576 mice. Number of rearings

and total distance travelled in the OF test (A), number and time of head-dipping

events in the HB test (B), and percentages of entries and time in the OA in the EPM

test (C) in the IL-6 overexpression study, OA open arms. Similar results are

represented for the IL-6 deficiency study in the OF, HB and EPM tests (D-F). Time to

descend and number of failed trials in the pole test in the IL-6 overexpression study

(G). Nest-building scores for females in the IL-6 overexpression study (H). Results

are mean ± SEM and analyzed by GzLM statistical test; ▲ p ≤ .05 Tg2576 effect, * p

≤ .05 IL-6 effect, ♦ p ≤ .05 interaction between both factors. n = 11 - 14 in females, n

= 10 - 15 in males (A-C); n = 11 - 17 in females, n = 10 - 16 in males (D-F); n = 10 -

13 in females, n = 10 - 11 in males (G); n = 5 - 16 (H).

Figure 4. Effect of IL-6 on working and spatial memory of Tg2576 mice.
Percentage of alternations and number of total arm entries in the YM test in the IL-6
overexpression study (A) and in the IL-6 deficiency study. Entries and time in the NA
of the spatial memory YM test in the IL-6 overexpression study (C), NA novel arm.
Spatial acquisition (D, left) and reversal (E, left) curves, with respective AUC
quantifications (inset). Latency to PA in the probe trials of spatial acquisition (D, right)
and reversal (E, right) phases at 1 h and 24 h of the MWM test in the IL-6 deficiency
study, PA platform area. Results are mean ± SEM and analyzed by GzLM and GEE
statistical tests; ▲ $p \leq .05$ Tg2576 effect, * $p \leq .05$ IL-6 effect, ♦ $p \leq .05$ interaction
between both factors. n = 13 - 16 in females, n = 13 - 14 in males (A); n = 11 - 16 in
females, n = 10 - 16 in males (B); n = 9 - 14 in females, n = 8 - 10 in males (C); n = 9
- 16 in females, n = 10 - 14 in males (D).

Figure 5. Effect of IL-6 on amyloidosis of Tg2576 mice. (A) Representative

immunofluorescence of 4G8 and quantification of 4G8 levels in the Cx and Hpc of the

IL-6 overexpression study (A). Representative immunoblot of 6E10 and quantification

of APP and Aβ levels relative to β-actin in the Cx and Hpc in the IL-6 overexpression

study (B). Aβ40 and Aβ42 levels in the Cx and Hpc, and relative Aβ42 / Aβ40 levels

in the IL-6 overexpression study (C). Representative immunoblot of 6E10 and

quantification of APP levels relative to β-actin in the Cx and Hpc of the IL-6 deficiency

study (D). Aβ40 and Aβ42 levels in Cx and Hpc in the IL-6 deficiency study. (F) Aβ40

and Aβ42 levels in serum samples in the IL-6 deficiency study. Results are mean ±

SEM and analyzed by Student's t-test; * p ≤ .05 IL-6 effect. Cx cortex, Hpc

hippocampus.

Figure 6. Effect of IL-6 on gliosis of Tg2576 mice. (A) Representative immunoblot

of GFAP and Iba1 in Cx, and quantification of GFAP and Iba1 levels in Tg2576 mice

relative to β-actin in Cx and Hpc in the IL-6 overexpression study. (B) Representative

immunoblot of GFAP and Iba-1 in Cx, and quantification of GFAP and Iba1 levels in

Tg2576 mice relative to β-actin in Cx and Hpc in the IL-6 deficiency study. Results

are mean ± SEM and analyzed by GzLM statistical test; ▲ p ≤ .05 Tg2576 effect, * p

≤ .05 IL-6 effect, ♦ p ≤ .05 interaction between both factors. Cx cortex, Hpc

hippocampus.

Figure 7. Effect of IL-6 overexpression on gliosis surrounding amyloid plaques of Tg2576 mice. (A) Representative immunostaining of GFAP and 4G8, higher magnifications of Cx (middle) and Hpc (right). (B) Quantification of GFAP levels in the Cx and Hpc. (C) Analysis of the percentage of GFAP surrounding amyloid plaques in the Cx and Hpc of Tg2576 mice. (D) Representative immunostaining of Iba1 and 4G8, higher magnifications of Cx (middle) and Hpc (right). (E) Quantification of Iba1 levels in the Cx and Hpc. (C) Analysis of the percentage of Iba1 surrounding amyloid plaques in the Cx and Hpc of Tg2576 mice. Results are mean ± SEM and analyzed by GzLM statistical test; ▲ p ≤ .05 Tg2576 effect, * p ≤ .05 IL-6 effect, ♦ p ≤ .05 interaction between both factors. Cx cortex, Hpc hippocampus.