Production of human papillomavirus type 16 virus-like particles in Physcomitrella photobioreactors

Paul Alexander Niederau[1], Maria Caroline Weilguny[1,2], Sarah Chamas[1], Caitlin Elizabeth Turney[1], Juliana Parsons[1], Marta Rodríguez-Franco[3], Sebastian N.W. Hoernstein[1], Eva L. Decker[1], Henrik Toft Simonsen[2,4], Ralf Reski[1,5,6] *

[1] Plant Biotechnology, Faculty of Biology, University of Freiburg, Schänzlestr. 1, 79104 Freiburg, Germany

[2] Department of Biotechnology and Biomedicine, Technical University of Denmark, 2800 Kongens Lyngby, Denmark

[3] Cell Biology, Faculty of Biology, University of Freiburg, Schänzlestr. 1, 79104 Freiburg, Germany

[4] Laboratoire de Biotechnologies Végétales appliquées aux Plantes Aromatiques et Médicinales, CNRS, UMR 5079, Université Jean Monnet, 42023 Saint-Etienne Cedex 2, France

[5] Signalling Research Centre BIOSS and CIBSS, University of Freiburg, Schänzlestr. 18, 79104 Freiburg, Germany

[6] Cluster of Excellence *liv*MatS @ FIT – Freiburg Centre for Interactive Materials and Bioinspired Technologies, University of Freiburg, Georges-Köhler-Allee 105, 79110 Freiburg, Germany

*Corresponding author: ralf.reski@biologie.uni-freiburg.de

ORCID:

P.A.N.	0009-0004-0194-8570
M.C.W.	0009-0009-9440-6872
S.C.	0009-0007-5704-5639
C.E.T.	0009-0001-3113-4648
J.P.	0000-0001-6261-2342
M.R.F.	0000-0003-1183-2075
S.N.W.H.	0000-0002-2095-689X
E.L.D.	0000-0002-9151-1361
H.T.S.	0000-0003-3070-807X
R.R.	0000-0002-5496-6711

Author E-mails:

P.A.N.   alexander.niederau@biologie.uni-freiburg.de

M.C.W.   mariaweilguny@gmail.com

S.C.   sarah-chamas@web.de

C.E.T.   caitlin.turney@email.uni-freiburg.de

J.P.   juliana.parsons@biologie.uni-freiburg.de

M.R.F.   marta.rodriguez@biologie.uni-freiburg.de

S.N.W.H.   sebastian.hoernstein@biologie.uni-freiburg.de

E.L.D.   eva.decker@biologie.uni-freiburg.de

H.T.S.   henrik.toft.simonsen@univ-st-etienne.fr

R.R.   ralf.reski@biologie.uni-freiburg.de

Abstract

Human papillomavirus (HPV) infections are the main causal agent of cervical and other anogenital cancers. Modern HPV vaccines are based on virus-like particles (VLPs), which are self-assembling nanoparticles composed of viral structural proteins. VLPs mimic native virions but lack viral DNA, rendering them non-infectious while retaining high immunogenicity. Despite available vaccines, the majority of cervical cancer deaths are due to HPV infections, particularly in low- and middle-income countries where inequalities in access to vaccination, screening, and treatment persist. While current HPV vaccines are produced in yeast or insect cells, alternative production platforms are explored, focusing on cost-effectiveness and scalability. Here, we used the moss plant Physcomitrella (*Physcomitrium patens*) to produce HPV-16 VLPs *via* expression of the major structural protein L1. We characterized Physcomitrella chloroplast transit peptides, which we used for targeting of L1 to chloroplasts, leading to higher recombinant protein accumulation compared to nuclear or cytoplasmic localization. We confirmed subcellular localization with confocal laser scanning microscopy and found L1 to accumulate within the chloroplast stroma. Production in 5 L-photobioreactors yielded more than 0.3 mg L1 per g fresh weight. We established a purification protocol for moss-produced L1 using a combination of ammonium sulphate precipitation and cation exchange chromatography. Purified samples were subjected to a controlled dis- and reassembly, ultimately yielding fully assembled HPV-16 L1 VLPs. This is the first report of production, purification and assembly of VLPs in a non-vascular plant.

Keywords: bryophyte, bioreactor, molecular farming, nanoparticle, plant biotechnology, vaccine candidate

Key message: First production of virus-like particles as a vaccine candidate in a non-vascular plant.

Word count: 6809

Introduction

Human papillomavirus (HPV) infections are a major source of cancer, causing approximately 690,000 cases annually (zur Hausen et al., 1981; zur Hausen, 2002; de Martel et al., 2020). The most prevalent type is cervical cancer, accounting for about 80% of HPV-attributable cancers. Other HPV-related cancers include anogenital and oropharyngeal cancers (Wu et al., 2024). The burden of HPV infection is not evenly distributed, with the highest prevalence in low- and middle-income countries, as a result of inequities in access to vaccination, screening, and treatment services (World Health Organisation, 2024a).

HPV belongs to the *Papillomaviridae*, comprising nearly 200 different types, which are commonly transmitted sexually (Ljubojevic & Skerlev, 2014). The majority of HPV-attributable cancers are linked to two types in particular, namely HPV-16 and HPV-18 (de Martel et al., 2020; Wu et al., 2024). HPV is a non-enveloped double stranded DNA virus of 8000 bp, encoding six early regulatory proteins (E1, E2, E4 – E7) and two late structural proteins (L1, L2) (Roden & Stern, 2018). L1 proteins multimerize into pentamers, the capsomers, and L1 capsomers and L2 monomers are imported into the nucleus *via* nuclear localization signals (NLS), where HPV virions assemble (Cerqueira & Schiller, 2017). While the role of L2 in virion assembly differs from type to type, L1 can self-assemble into virus-like particles (VLPs) in the absence of L2 (Kirnbauer et al., 1992). The final HPV-16 virion is composed of 360 L1 monomers, organized in 72 pentamers, and 72 L2 monomers and is reinforced by inter-pentameric disulfide bonds (Baker et al., 1991; Buck et al., 2005). Because the viral life cycle depends on differentiated epithelial cells, it is difficult to obtain large quantities of virions. Therefore, today's HPV vaccines are based on L1 VLPs.

VLPs are nanoparticles consisting solely of viral structural proteins (Marsian & Lomonossoff, 2016; Nooraei et al., 2021). Therefore, they cannot replicate and infect cells. These nanoparticles resemble native viruses in size, shape, and surface structure, presenting highly repetitive immunogenic epitopes. Although VLP-based HPV vaccines are currently produced in microbes (World Health Organization, 2024b), plant-based expression systems are gaining attention in this respect (Scotti & Rybicki, 2013; Rybicki, 2020). In *Nicotiana benthamiana*, expression of L1 and L2 in combination with a reporter gene generated pseudovirions (PsVs), which could substitute mammalian cell culture-derived PsVs in neutralisation assays (Lamprecht et al., 2016; Adams et al., 2023). Alternatively, Tobacco Mosaic Virus VLPs produced in *N. benthamiana* were used for encapsidation and targeted delivery of mRNA encoding oncogenic E6, a crucial target in therapeutic HPV vaccine development (Karan et al., 2024).

While first attempts to produce L1-based VLPs in plants had low yields (Biemelt et al., 2003;
Varsani et al., 2003), considerable improvements have been achieved, e.g. by codon optimization and
subcellular targeting strategies in transiently transformed tobacco (Maclean et al., 2007). A human codon-
optimized version (62% GC content) resulted in highest yields as compared to *N. benthamiana* optimized
(49% GC) and the native sequence (37% GC), presumably not due to the GC content but due to inhibitory
5' sequences, which act on transcription and mRNA processing (Maclean et al., 2007; Hitzeroth et al.,
2018). Further, these authors found chloroplast-targeted L1 yield highest at 0.5 mg / g fresh weight (FW)
followed by cytoplasmic L1 (without the C-terminal 22 aa NLS), whereas L1 targeted to the ER was not
detectable. Similarly, Zahin et al. (2016) reported 0.25 mg L1 / g FW *via* chloroplast targeting while leaving
the NLS intact. Although tobacco remains a prominent option, other plants are increasingly explored for
the efficient production of biopharmaceuticals.

The moss Physcomitrella has emerged as an alternative for biopharmaceutical production,
offering several advantages over traditional systems (Reski et al., 2015; Reski, 2018; Decker & Reski, 2020).
Its haploid dominant phase and high rate of homologous recombination enable precise gene targeting
(Strepp et al., 1998; Wiedemann et al., 2018; Rempfer et al., 2022). This unique feature facilitated the
modification of the N-glycosylation pathway, eliminating plant-specific modifications, such as β1,2-
xylosylation, α1,3-fucosylation, and Lewis A epitopes (Koprivova et al., 2004; Parsons et al., 2012; Decker
et al., 2014). Moreover, enzymes necessary for sialic acid synthesis, activation, and linkage to protein N-
glycans were introduced (Bohlender et al., 2020). These modifications have resulted in more
homogeneous and human-like N-glycan structures on recombinant proteins. The potential of
Physcomitrella as a biopharmaceutical production platform is demonstrated by several products in
development. For example, moss-produced α-galactosidase A has successfully completed clinical phase
Ib (Shen et al., 2016; Hennermann et al., 2019). Other moss-produced proteins include human factor H
(FH) and synthetic FH-related multitarget regulators MFHR1 and MFHR13, which are potential treatments
for complement-related disorders (Büttner-Mainik et al., 2011; Michelfelder et al., 2017, 2018; Top et al.,
2019; Ruiz-Molina et al., 2022a). Importantly, Physcomitrella cultivation in photobioreactors is tightly
controlled (Hohe et al., 2002), and thus compliant with Good Manufacturing Practice (GMP) standards,
ensuring high-quality production of recombinant proteins (Reski et al., 2018; Decker & Reski, 2020). Plants
in general (Rosales-Mendoza et al., 2016) and Physcomitrella in particular (Rosales-Mendoza et al., 2014)
are considered promising production hosts for innovative vaccines. The first moss-produced vaccine
candidate, an ENV-derived multi-epitope HIV chimeric protein, was immunogenic in mice (Orellana-

Escobedo et al., 2015). However, until now, there was no record of VLP-production in Physcomitrella, or
any other non-vascular plant.

Here, we analysed the expression of HPV-16 L1 in Physcomitrella. We characterized five
Physcomitrella chloroplast transit peptides and investigated the effect of subcellular L1 localization on
recombinant protein yields. We further established L1 production in photobioreactors and subsequent
purification of the product. Finally, the assembly of moss-produced HPV-16 L1 to VLPs was demonstrated
with transmission electron microscopy.

Materials and methods

Plant material

139 Physcomitrella (new species name: *Physcomitrium patens* (Hedw.) Mitt.; IMSC acc. no. 41269),
140 was cultivated according to Reski & Abel (1985) and Decker et al. (2015).

Optimization of HPV-16 L1 coding sequence

The CDS of h(human codon-optimized)L1 (GenBank acc. no. DQ067889, Maclean et al., 2007) was
used for optimization using physCO (https://www.plant-biotech.uni-freiburg.de/tools/physco/, Top et al.,
2021). Putative alternative splicing motifs AGGT were identified *via* search function. MicroRNA-binding
sites were predicted *via* psRNATarget with the library of 280 published Physcomitrella microRNAs and the
default settings *Schema V2* (2017 release) (http://plantgrn.noble.org/psRNATarget/, Dai & Zhao, 2011;
Dai et al., 2018). Alternative splice sites and microRNA-binding sites were removed and substituted by
synonymous codons based on http://www.kazusa.or.jp/codon/. Sequence editing was done in Benchling
(https://www.benchling.com/).

Expression vectors

Chloroplast transit peptides (CTPs) were selected from a list of proteins with experimentally
confirmed cleavage of a predicted CTP (Hoernstein et al., 2024). CTP-eGFP vectors were constructed based
on a vector containing an eGFP CDS with an N-terminal linker (sequence: GGGGGA) under control of the
PpActin5 promoter (Weise et al., 2006) and nos terminator (Hoernstein et al., 2023). CTP candidates were
amplified from Physcomitrella cDNA and inserted at the N-terminus of the linker using Gibson Assembly

(Gibson et al., 2009). The Citrine-L1 fusion constructs are based on a vector containing the PpActin5

promoter, nos terminator and citrine CDS (Wiedemann et al., 2018). The CDS of pL1, pL1Δ22 and CTP-pL1

were amplified using Phusion^{TM} polymerase (Thermo Fisher Scientific, Waltham, USA), cloned into the

pJet1.2 backbone (Thermo Fisher Scientific) and ligated into the expression vector using T4 Ligase (Thermo

Fisher Scientific), *via Xho*I and *BamH*I sites. The pL1 expression vectors were constructed based on a vector

containing the PpActin5 promoter, nos terminator and hpt cassette (Ruiz-Molina et al., 2022a). The pL1

sequence was synthesized by Genewiz (Leipzig, Germany). For constructs pL1 and pL1Δ22, the pL1 CDS

was amplified using Phusion^{TM} polymerase (Thermo Fisher Scientific) and cloned as described above. For

construct CTP-pL1, the pL1 and CTPc5 CDS were amplified and inserted into the backbone using Gibson

Assembly. Vectors were sequenced by Eurofins Genomics (Ebersberg, Germany). Vector maps are

compiled in **Supplementary Figure S1**. Primers were synthesized by Eurofins Genomics and are compiled

in **Supplementary Table S1**.

Protoplast preparation and transformation

Physcomitrella protoplasts were prepared and transformed based on existing protocols (Hohe &

Reski, 2002; Schween et al., 2003; Hohe et al., 2004; Decker et al., 2015). Transformed protoplasts were

incubated at 22 °C for 24 h in the dark and subsequently for 72 h at a 16/8-h light/dark photoperiod and

light intensity of 50–70 µmol m^{-2} s^{-1} before plating on KnopME + 25 mg/L hygromycin agar plates covered

with cellophane for two weeks. Surviving colonies were transferred to non-selective conditions for two

weeks before a second selection on KnopME + 50 mg/L hygromycin.

Fluorescence microscopy

Protoplasts were analysed 2-3 days after transfection with an AxioCam MRc5 camera on an

Axioplan2 stereo microscope (Zeiss, Oberkochen, Germany). Pictures were taken with a 40x objective or

a 63x objective with immersion oil. A FITC filter was used for eGFP, a rhodamine filter for chlorophyll

autofluorescence, and a FITC_LP filter for the combination of the two. Images were edited with GIMP

2.10.22 and ImageJ. For confocal microscopy, an AXIO Observer.Z1 Inverted Fluorescence Motorized

Phase Contrast Microscope Pred 7 with the objective Plan-Apochromat 63x/1.4 Oil DIC M27 (Zeiss) was

used. EGFP signals were detected at 488 / 509 nm, and chlorophyll autofluorescence at 587 / 610 nm.

Images were edited in ZEN 3.5 (Blue edition, ZEN Lite), ImageJ, and GIMP 2.10.22.

For analysis of subcellular targeting, an inverted confocal laser scanning microscope LSM 880 with

the objective LD LCI Plan-Apochromat 40x/1.2 H_2O autocorr and FastAiryScan mode (Zeiss) was used.

Citrine signals were detected at 488 / 516 nm, and chlorophyll autofluorescence at 561 / 595 nm. Images

were edited in ZEN 3.9 (Blue edition) and processed *via* deconvolution using default setting and

orthogonal projection using maximum intensity projection.

Biomass accumulation

Flasks containing 30 mL KnopME medium were inoculated at a density of 60 mg DW/L and

cultured for 4 weeks at the conditions described above. Biomass DW was measured by filtering 3x 10 mL

of culture through miracloth and subsequent drying at 105 °C for 2 hours. Biomass accumulation was

analysed by linear regression analysis. Slopes were compared using ANCOVA and post hoc multiple

comparisons test (TukeyHSD). The analysis was conducted in RStudio (version 4.2.2) and GraphPad Prism

(version 8.1).

Bioreactor operation

Stirred tank bioreactors (Applikon, Schiedam, Netherlands) were inoculated at 60 mg DW/L moss

density in 5 L KnopME. The light intensity was set to 160 µmol m^{-2} s^{-1} and increased to 350 µmol m^{-2} s^{-1}

after 4 days. At day 4, auxin 1-Naphtalene acetic acid (NAA, Sigma, St. Louis, USA) dissolved in 0.5 M KOH

was added to a final concentration of 5 µM. Throughout the bioreactor run, the pH was kept at pH 5.8,

temperature at 22 °C, and aeration at 0.3 vvm with 2% CO_2. The culture was continuously agitated with a

pitched 3 blade impeller at 500 rpm.

Protein extraction, purification and VLP assembly

For small scale extraction, 30 – 60 mg FW of vacuum-filtered moss material was combined with a

glass and a steel bead (each 3 mm in diameter) in a 2 mL reaction tube, frozen in liquid nitrogen and the

tissue lysed using a tissue lyser for 2 min at 28 Hz. Homogenized material was resuspended in PBS (2.7

mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 0.137 M NaCl, pH 7.2), 0.01% Tween20, and protease inhibitor

(P9599, Sigma) in a ratio 3 mL/g. Samples were placed in an ice-cold sonication bath for 15 min,

centrifuged for 30 min at 14,000 x g and 4 °C, and the supernatant taken off for further analysis.

For large scale protein extraction, 5 g FW of vacuum-filtered moss material were frozen in liquid

nitrogen and ground to a fine powder. The material was combined with 22 mL PBS and 0.01% Tween20,

50 mM DTT and 100 µL protease inhibitor. The mixture was homogenized using an ULTRA-TURRAX (IKA,

Staufen, Germany) at 10,000 rpm for 10 min on ice. Next, the extract was treated with a Q500 sonicator

using a CL-334 converter (QSonica, Newtown, USA) at amplitude 55%, 10s on & 40s off for 20 min at 8 °C.

Debris was removed by two centrifugations at 4500 x g for 10 min and 20,000 x g for 20 min, both at 4 °C.

The sample was combined with solid $(NH_4)_2SO_4$ to a final concentration of 45% and proteins precipitated

under stirring for 1 h at 8 °C. Precipitated proteins were pelleted by centrifugation at 12,000 x g for 10

min at 8 °C. Pellets were dissolved in 4 mL CaEx chromatography binding buffer (2.7 mM KCl, 10 mM

Na_2HPO_4, 1.8 mM KH_2PO_4, 0.068 M NaCl, 0.01% Tween 20, 5% Glycerol, 5 mM DTT, pH 6.8) and dialysed

(20k MW cut off, 88405, Thermo Fisher Scientific) overnight against the same buffer at 8 °C. The sample

was diluted to a final volume of 10 mL using binding buffer, and passed through a 0.22 µm filter (Roth,

Karlsruhe, Germany) before loading onto a 1 mL HiTrap SP HP cation exchange column (Cytiva,

Marlborough, USA), connected to an ÄKTA^{TM} start (Cytiva). The column was washed with 5 column

volumes (CV) of 0.3 M NaCl (based on binding and elution buffer) and L1 eluted *via* 4 CV of elution buffer

(2.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 1 M NaCl, 0.01% Tween 20, 5% Glycerol, 5 mM DTT, pH

6.8)

Eluted fractions containing L1 were pooled and dialysed against disassembly buffer (22.7 mM KCl,

10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 0.068 M NaCl, 0.05% Tween20, 5 mM DTT, pH 8.5) overnight at 8 °C

with 3x buffer exchange. Samples were concentrated 5x using a vacuum concentrator (Concentrator plus,

Eppendorf, Hamburg, Germany). For reassembly, samples were dialysed against reassembly buffer (22.7

mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 1 M NaCl, 0.05% Tween20, pH 6) overnight at 8 °C with 3x

buffer exchange.

SDS-PAGE and Western blot

SDS-PAGE and Western blots were performed based on Bohlender et al. (2022). Two systems

were used for protein separation, either 7.5% polyacrylamide gels (Mini-PROTEAN® TGX^{TM} Precast Gels,

Rio-Rad, Munich, Germany) in TGS buffer at 120 V, or 4 – 12% Bis-Tris gels (mPAGE^{TM} Precast Gels, Merck,

Darmstadt, Germany) in MOPS buffer at 150 V. As first antibody served 1:5000 diluted CamVir1 (Thermo

Fisher Scientific), as second antibody peroxidase-linked rabbit anti-mouse secondary antibody (Cytiva)

diluted 1:12,500. For dot blots, protein samples were blotted on a PVDF membrane (Cytiva) using a

filtration   manifold   kit   (SRC-96/1,   Schleicher&Schuell,   Dassel,   Germany).   Antibody   H16.V5

(AntibodySystems, Schiltigheim, France) was diluted 1:5000. Dot blots were developed as described for Western blots. Yeast-produced HPV-16 L1 (Abcam, Cambridge, UK) was used as a positive control.

L1 quantification

The ELISA protocol was based on Studentsov et al. (2002). In short, 96-well microtiter plates (Greiner, Kremsmünster, Austria) were coated with 1:500 CamVir1 antibody (Thermo Fisher Scientific) in PBS overnight at 8 °C. Wells were washed twice with washing buffer (1x PBS, 0.05% Tween20) and blocked with blocking buffer (1x PBS, 1% polyvinyl alcohol MW 47,000, 0.05% Tween20) for 3 h at 37 °C. After 3x washing, samples were diluted in blocking buffer and incubated in the wells for 1.5 h at 37 °C. Wells were washed 3x before adding 1:1000 diluted polyclonal anti-HPV-16 L1 antibody from rabbit (Cusabio, Houston, USA) for 1 h at 37 °C. After 5x washing, wells were incubated with 1:1000 diluted anti-rabbit HRP-conjugated antibody NA934 (Cytiva) for 1 h at 37°C. Wells were washed 3x before adding 100 µL TMB substrate (Thermo Fisher Scientific). The reaction was stopped with 50 µL of 2 M H_2SO_4. Absorbance was measured at 450 / 595 nm for reference. The signal of a wildtype sample was subtracted to account for background signal.

Transmission electron microscopy

TEM protocols are based on Abel et al. (1989). Negative staining of purified VLPs and moss extracts was performed using 300 Mesh glow discharged formvar/carbon-coated copper grids (Electron Microscopy Sciences, Hatfield, USA). Five µL of sample were applied on a grid and incubated for 5 min at RT. Grids were washed 3x with water and stained with 2% (w/v) uranyl acetate. After 30 sec, uranyl acetate was removed with a filter paper, grids were air dried, and inspected with a Hitachi HT7800 TEM (Tokyo, Japan) coupled to a Xarosa CMOS camera (Emsis, Münster, Germany).

Results

Adaptation of L1 CDS for Physcomitrella-optimized expression

The CDS of HPV-16 L1 was optimized for expression in Physcomitrella. The goal was to adapt the CDS regarding codon usage and preventing potential degradation through microRNAs and splicing. A human codon-optimized version (hL1, **Supplementary Figure S2**) was used as a basis for optimization since it had led to higher yields than plant-optimized versions or the native viral sequence in *Nicotiana benthamiana* (Maclean et al., 2007).

For codon optimization, physCO (Top et al., 2021) was used, which changed nine out of 505
codons and slightly increased the GC content from 62.25 % to 62.85 % (**Supplementary Figure S2**). Next,
the CDS was scanned for the motif AGGT, which is the most common motif at exon-intron borders for
alternative splicing in Physcomitrella (Top et al., 2021). Transformation of a heterologous cDNA containing
the AGGT motif led to alternative splicing in Physcomitrella, a process referred to as heterosplicing (Top
et al., 2021). Scanning the physCO-optimized hL1 sequence yielded 5 AGGT-motifs which were replaced
by synonymous codons (**Supplementary Figure S2**). MicroRNAs bind complementary sequences in the
mRNA and thereby cause inhibition of translation or degradation of mRNA (Jones-Rhoades et al., 2006;
Khraiwesh et al., 2010). One microRNA binding site was predicted within the CDS using psRNATarget (Dai
& Zhao, 2011; Dai et al., 2018). Therefore, we changed three bases within the predicted binding region
using synonymous codon substitutions. The final Physcomitrella-optimized CDS was named pL1
(**Supplementary Figure S2**) and differs from hL1 by 17 out of 1515 nucleotides without changing the aa
sequence.

Characterization of chloroplast transit peptides

Different constructs were designed to target L1 to specific cellular compartments, including
chloroplasts. Fusing chloroplast transit peptides (CTPs) to the L1 CDS led to increased protein yields in
tobacco as compared to constructs lacking CTPs (Maclean et al., 2007; Zahin et al., 2016). To assess
whether this is true for L1 production in Physcomitrella, a selection of Physcomitrella-derived CTPs was
characterized.

Initially, the CTP of FtsZ1-1 (Gremillon et al., 2007) was considered. However, cleavage site
analysis *via* TargetP-2.0 (Armenteros et al., 2019) predicted 37 aa of this CTP to remain at the N-terminus
of L1 after chloroplast import. This CTP was rejected because the additional aa could change L1 properties
and hamper VLP assembly. A list of alternative CTPs was derived from the Physcomitrella N-terminome
(Hoernstein et al., 2024) (**Supplementary Table S2**). Here, experimentally detected protein N-termini
were compared with TargetP-2.0 predictions for cleavable N-terminal targeting sequences. Five CTPs were
selected where an exact match between prediction and experimentally validated N-terminus was
observed across different experiments (Hoernstein al., 2024), namely CTPc5, CTPc10.1, CTPc10.2, CTPc21,
CTPc22 (**Table 1**). Their sequences were fused *in silico* to the N-terminus of L1 and analysed with TargetP-
2.0 to validate the correct cleavage of the fusion.

Our *in-silico* analysis indicated CTPc22 to end with a methionine, which could introduce an

alternative translation initiation site, resulting in translation of L1 without the CTP (**Table 1**). Therefore,

an alternative version lacking the C-terminal methionine (CTPc22noMet) was included. The likelihoods for

chloroplast localization were above 70 % for all candidates except CTPc10.1 (**Table 1**), which was rejected.

Table 1: *In-silico* analysis of Physcomitrella-derived chloroplast transit peptides. CTPs were selected based on mass

spectrometry datasets (Hoernstein et al., 2024). *In-silico* predictions of likelihood of chloroplast localization target

sequence cleavage were performed with TargetP-2.0.

Name of CTP	Originating from protein model	Predicted cleavage of CTP-L1	Length (aa)	Likelihood of GFP chloroplast localization	Likelihood of L1 chloroplast localization
FtsZ1-1	Pp3c22_4940V3.1	37 aa remain on N-terminus	84	0.95	0.92
CTPc5	Pp3c5_1370V3.1	-	74	0.987	0.943
CTPc10.1	Pp3c10_10490V3.1	-	63	0.456	0.229
CTPc10.2	Pp3c10_2940V3.1	-	70	0.908	0.845
CTPc21	Pp3c21_9980V3.1	-	43	0.956	0.891
CTPc22	Pp3c22_5470V3.1	-	62	0.939	0.739
CTPc22noMet	Pp3c22_5470V3.1	-	61	0.913	0.727

The remaining five CTPs (CTPc5, CTPc10.2, CTPc21, CTPc22, CTPc22noMet) were fused to the CDS

of eGFP (**Supplementary Figure S1**) and expressed in Physcomitrella protoplasts. Analysis *via* fluorescence

microscopy confirmed localization of eGFP signals in chloroplasts for CTPc5, CTPc10.2, CTPc22, and

CTPc22noMet, but not for CTPc21 (**Supplementary Figure S3**). Of the four remaining CTPs, CTPc5 had the

highest likelihood for plastid localization (**Table 1**) and were examined *via* confocal microscopy

(**Supplementary Figure S3**). Finally, we selected CTPc5 for the chloroplast-targeting of L1.

Targeting of L1 to nuclei and chloroplasts

To evaluate the impact of L1 localization on the yield, we generated three variants (pL1, pL1Δ22,

CTP-pL1). pL1 contains the full-length pL1 CDS, which includes the native HPV-16 NLS (Cerqueira & Schiller,

2017). For pL1Δ22 the NLS was removed. In CTP-pL1 the Physcomitrella CTPc5 was fused to the 5' of the

pL1 CDS, including the NLS.

After generation of these CDS variants, we aimed to evaluate L1 targeting to subcellular

compartments. The CDS of pL1, pL1Δ22 and CTP-pL1 were fused to the 5' end of Citrine CDS

(**Supplementary Figure S1**). Physcomitrella protoplasts were transfected with the plasmids and Citrine

localization was analysed using fluorescence microscopy. A vector encoding Citrine without any L1 variant

fusion served as a control.

Cells transformed with the control vector exhibited predominantly cytoplasmic Citrine signals,

whereas diffuse signals also occurred in nuclei (**Figure 1**). For pL1-Citrine, the signal localized to nuclei,

demonstrating that the HPV-16 NLS is functional in Physcomitrella. This is further supported by the fact

that removal of the NLS in pL1Δ22-Citrine results in cytoplasmic localization. For CTP-pL1-Citrine, the

signal is predominantly in chloroplasts, but is not evenly distributed, as seen for CTP-eGFP fusions lacking

L1 (**Supplementary Figure S3**). Instead, the pL1-Citrine fusion concentrates in several spots of high

fluorescence intensity and <1 μm in diameter. Overall, we confirmed successful L1 targeting to the

intended compartments.

Figure 1: Subcellular localization of L1 variants. Confocal microscopy images of Physcomitrella protoplasts transformed with

constructs encoding (CTP-)pL1(Δ22)-Citrine fusion proteins or Citrine alone. For each sample, chlorophyll autofluorescence in

magenta is seen on the left, Citrine signals in the middle and the merged signals on the right. In the control and pL1Δ22-Citrine,

the Citrine signals do not enter the chloroplasts and are mainly in the cytoplasm. For pL1-Citrine, the Citrine signals are in the

nucleus. In the CTP-pL1-Citrine sample, the Citrine signals form small spots of high intensity in the chloroplasts.

Characterization of L1 production

After validating the subcellular localization, pL1, pL1Δ22 and CTP-pL1 expression constructs were

cloned (**Supplementary Figure S1**). The linearized constructs were transferred into Physcomitrella

protoplasts and obtained plants were screened for productivity by Western blot and ELISA.

Western blot analysis revealed the L1 monomer of 56 kDa (pL1, CTP-pL1) and 54 kDa (pL1Δ22)

(**Figure 2**). In addition, pL1 and CTP-pL1 lines exhibited a band at approximately 54 kDa. All lines showed

another band at around 45 kDa, even though the signal was relatively weak in CTP lines and more

prominent in pL1Δ22 lines. Smaller protein isoforms can originate from degradation or DNA or RNA,

resulting in truncated transcripts and, subsequently, truncated L1 isoforms (Murén et al., 2009; Top et al.,

2021). The latter possibilities were ruled out by amplifying the L1 CDS *via* PCR from cDNA, which yielded

amplicons of the expected sizes, corresponding to the Western blot bands of 56 kDa (pL1, CTP-pL1) and

54 kDa (pL1Δ22), but not additional amplicons (**Supplementary Figure S4**). Also, two bands are visible at

130 – 180 kDa, of which the highest band at 180 kDa is also present in the yeast-produced L1 control. This

pattern of two close-by bands repeats another two times at higher molecular weight outside of the marker

range (**Figure 2**). ELISA analysis indicates higher yields in CTP-pL1 lines as compared to pL1 or pL1Δ22 lines.

In shaking flasks, the best producer is CTP-pL1 30 with 142 µg L1/g FW (**Figure 2**).

Figure 2: Analysis of HPV-16 L1 production *via* Western blot and ELISA. (**a**) For each of the constructs pL1, pL1Δ22, and CTP-pL1 three lines were tested *via* Western blot for expression of HPV-16 L1. 50 ng *S. cerevisiae*-produced L1 served as a positive control and Physcomitrella wildtype as a negative control. The proteins for the Western blot were separated on a 7.5% TGS polyacrylamide gel under reducing conditions before being transferred to a membrane and developed using CamVir1 antibody (1:5000). (**b**) Production levels of L1 were quantified *via* ELISA. Wildtype background signal was subtracted before analysis *via* one-way ANOVA and post hoc multiple comparisons test (TukeyHSD). Means and standard deviations are calculated based on n ≥ 3 technical replicates from two dilutions. Statistical significance (α = 0.05) is indicated by Compact Letter Display.

Besides the amount of recombinant protein per g FW biomass, the accumulation of biomass is important in biomanufacturing as high product yields per biomass can be trivial if biomass accumulation is slow. To study if recombinant protein amount or subcellular localization of L1 hampers plant growth, we studied the time course of biomass accumulation of the nine lines in shaking flasks for 4 weeks.

In our conditions, exponential growth could not be measured. This might be due to the culture conditions, to the rather long intervals between sampling, and/or to the intrinsic growth characteristics

of Physcomitrella protonema, which grows by apical cell division (Reski, 1998). Therefore, biomass density

was fitted using linear regression analysis model (**Supplementary Figure S5**), using the calculated slope to

approximate biomass accumulation rates. Comparison of the slopes between the lines pL1, pL1Δ22 and

CTP-pL1 showed pL1 lines accumulated biomass faster than pL1Δ22 (p = <0.0001) and CTP-pL1 (p = 0.0001)

lines. If this difference is due to the distinct L1 localization or to lower L1 production in pL1 lines (**Figure**

2) is unclear. Line pL1 47B showed the highest slope of 255 mg DW/L/week, as compared to line CTP-pL1

30 with a slope of 211 mg DW/L/week. However, this difference is negligible considering the yield of pL1

47B is 8 µg L1/g FW as to 142 µg L1/g FW for CTP-pL1 30 (**Figure 2**). Consequently, line CTP-pL1 30 was

chosen for VLP production.

Upscaled L1 production

For production of VLPs, the cultivation of line CTP-pL1 30 was upscaled to 5 L-photobioreactors.

The goal was to identify the best time point for harvesting material used for subsequent L1 purification.

Biomass increased from initially 60 mg DW/L to around 870 mg DW/L at day 4 (**Figure 3**). L1

production remained constant during the first 4 days, in the range of the yield observed for flasks. Upon

addition of the phytohormone auxin on day 4, L1 levels increased on day 6 and peaked at day 7 at 332 µg

/ g FW, 130% higher as compared to day 0. Afterwards, L1 production dropped sharply to levels prior to

auxin addition. Biomass continued to accumulate until day 11, after which it dropped slightly. These

conditions, including day 7 as the ideal harvesting time point, were used for further L1 production.

Figure 3: Analysis of biomass accumulation and L1 production of line CTP-pL1 30 in a 5 L-photobioreactor. The bioreactor was

inoculated with 60 mg dry weight (DW)/L and monitored over a course of 13 days. On day 4, 5 µM NAA was added. L1 production

peaked at day 7 at 332 µg / g fresh weight (FW). Biomass and L1 measurements are based on n ≥ 3 technical replicates. Production

levels of L1 were quantified *via* ELISA.

Two-step purification removes bulk of cellular proteins

Purification of L1 was performed using ammonium-sulphate (AMS) precipitation, followed by cation exchange (CaEx) chromatography. While AMS removes the bulk of cellular proteins, CaEx removes residual proteins and concentrates the sample.

We tested the efficacy of 40%, 45% and 50% AMS to precipitate L1 from an extract containing the soluble protein fraction from CTP-pL1 30 plant material produced in the bioreactor. We found 45% AMS to be sufficient to effectively remove L1 from the supernatant, whereas at 40% AMS small amounts of L1 remained in the supernatant and at 50% AMS higher amounts of cellular protein were precipitated together with L1 (**Supplementary Figure S6**). In preparation for chromatographic purification, the precipitated proteins were resuspended in CaEx binding buffer and dialysed overnight in binding buffer to remove residual AMS. Cloudiness of samples indicated a reduced solubility of precipitated proteins, and undissolved matter was removed by filtration. The relative amount of L1 and host protein was evaluated *via* Western blots and Coomassie-staining, respectively (**Supplementary Figure S6**). Samples after filtration showed a slight reduction in host protein content, but, except for the sample at 40% AMS, no reduction in L1. Overall, precipitation at 45% AMS was selected, which is in line with purifications of yeast-produced L1 (Park et al., 2008; Kim et al., 2010).

Next, CaEx chromatography was used to remove residual cellular proteins. The bulk of cellular proteins did not bind to the column and was detected in the flow through (**Figure 4**, **Supplementary Figure S7**). Notably, a large amount of L1 did not bind to the column and was detected in the flow through. Different binding buffer compositions were tested to increase L1 binding to the column, e.g. lowering the pH from 7.2 to 6.8, lowering the concentration of NaCl from 0.137 M to 0.068 M, addition of 5% glycerol, and re-loading of flow-through fraction. Nevertheless, a large part of L1 did not bind to the column and was found in the flow through. During column wash with 5 mL at 0.3 M NaCl, small amounts of contaminating proteins and L1 were washed off the column. For elution, NaCl was increased to 1 M NaCl and 0.5 mL fractions were collected, in total 4 mL. L1 eluted from the column in fractions E4 – E8 (**Figure 4**), which were pooled for further analysis.

In contrast to the confirmation of L1 expression (**Figure 2**) and the establishing of AMS precipitation (**Supplementary Figure S6**), DTT was used throughout the extraction and purification leading to CaEx chromatography. In comparison, the presence of DTT markedly reduces the Western blot signal intensity above 180 kDa (**Figure 4**). We hypothesize that in the absence of DTT, reactive oxygen species in

438   the plant extract cross-link L1 with itself and/or host proteins. Irrespective of the presence or absence of

439   DTT during extraction and purification, two prominent bands in the range of 130 and 180 kDa are present

440   in all Western blots (**Figures 2 and 4, Supplementary Figure S6**).

Figure 4: Two-step purification of L1 removes bulk of cellular protein. Analysis of L1 and residual cellular host proteins during

444   purification under reducing conditions *via* Western blot and Coomassie-stained SDS-PAGE (12% Bis-Tris gel). After precipitation

445   with 45% AMS, the filtered sample was subjected to cation exchange chromatography (CaEx) for final purification. Column-bound

446   L1 eluted at 1 M NaCl in fractions E4 – E8. The asterisk in the Coomassie-stained SDS-PAGE marks the band referring to HPV-16

447   L1. 50 ng *S. cerevisiae*-produced L1 served as a positive control and Physcomitrella wildtype as a negative control.

Capsomers and VLPs

Next, we performed a dis- and reassembly protocol on the pooled elution fractions, as it yields

more uniform and stable VLPs, which are less prone to aggregation (McCarthy et al., 1998; Mach et al.,

2006). First, pooled elution fractions were dialysed at low salt concentrations, high pH and DTT, that

results in the disassembly of VLPs or other multi-pentameric L1-based structures into capsomers. Second,

samples were dialysed at high salt, low pH and in the absence of reducing agent, that leads to the

reassembly of capsomers into VLPs.

To assess the conformation of L1 throughout the process, dot blots were performed using

different monoclonal antibodies. CamVir1 detects a linear epitope in both denatured and native L1 and

served as a control. In contrast, H16.V5 detects only conformational epitopes in both, L1 capsomers and

VLPs, but not monomeric or denatured L1.

The dot blot showed the presence of L1 in all CTP-pL1 30-derived samples throughout the

purification process, as well as yeast-derived denatured and native L1 (**Figure 5**). In comparison, H16.V5

detects L1 in the native positive control and in all CTP-pL1 30-derived samples, indicating the presence of

at least capsomers, if not VLPs, already in the raw extract. Further, capsomers seem to be stable

throughout AMS precipitation, CaEx chromatography, disassembly and reassembly.

Figure 5: Moss-produced HPV-16 L1 assembles into capsomers, which are stable throughout purification. Dot blots were

performed using two different antibodies to analyse the formation of capsomers in yeast-produced L1 and samples from CTP-

pL1 30 purification (extract, after AMS precipitation, elution) and dis- and reassembly. The CamVir1 antibody detects a linear

epitope in both denatured and native L1. The neutralizing antibody H16.V5 detects specifically conformational epitopes in L1

capsomers and VLPs. Positive H16.V5 signals indicate HPV-16 L1 monomers to assemble at least into capsomers, *in vivo* or after

extraction, which are stable throughout the purification. The experiment only allows qualitative but not quantitative comparison

since sample concentrations were adjusted individually. `L1 denatured´ was incubated at 95 °C for 5 min, all other samples were

kept at 4 °C. A wildtype sample was used as a negative control.

The assembly of L1 capsomers into VLPs was verified with Transmission Electron Microscopy

(TEM). As a control, yeast-produced L1 was analysed without prior treatment, after disassembly, and after

reassembly. In the untreated sample, particles of ca. 60 nm in size were detected. After disassembly,

structures of 5 – 10 nm in size were observed, corresponding to capsomers. VLPs reshape upon
reassembly treatment, demonstrating the successful dis- and reassembly of yeast-derived L1 VLPs.
Notably, reassembled VLPs are heterogenous in size and shape, despite the controlled dis- and reassembly
(**Figure 6**).

Figure 6: Analysis of dis- and reassembled VLPs from moss and yeast *via* transmission electron microscopy (TEM). Yeast-derived
L1 was analysed untreated (**a**), after disassembly (**b**), and after reassembly (**c**) *via* TEM. CTP-pL130 (**d** – **f**) and wildtype (**g** – **i**)
extracts were purified *via* AMS precipitation and CaEx chromatography (**d** & **g**), after disassembly (**e** & **h**), and after reassembly (**f**
& **i**). Untreated yeast-derived VLPs (**a**) disintegrate into capsomers upon disassembly (**b**) and reshape into VLPs upon reassembly
treatment (**c**). TEM shows the presence of moss-derived HPV-16 L1 capsomers in the elution (**d**) and disassembly sample (**e**) and
the assembly into VLPs in the reassembly sample (**f**). Capsomer and VLP-like structures are absent from the wildtype control
samples. VLPs are indicated by black arrows.

493   To analyse L1 assembly into VLPs, samples were taken from the pooled CaEx chromatography
elution fractions, after disassembly, and after reassembly.

In the elution sample, capsomeric structures comparable to those observed in the disassembled
yeast sample were observed, which are not present in the corresponding wildtype control (**Figure 6**). The
abundance of capsomers in the elution fractions was surprising given the fact that the elution buffer has
a high ionic strength, which favours reassembly. However, before column loading, the sample was
dialysed overnight in binding buffer, which favours disassembly. The exposure to high ionic strength
during elution might have been too short to allow VLP assembly. In addition, both binding and elution
buffer contained DTT, which induces disassembly (McCarthy et al., 1998; Mach et al., 2006). The
disassembled CTP-pL1 30-derived sample depicted capsomers as seen in the elution fraction, and for
disassembled yeast-derived L1 (**Figure 6**). **Figure 6f** displays fully assembled, moss-derived HPV-16 L1
VLPs. Again, VLPs were heterogenous in size and shape, as seen for reassembled yeast-derived VLPs.
Moss-produced VLPs tended to aggregate, which was not observed for reassembled yeast-derived VLPs,
but is common for HPV-16 L1 VLPs (Shi et al., 2005). No VLPs were detected in wildtype samples (**Figure**
6).

Discussion

Various strategies were explored for increasing production yields of HPV-16 L1 VLPs in plants, one
of which is optimization of codon usage (Maclean et al., 2007; Hitzeroth et al., 2018). The latter found that
codon optimization of L1 improves transcription, rather than translation. This is due to RNA motifs in the
L1 CDS, which posttranscriptionally regulate the virus life cycle and lead to pre-mRNA degradation (Mori
et al., 2006; Zhao & Schwartz, 2008). It is argued, that codon optimization removes those motifs, thereby
increasing mRNA and protein levels, irrespective of whether the CDS was optimised for expression in
humans, tobacco (Hitzeroth et al., 2018) or Physcomitrella (this study). Our Physcomitrella-optimized CDS
has a GC content of 62.85% as compared to hL1 with 62.25% (Maclean et al., 2007) and the wildtype
sequence with 38.27% (Varsani et al., 2003). Combined with the removal of microRNA binding sites and
heterosplice sites we obtained Physcomitrella-optimized pL1 CDS. We argue that pL1 addresses the issues
of L1 intrinsic degradation motifs as well as Physcomitrella-specific posttranscriptional degradation.
Besides codon optimization, research focused on localization signals or transit peptides. The most
common strategy is sequestration to chloroplasts, either *via* nuclear transformation and posttranslational
import into chloroplasts (Maclean et al., 2007; Pineo et al., 2013; Zahin et al., 2016; Hitzeroth et al., 2018;
Naupu et al., 2020; Muthamilselvan et al., 2023) or *via* transplastomic lines (Fernández-San Millán et al.,
2008; Lenzi et al., 2008). Here, we experimentally characterized five Physcomitrella CTPs, of which we

used CTPc5 for sequestration of L1 to chloroplasts. Employing the HPV-16 native NLS, L1 was targeted to

nuclei, whereas the removal of the NLS resulted in cytoplasmic L1. Earlier studies confirmed functionality

of the NLS in mammalian cells (Leder et al., 2001; Bissa et al., 2015). However, the NLS failed to localize

L1 to nuclei in tobacco (Šmídková et al., 2010). Our observation of the HPV-16 native NLS targeting L1 to

nuclei is another example for a conserved cross-kingdom functionality of localization signals between

Physcomitrella and mammals or human-infecting viruses (Gitzinger et al., 2009).

Western blots of L1-producing lines showed a variety of bands, both smaller and larger than the

expected 56 kDa or 54 kDa. A comparable banding pattern was observed when expressing L1 in tobacco

chloroplasts (Fernández-San Millán et al., 2008). Likewise, yeast-produced L1 exhibited bands with the

size of L1 dimers and trimers (Kim et al., 2007; Kim et al., 2010). L1 dimers and trimers are based on inter-

capsomeric disulfide bridges which stabilize virions and VLPs (Sapp et al., 1998; Buck et al., 2005; Buck et

al., 2013). McCarthy et al. (1998) resolved L1 multimers with DTT. Likewise, Buck et al. (2005)

demonstrated absence and presence of L1 dimers and trimers in a comparison of reducing and non-

reducing SDS-PAGE and Western blots, respectively. An investigation of cysteine 175 and cysteine 428 L1

mutants resulted in the absence of bands at 130 kDa and 180 kDa, thereby linking disulfide bridge-based

dimers and trimers to the observed bands (Buck et al., 2005). Surprisingly, reducing agents such as DTT or

β-mercaptoethanol did not result in the disappearance of these bands in our and other studies (Kim et al.,

2007; Fernández-San Millán et al., 2008; Kim et al., 2010). This indicates a tolerance of L1 dimers and

trimers to reducing agents or a different origin of multimerization observed at 130 kDa and 180 kDa.

Castells-Graells & Lomonossoff (2021) observed multimerization of *Nudaurelia capensis* omega virus α-

peptide produced in tobacco, irrespective of DTT, but not when produced in insect cells. These authors

hypothesize that cross-linkage of α-peptide occurs during extraction and purification and occurs in plant-

based expression systems. Regarding the observed bands <56 kDa, proteolytic degradation was proposed

in other studies observing the same phenomenon (Buck et al., 2005; Fernández-San Millán et al., 2008).

In tobacco, a shift from bands above 100 kDa in young leaves to more prominent bands at 45 kDa and 20

kDa in older leaves indicated proteolytic degradation of L1 over time (Fernández-San Millán et al., 2008).

In our study, bands at 45 kDa are most prominent in pL1Δ22 lines with cytoplasmic L1 and less prominent

for plastidic L1. This might result from organelle-specific proteases (Coates at al., 2022).

In Physcomitrella, combining the full-length L1 with a CTP significantly increases L1 yield as

compared to full length L1 or L1 without NLS, which is in line with findings in tobacco (Maclean et al.,

2007; Zahin et al., 2016). In our study, L1 yields above 0.3 mg/g FW were achieved in the photobioreactor.

In tobacco, yields of 0.1 – 1.2 mg/g FW (Maclean et al., 2007; Pineo et al., 2013; Zahin et al., 2016;

Hitzeroth et al., 2018) and up to 3 mg/g FW in transplastomic lines have been reported (Fernández-San

Millán et al., 2008). The success of targeting L1 to chloroplasts over localization to other organelles may

have several reasons. One is that plastid L1 may be protected from proteases absent in chloroplasts (van

Wijk, 2024). Besides, CTPs also act on the transcriptional level. Upon expression in tobacco, chimeric L1/L2

transcripts containing a CTP were more abundant than transcripts lacking a CTP, and consequently yielded

more protein (Hitzeroth et al., 2018). So far, it is unclear whether this difference is due to enhanced

transcription, increased mRNA half-life or changes in mRNA secondary structure.

For purification of L1 from Physcomitrella extracts, a combination of AMS precipitation and CaEx

chromatography was employed. We found 45% AMS to be sufficient to efficiently precipitate L1. Studies

from yeast (Park et al., 2008; Kim et al., 2010) and tobacco (Zahin et al., 2016) also used 45% AMS, whereas

sometimes 50% AMS was used (Kim et al., 2018). CaEx chromatography showed that a large part of L1

does not bind to the column. Likewise, Zahin et al. (2016) reported tobacco-produced L1 to bind poorly

to CaEx columns. Kim et al. (2010) purified yeast-produced L1 by either CaEx or heparin chromatography.

CaEx chromatography showed unbound L1 in the flow through, as observed in our study, whereas no L1

was found in the flow through employing heparin chromatography. Pineo et al. (2013) produced L1/L2

chimaeras in tobacco and experienced loss of L1 and insufficient removal of contaminating proteins using

heparin chromatography. Heparin is a highly sulphated form of heparan sulphate, a proteoglycan found

on host cell surfaces and a requirement for HPV virion attachment and infection (Knappe et al., 2007).

However, a prerequisite for heparin binding is the assembly of L1 to capsomers or VLPs (Rommel et al.,

2005), meaning monomers and not properly folded L1 are lost. Overall, chromatography yields for VLPs

differ and are influenced by different factors, including the production host, the used materials and

buffers, and purification prior to chromatography, making it difficult to compare purification strategies

across publications. Here, we achieved to remove the bulk of cellular proteins from moss extracts, but

experienced a substantial loss of L1 in the flow through. To address this, further optimization of

purification might be achieved by combinations with other methods, such as heparin chromatography.

Our dot blot analysis confirmed the presence of intact capsomers or VLPs in extracts and after AMS

precipitation, suggesting heparin chromatography as an option for the second step in purification.

TEM analysis of purified and reassembled CTP-pL1 samples shows VLPs of ca. 60 nm in size. These

nanoparticles assembled from capsomers as a result of the reassembly treatment. To the best of our

knowledge, this is the first report of VLPs produced in a non-vascular plant and of *in vitro* reassembly of

HPV-16 L1 VLPs from capsomers produced in any plant system. The VLPs are heterogenous in shape and

size. The same is true for yeast-derived VLPs after reassembly, whereas untreated VLPs were not

heterogenous. This is surprising, since controlled dis- and reassembly of VLPs can enhance stability and
uniformity (McCarthy et al., 1998; Mach et al., 2006; Shi et al., 2007). Besides high ionic strength and low
pH buffer, inter-pentameric disulfide bridges positively influence VLP stability and uniformity (Buck et al.,
2005). This could be achieved through prolonged incubation of cleared cell lysates before purification or
addition of oxidizing agents, which accelerate disulfide bridge formation. However, prolonged incubation
causes aggregation of HPV-16 L1 VLPs, as observed here, but could be addressed by improved formulation
(Shi et al., 2005).

In conclusion, we generated a CDS optimized for expression of HPV-16 L1 in Physcomitrella. We
further characterized five CTPs and confirmed the localization of L1 in chloroplasts. In addition, we
demonstrated chloroplast localization to yield higher L1 levels than targeting to nuclei or cytoplasm.
Cultivation in photobioreactors yielded more than 0.3 mg/g FW, which is in the range of other plant-based
expression platforms. We established a purification protocol and demonstrated the successful assembly
of moss-produced HPV-16 L1 VLPs. Future efforts should focus on further enhancing purification and
formulation of VLPs for immunological studies of this vaccine candidate.

Acknowledgements

606 We thank the Life Imaging Centre (LIC) of the University of Freiburg for support with confocal microscopy.
607 We are grateful for technical assistance by Agnes Novakovic, Britta Rothgänger, Christine Glockner and
608 Richard Haas and to Anne Katrin Prowse for language editing.

Author Contributions

611 PAN acquired funding, designed research, performed experiments, analysed data and wrote the
612 manuscript. MCW and SC performed experiments. CET, MRF and SNWH performed experiments and
613 analysed data. JP and ELD analysed data and helped writing the manuscript. HTS acquired funding and
614 conceptualized the idea. RR designed and supervised research, acquired funding and wrote the
615 manuscript. All authors read and approved the final version of the manuscript.

Conflict of interest disclosure

618 All authors declare no conflict of interest.

Data availability statement

Nucleotide sequence data was submitted to GenBank under accession numbers PV029542 (pL1) and PV068547(CTPc5). Moss lines and vectors were submitted to the International Moss Stock Centre (IMSC) under accession numbers 40963 (pL1 5A), 40964 (pL1 23A), 40965 (pL1 47B), 40966 (pL1Δ22 8A), 40967 (pL1Δ22 11B), 40968 (pL1Δ22 102B), 40969 (CTP-pL1 30), 40970 (CTP-pL1 103), 40971 (CTP-pL1 143), 1840 (pAct5_linker_eGFP_35ST_hpt), 2038 (pAct5_CTP_linker_eGFP_35ST_hpt), 1193 (pAct5_Citrine_nosT_hpt), 2249 (pAct5_pL1-Citrine_nosT_hpt), 2250 (pAct5_pL1Δ22-Citrine_nosT_hpt), 2251 (pAct5_CTP-pL1-Citrine_nosT_hpt), 1997 (pAct5_pL1_35ST_hpt), 1998 (pAct5_pL1Δ22_35ST_hpt) and 2048 (pAct5_CTP-pL1_35ST_hpt).

Funding statement

This work was supported by Maria-Sklodowska-Curie Actions Innovative Training Network under the Horizon 2020 programme under Grant Agreement No. 765115 (MossTech), by the German Research Foundation DFG under Germany's Excellence Strategy (CIBSS – EXC-2189 – Project ID 390939984), and the International Network Programme by the Danish Agency for Higher Education and Science (grant number 1113-00016B). The TEM (Hitachi HT7800) was funded by a DFG grant (project number 426849454) and is operated by the University of Freiburg, Faculty of Biology, as a partner unit within the Microscopy and Image Analysis Platform MIAP and the Life Imaging Center LIC, Freiburg. PAN gratefully acknowledges financial support by Studienstiftung des deutschen Volkes.

Short legends for Supporting Information

Supplementary Figure S1   **Vector maps**

Supplementary Figure S2   **Generation of Physcomitrella-optimized (p)L1**

Supplementary Figure S3   **Characterization of CTPs by analysis of subcellular eGFP localization**

Supplementary Figure S4   **Analysis of CDS integrity in L1-producing lines**

Supplementary Figure S5   **Comparison of biomass accumulation rates**

Supplementary Figure S6   **Ammonium sulphate precipitation**

Supplementary Figure S7   **Chromatogram of Cation Exchange purification**

Supplementary Table S1   **List of primers**

Supplementary Table S2   **Predicted chloroplast transit peptides for six selected proteins in**

651   **Physcomitrella**

References

Abel WO, Knebel W, Koop HU, Marienfeld JR, Quader H, Reski R, Schnepf E, Sporlein B (1989) A

cytokinin-sensitive mutant of the moss, *Physcomitrella patens*, defective in chloroplast division.

Protoplasma 152, 1-13. https://doi.org/10.1007/BF01354234

Adams A, Hendrikse M, Rybicki EP, Hitzeroth II (2023) Optimal size of DNA encapsidated by plant

produced human papillomavirus pseudovirions. *Virology* 580, 88 – 97.

https://doi.org/10.1016/j.virol.2023.02.003

Armenteros JJA, Salvatore M, Emanuelsson O, Wither O, von Heijne G, Elofsson A, Nielsen H (2019)

Detecting sequence signals in targeting peptides using deep learning. *Life Science Alliance* 2,

e201900429. https://doi.org/ 10.26508/lsa.201900429

Baker TS, Newcomb WW, Olson NH, Cowsert LM, Olson C, Brown JC (1991) Structures of bovine and

human papillomaviruses. Analysis by cryoelectron microscopy and three-dimensional image

reconstruction. *Biophysical Journal* 60, 1445–1456. https://doi.org/10.1016/S0006-3495(91)82181-6

Biemelt S, Sonnewald U, Galmbacher P, Willmitzer L, Müller M (2003) Production of human

papillomavirus type 16 virus-like particles in transgenic plants. *Journal of Virology* 77, 9211–9220.

https://doi.org/10.1128/JVI.77.17.9211-9220.2003

Bissa M, Zanotto C, Pacchioni S, Volonté L, Venuti A, Lembo D, De Giuli Morghen C, Radaelli A (2015) The

L1 protein of human papilloma virus 16 expressed by a fowlpox virus recombinant can assemble into

virus-like particles in mammalian cell lines but elicits a non-neutralising humoral response. *Antiviral*

Research 116, 67–75. https://doi.org/10.1016/j.antiviral.2015.01.012

Bohlender LL, Parsons J, Hoernstein SNW, Rempfer C, Ruiz-Molina N, Lorenz T, Rodríguez Jahnke F, Figl

R, Fode B, Altmann F, Reski R, Decker EL (2020) Stable protein sialylation in Physcomitrella. *Frontiers in*

Plant Science 11, 610032. https://doi.org/10.3389/FPLS.2020.610032

Bohlender LL, Parsons J, Hoernstein SNW, Bangert N, Rodriguez-Jahnke F, Reski R, Decker EL (2022)

Unexpected arabinosylation after humanization of plant protein N-glycosylation. *Frontiers in*

Bioengineering and Biotechnology 10, 838365. https://doi.org/10.3389/fbioe.2022.838365

Buck CB, Thompson CD, Pang YYS, Lowy DR, Schiller JT (2005) Maturation of papillomavirus capsids.

Journal of Virology 79, 2839–2846. https://doi.org/10.1128/JVI.79.5.2839-2846.2005

Buck CB, Day PM, Trus BL (2013) The papillomavirus major capsid protein L1. *Virology* 445, 169-174.

https://doi.org/ 10.1016/j.virol.2013.05.038

Büttner-Mainik A, Parsons J, Jérôme H, Hartmann A, Lamer S, Schaaf A, Schlosser A, Zipfel PF, Reski R,

Decker EL (2011) Production of biologically active recombinant human factor H in Physcomitrella. *Plant*

Biotechnology Journal 9, 373–383. https://doi.org/10.1111/J.1467-7652.2010.00552.X

Castells-Graells R, Lomonossoff GP (2021) Plant-based production can result in covalent cross-linking of

proteins. *Plant Biotechnology Journal* 19, 1095–1097. https://doi.org/10.1111/pbi.13598

Cerqueira C, Schiller JT (2017) Papillomavirus assembly: An overview and perspectives. *Virus Research*

231: 103–107. https://doi.org/10.1016/j.virusres.2016.11.010

Coates RJ, Young MT, Scofield S (2022) Optimising expression and extraction of recombinant proteins in

plants. *Frontiers in Plant Science* 13, 1074531. https://doi.org/10.3389/fpls.2022.1074531

Dai X, Zhao PX (2011) psRNATarget: a plant small RNA target analysis server. *Nucleic Acids Research* 39,

W155-W159. https://doi.org/10.1093/nar/gkr319

Dai X, Zhuang, Z, Zhao, PX (2018) psRNATarget: a plant small RNA target analysis server (2017 release).

Nucleic Acids Research 46, W49–W54. https://doi.org/10.1093/NAR/GKY316

de Martel C, Georges D, Bray F, Ferlay J, Clifford GM (2020) Global burden of cancer attributable to

infections in 2018: a worldwide incidence analysis. *The Lancet Global Health* 8, e180–e190.

https://doi.org/10.1016/S2214-109X(19)30488-7

Decker EL, Reski R (2020) Mosses in biotechnology. *Current Opinion in Biotechnology* 61, 21–27.

https://doi.org/10.1016/j.copbio.2019.09.021

Decker EL, Wiedemann G, Reski R (2015) Gene targeting for precision glyco-engineering: Production of

biopharmaceuticals devoid of plant-typical glycosylation in moss bioreactors. *Methods in Molecular*

Biology 1321, 213-224. https://doi.org/10.1007/978-1-4939-2760-9_15

Decker EL, Parsons J, Reski R (2014) Glyco-engineering for biopharmaceutical production in moss

bioreactors. *Frontiers in Plant Science* 5, 346. https://doi.org/10.3389/fpls.2014.00346

Fernández-San Millán A, Ortigosa SM, Hervás-Stubbs S, Corral-Martínez P, Seguí-Simarro JM, Gaétan J,

Coursaget P, Veramendi J (2008) Human papillomavirus L1 protein expressed in tobacco chloroplasts

self-assembles into virus-like particles that are highly immunogenic. *Plant Biotechnology Journal* 6, 427–

441. https://doi.org/10.1111/j.1467-7652.2008.00338.x

Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO (2009) Enzymatic assembly of DNA

molecules up to several hundred kilobases. *Nature Methods* 6, 343–345.

https://doi.org/10.1038/nmeth.1318

Gitzinger M, Parsons J, Reski R, Fussenegger M (2009) Functional cross-kingdom conservation of

mammalian and moss (*Physcomitrella patens*) transcription, translation and secretion machineries. *Plant*

Biotechnology Journal 7, 73–86. https://doi.org/10.1111/J.1467-7652.2008.00376.X

Gremillon L, Kiessling J, Hause B, Decker EL, Reski R, Sarnighausen E (2007) Filamentous temperature-

sensitive Z (FtsZ) isoforms specifically interact in the chloroplasts and in the cytosol of *Physcomitrella*

patens. *New Phytologist* 176, 299–310. https://doi.org/10.1111/j.1469-8137.2007.02169.x

Hennermann JB, Arash-Kaps L, Fekete G, Schaaf A, Busch A, Frischmuth T (2019) Pharmacokinetics,

pharmacodynamics, and safety of moss-aGalactosidase A in patients with Fabry disease. *Journal of*

Inherited Metabolic Disease 42, 527–533. https://doi.org/10.1002/JIMD.12052

Hitzeroth II, Chabeda A, Whitehead MP, Graf M, Rybicki EP (2018) Optimizing a human papillomavirus

type 16 L1-based chimaeric gene for expression in plants. *Frontiers in Bioengineering and Biotechnology*

6, 101. https://doi.org/10.3389/fbioe.2018.00101

Hoernstein SNW, Oezdemir B, van Gessel N, Miniera AA, Rogalla von Bieberstein B, Nilges L, Schweikert

Farinha J, Komoll R, Glauz S, Weckerle T, Scherzinger F, Rodriguez-Franco M, Mueller-Schuessele S, Reski

R (2023) A deeply conserved protease, acylamino acid-releasing enzyme (AARE), acts in ageing in

Physcomitrella and Arabidopsis. *Communications Biology* 6, 61. https://doi.org/10.1038/s42003-023-

04428-7

Hoernstein SNW, Schlosser A, Fiedler K, van Gessel N, Igloi GL, Lang D, Reski R (2024) A snapshot of the

Physcomitrella N-terminome reveals N-terminal methylation of organellar proteins. *Plant Cell Reports*

43, 250. https://doi.org/10.1007/s00299-024-03329-1

Hohe A, Reski R (2002) Optimisation of a bioreactor culture of the moss *Physcomitrella patens* for mass

production of protoplasts. *Plant Science* 163, 69-74. https://doi.org/10.1016/S0168-9452(02)00059-6

Hohe A, Decker EL, Gorr G, Schween G, Reski R (2002) Tight control of growth and cell differentiation in

photoautotrophically growing moss (*Physcomitrella patens*) bioreactor cultures. *Plant Cell Reports* 20,

1135-1140. https://doi.org/10.1007/s00299-002-0463-y

Hohe A, Egener T, Lucht JM, Holtorf H, Reinhard C, Schween G, Reski R (2004) An improved and highly

standardized transformation procedure allows efficient production of single and multiple targeted gene-

knockouts in a moss, *Physcomitrella patens*. *Current Genetics* 44, 339-347.

https://doi.org/10.1007/s00294-003-0458-4

Jones-Rhoades MW, Bartel DP, Bartel B (2006) MicroRNAs and their regulatory roles in plants. *Annual*

Review of Plant Biology 57, 19-53. https://doi.org/10.1146/annurev.arplant.57.032905.105218

Karan S, Affonso De Oliveira JF, Moreno-Gonzalez MA, Steinmetz NF (2024) A self-amplifying human

papillomavirus 16 vaccine candidate delivered by tobacco mosaic virus-like particle. *ACS Applied Bio*

Materials 7, 7675 – 7683. https://doi.org/10.1021/acsabm.4c01239

Khraiwesh B, Arif MA, Seumel GI, Ossowski S, Weigel D, Reski R, Frank W (2010) Transcriptional control

of gene expression by microRNAs. *Cell* 140, 111-122. https://doi.org/ 10.1016/j.cell.2009.12.023

Kim SN, Jeong HS, Park SN, Kim HJ (2007) Purification and immunogenicity study of human

papillomavirus type 16 L1 protein in *Saccharomyces cerevisiae*. *Journal of Virological Methods* 139, 24-

30. https://doi.org/ 10.1016/j.jviromet.2006.09.004

Kim HJ, Kim SY, Lim SJ, Kim JY, Lee SJ, Kim HJ (2010) One-step chromatographic purification of human

papillomavirus type 16 L1 protein from *Saccharomyces cerevisiae*. *Protein Expression and Purification* 70,

68–74. https://doi.org/10.1016/j.pep.2009.08.005

Kim HJ, Cho SY, Park MH, Kim HJ (2018) Comparison of the size distributions and immunogenicity of

human papillomavirus type 16 L1 virus-like particles produced in insect and yeast cells. *Archives of*

Pharmacal Research 41, 544–553. https://doi.org/10.1007/s12272-018-1024-4

Kirnbauer R, Booy F, Cheng N, Lowy DR, Schiller JT (1992) Papillomavirus L1 major capsid protein self-

assembles into virus-like particles that are highly immunogenic. *Proceedings of the National Academy of*

Sciences USA 89, 12180–12184. https://doi.org/10.1073/pnas.89.24.12180

Knappe M, Bodevin S, Selinka HC, Spillmann D, Streeck RE, Chen XS, Lindahl U, Sapp M (2007) Surface-

exposed amino acid residues of HPV16 L1 protein mediating interaction with cell surface heparan

sulfate. *Journal of Biological Chemistry* 282, 27913–27922. https://doi.org/10.1074/jbc.M705127200

Koprivova A, Stemmer C, Altmann F, Hoffmann A, Kopriva S, Gorr G, Reski R, Decker EL (2004) Targeted

knockouts of *Physcomitrella* lacking plant-specific immunogenic N-glycans. *Plant Biotechnology Journal*

2, 517–523. https://doi.org/10.1111/J.1467-7652.2004.00100.X

Lamprecht RL, Kennedy P, Huddy SM, Bethke S, Hendrikse M, Hitzeroth II, Rybicki EP (2016) Production

of human papillomavirus pseudovirions in plants and their use in pseudovirion-based neutralisation

assays in mammalian cells. *Scientific Reports* 6, 20431. https://doi.org/10.1038/srep20431

Leder C, Kleinschmidt JA, Wiethe C, Müller M (2001) Enhancement of capsid gene expression: Preparing

the human papillomavirus type 16 major structural gene L1 for DNA vaccination purposes. *Journal of*

Virology 75, 9201–9209. https://doi.org/10.1128/JVI.75.19.9201-9209.2001

Lenzi P, Scotti N, Alagna F, Tornesello ML, Pompa A, Vitale A, De Stradis A, Monti L, Grillo S, Buonaguro

FM, Maliga P, Cardi T (2008) Translational fusion of chloroplast-expressed human papillomavirus type 16

L1 capsid protein enhances antigen accumulation in transplastomic tobacco. *Transgenic Research* 17,

1091–1102. https://doi.org/10.1007/s11248-008-9186-3

Ljubojevic S, Skerlev M (2014) HPV-associated diseases. *Clinics in Dermatology* 32, 227-234.

https://doi.org/10.1016/j.clindermatol.2013.08.007

Mach H, Volkin DB, Troutman RD, Wang B, Luo Z, Jansen KU, Shi L (2006) Disassembly and reassembly of

yeast-derived recombinant human papillomavirus virus-like particles (HPV VLPs). *Journal of*

Pharmaceutical Sciences 95, 2195–2206. https://doi.org/10.1002/jps.20696

Maclean J, Koekemoer M, Olivier AJ, Stewart D, Hitzeroth II, Rademacher T, Fischer R, Williamson AL,

Rybicki EP (2007) Optimization of human papillomavirus type 16 (HPV-16) L1 expression in plants:

comparison of the suitability of different HPV-16 L1 gene variants and different cell-compartment

localization. *Journal of General Virology* 88, 1460–1469. https://doi.org/10.1099/vir.0.82718-0

Marsian J, Lomonossoff GP (2016) Molecular pharming – VLPs made in plants. *Current Opinion in*

Biotechnology 37, 201-206. https://doi.org/10.1016/j.copbio.2015.12.007

McCarthy MP, White WI, Palmer-Hill F, Koenig S, Suzich JA (1998) Quantitative disassembly and

reassembly of human papillomavirus type 11 virus like particles *in vitro*. *Journal of Virology* 72, 32–41.

https://doi.org/10.1128/JVI.72.1.32-41.1998

Michelfelder S, Parsons J, Bohlender LL, Hoernstein SNW, Niederkrüger H, Busch A, Krieghoff N, Koch J,

Fode B, Schaaf A, Frischmuth T, Pohl M, Zipfel PF, Reski R, Decker EL, Häffner K (2017) Moss-produced,

glycosylation-optimized human factor H for therapeutic application in complement disorders. *Journal of*

the American Society of Nephrology 28, 1462–1474. https://doi.org/10.1681/ASN.2015070745

Michelfelder S, Fischer F, Wäldin A, Hörle KV, Pohl M, Parsons J, Reski R, Decker EL, Zipfel PF, Skerka C,

Häffner K (2018) The MFHR1 fusion protein is a novel synthetic multitarget complement inhibitor with

therapeutic potential. *Journal of the American Society of Nephrology* 29, 1141–1153.

https://doi.org/10.1681/ASN.2017070738

Mori S, Ozaki S, Yasugi T, Yoshikawa H, Taketani Y, Kanda T (2006) Inhibitory cis-element-mediated

decay of human papillomavirus type 16 L1-transcript in undifferentiated cells. *Molecular and Cellular*

Biochemistry 288, 47–57. https://doi.org/10.1007/s11010-006-9117-7

Murén E, Nilsson A, Ulfstedt M, Johansson M, Ronne H (2009) Rescue and characterization of episomally

replicating DNA from the moss *Physcomitrella*. *Proceedings of the National Academy of Sciences USA*

106, 19444–19449

Muthamilselvan T, Khan MRI, Hwang I (2023) Assembly of human papillomavirus 16 L1 protein in

Nicotiana benthamiana chloroplasts into highly immunogenic Virus-Like Particles. *Journal of Plant*

Biology 66, 331–340. https://doi.org/10.1007/s12374-023-09393-6

Naupu PN, van Zyl AR, Rybicki EP, Hitzeroth II (2020) Immunogenicity of plant-produced human

papillomavirus (HPV) Virus-Like Particles (VLPs). *Vaccines* 8, 740.

https://doi.org/10.3390/VACCINES8040740

Nooraei S, Bahrulolum H, Hoseini ZS, Katalani C, Hajizade A, Easton AJ, Ahmadian G (2021) Virus-like

particles: preparation, immunogenicity and their roles as nanovaccines and drug nanocarriers. *Journal of*

Nanobiotechnology 19, 59. https://doi.org/10.1186/s12951-021-00806-7

Orellana-Escobedo L, Rosales-Mendoza S, Romero-Maldona A, Parsons J, Decker EL, Monreal-Escalante

E, Moreno-Fierros L, Reski R (2015) An Env-derived multi-epitope HIV chimeric protein produced in the

moss *Physcomitrella patens* is immunogenic in mice. *Plant Cell Reports* 34, 425-433.

https://doi.org/10.1007/s00299-014-1720-6

Park MA, Kim HJ, Kim HJ (2008) Optimum conditions for production and purification of human

papillomavirus type 16 L1 protein from *Saccharomyces cerevisiae*. *Protein Expression and Purification* 59,

175–181. https://doi.org/10.1016/j.pep.2008.01.021

Parsons J, Altmann F, Arrenberg CK, Koprivova A, Beike AK, Stemmer C, Gorr G, Reski R, Decker EL (2012)

Moss-based production of asialo-erythropoietin devoid of Lewis A and other plant-typical carbohydrate

determinants. *Plant Biotechnology Journal* 10, 851–861. https://doi.org/10.1111/J.1467-

7652.2012.00704.X

Pineo CB, Hitzeroth II, Rybicki EP (2013) Immunogenic assessment of plant-produced human

papillomavirus type 16 L1/L2 chimaeras. *Plant Biotechnology Journal* 11, 964–975.

https://doi.org/10.1111/pbi.12089

Rempfer C, Wiedemann G, Schween G, Kerres KL, Lucht JM, Horres R, Decker EL, Reski R (2022)

Autopolyploidization affects transcript patterns and gene targeting frequencies in Physcomitrella. *Plant

Cell Reports* 41, 153-173. https://doi.org/10.1007/s00299-021-02794-2

Reski R (1998) Development, genetics and molecular biology of mosses. *Botanica Acta* 111, 1–15.

https://doi.org/10.1111/j.1438-8677.1998.tb00670.x

Reski R (2018) Quantitative moss cell biology. *Current Opinion in Plant Biology* 46, 39–47.

https://doi.org/10.1016/J.PBI.2018.07.005

Reski R, Abel WO (1985) Induction of budding on chloronemata and caulonemata of the moss,

Physcomitrella patens, using isopentenyladenine. *Planta* 165, 354-358.

https://doi.org/1007/BF00392232

Reski R, Parsons J, Decker EL (2015) Moss-made pharmaceuticals: from bench to bedside. *Plant

Biotechnology Journal* 13, 1191–1198. https://doi.org/10.1111/pbi.12401

Reski R, Bae H, Simonsen HT (2018) *Physcomitrella patens*, a versatile synthetic biology chassis. *Plant

Cell Reports* 37, 1409–1417. https://doi.org/10.1007/S00299-018-2293-6

Roden RBS, Stern PL (2018) Opportunities and challenges for human papillomavirus vaccination in

cancer. *Nature Reviews Cancer* 18, 240-254. https://doi.org/10.1038/nrc.2018.13

Rommel O, Dillner J, Fligge C, Bergsdorf C, Wang X, Selinka HC, Sapp M (2005) Heparan sulfate

proteoglycans interact exclusively with conformationally intact HPV L1 assemblies: Basis for a virus-like

particle ELISA. *Journal of Medical Virology* 75, 114–121. https://doi.org/10.1002/jmv.20245

Rosales-Mendoza S, Orellana-Escobedo L, Romero-Maldonado A, Decker EL, Reski R (2014) The potential

of *Physcomitrella patens* as a platform for the production of plant-based vaccines. *Expert Reviews of*

Vaccines 13, 203-212. https://doi.org/10.1586/14760584.2014.872987

Rosales-Mendoza S, Salazar-Gonzalez JA, Decker EL, Reski R (2016) Implications of plant glycans in the

development of innovative vaccines. *Expert Reviews of Vaccines* 15, 915-925.

https://doi.org/10.1586/14760584.2016.1155987

Ruiz-Molina N, Parsons J, Müller M, Hoernstein SNW, Bohlender LL, Pumple S, Zipfel PF, Häffner K, Reski

R, Decker EL (2022a) A synthetic protein as efficient multitarget regulator against complement over-

activation. *Communications Biology* 5, 152. https://doi.org/10.1038/s42003-022-03094-5

Ruiz-Molina N, Parsons J, Schroeder S, Posten C, Reski R, Decker EL (2022b) Process engineering of

biopharmaceutical production in moss bioreactors *via* model-based description and evaluation of

phytohormone impact. *Frontiers in Bioengineering and Biotechnology* 10, 837965.

https://doi.org/10.3389/fbioe.2022.837965

Rybicki EP (2020) Plant molecular farming of virus-like nanoparticles as vaccines and reagents. *WIREs*

Nanomedicine and Nanobiotechnology 12, e1587. https://doi.org/10.1002/wnan.1587

Sapp M, Fligge C, Petzak I, Robin Harris J, Streeck RE (1998) Papillomavirus assembly requires

trimerization of the major capsid protein by disulfides between two highly conserved cysteines. *Journal*

of Virology 72, 6186 – 6189. https://doi.org/10.1128/jvi.72.7.6186-6189.1998

Schween G, Hohe A, Koprivova A, Reski R (2003) Effects of nutrients, cell density and culture techniques

on protoplast regeneration and early protonema development in a moss, *Physcomitrella patens*. *Journal*

of Plant Physiology 160, 209-212. https://doi.org/10.1078/0176-1617-00855

Scotti N, Rybicki EP (2013) Virus-like particles produced in plants as potential vaccines. *Expert Review of*

Vaccines 12, 211–224. https://doi.org/10.1586/erv.12.147

Shen JS, Busch A, Day TS, Meng XL, Yu CI, Dabrowska-Schlepp P, Fode B, Niederkrüger H, Forni S, Chen S,

Schiffmann R, Frischmuth T, Schaaf A (2016) Mannose receptor-mediated delivery of moss-made α-

galactosidase A efficiently corrects enzyme deficiency in Fabry mice. *Journal of Inherited Metabolic*

Disease 39, 293–303. https://doi.org/10.1007/S10545-015-9886-9

Shi L, Sanyal G, Ni A, Luo Z, Doshna S, Wang B, Graham TL, Wang N, Volkin DB (2005) Stabilization of

human papillomavirus virus-like particles by non-ionic surfactants. *Journal of Pharmaceutical Sciences*

94, 1538–1551. https://doi.org/10.1002/jps.20377

Shi L, Sings HL, Bryan JT, Wang B, Wang Y, Mach H, Kosinski M, Washabaugh MW, Sitrin R, Barr E (2007)

GARDASIL®: Prophylactic human papillomavirus vaccine development – From bench top to bed-side.

Clinical Pharmacology & Therapeutics 81, 259–264. https://doi.org/10.1038/SJ.CLPT.6100055

Šmídková M, Müller M, Thönes N, Piuko K, Angelisová P, Velemínský J, Angelis KJ (2010) Transient

expression of human papillomavirus type 16 virus-like particles in tobacco and tomato using a tobacco

rattle virus expression vector. *Biologia Plantarum* 54, 451–460. https://doi.org/10.1007/s10535-010-

0081-4

Studentsov YY, Schiffman M, Strickler HD, Ho GYF, Pang YYs, Schiller J, Herrero R, Burk RD (2002)

Enhanced enzyme-linked immunosorbent assay for detection of antibodies to virus-like particles of

human papillomavirus. *Journal of Clinical Microbiology* 40, 1755-1760.

https://doi.org/10.1128/JCM.40.5.1755-1760.2002

Strepp R, Scholz S, Kruse S, Speth V, Reski R (1998) Plant nuclear gene knockout reveals a role in plastid

division for the homolog of the bacterial cell division protein FtsZ, an ancestral tubulin. *Proceedings of*

the National Academy of Sciences USA 95, 4368–4373. https://doi.org/10.1073/PNAS.95.8.4368

Top O, Parsons J, Bohlender LL, Michelfelder S, Kopp P, Busch-Steenberg C, Hoernstein SNW, Zipfel PF,

Häffner K, Reski R, Decker EL (2019) Recombinant production of MFHR1, a novel synthetic multitarget

complement inhibitor, in moss bioreactors. *Frontiers in Plant Science* 10, 260.

https://doi.org/10.3389/FPLS.2019.00260

Top O, Milferstaedt SWL, van Gessel N, Hoernstein SNW, Özdemir B, Decker EL, Reski R (2021)

Expression of a human cDNA in moss results in spliced mRNAs and fragmentary protein isoforms.

Communications Biology 4, 964. https://doi.org/10.1038/s42003-021-02486-3

U.S. Food and Drug Administration (2018) FDA approves expanded use of Gardasil 9 to include

individuals 27 through 45 years old | FDA. https://www.fda.gov/news-events/press-

announcements/fda-approves-expanded-use-gardasil-9-include-individuals-27-through-45-years-old

van Wijk KJ (2024) Intra-chloroplast proteases: A holistic network view of chloroplast proteolysis. *Plant*

Cell 36, 3116–3130. https://doi.org/10.1093/plcell/koae178

Varsani A, Williamson AL, Rose RC, Jaffer M, Rybicki EP (2003) Expression of Human papillomavirus type

16 major capsid protein in transgenic *Nicotiana tabacum* cv. Xanthi. *Archives of Virology* 148, 1771–

1786. https://doi.org/10.1007/s00705-003-0119-4

Weise A, Rodriguez-Franco M, Timm B, Hermann M, Link S, Jost W, Gorr G (2006) Use of *Physcomitrella*

patens actin 5' regions for high transgene expression: importance of 5' introns. *Applied Microbiology*

and Biotechnology 70: 337–345. https://doi.org/10.1007/s00253-005-0087-6

Wiedemann G, van Gessel N, Köchl F, Hunn L, Schulze K, Maloukh L, Nogué F, Decker EL, Hartung F, Reski

R (2018) RecQ helicases function in development, DNA repair, and gene targeting in *Physcomitrella*

patens. *Plant Cell* 30, 717–736. https://doi.org/10.1105/tpc.17.00632

World Health Organisation (2024a) Human papillomavirus and cancer. https://www.who.int/news-

room/fact-sheets/detail/human-papilloma-virus-and-cancer, accessed 16/12/2024

1001

1002   World Health Organization (2024b) Considerations for human papillomavirus (HPV) vaccine product

1003   choice (Issue 2). https://iris.who.int/bitstream/handle/10665/376546/9789240089167-eng.pdf

1004

1005   Wu M, Huang H, Tang Y, Ren X, Jiang X, Tian M, Li W (2024) Unveiling the multifaceted realm of human

1006   papillomavirus: a comprehensive exploration of biology, interactions, and advances in cancer

1007   management. *Frontiers in Immunology* 15, 1430544. https://doi.org/10.3389/fimmu.2024.1430544

1008

1009   Zahin M, Joh J, Khanal S, Husk A, Mason H, Warzecha H, Ghim SJ, Miller DM, Matoba N, Jenson AB

1010   (2016) Scalable production of HPV16 L1 protein and VLPs from tobacco leaves. *PLOS ONE* 11, e 0160995.

1011   https://doi.org/10.1371/JOURNAL.PONE.0160995

1012

1013   Zhao X, Schwartz S (2008) Inhibition of HPV-16 L1 expression from L1 cDNAs correlates with the

1014   presence of hnRNP A1 binding sites in the L1 coding region. *Virus Genes* 36, 45–53.

1015   https://doi.org/10.1007/s11262-007-0174-0

1016

1017   zur Hausen H (2002) Papillomaviruses and cancer: From basic studies to clinical application. *Nature*

1018   *Reviews Cancer* 2, 342-350. https://doi.org/10.1038/nrc798

1019

1020   zur Hausen H, de Villiers EM, Gissmann L (1981) Papillomavirus infections and human genital cancer.

1021   *Gynecologic Oncology* 12, S124–S128. https://doi.org/10.1016/0090-8258(81)90067-6

1022 # Supplementary Information

1023 ## Supplementary Figure S1 – Vector maps

1024

List of all vectors used in this study. Vectors **a** & **b** were used for the characterization of chloroplast transit peptides (CTPs).

Vector **a** served as a control and was used as a cloning template for inserting different CTP candidates. Vectors **c**, **d**, **e** & **f** were

used for the analysis of L1 subcellular targeting. Vector **c** served as a control and was used as a cloning template for inserting

either pL1, pL1Δ22 or CTP-pL1 CDS (**d**, **e** & **f**, respectively). Vectors **g**, **h** & **i** were used for generation of stable transgenic

Physcomitrella lines. The vectors contain either the full length pL1 including the native nuclear localization signal (NLS) (**g**), a

truncated version lacking the NLS (**h**), or the full-length version including the native NLS and an N-terminal chloroplast transit

peptide (**i**). All vectors contain the PpActin5 promoter (yellow), the CaMV 35S terminator (grey), and a hygromycin

(hpt)(turquoise) resistance cassette for selection. Other elements are the GGGGGA linker (red), CTP (orange), NLS (blue) and eGFP

or Citrine (green). Genetic elements are not depicted true to scale. Vector sizes: 7260 bp (**a**), 7461 bp (**b**), 7299 bp (**c**), 8802 bp

(**d**), 8736 bp (**e**), 9021 bp (**f**), 7998 bp (**g**), 7932 bp (**h**), 8217 bp (**i**).

Supplementary Figure S2 - Generation of Physcomitrella-optimized (p)L1

HPV-16 hL1

ATGTCCCTGTGGCTGCCCAGCGAGGCCACCGTGTACCTGCCCCCCGTGCCCGTGAGCAAGGTGGTGAGCACCGATGAGTACGTGGCCCGGA

CCAACATCTACTACCACGCCGGCACCTCCAGACTGCTGGCCGTGGGCCACCCCTACTTCCCCATCAAGAAGCCCAACAACAACAAGATCCTGG

TGCCCAAGGTGAGCGGCCTGCAATACCGGGTGTTCAGAATCCACCTGCCCGACCCCAATAAGTTCGGCTTCCCCGACACCAGCTTCTACAACC

CCGACACCCAGAGACTGGTGTGGGCCTGCGTGGGCGTGGAGGTGGGCAGAGGCCAGCCTCTGGGCGTGGGCATCAGCGGCCACCCTCTGC

TGAACAAGCTGGACGACACCGAGAACGCCAGCGCCTACGCCGCCAACGCCGGCGTGGATAACAGAGAATGCATCAGCATGGACTACAAGCA

GACCCAGCTGTGCCTCATCGGCTGCAAGCCCCCCATCGGCGAGCACTGGGGCAAGGGCAGCCCCTGCACCAACGTGGCCGTGAATCCTGGC

GACTGTCCTCCCCTGGAACTCATCAACACCGTGATCCAGGACGGCGACATGGTGGACACCGGCTTCGGCGCCATGGACTTCACCACCCTCCAG

GCCAATAAGAGCGAGGTGCCCCTGGACATCTGCACCAGCATCTGCAAGTACCCCGACTACATCAAGATGGTGAGCGAGCCCTACGGCGATAG

CCTGTTCTTCTACCTGCGGCGGGAGCAGATGTTCGTGCGGCACCTGTTCAACAGAGCCGGCGCCGTGGGCGAGAACGTGCCCGACGACCTGT

ACATCAAGGGCAGCGGCAGCACCGCCAACCTGGCCAGCAGCAACTACTTCCCTACCCCCAGCGGCTCCATGGTGACCAGCGACGCCCAGATC

TTCAACAAGCCCTACTGGCTCCAGAGAGCCCAGGGCCACAACAATGGCATCTGCTGGGGCAACCAGCTGTTCGTGACCGTGGTGGATACCAC

CCGGAGCACCAACATGTCCCTGTGCGCCGCCATCAGCACCAGCGAGACCACCTACAAGAACACCAACTTCAAGGAGTACCTGAGGCACGGCG

AGGAGTACGACCTCCAGTTCATCTTCCAGCTGTGCAAGATCACCCTCACCGCCGACGTGATGACCTACATCCACAGCATGAACAGCACCATCC

TGGAGGACTGGAACTTCGGCCTGCAGCCCCCTCCTGGCGGCACCCTGGAGGACACCTACAGATTCGTGACCAGCCAGGCCATCGCATGCCAG

AAGCACACCCCTCCCGCCCCTAAGGAGGACCCCCTGAAGAAGTACACCTTCTGGGAGGTGAACCTGAAGGAGAAGTTCAGCGCCGACCTGG

ACCAGTTCCCTCTGGGCAGAAAGTTCCTGCTGCAAGCCGGCCTGAAGGCCAAGCCTAAGTTCACCCTGGGCAAGAGAAAGGCCACCCCCACC

ACAAGCAGCACCAGCACCACCGCCAAGCGGAAGAAGCGCAAGCTGTGA

HPV-16 pL1

ATGTCCCTGTGGCTGCCCAGCGAGGCCACCGTGTACCTGCCCCCCGTGCCCGTGAGCAAAGTGGTGAGCACCGATGAGTACGTGGCCCGGAC

CAACATCTACTACCACGCCGGCACCTCCAGACTGCTGGCCGTGGGCCACCCCTACTTCCCCATCAAGAAGCCCAACAACAACAAGATCCTGGT

GCCCAAAGTGAGCGGCCTGCAGTACCGGGTGTTCAGAATCCACCTGCCCGACCCCAACAAGTTCGGCTTCCCCGACACCAGCTTCTACAACCC

CGACACCCAGAGACTGGTGTGGGCCTGCGTGGGCGTGGAAGTGGGCAGAGGCCAGCCTCTGGGCGTGGGCATCAGCGGCCACCCTCTGCT

GAACAAGCTGGACGACACCGAGAACGCCAGCGCCTACGCCGCCAACGCCGGCGTGGATAACAGAGAGTGCATCAGCATGGACTACAAGCA

GACCCAGCTGTGCCTCATCGGCTGCAAGCCCCCCATCGGCGAGCACTGGGGCAAGGGCAGCCCCTGCACCAACGTGGCCGTGAACCCTGGC

GACTGCCCTCCCCTGGAGCTCATCAACACCGTGATCCAGGACGGCGACATGGTGGACACCGGCTTCGGCGCCATGGACTTCACCACCCTCCA

GGCCAACAAGAGCGAAGTGCCCCTGGACATCTGCACCAGCATCTGCAAGTACCCCGACTACATCAAGATGGTGAGCGAGCCCTACGGCGATA

GCCTGTTCTTCTACCTGCGGCGCGAGCAAATGTTTGTGCGGCACCTGTTCAACAGAGCCGGCGCCGTGGGCGAGAACGTGCCCGACGACCTG

TACATCAAGGGCAGCGGCAGCACCGCCAACCTGGCCAGCAGCAACTACTTCCCTACCCCCAGCGGCTCCATGGTGACCAGCGACGCCCAGAT

CTTCAACAAGCCCTACTGGCTCCAGAGAGCCCAGGGCCACAACAACGGCATCTGCTGGGGCAACCAGCTGTTCGTGACCGTGGTGGATACCA

CCCGGAGCACCAACATGTCCCTGTGCGCCGCCATCAGCACCAGCGAGACCACCTACAAGAACACCAACTTCAAGGAGTACCTGAGGCACGGC

GAGGAGTACGACCTCCAGTTCATCTTCCAGCTGTGCAAGATCACCCTCACCGCCGACGTGATGACCTACATCCACAGCATGAACAGCACCATC

CTGGAGGACTGGAACTTCGGCCTGCAGCCCCCTCCTGGCGGCACCCTGGAGGACACCTACAGATTCGTGACCAGCCAGGCCATCGCATGCCA

GAAGCACACCCCTCCCGCCCCTAAGGAGGACCCCCTGAAGAAGTACACCTTCTGGGAAGTGAACCTGAAGGAGAAGTTCAGCGCCGACCTG

GACCAGTTCCCTCTGGGCAGAAAGTTCCTGCTGCAGGCCGGCCTGAAGGCCAAGCCTAAGTTCACCCTGGGCAAGAGAAAGGCCACCCCCAC

CACAAGCAGCACCAGCACCACCGCCAAGCGGAAGAAGCGCAAGCTGTGA

Human codon-optimized HPV-16 (h)L1 CDS and Physcomitrella-optimized HPV-16 (p)L1 CDS. Grey marked nucleotides were

changed in HPV-16 pL1 through codon usage optimization by the tool physCO (Top et al., 2021). Turquoise marked motifs refer

to the Physcomitrella alternative splicing recognition sites. The green marked segment refers to a predicted miRNA binding site

identified by the tool psRNATarget (Dai & Zhao, 2011; Dai et al., 2018). Light orange coloured segments represent the nuclear

localization signal. Nucleotide changes are underlined in the HPV-16 pL1 sequence. (hL1, GenBank accession no. DQ067889,

(Maclean et al., 2007))

 Supplementary Figure S3 - Characterization of CTPs by analysis of subcellular eGFP

1080 localization

1081

1082

1083
1084

Microscopy pictures of Physcomitrella protoplasts transformed with CTP-eGFP constructs. Samples were analysed using a

rhodamine and a FITC filter to identify chloroplast (magenta) and eGFP localization, respectively, as well as a merge of the two

signals. For sample CTPc21-eGFP no eGFP signal could be observed in chloroplasts. For samples CTPc5-eGFP, CTPc10-eGFP,

CTPc22-eGFP and CTPc22noMet-eGFP green fluorescence signal is observed in chloroplasts. eGFP lacking a CTP localized to the

cytoplasm and served as a negative control. The experiment was repeated two times. CTPc5 was examined a third time using

confocal microscopy. Scale bars represent 10 µm.

Supplementary Figure S4 – Analysis of CDS integrity in L1-producing lines

1092

1093
1094

Analysis of rearrangements in the pL1 CDS using RT PCR. Lines pL1 23A, pL1Δ22 8A and CTP-pL1 30 were checked for DNA or

RNA rearrangements by performing PCR on cDNA using four different primer pairs. Primer pairs were selected to amplify the

entire cDNA. The forward primer binds at the start codon and is the same in each primer pair (pL1_fwd_1 for lines pL1 23A &

pL1Δ22 8A, CTP_pL1_fwd_1 for line CTP-pL1 30). The reverse primers are different in each pair and bind at the 3' end of the CDS

or within the terminator (**Supplementary Table S1**). For line pL1 23A usage of 25 ng or 200 ng cDNA in PCR reactions was

compared. Following experiments used 200 ng. Absence of distinct bands of a size smaller than the expected size is indicative for

no larger rearrangements on both DNA and RNA level. No amplicon is visible for lines pL1 23A using Primer Pair 2 and CTP-pL1 30

using Primer Pair 3. The constructs used for transformation of Physcomitrella protoplasts (**Supplementary Figure S1 g - i**) served

as a positive control.

 Supplementary Figure S5 – Comparison of biomass accumulation rates

1106

Analysis of biomass accumulation rates among nine Physcomitrella lines of three different constructs. Biomass accumulation
1107

rates were observed over a course of 4 weeks for lines pL1 5A, pL1 23A, pL1 47B (blue); pL1Δ22 8A, pL1Δ22 11B, pL1Δ22 102B

(green) and CTP-pL1 30, CTP-pL1 103, CTP-pL1 143 (purple). The data was plotted using linear regression analysis and all slopes

were analysed using ANOVA and post hoc multiple comparisons test (TukeyHSD). Comparison of the slopes between the

constructs of pL1, pL1Δ22 and CTP-pL1 showed pL1 lines accumulated biomass faster than pL1Δ22 (p = <0.0001) and CTP-pL1 (p

= 0.0001) lines. Per week n = 3 biological samples were taken.

Supplementary Figure S6 – Ammonium sulphate precipitation

1114

1115

1116

Ammonium sulphate precipitation of L1 from moss extracts. Samples from the supernatant and the resuspended and dialysed

pellet before and after filtration were analysed for the presence of L1 content and residual host proteins under reducing

conditions *via* Western blot and Coomassie-stained gel, respectively. CTP-pL1 30 extracts were precipitated with either 40%, 45%,

or 50% ammonium sulphate (AMS). After centrifugation, the supernatant and the precipitate were taken up and dialysed against

cation exchange chromatography binding buffer to remove excess AMS. Non-dissolved protein was cleared from the sample using

0.22 μm filters. 45% AMS was found to be sufficient for full precipitation of L1 from crude extracts. 30 ng *S. cerevisiae*-produced

L1 served as a positive control and Physcomitrella wildtype as a negative control.

Supplementary Figure S7 – Chromatogram of Cation Exchange purification

1125

1126

1127   **Chromatogram of HPV-16 L1 purification from moss using cation exchange chromatography.** The chromatogram

1128   shows the UV absorbance, indicative for protein concentration, in blue, gradient concentration in pink, and conductivity,

1129   indicative for salt concentration, in red. The y axis depicts milli-Absorbance Units and the x axis depicts the volume in mL. The

1130   large peak between 16 – 26 mL shows the sample application or flow through, the following small peak at 28 mL the wash fraction

1131   and the larger peak at 33 – 35 mL the elution fractions.

 ## Supplementary Table S1 – List of primers

1133 Capital letters refer to the part of the primer binding to the template whereas lower case letters represent overhangs added to
1134 the template via PCR. Abbreviations: fwd or fw (forward), rev or rv (reverse), CTP (chloroplast transit peptide), MMEJ (micro
1135 homology-mediated end joining), GFP (green fluorescent protein), OH (overhang (lower case letters)).

Primer name	Sequence (5'→3')	Application
AA2_3integFP (hpt3'seq)	ACTCGCCGATAGTGGAAACC	Amp of linker-eGFP vector part 1
PAN_PpActin-Pro_Rv	CTGCACAACAACCACCAATATTTAATTTC	Amp of linker-eGFP vector part 1
linker:EGFP_p1840_Fw	GGAGGTGGAGGTGGAGC	Amp of linker-eGFP vector part 2
HPTLUCfwd2	TTGCCCTCGGACGAGTGC	Amp of linker-eGFP vector part 2
CTP_Pp3c10_Fw	aattaaatattggtggttgttgtgcagATGGCGTTATTGGCGCGC	Amp of CTPc10
CTP_Pp3c10_Rv	gctccacctccacctccATTGGTGACCCTCAAAGGTCCAG	Amp of CTPc10
CTP_Pp3c5_Fw	aattaaatattggtggttgttgtgcagATGGCTACGATGAAGATTGCTGTGG	Amp of CTPc5
CTP_Pp3c5_Rv	gctccacctccacctccAGCTCGCACGGAGCTGTTC	Amp of CTPc5
CTP_Pp3c22_Fw	aattaaatattggtggttgttgtgcagATGCGGGCTCTCTGTGGAG	Amp of CTPc22 & CTPc22_noMet
CTP_Pp3c22_Rv	gctccacctccacctccCATAGCGACAATGCACCGTCTTC	Amp of CTPc22
CTP_Pp3c22_noMet_Rv	gctccacctccacctccAGCGACAATGCACCGTCTTCC	Amp of CTPc22_noMet
CTP_Pp3c21_Fw	aattaaatattggtggttgttgtgcagATGGCCGCCGTCGGAATG	Amp of CTPc21
CTP_Pp3c21_Rv	gctccacctccacctccGGCACGGACAGCGAACGC	Amp of CTPc21
Act5_seq	TCCACATGGCTACAGCTG	Colony PCR CTP-eGFP constructs
EGFP_NoLinker_Rev	CTTGTACAGCTCGTCCATG	Colony PCR CTP-eGFP constructs
XhoI-pL1-BamHI_fw	gtgcagctcgagATGTCCCTGTGGCTGCC	Amp of pL1 and pL1Δ22
XhoI-pL1-BamHI _Rv	tctagaggatccTCACAGCTTGCGCTTCTTCC	Amp of pL1
XhoI-pL1delta22-BamHI _Rv	tctagaggatcctcaGCCCAGGGTGAACTTAGGC	Amp of pL1Δ22
pL1_w/oATG_Fw	TCCCTGTGGCTGCCCAG	Amp of CTP-pL1 (no Start codon)
pL1_Rv_35St-OH	GACTGGTGATTTTTGCGGACTCTAG	Amp of pL1, Heterosplicing and MMEJ check of pL1 lines

CTPc5_Fw_Act5p-OH	tattggtggttgttgtgcagctcgagATGGCTACGATGAAGATTGCTGTG	Amp of CTPc5
CTPc5_Rv_pL1-OH	ctcgctgggcagccacagggaAGCTCGCACGGAGCTGTTC	Amp of CTPc5
Seq_hL1+HSP_Fw	CGTGATGACCTACATCCACAG	Colony PCR pL1 constructs
pL1_Rv_35St-OH	GACTGGTGATTTTTGCGGACTCTAG	Colony PCR pL1 constructs
pL1_fwd_1	ATGTCCCTGTGGCTGCCC	Heterosplicing and MMEJ check of pL1, pL1Δ22 lines (primer pair 1-4, forward)
pL1_rev_1	TCACAGCTTGCGCTTCTTCC	Heterosplicing and MMEJ check of pL1, CTP-pL1 lines (primer pair 1, reverse)
pL1d22_rev_1	TCAGCCCAGGGTGAACTTAGG	Heterosplicing and MMEJ check of pL1Δ22 lines (primer pair 1, reverse)
CTP-pL1_fwd_1	ATGGCTACGATGAAGATTGCTGTG	Heterosplicing and MMEJ check of CTP-pL1 lines (primer pair 1-4, forward)
pL1_Rv_35St-OH	GACTGGTGATTTTTGCGGACTCTAG	Heterosplicing and MMEJ check of pL1, pL1Δ22, CTP-pL1 lines (primer pair 2, reverse)
101_35S-T_rev	CCCTTATCTGGGAACTACTCACACATTATTCTGGA	Heterosplicing and MMEJ check of pL1, pL1Δ22, CTP-pL1 lines (primer pair 3, reverse)
3342_LdC-HATag-35ST-rev	GGGTTTCGCTCATGTGTTGAGCA	Heterosplicing and MMEJ check of pL1, pL1Δ22, CTP-pL1 lines (primer pair 4, reverse)
pL1-fwd	GCTCGAGATGTCCCTGTG	pL1(Δ22)-GFP fusion construct
pL1-rev	AGGATCCCAGCTTGCGCTTCTTC	(CTP-)pL1-GFP fusion construct
pL1d22-rev	AGGATCCGCCCAGGGTGAACTTAG	pL1Δ22-GFP fusion construct
CTPc5-pL1 fwd	GCTCGAGATGGCTACGATGA	CTP-pL1-GFP fusion construct

1136

1137

1138 Supplementary Table S2 – Predicted chloroplast transit peptides for six selected proteins
1139 in Physcomitrella (the predictions were performed with TargetP-2.0; Armenteros et al., 2019)
1140

Name: CTPc5

Protein: **Pp3c5_1370V3.1**

Predicted chloroplast transit peptide (74 aa):

MATMKIAVAAVAGTPAAAVGTRSAAATETARLCASSAQGLGLSTFVGLRAGSNASRGVTPRVRPLKAENSSVRA

Name: CTPc10.1

Protein: **Pp3c10_10490V3.1**

Predicted chloroplast transit peptide (63 aa):

MEVLPRAIATSHVVVSSSSAPKESSESGSKNNSQVSLSSVPFLRPASALAASIVSSATPAALA

Name: CTPc10.2

Protein: **Pp3c10_2940V3.1**

Predicted chloroplast transit peptide (70 aa):

MALLARGAVAAVAAVGFSNTTSKQGHQNGGSNGVLAFRGNAFCRAVAARTSSVGCQVRVKRGAGPLRVTN

Name: CTPc21

Protein: **Pp3c21_9980V3.1**

Predicted chloroplast transit peptide (43 aa):

MAAVGMTMSLKSSTALKGDFLGAALKQTVAAPKAANVAFAVRA

Name: CTPc22

Protein: **Pp3c22_5470V3.1**

Predicted chloroplast transit peptide (62 aa):

MRALCGVPSCASLSSHRLASPLLAYRACEFALGSSPGLRLEFVGRAVGLQSRSIGRRCIVAM

1141