Production of human papillomavirus type 16 virus-like particles in Physcomitrella photobioreactors

Paul Alexander Niederau[1], Maria Caroline Weilguny[1,2], Sarah Chamas[1], Caitlin Elizabeth Turney[1], Juliana Parsons[1], Marta Rodríguez-Franco[3], Sebastian N.W. Hoernstein[1], Eva L. Decker[1], Henrik Toft Simonsen[2,4], Ralf Reski[1,5,6] *

[1] Plant Biotechnology, Faculty of Biology, University of Freiburg, Schänzlestr. 1, 79104 Freiburg, Germany

[2] Department of Biotechnology and Biomedicine, Technical University of Denmark, 2800 Kongens Lyngby, Denmark

[3] Cell Biology, Faculty of Biology, University of Freiburg, Schänzlestr. 1, 79104 Freiburg, Germany

[4] Laboratoire de Biotechnologies Végétales appliquées aux Plantes Aromatiques et Médicinales, CNRS, UMR 5079, Université Jean Monnet, 42023 Saint-Etienne Cedex 2, France

[5] Signalling Research Centre BIOSS and CIBSS, University of Freiburg, Schänzlestr. 18, 79104 Freiburg, Germany

[6] Cluster of Excellence *liv*MatS @ FIT – Freiburg Centre for Interactive Materials and Bioinspired Technologies, University of Freiburg, Georges-Köhler-Allee 105, 79110 Freiburg, Germany

*Corresponding author: ralf.reski@biologie.uni-freiburg.de

ORCID

P.A.N.	0009-0004-0194-8570
M.C.W.	0009-0009-9440-6872
S.C.	0009-0007-5704-5639
C.E.T.	0009-0001-3113-4648
J.P.	0000-0001-6261-2342
M.R.F.	0000-0003-1183-2075
S.N.W.H.	0000-0002-2095-689X
E.L.D.	0000-0002-9151-1361
H.T.S.	0000-0003-3070-807X
R.R.	0000-0002-5496-6711

Author E-mails

P.A.N.   alexander.niederau@biologie.uni-freiburg.de

M.C.W.   mariaweilguny@gmail.com

S.C.   sarah-chamas@web.de

C.E.T.   caitlin.turney@email.uni-freiburg.de

J.P.   juliana.parsons@biologie.uni-freiburg.de

M.R.F.   marta.rodriguez@biologie.uni-freiburg.de

S.N.W.H.   sebastian.hoernstein@biologie.uni-freiburg.de

E.L.D.   eva.decker@biologie.uni-freiburg.de

H.T.S.   henrik.toft.simonsen@univ-st-etienne.fr

R.R.   ralf.reski@biologie.uni-freiburg.de

Word count: 7830

Figures: 6

Tables: 1

Abstract

Virus-like particles (VLPs) are self-assembling nanoparticles composed of viral structural proteins which mimic native virions but lack viral DNA and infectivity. VLPs are a resourceful class of biopharmaceuticals applied as subunit vaccines or as delivery vehicles for drugs and nucleic acids. Similar to viruses, VLPs are diverse in structure, composition, and assembly, requiring a tailored production platform aligned with the intended application. The moss plant Physcomitrella (*Physcomitrium patens*) is an emerging expression system offering humanized N-glycosylation, scalability, and adaptability to existing industry settings. Here, we used Physcomitrella to produce human papillomavirus (HPV) 16 VLPs. HPV VLPs are composed of the major structural protein L1 and are used as vaccines against HPV infections which are the main causal agent of cervical and other anogenital cancers. We characterized Physcomitrella chloroplast transit peptides, which we used for targeting of moss-produced L1 to chloroplasts, leading to higher recombinant protein yield compared to nuclear or cytoplasmic localization. We confirmed subcellular localization with confocal laser scanning microscopy and found L1 to accumulate within the chloroplast stroma. Production in 5-liter photobioreactors yielded over 0.3 mg L1 per gram fresh weight. We established a purification protocol for moss-produced L1 using a combination of ammonium sulfate precipitation and cation exchange chromatography. Purified samples were subjected to a controlled dis- and reassembly, yielding fully assembled HPV-16 L1 VLPs. This is the first report of production, purification, and assembly of VLPs in a non-vascular plant.

Keywords: bryophyte, bioreactor, molecular farming, nanoparticle, plant biotechnology, vaccine candidate

Key message: First production of virus-like particles as a vaccine candidate in a non-vascular plant.

Introduction

Virus-like particles (VLPs) are nanoparticles consisting of viral structural proteins (Marsian and Lomonossoff, 2016; Nooraei et al., 2021). VLPs cannot replicate and infect cells, but resemble native viruses in size, shape, and surface structure, presenting highly repetitive immunogenic epitopes. As a result, VLPs have a wide range of applications as vaccines and nanocarriers and are of high interest in biopharmaceutical research and development.

There is currently a range of VLP-based vaccines licensed and available against infectious diseases such as hepatitis B, hepatitis E, human papillomavirus (HPV), malaria, and human Norwalk virus with many more vaccine candidates in clinical trials (Nooraei et al., 2021; Gupta et al., 2023). Viral capsid proteins stimulate both the cellular and the humoral immune response, leading to the generation of cytotoxic and memory T cells as well as the production of specific antibodies *via* B cells (Ross et al., 2009; Song et al., 2011; Serradell et al., 2019; Chi et al., 2024). VLP-based vaccines offer improved safety as compared to live-attenuated or inactivated vaccines, which carry the risk of reverting to a pathogenic form (Burns et al., 2014). At the same time, VLPs don't suffer the limitations of DNA-based vaccines, including risk of genome integration or of appropriate delivery which ultimately lowers immunogenicity (Lu et al., 2024). VLPs are not only used for their intrinsic antigenic properties but serve as nanoparticles for drug delivery by loading their inner space with cargo such as nucleic acids (Lamprecht et al., 2016; Adams et al., 2023). Besides the inner space, VLPs can also be used to display antigens on the capsid's surface. For the treatment of pancreatic cancers, disease-unrelated VLPs were coated with respective antigens and were demonstrated to inhibit cancer growth (Cubas et al., 2011; Zhang et al., 2013). Yet, empty VLPs of Cowpea Mosaic Virus, a rod-like shaped plant virus, were shown to reduce cancer growth, despite the absence of disease-specific antigens in the vaccine (Lizotte et al., 2015). The ability of VLPs to induce strong T cell-mediated immune responses is essential in VLP-based cancer vaccines (Nooraei et al., 2021). Overall, VLPs possess characteristics that make them highly versatile for applications in vaccine development, targeted drug delivery, gene therapy, and immunotherapy, offering significant potential in modern biomedicine.

VLPs are classified based on various structural characteristics, one of which is the presence or absence of a lipid envelope (Nooraei et al., 2021). In enveloped VLPs (eVLPs) the viral capsid proteins are imbedded within a lipid membrane acquired during particle assembly, typically from intracellular membranes in the secretory pathway or the outer plasma membrane. Additional structural characteristics include the number and types of capsid proteins, as well as their organization into single, double, or triple-layered architectures. In some instances, capsid proteins require proteolytic processing or N-glycosylation

to enable budding of assembled particles from the host cell's membrane and acquisition of the lipid
envelope (Welsch et al., 2007; Zlotnick and Mukhopadhyay, 2011; Kim et al., 2014). Furthermore, viral
antigens displayed on the surface of the lipid membrane are often glycosylated, which significantly
contributes to the immunogenicity of eVLP-based vaccines (Mortola and Roy, 2004). One example is the
major glycoprotein of Ebola virus for which the distinct glycosylation pattern is crucial for the formation
of neutralizing antibodies (Peng et al., 2022). Other notable cases include hepatitis C virus and Zika virus
– both of which currently lack a licensed vaccine – where the induction of neutralizing antibodies depends
on viral glycoproteins (Torresi, 2017; Hasan et al., 2018). Consequently, the choice of VLP, its structural
complexity, and intended application play a crucial role in determining the appropriate expression system
and guiding cell line development.

The moss Physcomitrella has emerged as an alternative for biopharmaceutical production,
offering several advantages over traditional systems (Reski et al., 2015; Reski, 2018; Decker and Reski,
2020). Its haploid dominant phase and high rate of homologous recombination enable precise gene
targeting (Strepp et al., 1998; Wiedemann et al., 2018; Rempfer et al., 2022). This feature facilitated the
modification of the N-glycosylation pathway, eliminating plant-specific modifications, such as β1,2-
xylosylation, α1,3-fucosylation, and Lewis A epitopes (Koprivova et al., 2004; Parsons et al., 2012; Decker
et al., 2014). Moreover, enzymes necessary for sialic acid synthesis, activation, and linkage to protein N-
glycans were introduced (Bohlender et al., 2020; Bohlender et al., 2022). These modifications have
resulted in more homogeneous and human-like N-glycan structures on recombinant proteins. The
potential of Physcomitrella as a biopharmaceutical production platform is demonstrated by several
products in development. For example, moss-produced α-galactosidase A has successfully completed
clinical phase Ib (Shen et al., 2016; Hennermann et al., 2019). Other moss-produced proteins include
human factor H (FH) and synthetic FH-related multitarget regulators MFHR1 and MFHR13, which are
potential treatments for complement-related disorders (Büttner-Mainik et al., 2011; Michelfelder et al.,
2017, 2018; Top et al., 2019; Ruiz-Molina et al., 2022a) such as age-dependent macular degeneration
(Hector et al., 2025). Factor H produced in other systems faced difficulties in its application, such as *Pichia*
pastoris-produced factor H exhibiting non-optimal glycosylation and reduced half-life (Schmidt et al.,
2011; Kerr et al., 2021). Importantly, Physcomitrella cultivation in photobioreactors is tightly controlled
(Hohe et al., 2002), and thus compliant with Good Manufacturing Practice (GMP) standards, ensuring
high-quality production of recombinant proteins (Reski et al., 2018; Decker and Reski, 2020). As opposed
to transiently transfected *Nicotiana benthamiana* plants cultivated in green houses, cultivation of stable
transgenic Physcomitrella cell lines in bioreactors is more in line with current production processes

applied in industry, thereby facilitating adaption to industry settings (Benvenuto et al., 2023). In addition, cultivation of plant material in bioreactors allows greater scalability, which is a relevant aspect in transitioning to industrial scale production. The company Medicago, focusing on VLP production in transiently transfected *N. benthamiana*, announced challenges in transitioning with their platform to industrial scale production as a reason for seizing further operations in 2023 (Mitsubishi Chemical Group Corporation, 2023). In contrast, the company Eleva upscaled protein production in Physcomitrella from 200 L single-use bioreactors (Niederkrüger et al., 2019) to 1000 L photobioreactors (Eleva, 2025). In summary, Physcomitrella is an ideal production host for complex biopharmaceuticals, including viral antigens or glycosylated proteins, with great prospects regarding scalability and transition to industrial production (Rosales-Mendoza et al., 2014). The first moss-produced vaccine candidate, an ENV-derived multi-epitope HIV chimeric protein, was immunogenic in mice (Orellana-Escobedo et al., 2015). However, until now, there was no record of VLP-production in Physcomitrella, or any other non-vascular plant.

VLPs of the human papillomavirus (HPV) are of great medicinal relevance due to their use as vaccines to prevent HPV infection. HPV infections are a major source of cancer, causing approximately 690,000 cases annually (zur Hausen et al., 1981; zur Hausen, 2002; de Martel et al., 2020). The most prevalent type is cervical cancer, accounting for about 80% of HPV-attributable cancers. Other HPV-related cancers include anogenital and oropharyngeal cancers (Wu et al., 2024). HPV belongs to the *Papillomaviridae*, comprising nearly 200 different types, which are commonly transmitted sexually (Ljubojevic and Skerlev, 2014). The majority of HPV-attributable cancers are linked to two types in particular, namely HPV-16 and HPV-18 (de Martel et al., 2020; Wu et al., 2024). HPV is a non-enveloped double stranded DNA virus with a genome size of 8000 bp, encoding six early regulatory proteins (E1, E2, E4 – E7) and two late structural proteins (L1, L2) (Roden and Stern, 2018). L1 proteins multimerize into pentamers, the capsomers, and L1 capsomers and L2 monomers are imported into the nucleus *via* nuclear localization signals (NLS), where HPV virions assemble (Cerqueira and Schiller, 2017). The final HPV-16 virion is composed of 360 L1 monomers, organized in 72 pentamers, and 72 L2 monomers and is reinforced by inter-pentameric disulfide bonds (Baker et al., 1991; Buck et al., 2005). While the role of L2 in virion assembly differs from type to type, L1 can self-assemble into VLPs in the absence of L2 (Kirnbauer et al., 1992). HPV L1 VLPs are mono-layered and do not contain a lipid envelope. Because the viral life cycle depends on differentiated epithelial cells, it is difficult to obtain large quantities of virions. Therefore, current HPV vaccines are based on L1 VLPs. Although VLP-based HPV vaccines are currently produced in *Escherichia coli*, *Saccharomyces cerevisiae*, and baculovirus-infected insect cells (World Health

Organization, 2024), plant-based expression systems are gaining attention in this respect (Scotti and
Rybicki, 2013; Rybicki, 2020).

While first attempts to produce L1-based VLPs in plants had low yields (Biemelt et al., 2003;
Varsani et al., 2003), considerable improvements have been achieved, e.g. by codon optimization and
subcellular targeting strategies in transiently transformed tobacco (Maclean et al., 2007). A human codon-
optimized version (62% GC content) resulted in highest yields as compared to *N. benthamiana* optimized
(49% GC) and the native sequence (37% GC), presumably not due to the GC content but due to inhibitory
5' sequences, which act on transcription and mRNA processing (Maclean et al., 2007; Hitzeroth et al.,
2018). Further, these authors found chloroplast-targeted L1 yield highest at 0.5 mg / g fresh weight (FW)
followed by cytoplasmic L1 (without the C-terminal 22 aa NLS), whereas L1 targeted to the ER was not
detectable. Similarly, Zahin et al. (2016) reported 0.25 mg L1 / g FW *via* chloroplast targeting while leaving
the NLS intact. Although *N. benthamiana* remains a prominent option, other plants are increasingly
explored for the efficient production of biopharmaceuticals.

Here, we analyzed the expression of HPV-16 L1 in Physcomitrella. We characterized five
Physcomitrella chloroplast transit peptides and investigated the effect of subcellular L1 localization on
recombinant protein yields. We further established L1 production in photobioreactors and subsequent
purification of the product. Finally, the assembly of moss-produced HPV-16 L1 to VLPs was demonstrated
with transmission electron microscopy. Our goal was to obtain proof-of-concept to produce a simple but
medicinally relevant VLP before progressing towards more complex targets such as eVLPs consisting of
multiple proteins and layers.

Materials and methods

Plant material

189   Physcomitrella (new species name: *Physcomitrium patens* (Hedw.) Mitt.; IMSC accession number
190 41269), was cultivated according to Reski and Abel (1985) and Decker et al. (2015).
191

Optimization of HPV-L16 L1 coding sequence

The coding sequence (CDS) of h(human codon-optimized)L1 (GenBank acc. no. DQ067889,
Maclean et al., 2007) was used for optimization using physCO (https://www.plant-biotech.uni-

freiburg.de/tools/physco/; Top et al., 2021). Putative alternative splicing motifs AGGT were identified *via*

search function. MicroRNA-binding sites were predicted *via* psRNATarget with the library of 280 published

Physcomitrella microRNAs and the default settings *Schema V2* (2017 release)

(http://plantgrn.noble.org/psRNATarget/; Dai and Zhao, 2011; Dai et al., 2018). Alternative splice sites

and microRNA-binding sites were removed and substituted by synonymous codons based on

http://www.kazusa.or.jp/codon/. Sequence editing was done in Benchling

(https://www.benchling.com/).

Expression vectors

Chloroplast transit peptides (CTPs) were selected from a list of proteins with experimentally

confirmed cleavage of a predicted CTP (Hoernstein et al., 2024). CTP-eGFP vectors were constructed based

on a vector containing an eGFP CDS with an N-terminal linker (sequence: GGGGGA) under control of the

PpActin5 promoter (Weise et al., 2006) and nos terminator (Hoernstein et al., 2023). CTP candidates were

amplified from Physcomitrella cDNA and inserted at the N-terminus of the linker using Gibson Assembly

(Gibson et al., 2009). The Citrine-L1 fusion constructs are based on a vector containing the PpActin5

promoter, nos terminator and citrine CDS (Wiedemann et al., 2018). The CDS of pL1, pL1Δ22 and CTP-pL1

were amplified using Phusion™ polymerase (Thermo Fisher Scientific, Waltham, USA), cloned into the

pJet1.2 backbone (Thermo Fisher Scientific) and ligated into the expression vector using T4 Ligase (Thermo

Fisher Scientific), *via Xho*I and *BamH*I sites. The pL1 expression vectors were constructed based on a vector

containing the PpActin5 promoter, nos terminator and hpt cassette (Ruiz-Molina et al., 2022a). The pL1

sequence was synthesized by Genewiz (Leipzig, Germany). For constructs pL1 and pL1Δ22, the pL1 CDS

was amplified using Phusion™ polymerase (Thermo Fisher Scientific) and cloned as described above. For

construct CTP-pL1, the pL1 and CTPc5 CDS were amplified and inserted into the backbone using Gibson

Assembly. Vectors were sequenced by Eurofins Genomics (Ebersberg, Germany). Vector maps are

compiled in **Supplementary Figure S1**. Primers were synthesized by Eurofins Genomics and are compiled

in **Supplementary Table S1**.

Protoplast preparation and transformation

Physcomitrella protoplasts were prepared and transformed based on existing protocols (Hohe

and Reski, 2002; Schween et al., 2003; Hohe et al., 2004; Decker et al., 2015). Transformed protoplasts

were incubated at 22 °C for 24 h in the dark and subsequently for 72 h at a 16/8-h light/dark photoperiod

and light intensity of 50–70 µmol m^{-2} s^{-1} before plating on KnopME + 25 mg/L hygromycin agar plates

covered with cellophane for two weeks. Surviving colonies were transferred to non-selective conditions

for two weeks before a second selection on KnopME + 50 mg/L hygromycin.

Fluorescence microscopy

Protoplasts were analyzed 2-3 days after transfection with an AxioCam MRc5 camera on an

Axioplan2 stereo microscope (Zeiss, Oberkochen, Germany). Pictures were taken with a 40x objective or

a 63x objective with immersion oil. A FITC filter was used for eGFP, a rhodamine filter for chlorophyll

autofluorescence, and a FITC_LP filter for the combination of the two. Images were edited with GIMP

2.10.22 and ImageJ. For confocal microscopy, an AXIO Observer.Z1 Inverted Fluorescence Motorized

Phase Contrast Microscope Pred 7 with the objective Plan-Apochromat 63x/1.4 Oil DIC M27 (Zeiss) was

used. EGFP signals were detected at 488 / 509 nm, and chlorophyll autofluorescence at 587 / 610 nm.

Images were edited in ZEN 3.5 (Blue edition, ZEN Lite), ImageJ, and GIMP 2.10.22.

For analysis of subcellular targeting, an inverted confocal laser scanning microscope LSM 880 with

the objective LD LCI Plan-Apochromat 40x/1.2 H$_2$O autocorr and FastAiryScan mode (Zeiss) was used.

Citrine signals were detected at 488 / 516 nm, and chlorophyll autofluorescence at 561 / 595 nm. Images

were edited in ZEN 3.9 (Blue edition) and processed *via* deconvolution using default setting and

orthogonal projection using maximum intensity projection.

Biomass accumulation

Flasks containing 30 mL KnopME medium were inoculated at a density of 60 mg dry weight (DW)/L

and cultured for 4 weeks at the conditions described above. Biomass DW was measured by filtering 3x 10

mL of culture through miracloth and subsequent drying at 105 °C for 2 hours. Biomass accumulation was

analyzed by linear regression analysis. Slopes were compared using ANCOVA and post hoc multiple

comparisons test (TukeyHSD). The analysis was conducted in RStudio (version 4.2.2) and GraphPad Prism

(version 8.1).

Bioreactor operation

Stirred tank bioreactors (Getinge, Sweden) were inoculated at 60 mg DW/L moss density in 5 L

KnopME. The light intensity was set to 160 µmol m^{-2} s^{-1} and increased to 350 µmol m^{-2} s^{-1} after 4 days. At

day 4, auxin 1-Naphtalene acetic acid (NAA, Sigma, St. Louis, USA) dissolved in 0.5 M KOH was added to a

final concentration of 5 µM. Throughout the bioreactor run, the pH was kept at pH 5.8, temperature at
22 °C, and aeration at 0.3 vvm (volume air per volume media per minute) with 2% CO_2. The culture was
continuously agitated with a pitched 3 blade impeller at 500 rpm.

Protein extraction, purification and VLP assembly

For small scale extraction, 30 – 60 mg fresh weight (FW) of vacuum-filtered moss material was
combined with a glass and a steel bead (each 3 mm in diameter) in a 2 mL reaction tube, frozen in liquid
nitrogen and the tissue lysed using a tissue lyser for 2 min at 28 Hz. Homogenized material was
resuspended in PBS (2.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 0.137 M NaCl, pH 7.2), 0.01% (v/v)
Tween20, and protease inhibitor (P9599, Sigma) in a ratio 3 mL/g. Samples were placed in an ice-cold
sonication bath for 15 min, centrifuged for 30 min at 14,000 x g and 4 °C, and the supernatant taken off
for further analysis.

For large scale protein extraction, 5 g FW of vacuum-filtered moss material were frozen in liquid
nitrogen and ground to a fine powder. The material was combined with 22 mL PBS and 0.01% (v/v)
Tween20, 50 mM DTT and 100 µL protease inhibitor. The mixture was homogenized using an ULTRA-
TURRAX (IKA, Staufen, Germany) at 10,000 rpm for 10 min on ice. Next, the extract was treated with a
Q500 sonicator using a CL-334 converter (QSonica, Newtown, USA) at amplitude 55%, 10s on & 40s off
for 20 min at 8 °C. Debris was removed by two centrifugations at 4500 x g for 10 min and 20,000 x g for
20 min, both at 4 °C. The sample was combined with solid $(NH_4)_2SO_4$ to a final concentration of 45% (w/v)
and proteins precipitated under stirring for 1 h at 8 °C. Amounts of $(NH_4)_2SO_4$ were calculated using an
online tool (https://files.encorbio.com/protocols/AM-SO4.htm) (EnCor Biotechnology Inc., Gainesville,
USA). Precipitated proteins were pelleted by centrifugation at 12,000 x g for 10 min at 8 °C. Pellets were
dissolved in 4 mL cation exchange (CaEx) chromatography binding buffer (2.7 mM KCl, 10 mM Na_2HPO_4,
1.8 mM KH_2PO_4, 0.068 M NaCl, 0.01% (v/v) Tween 20, 5% (v/v) Glycerol, 5 mM DTT, pH 6.8) and dialyzed
(20k MW cut off, 88405, Thermo Fisher Scientific) overnight against the same buffer at 8 °C. The sample
was diluted to a final volume of 10 mL using binding buffer, and passed through a 0.22 µm filter (Roth,
Karlsruhe, Germany) before loading onto a 1 mL HiTrap SP HP cation exchange column (Cytiva,
Marlborough, USA), connected to an ÄKTA^{TM} start (Cytiva). The column was washed with 5 column
volumes (CV) of 0.3 M NaCl (based on binding and elution buffer) and L1 eluted *via* 4 CV of elution buffer
(2.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 1 M NaCl, 0.01% (v/v) Tween 20, 5% (v/v) Glycerol, 5 mM
DTT, pH 6.8)

Eluted fractions containing L1 were pooled and dialyzed against disassembly buffer (22.7 mM KCl,

10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 0.068 M NaCl, 0.05% (v/v) Tween20, 5 mM DTT, pH 8.5) overnight at 8

°C with 3x buffer exchange. Samples were concentrated 5x using a vacuum concentrator (Concentrator

plus, Eppendorf, Hamburg, Germany). For reassembly, samples were dialyzed against reassembly buffer

(22.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, 1 M NaCl, 0.05% (v/v) Tween20, pH 6) overnight at 8 °C

with 3x buffer exchange.

SDS-PAGE and western blot

SDS-PAGE and western blots were performed based on Bohlender et al. (2022). Two systems were

used for protein separation, either 7.5% polyacrylamide gels (Mini-PROTEAN® TGX™ Precast Gels, Rio-

Rad, Munich, Germany) in TGS buffer at 120 V, or 4 – 12% Bis-Tris gels (mPAGE™ Precast Gels, Merck,

Darmstadt, Germany) in MOPS buffer at 150 V. As first antibody served 1:5000 diluted CamVir1 (Thermo

Fisher Scientific), as second antibody peroxidase-linked rabbit anti-mouse secondary antibody (Cytiva)

diluted 1:12,500. For dot blots, protein samples were blotted on a PVDF membrane (Cytiva) using a

filtration manifold kit (SRC-96/1, Schleicher&Schuell, Dassel, Germany). Antibody H16.V5

(AntibodySystems, Schiltigheim, France) was diluted 1:5000. Dot blots were developed as described for

western blots. Yeast-produced HPV-16 L1 (Abcam, Cambridge, UK) was used as a positive control.

L1 quantification

The ELISA protocol was based on Studentsov et al. (2002). In short, 96-well microtiter plates

(Greiner, Kremsmünster, Austria) were coated with 1:500 CamVir1 antibody (Thermo Fisher Scientific) in

PBS overnight at 8 °C. Wells were washed twice with washing buffer (1x PBS, 0.05% (v/v) Tween20) and

blocked with blocking buffer (1x PBS, 1% (w/v) polyvinyl alcohol MW 47,000, 0.05% (v/v) Tween20) for 3

h at 37 °C. After 3x washing, samples were diluted in blocking buffer and incubated in the wells for 1.5 h

at 37 °C. Wells were washed 3x before adding 1:1000 diluted polyclonal anti-HPV-16 L1 antibody from

rabbit (Cusabio, Houston, USA) for 1 h at 37 °C. After 5x washing, wells were incubated with 1:1000 diluted

anti-rabbit HRP-conjugated antibody NA934 (Cytiva) for 1 h at 37°C. Wells were washed 3x before adding

100 µL TMB substrate (Thermo Fisher Scientific). The reaction was stopped with 50 µL of 2 M H_2SO_4.

Absorbance was measured at 450 / 595 nm for reference. The signal of a wildtype sample was subtracted

to account for background signal.

Transcription electron microscopy

The TEM protocol was based on Abel et al. (1989). Negative staining of purified VLPs and moss extracts was performed using 300 Mesh glow discharged formvar/carbon-coated copper grids (Electron Microscopy Sciences, Hatfield, USA). Five µL of sample were applied on a grid and incubated for 5 min at RT. Grids were washed 3x with water and stained with 2% (w/v) uranyl acetate. After 30 sec, uranyl acetate was removed with a filter paper, grids were air dried, and inspected with a Hitachi HT7800 TEM (Tokyo, Japan) coupled to a Xarosa CMOS camera (Emsis, Münster, Germany).

Results

Adaptation of L1 coding sequence for Physcomitrella-optimized expression

The coding sequence (CDS) of HPV-16 L1 was optimized for expression in Physcomitrella. The goal was to adapt the CDS regarding codon usage and preventing potential degradation through microRNAs and splicing. A human codon-optimized version (hL1, **Supplementary Figure S2**) was used as a basis for optimization since it had led to higher yields than plant-optimized versions or the native viral sequence in *Nicotiana benthamiana* (Maclean et al., 2007).

For codon optimization, physCO (Top et al., 2021) was used, which changed nine out of 505 codons and slightly increased the GC content from 62.25% to 62.85% (**Supplementary Figure S2**). Next, the CDS was scanned for the motif AGGT, which is the most common motif at exon-intron borders for alternative splicing in Physcomitrella (Top et al., 2021). Transformation of a heterologous cDNA containing the AGGT motif led to alternative splicing in Physcomitrella, a process referred to as heterosplicing (Top et al., 2021). Scanning the physCO-optimized hL1 sequence yielded 5 AGGT-motifs which were replaced by synonymous codons (**Supplementary Figure S2**). MicroRNAs bind complementary sequences in the mRNA and thereby cause inhibition of translation or degradation of mRNA (Jones-Rhoades et al., 2006; Khraiwesh et al., 2010). One microRNA binding site was predicted within the CDS using psRNATarget (Dai and Zhao, 2011; Dai et al., 2018). Therefore, we changed three bases within the predicted binding region using synonymous codon substitutions. The final Physcomitrella-optimized CDS was named pL1 (**Supplementary Figure S2**) and differs from hL1 by 17 out of 1515 nucleotides without changing the amino acid (aa) sequence.

Characterization of chloroplast transit peptides

Different constructs were designed to target L1 to specific cellular compartments, including chloroplasts. Fusing chloroplast transit peptides (CTPs) to the L1 CDS led to increased protein yields in tobacco as compared to constructs lacking CTPs (Maclean et al., 2007; Zahin et al., 2016). To assess whether this is also true for L1 production in Physcomitrella, a selection of Physcomitrella-derived CTPs was characterized.

Initially, the CTP of FtsZ1-1 (Gremillon et al., 2007) was considered. However, cleavage site analysis *via* TargetP-2.0 (Armenteros et al., 2019) predicted 37 aa of this CTP to remain at the N-terminus of L1 after chloroplast import. This CTP was rejected because the additional aa could change L1 properties and hamper VLP assembly. A list of alternative CTPs was derived from the Physcomitrella N-terminome (Hoernstein et al., 2024) (**Supplementary Table S2**). Here, experimentally detected protein N-termini were compared with TargetP-2.0 predictions for cleavable N-terminal targeting sequences. Five CTPs were selected where an exact match between prediction and experimentally validated N-terminus was observed across different experiments (Hoernstein al., 2024), namely CTPc5, CTPc10.1, CTPc10.2, CTPc21, CTPc22 (**Table 1**). Their sequences were fused *in silico* to the N-terminus of L1 and analyzed with TargetP-2.0 to validate the correct cleavage of the fusion.

Our *in-silico* analysis indicated CTPc22 to end with a methionine, which could introduce an alternative translation initiation site, resulting in translation of L1 without the CTP (**Table 1**). Therefore, an alternative version lacking the C-terminal methionine (CTPc22noMet) was included. The likelihoods for chloroplast localization were above 70% for all candidates except CTPc10.1 (**Table 1**), which was rejected.

The remaining five CTPs (CTPc5, CTPc10.2, CTPc21, CTPc22, CTPc22noMet) were fused to the CDS of eGFP (**Supplementary Figure S1**) and expressed in Physcomitrella protoplasts. Analysis *via* fluorescence microscopy confirmed localization of eGFP signals in chloroplasts for CTPc5, CTPc10.2, CTPc22, and CTPc22noMet, but not for CTPc21 (**Supplementary Figure S3**). Of the four remaining CTPs, CTPc5 had the highest likelihood for plastid localization (**Table 1**) and were examined *via* confocal microscopy (**Supplementary Figure S3**). Finally, we selected CTPc5 for the chloroplast-targeting of L1.

Table 1. *In-silico* analysis of Physcomitrella-derived chloroplast transit peptides. CTPs were selected based on mass spectrometry datasets (Hoernstein et al., 2024). *In-silico* predictions of likelihood of chloroplast localization target sequence cleavage were performed with TargetP-2.0.

Name of CTP	Originating from protein model	Predicted cleavage of CTP-L1	Length (aa)	Likelihood of GFP chloroplast localization	Likelihood of L1 chloroplast localization
FtsZ1-1	Pp3c22_4940V3.1	37 aa remain on N-terminus	84	0.95	0.92
CTPc5	Pp3c5_1370V3.1	-	74	0.987	0.943
CTPc10.1	Pp3c10_10490V3.1	-	63	0.456	0.229
CTPc10.2	Pp3c10_2940V3.1	-	70	0.908	0.845
CTPc21	Pp3c21_9980V3.1	-	43	0.956	0.891
CTPc22	Pp3c22_5470V3.1	-	62	0.939	0.739
CTPc22noMet	Pp3c22_5470V3.1	-	61	0.913	0.727

Targeting of L1 to nuclei and chloroplasts

To evaluate the impact of L1 localization on product yield, we generated three variants (pL1, pL1Δ22, CTP-pL1). pL1 contains the full-length pL1 CDS, which includes the native HPV-16 NLS (Cerqueira and Schiller, 2017). For pL1Δ22 the NLS was removed. In CTP-pL1 the Physcomitrella CTPc5 was fused to the 5' of the pL1 CDS, including the NLS.

After generation of these CDS variants, we aimed to evaluate L1 targeting to subcellular compartments. The CDS of pL1, pL1Δ22 and CTP-pL1 were fused to the 5' end of Citrine CDS (**Supplementary Figure S1**). Physcomitrella protoplasts were transfected with the plasmids and Citrine localization was analyzed using fluorescence microscopy. A vector encoding Citrine without any L1 variant fusion served as a control.

Cells transformed with the control vector exhibited predominantly cytoplasmic Citrine signals, whereas diffuse signals also occurred in nuclei (**Figure 1**). For pL1-Citrine, the signal localized to nuclei, demonstrating that the HPV-16 NLS is functional in Physcomitrella. This is further supported by the fact that removal of the NLS in pL1Δ22-Citrine results in cytoplasmic localization of the signal. For CTP-pL1-

394 Citrine, the signal is predominantly in chloroplasts, but is not evenly distributed, as seen for CTP-eGFP

395 fusions lacking L1 (**Supplementary Figure S3**). Instead, the pL1-Citrine fusion concentrates in several spots

396 of high fluorescence intensity and less than 1 μm in diameter. It is unclear whether this difference is due

397 to distinct features of GFP and Citrine or due to the presence of L1 and potential formation of L1

398 multimeric structures. Overall, we confirmed successful L1 targeting to the intended compartments.

400
401

Figure 1. Subcellular localization of L1 variants. Confocal microscopy images of Physcomitrella

protoplasts transformed with constructs encoding (CTP-)pL1(Δ22)-Citrine fusion proteins or Citrine alone.

For each sample, chlorophyll autofluorescence in magenta is seen on the left, Citrine signals in the middle

and the merged signals on the right. In the control and pL1Δ22-Citrine, the Citrine signals do not enter the

chloroplasts and are mainly in the cytoplasm. For pL1-Citrine, the Citrine signals are in the nucleus. In the

CTP-pL1-Citrine sample, the Citrine signals form small spots of high intensity in the chloroplasts.

Characterization of L1 production

After validating the subcellular localization, pL1, pL1Δ22 and CTP-pL1 expression constructs were

cloned (**Supplementary Figure S1**). The linearized constructs were transferred into Physcomitrella

protoplasts and obtained plants were screened for productivity by western blot and ELISA.

Western blot analysis revealed the L1 monomer of 56 kDa (pL1, CTP-pL1) and 54 kDa (pL1Δ22)
(**Figure 2**). In addition, pL1 and CTP-pL1 lines exhibited a band at approximately 54 kDa. All lines showed
another band at around 45 kDa, even though the signal was relatively weak in CTP lines and more
prominent in pL1Δ22 lines. Smaller protein isoforms can originate from degradation or DNA or RNA,
resulting in truncated transcripts and, subsequently, truncated protein isoforms (Murén et al., 2009; Top
et al., 2021). The latter possibilities were ruled out by amplifying the L1 CDS *via* PCR from cDNA, which
yielded amplicons of the expected sizes, corresponding to the western blot bands of 56 kDa (pL1, CTP-
pL1) and 54 kDa (pL1Δ22), but not additional amplicons (**Supplementary Figure S4**). Also, two bands are
visible at 130 – 180 kDa, of which the highest band at 180 kDa is also present in the yeast-produced L1
control. This pattern of two close-by bands repeats another two times at higher molecular weight outside
of the marker range (**Figure 2**). Addition of reducing agents DTT or ascorbic acid to the extraction buffer
markedly reduces the western blot signal intensity above 180 kDa (**Supplementary Figure S5**). We
hypothesize that in the absence of reducing agents, reactive oxygen species (ROS) in the plant extract
cross-link L1 with itself and/or host proteins. Irrespective of the presence or absence of reducing agents
during extraction and purification, two prominent bands in the range of 130 and 180 kDa are present in
all western blots (**Figures 2 and 4, Supplementary Figure S7**). Western blot and ELISA analysis indicate
higher yields in CTP-pL1 lines as compared to pL1 or pL1Δ22 lines. In shaking flasks, the best producer is
CTP-pL1 30 with 142 µg L1/g FW (**Figure 2**).

Besides the amount of recombinant protein per g FW biomass, the accumulation of biomass is
important in biomanufacturing as high product yields per biomass can be trivial if biomass accumulation
is slow. To study if recombinant protein amount or subcellular localization of L1 hampers plant growth,
we analyzed the time course of biomass accumulation of the nine lines in shaking flasks for 4 weeks.

In our conditions, exponential growth could not be measured. This might be due to the culture
conditions, to the rather long intervals between sampling, and/or to the intrinsic growth characteristics
of Physcomitrella protonema, which grows by apical cell division (Reski, 1998). Therefore, biomass density
was fitted using linear regression analysis model (**Supplementary Figure S6**), using the calculated slope to
approximate biomass accumulation rates. Comparison of the slopes between the lines pL1, pL1Δ22 and
CTP-pL1 showed pL1 lines accumulated biomass faster than pL1Δ22 (p = <0.0001) and CTP-pL1 (p = 0.0001)
lines. If this difference is due to the distinct L1 localization or to lower L1 production in pL1 lines (**Figure
2**) is unclear. Line pL1 47B showed the highest slope of 255 mg dry weight (DW)/L/week, as compared to
line CTP-pL1 30 with a slope of 211 mg DW/L/week. However, this difference is negligible considering the

yield of pL1 47B is 8 µg L1/g FW as to 142 µg L1/g FW for CTP-pL1 30 (**Figure 2**). Consequently, line CTP-

pL1 30 was chosen for VLP production.

447
448

Figure 2: Analysis of HPV-16 L1 production *via* western blot and ELISA. (**a**) For each of the constructs pL1,

pL1Δ22, and CTP-pL1 three lines were tested *via* western blot for expression of HPV-16 L1. 50 ng *S.*

cerevisiae-produced L1 served as a positive control and Physcomitrella wildtype as a negative control. The

proteins for the western blot were separated on a 7.5% TGS polyacrylamide gel under reducing conditions

before being transferred to a membrane and developed using CamVir1 antibody (1:5000). (**b**) Production

levels of L1 were quantified *via* ELISA. Wildtype background signal was subtracted before analysis *via* one-

way ANOVA and post hoc multiple comparisons test (TukeyHSD). Means and standard deviations are

calculated based on n ≥ 3 technical replicates from two dilutions and are given per biomass fresh weight

(FW). Statistical significance (α = 0.05) is indicated by Compact Letter Display.

Upscaled L1 production

For production of VLPs, the cultivation of line CTP-pL1 30 was upscaled to 5 L-photobioreactors. The goal here was to identify the best time point for harvesting material used for subsequent L1 purification.

Biomass increased from initially 60 mg DW/L to around 870 mg DW/L at day 4 (**Figure 3**). L1 production remained constant during the first 4 days, in the range of the yield observed for flasks. Upon addition of the phytohormone auxin (NAA) on day 4, L1 levels increased on day 6 and peaked at day 7 at 332 µg/g FW, 130% higher as compared to day 0. Afterwards, L1 production dropped sharply to levels prior to auxin addition. Biomass continued to accumulate until day 11, after which it dropped slightly. These conditions, including day 7 as the ideal harvesting time point, were used for further L1 production.

Figure 3. Analysis of biomass accumulation and L1 production of line CTP-pL1 30 in a 5 L-photobioreactor. The bioreactor was inoculated with 60 mg dry weight (DW)/L and monitored over a course of 13 days. On day 4, 5 µM auxin (NAA) was added. L1 production peaked at day 7 at 332 µg/g fresh weight (FW). Biomass and L1 measurements are based on n ≥ 3 technical replicates. Production levels of L1 were quantified *via* ELISA.

Two-step purification removes bulk of cellular proteins

Purification of L1 was performed using ammonium-sulfate (AMS) precipitation, followed by cation exchange (CaEx) chromatography. While AMS removes the bulk of cellular proteins, CaEx removes residual proteins and concentrates the sample.

We tested the efficacy of 40%, 45% and 50% (w/v) AMS to precipitate L1 from an extract
containing the soluble protein fraction from CTP-pL1 30 plant material produced in the bioreactor. We
found 45% AMS to be sufficient to effectively remove L1 from the supernatant, whereas at 40% AMS small
amounts of L1 remained in the supernatant and at 50% AMS higher amounts of cellular protein were
precipitated together with L1 (**Supplementary Figure S7**). In preparation for chromatographic purification,
the precipitated proteins were resuspended in CaEx binding buffer and dialyzed overnight in binding
buffer to remove residual AMS. Cloudiness of samples indicated a reduced solubility of precipitated
proteins, and undissolved matter was removed by filtration. The relative amount of L1 and host protein
was evaluated *via* western blots and Coomassie-staining, respectively (**Supplementary Figure S7**).
Samples after filtration showed a slight reduction in host protein content, but, except for the sample at
40% AMS, no reduction in L1. Overall, precipitation at 45% AMS was selected, which is in line with
purifications of yeast-produced L1 (Park et al., 2008; Kim et al., 2010).

Next, CaEx chromatography was used to remove residual cellular proteins. The bulk of cellular
proteins did not bind to the column and was detected in the flow through (**Figure 4**, **Supplementary Figure
S8**). Notably, a large amount of L1 did not bind to the column and was detected in the flow through.
Different binding buffer compositions were tested to increase L1 binding to the column, e.g. lowering the
pH from 7.2 to 6.8, lowering the concentration of NaCl from 0.137 M to 0.068 M, addition of 5% (v/v)
glycerol, and re-loading of flow-through fraction. Nevertheless, a large part of L1 did not bind to the
column and was found in the flow through. During column wash with 5 mL at 0.3 M NaCl, small amounts
of contaminating proteins and L1 were washed off the column. For elution, NaCl was increased to 1 M
NaCl and 0.5 mL fractions were collected, in total 4 mL. L1 eluted from the column in fractions E4 − E8
(**Figure 4**), which were pooled for further analysis.

Figure 4. Two-step purification of L1 removes bulk of cellular protein. Analysis of L1 and residual cellular

host proteins during purification under reducing conditions *via* western blot and Coomassie-stained SDS-

PAGE (12% Bis-Tris gel). After precipitation with 45% (w/v) AMS, the filtered sample was subjected to

cation exchange chromatography (CaEx) for final purification. Column-bound L1 eluted at 1 M NaCl in

fractions E4 – E8. The asterisk in the Coomassie-stained SDS-PAGE marks the band referring to HPV-16 L1.

50 ng *S. cerevisiae*-produced L1 served as a positive control and Physcomitrella wildtype as a negative

control.

Capsomers and VLPs

Next, we performed a dis- and reassembly protocol on the pooled elution fractions, as it yields

more uniform and stable VLPs, which are less prone to aggregation (McCarthy et al., 1998; Mach et al.,

2006). First, pooled elution fractions were dialyzed at low salt concentrations, high pH and DTT, that

results in the disassembly of VLPs or other multi-pentameric L1-based structures into capsomers. Second,

samples were dialyzed at high salt, low pH and in the absence of reducing agent, that leads to the

reassembly of capsomers into VLPs.

To assess the conformation of L1 throughout the process, dot blots were performed using

different monoclonal antibodies. CamVir1 detects a linear epitope in both denatured and native L1 and

served as a control. In contrast, H16.V5 detects only conformational epitopes in both, L1 capsomers and

VLPs, but not monomeric or denatured L1.

The dot blot showed the presence of L1 in all CTP-pL1 30-derived samples throughout the

purification process, as well as yeast-derived denatured and native L1 (**Figure 5**). In comparison, H16.V5

detects L1 in the native positive control and in all CTP-pL1 30-derived samples, indicating the presence of

at least capsomers, if not VLPs, already in the raw extract. Further, conformational epitopes seem to be

stable throughout AMS precipitation, CaEx chromatography, disassembly, and reassembly. However, the

reassembly sample exhibits a weaker signal for H16.V5 as compared to CamVir1.

Figure 5. Moss-produced HPV-16 L1 assembles into capsomers, which are stable throughout

purification. Dot blots were performed using two different antibodies to analyze the formation of

capsomers in yeast-produced L1 and samples from CTP-pL1 30 purification (extract, after AMS

precipitation, elution) and dis- and reassembly. The CamVir1 antibody detects a linear epitope in both

denatured and native L1. The neutralizing antibody H16.V5 detects specifically conformational epitopes

in L1 capsomers and VLPs. Positive H16.V5 signals indicate HPV-16 L1 monomers to assemble at least into

capsomers, *in vivo* or after extraction, which are stable throughout the purification. The experiment only

allows qualitative but not quantitative comparison since sample concentrations were adjusted

individually. `L1 denatured´ was incubated at 95 °C for 5 min, all other samples were kept at 4 °C. A wildtype sample was used as a negative control.

The assembly of L1 into VLPs was verified with Transmission Electron Microscopy (TEM). As a control, yeast-produced L1 was analyzed without prior treatment, after disassembly, and after reassembly. In the untreated sample, particles of approx. 60 nm in size were detected. After disassembly, structures of 5 – 10 nm in size were observed, corresponding in size and appearance to capsomers observed for yeast-derived L1 as part of this study (**Figure 6b**) and elsewhere (McCarthy et al., 1996). VLPs reshape upon reassembly treatment, demonstrating the successful dis- and reassembly of yeast-derived L1 VLPs. Notably, reassembled VLPs are heterogenous in size and shape, despite the controlled dis- and reassembly (**Figure 6**).

Figure 6. Analysis of dis- and reassembled VLPs from moss and yeast *via* transmission electron

microscopy (TEM). Yeast-derived L1 was analyzed untreated (**a**), after disassembly (**b**), and after

reassembly (**c**) *via* TEM. CTP-pL130 (**d** – **f**) and wildtype (**g** – **i**) extracts were purified *via* AMS precipitation

and CaEx chromatography (**d** & **g**), after disassembly (**e** & **h**), and after reassembly (**f** & **i**). Untreated yeast-

derived VLPs (**a**) disintegrate into capsomers upon disassembly (**b**) and reshape into VLPs upon reassembly

treatment (**c**). TEM shows the presence of moss-derived HPV-16 L1 capsomers in the elution (**d**) and

disassembly sample (**e**) and the assembly into VLPs in the reassembly sample (**f**). Capsomer and VLP-like

structures are absent from the wildtype control samples. VLPs are indicated by black arrows.

567   To analyze L1 assembly into VLPs, samples were taken from the pooled CaEx chromatography

568 elution fractions, after disassembly, and after reassembly.

569   In the elution sample, no VLPs but capsomeric structures were observed, comparable to those in

the disassembled yeast sample, which are not present in the corresponding wildtype control (**Figure 6**).

The abundance of capsomers in the elution fractions was surprising given the fact that the elution buffer

has a high ionic strength, which favors reassembly. However, before column loading, the sample was

dialyzed overnight in binding buffer, which favors disassembly. The exposure to high ionic strength during

elution might have been too short to allow VLP assembly. In addition, both binding and elution buffer

contained DTT, which induces disassembly (McCarthy et al., 1998; Mach et al., 2006). The disassembled

CTP-pL1 30-derived sample depicted capsomers as seen in the elution fraction, and for disassembled

yeast-derived L1 (**Figure 6**). **Figure 6f** displays fully assembled, moss-derived HPV-16 L1 VLPs. Again, VLPs

were heterogenous in size and shape, as seen for reassembled yeast-derived VLPs. Moss-produced VLPs

tended to aggregate, which was not observed for reassembled yeast-derived VLPs, but is common for

HPV-16 L1 VLPs (Shi et al., 2005). Aggregation might also explain the relatively low signal for the

conformational antibody H16.V5 in the sample of reassembled moss-derived VLPs (**Figure 5**). No VLPs

were detected in wildtype samples (**Figure 6**).

Discussion

585   The goal of this study was to gain proof-of-concept for the production of virus-like particles (VLPs)

in the moss plant Physcomitrella by producing HPV-16 L1 VLPs. For this purpose, we generated a coding

sequence (CDS) optimized for expression of HPV-16 L1 in Physcomitrella. We further characterized five

chloroplast transit peptides (CTPs) and confirmed the localization of L1 in chloroplasts. In addition, we

demonstrated chloroplast localization to yield higher L1 levels than targeting the protein to nuclei or

cytoplasm. Cultivation in photobioreactors yielded more than 0.3 mg / g FW, which is in the range of other

plant-based expression platforms. We established a purification protocol and demonstrated the

successful assembly of moss-produced HPV-16 L1 VLPs.

In tobacco plants (*Nicotiana benthamiana*), various strategies were explored for increasing

production yields of HPV-16 L1 VLPs, one of which is optimization of codon usage (Maclean et al., 2007;

Hitzeroth et al., 2018). Hitzeroth et al. (2008) found that codon optimization of L1 improves transcription,

rather than translation. This is due to RNA motifs in the L1 CDS, which post-transcriptionally regulate the

virus life cycle and lead to pre-mRNA degradation (Mori et al., 2006; Zhao and Schwartz, 2008). It is argued

that codon optimization removes those motifs, thereby increasing mRNA and protein levels, irrespective

of whether the CDS was optimized for expression in humans, tobacco (Hitzeroth et al., 2018) or

Physcomitrella (this study). Our Physcomitrella-optimized CDS has a GC content of 62.85% as compared

to hL1 with 62.25% (Maclean et al., 2007) and the wildtype sequence with 38.27% (Varsani et al., 2003).

Combined with the removal of microRNA binding sites and heterosplice sites we obtained the

Physcomitrella-optimized pL1 CDS. We argue that pL1 addresses the issues of L1 intrinsic degradation

motifs as well as Physcomitrella-specific posttranscriptional degradation.

Besides codon optimization, research focused on localization signals and transit peptides. The

most common strategy is sequestration to chloroplasts, either *via* nuclear transformation and

posttranslational import into chloroplasts (Maclean et al., 2007; Matic et al., 2011; Pineo et al., 2013;

Zahin et al., 2016; Hitzeroth et al., 2018; Naupu et al., 2020; Muthamilselvan et al., 2023) or *via*

transplastomic lines (Fernández-San Millán et al., 2008; Lenzi et al., 2008). Here, we experimentally

characterized five Physcomitrella CTPs, of which we used one (CTPc5) for sequestration of L1 to

chloroplasts. Employing the HPV-16 native NLS, L1 was targeted to nuclei, whereas the removal of the NLS

resulted in cytoplasmic L1. Earlier studies confirmed functionality of the NLS in mammalian cells (Leder et

al., 2001; Bissa et al., 2015). However, the NLS failed to localize L1 to nuclei in tobacco (Šmídková et al.,

2010). Our observation of the HPV-16 native NLS targeting L1 to nuclei in Physcomitrella is another

example for a conserved cross-kingdom functionality of localization signals between moss and mammals

(Gitzinger et al., 2009).

Western blots of L1-producing Physcomitrella lines showed a variety of bands, both smaller and

larger than the expected 56 kDa or 54 kDa. A comparable banding pattern was observed when expressing

L1 in tobacco chloroplasts (Fernández-San Millán et al., 2008). Likewise, yeast-produced L1 exhibited

bands with the size of L1 dimers and trimers (Kim et al., 2007; Kim et al., 2010). L1 dimers and trimers are

based on inter-capsomeric disulfide bridges which stabilize virions and VLPs (Sapp et al., 1998; Buck et al.,

2005; Buck et al., 2013). McCarthy et al. (1998) resolved L1 multimers with DTT. Likewise, Buck et al.

(2005) demonstrated absence and presence of L1 dimers and trimers in a comparison of reducing and

non-reducing SDS-PAGE and western blots, respectively. An investigation of cysteine 175 and cysteine 428

L1 mutants resulted in the absence of bands at 130 kDa and 180 kDa, thereby linking disulfide bridge-

based dimers and trimers to the observed bands (Buck et al., 2005). Surprisingly, reducing agents such as

DTT, ascorbic acid or β-mercaptoethanol did not result in the disappearance of these bands in our and

other studies (Kim et al., 2007; Fernández-San Millán et al., 2008; Kim et al., 2010). This indicates a

tolerance of L1 dimers and trimers to reducing agents or a different origin of multimerization observed at

130 kDa and 180 kDa. Castells-Graells and Lomonossoff (2021) observed multimerization of *Nudaurelia*

capensis omega virus α-peptide produced in tobacco, irrespective of DTT, but not when produced in insect

cells. These authors hypothesize that cross-linkage of α-peptide occurs during extraction and purification

and occurs in plant-based expression systems. Regarding the observed bands below 56 kDa, proteolytic

degradation was proposed in other studies observing the same phenomenon (Buck et al., 2005;

Fernández-San Millán et al., 2008). In tobacco, a shift from bands above 100 kDa in young leaves to more

prominent bands at 45 kDa and 20 kDa in older leaves indicated proteolytic degradation of L1 over time

(Fernández-San Millán et al., 2008). In our study, bands at 45 kDa are most prominent in pL1Δ22 lines with

cytoplasmic L1 and less prominent for plastidic L1. This might result from organelle-specific proteases

(Coates at al., 2022).

In Physcomitrella, combining the full-length L1 with a CTP significantly increases L1 yield as

compared to full length L1 or L1 without NLS, which is in line with findings in tobacco (Maclean et al.,

2007; Zahin et al., 2016). In our study, L1 yields above 0.3 mg / g FW were achieved in the photobioreactor.

In tobacco, yields of 0.1 – 1.2 mg / g FW (Maclean et al., 2007; Pineo et al., 2013; Zahin et al., 2016;

Hitzeroth et al., 2018) and up to 3 mg / g FW in transplastomic lines have been reported (Fernández-San

Millán et al., 2008). The success of targeting L1 to chloroplasts over localization to other organelles may

have several reasons. One is that plastid L1 may be protected from proteases absent in chloroplasts (van

Wijk, 2024). Besides, CTPs also act on the transcriptional level. Upon expression in tobacco, chimeric L1/L2

transcripts containing a CTP were more abundant than transcripts lacking a CTP, and consequently yielded

more protein (Hitzeroth et al., 2018). So far, it is unclear whether this difference is due to enhanced

transcription, increased mRNA half-life or changes in mRNA secondary structure.

For purification of L1 from Physcomitrella extracts, we employed a combination of AMS

precipitation and CaEx chromatography. We found 45% AMS to be sufficient to efficiently precipitate L1.

Studies from yeast (Park et al., 2008; Kim et al., 2010) and tobacco (Zahin et al., 2016) also used 45% AMS,

whereas sometimes 50% AMS was used (Kim et al., 2018). CaEx chromatography showed that a large part

of L1 does not bind to the column. Likewise, Zahin et al. (2016) reported tobacco-produced L1 to bind

poorly to CaEx columns. Kim et al. (2010) purified yeast-produced L1 by either CaEx or heparin

chromatography. CaEx chromatography showed unbound L1 in the flow through, as observed in our

study, whereas no L1 was found in the flow through employing heparin chromatography. Pineo et al.

(2013) produced L1/L2 chimaeras in tobacco and experienced loss of L1 and insufficient removal of

contaminating proteins using heparin chromatography. Heparin is a highly sulfated form of heparan

sulfate, a proteoglycan found on host cell surfaces and a requirement for HPV virion attachment and

infection (Knappe et al., 2007). However, a prerequisite for heparin binding is the assembly of L1 to

capsomers or VLPs (Rommel et al., 2005), meaning monomers and not properly folded L1 are lost. Overall,

chromatography yields for VLPs differ and are influenced by different factors, including the production

host, the used materials and buffers, and purification prior to chromatography, making it difficult to

compare purification strategies across publications. Here, we achieved to remove the bulk of cellular

proteins from moss extracts, but experienced a substantial loss of L1 in the flow through. To address this,

further optimization of purification might be achieved by combinations with other methods such as

heparin chromatography. Our dot blot analysis confirmed the presence of intact capsomers or VLPs in

extracts and after AMS precipitation, suggesting heparin chromatography as an option for the second step

in purification.

TEM analysis of purified and reassembled CTP-pL1 samples shows VLPs of approx. 60 nm in size.

These nanoparticles assembled from capsomers as a result of the reassembly treatment. The absence of

VLPs in the elution fractions does not necessarily indicate that dis- and reassembly treatment is a

prerequisite for VLP formation from moss-produced L1, but rather that the buffer conditions did not favor

assembly after elution. To the best of our knowledge, this is the first report of VLPs produced in a non-

vascular plant and of *in vitro* reassembly of HPV-16 L1 VLPs from capsomers produced in any plant system.

The VLPs are heterogenous in shape and size. The same is true for yeast-derived VLPs after reassembly,

whereas untreated VLPs were not heterogenous. This is surprising, since controlled dis- and reassembly

of VLPs can enhance stability and uniformity (McCarthy et al., 1998; Mach et al., 2006; Shi et al., 2007).

Besides high ionic strength and low pH buffer, inter-pentameric disulfide bridges positively influence VLP

stability and uniformity (Buck et al., 2005). This could be achieved through prolonged incubation of cleared

cell lysates before purification or addition of oxidizing agents, which accelerate disulfide bridge formation.

However, prolonged incubation causes aggregation of HPV-16 L1 VLPs, as observed here, but could be
addressed by improved formulation (Shi et al., 2005).

While the moss plant Physcomitrella is an established production host for biopharmaceuticals, we
aim to expand its application to other markets such as the cosmetics industry (Munoz et al., 2024) and
material sciences at the interface with biomedicine (Milferstaedt et al., 2025; Ramezaniaghdam et al.,
2025). In this context, the present study serves as a proof-of-concept for the production of a relatively
simple but medicinally relevant VLP in Physcomitrella. Future efforts may focus on more complex eVLPs
that require proper N-glycosylation or complex folding and assembly; a specialty niche well suited for the
moss-based expression platform.

Acknowledgements

We thank the Life Imaging Centre (LIC) of the University of Freiburg for support with confocal microscopy.
We are grateful for technical assistance by Agnes Novakovic, Britta Rothgänger, Christine Glockner and
Richard Haas and to Anne Katrin Prowse for language editing.

Author contributions

PAN: Conceptualization, Formal analysis, Funding acquisition, Investigation, Methodology, Visualization,
Writing – original draft, Writing – review & editing
MCW: Investigation, Methodology, Writing – review & editing
SC: Investigation, Methodology, Writing – review & editing
CET: Investigation, Methodology, Formal analysis, Writing – review & editing
MRF: Investigation, Formal analysis, Writing – review & editing
SNWH: Investigation, Formal analysis, Writing – review & editing
JP: Writing – original draft, Writing – review & editing
ELD: Writing – original draft, Writing – review & editing
HTS: Conceptualization, Resources, Writing – review & editing
RR: Supervision, Resources, Writing – original draft, Writing – review & editing

Conflict of interest disclosure

715   RR is a founder of Greenovation Biotech, now Eleva Biologics, an inventor of the moss bioreactor, and
716   currently serves as a member of Eleva's advisory board. HTS is a co-founder of Tailwind Biotech, a
717   company designing DNA regulatory parts, and currently serves as a member of Tailwind's executive board.
718   All other authors declare no conflict of interest.

Data availability statement

Nucleotide sequence data was submitted to GenBank under accession numbers PV029542 (pL1) and
PV068547(CTPc5). Moss lines and vectors were submitted to the International Moss Stock Centre (IMSC)
under accession numbers 40963 (pL1 5A), 40964 (pL1 23A), 40965 (pL1 47B), 40966 (pL1Δ22 8A), 40967
(pL1Δ22 11B), 40968 (pL1Δ22 102B), 40969 (CTP-pL1 30), 40970 (CTP-pL1 103), 40971 (CTP-pL1 143), 1840
(pAct5_linker_eGFP_35ST_hpt), 2038 (pAct5_CTP_linker_eGFP_35ST_hpt), 1193
(pAct5_Citrine_nosT_hpt), 2249 (pAct5_pL1-Citrine_nosT_hpt), 2250 (pAct5_pL1Δ22-Citrine_nosT_hpt),
2251 (pAct5_CTP-pL1-Citrine_nosT_hpt), 1997 (pAct5_pL1_35ST_hpt), 1998 (pAct5_pL1Δ22_35ST_hpt)
and 2048 (pAct5_CTP-pL1_35ST_hpt).

Funding statement

This work was supported by Maria-Sklodowska-Curie Actions Innovative Training Network under the
Horizon 2020 program under Grant Agreement No. 765115 (MossTech), by the German Research
Foundation DFG under Germany's Excellence Strategy (CIBSS – EXC-2189 – Project ID 390939984), and
the International Network Program by the Danish Agency for Higher Education and Science (grant number
1113-00016B). The TEM (Hitachi HT7800) was funded by a DFG grant (project number 426849454) and is
operated by the University of Freiburg, Faculty of Biology, as a partner unit within the Microscopy and
Image Analysis Platform MIAP and the Life Imaging Center LIC, Freiburg. PAN gratefully acknowledges
financial support by Studienstiftung des deutschen Volkes.

Short legends for Supporting Information

740		
741	Supplementary Figure S1	**Vector maps**
742	Supplementary Figure S2	**Generation of Physcomitrella-optimized (p)L1**
743	Supplementary Figure S3	**Characterization of CTPs by analysis of subcellular eGFP localization**
744	Supplementary Figure S4	**Analysis of CDS integrity in L1-producing lines**
745	Supplementary Figure S5	**Analysis of the effect of reducing agents on L1 band patterning**
746	Supplementary Figure S6	**Comparison of biomass accumulation rates**
747	Supplementary Figure S7	**Ammonium sulfate precipitation**
748	Supplementary Figure S8	**Chromatogram of Cation Exchange purification**
749	Supplementary Table S1	**List of primers**
750	Supplementary Table S2	**Predicted chloroplast transit peptides for six selected proteins in**
751		**Physcomitrella**

References

Abel WO, Knebel W, Koop HU, Marienfeld JR, Quader H, Reski R, Schnepf E, Sporlein B. A cytokinin-sensitive mutant of the moss, *Physcomitrella patens*, defective in chloroplast division. Protoplasma (1989) 152:1-13. doi: 10.1007/BF01354234

Adams A, Hendrikse M, Rybicki EP, Hitzeroth II. Optimal size of DNA encapsidated by plant produced human papillomavirus pseudovirions. Virology (2023) 580:88–97. doi: 10.1016/j.virol.2023.02.003

Armenteros JJA, Salvatore M, Emanuelsson O, Wither O, von Heijne G, Elofsson A, Nielsen H. Detecting sequence signals in targeting peptides using deep learning. Life Sci All (2019) 2:e201900429. doi: 10.26508/lsa.201900429

Baker TS, Newcomb WW, Olson NH, Cowsert LM, Olson C, Brown JC. Structures of bovine and human papillomaviruses. Analysis by cryoelectron microscopy and three-dimensional image reconstruction. Biophys J (1991) 60:1445–1456. doi: 10.1016/S0006-3495(91)82181-6

Benvenuto E, Broer I, D'Aoust MA, Hitzeroth I, Hundleby P, Menassa R, Oksman-Caldentey KM, Peyret H, Salgueiro S, Saxena P, Stander J, Warzecha H, Ma J. Plant molecular farming in the wake of the closure of Medicago Inc. Nature Biotech (2023) 41:893–894. doi: 10.1038/s41587-023-01812-w

Biemelt S, Sonnewald U, Galmbacher P, Willmitzer L, Müller M. Production of human papillomavirus type 16 virus-like particles in transgenic plants. J Virol (2003) 77:9211–9220. doi: 10.1128/JVI.77.17.9211-9220.2003

Bissa M, Zanotto C, Pacchioni S, Volonté L, Venuti A, Lembo D, De Giuli Morghen C, Radaelli A. The L1 protein of human papilloma virus 16 expressed by a fowlpox virus recombinant can assemble into virus-like particles in mammalian cell lines but elicits a non-neutralising humoral response. Antiviral Res (2015) 116:67–75. doi: 10.1016/j.antiviral.2015.01.012

Bohlender LL, Parsons J, Hoernstein SNW, Rempfer C, Ruiz-Molina N, Lorenz T, Rodríguez Jahnke F, Figl R,

Fode B, Altmann F, Reski R, Decker EL. Stable protein sialylation in Physcomitrella. Front Plant Sci (2020)

11:610032. doi: 10.3389/FPLS.2020.610032

Bohlender LL, Parsons J, Hoernstein SNW, Bangert N, Rodriguez-Jahnke F, Reski R, Decker EL. Unexpected

arabinosylation after humanization of plant protein N-glycosylation. Front Bioeng Biotech (2022)

10:838365. https://doi.org/10.3389/fbioe.2022.838365

Buck CB, Thompson CD, Pang YYS, Lowy DR, Schiller JT. Maturation of papillomavirus capsids. J Virol (2005)

79:2839–2846. doi: 10.1128/JVI.79.5.2839-2846.2005

Buck CB, Day PM, Trus BL. The papillomavirus major capsid protein L1. Virol (2013) 445:169-174. doi:

10.1016/j.virol.2013.05.038

Burns CC, Diop OM, Sutter RW, Kew OM. Vaccine-derived polioviruses. J Infect Dis (2014) 210:S283-S293.

doi: 10.1093/INFDIS/JIU295

Büttner-Mainik A, Parsons J, Jérôme H, Hartmann A, Lamer S, Schaaf A, Schlosser A, Zipfel PF, Reski R,

Decker EL. Production of biologically active recombinant human factor H in Physcomitrella. Plant Biotech

J (2011) 9:373–383. doi: 10.1111/J.1467-7652.2010.00552.X

Castells-Graells R, Lomonossoff GP. Plant-based production can result in covalent cross-linking of proteins.

Plant Biotech J (2021) 19:1095–1097. doi: 10.1111/pbi.13598

Cerqueira C, Schiller JT. Papillomavirus assembly: An overview and perspectives. Virus Res (2017)

231:103–107. doi: 10.1016/j.virusres.2016.11.010

Chi H, Pepper M, Thomas PG. Principles and therapeutics of adaptive immunity. Cell (2024) 187:2052-

2078. doi: doi.org/10.1016/j.cell.2024.03.037

Coates RJ, Young MT, Scofield S. Optimising expression and extraction of recombinant proteins in plants.

Front Plant Sci (2022) 13:1074531. doi: 10.3389/fpls.2022.1074531

Cubas R, Zhang S, Li M, Chen C, Yao Q. Chimeric Trop2 virus-like particles: A potential immunotherapeutic

approach against pancreatic cancer. J Immunoth (2011) 34:251–263. doi: CJI.0B013E318209EE72

Dai X, Zhao PX. psRNATarget: a plant small RNA target analysis server. Nucl Acids Res (2011) 39:W155-

W159. doi: 10.1093/nar/gkr319

Dai X, Zhuang, Z, Zhao, PX. psRNATarget: a plant small RNA target analysis server (2017 release). Nucl

Acids Res (2018) 46:W49–W54. doi: 10.1093/NAR/GKY316

de Martel C, Georges D, Bray F, Ferlay J, Clifford GM. Global burden of cancer attributable to infections in

2018: a worldwide incidence analysis. Lancet Glob Health (2020) 8:e180–e190. doi: 10.1016/S2214-

109X(19)30488-7

Decker EL, Reski R. Mosses in biotechnology. Curr Op Biotech (2020) 61:21–27. doi:

10.1016/j.copbio.2019.09.021

Decker EL, Wiedemann G, Reski R. Gene targeting for precision glyco-engineering: Production of

biopharmaceuticals devoid of plant-typical glycosylation in moss bioreactors. Meth Mol Biol (2015)

1321:213-224. doi: 10.1007/978-1-4939-2760-9_15

Decker EL, Parsons J, Reski R. Glyco-engineering for biopharmaceutical production in moss bioreactors.

Front Plant Sci (2014) 5:346. doi: 10.3389/fpls.2014.00346

Eleva (2025). Eleva and 3PBIOVAN sign strategic GMP manufacturing and technology alliance to boost

capacity for Eleva's platform and pipeline. https://elevabiologics.com/eleva-and-3pbiovian-sign-

strategic-alliance [Accessed May 21, 2025].

Fernández-San Millán A, Ortigosa SM, Hervás-Stubbs S, Corral-Martínez P, Seguí-Simarro JM, Gaétan J,

Coursaget P, Veramendi J. Human papillomavirus L1 protein expressed in tobacco chloroplasts self-

assembles into virus-like particles that are highly immunogenic. Plant Biotech J (2008) 6:427–441. doi:

10.1111/j.1467-7652.2008.00338.x

Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO. Enzymatic assembly of DNA molecules

up to several hundred kilobases. Nature Met (2009) 6:343–345. doi: 10.1038/nmeth.1318

Gitzinger M, Parsons J, Reski R, Fussenegger M. Functional cross-kingdom conservation of mammalian and

moss (*Physcomitrella patens*) transcription, translation and secretion machineries. Plant Biotech J (2009)

7:73–86. doi: 10.1111/J.1467-7652.2008.00376.X

Gremillon L, Kiessling J, Hause B, Decker EL, Reski R, Sarnighausen E. Filamentous temperature-sensitive

Z (FtsZ) isoforms specifically interact in the chloroplasts and in the cytosol of *Physcomitrella patens*. New

Phyt (2007) 176:299–310. doi: 10.1111/j.1469-8137.2007.02169.x

Gupta R, Arora K, Roy SS, Joseph A, Rastogi R, Arora NM, Kundu PK. Platforms, advances, and technical

challenges in virus-like particles-based vaccines. Front Immun (2023) 14:1123805. doi:

10.3389/FIMMU.2023.1123805

Hasan SS, Sevvana M, Kuhn RJ, Rossmann MG. Structural biology of Zika virus and other flaviviruses. Nat

Struc Mol Biol (2018) 25:13–20. doi: 10.1038/s41594-017-0010-8

Hector M, Behnke V, Dabrowska-Schlepp, Busch A, Schaaf A, Langmann T, Wolf A. Moss-derived human

complement factor H modulates retinal immune response and attenuates retinal degeneration. J

Neuroinflam (2025) 22:104. doi: 10.1186/s12974-025-03418-2

Hennermann JB, Arash-Kaps L, Fekete G, Schaaf A, Busch A, Frischmuth T. Pharmacokinetics,

pharmacodynamics, and safety of moss-aGalactosidase A in patients with Fabry disease. J Inher Metab

Dis (2019) 42:527–533. doi: 10.1002/JIMD.12052

Hitzeroth II, Chabeda A, Whitehead MP, Graf M, Rybicki EP. Optimizing a human papillomavirus type 16

L1-based chimaeric gene for expression in plants. Front Bioeng Biotech (2018) 6:101. doi:

10.3389/fbioe.2018.00101

Hoernstein SNW, Oezdemir B, van Gessel N, Miniera AA, Rogalla von Bieberstein B, Nilges L, Schweikert

Farinha J, Komoll R, Glauz S, Weckerle T, Scherzinger F, Rodriguez-Franco M, Mueller-Schuessele S, Reski

R. A deeply conserved protease, acylamino acid-releasing enzyme (AARE), acts in ageing in Physcomitrella

and Arabidopsis. Comm Biol (2023) 6:61. doi: 10.1038/s42003-023-04428-7

Hoernstein SNW, Schlosser A, Fiedler K, van Gessel N, Igloi GL, Lang D, Reski R. A snapshot of the

Physcomitrella N-terminome reveals N-terminal methylation of organellar proteins. Plant Cell Rep (2024)

43:250. doi: 10.1007/s00299-024-03329-1

Hohe A, Reski R. Optimisation of a bioreactor culture of the moss *Physcomitrella patens* for mass

production of protoplasts. Plant Sci (2002) 163: 69-74. doi: 10.1016/S0168-9452(02)00059-6

Hohe A, Decker EL, Gorr G, Schween G, Reski R. Tight control of growth and cell differentiation in

photoautotrophically growing moss (*Physcomitrella patens*) bioreactor cultures. Plant Cell Rep (2002)

20:1135-1140. doi: 10.1007/s00299-002-0463-y

Hohe A, Egener T, Lucht JM, Holtorf H, Reinhard C, Schween G, Reski R. An improved and highly

standardized transformation procedure allows efficient production of single and multiple targeted gene-

knockouts in a moss, *Physcomitrella patens*. Curr Genet (2004) 44:339-347. doi: 10.1007/s00294-003-

0458-4

Jones-Rhoades MW, Bartel DP, Bartel B. MicroRNAs and their regulatory roles in plants. Ann Rev Plant

Biol (2006) 57:19-53. doi: 10.1146/annurev.arplant.57.032905.105218

Kerr H, Herbert AP, Makou E, Abramczyk D, Malik TH, Lomax-Browne H, Yang Y, Pappworth IY, Denton H,

Richards A, Marchbank KJ, Pickering MC, Barlow PN. Murine factor H co-produced in yeast with protein

disulfide isomerase ameliorated C3 dysregulation in factor H-deficient mice. Front Immun (2021)

12:681098. https://doi.org/10.3389/FIMMU.2021.681098

Khraiwesh B, Arif MA, Seumel GI, Ossowski S, Weigel D, Reski R, Frank W. Transcriptional control of gene

expression by microRNAs. Cell (2010) 140:111-122. doi: 10.1016/j.cell.2009.12.023

Kim SN, Jeong HS, Park SN, Kim HJ. Purification and immunogenicity study of human papillomavirus type
16 L1 protein in *Saccharomyces cerevisiae*. J Virol Met (2007) 139:24-30. doi:
10.1016/j.jviromet.2006.09.004

Kim HJ, Kim SY, Lim SJ, Kim JY, Lee SJ, Kim HJ. One-step chromatographic purification of human
papillomavirus type 16 L1 protein from *Saccharomyces cerevisiae*. Prot Expr Pur (2010) 70:68–74. doi:
10.1016/j.pep.2009.08.005

Kim HS, Jeon JH, Lee KJ, Ko K. N-glycosylation modification of plant-derived virus-like particles: An
application in vaccines. BioMed Res Int (2014) 2014:249519. doi: 10.1155/2014/249519

Kim HJ, Cho SY, Park MH, Kim HJ. Comparison of the size distributions and immunogenicity of human
papillomavirus type 16 L1 virus-like particles produced in insect and yeast cells. Arch Pharm Res (2018)
41:544–553. doi: 10.1007/s12272-018-1024-4

Kirnbauer R, Booy F, Cheng N, Lowy DR, Schiller JT. Papillomavirus L1 major capsid protein self-assembles
into virus-like particles that are highly immunogenic. Proc Nat Acad Sci USA (1992) 89:12180–84. doi:
10.1073/pnas.89.24.12180

Knappe M, Bodevin S, Selinka HC, Spillmann D, Streeck RE, Chen XS, Lindahl U, Sapp M. Surface-exposed
amino acid residues of HPV16 L1 protein mediating interaction with cell surface heparan sulfate. J Biol
Chem (2007) 282:27913–22. https://doi.org/10.1074/jbc.M705127200

Koprivova A, Stemmer C, Altmann F, Hoffmann A, Kopriva S, Gorr G, Reski R, Decker EL. Targeted
knockouts of *Physcomitrella* lacking plant-specific immunogenic N-glycans. Plant Biotech J (2004) 2:517–
523. doi: 10.1111/J.1467-7652.2004.00100.X

Lamprecht RL, Kennedy P, Huddy SM, Bethke S, Hendrikse M, Hitzeroth II, Rybicki EP. Production of human
papillomavirus pseudovirions in plants and their use in pseudovirion-based neutralisation assays in
mammalian cells. Sci Rep (2016) 6:20431. doi: 10.1038/srep20431

Leder C, Kleinschmidt JA, Wiethe C, Müller M. Enhancement of capsid gene expression: Preparing the

human papillomavirus type 16 major structural gene L1 for DNA vaccination purposes. J Virol (2001)

75:9201–09. doi: 10.1128/JVI.75.19.9201-9209.2001

Lenzi P, Scotti N, Alagna F, Tornesello ML, Pompa A, Vitale A, De Stradis A, Monti L, Grillo S, Buonaguro

FM, Maliga P, Cardi T. Translational fusion of chloroplast-expressed human papillomavirus type 16 L1

capsid protein enhances antigen accumulation in transplastomic tobacco. Transgen Res (2008) 17:1091–

1102. doi: 10.1007/s11248-008-9186-3

Lizotte PH, Wen AM, Sheen MR, Fields J, Rojanasopondist P, Steinmetz NF, Fiering S. *In situ* vaccination

with Cowpea Mosaic Virus nanoparticles suppresses metastatic cancer. Nat Nanotech (2015) 11:295–303.

doi: 10.1038/nnano.2015.292

Ljubojevic S, Skerlev M. HPV-associated diseases. Clin Derm (2014) 32:227-234. doi:

10.1016/j.clindermatol.2013.08.007

Lu B, Lim JM, Yu B, Song S, Neeli P, Sobhani N, K P, Bonam SR, Kurapati R, Zheng J, Chai D. The next-

generation DNA vaccine platforms and delivery systems: advances, challenges and prospects. Front

Immunol (2024) 15:1332939. doi: 10.3389/FIMMU.2024.1332939

Mach H, Volkin DB, Troutman RD, Wang B, Luo Z, Jansen KU, Shi L. Disassembly and reassembly of yeast-

derived recombinant human papillomavirus virus-like particles (HPV VLPs). J Pharm Sci (2006) 95:2195–

2206. doi: org/10.1002/jps.20696

Maclean J, Koekemoer M, Olivier AJ, Stewart D, Hitzeroth II, Rademacher T, Fischer R, Williamson AL,

Rybicki EP. Optimization of human papillomavirus type 16 (HPV-16) L1 expression in plants: comparison

of the suitability of different HPV-16 L1 gene variants and different cell-compartment localization. J Gen

Virol (2007) 88:1460–69. doi: org/10.1099/vir.0.82718-0

Marsian J, Lomonossoff GP. Molecular pharming – VLPs made in plants. Curr Op Biotech (2016) 37:201-

206. doi: 10.1016/j.copbio.2015.12.007

Matic S, Rinaldi R, Masenga V, Noris E. Efficient production of chimeric Human papillomavirus 16 L1

protein bearing the M2e influenza epitope in *Nicotiana benthamiana* plants. BMC Biotech (2011) 11:106.

doi: 10.1186/1472-6750-11-106

McCarthy MP, White WI, Palmer-Hill F, Koenig S, Suzich JA. Quantitative disassembly and reassembly of

human papillomavirus type 11 virus like particles *in vitro*. J Virol (1998) 72:32–41. doi:

10.1128/JVI.72.1.32-41.1998

Michelfelder S, Parsons J, Bohlender LL, Hoernstein SNW, Niederkrüger H, Busch A, Krieghoff N, Koch J,

Fode B, Schaaf A, Frischmuth T, Pohl M, Zipfel PF, Reski R, Decker EL, Häffner K. Moss-produced,

glycosylation-optimized human factor H for therapeutic application in complement disorders. J Am Soc

Nephrol (2017) 28:1462–74. doi: org/10.1681/ASN.2015070745

Michelfelder S, Fischer F, Wäldin A, Hörle KV, Pohl M, Parsons J, Reski R, Decker EL, Zipfel PF, Skerka C,

Häffner K. The MFHR1 fusion protein is a novel synthetic multitarget complement inhibitor with

therapeutic potential. J Am Soc Nephrol (2018) 29:1141–53. doi: 10.1681/ASN.2017070738

Milferstaedt SWL, Joest M, Bohlender LL, Hoernstein SNW, Özdemir B, Decker EL, van der Does C, Reski

R. Differential GTP-dependent in-vitro polymerization of recombinant Physcomitrella FtsZ proteins. Sci

Rep (2025) 15:3095. doi: 10.1038/s41598-024-85077-6

Mitsubishi Chemical Group Corporation. Overseas consolidated subsidiary, Medicago to cease operations

(2023). https://www.mcgc.com/english/news_release/pdf/01468/01708.pdf [Accessed May 21, 2018].

Munoz C, Schröder K, Henes B, Hubert J, Leblond S, Poigny S, Reski R, Wandrey F. Phytochemical

exploration of ceruchinol in moss: A multidisciplinary study in biotechnological cultivation of

Physcomitrium patens (Hedw.) Mitt. Appl Sci (2024) 14:1274. doi: 10.3390/app14031274

Mori S, Ozaki S, Yasugi T, Yoshikawa H, Taketani Y, Kanda. Inhibitory cis-element-mediated decay of

human papillomavirus type 16 L1-transcript in undifferentiated cells. Mol Cell Biochem (2006) 288:47–57.

doi: 10.1007/s11010-006-9117-7

1002

1003   Mortola E, Roy P. Efficient assembly and release of SARS coronavirus-like particles by a heterologous

1004   expression system. FEBS Lett (2004) 576:174–178. doi: 10.1016/J.FEBSLET.2004.09.009

1005

1006   Murén E, Nilsson A, Ulfstedt M, Johansson M, Ronne H. Rescue and characterization of episomally

1007   replicating DNA from the moss *Physcomitrella*. Proc Nat Acad Sci USA (2009) 106:19444–49. doi:

1008   10.1073/pnas.0908037106

1009

1010   Muthamilselvan T, Khan MRI, Hwang I. Assembly of human papillomavirus 16 L1 protein in *Nicotiana*

1011   *benthamiana* chloroplasts into highly immunogenic Virus-Like Particles. J Plant Biol (2023) 66:331–340.

1012   doi: 10.1007/s12374-023-09393-6

1013

1014   Naupu PN, van Zyl AR, Rybicki EP, Hitzeroth II. Immunogenicity of plant-produced human papillomavirus

1015   (HPV) virus-like particles (VLPs). Vacc (2020) 8:740. doi: 10.3390/VACCINES8040740

1016

1017   Niederkrüger H, Busch A, Dabrowska-Schlepp P, Krieghoff N, Schaaf A, Frischmuth T. „Single-use

1018   processing as a safe and convenient way to develop and manufacture moss-derived biopharmaceuticals".

1019   In: Eibl R, Eibl D, editors. Single-use technology in biopharmaceutical manufacture, 2nd edition. Hoboken

1020   NJ: Wiley (2019). p.311-318. https://doi.org/10.1002/9781119477891.ch28

1021

1022   Nooraei S, Bahrulolum H, Hoseini ZS, Katalani C, Hajizade A, Easton AJ, Ahmadian G. Virus-like particles:

1023   preparation, immunogenicity and their roles as nanovaccines and drug nanocarriers. J Nanobiotech (2021)

1024   19:59. doi: 10.1186/s12951-021-00806-7

1025

1026   Orellana-Escobedo L, Rosales-Mendoza S, Romero-Maldona A, Parsons J, Decker EL, Monreal-Escalante E,

1027   Moreno-Fierros L, Reski R. An Env-derived multi-epitope HIV chimeric protein produced in the moss

1028   *Physcomitrella patens* is immunogenic in mice. Plant Cell Rep (2015) 34:425-433. doi: 10.1007/s00299-

1029   014-1720-6

1030

1031   Park MA, Kim HJ, Kim HJ. Optimum conditions for production and purification of human papillomavirus

1032   type 16 L1 protein from *Saccharomyces cerevisiae*. Prot Expr Pur (2008) 59:175–181. doi:

1033   10.1016/j.pep.2008.01.021

1034

1035 Parsons J, Altmann F, Arrenberg CK, Koprivova A, Beike AK, Stemmer C, Gorr G, Reski R, Decker EL. Moss-
1036 based production of asialo-erythropoietin devoid of Lewis A and other plant-typical carbohydrate
1037 determinants. Plant Biotech J (2012) 10:851–861. doi: 10.1111/J.1467-7652.2012.00704.X

1038

1039 Peng W, Rayaprolu V, Parvate AD, Proner MF, Hui S, Parekh D, Shaffer K, Yu X, Saphire EO, Snijder J. Glycan
1040 shield of the ebolavirus envelop glycoprotein GP. Comm Biol (2022) 5:785. doi: 10.1038/s42003-022-
1041 03767-1

1042

1043 Pineo CB, Hitzeroth II, Rybicki EP. Immunogenic assessment of plant-produced human papillomavirus type
1044 16 L1/L2 chimaeras. Plant Biotech J (2013) 11:964–975. https://doi.org/10.1111/pbi.12089

1045

1046 Ramezaniaghdam M, Bohlender LL, Parsons J, Hoernstein SNW, Decker EL, Reski R. Recombinant
1047 production of spider silk protein in Physcomitrella photobioreactors. Plant Cell Rep (2025) 44:103. doi:
1048 10.1007/s00299-025-03485-y

1049

1050 Rempfer C, Wiedemann G, Schween G, Kerres KL, Lucht JM, Horres R, Decker EL, Reski R.
1051 Autopolyploidization affects transcript patterns and gene targeting frequencies in Physcomitrella. Plant
1052 Cell Rep (2022) 41:153-173. doi: 10.1007/s00299-021-02794-2

1053

1054 Reski R. Development, genetics and molecular biology of mosses. Bot Acta (1998) 111:1–15. doi:
1055 10.1111/j.1438-8677.1998.tb00670.x

1056

1057 Reski R. Quantitative moss cell biology. Curr Op Plant Biol (2018) 46:39–47. doi:
1058 10.1016/J.PBI.2018.07.005

1059

1060 Reski R, Abel WO. Induction of budding on chloronemata and caulonemata of the moss, *Physcomitrella*
1061 *patens*, using isopentenyladenine. Planta (1985) 165:354-358. doi: 1007/BF00392232

1062

1063 Reski R, Parsons J, Decker EL. Moss-made pharmaceuticals: from bench to bedside. Plant Biotech J (2015)
1064 13:1191–98. doi: 10.1111/pbi.12401

1065

1066   Reski R, Bae H, Simonsen HT. *Physcomitrella patens*, a versatile synthetic biology chassis. Plant Cell Rep
1067   (2018) 37:1409–17. doi: 10.1007/S00299-018-2293-6

1068

1069   Roden RBS, Stern PL. Opportunities and challenges for human papillomavirus vaccination in cancer. Nat
1070   Rev Cancer (2018) 18:240-254. doi: 10.1038/nrc.2018.13

1071

1072   Rommel O, Dillner J, Fligge C, Bergsdorf C, Wang X, Selinka HC, Sapp M. Heparan sulfate proteoglycans
1073   interact exclusively with conformationally intact HPV L1 assemblies: Basis for a virus-like particle ELISA. J
1074   Med Virol (2005) 75:114–121. doi: 10.1002/jmv.20245

1075

1076   Rosales-Mendoza S, Orellana-Escobedo L, Romero-Maldonado A, Decker EL, Reski R. The potential of
1077   *Physcomitrella patens* as a platform for the production of plant-based vaccines. Exp Rev Vacc (2014)
1078   13:203-212. doi: 10.1586/14760584.2014.872987

1079

1080   Ross TM, Mahmood K, Crevar CJ, Schneider-Ohrum K, Heaton PM, Bright RA. A trivalent virus-like particle
1081   vaccine elicits protective immune responses against seasonal influenza strains in mice and ferrets. PLOS
1082   ONE (2009) 4:e6032. doi: 10.1371/JOURNAL.PONE.0006032

1083

1084   Ruiz-Molina N, Parsons J, Müller M, Hoernstein SNW, Bohlender LL, Pumple S, Zipfel PF, Häffner K, Reski
1085   R, Decker EL. A synthetic protein as efficient multitarget regulator against complement over-activation.
1086   Comm Biol (2022a) 5:152. doi: 10.1038/s42003-022-03094-5

1087

1088   Ruiz-Molina N, Parsons J, Schroeder S, Posten C, Reski R, Decker EL. Process engineering of
1089   biopharmaceutical production in moss bioreactors *via* model-based description and evaluation of
1090   phytohormone impact. Front Bioeng Biotech (2022b) 10:837965. doi: 10.3389/fbioe.2022.837965

1091

1092   Rybicki EP. Plant molecular farming of virus-like nanoparticles as vaccines and reagents. WIREs Nanomed
1093   Nanobiotech (2020) 12:e1587. doi: 10.1002/wnan.1587

1094

1095   Sapp M, Fligge C, Petzak I, Robin Harris J, Streeck RE. Papillomavirus assembly requires trimerization of
1096   the major capsid protein by disulfides between two highly conserved cysteines. J Virol (1998) 72:6186-
1097   6189. doi: 10.1128/jvi.72.7.6186-6189.1998

1098

1099 Schmidt CQ, Slingsby FC, Richards A, Barlow PN. Production of biologically active complement factor H in

1100 therapeutically useful quantities. Prot Expr Pur (2011) 76:254–263. doi: 10.1016/J.PEP.2010.12.002

1101

1102 Schween G, Hohe A, Koprivova A, Reski R. Effects of nutrients, cell density and culture techniques on

1103 protoplast regeneration and early protonema development in a moss, *Physcomitrella patens*. J Plant Phys

1104 (2003) 160:209-212. doi: 10.1078/0176-1617-00855

1105

1106 Scotti N, Rybicki EP. Virus-like particles produced in plants as potential vaccines. Exp Rev Vacc (2013)

1107 12:211–224. doi: 10.1586/erv.12.147

1108

1109 Serradell MC, Rupil LL, Martino RA, Prucca CG, Carranza PG, Saura A, Fernández EA, Gargantini PR, Tenaglia

1110 AH, Petiti JP, Tonelli RR, Reinoso-Vizcaino N, Echenique J, Berod L, Piaggio E, Bellier B, Sparwasser T,

1111 Klatzmann D, Luján HD (2019) Efficient oral vaccination by bioengineering virus-like particles with

1112 protozoan surface proteins. Nat Comm (2019) 10:361. doi: 10.1038/s41467-018-08265-9

1113

1114 Shen JS, Busch A, Day TS, Meng XL, Yu CI, Dabrowska-Schlepp P, Fode B, Niederkrüger H, Forni S, Chen S,

1115 Schiffmann R, Frischmuth T, Schaaf A. Mannose receptor-mediated delivery of moss-made α-

1116 galactosidase A efficiently corrects enzyme deficiency in Fabry mice. J Inher Metab Dis (2016) 39:293–303.

1117 Doi: 10.1007/S10545-015-9886-9

1118

1119 Shi L, Sanyal G, Ni A, Luo Z, Doshna S, Wang B, Graham TL, Wang N, Volkin DB. Stabilization of human

1120 papillomavirus virus-like particles by non-ionic surfactants. J Pharm Sci (2005) 94:1538–51. doi:

1121 10.1002/jps.20377

1122

1123 Shi L, Sings HL, Bryan JT, Wang B, Wang Y, Mach H, Kosinski M, Washabaugh MW, Sitrin R, Barr E.

1124 GARDASIL®: Prophylactic human papillomavirus vaccine development – From bench top to bed-side. Clin

1125 Pharm Therap (2007) 81:259–264. doi: 10.1038/SJ.CLPT.6100055

1126

1127 Šmídková M, Müller M, Thönes N, Piuko K, Angelisová P, Velemínský J, Angelis KJ. Transient expression of

1128 human papillomavirus type 16 virus-like particles in tobacco and tomato using a tobacco rattle virus

1129 expression vector. Biol Plant (2010) 54:451–460. doi: 10.1007/s10535-010-0081-4

1130

1131 Song JM, Wang BZ, Park KM, van Rooijen N, Quan FS, Kim MC, Jin HT, Pekosz A, Compans RW, Kang SM.

1132 Influenza virus-like particles containing M2 induce broadly cross protective immunity. PLOS ONE (2011)

1133 6:e14538. doi: 10.1371/JOURNAL.PONE.0014538

1134

1135 Studentsov YY, Schiffman M, Strickler HD, Ho GYF, Pang YYs, Schiller J, Herrero R, Burk RD. Enhanced

1136 enzyme-linked immunosorbent assay for detection of antibodies to virus-like particles of human

1137 papillomavirus. J Clin Microbiol (2002) 40:1755-60. doi: 10.1128/JCM.40.5.1755-1760.2002

1138

1139 Strepp R, Scholz S, Kruse S, Speth V, Reski R. Plant nuclear gene knockout reveals a role in plastid division

1140 for the homolog of the bacterial cell division protein FtsZ, an ancestral tubulin. Proc Nat Acad Sci USA

1141 (1998) 95:4368–73. doi: 10.1073/PNAS.95.8.4368

1142

1143 Top O, Parsons J, Bohlender LL, Michelfelder S, Kopp P, Busch-Steenberg C, Hoernstein SNW, Zipfel PF,

1144 Häffner K, Reski R, Decker EL. Recombinant production of MFHR1, a novel synthetic multitarget

1145 complement inhibitor, in moss bioreactors. Front Plant Sci (2019) 10:260. doi: 10.3389/FPLS.2019.00260

1146

1147 Top O, Milferstaedt SWL, van Gessel N, Hoernstein SNW, Özdemir B, Decker EL, Reski R. Expression of a

1148 human cDNA in moss results in spliced mRNAs and fragmentary protein isoforms. Comm Biol (2021) 4:964.

1149 doi: 10.1038/s42003-021-02486-3

1150

1151 Torresi J. The rationale for a preventative HCV virus-like particle (VLP) vaccine. Front Microbiol (2017)

1152 8:2163. doi: 10.3389/fmicb.2017.02163

1153

1154 van Wijk KJ. Intra-chloroplast proteases: A holistic network view of chloroplast proteolysis. Plant Cell

1155 (2024) 36:3116–30. https://doi.org/10.1093/plcell/koae178

1156

1157 Varsani A, Williamson AL, Rose RC, Jaffer M, Rybicki EP. Expression of Human papillomavirus type 16

1158 major capsid protein in transgenic *Nicotiana tabacum* cv. Xanthi. Arch Virol (2003) 148:1771–86. doi:

1159 10.1007/s00705-003-0119-4

1160

1161 Weise A, Rodriguez-Franco M, Timm B, Hermann M, Link S, Jost W, Gorr G. Use of *Physcomitrella patens*

1162 actin 5' regions for high transgene expression: importance of 5' introns. Appl Microbiol Biotech (2006)

1163 70:337–345. doi: 10.1007/s00253-005-0087-6

1164

1165 Welsch S, Müller B, Kräusslich HG. More than one door - Budding of enveloped viruses through cellular

1166 membranes. FEBS Lett (2007) 581:2089–97. doi: 10.1016/J.FEBSLET.2007.03.060

1167

1168 Wiedemann G, van Gessel N, Köchl F, Hunn L, Schulze K, Maloukh L, Nogué F, Decker EL, Hartung F, Reski

1169 R. RecQ helicases function in development, DNA repair, and gene targeting in *Physcomitrella patens*. Plant

1170 Cell (2018) 30:717–736. doi: 10.1105/tpc.17.00632

1171

1172 World Health Organization. Considerations for human papillomavirus (HPV) vaccine product choice

1173 (2024). https://iris.who.int/bitstream/handle/10665/376546/9789240089167-eng.pdf [Accessed May

1174 21, 2025].

1175

1176 Wu M, Huang H, Tang Y, Ren X, Jiang X, Tian M, Li W. Unveiling the multifaceted realm of human

1177 papillomavirus: a comprehensive exploration of biology, interactions, and advances in cancer

1178 management. Front Immunol (2024) 15:1430544. doi: 10.3389/fimmu.2024.1430544

1179

1180 Zahin M, Joh J, Khanal S, Husk A, Mason H, Warzecha H, Ghim SJ, Miller DM, Matoba N, Jenson AB. Scalable

1181 production of HPV16 L1 protein and VLPs from tobacco leaves. PLOS ONE (2016) 11:e 0160995. doi:

1182 10.1371/JOURNAL.PONE.0160995

1183

1184 Zhang S, Yong LK, Li D, Cubas R, Chen C, Yao Q. Mesothelin virus-like particle immunization controls

1185 pancreatic cancer growth through CD8+ T cell induction and reduction in the frequency of CD4+ foxp3+

1186 ICOS- regulatory T cells. PLOS ONE (2013) 8:e68303 doi: 10.1371/JOURNAL.PONE.0068303

1187

1188 Zhao X, Schwartz S. Inhibition of HPV-16 L1 expression from L1 cDNAs correlates with the presence of

1189 hnRNP A1 binding sites in the L1 coding region. Virus Genes (2008) 36:45–53. doi: 10.1007/s11262-007-

1190 0174-0

1191

1192   Zlotnick A, Mukhopadhyay S. Virus assembly, allostery and antivirals. Trends Microbiol (2011) 19:14–

1193   23.doi: 10.1016/J.TIM.2010.11.003

1194

1195   zur Hausen H (2002) Papillomaviruses and cancer: From basic studies to clinical application. Nat Rev

1196   Cancer (2002) 2:342-350. doi: 10.1038/nrc798

1197

1198   zur Hausen H, de Villiers EM, Gissmann L. Papillomavirus infections and human genital cancer. Gyn Oncol

1199   (1081) 12:S124–S128. doi 10.1016/0090-8258(81)90067-6

Supplementary Information

Supplementary Figure S1 – Vector maps

List of all vectors used in this study. Vectors **a** & **b** were used for the characterization of chloroplast transit

peptides (CTPs). Vector **a** served as a control and was used as a cloning template for inserting different

CTP candidates. Vectors **c**, **d**, **e** & **f** were used for the analysis of L1 subcellular targeting. Vector **c** served

as a control and was used as a cloning template for inserting either pL1, pL1Δ22 or CTP-pL1 CDS (**d**, **e** & **f**,

respectively). Vectors **g**, **h** & **i** were used for generation of stable transgenic Physcomitrella lines. The

vectors contain either the full length pL1 including the native nuclear localization signal (NLS) (**g**), a

truncated version lacking the NLS (**h**), or the full-length version including the native NLS and an N-terminal

chloroplast transit peptide (**i**). All vectors contain the PpActin5 promoter (yellow), the CaMV 35S

terminator (grey), and a hygromycin (hpt)(turquoise) resistance cassette for selection. Other elements are

the GGGGGA linker (red), CTP (orange), NLS (blue) and eGFP or Citrine (green). Genetic elements are not

depicted true to scale. Vector sizes: 7260 bp (**a**), 7461 bp (**b**), 7299 bp (**c**), 8802 bp (**d**), 8736 bp (**e**), 9021

bp (**f**), 7998 bp (**g**), 7932 bp (**h**), 8217 bp (**i**).

Supplementary Figure S2 - Generation of Physcomitrella-optimized (p)L1

HPV-16 hL1

ATGTCCCTGTGGCTGCCCAGCGAGGCCACCGTGTACCTGCCCCCCGTGCCCGTGAGCAAGGTGGTGAGCACCGATGAGTACGTGGCCCGGA

CCAACATCTACTACCACGCCGGCACCTCCAGACTGCTGGCCGTGGGCCACCCCTACTTCCCCATCAAGAAGCCCAACAACAACAAGATCCTGG

TGCCCAAGGTGAGCGGCCTGCAATACCGGGTGTTCAGAATCCACCTGCCCGACCCCAATAAGTTCGGCTTCCCCGACACCAGCTTCTACAACC

CCGACACCCAGAGACTGGTGTGGGCCTGCGTGGGCGTGGAGGTGGGCAGAGGCCAGCCTCTGGGCGTGGGCATCAGCGGCCACCCTCTGC

TGAACAAGCTGGACGACACCGAGAACGCCAGCGCCTACGCCGCCAACGCCGGCGTGGATAACAGAGAATGCATCAGCATGGACTACAAGCA

GACCCAGCTGTGCCTCATCGGCTGCAAGCCCCCCATCGGCGAGCACTGGGGCAAGGGCAGCCCCTGCACCAACGTGGCCGTGAATCCTGGC

GACTGTCCTCCCCTGGAACTCATCAACACCGTGATCCAGGACGGCGACATGGTGGACACCGGCTTCGGCGCCATGGACTTCACCACCCTCCAG

GCCAATAAGAGCGAGGTGCCCCTGGACATCTGCACCAGCATCTGCAAGTACCCCGACTACATCAAGATGGTGAGCGAGCCCTACGGCGATAG

CCTGTTCTTCTACCTGCGGGCGGGAGCAGATGTTCGTGCGGCACCTGTTCAACAGAGCCGGCGCCGTGGGCGAGAACGTGCCCGACGACCTGT

ACATCAAGGGCAGCGGCAGCACCGCCAACCTGGCCAGCAGCAACTACTTCCCTACCCCCAGCGGCTCCATGGTGACCAGCGACGCCCAGATC

TTCAACAAGCCCTACTGGCTCCAGAGAGCCCAGGGCCACAACAATGGCATCTGCTGGGGCAACCAGCTGTTCGTGACCGTGGTGGATACCAC

CCGGAGCACCAACATGTCCCTGTGCGCCGCCATCAGCACCAGCGAGACCACCTACAAGAACACCAACTTCAAGGAGTACCTGAGGCACGGCG

AGGAGTACGACCTCCAGTTCATCTTCCAGCTGTGCAAGATCACCCTCACCGCCGACGTGATGACCTACATCCACAGCATGAACAGCACCATCC

TGGAGGACTGGAACTTCGGCCTGCAGCCCCCTCCTGGCGGCACCCTGGAGGACACCTACAGATTCGTGACCAGCCAGGCCATCGCATGCCAG

AAGCACACCCCTCCCGCCCCTAAGGAGGACCCCCTGAAGAAGTACACCTTCTGGGAGGTGAACCTGAAGGAGAAGTTCAGCGCCGACCTGG

ACCAGTTCCCTCTGGGCAGAAAGTTCCTGCTGCAAGCCGGCCTGAAGGCCAAGCCTAAGTTCACCCTGGGCAAGAGAAAGGCCACCCCCACC

ACAAGCAGCACCAGCACCACCGCCAAGCGGAAGAAGCGCAAGCTGTGA

1234

HPV-16 pL1

ATGTCCCTGTGGCTGCCCAGCGAGGCCACCGTGTACCTGCCCCCCGTGCCCGTGAGCAAAGTGGTGAGCACCGATGAGTACGTGGCCCGGAC

CAACATCTACTACCACGCCGGCACCTCCAGACTGCTGGCCGTGGGCCACCCCTACTTCCCCATCAAGAAGCCCAACAACAACAAGATCCTGGT

GCCCAAAGTGAGCGGCCTGCAGTACCGGGTGTTCAGAATCCACCTGCCCGACCCCAACAAGTTCGGCTTCCCCGACACCAGCTTCTACAACCC

CGACACCCAGAGACTGGTGTGGGCCTGCGTGGGCGTGGAAGTGGGCAGAGGCCAGCCTCTGGGCGTGGGCATCAGCGGCCACCCTCTGCT

GAACAAGCTGGACGACACCGAGAACGCCAGCGCCTACGCCGCCAACGCCGGCGTGGATAACAGAGAGTGCATCAGCATGGACTACAAGCA

GACCCAGCTGTGCCTCATCGGCTGCAAGCCCCCCATCGGCGAGCACTGGGGCAAGGGCAGCCCCTGCACCAACGTGGCCGTGAACCCTGGC

GACTGCCCTCCCCTGGAGCTCATCAACACCGTGATCCAGGACGGCGACATGGTGGACACCGGCTTCGGCGCCATGGACTTCACCACCCTCCA

GGCCAACAAGAGCGAAGTGCCCCTGGACATCTGCACCAGCATCTGCAAGTACCCCGACTACATCAAGATGGTGAGCGAGCCCTACGGCGATA

GCCTGTTCTTCTACCTGCGGGCGCGAGCAAATGTTTGTGCGGCACCTGTTCAACAGAGCCGGCGCCGTGGGCGAGAACGTGCCCGACGACCTG

TACATCAAGGGCAGCGGCAGCACCGCCAACCTGGCCAGCAGCAACTACTTCCCTACCCCCAGCGGCTCCATGGTGACCAGCGACGCCCAGAT

CTTCAACAAGCCCTACTGGCTCCAGAGAGCCCAGGGCCACAACAACGGCATCTGCTGGGGCAACCAGCTGTTCGTGACCGTGGTGGATACCA

CCCGGAGCACCAACATGTCCCTGTGCGCCGCCATCAGCACCAGCGAGACCACCTACAAGAACACCAACTTCAAGGAGTACCTGAGGCACGGC

GAGGAGTACGACCTCCAGTTCATCTTCCAGCTGTGCAAGATCACCCTCACCGCCGACGTGATGACCTACATCCACAGCATGAACAGCACCATC

CTGGAGGACTGGAACTTCGGCCTGCAGCCCCCTCCTGGCGGCACCCTGGAGGACACCTACAGATTCGTGACCAGCCAGGCCATCGCATGCCA

GAAGCACACCCCTCCCGCCCCTAAGGAGGACCCCCTGAAGAAGTACACCTTCTGGGAAGTGAACCTGAAGGAGAAGTTCAGCGCCGACCTG

GACCAGTTCCCTCTGGGCAGAAAGTTCCTGCTGCAGGCCGGCCTGAAGGCCAAGCCTAAGTTCACCCTGGGCAAGAGAAAGGCCACCCCCAC

CACAAGCAGCACCAGCACCACCGCCAAGCGGAAGAAGCGCAAGCTGTGA

Human codon-optimized HPV-16 (h)L1 CDS and Physcomitrella-optimized HPV-16 (p)L1 CDS. Grey

marked nucleotides were changed in HPV-16 pL1 through codon usage optimization by the tool physCO

(Top et al., 2021). Turquoise marked motifs refer to the Physcomitrella alternative splicing recognition

sites. The green marked segment refers to a predicted miRNA binding site identified by the tool

psRNATarget (Dai and Zhao, 2011; Dai et al., 2018). Light orange colored segments represent the nuclear

localization signal. Nucleotide changes are underlined in the HPV-16 pL1 sequence. (hL1, GenBank

accession number DQ067889; Maclean et al., 2007).

Supplementary Figure S3 - Characterization of CTPs by analysis of subcellular eGFP localization

1261

1262

1263

1264
1265

Microscopy pictures of Physcomitrella protoplasts transformed with CTP-eGFP constructs. Samples

were analyzed using a rhodamine and a FITC filter to identify chloroplast (magenta) and eGFP localization,

respectively, as well as a merge of the two signals. For sample CTPc21-eGFP no eGFP signal could be

observed in chloroplasts. For samples CTPc5-eGFP, CTPc10-eGFP, CTPc22-eGFP and CTPc22noMet-eGFP

green fluorescence signal is observed in chloroplasts. eGFP lacking a CTP localized to the cytoplasm and

served as a negative control. The experiment was repeated two times. CTPc5 was examined a third time

using confocal microscopy. Scale bars represent 10 μm.

Supplementary Figure S4 – Analysis of CDS integrity in L1-producing lines

1274

1275

1276

Analysis of rearrangements in the pL1 CDS using RT PCR. Lines pL1 23A, pL1Δ22 8A and CTP-pL1 30 were

checked for DNA or RNA rearrangements by performing PCR on cDNA using four different primer pairs.

Primer pairs were selected to amplify the entire cDNA. The forward primer binds at the start codon and

is the same in each primer pair (pL1_fwd_1 for lines pL1 23A & pL1Δ22 8A, CTP_pL1_fwd_1 for line CTP-

pL1 30). The reverse primers are different in each pair and bind at the 3' end of the CDS or within the

terminator (**Supplementary Table S1**). For line pL1 23A usage of 25 ng or 200 ng cDNA in PCR reactions

was compared. Following experiments used 200 ng. Absence of distinct bands of a size smaller than the

expected size is indicative for no larger rearrangements on both DNA and RNA level. No amplicon is visible

for lines pL1 23A using Primer Pair 2 and CTP-pL1 30 using Primer Pair 3. The constructs used for

transformation of Physcomitrella protoplasts (**Supplementary Figure S1 g - i**) served as a positive control.

1287

1288 **Supplementary Figure S5 – Analysis of the effect of reducing agents on L1 band patterning**

1289

1290

1291

1292 **Analysis of the effect of reducing agents on L1 band patterning in western blots.** Samples of line CTP-

1293 pL1 30 and wildtype were extracted without reducing agent, with 50 mM DTT or 40 mM ascorbic acid.

1294 The proteins for the western blot were separated on a 7.5% TGS polyacrylamide gel under reducing

1295 conditions (5 mM DTT) before being transferred to a membrane and developed using CamVir1 antibody

1296 (1:5000).

1297

1298 **Supplementary Figure S6 – Comparison of biomass accumulation rates**
1299

1300
Analysis of biomass accumulation rates among nine Physcomitrella lines of three different constructs.
Biomass accumulation rates were observed over a course of 4 weeks for lines pL1 5A, pL1 23A, pL1 47B
(blue); pL1Δ22 8A, pL1Δ22 11B, pL1Δ22 102B (green) and CTP-pL1 30, CTP-pL1 103, CTP-pL1 143 (purple).
The data was plotted using linear regression analysis and all slopes were analysed using ANOVA and post
hoc multiple comparisons test (TukeyHSD). Comparison of the slopes between the constructs of pL1,
pL1Δ22 and CTP-pL1 showed pL1 lines accumulated biomass faster than pL1Δ22 (p = <0.0001) and CTP-
pL1 (p = 0.0001) lines. Per week n = 3 biological samples were taken.

Supplementary Figure S7 – Ammonium sulfate precipitation
1309

1310
1311
Ammonium sulfate precipitation of L1 from moss extracts. Samples from the supernatant and the

resuspended and dialyzed pellet before and after filtration were analyzed for the presence of L1 content

and residual host proteins under reducing conditions *via* western blot and Coomassie-stained gel,

respectively. CTP-pL1 30 extracts were precipitated with either 40%, 45%, or 50% (w/v) ammonium sulfate

(AMS). After centrifugation, the supernatant and the precipitate were taken up and dialyzed against cation

exchange chromatography binding buffer to remove excess AMS. Non-dissolved protein was cleared from

the sample using 0.22 μm filters. 45% AMS was found to be sufficient for full precipitation of L1 from

crude extracts. 30 ng *S. cerevisiae*-produced L1 served as a positive control and Physcomitrella wildtype

as a negative control.

1321

1322 **Supplementary Figure S8 – Chromatogram of Cation Exchange purification**

1323

1324

1325

Chromatogram of HPV-16 L1 purification from moss using cation exchange chromatography. The
chromatogram shows the UV absorbance, indicative for protein concentration, in blue, gradient
concentration in pink, and conductivity, indicative for salt concentration, in red. The y axis depicts milli-
Absorbance Units and the x axis depicts the volume in mL. The large peak between 16 – 26 mL shows the
sample application or flow through, the following small peak at 28 mL the wash fraction and the larger
peak at 33 – 35 mL the elution fractions.

 Supplementary Table S1 – List of primers

1333 Capital letters refer to the part of the primer binding to the template whereas lower case letters represent
1334 overhangs added to the template via PCR. Abbreviations: fwd or fw (forward), rev or rv (reverse), CTP
1335 (chloroplast transit peptide), MMEJ (micro homology-mediated end joining), GFP (green fluorescent
1336 protein), OH (overhang (lower case letters)).

1337

Primer name	Sequence (5'→3')	Application
AA2_3integFP (hpt3'seq)	ACTCGCCGATAGTGGAAACC	Amp of linker-eGFP vector part 1
PAN_PpActin-Pro_Rv	CTGCACAACAACCACCAATATTTAATTTC	Amp of linker-eGFP vector part 1
linker:EGFP_p1840_Fw	GGAGGTGGAGGTGGAGC	Amp of linker-eGFP vector part 2
HPTLUCfwd2	TTGCCCTCGGACGAGTGC	Amp of linker-eGFP vector part 2
CTP_Pp3c10_Fw	aattaaatattggtggttgttgtgcagATGGCGTTATTGGCGCGC	Amp of CTPc10
CTP_Pp3c10_Rv	gctccacctccacctccATTGGTGACCCTCAAAGGTCCAG	Amp of CTPc10
CTP_Pp3c5_Fw	aattaaatattggtggttgttgtgcagATGGCTACGATGAAGATTGCTGTGG	Amp of CTPc5
CTP_Pp3c5_Rv	gctccacctccacctccAGCTCGCACGGAGCTGTTC	Amp of CTPc5
CTP_Pp3c22_Fw	aattaaatattggtggttgttgtgcagATGCGGGCTCTCTGTGGAG	Amp of CTPc22 & CTPc22_noMet
CTP_Pp3c22_Rv	gctccacctccacctccCATAGCGACAATGCACCGTCTTC	Amp of CTPc22
CTP_Pp3c22_noMet_Rv	gctccacctccacctccAGCGACAATGCACCGTCTTCC	Amp of CTPc22_noMet
CTP_Pp3c21_Fw	aattaaatattggtggttgttgtgcagATGGCCGCCGTCGGAATG	Amp of CTPc21
CTP_Pp3c21_Rv	gctccacctccacctccGGCACGGACAGCGAACGC	Amp of CTPc21
Act5_seq	TCCACATGGCTACAGCTG	Colony PCR CTP-eGFP constructs
EGFP_NoLinker_Rev	CTTGTACAGCTCGTCCATG	Colony PCR CTP-eGFP constructs
XhoI-pL1-BamHI_fw	gtgcagctcgagATGTCCCTGTGGCTGCC	Amp of pL1 and pL1Δ22
XhoI-pL1-BamHI _Rv	tctagaggatccTCACAGCTTGCGCTTCTTCC	Amp of pL1
XhoI-pL1delta22-BamHI _Rv	tctagaggatcctcaGCCCAGGGTGAACTTAGGC	Amp of pL1Δ22
pL1_w/oATG_Fw	TCCCTGTGGCTGCCCAG	Amp of CTP-pL1 (no Start codon)

pL1_Rv_35St-OH	GACTGGTGATTTTTGCGGACTCTAG	Amp of pL1, Heterosplicing and MMEJ check of pL1 lines
CTPc5_Fw_Act5p-OH	tattggtggttgttgtgcagctcgagATGGCTACGATGAAGATTGCTGTG	Amp of CTPc5
CTPc5_Rv_pL1-OH	ctcgctgggcagccacagggaAGCTCGCACGGAGCTGTTC	Amp of CTPc5
Seq_hL1+HSP_Fw	CGTGATGACCTACATCCACAG	Colony PCR pL1 constructs
pL1_Rv_35St-OH	GACTGGTGATTTTTGCGGACTCTAG	Colony PCR pL1 constructs
pL1_fwd_1	ATGTCCCTGTGGCTGCCC	Heterosplicing and MMEJ check of pL1, pL1Δ22 lines (primer pair 1-4, forward)
pL1_rev_1	TCACAGCTTGCGCTTCTTCC	Heterosplicing and MMEJ check of pL1, CTP-pL1 lines (primer pair 1, reverse)
pL1d22_rev_1	TCAGCCCAGGGTGAACTTAGG	Heterosplicing and MMEJ check of pL1Δ22 lines (primer pair 1, reverse)
CTP-pL1_fwd_1	ATGGCTACGATGAAGATTGCTGTG	Heterosplicing and MMEJ check of CTP-pL1 lines (primer pair 1-4, forward)
pL1_Rv_35St-OH	GACTGGTGATTTTTGCGGACTCTAG	Heterosplicing and MMEJ check of pL1, pL1Δ22, CTP-pL1 lines (primer pair 2, reverse)
101_35S-T_rev	CCCTTATCTGGGAACTACTCACACATTATTCTGGA	Heterosplicing and MMEJ check of pL1, pL1Δ22, CTP-pL1 lines (primer pair 3, reverse)
3342_LdC-HATag-35ST-rev	GGGTTTCGCTCATGTGTTGAGCA	Heterosplicing and MMEJ check of pL1, pL1Δ22, CTP-pL1 lines (primer pair 4, reverse)
pL1-fwd	GCTCGAGATGTCCCTGTG	pL1(Δ22)-GFP fusion construct
pL1-rev	AGGATCCCAGCTTGCGCTTCTTC	(CTP-)pL1-GFP fusion construct
pL1d22-rev	AGGATCCGCCCAGGGTGAACTTAG	pL1Δ22-GFP fusion construct
CTPc5-pL1 fwd	GCTCGAGATGGCTACGATGA	CTP-pL1-GFP fusion construct

1338

1339

1340 **Supplementary Table S2 – Predicted chloroplast transit peptides for six selected proteins in**
1341 **Physcomitrella. The predictions were performed with TargetP-2.0; Armenteros et al. (2019).**
1342

Name: CTPc5

Protein: **Pp3c5_1370V3.1**

Predicted chloroplast transit peptide (74 aa):

MATMKIAVAAVAGTPAAAVGTRSAAATETARLCASSAQGLGLSTFVGLRAGSNASRGVTPRVRPLKAENSSVRA

Name: CTPc10.1

Protein: **Pp3c10_10490V3.1**

Predicted chloroplast transit peptide (63 aa):

MEVLPRAIATSHVVVSSSSAPKESSESGSKNNSQVSLSSVPFLRPASALAASIVSSATPAALA

Name: CTPc10.2

Protein: **Pp3c10_2940V3.1**

Predicted chloroplast transit peptide (70 aa):

MALLARGAVAAVAAVGFSNTTSKQGHQNGGSNGVLAFRGNAFCRAVAARTSSVGCQVRVKRGAGPLRVTN

Name: CTPc21

Protein: **Pp3c21_9980V3.1**

Predicted chloroplast transit peptide (43 aa):

MAAVGMTMSLKSSTALKGDFLGAALKQTVAAPKAANVAFAVRA

Name: CTPc22

Protein: **Pp3c22_5470V3.1**

Predicted chloroplast transit peptide (62 aa):

MRALCGVPSCASLSSHRLASPLLAYRACEFALGSSPGLRLEFVGRAVGLQSRSIGRRCIVAM

1343