TITLE

mRNA stability factor HuR promotes immune evasion in pancreatic ductal adenocarcinoma

AUTHORS

Yifei Guo[1,2,3], Jennifer M. Finan[1,2,3], Alexandra Q. Bartlett[1,2,3], Shamilene Sivagnanam[2,3], Katie E. Blise[3], Nell Kirchberger[2,3],

Konjit Betre[2,3], Kevin Hawthorne[3,4], Canping Chen[5,6], Aaron Grossberg[3,7], Zheng Xia[5,6], Lisa M. Coussens[2,3], Rosalie C.

Sears[3,4], Jonathan R. Brody[1,2,3], Robert Eil[1,2,3]

[1]Department of Surgery, Oregon Health & Science University, Portland, OR, USA;

[2]Department of Cell, Developmental & Cancer Biology, Oregon Health & Science University, Portland, OR, USA;

[3]Brenden-Colson Center for Pancreatic Care, Knight Cancer Institute, Oregon Health & Science University, Portland, OR,

USA;

[4]Department of Molecular and Medical Genetics, Oregon Health & Science University, Portland, OR, USA;

[5]Computational Biology Program, Oregon Health & Science University, Portland, OR, USA;

[6]Department of Biomedical Engineering, Oregon Health & Science University, Portland, OR, USA;

[7]Department of Radiation Medicine, Oregon Health and Science University, Portland, OR, USA.

ABSTRACT

The tumor microenvironment (TME) of pancreatic ductal adenocarcinoma (PDAC) is characterized by a limited

infiltration of tumor-specific T cells and anti-tumor T cell activity. Extracellular factors in the PDAC TME have been widely

reported to mediate immune suppression, but the contribution from tumor-intrinsic factors is not well understood. The RNA-

binding protein, HuR (*ELAVL1*), is enriched in PDAC and negatively correlates with T cell infiltration. In an

immunocompetent Kras-p53-Cre (KPC) orthotopic model of PDAC, we found that genetic disruption of HuR impaired

tumor growth. Importantly, we found that HuR depletion in tumors enhanced both T cell number and activation states.

Mechanistically, HuR mediated the stabilization of mTOR pathway transcripts, and inhibition of mTOR activity rescued the

impaired function of local T cells. Phenotypically, we found that HuR induced T-cell suppression in PDAC, as HuR depletion

sensitize PDAC tumors to immune checkpoint blockade, while isogenic, wildtype tumors are resistant. Our findings describe

a novel role of HuR in facilitating tumor immune suppression in the PDAC TME by inhibiting T cell infiltration and

function, and implicate HuR inhibition as a potential therapeutic combination with immunotherapy.

SIGNIFICANCE

This study identified a novel mechanism that HuR supports pancreatic tumor growth by restricting T cell infiltration,

promoting immune evasion. Our work supports targeting strategies against HuR in PDAC with the goal of enhancing PDAC

sensitivity to immune-based cancer therapies, such as checkpoint blockade and T cell transfer.

INTRODUCTION

Pancreatic ductal adenocarcinoma (PDAC) is one of the leading causes of cancer-related death worldwide with a 5-

year survival rate of 13%, underscoring the need to develop improved therapeutic strategies[1]. The complex tumor

microenvironment (TME) of PDAC is thought to hinder both chemo- and immunotherapy efficacy[2]. Despite the presence

of immunogenic neoantigens that are predicted to be recognized by endogenous T cells, PDAC has remained resistant

to immune checkpoint blockade (ICB)[3–5]. Some have proposed that the ability of cancer cells to downregulate antigen

presentation machinery is a major driver of evading T cell recognition in PDAC. However, only a small proportion (~1-2%)

of PDAC contain mutations within antigen presentation or processing machinery[6]. Even in patients with mismatch repair

deficient (MMR-D) driven PDAC, where neoantigen frequency is equivalent to other MMR-D cancers, ICB had

compromised rates of objective response in PDAC in comparison to other MMR-D cancers[7]. Thus, we and others have

concluded that unresolved tumor-intrinsic sources of T cell suppression exist in PDAC. Specifically, cytotoxic T cells in the

PDAC TME have characteristics of T cell dysfunction, such as an inadequate response to stimulation (hypo-responsiveness),

which is recognized as a major source of tumor immune escape[8–10]. Therefore, in PDAC, there remains a need to understand

and reverse the tumor specific mechanisms that drive resistance to cancer immunotherapies[11].

Investigators continue to appreciate that TME features, including cytokines and cell-cell interactions, can restrain

the antitumor functions of CD4[+] and CD8[+] T cells. In tumors, PDAC especially, a combination of rapid cellular proliferation

and pathologic extracellular architecture contribute to a microenvironment that is low in both glucose and oxygen[12,13].

Meanwhile, T cells depend on consumption of nutrients to effectively respond to TCR ligation with acquisition of effector

programs, production of cytokines, and target cell cytolysis. Thus, the tumor-imposed nutrient deficit results in a metabolic

competition with local immune cells, contributing to T cell hypo-responsiveness and resistance to immunotherapies[14]. As

others have increasingly appreciated the importance of nutrient availability and metabolism for effective T cell antitumor function, interest continues to grow in therapies that could increase nutrient availability and consumption by tumor-specific T cells[15].

In human PDAC and preclinical models, the RNA-binding protein, human antigen R (HuR, *ELAVL1*) is known to be hyperactive, supporting multiple programmatic hallmarks of cancer[12,13,16,17]. Functionally, HuR recognizes and binds to 3' untranslated regions (UTRs) of transcripts, enhancing their stability and translation[18]. For instance, HuR reshapes the metabolic profile of PDAC, in part, by stabilizing *IDH1* transcripts and reprogramming both glycolytic and lipogenic metabolism[13,19,20]. Broadly, HuR activity is associated with Myc programs and glycolytic metabolism, enhancing the activation of c-Myc programs and consumption of surrounding nutrients for growth in size and cell division[12,21,22]. Additionally, pancreas-specific HuR overexpression alone caused fibro-inflammation in the mouse pancreas and led to an increase in precursor lesions of PDAC in cooperation with a $Kras^{G12D}$ mutation[23]. Our prior observations support a model wherein tumor-intrinsic HuR can mediate cell-cell communication and alter the PDAC TME in immune compromised models[13,24]. In the context of metabolic competition between tumor and immune cells, we hypothesized that tumor-intrinsic HuR activity may drive a competitive metabolic advantage over neighboring T cells. Yet, due to a lack of murine PDAC model that can capture the immune contexture with HuR depletion, the role of HuR in the PDAC-immune crosstalk has not been examined[16,17,24].

Taken together, our prior works point to a hypothesis that HuR may have a substantial role in mediating immune evasion in the PDAC TME. Moreover, we report here that HuR expression is negatively correlated with both T cell infiltration and outcome in PDAC patients. Thus, we investigated whether depletion of HuR in PDAC would impact tumor growth and anti-tumor immune activity in an immunocompetent model of PDAC. To this end, we developed an isogenic $Kras^{G12D},Trp53^{R172H},Pdx1\text{-}Cre$ (KPC) mutant-driven murine PDAC model, with and without HuR, to further elucidate the impact of tumor-intrinsic HuR on the immune components of the PDAC TME[25]. We observed that tumors with disruption in the *Elavl1* locus (HuR) had impaired tumor growth *in vivo*. To investigate the impact of the immune cells on tumor growth, we analyzed the infiltrated immune population in tumors and found that HuR depletion in tumors led to increased T cell infiltration and activation in the PDAC TME. Moreover, antibody-based depletion of CD4[+] and CD8[+] T cells restored the *in vivo* growth of HuR-deficient KPC tumors, demonstrating that HuR is required for intact PDAC growth in the immunocompetent setting. Introduction of a model antigen and assessment of tumor specific T cells identified that tumor-intrinsic HuR deterred the infiltration of antigen-specific CD8[+] T cells and their acquisition of effector functionality.

Mechanistically, HuR mediated stabilization of mTOR pathway-related transcripts and enhanced PDAC nutrient

consumption to the detriment of local T-cell antitumor function. Consistent with such a model, inhibition of the mTOR

pathway in PDAC cells enhanced the cytokine production capacity of co-cultured CD8[+] T cells in an HuR-dependent

manner. Therapeutically, KPC tumor's resistance to ICB treatment was dependent on the presence of HuR, and HuR

depletion combined with ICB led to tumor regression. Herein, we provide a deeper understanding of a potent immune

suppression mechanism driven by the pro-oncogenic protein, HuR, in PDAC.

METHODS

Single-cell data collection and analysis

Single-cell RNA sequencing (scRNA-seq) data were sourced from the Deeply Integrated Single-Cell Omics

(DISCO) database[26]. A total of 115 PDAC samples, encompassing 557,304 cells, were utilized to investigate the relationship

between *ELAVL1* gene expression in malignant cells and lymphocyte infiltration. Malignant cells were identified using the

R package scATOMIC (v2.0.2)[27] with the parameter "known_cancer_type = PAAD cell" and default settings for other

parameters. Spearman's correlation analysis was conducted to assess the association between gene expression levels in

tumor cells and cell-type infiltration in each sample.

HuR signature survival analysis

Published HuR activity signature[17] was assessed for Kaplan-Meier analysis of overall survival in surgical-resected

PDAC patient dataset[28]. The dataset is available in dbGaP database under accession phs003597.v1.p1.

Immunohistochemistry staining of tumors

Mouse tissues were fixed in 4% PFA overnight at 4°C before embedding and 5 μm sectioning. As previously

described[24], paraffin-embedded histological sections were stained for HuR (Santa Cruz Biotechnology; sc-5261) overnight

at 4°C, followed by incubation with a biotinylated secondary antibody for 1 hour at room temperature. Color was developed

with DAB substrate (Vector Laboratories; SK-4100), and mount for imaging. Brightfield images were acquired by Zeiss

Axio slide scanner.

Cell lines and virus

The PDAC murine cell line, KPC8069 (Kras^{G12D}-Trp53^{R172H}-Pdx1Cre), was a gift from Dr. Michael Hollingsworth. Cells were cultured in normal DMEM with 10% fetal bovine serum (Gibco) and 1% penicillin/streptomycin (Thermo Scientific; 15070063). Cell lines were maintained at 37°C in 5% humidified CO_2. Cell lines were *Mycoplasma* tested routinely using Mycoplasma Detection Kit (SouthernBiotech; 13100-01). Before any experiment, cells were passaged at least twice after thawing.

For HuR re-expression, KPC8069 cells were transfected with ecotropic retrovirus either carrying codon-enhanced *Elavl1* sequence or empty vector and supplemented with 1 μg/ml polybrene (Millipore Sigma; TR1003G). Cells containing the target vectors were selected by hygromycin (Millipore Sigma; H3274) and validated for HuR expression with immunoblotting. For ovalbumin (OVA) expression, KPC8069 cells were transfected with ecotropic retrovirus carrying the full-length OVA sequence and supplemented with 1 μg/ml polybrene. Cells containing the target vectors were selected by blasticidin (Alfa Aesar; J67216-XF) and validated for OVA expression with flow cytometry.

All ecotropic retroviruses are generated by packaging 20 μg of plasmid DNA synthesized by Genscript, Inc. containing genes of interest in an MSCV derived γ-retrovirus backbone (derived from Addgene; 122727) along with 3 μg pCL-Eco plasmid DNA (Addgene; 12371) with platinum-E ecotropic packaging cells (Cell Biolabs; RV-101) and jetOPTIMUS transfection reagent (Avantor; 101000006).

INDEL assessment

For INDEL assessment, genomic DNA was harvested from cells at time of analysis using the DNeasy Blood and Tissue DNA kit (Qiagen; 69504), and the region containing *Elavl1* short guide (sg) RNA target was PCR-amplified and Sanger sequenced by Eurofins Genomics. For each sample, a 493-bp region was amplified for sequencing (forward primer: 5′-ACTGGGGCTGAAGAAAAGTAGT-3′, reverse: 5′-GAACTGAATCATCTCCCCGGT-3′). 100 to 200 ng of each amplicon was sent for sequencing with 2 pmol of its respective sequencing primer (5′-TAGTCTGAGGTTAAAGAACAAAACTTCATT-3′). The indel percentage of amplicons derived from HuR-KO cells was deduced using the Synthego ICE tool (Synthego; ICE Analysis v3.0).

Immunoblot analysis

Cells were lysed using ice-cold RIPA buffer (Thermo Scientific; 89900) supplemented with protease inhibitors (Thermo Scientific; 87786). Protein concentration was quantified using the Pierce BCA Protein Assay kit (Thermo

Scientific; 23225). Equal amounts of whole protein extracts were loaded and separated by electrophoresis on a 4% to 12%

Bis-Tris gel (Invitrogen; NP0321BOX) and transferred to a PVDF membrane (Bio-Rad; 1704156). Blots were blocked in

Intercept® (TBS) Blocking Buffer (LI-COR Biosciences; 927-70003) and then probed with antibodies against anti-HuR

(1:1000, Santa Cruz Biotechnology; sc-5261) and anti-α-tubulin (1:5000, Invitrogen; A11126), followed by Thermo Fisher

Scientific Alexa secondary antibodies. Immunofluorescence was measured using a digital imager (Invitrogen; iBright 1500).

RNA extraction and RT-qPCR

RNA was extracted using RNeasy Mini Kit (Qiagen; 74106). RNA concentrations and quality were assessed by

NanoDrop spectrophotometer (Thermo Scientific). Complementary DNA (cDNA) was made from 500 ng RNA using High

Capacity cDNA Reverse Transcriptase Kit (Thermo Scientific; 4368813). RT-qPCR was performed with SYBR green

master mix (Thermo Scientific; 4385617) with HuR primers and cDNA. Amplification signal was captured by RT-qPCR

machine (Thermo Scientific; QuantStudio 7 Flex). Relative quantifications were assessed using the Log_2–$\Delta\Delta Ct$ method,

using GAPDH for normalization.

Mouse experiments

Animal protocol # IP00003322 and #IP00003155 were approved by Institutional Animal Care Regulations and Use

Committee of OHSU. These protocols are specifically applicable to the experiments reported in this manuscript. All mice

were obtained from The Jackson Laboratory. C57BL/6 mice (Cat. 000664) of 8–9 weeks of age were used as recipient hosts

for KPC tumor implantation. OT-I transgenic mice were used as OT-I T cell donors.

For T cell depletion studies, mice were intraperitoneally injected with αCD8 blocking antibody (200 μg, BioXcell;

BE0061) and αCD4 blocking antibody (200 μg; BioXcell; BE0003-1) two days before tumor implantation. The treatment

was maintained for the duration of the experiment by injecting every four days of αCD8 and αCD4 (100 μg each). For

isotype controls, rat IgG2b (BioXcell; BE0090) at matching amount was used. This approach achieved >99% depletion of

$CD8^+$ and $CD4^+$ T cells in spleens of T cell-depleted mice compared to control mice at endpoint.

For αPD-1 treatment, mice were intraperitoneally injected with αPD-1 blocking antibody (200 μg, BioXcell;

BP0146) or isotype controls, rat IgG2a (BioXcell; BE0089) every 4 days. The first injections happened 5 days after tumor

implantation, and each mouse received a total of 4 injections.

Mouse PDAC tumor model

For implantable tumor experiments, KPC tumor cells (1×10^6) were resuspended in 100 μl of 1:1 mixture of PBS (Fisher Scientific; 14-190-250) and Matrigel (Fisher Scientific; 354234), injected subcutaneously into the flanks of mice, and allowed to grow for 37 days. Tumor length and width were measured by caliper every four days.

For orthotopic tumor growth experiments, KPC tumor cells (4×10^4) were resuspended in 20 μl of 1:1 mixture of PBS and Matrigel, orthotopically injected into the tail of mouse pancreas, and allowed to grow for 20 days. For orthotopic tumor αPD-1 experiments, KPC tumor cells (9×10^4) were injected and allowed to grow for 15 days. Earlier endpoint criteria included severe weight loss, or extreme weakness and inactivity. Tumor weight was measured by chemical scale (Sartorius) at the endpoint.

Flow cytometry and analysis

Following euthanasia, tumors and spleens were promptly dissected, weighed, and stored in ice-cold PBS. Tumors were subsequently minced with surgical scissors and dissociated with GentleMACS octo-dissociator and preset program (m_impTumor_01, Miltenyi Biotec). The dissociated tumor was then filtered through a 40 μm filter. Spleens were mashed and filtered through a 40 μm filter. All samples were ACK lysed for 60 seconds using ACK lysing buffer (Gibco; A1049201), and then consolidated for staining.

Single-cell suspension of each sample was incubated with Live/Dead Fixable Aqua (Thermo Scientific; L34966) and Fc block (BD; 553142) in PBS for 30 min at 4°C, followed by incubation with primary fluorochrome-labeled antibodies at 4°C for 30 min in PBS with 0.5% BSA and 2 mM EDTA. For *ex vivo* cytokine production by T cell, samples were incubated for 4 hours at 37°C with PMA (20 ng/mL, Sigma; P1585), ionomycin (1 μg/mL, Sigma; I0634), and GolgiStop (BD; 554724). Intracellular staining was done using the Fixation/Permeabilization Kit (BD; 554722). Antibodies used in flow analysis are listed in Supplementary Table 1. Flow cytometric analysis was performed on Cytek Aurora spectral cytometer. Collected data were analyzed using FlowJo v10 (TreeStar).

In vitro T cell activation and co-cultures

Murine OT-I CD8+ T cells were isolated from the spleens of OT-I transgenic mice (The Jackson Laboratory; #003831) using the EasySep Mouse CD8+ T cell Isolation Kit (Stemcell; 19853), which yielded an enrichment of 96%–100% purity. Isolated CD8+ T cells were stimulated *in vitro* with plate-bound anti-CD3 and anti-CD28 (5 μg/mL; BioXcell)

for 24 hours and expanded in RPMI-1640 (Gibco) completed with 10% FBS (Gibco), 1% nonessential amino acids (Gibco),

1% sodium pyruvate (Gibco), 1% penicillin/streptomycin (Gibco), 57 μmol/L 2-BME (Sigma), and 100 IU/mL of

recombinant human IL-2 (Peprotech).

For CD8[+] T cell co-culture with tumor cells, 50,000 CD8[+] T cells were incubated with 50,000 tumor cells in 96-

well round-bottom plates for 24 hours at 37°C with phorbol 12-myristate 13-acetate (PMA) (20 ng/mL, Sigma; P1585),

ionomycin (1 μg/mL, Sigma; I0634), and GolgiStop (BD; 554724). Intracellular staining was done using the

Fixation/Permeabilization Kit and antibodies used are listed in Supplementary Table 1. SIINFEKL tetramers were obtained

from the NIH tetramer core facility.

RNA-sequencing

RNA was isolated from frozen cells using RNeasy Mini Kit (Qiagen; 74106). RNA concentrations and quality were

assessed by NanoDrop spectrophotometer. Preparation of RNA library and transcriptome sequencing were conducted by

Novogene Co., Ltd using the HiSeq Illumina platform. Novogene analysis service aligned raw reads to GRCm38 genome

and performed DESeq2 differential expression analysis. Gene set enrichment analysis (GSEA) was done using GSEA v4.2.3

software with the MSigDB hallmark and curated gene sets[29,30].

Ribonucleoprotein immunoprecipitation assay

KPC cells were washed and harvested with cell scrapper in cold-PBS. Ribonucleoprotein immunoprecipitation

assay was done according to manufacturer's instructions (MilliporeSigma; 17-701) with modification to RNA purification.

Lysates were immunoprecipitated by using 10 μg of either anti-HuR antibody (Santa Cruz Biotechnology; sc-5261) or anti-

IgG control (MilliporeSigma; 12-371). For RNA purification from immunoprecipitation, the protocol was followed up to

the precipitation of RNA in ethanol, after which the entire mixture was transferred to an RNA column using the RNeasy

Micro Kit (Qiagen; 74004), and the manufacturer's instructions were followed thereafter. RNA quality was analyzed via

Bioanalyzer Pico chip by the Massively Parallel Sequencing Shared Resource Core at OHSU.

Preparation of RNA library and transcriptome sequencing were conducted by Novogene Co., Ltd using the HiSeq

Illumina platform. Novogene analysis service aligned raw reads to GRCm38 genome and performed HuR-enriched peak

annotation with MACS2 software[31]. KEGG (Kyoto Encyclopedia of Genes and Genomes) pathway enrichment analysis

was done on HuR-enriched peaks.

Statistical analysis

Data were analyzed using unpaired two-tailed Student's *t*-test and one-way ANOVA test. In all cases, two-tailed

tests with P values less than 0.05 were considered significant. Statistics were calculated using GraphPad Prism 10 software

(GraphPad Software Inc). Results were graphed as mean ± standard deviation for *in vitro* experiments and ± standard error

of mean for animal experiments. Experimental sample sizes were chosen using power calculations and preliminary

experiments. Samples that had undergone technical failure during processing were excluded from subsequent analysis.

Data availability

RNA-sequencing dataset is deposited at the GEO (Gene Expression Omnibus) database as GSE287307. RIP-

sequencing dataset is deposited as GSE287349.

RESULTS

Tumor ELAVL1 (HuR) expression correlates with patient outcome and T cell infiltration in PDAC

We previously demonstrated that the RNA-binding protein HuR is highly expressed and activated (i.e. translocated

to the cytoplasm) in PDAC compared to adjacent normal pancreas[24,32]. First, we applied a validated HuR activity signature

highlighting HuR-dependent transcripts[17] to a large RNA-sequencing dataset including surgical-resected PDAC patients[28],

and showed that a lower HuR target activity signature predicted significantly prolonged patient survival (**Fig. 1A**).

Separately, baseline immune infiltration has been demonstrated to be associated with overall prognosis and response to ICB

in a variety of cancers[33]. To further understand the relationship between HuR activity in PDAC and antitumor immune

activity, we assayed DISCO single-cell RNA sequencing datasets and found tumor-intrinsic HuR expression levels

negatively correlated with total CD3+ lymphocyte infiltration, i.e. lower tumor-intrinsic *ELAVL1* mRNA levels were

associated with more abundant CD3+ lymphocyte infiltration in patient samples (**Fig. 1B** and **C**). To determine the causative

relationship between tumor-intrinsic HuR and T cell infiltration, and antitumor function in PDAC, we used a syngenetic

implantable mouse model of PDAC (KPC)[25]. To first assess whether HuR activity is recapitulated in our KPC model as in

human PDAC, HuR protein expression and localization was visualized with IHC staining. In concordance with patient data,

the KPC tumors had higher expression of whole-cell HuR compared to the healthy mouse pancreas (**Fig. 1D** and **E**). These

findings support previous work that tumor-intrinsic HuR is a poor prognostic marker in PDAC and pro-oncogenic[17,34], and

in addition, negatively correlates with lymphocyte infiltration in PDAC. HuR is similarly elevated in the KPC model as in

human PDAC providing a strong rationale for the investigation of tumor-intrinsic HuR in this immunocompetent murine

PDAC model. To our knowledge, this is the first genetic assessment of the role of HuR in PDAC tumor *in vivo* in an immune

intact setting.

Tumor-intrinsic HuR supports tumor growth in KPC model

We first established KPC cells with CRISPR-Cas9 mediated *Elavl1* genetic deletion using electroporation of

scramble or *Elavl1* targeted sgRNAs complexed to purified recombinant Cas9 protein, as previously described[35]. We isolated

and expanded single cell clones from these KPC HuR knockout (KO) pooled cell lines and three clones with stable knockout

and proliferation performance were selected and used for subsequent analyses. We then validated and quantified CRISPR-

Cas9 mediated disruption of the *Elavl1* locus via insertion or deletion (INDEL) via Sanger sequencing for our isolated

clones (**Fig. 2A**). Accordingly, the clones had a significant reduction in HuR mRNA and lost HuR protein expression (**Fig.**

2B and **C**).

To determine the impact of HuR on PDAC progression *in vivo*, we subcutaneously implanted KPC cells subjected

to electroporation with scramble sgRNAs wildtype (WT) or sgRNAs targeting *Elavl1* (HuR-KO) into syngeneic C57BL/6

mice and tracked tumor growth over time. We found that the loss of tumor-intrinsic HuR significantly delayed growth over

time by area and by end point weight (**Fig. 2D** and **E**). However, Ki67 staining showed no difference in tumor cell

proliferation between these WT and HuR-KO tumors, suggesting that the decrease in tumor growth that resulted from the

loss of PDAC HuR was dependent on a tumor-host interaction (**Fig. 2F**)*.* Importantly, re-expression of the *Elavl1* gene via

durable retroviral transduction in these HuR-KO cells rescued the tumor growth defect (**Fig. 2F** and **G**). Cumulatively, these

data causatively demonstrate, in genetic gain and loss of function, that HuR is required for PDAC progression *in vivo*.

HuR-dependent growth of PDAC is immune mediated in vivo

Our compiled analysis of TCGA cohorts identified a negative correlation between *ELAVL1* abundance and T cell

infiltration, suggesting that tumor-intrinsic HuR may act to constrain local immune activity and promote tumor progression.

To define the functional relationship between endogenous antitumor immunity and tumor-intrinsic HuR, we orthotopically

implanted KPC WT or HuR-KO into the pancreas of C57BL/6 hosts. We again observed a significant delay in growth in

the HuR-KO tumors in comparison to KPC WT tumors in this physiological-relevant stromal and immune

microenvironment (**Fig. 3A**). To provide a more granular understanding of the endogenous immune response, we performed enumerative and phenotypic assessment of intratumoral immune populations via flow cytometry. Our analysis revealed an increase in the abundance of CD4$^+$ T cells in KPC HuR-KO tumors as compared to WT tumors (**Fig. 3B**). We also observed an increase in NK1.1$^+$ T cells and CD19$^+$ B cells as a proportion of all CD45$^+$ leukocytes within HuR-KO tumors (Supplementary Fig. S1A). Phenotypic analysis of the tumor-infiltrating T cells demonstrated that both CD4$^+$ and CD8$^+$ T cells from HuR-KO tumors had a greater proportion expressing the transcription factor T-bet (*Tbx21*) (**Fig. 3C**). This is consistent with ongoing T cell effector function and polarization towards T helper Type 1 (T$_H$1) effector specification, the cell population responsible for antitumor activity[36]. Additionally, the percentage of CD4$^+$ Foxp3$^+$ regulatory T (T$_{Reg}$) cells trended lower in HuR-KO tumors compared to WT tumors, although not statistically significant (Supplementary Fig. S1B). Additionally, T cell surface phenotype markers CD25 and CD69 were more abundant on CD4$^+$ and CD8$^+$ T cells from HuR-KO tumors than WT tumors, suggestive of a more activated state in the HuR-KO tumors (**Fig. 3D**). Surface expression of antigen presentation major histocompatibility complex (MHC) class I, H2-kb, and immune checkpoint protein, PD-L1, on tumor cells was not significantly different between WT and HuR-KO tumors, suggesting HuR mediates immune suppression in this model independently of antigen presentation or the expression of co-inhibitory surface ligands (Supplementary Fig. S1C).

Lastly, to test the causal relationship between endogenous antitumor immunity and HuR-mediated tumor progression, we performed antibody-mediated depletion of CD4$^+$ and CD8$^+$ T cells from C57BL/6 hosts followed by orthotopic implantation of KPC tumors with or without HuR depletion. We found that depletion of CD4$^+$ and CD8$^+$ T cells restored the tumor growth of HuR-KO tumors, suggesting that the defect in tumor growth of HuR-KO PDAC is due to a comparative increase in endogenous T cell-mediated tumor control in the absence of tumor-intrinsic HuR (**Fig. 3A**). Together these data indicate that the tumor regression in the KPC HuR-KO tumors is mediated by presence of activated T cells, which are otherwise suppressed in WT conditions.

HuR restricts the effector functions of CD4$^+$ and CD8$^+$ T cells in PDAC

Drawing from our observation that tumor-intrinsic HuR suppressed the activation levels of PDAC-infiltrating T cells, we next asked whether HuR constrained the acquisition of effector functions of tumor-infiltrating T cells. To that end, we performed direct *ex vivo* reactivation of T cells infiltrating WT or HuR-KO PDAC tumors and captured and assessed their cytokine production via intracellular cytokine staining quantified by flow cytometry. By unsupervised clustering all

CD4[+] T cells based on their cytokine production, CD4[+] T cells derived from WT and HuR-KO tumors naturally separated

in two clusters demonstrating their distinct phenotypes (**Fig. 4A**). Consistent with increase in activation markers, CD4[+] T

cells derived from HuR-KO tumors exhibited increase in cytokine production and polyfunctionality (**Fig. 4A** and **B**).

Additionally, CD8[+] T cell derived from HuR-KO tumors displayed an increase in IFN-γ production compared to CD8[+] T

cells derived from WT tumors (Supplementary Fig. S2A).

HuR restricts the infiltration of antigen-specific CD4[+] and CD8[+] T cells in PDAC

To determine the impact of HuR loss in PDAC on the function and abundance of tumor-specific T cells, we

retrovirally transduced KPC cells with an ovalbumin (OVA)-expressing construct to make KPC cells express full-length

OVA for processing and presentation on MHC (**Fig. 4D**, left). When co-cultured KPC-OVA cells with *ex vivo* expanded and

activated T cell receptor (TCR) transgenic (OT-I) effector CD8[+] T cells specific for the ovalbumin SIINFEKL epitope

presented via H2kb MHC restriction. We found that KPC WT OVA cells exhibited suppression of OT-1 T cell cytokine

production (IFN-γ and TNF-α production) relative to KPC HuR-KO OVA cells (**Fig. 4C**), despite there being no difference

in MHC-I abundance or antigen presentation in KPC HuR-KO cells in response to interferon stimulation (**Fig. 4D**). This

co-culture model continues to support the notion that tumor-intrinsic HuR suppresses local T cell function in a manner

independent of antigen presentation or interferon response. To evaluate the endogenous T cell responses against the OVA

antigen in the presence or absence of tumor-intrinsic HuR, HuR WT or HuR-KO OVA cells were orthotopically implanted

into C57BL/6 mice. Notably, KPC HuR-KO OVA tumors were rejected in 80% of mice, with significantly smaller tumors

than KPC WT OVA tumors in those instances where a tumor was indeed evident (**Fig. 5D**). Given the absence of a pancreatic

tumor in most instances of HuR-KO KPC OVA, we evaluated the peritumoral tissues in both groups of tumor-bearing hosts.

We found statistically significant increases in the amount of CD8[+] T cells, and OVA tetramer[+] CD8[+] T cells in hosts bearing

HuR-KO OVA tumors than WT OVA tumors (**Fig. 4E**). We observed much lower SIINFEKL tetramer staining in the spleen

of tumor bearing mice, and no difference in the amounts of OVA tetramer[+] CD8[+] T cells between spleens from WT tumor-

bearing hosts compared to HuR-KO tumor-bearing hosts (Supplementary Fig. S2B). Together these data indicate that tumor-

intrinsic HuR suppresses the number of functional populations (cytokine producing, and antigen-specific) of T cells in the

PDAC TME, thus promoting tumor immune evasion.

HuR stabilizes anabolic transcripts and metabolic pathways

To identify the mechanism underlying HuR-dependent immune suppression and tumor progression in our immunocompetent model, we performed bulk-RNA sequencing on KPC WT and HuR-KO cells (Supplementary Fig. 3A). GSEA analysis identified established signatures involved in MYC pathways and metabolic reprogramming, including adipogenesis and cholesterol homeostasis, were higher in the KPC WT cells compared to the HuR-KO cells (Supplementary Fig. S3A and B). Given that HuR is known to stabilize RNA by directly binding to the 3'UTR regions of transcripts, we sought to identify those RNAs in the KPC WT cells using ribonucleoprotein immunoprecipitation (RIP) followed by high-throughput RNA sequencing (**Fig. 5A**). We validated HuR-RIP pulldown with immunoblot analysis from the bead-bound lysate and flow-through. Validating our protocol, motif analysis showed preferential binding to AU-rich sequences as previously described for HuR (**Fig. 6C** and Supplementary Fig. 3C)[37]. Additionally, established HuR metabolism targets, the *IDH1* 3' UTR[13], and the *Rictor* 3' UTR, demonstrated significantly peak enrichment in comparison to reference control *GAPDH* loci (**Fig. 5B** and Supplementary Fig. 3D). KEGG pathway analysis of the 1847 bound mRNAs revealed enrichment of transcripts involved in metabolism, including MAPK signaling, mTOR signaling, HIF1-α pathway and proteoglycan pathways, consistent with a model wherein HuR drives PDAC nutrient consumption to the potential detriment of adjacent cell populations (**Fig. 5D**).

To test our hypothesis that HuR increases mTOR activity and drives metabolic competition to suppress local T-cell function, we pretreated KPC OVA cells with mTOR inhibitors (AZD8055 and Torin1) and IFN-γ, and then co-cultured the pretreated KPC OVA cells with OVA-antigen specific OT-I cells (**Fig. 5E**). The pretreated KPC OVA cells had decreased mTOR activity by the measurement of decreased levels of phosphorylated-ribosome subunit 6 (p-S6). Of note, we validated that the inhibitors did not impair H2kb levels in these tumor cells (**Fig. 5F**). Confirming our hypothesis, we observed that OT-I T cells co-cultured with mTOR-inhibited KPC WT OVA cells had increased cytokine production (IFN-γ and TNF) in comparison to T cells co-cultured with vehicle-treated KPC WT OVA cells (**Fig. 5G and H**). In comparison, T cells co-cultured with KPC HuR-KO OVA cells demonstrated higher cytokine production *de novo,* independent of additional mTOR inhibition – suggesting overlap between the two pathways (**Fig. 5G** and **H**). This data further supports our hypothesis that HuR-KO tumor cells have lower mTOR activity at baseline and as a result, less nutrient competition with T cells, thus mTOR inhibition in HuR-KO tumor cells could not increase T cell function further. In addition, because of the different inhibition capacity of different inhibitors, T cells co-cultured with tumor cells treated with mTOR1 and mTOR2 inhibitor, Torin1, displayed increase in both IFN-γ and TNF production, while mTOR1 inhibitor, rapamycin, only displayed increase in TNF production (**Fig. 5G** and **H**). Taken together, these data indicate mechanistically tumor-intrinsic HuR directly binds

to and stabilizes transcripts involved in mTOR pathways, and the HuR-mediated increase of mTOR activity in tumor

metabolic competes with local T cells and ultimately suppress T cell function.

PDAC resistance to Immune checkpoint blockade is dependent on HuR

As previous discussed, ICB showed limited response in PDAC patients with deficiency in DNA mismatch repair

machinery[7], which suggested that the PDAC TME is inherently immune suppressive. Drawing from the observation that

PDAC HuR is responsible for constraining the infiltration and function of tumor specific T cells, we next asked whether

PDAC therapeutic resistance to ICB treatment was dependent on HuR function. Following a published treatment regimen[3],

we treated KPC WT and HuR-KO tumors-bearing mice with αPD-1 or control αIgG (**Fig. 6A**). Consistent with prior reports,

αPD-1 treatment alone in KPC WT tumors did not lead to tumor regression. However, αPD-1 treatment in KPC HuR-KO

tumors induced significant and almost total tumor regression (**Fig. 6A**). The αPD-1 treatment did not induce significant loss

of body weight in mice, indicating the treatment was well-tolerated in both groups (**Fig. 6B**). Consistent with previous

observations (**Fig. 3A**), HuR depletion led to increased $CD4^+$ T cell infiltration in tumors. While αPD-1 treatment in HuR-

KO tumors did not further boost the $CD4^+$ T cell infiltration compared to the IgG treatment, interestingly, αPD-1 treatment

in HuR-KO tumors induced higher infiltration of $CD8^+$ T cells with no effect observed in the WT tumors (**Fig. 6C**). Together,

these data indicate that $CD8^+$ T cells are the main responder to αPD-1 treatment in HuR-KO tumors, and support our

hypothesis that HuR depletion in PDAC tumors provides a greater proportion of T cells in the TME, which increased the

capacity for enhanced antitumor effector activity.

DISCUSSION

The immune landscape of PDAC TME is highly immunosuppressive and is a major barrier to effective

immunotherapy response[2,8]. Hence, understanding the underlying molecular mechanisms is key to advancing

immunotherapy strategies. From multi-omics analysis on large patient databases, the RNA-binding protein, HuR, negatively

correlates with both patient survival and total lymphocytes infiltration in PDAC. Herein, we found that HuR expressed by

PDAC promotes tumor growth in the immunocompetent PDAC model. Within PDAC cells, HuR directly binds to and

stabilizes anabolic transcripts, ultimately limiting the abundance and function of $CD4^+$ and $CD8^+$ T cells. As the primary

mediators of antitumor immunity, the T cells in the HuR-proficient PDAC TME had limited effector functions in cytokine

production and antigen-specific response. This mechanism could be exploited with therapeutic potential, as depletion of HuR sensitizes PDAC tumor to ICB treatment with increase efficacy.

In previous studies, loss of HuR in human PDAC xenografts in immunocompromised hosts had no impact on tumor growth[24], which complements our findings that HuR has an immune-mediated tumor suppressive role. In addition, previous work targeting overexpressing HuR in the mouse pancreas lead to chronic inflammation in the TME and promoted tumor development, indicating HuR contributes to an unbalanced inflamed environment[23]. Herein, this is the first report of PDAC tumor-intrinsic HuR having a potent extrinsic effect on the immune cells in the TME. This new discovery of HuR function fits with its known role in supporting tumor initial and tumor progression, and future study should investigate how HuR-mediated immune suppression promotes PDAC initiation from pancreatitis. Similarly, another key transcriptional regulator in PDAC, Myc, has been shown to control immune evasion in PDAC[11]. Therefore, studying pro-oncogenic molecules like HuR and Myc in immune-competent models may provide a much-needed understanding of the post-transcriptional and transcriptional tumor-intrinsic networks involved in immune suppression in the PDAC TME.

Interestingly, we did not observe a high correlation between gain/loss of RNA transcripts by RNA-sequencing and alignment abundance on RNA-immunoprecipitation. Overlapping these HuR-bound targets with those whose expression was reduced upon HuR-KO in the same cell lines identified 141 RNAs that are dependent on HuR binding to maintain transcript levels (data not shown). While the low percentage of overlap was surprising, published works have observed similar non-linear relationships between RNA-binding proteins and their targets[37,38]. These results further illustrate the complicated regulation mechanisms of RNA-binding proteins with its binding targets in general. A more granular understanding of the relationship between HuR 3' UTR binding and the impact on protein abundance will inform our ability to leverage HuR function for translational purposes.

Transcriptomic analyses in our KPC model confirmed multiple dysregulated pathways, which have been previously illustrated to be regulated by HuR[16,17]. Specifically, HuR directly upregulates multiple pathways in tumor remodeled metabolism[13]. Metabolic reprogramming in the tumor through the Warburg effect, characterized by aerobic glycolysis, not only enhances tumor proliferation but also depletes the limited resources within the TME[14,39]. Recent metabolite profiling of the PDAC interstitial fluid showed altered nutrient availability and nutrient deprivation in the PDAC TME[40], suggesting metabolic competition could be one of the mechanisms that tumor cells use to suppress immune cell function. Particularly, reports have shown that activated T cells need to reprogram their glucose and lipid metabolism to support their effector functions[41,42], and Tregs are more adapted to glucose deprivation with an increase in fatty-acid oxidation[43]. These required

nutrients are in concordance with the HuR-mediated dysregulated metabolic pathways, suggesting HuR mediates immune suppression via metabolic competition in the PDAC TME. In lieu of this, previous finding that PDAC WT and HuR-KO tumors grow similarly in immunocompromised model further supports the notion that HuR-mediated anabolic pathways do not support tumor growth by proliferation[24], instead by inducing immune suppression. In total, our results warrant further exploration to define the mechanistic connection between HuR-mediated PDAC metabolism and T cell function and differentiation in the TME.

In the PDAC immune landscape, our findings once again demonstrate the potential of CD8$^+$ T cells to be harnessed for PDAC tumor control, while also highlight that CD4$^+$ T cells can play a crucial role in addition to well-described CD8$^+$ T cells. Similar observations of CD4$^+$ T cell having anti-PDAC response were found in the setting of αCD40 stimulation[44]. These findings support the potential of targeting CD4$^+$ T cells to augment anti-tumor responses. Other than T cells, tumor-intrinsic HuR has broader impact on the immune landscape of the PDAC TME by increasing B cell infiltration and decreasing dendritic cell populations (data not shown). One recent study identified an immunosuppressive subset of dendritic cells in the PDAC TME, which could account for the phenotype we observed[45]. Still, the implications of these findings highlight the need for further investigation into the role of non-T cell immune cells.

Broadly, these findings advance the paradigm that tumor-intrinsic mechanisms limit T cell infiltration and the efficacy of anti-tumor immunity within the TME. Specifically, we highlight a novel role of tumor-intrinsic HuR as a modulator of PDAC immune evasion. Currently, targeting strategies against HuR are under development (e.g., siHuR nanotherapies and small molecule inhibitors)[46], and could potentially break PDAC resistance to immune-based anti-cancer therapies (i.e., checkpoint blockade and T cell transfer) as demonstrated here in our pre-clinical KPC model. Because HuR also has cell-intrinsic functions in intratumoral immune populations, further work will be required to understand the combined impact of HuR inhibition on tumor control.

FIGURE LEGENDS

Figure 1. Tumor ELAVL1 (HuR) expression correlates with patient outcome and T cell infiltration in PDAC

(A) Kaplan-Meier plots demonstrating proportional overall survival in PDAC patients (n = 199) distinguished by low versus high HuR Activity Signature.

(**B**) Scatter plot showing the Spearman's correlation between ELAVL1 expression in tumor cells and lymphocyte infiltration, including CD8+ T cells, CD4+ T cells, and NK cells (n = 115 patients) and (**C**) violin plot demonstrating the difference in lymphocyte infiltration in HuR- low and HuR-high tumors from the same samples.

(**D**) Representative images from patient tumor microarrays and KPC tumors stained for HuR expression and (**E**) quantification of HuR protein staining intensity of normal pancreas, KPC GEMM (n=3) and xenograft tumors (n = 8). Error bars represent standard error of the mean. ns, not significant; **P < 0.01; ***P < 0.005; ****P < 0.001, unpaired 2-tailed Student's t tests (**C,E**).

<u>Figure 2</u>. **Tumor-intrinsic HuR supports tumor growth in KPC model**

(**A**) INDEL quantification of single-cell clones isolated from CRISPR-Cas9 mediated disruption of the *Elavl1* locus in KPC8069 tumor cells and (**B**) quantification of mRNA abundance by RT-qPCR and (**C**) protein abundance by immunoblot of the same clones.

(**D**) Rates of subcutaneous KPC tumor growth represented over time following implant of KPC WT and HuR-KO tumor cells (n = 5) . (**E**) Summary quantification and pictures of the final tumor weight and size on day 34.

(**F**) Quantification of the percentages of Ki67$^+$ cells in total non-immune cells in subcutaneous KPC tumors by flow cytometry (n = 5) .

(**G**) Histogram and quantification of flow cytometry analysis of HuR expression in KPC WT and HuR-KO tumor cells retrovirally transduced with empty vector or HuR-overexpression vector prior to subcutaneous implantation to C57BL/6 hosts (n = 10). (**H**) Summary quantification of the final tumor weight on day 30.

Error bars represent stand deviation (**B,F**) and standard error of the mean (**D,E,F,G,H**). *P < 0.05; **P < 0.01; ****P < 0.001, one-way ANOVA test (**B,G,H**), unpaired 2-tailed Student's t tests (**D,E,F**).

<u>Figure 3</u>. **HuR-dependent growth of PDAC is immune mediated *in vivo***

(**A**) Representative flow cytometry illustrating enumerating T cells in host spleens treated with IgG or αCD4/8 before and during orthotopic tumor implantation and summary analysis of final KPC WT and HuR-KO tumor weight in host with or without T cell depletion (n = 6) .

(**B**) Summary analysis enumerating tumor infiltrating lymphocytes (TIL) in orthotopic KPC WT and HuR-KO tumors (n = 5).

(**C**) Summary data depicting CD4 and CD8 T-cell Tbet$^+$ percentages and (**D**) representative flow cytometry plot and

summary data depicting CD25 and CD69 phenotypic states on CD4 and CD8 T cells as in (**B**).

Error bars represent standard error of the mean. *P < 0.05; **P < 0.01; ***P < 0.005; ****P < 0.001, one-way ANOVA test

(**A**), unpaired 2-tailed Student's t tests (**B,C,D**),

<u>**Figure 4**</u>. **HuR restricts the function and infiltration CD4 and CD8 T cells in PDAC**

(**A**) Concatenated single-cell overlay and colorimetric density tSNE based depiction of the indicated proteins by flow

cytometry, and (**B**) representative and summary quantification of cytokines following *ex vivo* stimulation of isolated CD4$^+$

T cells from KPC WT and HuR-KO tumors (n = 5).

(**C**) Representative flow cytometry plot and summary data depicting cytokine productions of OT-I CD8$^+$ T cells co-cultured

with KPC WT OVA or KPC HuR-KO OVA tumor cells (n = 3).

(**D**) Representative flow cytometry plot depicting HuR expression and Kb-SIIN staining of KPC WT OVA or KPC HuR-

KO OVA tumor cells prior to orthotopic implantation, and summary data of tumor rejection rate and final tumor weight at

day 20. (**E**) Representative flow cytometry plot and summary data enumerating spleen and pancreas infiltrating CD8$^+$ T

cells and SIIN$^+$ CD8$^+$ T cells from KPC WT OVA and KPC HuR-KO OVA tumor-bearing mice (n = 5).

Error bars represent stand deviation (**C**) and standard error of the mean (**B,D,E**). *P < 0.05; **P < 0.01; ***P < 0.005; ****P

< 0.001, unpaired 2-tailed Student's t tests.

<u>**Figure 5**</u>. **HuR stabilizes anabolic transcripts and metabolic pathways**

(**A**) Schematic depicting RIP-seq process to identify HuR-bound transcript (n = 3). (**B**) RIP-seq reads genome alignment

plot to *IDH1 and Rictor* loci (3'UTR region highlighted with red box). (**C**) Representative immunoblot validation of HuR

pulldown in KPC cells in comparison to IgG control. (**D**) Summary depiction of enrichment scores (RichFactor) of KEGG

pathways with transcripts comparatively enriched in HuR-IP samples.

(**E**) Schematic depicting the pre-treatment of tumor cells with mTOR inhibitors (Rapamycin and Torin1) and IFN-γ before

co-culture with OT-I T cells. (**F**) Representative flow cytometry plot depicting *p*-S6 and H2-Kb staining of KPC WT OVA

cells after mTOR inhibitors and IFN-γ treatments. (**G**, **H**) Summary data depicting cytokine productions of OT-I CD8$^+$ T

cells co-cultured with pre-treated KPC WT OVA or KPC HuR-KO OVA tumor cells (n = 3).

Error bars represent stand deviation (**G**, **H**). *P < 0.05; ***P < 0.005; ****P < 0.001, one-way ANOVA test.

Figure 6. PDAC resistance to Immune checkpoint blockade is dependent on HuR

(**A**) Schematic for αPD-1 treatment on KPC tumor bearing mice, and summary data on final tumor weight at day15. (**B**)

Body weight of mice in (**A**) by group tracked over time (n = 4).

(**C**) Summary analysis enumerating tumor infiltrating lymphocytes (TIL) of KPC WT and HuR-KO tumors in mice received

IgG or αPD-1 treatment in (**A**).

Error bars represent standard error of the mean. *P < 0.05; **P < 0.01, unpaired 2-tailed Student's t tests (**A**), one-way

ANOVA test (**C**).

Supplement Figure 1. **Tumor-intrinsic HuR mediates immune composition in the PDAC TME.**

(**A**) Summary analysis enumerating tumor infiltrating NK cells, B cells, dendritic cells, macrophages, and (**B**) Foxp3$^+$ CD4

T-cell percentages. (**C**) Summary data depicting H2-kb and PD-L1 expression on non-immune cells in orthotopic KPC WT

and HuR-KO tumors (n = 5).

Error bars represent standard error of the mean. *P < 0.05; ***P < 0.005, unpaired 2-tailed Student's t tests.

Supplement Figure 2. **HuR restricts the function and infiltration CD4 and CD8 T cells in PDAC**

(**A**) Summary quantification of IFN-γ production following *ex vivo* stimulation of isolated CD8+ T cells from KPC WT and

HuR-KO tumors (n = 5).

(**B**) Summary data enumerating spleen infiltrating CD8 T cells and SIIN tetramer$^+$ CD8 T cells from KPC WT OVA and

KPC HuR-KO OVA tumor-bearing mice (n = 5).

Error bars represent standard error of the mean. **P < 0.01, unpaired 2-tailed Student's t tests.

Supplement Figure 3. **HuR stabilizes anabolic transcripts and metabolic pathways**

(**A**) Schematic and volcano plot depicting whole-transcriptomic changes between KPC WT and KPC HuR-KO tumor cells

(n = 3). (**B**) Summary depiction of normalized enrichment scores (NES) of gene-sets with transcripts comparatively enriched

or lost between indicated groups.

(**C**)Top 2 motif enrichment sequences from HuR-IP targets. (**D**) RIP-seq reads genome alignment plot to *GAPDH* locus

(3'UTR region highlighted with red box).

529 FDR q-value < 0.05, weighted Kolmogorov–Smirnov statistic test (**B**).

REFERENCES

1. Siegel RL, Miller KD, Wagle NS, Jemal A. Cancer statistics, 2023. *CA Cancer J Clin*. 2023;73(1):17-48.
doi:10.3322/caac.21763

2. Polani F, Grierson PM, Lim KH. Stroma-targeting strategies in pancreatic cancer: Past lessons, challenges and
prospects. *World J Gastroenterol*. 2021;27(18):2105-2121. doi:10.3748/wjg.v27.i18.2105

3. Evans RA, Diamond MS, Rech AJ, et al. Lack of immunoediting in murine pancreatic cancer reversed with neoantigen.
JCI Insight. 2016;1(14). doi:10.1172/jci.insight.88328

4. Freed-Pastor WA, Lambert LJ, Ely ZA, et al. The CD155/TIGIT axis promotes and maintains immune evasion in
neoantigen-expressing pancreatic cancer. *Cancer Cell*. 2021;39(10):1342-1360.e14. doi:10.1016/j.ccell.2021.07.007

5. Bailey P, Chang DK, Forget MA, et al. Exploiting the neoantigen landscape for immunotherapy of pancreatic ductal
adenocarcinoma. *Sci Rep*. 2016;6(1):35848. doi:10.1038/srep35848

6. Concha-Benavente F, Srivastava R, Ferrone S, Ferris RL. Immunological and clinical significance of HLA class I
antigen processing machinery component defects in malignant cells. *Oral Oncol*. 2016;58:52-58.
doi:10.1016/j.oraloncology.2016.05.008

7. Marabelle A, Le DT, Ascierto PA, et al. Efficacy of Pembrolizumab in Patients With Noncolorectal High Microsatellite
Instability/Mismatch Repair–Deficient Cancer: Results From the Phase II KEYNOTE-158 Study. *J Clin Oncol*.
Published online January 1, 2020. doi:10.1200/JCO.19.02105

8. Ino Y, Yamazaki-Itoh R, Shimada K, et al. Immune cell infiltration as an indicator of the immune microenvironment of
pancreatic cancer. *Br J Cancer*. 2013;108(4):914-923. doi:10.1038/bjc.2013.32

9. Gu H, Deng W, Zheng Z, Wu K, Sun F. CCL2 produced by pancreatic ductal adenocarcinoma is essential for the
accumulation and activation of monocytic myeloid-derived suppressor cells. *Immun Inflamm Dis*. 2021;9(4):1686-
1695. doi:10.1002/iid3.523

10.   Blise KE, Sivagnanam S, Betts CB, et al. Machine Learning Links T-cell Function and Spatial Localization to
Neoadjuvant Immunotherapy and Clinical Outcome in Pancreatic Cancer. *Cancer Immunol Res*. 2024;12(5):544-558.
doi:10.1158/2326-6066.CIR-23-0873

11.   Krenz B, Gebhardt-Wolf A, Ade CP, et al. MYC- and MIZ1-Dependent Vesicular Transport of Double-Strand
RNA Controls Immune Evasion in Pancreatic Ductal Adenocarcinoma. *Cancer Res*. 2021;81(16):4242-4256.
doi:10.1158/0008-5472.CAN-21-1677

12.   Blanco FF, Jimbo M, Wulfkuhle J, et al. The mRNA-binding protein HuR promotes hypoxia-induced
chemoresistance through posttranscriptional regulation of the proto-oncogene PIM1 in pancreatic cancer cells.
Oncogene. 2016;35(19):2529-2541. doi:10.1038/onc.2015.325

13.   Zarei M, Lal S, Parker SJ, et al. Posttranscriptional Upregulation of IDH1 by HuR Establishes a Powerful
Survival Phenotype in Pancreatic Cancer Cells. *Cancer Res*. 2017;77(16):4460-4471. doi:10.1158/0008-5472.CAN-17-
0015

14.   Chang CH, Qiu J, O'Sullivan D, et al. Metabolic Competition in the Tumor Microenvironment Is a Driver of
Cancer Progression. *Cell*. 2015;162(6):1229-1241. doi:10.1016/j.cell.2015.08.016

15.   Li W, Pan X, Chen L, et al. Cell metabolism-based optimization strategy of CAR-T cell function in cancer
therapy. *Front Immunol*. 2023;14. doi:10.3389/fimmu.2023.1186383

16.   Lal S, Burkhart RA, Beeharry N, et al. HuR post-transcriptionally regulates WEE1: Implications for the DNA damage response in pancreatic cancer cells. *Cancer Res*. 2014;74(4):1128-1140. doi:10.1158/0008-5472.CAN-13-1915

17.   Brown SZ, McCarthy GA, Carroll JR, et al. The RNA-Binding Protein HuR Posttranscriptionally Regulates the Protumorigenic Activator YAP1 in Pancreatic Ductal Adenocarcinoma. *Mol Cell Biol*. 2022;42(7):e00018-22. doi:10.1128/mcb.00018-22

18.   Kim HH, Yang X, Kuwano Y, Gorospe M. Modification at HuR(S242) Alters HuR Localization and Proliferative Influence. *Cell Cycle Georget Tex*. 2008;7(21):3371-3377.

19.   Anand S, Khan MA, Zubair H, et al. MYB sustains hypoxic survival of pancreatic cancer cells by facilitating metabolic reprogramming. *EMBO Rep*. 2023;24(3):e55643. doi:10.15252/embr.202255643

20.   Huang F, Bu L, Li M, et al. HuR/miR-124-3p/VDR complex bridges lipid metabolism and tumor development in colorectal cancer. Published online March 1, 2023. doi:10.21203/rs.3.rs-2587383/v1

21.   Osma-Garcia IC, Capitan-Sobrino D, Mouysset M, et al. The RNA-binding protein HuR is required for maintenance of the germinal centre response. *Nat Commun*. 2021;12(1):6556. doi:10.1038/s41467-021-26908-2

22.   Shao Z, Tu Z, Shi Y, et al. RNA-Binding Protein HuR Suppresses Inflammation and Promotes Extracellular Matrix Homeostasis via NKRF in Intervertebral Disc Degeneration. *Front Cell Dev Biol*. 2020;8:611234. doi:10.3389/fcell.2020.611234

23.   Peng W, Furuuchi N, Aslanukova L, et al. Elevated HuR in Pancreas Promotes a Pancreatitis-Like Inflammatory Microenvironment That Facilitates Tumor Development. *Mol Cell Biol*. 2018;38(3):e00427-17. doi:10.1128/MCB.00427-17

24.   McCarthy GA, Di Niro R, Finan JM, et al. Deletion of the mRNA stability factor ELAVL1 (HuR) in pancreatic cancer cells disrupts the tumor microenvironment integrity. *NAR Cancer*. 2023;5(2):zcad016. doi:10.1093/narcan/zcad016

25.   Hingorani SR, Wang L, Multani AS, et al. Trp53R172H and KrasG12D cooperate to promote chromosomal instability and widely metastatic pancreatic ductal adenocarcinoma in mice. *Cancer Cell*. 2005;7(5):469-483. doi:10.1016/j.ccr.2005.04.023

26.   Li M, Zhang X, Ang KS, et al. DISCO: a database of Deeply Integrated human Single-Cell Omics data. *Nucleic Acids Res*. 2022;50(D1):D596-D602. doi:10.1093/nar/gkab1020

27.   Nofech-Mozes I, Soave D, Awadalla P, Abelson S. Pan-cancer classification of single cells in the tumour microenvironment. *Nat Commun*. 2023;14(1):1615. doi:10.1038/s41467-023-37353-8

28.   Link JM, Eng JR, Pelz C, et al. Ongoing replication stress tolerance and clonal T cell responses distinguish liver and lung recurrence and outcomes in pancreatic cancer. *Nat Cancer*. Published online January 9, 2025:1-22. doi:10.1038/s43018-024-00881-3

29.   Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A*. 2005;102(43):15545-15550. doi:10.1073/pnas.0506580102

30.   Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst*. 2015;1(6):417-425. doi:10.1016/j.cels.2015.12.004

31.   Zhang Y, Liu T, Meyer CA, et al. Model-based Analysis of ChIP-Seq (MACS). *Genome Biol*. 2008;9(9):R137. doi:10.1186/gb-2008-9-9-r137

32.   Lal S, Cheung EC, Zarei M, et al. CRISPR Knockout of the HuR Gene Causes a Xenograft Lethal Phenotype. *Mol Cancer Res*. 2017;15(6):696-707. doi:10.1158/1541-7786.MCR-16-0361

33.   Carstens JL, Correa de Sampaio P, Yang D, et al. Spatial computation of intratumoral T cells correlates with survival of patients with pancreatic cancer. *Nat Commun*. 2017;8(1):15095. doi:10.1038/ncomms15095

34.   Richards NG, Rittenhouse DW, Freydin B, et al. HuR Status is a Powerful Marker for Prognosis and Response to Gemcitabine-Based Chemotherapy for Resected Pancreatic Ductal Adenocarcinoma Patients. *Ann Surg*. 2010;252(3):499. doi:10.1097/SLA.0b013e3181f1fd44

35.   Collier C, Wucherer K, McWhorter M, et al. Intracellular K+ Limits T-cell Exhaustion and Preserves Antitumor Function. *Cancer Immunol Res*. 2024;12(1):36-47. doi:10.1158/2326-6066.CIR-23-0319

36.   Fridman WH, Pagès F, Sautès-Fridman C, Galon J. The immune contexture in human tumours: impact on clinical outcome. *Nat Rev Cancer*. 2012;12(4):298-306. doi:10.1038/nrc3245

37.   Rothamel K, Arcos S, Kim B, et al. ELAVL1 primarily couples mRNA stability with the 3′ UTRs of interferon-stimulated genes. *Cell Rep*. 2021;35(8):109178. doi:10.1016/j.celrep.2021.109178

38.   Antal CE, Oh TG, Aigner S, et al. A super-enhancer-regulated RNA-binding protein cascade drives pancreatic cancer. *Nat Commun*. 2023;14(1):5195. doi:10.1038/s41467-023-40798-6

39.   Hanahan D, Weinberg RA. Hallmarks of Cancer: The Next Generation. *Cell*. 2011;144(5):646-674. doi:10.1016/j.cell.2011.02.013

40.   Sullivan MR, Danai LV, Lewis CA, et al. Quantification of microenvironmental metabolites in murine cancers reveals determinants of tumor nutrient availability. DeBerardinis R, van Lohuizen M, DeBerardinis R, Frezza C, eds. *eLife*. 2019;8:e44235. doi:10.7554/eLife.44235

41.   Kidani Y, Elsaesser H, Hock MB, et al. Sterol regulatory element–binding proteins are essential for the metabolic programming of effector T cells and adaptive immunity. *Nat Immunol*. 2013;14(5):489-499. doi:10.1038/ni.2570

42.   Leone RD, Powell JD. Metabolism of immune cells in cancer. *Nat Rev Cancer*. 2020;20(9):516-531. doi:10.1038/s41568-020-0273-y

43.   Yan Y, Huang L, Liu Y, et al. Metabolic profiles of regulatory T cells and their adaptations to the tumor microenvironment: implications for antitumor immunity. *J Hematol OncolJ Hematol Oncol*. 2022;15:104. doi:10.1186/s13045-022-01322-3

44.   Morrison AH, Diamond MS, Hay CA, Byrne KT, Vonderheide RH. Sufficiency of CD40 activation and immune checkpoint blockade for T cell priming and tumor immunity. *Proc Natl Acad Sci U S A*. 2020;117(14):8022-8031. doi:10.1073/pnas.1918971117

45.   Kenkel JA, Tseng WW, Davidson MG, et al. An Immunosuppressive Dendritic Cell Subset Accumulates at Secondary Sites and Promotes Metastasis in Pancreatic Cancer. *Cancer Res*. 2017;77(15):4158-4170. doi:10.1158/0008-5472.CAN-16-2212

46.   Finan JM, Sutton TL, Dixon DA, Brody JR. Targeting the RNA-Binding Protein HuR in Cancer. *Cancer Res*. 2023;83(21):3507-3516. doi:10.1158/0008-5472.CAN-23-0972

Figure 1

A HuR Sig. GSVA Score
Survival (%) vs Time (days)
Top Quartile
Bottom Quartile
p-value: <0.001

B
slope: -0.3925
p-value: <0.0001
Lymphocytes of all (%) vs *ELAVL1* (log count)

C
Lymphocytes of all (%)
**
○ HuR Low ● HuR High

D
Human
Pancreas — Tumor
200 µm — 200 µm
Murine
Pancreas — Tumor
50 µm — 50 µm

E
KPC GEMM
HuR Intensity (pixel)
***
KPC Orthotopic
HuR Intensity (pixel)
****
Kras^{G12D}
Trp53^{R172H} −   +   −   +

Figure 2

A

Clone	ICE	KO-Score	R^2	Indels
KO6	80	74	0.8	{'1': 7.0, '-1': 67.0, '-3': 6.0}
KO8	97	97	0.97	{'-1': 97.0}
KO9	93	93	0.93	{'-1': 83.0, '-2': 10.0}

B *ELAVL1* mRNA

C

D

E

F

G

H

Figure 3

Figure 4

Figure 5

Figure 6

A

KPC WT ● KPC HuR KO ●

IgG or αPD-1 (200 μg/injection)

C57BL/6

Day 5 8 11 14

Tumor Weight (g) — ns

IgG	+	−
αPD-1	−	+

Tumor Weight (g) — *

	+	−
	−	+

B

Body Weight (g) vs Days

● KPC WT
● KPC HuR KO
— IgG
- - αPD-1

C

CD4⁺ of CD45⁺ (%) — **

IgG	+	−	+	−
αPD-1	−	+	−	+

CD8⁺ of CD45⁺ (%) — *

	+	−	+	−
	−	+	−	+

Supplement Figure 1

Supplement Figure 2

Supplement Figure 3

A

HuR WT   HuR KO

RNA Extraction

Bulk RNA-seq

Clu
Pa2g4
Gldc
Ilf2
Gm10705
Mal2
Hoxb9
Hoxa3
Hoxb6
Hoxb5
Hoxb8
Hoxb5os

P value (-log$_2$)

HuR KO : WT (log$_2$)

B

E2F Targets
Myc Targets
mRNA Splicing
G2M Checkpoint
Adipogenesis
Cholesterol Homeostasis
Inflammatory Response
Glutamate Receptors

NES HuR KO : WT

C   Motif Enrichment

D

HuR
RIP-seq
Input

GAPDH

Supplemental Table 1. Flow cytometry antibodies.

Marker	Flour
CD11c	BV650
CD19	BV711
CD25	BV421
CD4	PCP-Cy5.5
CD4	BUV805
CD45	AF700
CD62L	BUR395
CD69	APC-Cy7
CD8	FITC
CD8	APC-Cy7
CD90.2	BV786
F4/80	BV421
FasL	PE
Foxp3	PE
GranzB	PE
IFNg	APC
IL-2	FITC
Ki67	BUV805
LiveDead	Aqua
MHC-I	FITC
MHC-II	APC-Cy7
NK1.1	PE-Cy7
PD-1	APC
PD-L1	APC
TBET	PE-Cy7
Tim3	BV650
TNF	BV421