Motion processing in visual cortex of maculopathy patients

Abbreviated title: Cortical motion processing in maculopathy patients

Célia Michaud[1 #], Jade Guénot[1 #], Cynthia Faurite[2], Mathilde Gallice[3], Christophe Chiquet[3], Nathalie Vayssière[1], Isabelle Berry[1], Yves Trotter[1], Vincent Soler[1], Carole Peyrin[2] *, Benoit R. Cottereau[1, 4] *

[1] CerCo UMR 5549, CNRS – Université Toulouse III, 31059 Toulouse, France.
[2] Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LPNC, 38400 Saint-Martin d'Hères, France
[3] Department of Ophthalmology, Grenoble Alpes University Hospital, 38700 La Tronche, France.
[4] IPAL, CNRS IRL 2955, 117418 Singapore.
[#] These authors contributed equally to this work.
* Co-senior authors.

Corresponding authors: Célia Michaud ('celia.michaud@cnrs.fr') and Benoit Cottereau ('benoit.cottereau@cnrs.fr')

Number of pages: 51
Number of figures: 7 + 7 supplementary
Number of tables: 2 + 1 supplementary
Number of words
      Abstract: 239
      Introduction: 652
      Discussion: 1550

Competing interests: The authors have no competing interests or conflicts of interest to disclose.

Ethics Statement: All participants gave their written informed consent and the protocol was approved by the French Ethics Committee (CPP) of Sud Mediterranée IV (2022-00957-36).

Data and code availability: MRI and fMRI data are provided at the following link: These data have undergone spatial normalization, in compliance with the General Data Protection Regulation (GDPR) under which this project is carried out. Data used for the analyses reported in Figures 6 and 7 (differences of PSCs and motion direction discrimination thresholds) are provided here:
https://osf.io/gpxhz/?view_only=43a1f1993d8742e6ba85ed4889b99a75

Acknowledgments: This study was supported by a French grant from the Agence
Nationale de la Recherche (ANR-21-CE28-0021, ANR PRC ReViS-MD; awarded to C.P.
and B.R.C.). This work was performed on the Inserm/UPS UMR1214 Technical Platform of
Toulouse and the IRMaGe platform member of France Life Imaging network (grant ANR-11-
INBS-0006) for Grenoble. We thank Judith Eck for helping on fMRI analysis, and Ilia
Korjoukov and Louise Kauffmann for helping develop our MRI paradigm.
48
49

Abstract: Previous studies on animal models suggested that visual areas involved in motion processing could undergo important cortical reorganizations following retinal damages. This could have major implications for patients suffering from macular degeneration (MD), one leading cause of vision loss. Here, we performed fMRI recordings in a group of maculopathy patients (including individuals suffering from age-related macular degeneration or from Stargardt's Disease) and a control group to characterize the motion processing cortical network in MD patients and determine whether this network is modified following the onset of the scotoma. We used an experimental protocol based on random-dot kinematograms (RDKs) classically employed to characterize motion-selective areas in the brain. To ensure that the visual information processed by the two groups was equivalent, the visual field in each control participant was masked using an artificial scotoma directly derived from clinical measurements in their paired patient. We found that in MD patients, translational motion elicited significant and robust activations in a restricted cortical network which included the human V5/MT+ complex (hMT+), areas V3A and V6, and a portion of primary visual areas (V1, V2 and V3) connected to peripheral vision. Importantly, the same patterns of responses were also observed in control participants. Moreover, the extent and strength of activation within these motion-selective areas did not differ significantly between the two groups. Altogether, these results suggest that in humans, the motion-selective network does not undergo significant large-scale cortical reorganizations following the onset of MD.

Significance statement: Motion processing in the visual cortex of patients with macular degeneration has never been characterized. Here, we performed fMRI recordings in 7 maculopathy patients and found robust motion-selective activations in a cortical network which included the human V5/MT+ complex (hMT+), areas V3A and V6, and a portion of primary visual areas connected to peripheral vision. These activations closely align with

those reported in participants with normal vision in the literature and do not significantly differ

from those measured in a group of age and gender-matched control participants who viewed

the motion stimuli with a matched artificial scotoma. Altogether, our results suggest that the

motion-selective network does not undergo significant large-scale reorganizations in

maculopathy patients following the onset of the scotoma.

1. Introduction

Macular degeneration (MD) is a visual pathology that alters the outer layers of the central

retina, resulting in central vision loss. Its most common form is age-related macular

degeneration (AMD) which develops after 60 years and currently affects over 67 million of

patients in Europe (Li et al., 2020). Macular degeneration can also affect younger

populations under the form of Stargardt disease (STG) which usually onsets during

adolescence but can also appear later in life (Late-Onset Stargardt Disease; Westeneng-van

Haaften et al., 2012). Despite their different origins, both diseases result in a loss of visual

acuity and decreased contrast sensitivity, with severe consequences on patient's quality of

life as they impair recognition and mobility (Hassan et al., 2002; Kuyk & Elliott, 1999; Marron

& Bailey, 1982; Wood et al., 2009). MD patients usually rely on peripheral vision, which has

a much lower spatial resolution compared to central vision, to process their visual

environment (Musel et al., 2011; Peyrin et al., 2017; Tran et al., 2011). Most of them develop

a preferred retinal locus (PRL) outside the scotoma for ocular fixation (Fletcher & Schuchard,

1997). This overuse of peripheral vision was hypothesized to lead to cortical reorganizations

in visual cortex (Baker, 2005), notably in neural populations which previously processed

central vision. However, the associated neural mechanisms remain poorly understood. Most

studies on cortical reorganisation in MD patients focused on the primary visual cortex (see

however Little et al., 2008 or Ramanoël et al., 2018). Recordings using functional magnetic

resonance imaging (fMRI) demonstrated that stimuli presented in peripheral vision activated

portions of area V1 connected to the scotoma (Dilks et al., 2009), but it is now accepted that

this effect is dependent on the cognitive tasks and low-level visual features, and only reflect

unmasking of cortical feedbacks (Masuda et al., 2008, 2021; Morland, 2015). Interestingly,

recent studies on animal models suggested that cortical reorganization after MD can be

more pronounced in higher-level visual cortex, notably along the dorsal pathway. Using

fMRI, Shao et al., (2013) found that visual responses in area MT, a dorsal region processing

motion, were significantly modified in a macaque monkey with juvenile MD. In a cat model,

Burnat et al., (2017) showed that the posteromedial lateral suprasylvian area PMLS (the

homologue of primate cortical area V5/MT; Villeneuve et al., 2006) and to a lesser degree

area 7 (which is also motion sensitive; Pigarev & Rodionova, 1998) had enhanced activity

following the induction of a central retinal lesion in adult animals. These enhancements were

associated with better performances in a motion task. In humans, a recent study showed

that functional connectivity between the portions of early visual areas (V1, V2, and V3) that

respond to peripheral vision and MT was enhanced in individuals with central vision loss

(Fleming et al., 2024). Another study in psychophysics reported that self-induced motion

perception from peripheral stimuli was increased in MD patients (Tarita-Nistor et al., 2008),

suggesting some degree of plasticity in cortical areas processing motion. It is thus possible

that neural responses in the motion-selective cortical network of MD patients are enhanced

following the onset of their scotoma but this hypothesis has never been tested using

functional neuroimaging.

Here, we employed an fMRI protocol commonly used to identify cortical areas selective to

motion (see e.g., R. Tootell et al., 1995 or Zeki et al., 1991) in 7 MD patients and 7 age and

gender-matched controls, whose central visual field was masked using simulated scotomas

of the same positions and dimensions as those of their corresponding patients. Our main

objectives were to characterize the motion processing cortical network in MD patients and to

determine whether this network undergoes significant large-scale reorganisations following

the onset of the scotoma. We tested if the cortical areas within this network are similar or

different between the two populations and whether their associated BOLD responses are comparable. To complete our analysis, we performed psychophysical measurements to assess the motion discrimination thresholds in both groups.

2. Methods

2.1 Participants: Seven maculopathy patients (age range: 50 - 81) were recruited from the Retina Center of the Purpan Hospital of Toulouse and from the Ophthalmology center of the Grenoble Hospital (CHUGA) in France (see Table 1). Five of them were diagnosed with age-related macular degeneration (AMD), while the two others suffered from late-onset Stargardt disease. They all had dense central bilateral scotomas and a stable preferred retinal locus in at least one eye. None of them suffered from other visual illnesses. The experiments were also conducted in 7 age- and gender-matched controls with normal (or corrected to normal) vision. Age differences between patients and controls were all less than two years except for the control of MD2 (81) who was 76 (despite a broad screening procedure, we did not manage to find an older volunteer). Importantly, each control participant underwent the experiments in the same center as their paired patient. None of the participants suffered from general long-term illnesses. In order to evaluate their cognitive functions and their psychological states, they all underwent the Mini-Mental State Examination (MMSE) and the Beck Depression Inventory (BDI-II), respectively. All of them had a score above 23 for the MMSE and below 19 for the BDI-II. All participants gave their written informed consent and the protocol was approved by the French Ethics Committee (CPP) of Sud Mediterranée IV (2022-00957-36).

Subjects	Sex	Age	Diagnosis	Visual acuity LE	Visual acuity RE	Tested eye	Scotoma radius deg	PRL polar angle (θ), deg	PRL eccentricity (ρ), deg
MD1	F	75	AMD	0.14	0.15	L	9.72	-27.4	10.4
MD2	M	81	AMD	0.03	0.48	R	5.3	143.9	1.1
MD3	F	66	AMD	0.6	0.23	L	7.31	157.3	2.1
MD4	M	73	AMD	0.23	0.22	L	9.84	28.5	9.58
MD5	M	76	AMD	0.05	0.23	R	6.15 12.31*	-151,5	0.9
STG1	F	50	STG	0.4	0.7	L	5.18	0	0
STG2	M	69	STG	0.25	0.25	R	5.87	-90	5.22
CTRL1	F	74	-	1.54	1.80	R	-	-	-
CTRL2	M	76	-	1.03	1.13	R	-	-	-
CTRL3	F	67	-	1.11	0.61	L	-	-	-
CTRL4	M	71	-	1.27	1.73	L	-	-	-
CTRL5	M	77	-	0.96	1.71	R	-	-	-
CTRLSTG1	F	52	-	1.31	0.99	L	-	-	-
CTRLSTG2	M	68	-	1.43	2	R	-	-	-

Table 1: Chara*cteri*s*tics of MD patients and age-matched controls included in the study. Precisions*

about the diagnosis of patients (AMD for Age-related Macular Degeneration, STG for late-onset

Stargardt's disease), the decimal visual acuity for the left (L) and right (R) eye, as well as the polar

coordinates (angles and eccentricities) of the Preferred Retinal Locus (PRL) are included. Patients

and their controls carried out the experiments in the same centers, i.e in Toulouse or in Grenoble.

Controls for MD patients 1 to 5 are labelled CTRL1 to CTRL 5. Controls for STG patients STG1 and

STG2 are labelled CTRLSTG1 and CTRLSTG2. It should be noted that because MD5 had an

elongated scotoma, we used an ellipse rather than a disk for the artificial scotoma of his control. In

this case, the two values provided in the 'scotoma radius' column (marked by an asterisk) provide the

lengths of the ellipse semi-minor and semi-major axes.

For each participant, the eye with the better acuity was determined using the Sloan letters of

the Freiburg Visual Acuity Test (Bach, 2007). In most participants, this eye was stimulated

during the experiments and the other eye was patched. Only patients MD1 and STG1 were

stimulated in the other eye because it was the only one with a stable preferred retinal locus

(PRL, see below). In their case, the eye with the better acuity was patched.

2.2 Scotoma size: For each patient, an ophthalmologist delineated the border of the scotoma from the fundus of the eye captured by standard and autofluorescence imaging (see Figure 1-A for MD3, other examples are provided in Supplementary figure 1-1). The area of the scotoma (in square degrees of visual angles) was extracted using MATLAB (MathWorks, Natick, MA, USA). This area was used to compute the simulated scotoma of the age- and gender-matched controls (see below).

2.3 Position of the Preferred Retinal Locus (PRL) in maculopathy patients: MD patients usually develop a new locus of ocular fixation following the onset of their scotoma. To test for the presence of such preferred retinal locus (PRL) and to determine its location, we performed spectral-domain optical coherence tomography (see Figure 1-B). During the measure, patients were instructed to fixate a light produced by the OCT. Patients with a developed PRL naturally used it to direct their gaze, thereby allowing us to record its retinal coordinates. This process was repeated three times with a short break in between acquisitions, to ensure that the measure was stable (see Contemori et al., 2024). Table 1 reports the average PRL locations across these three measures (see Maniglia et al., 2018 or Guénot et al., 2022 for more details). It is important to note that two patients (MD5 and STG1) relied on central vision for fixation, indicating that they retained some residual visual function near their fovea despite significant damages to their maculas, as shown in Supplementary Figure 1-1. As will be demonstrated, the inclusion of these patients does not introduce biases into our fMRI results.

Figure 1: Estimation of the scotoma size and PRL position in one typical patient (MD3). These values are used to define the artificial scotoma masking the central visual field in the associated control participant. It should be noted that all retinal images in this figure have been inverted horizontally to be aligned with the visual field. The letters on panel A and B specify the orientation of the images (L: Left; R: Right; D: Down; U: Up). A) An autofluorescence imaging of the eye fundus is acquired. The border of the scotoma is delineated and the associated surface is used to calculate the radius of the scotoma in degrees of visual angle, given that 288µm on the image corresponds to 1 degree of visual angle. B) Patients were instructed to fixate a central light and we measured the x and y position of their PRL with spectral-domain optical coherence tomography (Spectralis Optical Coherence Tomography [OCT]; Heidelberg Engineering, Heidelberg, Germany). This process was repeated three times (leftward panel) and the final coordinates (green dot on the rightward panel) were the average across these three measures. Here, the position of the fovea is marked by a red dot. From the surface of the scotoma and the position of the PRL, it is possible to simulate the patient's visual field. C) During the fMRI experiments, MD patients were instructed to fixate a central cross (shown here in green) with their PRL. Controls had to fixate a red cross at the position corresponding to their paired patient's fovea while an artificial scotoma was displayed in dark gray. The dotted lines (not shown during the experiments) specify the four quadrants of the patients' and controls' visual fields in this case: upper-left ('UL'), upper-right ('UR'), lower-left ('LL') and ('LR'). The screen (in light gray) dimensions provided here (20° x 15°) are those used in the two MRI centers. The percentage displayed on the top of the graphics specifies the visible portion of the screen (i.e., the surface that is not obscured by the artificial scotoma) in this case.

%%%

2.4 Artificial scotoma and viewing conditions in control participants: For each control participant, we masked the central visual field using an artificial scotoma as in previous studies (Christen & Abegg, 2017; Gupta et al., 2018). This simulated scotoma had the same surface as the one measured in the associated patient (see the '*Scotoma size*' section) and was centered on the fovea (except for CTRL2 and CTRL 3 where the artificial was shifted to the left because MD2's and MD3's scotoma occupied a larger portion of the

left visual hemifield). Control participants were instructed to gaze on a fixation cross that we

displayed at the position of their paired patient's fovea (see the red cross on Figure 1-C). In

this case, the center of the screen fell on the visible part of their visual field, but at the PRL

location of their paired patient. For all control participants, artificial scotomas were simplified

as a disk except for CTRL5 for which we used an ellipse to more closely match the real

shape of MD5 scotoma, which was very elongated (see Supplementary figure 1-1). This

protocol permits patients and controls to receive comparable visual inputs. It notably

equalizes the position and surface of the occluded part of the visual field (see Guénot et al.,

2022).

2.5 Motion stimuli: Motion stimuli were adapted from the experimental protocol of a

previous electrophysiological study (Graziano et al., 1994). They consisted in translational

patterns defined from randomdot kinematograms (RDKs) (see Figure 2-A) generated with

Matlab (R2017a). The RDKs contained bright non-overlapping dots (diameter: 0.2°) moving

on a homogenous dark background with a high contrast (100%). They had a density of

0.3945 dots per square degree of visual angle. When a dot reached the border of the display

screen, it was immediately relocated at a random position. This process permits to equalize

mean luminance and dot density across the screen during the experiments. Each dot moved

at a velocity of 7°/s, which corresponds to the average preferred speed of neurons in

macaque MT (Priebe et al., 2006). This speed is in agreement with those used in previous

fMRI studies on motion processing (6°/s in Zeki et al., 1991, 8°/s in (Huk et al., 2002). These

translational motion stimuli were used for the fMRI recordings (see the next section) and for

the psychophysical measurements (see the '*Experimental protocol for the psychophysical*

measurements' section).

2.6 Experimental protocol for fMRI measurements: In both Toulouse and Grenoble

MRI platforms, stimuli were displayed on a large screen and covered 20°x15° of visual

angle. fMRI recordings were performed using a block-design. Each run consisted of 312 s

and were divided into 6 identical cycles of 52 s (see Figure 2-A). Each cycle started with a

baseline period of 10 s during which only a red fixation cross of 1° of visual angle was

displayed on a black screen. It was followed by a block of 16s of uniform motion, then by

another 10 s of baseline and finally by a block of 16 s with static dots. During motion blocks,

a new pattern of translational motion during which all the dots moved either upward or

downward was presented every 2 s. These patterns last 1.8 s and were separated by short

periods (0.2 s) during which the screen went black and only the fixation cross was

presented. The same principle was applied during the blocks of static dots (a new pattern

was displayed for 1.8 s every 2 s). During both translational motion and static dot blocks, an

artificial scotoma masked the central vision of control participants.

Figure 2: Experimental protocols. A) fMRI recordings were performed using a block-design. Runs

lasted 312 s and contained 6 identical cycles of 52 s. Each cycle started with a baseline period of 10 s

during which only a red fixation cross was displayed on a black screen. It was followed by a block of

16 s of uniform motion (shown here in blue), then by another 10 s of baseline and finally by a block of

16 s with static dots (in red). During motion blocks, a new pattern of translational motion during which

all the dots moved either upward or downward was presented every 2 s. These patterns last 1.8 s and

were separated by short periods (0.2 s) during which the screen went black and only the fixation cross

was presented. The same principle was applied during the blocks of static dots (a new pattern was

displayed for 1.8 s every 2 s and patterns were separated by 0.2 s). B) During psychophysical

recordings, participants had to report the perceived motion direction of the stimuli (either upward or

downward). We manipulated motion coherence and estimated thresholds corresponding to 80 percent

of correct detections. For illustration, an upward motion trial is shown at 20% of coherence. Signal

dots are shown in white and noise dots in gray for better visibility on the figure (all the dots were white

during the experiment). During all the recordings (fMRI and psychophysics), participants had to fixate

a red cross that was central for patients (who used their PRL) and at the position corresponding to

their paired patient's fovea for controls. This red fixation cross was always present on the screen.

%%

This protocol is classically employed to identify motion selective cortical areas (Zeki et al.,

1991; Watson et al., 1993; Tootell et al., 1995a; Huk et al., 2002). The visual display of the

stimuli and its temporal synchronization with the acquisition of the fMRI data was controlled

using the EventIDE software (Okazolab). A red fixation cross was displayed during all the

functional recordings, in the center of the screen for patients (who directed their gaze with

their PRLs) and at the position corresponding to their paired patient's fovea for controls (see

above). As shown in the central columns of Figures 3 and 4, the visible portion of the stimuli

covered most of the screen for the majority of participants. The exceptions were MD5 and

his control, for whom 42% of the screen was visible. In order to make sure that all

participants correctly directed their gaze and were focused on the visual stimuli, a white

circle (0.75° of radius) was randomly displayed for 200 ms on the center of the screen during

the runs (10 displays per run on average). Its size and duration were determined from pilot

experiments. Participants were instructed to press a button on a gamepad whenever they

detected it. Even though this task was not easy (notably for patients who were fixating with

their PRLs), they all performed it well (mean percentage of correct detection for patients: 80;

and controls: 90) and we thus kept all the runs for further analyses.

2.7 (f)MRI recordings: (f)MRI recordings were performed at 3T (Achieva dStream 3.0T

TX, Philips, NL) at Toulouse NeuroImaging Center (ToNIC, Toulouse, France) and at

IRMaGe MRI facility (Grenoble, France). Importantly, both MRI platforms are equipped with

the same MR hardware configuration (with Quasar Dual gradient system and dStream

architecture supporting MultiBand-accelerated acquisitions, 32 channel head coil). They

were operated using the same software version (R5.7.1). This permits to minimize the

differences between the recordings performed in the two cities, although we emphasize that

each patient from one center was matched with a control from the same center. We acquired

two high-resolution anatomical images of the participant's brain using a T1-weighted

magnetization-prepared rapid gradient-echo (MPRAGE) sequence (repetition time, TR = 8.1

ms; echo time, TE = 3.7 ms, flip angle = 8°; FOV: 240 x 240 mm; voxel size = 1 x 1 x 1 mm;

170 sagittal slices). These anatomical images were first co-registered and then averaged

together to be used as a reference to which the functional images from all the experiments

were aligned. Functional images were collected with a T2 * weighted gradient-echo

echoplanar imaging (EPI) sequence (TR = 2000 ms, TE = 30 ms, flip angle = 75°, SENSE

factor = 1.6; multi-band (MB) = 3; FOV: 224 x 224 mm; matrix size: 112 x 110 mm; voxel

size = 2 x 2 x 2 mm, 63 transverse slices, dyn = 156).

2.8 (f)MRI data analyses: MRIQC v.13.1.0 was used to check the quality of all our

functional recordings (Esteban et al., 2017). We did not observe any out-of-range runs in our

data, indicating negligible head-movement. We thus kept all the measurements for further

analyses. Pre-processing of fMRI data included slice scan time correction, 3D motion

correction using trilinear/sinc interpolation, and a linear trend followed by a high-pass filtering

using Fourier analysis with a cut-off at 3 cycles. For each participant, functional data were

co-registered on the anatomy. Functional and anatomical data were brought into anterior

commissure - posterior commissure (ACPC) space using cubic spline interpolation and then

transformed into standard Talairach (TAL) space (Talairach & Tournoux, 1988). For further

analyses, the volume time courses were spatially smoothed at 4 mm FWHM.

Individual analyses were performed on each node of the participant' cortical tessellations by

fitting a general linear model (GLM) to the corresponding blood oxygen level-dependent

(BOLD) signal. The model contained 2 main regressors, representing the 2 experimental

conditions: uniform motion (either upward or downward) and control (static dots). These

regressors were convolved with a model of the human hemodynamic response function. We

also introduced 8 regressors of non-interest in our model. 6 of them corresponded to the

translation and rotation of the head estimated during the motion-correction step of the pre-

processing (see above). The two last ones were time-courses captured within the ventricles

and the white-matter. This operation permits to reduce the noise in the data (see e.g.,

Audurier et al., 2022). For each participant, the beta weights obtained from the GLM were

subsequently used to perform univariate analyses (t-scores) at the whole-brain level. All

these analyses were realized using the BrainVoyager software (v23, Brain Innovation,

Maastricht, the Netherlands) (Goebel et al., 2006). Results are shown on inflated surfaces,

obtained by segmenting the white and gray matter, as well as the cerebrospinal fluid, and

reconstructing a surface at the boundary of the white and gray matter. Statistical maps are

mainly thresholded at p-value < 10^{-3} (uncorrected), as is often the case in individual-level

fMRI analysis (see e.g., Aedo-Jury et al., 2020).

Group-level analyses were performed within independently defined regions of interest

(ROIs). These ROIs were given by spheres in the volumetric space centered on the

Talairach coordinates provided in previous studies for areas hMT+ (Watson et al., 1993),

V3A (R. B. H. Tootell et al., 1997) and V6 (Pitzalis et al., 2010). Within each ROI, we

computed the differences between the averages percentages of signal changes (PSCs)

during the translational motion condition and during its control condition (static dots). For

each group, we first determined whether these differences of PSCs were significantly

different from zero across participants. We used Wilcoxon sign-rank tests because it is non-

parametric and thus does not assume that the data are normally distributed. Then, we tested

whether the differences of PSCS were different between the two groups (MD patients and

controls), also using Wilcoxon sign-rank tests. Here, data were paired between each patient

and his/her control to reflect the fact that each pair received the same visual stimulations. To

complete our statistical analyses and determine how much evidence favors the alternate

hypothesis (H_1: the differences of PSCs are not the same between the two groups) against

the null hypothesis (H_0) in a given ROI, we also computed the Bayes factor using the R-code

open source JASP program. This factor is given by the ratio of the likelihood of the alternate

hypothesis (H_1) to the likelihood of the null hypothesis (H_0). Values weaker than one favors

the null hypothesis whereas values greater than one favors its alternative (Raftery, 1995).

We used spheres with a radius of 10 mm, a value which leads to a cortical surface in line

with the one reported for area hMT+ (see Kolster et al., 2010). It should be noted that the

analyses were also repeated with radius values of 5 and 15 mm to make sure that the choice

of this parameter did not influence our results.

2.9 Experimental protocol for psychophysical measurements: The psychophysical

measurements were performed separately from the fMRI recordings, and outside the MRI

scanner. During the experiment, participants sat in a chair whose height was adapted in

order to adjust the height of the eyes to the center of the screen. Their head was placed on a

head-support device clamped on top of a table and equipped with both chin and forehead

supports. The chair and head-support devices were positioned so as to ensure a fine

alignment between the participants' head and trunk axes. Participants had to keep this

position as constant as possible. Dot stimuli of 0.2° in size were presented on computer

screens using a large portion of the visual field (58.1° × 43.8° of visual angle in Toulouse

and 48.6° x 30.7° in Grenoble) to maximize coverage of peripheral vision. As for the fMRI

recordings, during the whole experiment, participants had to fixate monocularly with the

same study eye (the other eye was patched) a red cross. This cross was central for patients

(who used their PRL) and at the position corresponding to their patient fovea for age-

matched controls.

We used the same random dot kinematogram stimuli as for the fMRI experiments (see the

'*Motion stimuli*' section), except that here, each dot had a limited lifetime of 200 ms (12

frames at 60 Hz). At the end of its lifetime, it was randomly reassigned to a new spatial

position. To avoid a coherent flickering of the stimulus every 200 ms, each dot initial age was

randomly picked between 0 and 166 ms (11 frames) at the beginning of each trial. We chose

to manipulate lifetime to permit a direct comparison of the psychophysical results with those

obtained in a previous study using the same paradigm, but performed on other MD patients

(Guénot et al., 2022). It should be noted that we used unlimited lifetimes for the fMRI

recordings in order to maximize the BOLD responses of motion selective areas, and hence

to facilitate their localization.

A forced binary decision task was used to estimate motion discrimination thresholds. Stimuli

were presented in blocks of 64 trials. Each block lasted about 3 min. Each trial started with

the presentation of the stimulus for 200 ms. Participants had to report whether the motion of

the stimulus was upward or downward. They were instructed to respond as quickly as

possible while maximizing their performances. Responses reported after 2 s were

considered as incorrect. After each trial, an auditory feedback indicated whether the chosen

direction was correct or not. During each block, we manipulated motion coherency (i.e., the

percentage of dots moving along the same direction while the other dots had random

directions) and estimated the thresholds corresponding to 80% of correct detection using an

adaptive Bayesian approach (QUEST, see below). Participants completed 4 blocks. The first

block was considered as a training and not included in the analyses. Breaks were included

(3 minutes on average) between blocks to reduce fatigue. The whole experiment was

completed in one session of about 20 minutes.

2.10 Robust estimation of motion coherence thresholds: Because it is complicated

for MD patients to perform long psychophysical experiments, we used the QUEST adaptive

procedure to obtain rapid, efficient and robust estimations of motion coherence thresholds

(Guénot et al., 2022, 2023). QUEST is based on a Bayesian approach assuming that the

psychophysics function underlying the participant performance follows a Weibull distribution.

During a block, the estimated parameters of this function were updated after each trial on the

basis of the participant's response. Coherence values corresponded to the current maximum

likelihood estimate of the threshold. We fixed the maximum number of trials at 64 as it was

previously shown that this value leads to robust thresholds in most circumstances (Watson &

Pelli, 1983). We used an initial threshold value of 58 % based on previous recordings using

the same stimuli in MD patients (Guénot et al., 2022).

3. Results

427   In this section, we first detail the fMRI activation patterns observed in our MD patients and

428   their controls. Next, we describe the results of the psychophysical experiment and their

429   relationship with BOLD responses.

3.1 Motion-selective responses in the cortex of MD patients and their controls

Figure 3 shows the motion-selective activations (t-scores, p-value < 0.001, uncorrected)

obtained in one typical MD patient (MD3, upper panel) and her control (CTRL3, lower panel)

displayed on enlarged lateral views of their inflated individual cortical surfaces. Motion-

selective activations in the other participants are shown in Figure 4. Activations displayed on

medial views for all participants are shown on Figure 5. In all these figures, cortical nodes

with significantly stronger responses to translational motion than to static dots appear in hot

colors (yellow/red) while nodes with significantly stronger responses to the opposite contrast

are shown in cold (blue) colors. Data were thresholded at p-value < 10^{-3} (uncorrected).

Figure 3: Comparison between BOLD responses to the translational motion condition and to its control condition (static dots) in one representative patient (MD3) and her age- and gender-matched control (CTRL3). Activations are shown on inflated individual cortical surfaces. Gyri appear in pale gray while sulci are shown in darker gray. Cortical nodes with significantly stronger responses to translational motion appear in yellow/red colors while nodes with significantly stronger responses to static dots are shown in blue. Data were thresholded at p-value < 10^{-3} (uncorrected). It should be noted that area V3A in the right hemisphere of MD3 was detected using a threshold p-value of 10^{-2} and it is thus not visible here. Colored lines provide the anatomical positions of the Inferior Occipital Sulcus (IOS, in olive green), of the Lateral Occipital Sulcus (LOS, in green), of the Transverse Occipital Sulcus (TOS, in purple), of the ventral part of the Anterior Occipital Sulcus (AOSv, in cyan), and of the Superior Temporal Sulcus (STS, in red). The central panel illustrates visual stimulation in this case. During the fMRI recordings, MD3 fixed the center of the screen (green cross) with her PRL while CTRL3 gazed at the position of MD3's fovea (red cross). The dark disk is the artificial scotoma (see Figure 1 for more details on its definition). The dotted lines (not shown during the recordings)

specify the positions of the horizontal and vertical meridians. We also provide the coherence

thresholds (CoTh) estimated through psychophysical measurements.

%%

In all participants, translational motion elicited significantly stronger responses in a cortical

network that is mainly localized within the dorsal visual pathway (i.e., in occipito-parietal

areas) and within early visual areas. It should be noted that this network is very restricted

when compared to the one obtained by contrasting responses to visual stimulation versus

baseline (ocular fixation) and which covers the majority of the posterior part of the brain and

notably includes areas along the ventral pathway (lateral and medial views of the

corresponding activations maps are provided in Supplementary Figures 4-1 and 4-2). In the

following, we first focus on responses within cortical regions that were consistently reported

as motion-selective in the literature. Then, we describe our results in early visual areas and

in other parts of the brain.

3.2 Activations in area hMT+: In all MD patients and their controls, we observed

significantly stronger responses to translational motion than to static dots within (or posterior

to) the ventral part of the anterior occipital sulcus (AOSv, in cyan in Figure 3), in between the

inferior (IOS, in olive green) and the lateral (LOS, in green) occipital sulci (see Figures 3 and

4). The Talairach coordinates of the local maxima (t-scores) measured in this region are

provided in Table 2. Averages values in the left and right hemispheres for patients ([-42±2; -

71±2; 5±2] and [41±2; -66±2; 2±2]) and controls ([-42±2; -64±2; 4±3] and [42±2; -63±2; 3±2])

were similar. These anatomical locations and coordinates are in very good agreement with

those reported in the literature for area hMT+ (e.g., [-41; -69; 2] and [41; -67; 2] in Watson et

al., 1993 or [-47; -76; 2] and [44; -67; 0] in Dumoulin, 2000). To confirm that our data

reflected motion-selective activations in this area, we used independently defined regions of

interest (ROIs) centered on the Talairach coordinates provided by Watson et al., (1993) for

hMT+ and tested whether the differences between the percentages of signal change (PSCs)

measured during the translational motion condition and during its control (static dots) were

significantly different from zeros across participants in this ROI (see the '*Methods*' section).

This was the case for both groups (Wilcoxon sign-rank tests, p-value = 0.0156 in MD

patients and p-value = 0.0078 in controls, see Figure 6-A). Importantly, we did not find

significant differences between the differences in PSCs observed in this ROI for the two

groups (Wilcoxon paired sign-rank tests, p-value = 0.1094, Bayes factor = 1.6). As an

additional analysis, we also considered the activation extent within this ROI and found that

they did not significantly differ between the two groups (see Supplementary figure 6-1).

Altogether, these results suggest that both MD patients and their controls have motion-

selective responses in area hMT+ and that these responses do not significantly differ

between the two populations.

3.3 Activations in area V3A: In all MD patients and their controls, we observed

significantly stronger responses to translational motion within the transverse occipital sulcus

(in purple on Figure 3, see also Figure 4). For some participants, we had to decrease our

threshold to a p-value of 10^{-2}, uncorrected) at similar Talairach coordinates between the two

groups ([-18±2 -81±2 16±3] and [21±3 -80±2 22±4] for patients and [-22±2 -80±1 25±3] and

[23±1 -76±1 20±2] for controls on average (see Table 2). These anatomical locations and

coordinates are in very good agreement with those reported in the literature for area V3A

(e.g., [±14; -84; 19] in Tootell et al., 1997 or [±18; -78; 25] in Pitzalis et al., 2006). For both

groups, we confirmed that our data reflected motion-selective activations in this area from

statistical analysis within an ROI centered on the Talairach coordinates provided by Tootell

et al. (1997) (Wilcoxon sign-rank tests, p-value = 0.0078 in MD patients and p-value =

0.0234 in controls, see Figure 6-B and above). We did not find significant differences

between the differences of PSCs observed in this ROI for the two groups (Wilcoxon paired

sign-rank tests, p-value = 0.469, Bayes factor = 0.5), nor between their activation extents

(see Supplementary figure 6-1). Altogether, these results suggest that both MD patients and

their controls have motion-selective responses in area V3A and that these responses do not

significantly differ between the two populations.

3.4 Activations in area V6: In six MD patients and their controls (all participants except

MD3 and CTRL3), we observed significantly stronger responses to translational motion in

the upper part of the cuneus, within the posterior branch of the parieto-occipital sulcus (POS)

at similar Talairach coordinates between the two groups ([-12±2 -76±2 29±3] and [12±3 -

73±3 31±3] for patients and [-14±2 -74±2 32±3] and [11±1 -75±1 35±3] for controls on

average, see Table 2). These anatomical locations and coordinates are in very good

agreement with those reported in the literature for area V6 (e.g., [±9; -78; 34] in Pitzalis et

al., 2010). For both groups, we confirmed that our data reflected selectivity to translational

motion in this area from statistical analysis within an ROI centered on the Talairach

coordinates provided by Pitzalis et al. (2010) (Wilcoxon sign-rank tests, p-value = 0.0156 in

MD patients and p-value = 0.00781 in controls, see Figure 6-B and above). We did not find

significant differences between the differences of PSCs observed in this ROI for the two

groups (Wilcoxon paired sign-rank tests, p-value = 0.297, Bayes factor = 0.7), nor between

their activation extents (see Supplementary figure 6-1). Altogether, these results suggest

that both MD patients and their controls have motion-selective responses in area V6 and that

these responses do not significantly differ between the two populations.

3.5 Activations in early visual areas and relationship to retinotopic properties:

In all participants, we also observed motions selective activations below and above the most

anterior part of the calcarine sulcus (see Figure 5). These regions correspond to the portions

of primary visual areas (V1, V2 and V3) connected to peripheral vision. Their stronger

responses to translational motion are not surprising given that a vast proportion of their

neurons are motion-direction selective (Dumoulin et al., 2003). Here as well, we did not

observe different patterns of activations in statistical parametric maps between groups

(Figure 5). It should be noted that for patients whose scotoma covers a larger surface in one

visual hemifield (because their PRL is lateralized) and for their controls (MD1/CTRL1, MD2/CTRL2, MD3/CTRL3 and MD4/CTRL4), motion-selective activations were generally stronger in the hemisphere ipsilateral to the scotoma (i.e., contralateral to the visual hemifield that receives more visual inputs) which reflects the retinotopic properties of early visual areas (see e.g., Benson & Winawer, 2018). In V3A, hemispheric asymmetries were also observed for these participants but they were less pronounced, reflecting the fact that neural receptive fields in this area are bigger than in areas V1, V2 and V3 (e.g., Barton & Brewer, 2017 reported population receptive fields with radiuses between 4 and 6° at eccentricities comparable to those in our study). We did not observe hemispheric asymmetries in areas hMT+ and V6, probably because these areas have very large receptive fields which overlap with the ipsilateral visual hemifield (for hMT+, see Huk et al., 2002; for V6, see Pitzalis et al., 2010). For hMT+, this is notably true in its hMST functional subdivision.

3.6 Additional activations: Beyond the activations reported above, we also observed motion-selective responses in other parts of the brain but they were not as reliable and robust across participants. In some patients and their controls (MD2/CTRL2, MD3/CTRL3, MD4/CTRL4, STG1/CTRLSTG1 and STG2/CTRLSTG2, see also on the right hemisphere of CTRL5), we notably observed motion-selective activations in the fundus of the superior temporal sulcus (STS, in red on Figure 3) and also (but less frequently) on its lower bank (MD3 in the right hemisphere, MD4 in the right hemisphere, CTRL5 in the left hemisphere) or within the middle temporal sulcus (MTS, CTRL2). These activations could correspond to the human posterior temporal sulcus (pSTS) area, which is selective to biological motion (Grossman & Blake, 2002) and which preferentially encodes peripheral vision (Barton & Brewer, 2017). The associated Talairach coordinates are in good agreement with those reported for this area in the literature (see supplementary Table 2-1) but an ROI-based analysis using a sphere centered on the coordinates provided by Barton and Brewer (2017)

did not confirm its implication (Wilcoxon sign-rank tests, p-value = 0.109 in both MD patients

and controls, see Supplementary Figure 6-2).

In some participants (MD3, MD4, CTRL1 and CTRL2), additional motion-selective

activations were also observed in the parietal cortex and notably within the intra-parietal

sulcus, where is located the ventral intraparietal (VIP) area, which is motion selective

(Bremmer et al., 2001).

Figure 4: Comparison between BOLD responses to the translational motion condition and to its control condition (static dots) in all participants, except MD3 and her control (CTRL3), whose data are provided on enlarged views in Figure 3. Significantly stronger responses to motion are shown on lateral views of the cortical surfaces for MD patients (leftward panels) and their controls (rightward panels). See Figure 3 for the details of the legend.

%%

		hMT+	V3A	V6
MD1	LH	-47; -71; 4	-17; -86; 13	-9; -74; 33
	RH	49; -66; -4	22; -84; 12	7; -77; 34
MD2	LH	-43; -66; 7	-22; -77; 18	-6; -80; 24
	RH	32; -66; 12	13; -78; 22	4; -78; 21
MD3	LH	-47; -66; 5	-16; -82; 14	X
	RH	38; -64; -2	29; -73; 32*	X
MD4	LH	-43; -73; -2	-27; -72; 9*	-17; -72; 23
	RH	49; -64; -5	29; -78; 27	20; -69; 24
MD5	LH	-39; -73; 13	-11; -77; 29	-13; -70; 32
	RH	40; -66; 7	18; -81; 33	19; -69; 34
STG1	LH	-33; -78; 0	-11; -87; 7	-12; -81; 38*
	RH	37; -70; -1	11; -84; 8	6; -74; 34*
STG2	LH	-42; -73; 8	-22; -83; 21	-12; -77; 21
	RH	43; -67; 5	24; -80; 19	18; -72; 38
CTRL1	LH	-44; -64; 15	-20; -79; 31	-16; -69; 34
	RH	49; -65; 11	24; -75; 21	13; -67; 45
CTRL2	LH	-42; -58; 7	-22; -73; 32	-19; -68; 46
	RH	38; -54; 6	21; -66; 28	10; -70; 45
CTRL3	LH	-36; -60; 3	-20; -77; 31	-15; -71; 31
	RH	38; -63; -3	26; -68; 24	14; -69; 27*
CTRL4	LH	-40; -70; -4	-21; -79; 17	-13; -72; 31
	RH	48; -68; 3	19; -82; 23	7; -75; 29
CTRL5	LH	-40; -69; -7	-21; -79; 14	-12; -84; 15
	RH	38; -70; 0	29; -77; 19	9; -81; 27*
CTRLSTG1	LH	-47; -63; 13	-20; -89; 26	-6; -75; 34
	RH	45; -59; 2	23; -82; 15	12; -80; 33
CTRLSTG2	LH	-47; -69; -2	-29; -87; 22	-18; -78; 30
	RH	35; -63; -1	18; -81; 10	10; -81; 36
Average patients	LH	-42±2; -71±2; 5±2	-18±2; -81±2; 16±3	-12±2; -76±2; 29±3
	RH	41±2; -66±2; 3±2	21±3; -80±3; 22±4	12±3; -73±3; 31±3
Average controls	LH	-42±2; -64±2; 4±3	-22±1; -80±2; 25±3	-14±2; -74±2; 32±3
	RH	42±2; -63±2; 3±2	23±1; ±-76±1; 20±2	11±1; -75±1; 35±3
Literature	LH	-41; -69; 2	-14; -84; 19	-9; -78; 34

	RH	41; -67; 2	14; -84; 19	9; -78; 34

Table 2: Talairach coordinates of the different motion-selective areas observed in our study (hMT+,

V3A and V6). Coordinates are provided for each MD patient and his/her control. The '*' symbol

indicates the coordinates for which a more permissive threshold has been used, i.e $p < 10^{-2}$. The table

also provides average coordinates across each group as well as average coordinates reported in

previous studies for these 3 areas (Watson et al., 1993 for hMT+; Tootell et al., 1997 for V3A and

Pitzalis et al., 2010 for V6).

%%%

We saw that two patients (MD5 and STG1) relied on central vision for fixation, indicating that

they retained some residual visual function near their fovea (see the 'Methods'). To ensure

that their inclusion did not bias our results, we replicated all statistical analyses described

above excluding these patients (and their matched controls) and found consistent effects

(see Supplementary Figure 6-3).

Figure 5: Comparison between BOLD responses to the translational motion condition and to its control condition (static dots) in all participants. Significantly stronger responses to motion are shown on medial views of the cortical surfaces for MD patients (leftward panels) and their controls (rightward panels). See Figure 3 for the details of the legend.

%%

●/▲MD1/CTRL1 ●/▲MD2/CTRL2 ●/▲MD3/CTRL3 ●/▲MD4/CTRL4 ●/▲MD5/CTRL5 ●/▲STG1/CTRLSTG1 ●/▲STG2/CTRLSTG2

Figure 6: Motion-selective activations in the hMT+ (A), V3A (B) and V6 (C) ROIs. For each patient

and their age-matched control (connected by lines), colored dots provide the differences between the

average (across voxels and hemispheres) percentages of signal changes (PSCs) during the

translational motion and control (static dots) conditions. For each group, the boxplots provide the first

quartile, median, last quartile, and the extreme PSC difference. Stars (* or **) indicate that the PSC

difference is significantly different from zero (p-values < 0.05 or <0.01, respectively). Statistical

analyses based on paired Wilcoxon sign-rank tests did not reveal significant differences between

activations in the two groups (p-values > 0.05 for the three ROIs).

%%%

3.7 Motion-direction discrimination thresholds and their relationship with

BOLD responses

The motion direction discrimination thresholds estimated in each MD patient and their age-

matched control (see the 'Material and Methods' section) are provided in Figure 7-A.

Thresholds varied between 0.37 and 0.99 in patients (0.37 for MD3 whose fMRI data are

also shown on Figure 3; 0.67 on average) and between 0.21 and 0.98 in control participants

(0.21 for CTRL3, see Figure 3; 0.58 on average). We did not find significant differences

between the thresholds measured in the two groups (Wilcoxon signed-rank test, p-value =

0.1698, Bayes factor = 0.25). Altogether, these results confirm Guénot et al.'s (2022) results

where motion direction discrimination thresholds did not significantly differ between MD

patients and their age-matched controls under identical experimental conditions (see Figure

5 in Guénot et al., 2022).

Figure 7: Behavioral results and relationship with cortical responses in motion-selective ROIs. A)

Motion-direction discrimination thresholds estimated in the two groups. The boxplots show the

different values for the first quartile, median, last quartile, and the extreme data points. The cross

represents the means for both groups. Dots provide the individual data points of the distributions. It

should be noted that they were slightly offset horizontally to improve their visibility. B) Differences

between the average percentages of signal changes (PSCs) during the translational motion and

control (static dots) conditions in the hMT+ ROI as a function of coherence thresholds. Dots and

triangles respectively correspond to MD patients and controls. Spearman correlation coefficients rho

(ρ) are shown (the associated p-values were respectively 0.1389 for patients and 0.5560 for controls).

Data in the V3A and V6 ROIs are respectively provided in panels C and D. The p-values associated

with the Spearman correlation coefficients were 0.1667 and 0.2357 in area V3A and 0.8397 and

0.5560 in area V6 for patients and control participants, respectively. The x-axis was cropped at 0.2 for

better visibility.

%%%

Next, we examine whether the performances measured in psychophysics are correlated with

fMRI activations in motion-selective areas. For each participant, panels B, C and D of Figure

7 show the differences between the averages percentages of signal changes (PSCs) during

the translational motion condition and during its control condition (static dots) in areas hMT+,

V3A and V6 as a function of motion-discrimination thresholds. For each group, we provide

the corresponding Spearman correlation coefficients. This measure tests whether the data

are monotonically related and do not assume normality. Although stronger responses to

motion stimuli were generally associated with lower perceptual thresholds (Spearman

coefficients were mostly negative), we did not observe any significant correlations in the data

(p-values > 0.5, see details in the legend of figure 7).

To complete these analyses, we also tested whether the size of the scotoma could influence

the percentages of signal changes measured in our ROIs but we did not observe significant

effects in both groups (see Supplementary Figure 7-1). This is probably because receptive

fields in these areas are large enough to capture visual inputs outside the scotoma (see also

the '*Activations in early visual areas and relationship to retinotopic properties*' section**).**

3.8 Control analyses to verify the robustness of our ROI-based approach

All the ROIs-based analyses in our study used spheres of 10 mm. To make sure that this

choice did not impact our results, we reproduced these analyses with spheres of 5 and 15

mm of radius and found the same patterns of results.

4. Discussion

The aim of this study was to characterize the cortical network processing translational motion in maculopathy patients. We recorded fMRI data in 7 patients and in 7 age- and gender-matched controls using an experimental paradigm based on RDKs classically employed to identify motion selective areas (Zeki et al., 1991; Watson et al., 1993; Tootell et al., 1995a; Huk et al., 2002). We also estimated motion-direction discrimination thresholds in each participant. Stimuli in each control participant were masked using an artificial scotoma of the same dimension as the scotoma of their paired patients, in order to equalize the visual inputs received by the two populations.

A previous study on an animal model (Burnat et al., 2017) demonstrated that inducing a central retinal lesion in adult cats enhanced neuronal activity in the posteromedial lateral suprasylvian area (PMLS), the homologue of primate cortical area V5/MT (Villeneuse et al., 2006), and to a lesser extent in area 7, which is also motion-sensitive (Pigarev & Rodionova, 1998). Using a population receptive field analysis, Shao et al. (2013) found that fMRI activations in area V5/MT were more extensive in a macaque with juvenile MD compared to control monkeys with a 10° artificial scotoma. Additionally, a recent human study revealed enhanced functional connectivity between MT and early visual areas responsive to peripheral vision in individuals with central vision loss (Fleming et al., 2024). Based on these findings, we hypothesized that the cortical network processing motion might reorganize in MD patients in the presence of the scotoma through enhanced BOLD signals in motion-selective areas and eventually through an expansion of these areas. However, our observations indicated that the cortical networks activated by motion were similar between

MD patients and controls, although with only 2 patients suffering from late-onset Stargardt

disease, we cannot make a definitive statement here. In both populations, our protocol

elicited significant and robust responses in areas hMT+, V3A and V6, as well as within the

portion of early visual areas connected to peripheral vision. An ROI-based analysis

confirmed the implication of these regions and the associated Talairach coordinates are in

good agreement with those reported in the literature. Importantly, we did not observe

significant motion-selective activations outside of this network, which thus rules out the

possibility that new motion areas emerged in patients after the disease apparition.

Furthermore, we found no significant differences in BOLD response amplitudes and extents

between patients and controls in hMT+, V3A, and V6. In V3A and V6, the null hypothesis

was supported by Bayes factors (<1), although a larger sample is needed for firmer

conclusions. In hMT+, a slight trend toward weaker and less extensive activations in patients

compared to controls was noted (Bayes factors = 1.6 and 1.5), but this trend was not

significant and actually goes against our hypotheses about reorganization. Overall, our

results do not indicate any significant large-scale enhancement of the motion-selective

network in MD patients. It is possible that the effects observed by Burnat et al. (2017) in cats

(and also in macaques by Shao et al., 2013) are not transferable to humans. Indeed, it has

been suggested that retinal damage in MD may trigger cortical degeneration in high-level

areas (Hernowo et al., 2014). This raises the possibility that some of the motion-selective

regions studied here could be affected by atrophic processes secondary to sensory

deprivation, limiting their capacity to reorganize. Alternatively, maybe reorganizations exist in

humans but task-based fMRI is not an appropriate neuroimaging technique to capture them

since it only provides an indirect measure of cortical activity (see Calford et al., 2005).

From their fMRI recordings performed in macaques, Shao et al. (2013) also found

that population receptive field (pRF) size in area V5/MT was significantly smaller in an

animal with juvenile MD. To better understand whether this effect is also present in humans,

it will be important to reproduce retinotopic mapping experiments in MD patients and to

perform a pRF analysis. Although several research groups demonstrated that such an

approach is feasible in patients suffering from different kinds of visual pathology (Carvalho et

al., 2022; Papanikolaou et al., 2014; Silson et al., 2023), it remains a challenge in

maculopathy patients because their eye fixation is usually not stable enough to obtain

reliable signals.

While numerous neuroimaging studies on cortical reorganization in MD patients

focused on primary visual areas (see e.g., Morland, 2015 for a complete review), much less

investigated how activations in higher-level cortical areas are modified in maculopathy. Little

et al. (2008) found that age-related MD patients had stronger BOLD responses than control

participants in the prefrontal cortex and in the intraparietal sulcus during visually guided

saccade and smooth-pursuit tasks, two cognitive functions that rely on a form of motion

processing. Important modifications in brain activity were also observed in MD patients

during word recognition (Szlyk & Little, 2009) and scene processing (Ramanoël et al., 2018),

maybe because the visual field in control participants was not masked by an artificial

scotoma. It was suggested that enhanced fMRI activations measured in higher-level cortical

areas in these cases reflected a form of compensation for the decreased sensory functions.

If we did not observe significant modifications of brain activity in our case, it might be

because motion perception in MD patients is not significantly impaired in our task and no

compensation is thus needed.

We estimated motion direction discrimination thresholds in all our participants

because we wanted to replicate one previous result from Guénot et al. (2022) and also to

determine whether fMRI activations in some cortical areas were correlated with motion

perception. We found that discrimination thresholds for lateral motion do not significantly

differ between patients and controls under similar visual conditions. This confirms and further

supports the results of Guénot et al. (2022) which were obtained using the same protocol but

with other groups of patients and controls. While TMS studies suggested that hMT+ is

causally involved in motion perception (Chakraborty et al., 2021; McKeefry et al., 2008;

Strong et al., 2017) and fMRI activations in this area under different motion conditions were

shown to correlate with behavioral reports (Tootell et al., 1995), we did not observe

significant correlations between BOLD activations in our different motion-selective areas

(notably hMT+, V3A and V6) and perceptual thresholds. It should be noted that in our case,

correlations were estimated across MD patients (or across controls), while Tootell et al.

(1995b) examined intra-individual correlations across experimental conditions. The important

inter-individual variability observed in motion-discrimination thresholds and in brain

activations in our data as well as our limited number of participants might explain why our

results differ. Another explanation could also lie in the difference between the protocols used

during our psychophysical and fMRI experiments. Indeed, in order to maximize BOLD

responses, our stimuli alternated between dots moving coherently during 1.8s (with an

unlimited lifetime) and static dots under the MRI scanner while our behavioural protocol

involved dots moving for 0.2s with a limited lifetime. It was reported that small changes in

paradigm based on random dots can influence the results (Pilly & Seitz, 2009).

A limitation of our study is the limited number of participants (7 patients and their

controls). Given this small sample size and our methodology specifically designed to be

suitable in this case, it should be viewed as a case study with matched controls. It is very

challenging to recruit MD patients for neuroimaging experiments and this number can be

explained by the inclusion criteria that we applied. We only recruited patients with no long-

term illnesses, with a stable PRL on at least one eye, with typical cognitive functions and

psychological states (as evaluated by the MMSE and BDI-II), and with no contraindication for

fMRI recordings. The present study is nonetheless in line with previous neuroimaging works

in maculopathy patients (e.g., 4 patients were scanned in Ramanoël et al., 2018, 6 in Little et

al., 2008, 8 in Szlyk & Little, 2009 and 6 in Masuda et al., 2021). Because of our small

number of participants, our analyses lack the statistical power to definitively rule out all forms

of reorganization in MD patients (notably, our Bayes factors only indicate a weak evidence

against enhanced responses in patients). The reliable activation maps that we obtained in all

our patients and their controls (with no significant differences between the response

strengths and extent observed within motion selective areas in the two populations)

nonetheless suggest that the motion processing network is preserved and does not undergo

large-scale enhancements in MD patients following the onset of the scotoma.

We did not perform whole brain analyses because there was an important variability

between the scotomas and PRL positions of our MD patients and hence between their visual

inputs and between the associated cortical activations. It makes it difficult to obtain reliable

results once fMRI responses across patients (or across controls) are projected onto a global

brain template. This problem is further amplified by the important variability that exists

between the position and extent of functional visual areas across individuals (see e.g.,

Kolster et al., 2010 for the hMT+ cluster). We rather chose to analyze the data at the

individual level and to characterize the motion-selective network in each patient and their

age-matched controls. We performed group-level pairwise statistics using independently

defined regions-of-interest (ROIs), which permits us to get rid of the multiple comparisons

problem (see e.g., Poldrack, 2007).

References

Aedo-Jury, F., Cottereau, B. R., Celebrini, S., & Séverac Cauquil, A. (2020). Antero-

Posterior vs. Lateral Vestibular Input Processing in Human Visual Cortex. *Frontiers in*

Integrative Neuroscience, *14*, 43. https://doi.org/10.3389/fnint.2020.00043

Audurier, P., Héjja-Brichard, Y., De Castro, V., Kohler, P. J., Norcia, A. M., Durand, J.-B., &

Cottereau, B. R. (2022). Symmetry Processing in the Macaque Visual Cortex.

Cerebral Cortex, *32*(10), 2277⎕2290. https://doi.org/10.1093/cercor/bhab358

Bach, M. (2007). The Freiburg Visual Acuity Test-variability unchanged by post-hoc re-

analysis. *Graefe's Archive for Clinical and Experimental Ophthalmology = Albrecht*

Von Graefes Archiv Fur Klinische Und Experimentelle Ophthalmologie, *245*(7),

965□971. https://doi.org/10.1007/s00417-006-0474-4

Baker, C. I. (2005). Reorganization of Visual Processing in Macular Degeneration. *Journal of*

Neuroscience, *25*(3), 614□618. https://doi.org/10.1523/JNEUROSCI.3476-04.2005

Barton, B., & Brewer, A. A. (2017). Visual Field Map Clusters in High-Order Visual

Processing□: Organization of V3A/V3B and a New Cloverleaf Cluster in the Posterior

Superior Temporal Sulcus. *Frontiers in Integrative Neuroscience*, *11*.

https://doi.org/10.3389/fnint.2017.00004

Benson, N. C., & Winawer, J. (2018). Bayesian analysis of retinotopic maps. *eLife*, *7*,

e40224. https://doi.org/10.7554/eLife.40224

Bremmer, F., Schlack, A., Shah, N. J., Zafiris, O., Kubischik, M., Hoffmann, K.-P., Zilles, K.,

& Fink, G. R. (2001). Polymodal Motion Processing in Posterior Parietal and

Premotor Cortex. *Neuron*, *29*(1), 287□296. https://doi.org/10.1016/S0896-

6273(01)00198-2

Burnat, K., Hu, T.-T., Kossut, M., Eysel, U. T., & Arckens, L. (2017). Plasticity Beyond V1□:

Reinforcement of Motion Perception upon Binocular Central Retinal Lesions in

Adulthood. *Journal of Neuroscience*, *37*(37), 8989□8999.

https://doi.org/10.1523/JNEUROSCI.1231-17.2017

Calford, M. B., Chino, Y. M., Das, A., Eysel, U. T., Gilbert, C. D., Heinen, S. J., Kaas, J. H., &

Ullman, S. (2005). Rewiring the adult brain. *Nature*, 438(7065), E3□E3.

https://doi.org/10.1038/nature04359

Carvalho, J., Invernizzi, A., Martins, J., Renken, R. J., & Cornelissen, F. W. (2022). Local

neuroplasticity in adult glaucomatous visual cortex. *Scientific Reports*, *12*(1), 21981.

https://doi.org/10.1038/s41598-022-24709-1

Chakraborty, A., Tran, T. T., Silva, A. E., Giaschi, D., & Thompson, B. (2021). Continuous

theta burst TMS of area MT+ impairs attentive motion tracking. *The European*

Journal of Neuroscience, 54(9), 7289‑7300. https://doi.org/10.1111/ejn.15480

Christen, M., & Abegg, M. (2017). The effect of magnification and contrast on reading

performance in different types of simulated low vision. *Journal of Eye Movement*

Research, 10(2). https://doi.org/10.16910/jemr.10.2.5

Contemori, G., Maniglia, M., Guénot, J., Soler, V., Cherubini, M., Cottereau, B. R., & Trotter,

Y. (2024). tRNS boosts visual perceptual learning in participants with bilateral

macular degeneration. *Frontiers in Aging Neuroscience, 16*, 1326435.

https://doi.org/10.3389/fnagi.2024.1326435

Dilks, D. D., Baker, C. I., Peli, E., & Kanwisher, N. (2009). Reorganization of visual

processing in macular degeneration is not specific to the « preferred retinal locus ».

The Journal of Neuroscience: The Official Journal of the Society for Neuroscience,

29(9), 2768‑2773. https://doi.org/10.1523/JNEUROSCI.5258-08.2009

Dumoulin, S. O. (2000). A New Anatomical Landmark for Reliable Identification of Human

Area V5/MT : A Quantitative Analysis of Sulcal Patterning. *Cerebral Cortex, 10*(5),

454‑463. https://doi.org/10.1093/cercor/10.5.454

Dumoulin, S. O., Baker, C. L., Jr, Hess, R. F., & Evans, A. C. (2003). Cortical Specialization

for Processing First- and Second-order Motion. *Cerebral Cortex, 13*(12), 1375‑1385.

https://doi.org/10.1093/cercor/bhg085

Esteban, O., Birman, D., Schaer, M., Koyejo, O. O., Poldrack, R. A., & Gorgolewski, K. J.

(2017). MRIQC : Advancing the automatic prediction of image quality in MRI from

unseen sites. *PLOS ONE, 12*(9), e0184661.

https://doi.org/10.1371/journal.pone.0184661

Fleming, L. L., Defenderfer, M. K., Demirayak, P., Stewart, P., Decarlo, D. K., & Visscher, K.

M. (2024). Impact of Deprivation and Preferential Usage on Functional Connectivity

Between Early Visual Cortex and Category Selective Visual Regions. *Human Brain*

Mapping, 45(17), e70064. https://doi.org/10.1002/hbm.70064

Fletcher, D. C., & Schuchard, R. A. (1997). Preferred Retinal Loci Relationship to Macular

Scotomas in a Low-vision Population. *Ophthalmology, 104*(4), 632‑638.

https://doi.org/10.1016/S0161-6420(97)30260-7

Goebel, R., Esposito, F., & Formisano, E. (2006). Analysis of functional image analysis

contest (FIAC) data with brainvoyager QX□: From single□subject to cortically aligned

group general linear model analysis and self□organizing group independent

component analysis. *Human Brain Mapping*, *27*(5), 392□401.

https://doi.org/10.1002/hbm.20249

Graziano, M., Andersen, R., & Snowden, R. (1994). Tuning of MST neurons to spiral

motions. *The Journal of Neuroscience*, *14*(1), 54□67.

https://doi.org/10.1523/JNEUROSCI.14-01-00054.1994

Grossman, E. D., & Blake, R. (2002). Brain Areas Active during Visual Perception of

Biological Motion. *Neuron*, *35*(6), 1167□1175. https://doi.org/10.1016/S0896-

6273(02)00897-8

Guénot, J., Trotter, Y., Delaval, A., Baurès, R., Soler, V., & Cottereau, B. R. (2023).

Processing of translational, radial and rotational optic flow in older adults. *Scientific*

Reports, *13*(1), 15312. https://doi.org/10.1038/s41598-023-42479-2

Guénot, J., Trotter, Y., Fricker, P., Cherubini, M., Soler, V., & Cottereau, B. R. (2022). Optic

Flow Processing in Patients With Macular Degeneration. *Investigative*

Ophthalmology & Visual Science, *63*(12), 21. https://doi.org/10.1167/iovs.63.12.21

Gupta, A., Mesik, J., Engel, S. A., Smith, R., Schatza, M., Calabrèse, A., van Kuijk, F. J.,

Erdman, A. G., & Legge, G. E. (2018). Beneficial Effects of Spatial Remapping for

Reading With Simulated Central Field Loss. *Investigative Ophthalmology & Visual*

Science, *59*(2), 1105□1112. https://doi.org/10.1167/iovs.16-21404

Hassan, S. E., Lovie-Kitchin, J. E., & Woods, A. R. L. (2002). Vision and Mobility

Performance of Subjects with Age-Related Macular Degeneration: *Optometry and*

Vision Science, *79*(11), 697□707. https://doi.org/10.1097/00006324-200211000-

00007

Hernowo, A. T., Prins, D., Baseler, H. A., Plank, T., Gouws, A. D., Hooymans, J. M. M.,

Morland, A. B., Greenlee, M. W., & Cornelissen, F. W. (2014). Morphometric

analyses of the visual pathways in macular degeneration. *Cortex*, *56*, 99□110.

https://doi.org/10.1016/j.cortex.2013.01.003

Huk, A. C., Dougherty, R. F., & Heeger, D. J. (2002). Retinotopy and Functional Subdivision

of Human Areas MT and MST. *The Journal of Neuroscience*, *22*(16), 7195□7205.

https://doi.org/10.1523/JNEUROSCI.22-16-07195.2002

Kolster, H., Peeters, R., & Orban, G. A. (2010). The Retinotopic Organization of the Human

Middle Temporal Area MT/V5 and Its Cortical Neighbors. *Journal of Neuroscience*,

30(29), 9801□9820. https://doi.org/10.1523/JNEUROSCI.2069-10.2010

Kuyk, T., & Elliott, J. L. (1999). Visual factors and mobility in persons with age-related

macular degeneration. *Journal of Rehabilitation Research and Development*, *36*(4),

303□312.

Li, J. Q., Welchowski, T., Schmid, M., Mauschitz, M. M., Holz, F. G., & Finger, R. P. (2020).

Prevalence and incidence of age-related macular degeneration in Europe□: A

systematic review and meta-analysis. *The British Journal of Ophthalmology*, *104*(8),

1077□1084. https://doi.org/10.1136/bjophthalmol-2019-314422

Little, D., Thulborn, K., & Szlyk, J. (2008). An FMRI study of saccadic and smooth-pursuit

eye movement control in patients with age-related macular degeneration.

Investigative Ophthalmology & Visual Science, *49*(4). https://doi.org/10.1167/iovs.07-

0372

Maniglia, M., Soler, V., Cottereau, B., & Trotter, Y. (2018). Spontaneous and training-

induced cortical plasticity in MD patients□: Hints from lateral masking. *Scientific*

Reports, *8*(1), 90. https://doi.org/10.1038/s41598-017-18261-6

Marron, J. A., & Bailey, I. L. (1982). Visual Factors and Orientation-Mobility Performance:

Optometry and Vision Science, *59*(5), 413□426. https://doi.org/10.1097/00006324-

198205000-00009

Masuda, Y., Dumoulin, S. O., Nakadomari, S., & Wandell, B. A. (2008). V1 projection zone

signals in human macular degeneration depend on task, not stimulus. *Cerebral*

Cortex (New York, N.Y.: 1991), *18*(11), 2483□2493.

https://doi.org/10.1093/cercor/bhm256

Masuda, Y., Takemura, H., Terao, M., Miyazaki, A., Ogawa, S., Horiguchi, H., Nakadomari,

S., Matsumoto, K., Nakano, T., Wandell, B. A., & Amano, K. (2021). V1 Projection

Zone Signals in Human Macular Degeneration Depend on Task Despite Absence of

Visual Stimulus. *Current Biology*, *31*(2), 406-412.e3.

https://doi.org/10.1016/j.cub.2020.10.034

McKeefry, D. J., Burton, M. P., Vakrou, C., Barrett, B. T., & Morland, A. B. (2008). Induced

deficits in speed perception by transcranial magnetic stimulation of human cortical

areas V5/MT+ and V3A. *The Journal of Neuroscience: The Official Journal of the*

Society for Neuroscience, *28*(27), 6848□6857.

https://doi.org/10.1523/JNEUROSCI.1287-08.2008

Morland, A. B. (2015). Organization of the Central Visual Pathways Following Field Defects

Arising from Congenital, Inherited, and Acquired Eye Disease. *Annual Review of*

Vision Science, *1*(1), 329□350. https://doi.org/10.1146/annurev-vision-082114-

035600

Musel, B., Hera, R., Chokron, S., Alleysson, D., Chiquet, C., Romanet, J.-P., Guyader, N., &

Peyrin, C. (2011). Residual abilities in age-related macular degeneration to process

spatial frequencies during natural scene categorization. *Visual Neuroscience*, *28*(6),

529□541. https://doi.org/10.1017/S0952523811000435

Papanikolaou, A., Keliris, G. A., Papageorgiou, T. D., Shao, Y., Krapp, E., Papageorgiou, E.,

Stingl, K., Bruckmann, A., Schiefer, U., Logothetis, N. K., & Smirnakis, S. M. (2014).

Population receptive field analysis of the primary visual cortex complements

perimetry in patients with homonymous visual field defects. *Proceedings of the*

National Academy of Sciences, *111*(16). https://doi.org/10.1073/pnas.1317074111

Peyrin, C., Ramanoël, S., Roux-Sibilon, A., Chokron, S., & Hera, R. (2017). Scene

perception in age-related macular degeneration□: Effect of spatial frequencies and

contrast in residual vision. *Vision Research*, *130*, 36□47.

https://doi.org/10.1016/j.visres.2016.11.004

Pigarev, I. N., & Rodionova, E. I. (1998). Two visual areas located in the middle suprasylvian

gyrus (cytoarchitectonic field 7) of the cat's cortex. *Neuroscience*, *85*(3), 717□732.

https://doi.org/10.1016/S0306-4522(97)00642-8

Pilly, P. K., & Seitz, A. R. (2009). What a difference a parameter makes□: A psychophysical

comparison of random dot motion algorithms. *Vision Research*, 49(13), 1599□1612.

https://doi.org/10.1016/j.visres.2009.03.019

Pitzalis, S., Galletti, C., Huang, R.-S., Patria, F., Committeri, G., Galati, G., Fattori, P., &

Sereno, M. I. (2006). Wide-Field Retinotopy Defines Human Cortical Visual Area V6.

The Journal of Neuroscience, *26*(30), 7962□7973.

https://doi.org/10.1523/JNEUROSCI.0178-06.2006

Pitzalis, S., Sereno, M. I., Committeri, G., Fattori, P., Galati, G., Patria, F., & Galletti, C.

(2010). Human v6□: The medial motion area. *Cerebral Cortex (New York, N.Y.:*

1991), *20*(2), 411□424. https://doi.org/10.1093/cercor/bhp112

Poldrack, R. A. (2007). Region of interest analysis for fMRI. *Social Cognitive and Affective*

Neuroscience, *2*(1), 67□70. https://doi.org/10.1093/scan/nsm006

Priebe, N. J., Lisberger, S. G., & Movshon, J. A. (2006). Tuning for Spatiotemporal

Frequency and Speed in Directionally Selective Neurons of Macaque Striate Cortex.

The Journal of Neuroscience, *26*(11), 2941□2950.

https://doi.org/10.1523/JNEUROSCI.3936-05.2006

Raftery, A. E. (1995). Bayesian Model Selection in Social Research. *Sociological*

Methodology, *25*, 111. https://doi.org/10.2307/271063

Ramanoël, S., Chokron, S., Hera, R., Kauffmann, L., Chiquet, C., Krainik, A., & Peyrin, C.

(2018). Age-related macular degeneration changes the processing of visual scenes

in the brain. *Visual Neuroscience*, *35*, E006.

https://doi.org/10.1017/S0952523817000372

Shao, Y., Keliris, G. A., Papanikolaou, A., Fischer, M. D., Zobor, D., Jägle, H., Logothetis, N.

K., & Smirnakis, S. M. (2013). Visual cortex organisation in a macaque monkey with

macular degeneration. *European Journal of Neuroscience*, *38*(10), 3456□3464.

https://doi.org/10.1111/ejn.12349

Silson, E. H., Baker, C. I., Aleman, T. S., Maguire, A. M., Bennett, J., & Ashtari, M. (2023).

Motion-selective areas V5/MT and MST appear resistant to deterioration in

choroideremia. *NeuroImage: Clinical*, *38*, 103384.

https://doi.org/10.1016/j.nicl.2023.103384

Strong, S. L., Silson, E. H., Gouws, A. D., Morland, A. B., & McKeefry, D. J. (2017). A Direct

Demonstration of Functional Differences between Subdivisions of Human V5/MT+.

Cerebral Cortex (New York, NY), *27*(1), 1□10. https://doi.org/10.1093/cercor/bhw362

Szlyk, J. P., & Little, D. M. (2009). An FMRI study of word-level recognition and processing

in patients with age-related macular degeneration. *Investigative Ophthalmology &*

Visual Science, *50*(9), 4487□4495. https://doi.org/10.1167/iovs.08-2258

Talairach, J., & Tournoux, P. (1988). *Co-planar Stereotaxic Atlas of the Human Brain□: 3-*

dimensional Proportional System□: an Approach to Cerebral Imaging. G. Thieme.

Tarita-Nistor, L., González, E. G., Markowitz, S. N., Lillakas, L., & Steinbach, M. J. (2008).

Increased Role of Peripheral Vision in Self-Induced Motion in Patients with Age-

Related Macular Degeneration. *Investigative Ophthalmology & Visual Science*, *49*(7),

3253□3258. https://doi.org/10.1167/iovs.07-1290

Tootell, R. B. H., Mendola, J. D., Hadjikhani, N. K., Ledden, P. J., Liu, Arthur. K., Reppas, J.

B., Sereno, M. I., & Dale, A. M. (1997). Functional Analysis of V3A and Related

Areas in Human Visual Cortex. *The Journal of Neuroscience*, *17*(18), 7060□7078.

https://doi.org/10.1523/JNEUROSCI.17-18-07060.1997

Tootell, R. B. H., Reppas, J. B., Dale, A. M., Look, R. B., Sereno, M. I., Malach, R., Brady, T.

J., & Rosen, B. R. (1995). Visual motion aftereffect in human cortical area MT

revealed by functional magnetic resonance imaging. *Nature*, *375*(6527), 139□141.

https://doi.org/10.1038/375139a0

Tootell, R., Reppas, J., Kwong, K., Malach, R., Born, R., Brady, T., Rosen, B., & Belliveau, J.

(1995). Functional analysis of human MT and related visual cortical areas using

magnetic resonance imaging. *The Journal of Neuroscience*, *15*(4), 3215□3230.

989   https://doi.org/10.1523/JNEUROSCI.15-04-03215.1995

990 Tran, T. H. C., Guyader, N., Guerin, A., Despretz, P., & Boucart, M. (2011). Figure Ground

991   Discrimination in Age-Related Macular Degeneration. *Investigative Opthalmology &*

992   *Visual Science*, *52*(3), 1655. https://doi.org/10.1167/iovs.10-6003

993 Villeneuve, M. Y., Ptito, M., & Casanova, C. (2006). Global motion integration in the postero-

994   medial part of the lateral suprasylvian cortex in the cat. *Experimental Brain Research*,

995   *172*(4), 485□497. https://doi.org/10.1007/s00221-006-0357-2

996 Watson, A. B., & Pelli, D. G. (1983). Quest□: A Bayesian adaptive psychometric method.

997   *Perception & Psychophysics*, *33*(2), 113□120. https://doi.org/10.3758/BF03202828

998 Westeneng-van Haaften, S. C., Boon, C. J. F., Cremers, F. P. M., Hoefsloot, L. H., Den

999   Hollander, A. I., & Hoyng, C. B. (2012). Clinical and Genetic Characteristics of Late-

1000   onset Stargardt's Disease. *Ophthalmology*, *119*(6), 1199□1210.

1001   https://doi.org/10.1016/j.ophtha.2012.01.005

1002 Wood, J. M., Lacherez, P. F., Black, A. A., Cole, M. H., Boon, M. Y., & Kerr, G. K. (2009).

1003   Postural Stability and Gait among Older Adults with Age-Related Maculopathy.

1004   *Investigative Opthalmology & Visual Science*, *50*(1), 482.

1005   https://doi.org/10.1167/iovs.08-1942

1006 Zeki, S., Watson, J. D., Lueck, C. J., Friston, K. J., Kennard, C., & Frackowiak, R. S. (1991).

1007   A direct demonstration of functional specialization in human visual cortex. *The*

1008   *Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *11*(3),

1009   641□649. https://doi.org/10.1523/JNEUROSCI.11-03-00641.1991

1010

1011

1012

1013

1014

Supplementary Materials

Supplementary Figure 1-1: Images of the eye fundus and estimation of the scotoma size (left panel)

and PRL (middle panel) in all patients (except MD3, whose data are provided in figures 1 and 3). The

1019 right panels show the artificial scotomas used in the corresponding control participants. In each case,

1020 we provide the visible portion of the stimulus (i.e., the surface that is not covered by the artificial

1021 scotoma). It is to be noted that the fovea and the PRL are located at the same place for STG1. See

1022 Figure 1 for more details.

1023 %%%

1024

1025

1026 ***Supplementary Figure 4-1:*** Comparison between BOLD responses to visual stimulation and to a

1027 blank screen on the lateral cortical surfaces of all MD patients (left panels) and their respective

1028 controls (right panels). See Figure 3 for the details of the legend.

1029 %%

1030

1031

1032 **Supplementary Figure 4-2:** Comparison between BOLD responses to visual stimulation and to a

1033 blank screen on the medial cortical surfaces of all MD patients (left panels) and their respective

1034 controls (right panels). See Figure 3 for the details of the legend.

1035   %%

		pSTS
MD1	**LH**	X
	RH	X
MD2	**LH**	X
	RH	51; -48; 10*
MD3	**LH**	-47; -46; 2
	RH	53; -51; 5
MD4	**LH**	-47; -47; 9
	RH	56; -40; 5
MD5	**LH**	X
	RH	X
STG1	**LH**	-39; -49; 11*
	RH	51; -43; 12*
STG2	**LH**	-49; -50; 5
	RH	54; -41; 6
CTRL1	**LH**	X
	RH	X
CTRL2	**LH**	-51; -40; 3
	RH	52; -33; 5
CTRL3	**LH**	-46; -38; -1
	RH	43; -42; -1
CTRL4	**LH**	X
	RH	55; -35; 10
CTRL5	**LH**	X
	RH	44; -35; 1
CTRLSTG1	**LH**	-45; -49; 10
	RH	53; -37; 11
CTRLSTG2	**LH**	-47; -48; 5
	RH	40; -54; 12
Average patients	**LH**	-46±4; -48±8; 7±4
	RH	53±3; -45±3; 8±2
Average controls	**LH**	-47±2; -44±1; 4±2
	RH	48±1; -39±1; 6±1
Literature	**LH**	-56; -47; 8
	RH	52; -49; 15

1036   ***Supplementary Table 2-1:*** Talairach coordinates of pSTS observed in patients and controls, the

1037   calculated mean for both patients and controls, and the coordinates reported in Barton and Brewer

1038   (2017). The '*' symbol indicates the coordinates for which a more permissive threshold has been

1039   used, i.e p<10-2.

1040

[legend] MD1/CTRL1 MD2/CTRL2 MD3/CTRL3 MD4/CTRL4 MD5/CTRL5 STG1/CTRLSTG1 STG2/CTRLSTG2

Supplementary Figure 6-1: Activation extents with the hMT+ (A), V3A (B) and V6 (C) ROIs. These

extents were computed as the percentage of voxels passing a threshold of t-score = 3 within each

ROI. For each group, the boxplots provide the first quartile, median, last quartile, and the extreme

percentages of activated voxels. None of the comparison led to significant results for either hMT+ (p-

value > 0.05, Bayes factor = 1.5), V3A (p-value > 0.05, Bayes factor = 0.4) or V6 (p-value > 0.05,

Bayes factor = 0.4). ROIs were defined as spheres of 10 mm of radius centered on the Talairach

coordinates provided in the literature (see the '*Methods*' section) but similar results were obtained

when considering spheres of 5 and 15 mm of radius.

1050

1051

[legend] MD1/CTRL1 MD2/CTRL2 MD3/CTRL3 MD4/CTRL4 MD5/CTRL5 STG1/CTRLSTG1 STG2/CTRLSTG2

Supplementary Figure 6-2: Motion-selective activations in the pSTS ROI. For each patient and their

age-matched control (connected by lines), colored dots provide the differences between the average

(across voxels and hemispheres) percentages of signal changes (PSCs) during the translational

motion and control (static dots) conditions. For each group, the boxplots provide the first quartile,

median, last quartile, and the extreme PSC difference. These differences are not significantly different

from zero (p-values = 0.109 in both MD patients and controls, Wilcoxon test). Statistical analyses

based on paired Wilcoxon sign-rank tests did not reveal significant differences between activations in

the two groups (p-value > 0.05, Bayes factor = 0.47).

1061

Supplementary Figure 6-3: Motion-selective activations in the hMT+ (A), V3A (B) and V6 (C) ROIs

when removing MD5, STG1, or both. For each patient and their age-matched control (connected by

lines), colored dots provide the differences between the average (across voxels and hemispheres)

percentages of signal changes (PSCs) during the translational motion and control (static dots)

conditions. For each group, the boxplots provide the first quartile, median, last quartile, and the

extreme PSC difference. Stars (* or **) indicate that the PSC difference is significantly different from

zero (p-values < 0.05 or <0 .01, respectively). Statistical analyses based on paired Wilcoxon sign-rank

tests did not reveal significant differences between activations in the two groups (p-values > 0.05 for

the three ROIs).

1071

1072

1073

Supplementary Figure 7-1: Differences between the average percentages of signal changes (PSCs)

during the translational motion and control (static dots) conditions in the hMT+ (A), V3A (B) and V6

(C) ROIs as a function of the scotoma size, given by the radius of the associated disk (see the

'*Methods*' section). The corresponding Spearman correlation coefficients rho (ρ) are shown. None of

these correlations were significant (p-values > 0.05). For patient MD5 (whose scotoma is elongated,

see supplementary Figure 1) and his control, we used the semi minor axis of the associated ellipse for

the scotoma size in this Figure. We obtained similar results when using the semi major axis of the

ellipse.

1082

1083