Motion processing in visual cortex of maculopathy patients

Célia Michaud[1] [#], Jade Guénot[1] [#], Cynthia Faurite[2], Mathilde Gallice[3], Christophe Chiquet[3], Nathalie Vayssière[1], Isabelle Berry[1], Yves Trotter[1], Vincent Soler[1], Carole Peyrin[2] *, Benoit R. Cottereau[1, 4] *

[1] CerCo UMR 5549, CNRS – Université Toulouse III, 31059 Toulouse, France.
[2] Univ. Grenoble Alpes, Univ. Savoie Mont Blanc, CNRS, LPNC, 38400 Saint-Martin d'Hères, France
[3] Department of Ophthalmology, Grenoble Alpes University Hospital, 38700 La Tronche, France.
[4] IPAL, CNRS IRL 2955, 117418 Singapore.
[#] These authors contributed equally to this work.
* Co-senior authors.

Corresponding authors: Célia Michaud ('celia.michaud@cnrs.fr') and Benoit Cottereau ('benoit.cottereau@cnrs.fr')

Abbreviated title: Cortical motion processing in maculopathy patients

Number of pages: 47
Number of figures: 7
Number of tables: 2
Number of words
     Abstract: 241
     Introduction: 649
     Discussion: 1452

Abstract: Previous studies on animal models suggested that visual areas involved in motion processing could undergo important cortical reorganizations following retinal damages. This could have major implications for patients suffering from macular degeneration (MD), one leading cause of vision loss. Here, we performed fMRI recordings in a group of maculopathy patients (including individuals suffering from age-related macular degeneration or from Stargardt's Disease) and a control group to characterize the motion processing cortical network in MD patients and determine whether this network undergoes significant large-scale reorganizations following the onset of the scotoma. We used an experimental protocol based on random-dot kinematograms (RDKs) classically employed to characterize motion-selective

areas in the brain. To ensure that the visual information processed by the two groups was

equivalent, the visual field in each control participant was masked using an artificial scotoma

directly derived from clinical measurements in their paired patient. We found that in MD

patients, translational motion elicited significant and robust activations in a restricted cortical

network which included the human V5/MT+ complex (hMT+), areas V3A and V6, and a portion

of primary visual areas (V1, V2 and V3) connected to peripheral vision. Importantly, the same

patterns of responses were also observed in control participants. Moreover, the extent and

strength of activation within these motion-selective areas did not differ significantly between

the two groups. Altogether, these results suggest that in humans, the motion-selective network

does not undergo significant large-scale cortical reorganizations following the onset of MD.

Keywords: Macular degeneration, Motion processing, fMRI, Plasticity, Cortical

reorganizations

Significance statement: Motion processing in the visual cortex of patients with macular

degeneration has never been characterized. Here, we performed fMRI recordings in 7

maculopathy patients and found robust motion-selective activations in a cortical network which

included the human V5/MT+ complex (hMT+), areas V3A and V6, and a portion of primary

visual areas connected to peripheral vision. These activations are in very good agreement

with those reported in participants with normal vision in the literature and do not significantly

differ from those measured in a group of age and gender-matched control participants.

Altogether, our results suggest that the motion-selective network does not undergo significant

large-scale reorganizations in maculopathy patients following the onset of the scotoma.

1. Introduction

Macular degeneration (MD) is a visual pathology that alters the outer layers of the central retina, resulting in central vision loss. Its most common form is age-related macular degeneration (AMD) which develops after 60 years and currently affects over 67 million of patients in Europe (Li et al., 2020). Macular degeneration can also affect younger populations under the form of Stargardt's disease (STG) which usually onsets during adolescence but can also appear later in life (Late-Onset Stargardt Disease; Westeneng-van Haaften et al., 2012). Despite their different origins, both diseases result in a loss of visual acuity and decreased contrast sensitivity, with severe consequences on patient's quality of life as they impair recognition and mobility (Hassan et al., 2002; Kuyk & Elliott, 1999; Marron & Bailey, 1982; Wood et al., 2009). MD patients usually rely on peripheral vision, which has a much lower spatial resolution compared to central vision, to process their visual environment (Musel et al., 2011; Tran et al., 2011). Most of them develop a preferred retinal locus (PRL) outside the scotoma for ocular fixation (Fletcher & Schuchard, 1997). This overuse of peripheral vision was hypothesized to lead to cortical reorganizations in visual cortex (Baker, 2005), notably in neural populations which previously processed central vision. However, the associated neural mechanisms remain poorly understood. Most studies on cortical reorganization in MD patients focused on the primary visual cortex (see however Little et al., 2008 or Ramanoël et al., 2018). Recordings using functional magnetic resonance imaging (fMRI) demonstrated that stimuli presented in peripheral vision activated portions of area V1 connected to the scotoma (Dilks et al., 2009), but it is now accepted that this effect is dependent on the cognitive tasks and low-level visual features, and only reflect unmasking of cortical feedbacks (Masuda et al., 2008, 2021; Morland, 2015). Interestingly, recent studies on animal models suggested that cortical reorganization after MD can be more pronounced in higher-level visual cortex, notably along the dorsal pathway. Using fMRI, Shao et al., (2013) found that visual responses in area MT, a dorsal region processing motion, were significantly modified in a macaque monkey with

juvenile MD. In a cat model, Burnat et al., (2017) showed that the posteromedial lateral

suprasylvian area PMLS (the homologue of primate cortical area V5/MT; Villeneuve et al.,

2006) and to a lesser degree area 7 (which is also motion sensitive; Pigarev & Rodionova,

1998) had enhanced activity following the induction of a central retinal lesion in adult animals.

These enhancements were associated with better performances in a motion task. In humans,

a recent study showed that functional connectivity between the portions of early visual areas

(V1, V2, and V3) that respond to peripheral vision and MT was enhanced in individuals with

central vision loss (Fleming et al., 2024). Another study in psychophysics reported that self-

induced motion perception from peripheral stimuli was increased in MD patients (Tarita-Nistor

et al., 2008), suggesting some degree of plasticity in cortical areas processing motion. It is

thus possible that the motion-selective cortical network is modified in MD patients following

the onset of the scotoma but this hypothesis has never been tested using functional

neuroimaging.

Here, we employed an fMRI protocol commonly used to identify cortical areas selective to

motion (see e.g., R. Tootell et al., 1995 or Zeki et al., 1991) in 7 MD patients and 7 age and

gender-matched controls, whose central visual field was masked using simulated scotomas

of the same positions and dimensions as those of their corresponding patients. Our main

objectives were to characterize the motion processing cortical network in MD patients and to

determine whether this network undergoes significant large-scale reorganisations following

the onset of the scotoma. We tested if the cortical areas within this network are similar or

different between the two populations and whether their associated BOLD responses are

comparable. To complete our analysis, we performed psychophysical measurements to

assess the motion discrimination thresholds in both groups.

2. Methods

2.1 Participants: Seven maculopathy patients (age range: 50 - 81) were recruited from the

Retina Center of the Purpan Hospital of Toulouse and from the Ophthalmology center of the

Grenoble Hospital (CHUGA) in France (see Table 1). Five of them were diagnosed with age-

related macular degeneration (AMD), while the two others suffered from late-onset Stargardt'

disease. They all had dense central bilateral scotomas and a stable preferred retinal locus on

at least one eye. None of them suffered from other visual illnesses. The experiments were

also conducted in 7 age- and gender-matched controls with normal (or corrected to normal)

vision. Age differences between patients and controls were all lower than two years except for

the control of MD2 (81) who was 76 (despite a broad screening procedure, we did not manage

to find an older volunteer). Importantly, each control participant underwent the experiments in

the same center as their paired patient. None of the participants suffered from general long-

term illnesses. In order to evaluate their cognitive functions and their psychological states,

they all underwent the Mini-Mental State Examination (MMSE) and the Beck Depression

Inventory (BDI-II), respectively. All of them had a score superior to 23 for the MMSE and

inferior to 19 for the BDI-II. All participants gave their written informed consent and the protocol

was approved by the French Ethics Committee (CPP) of Sud Mediterranée IV (2022-00957-

36).

Subjects	Sex	Age	Diagnosis	Visual acuity		Tested eye	Scotoma radius deg	PRL polar angle (θ), deg	PRL eccentricity (ρ), deg
				LE	RE				
MD1	F	75	AMD	0.14	0.15	L	9.72	-27.4	10.4
MD2	M	81	AMD	0.03	0.48	R	5.3	143.9	1.1
MD3	F	66	AMD	0.6	0.23	L	7.31	157.3	2.1
MD4	M	73	AMD	0.23	0.22	L	9.84	28.5	9.58
MD5	M	76	AMD	0.05	0.23	R	6.15 12.31*	-151,5	0.9
STG1	F	50	STG	0.4	0.7	L	5.18	0	0
STG2	M	69	STG	0.25	0.25	R	5.87	-90	5.22
CTRL1	F	74	-	1.54	1.80	R	-	-	-
CTRL2	M	76	-	1.03	1.13	R	-	-	-
CTRL3	F	67	-	1.11	0.61	L	-	-	-
CTRL4	M	71	-	1.27	1.73	L	-	-	-
CTRL5	M	77	-	0.96	1.71	R	-	-	-
CTRLSTG1	F	52	-	1.31	0.99	L	-	-	-
CTRLSTG2	M	68	-	1.43	2	R	-	-	-

Table 1: Characteristics of MD patients and age-matched controls included in the study. Precisions about the diagnosis of patients (AMD for Age-related Macular Degeneration, STG for late-onset Stargardt's disease), the decimal visual acuity for the left (L) and right (R) eye, as well as the polar coordinates (angles and eccentricities) of the Preferred Retinal Locus (PRL) are included. Patients and their controls carried out the experiments in the same centers, i.e in Toulouse or in Grenoble. Controls for MD patients 1 to 5 are labelled CTRL1 to CTRL 5. Controls for STG patients STG1 and STG2 are labelled CTRLSTG1 and CTRLSTG2. It should be noted that because MD5 had an elongated scotoma, we used an ellipse rather than a disk for the artificial scotoma of his control. In this case, the two values provided in the 'scotoma radius' column (marked by an asterisk) provide the lengths of the ellipse semi-minor and semi-major axes.

For each participant, the eye with the better acuity was determined using the Sloan letters of the Freiburg Visual Acuity Test (Bach, 2007). In most participants, this eye was stimulated during the experiments and the other eye was patched. Only patients MD1 and ST1 were stimulated in the other eye because it was the only one with a stable preferred retinal locus (PRL, see below). In their case, the eye with the better acuity was patched.

2.2 Scotoma size: For each patient, an ophthalmologist delineated the border of the scotoma from the fundus of the eye captured by standard and autofluorescence imaging (see Figure 1-A for MD3, other examples are provided in Supplementary figure 1). The area of the scotoma (in square degrees of visual angles) was extracted using MATLAB (MathWorks, Natick, MA, USA). This area was used to compute the simulated scotoma of the age- and gender-matched controls (see below).

2.3 Position of the Preferred Retinal Locus (PRL) in maculopathy patients: MD patients usually develop a new locus of ocular fixation following the onset of their scotoma. To test for the presence of such preferred retinal locus (PRL) and to determine its location, we

performed spectral-domain optical coherence tomography (see Figure 1-B). During the

measure, patients were instructed to fixate a light produced by the OCT. Patients with a

developed PRL naturally used it to gaze, thereby allowing to record its retinal coordinates.

This process was repeated three times with a short break in between acquisitions, to ensure

that the measure was stable (see Contemori et al., 2024). Table 1 reports the average PRL

locations across these three measures (see Maniglia et al., 2018 or Guénot et al., 2022 for

more details). It should be noted that two patients (MD5 and STG1) had their PRLs within the

scotoma, even though these scotomas were clearly apparent on autofluorescence imaging.

Figure 1: Estimation of the scotoma size and PRL position in one typical patient (MD3). These values

are used to define the artificial scotoma masking the central visual field in the associated control

participant. It should be noted that all retinal images in this figure have been inverted horizontally to be

aligned with the visual field. The letters on panel A and B specify the orientation of the images (L: Left;

R: Right; D: Down; U: Up). A) An autofluorescence imaging of the eye fundus is acquired. The border

of the scotoma is delineated and the associated surface is used to calculate the radius of the scotoma

in degrees of visual angle, given that 288μm on the image corresponds to 1 degree of visual angle. B)

Patients were instructed to fixate a central light and we measured the x and y position of their PRL with

spectral-domain optical coherence tomography (Spectralis Optical Coherence Tomography [OCT];

Heidelberg Engineering, Heidelberg, Germany). This process was repeated three times (leftward panel)

and the final coordinates (green dot on the rightward panel) were the average across these three

measures. Here, the position of the fovea is marked by a red dot. From the surface of the scotoma and

the position of the PRL, it is possible to simulate the patient's visual field. C) During the fMRI

experiments, MD patients were instructed to fixate a central cross (shown here in green) with their PRL.

Controls had to fixate a red cross at the position corresponding to their paired patient's fovea while an

artificial scotoma was displayed in dark gray. The dotted lines (not shown during the experiments)

specify the four quadrants of the patients' and controls' visual fields in this case: upper-left ('UL'), upper-

right ('UR'), lower-left ('LL') and ('LR'). The screen (in light gray) dimensions provided here (20° x 15°)

are those used in the two MRI centers.

%%

2.4 Artificial scotoma and viewing conditions in control participants: For each

control participant, we masked the central visual field using an artificial scotoma as in previous

studies (Christen & Abegg, 2017; Gupta et al., 2018). This simulated scotoma had the same

surface as the one measured in the associated patient (see the '*Scotoma size*' section) and

was centered on the fovea (except for CTRL2 and CTRL 3 where the artificial was shifted to

the left because MD2's and MD3's scotoma occupied a larger portion of the left visual

hemifield). Control participants were instructed to gaze on a fixation cross that we displayed

at the position of their paired patient's fovea (see the red cross on Figure 1-C). In this case,

the center of the screen fell on their actual peripheral vision, but at the PRL location of their

paired patient. For all control participants, artificial scotomas were simplified as a disk except

for CTRL5 for which we used an ellipse to more closely match the real shape of MD5 scotoma,

which was very elongated (see Supplementary figure 1). This protocol permits patients and

controls to receive comparable visual inputs. It notably equalizes the position and surface of

the occluded part of the visual field (see Guénot et al., 2022).

2.5 Motion stimuli: Motion stimuli were adapted from the experimental protocol of a

previous electrophysiological study (Graziano et al., 1994). They consisted in translational

patterns defined from randomdot kinematograms (RDKs) (see Figure 2-A) generated with

Matlab (R2017a). The RDKs contained bright non-overlapping dots (diameter: 0.2°) moving

on a homogenous dark background with a high contrast (100%). They had a density of 0.3945

dots per degree of visual angle. When a dot reached the border of the display screen, it was

immediately relocated at a random position. This process permits to equalize mean luminance

and dot density across the screen during the experiments. Each dot moved at a velocity of

7°/s, which corresponds to the average preferred speed of neurons in macaque MT (Priebe et

al., 2006). This speed is in agreement with those used in previous fMRI studies on motion

processing (6°/s in Zeki et al., 1991, 8°/s in Huk & Heeger, 2002). These translational motion

stimuli were used for the fMRI recordings (see the next section) and for the psychophysical

measurements (see the '*Experimental protocol for the psychophysical measurements*'

section).

2.6 Experimental protocol for fMRI measurements: In both Toulouse and Grenoble

MRI platforms, stimuli were displayed on a large screen and covered 20°x15° of visual angle.

fMRI recordings were performed using a block-design. Each run consisted of 312 s and were

divided into 6 identical cycles of 52 s (see Figure 2-A). Each cycle started with a baseline

period of 10 s during which only the fixation point and the artificial scotoma (for controls) were

displayed on a black screen. It was followed by a block of 16s of uniform motion, then by

another 10 s of baseline and finally by a block of 16 s with static dots. During motion blocks,

a new pattern of translational motion during which all the dots moved either upward or

downward was presented every 2 s. These patterns last 1.8 s and were separated by short

periods (0.2 s) of ocular fixation. The same principle was applied during the blocks of statics

dots (a new pattern was displayed for 1,.8 s every 2 s and patterns were separated by 0.2 s

of fixation).

Figure 2: Experimental protocols. A) fMRI recordings were performed using a block-design. Runs lasted 312 s and contained 6 identical cycles of 52 s. Each cycle started with a baseline period of 10 s during which only the fixation cross and the artificial scotoma (for controls) were displayed on a black screen. It was followed by a block of 16 s of uniform motion (shown here in blue), then by another 10 s of baseline and finally by a block of 16 s with static dots (in red). During motion blocks, a new pattern of translational motion during which all the dots moved either upward or downward was presented every 2 s. These patterns last 1.8 s and were separated by short periods (0.2 s) of ocular fixation. The same principle was applied during the blocks of statics dots (a new pattern was displayed for 1.8 s every 2 s and patterns were separated by 0.2 s of fixation). B) During psychophysical recordings, participants had to report the perceived motion direction of the stimuli (either upward or downward). We manipulated motion coherence and estimated thresholds corresponding to 80 percent of correct detections. For illustration, an upward motion trial is shown at 20% of coherence. Signal dots are shown in white and noise dots in gray for better visibility on the figure (all the dots were white during the experiment). During all the recordings (fMRI and psychophysics), participants had to fixate a red cross that was central for patients (who used their PRL) and at the position corresponding to their paired patient's fovea for controls.

%%%

This protocol is classically employed to identify motion selective cortical areas (Zeki et al., 1991; Watson et al., 1993; Tootell et al., 1995a; Huk & Heeger, 2002). The visual display of the stimuli and its temporal synchronization with the acquisition of the fMRI data was controlled using the EventIDE software (Okazolab). A red fixation cross was displayed during all the

functional recordings, in the center of the screen for patients (who gazed with their PRLs) and

at the position corresponding to their paired patient's fovea for controls (see above). In order

to make sure that all participants correctly gazed and were focused on the visual stimuli, a

white circle was randomly displayed for 200 ms on top of this cross during the runs (10 displays

per run on average). Participants were instructed to press a button on a gamepad whenever

they detected it. Even though this task was not easy (notably for patients who were fixating

with their PRLs), they all performed it well (mean percentage of correct detection for patients:

80; and controls: 90) and we thus kept all the runs for further analyses.

2.7 (f)MRI recordings: (f)MRI recordings were performed at 3T (Achieva dStream 3.0T TX,

Philips, NL) at Toulouse NeuroImaging Center (ToNIC, Toulouse, France) and at IRMaGe

MRI facility (Grenoble, France). Importantly, both MRI platforms are equipped with the same

MR hardware configuration (with Quasar Dual gradient system and dStream architecture

supporting MultiBand-accelerated acquisitions, 32 channel head coil). They were operated

using the same software version (R5.7.1). This permits to minimize the differences between

the recordings performed in the two cities, although we emphasize that each patient from one

center was matched with a control from the same center. We acquired two high-resolution

anatomical images of the participant's brain using a T1-weighted magnetization-prepared

rapid gradient-echo (MPRAGE) sequence (repetition time, TR = 8.1 ms; echo time, TE = 3.7

ms, flip angle = 8°; FOV: 240 x 240 mm; voxel size = 1 x 1 x 1 mm; 170 sagittal slices). These

anatomical images were first co-registered and then averaged together to be used as a

reference to which the functional images from all the experiments were aligned. Functional

images were collected with a T2 * weighted gradient-echo echoplanar imaging (EPI) sequence

(TR = 2000 ms, TE = 30 ms, flip angle = 75°, SENSE factor = 1.6; multi-band (MB) = 3; FOV:

224 x 224 mm; matrix size: 112 x 110 mm; voxel size = 2 x 2 x 2 mm, 63 transverse slices,

dyn = 156).

2.8 (f)MRI data analyses: MRIQC v.13.1.0 was used to check the quality of all our functional recordings (Esteban et al., 2017). We did not observe any out-of-range runs in our data, indicating little to no head-movement. We thus kept all the measurements for further analyses. Pre-processing of fMRI data included slice scan time correction, 3D motion correction using trilinear/sinc interpolation, and a linear trend followed by a high-pass filtering using Fourier analysis with a cut-off at 3 cycles. For each participant, functional data were co-registered on the anatomy. Functional and anatomical data were brought into anterior commissure - posterior commissure (ACPC) space using cubic spline interpolation and then transformed into standard Talairach (TAL) space (Talairach & Tournoux, 1988). For further analyses, the volume time courses were spatially smoothed at 4 mm FWHM.

Individual analyses were performed on each node of the participant' cortical tessellations by fitting a general linear model (GLM) to the corresponding blood oxygen level-dependent (BOLD) signal. The model contained 2 main regressors, representing the 2 experimental conditions: uniform motion (either upward or downward) and control (static dots). These regressors were convolved with a model of the human hemodynamic response function. We also introduced 8 regressors of non-interest in our model. 6 of them corresponded to the translation and rotation of the head estimated during the motion-correction step of the pre-processing (see above). The two last ones were time-courses captured within the ventricles and the white-matter. This operation permits to reduce the noise in the data (see e.g., Audurier et al., 2022). For each participant, the beta weights obtained from the GLM were subsequently used to perform univariate analyses (t-scores) at the whole-brain level. All these analyses were realized using the BrainVoyager software (v23, Brain Innovation, Maastricht, the Netherlands) (Goebel et al., 2006). Results are shown on inflated surfaces, obtained by segmenting the white and gray matter, as well as the cerebrospinal fluid, and reconstructing a surface at the boundary of the white and gray matter. Statistical maps are mainly thresholded

at p-value $< 10^{-3}$ (uncorrected), as is often the case in individual-level fMRI analysis (see e.g.,

Aedo-Jury et al., 2020).

Group-level analyses were performed within independently defined regions of interest (ROIs).

These ROIs were given by spheres in the volumetric space centered on the Talairach

coordinates provided in previous studies for areas hMT+ (Watson et al., 1993), V3A (R. B. H.

Tootell et al., 1997) and V6 (Pitzalis et al., 2010). Within each ROI, we computed the

differences between the averages percentages of signal changes (PSCs) during the

translational motion condition and during its control condition (static dots). For each group, we

first determined whether these differences of PSCs were significantly different from zero

across participants. We used Wilcoxon sign-rank tests because it is non parametric and thus

does not assume that the data are normally distributed. Then, we tested whether the

differences of PSCS were different between the two groups (MD patients and controls), also

using Wilcoxon sign-rank tests. Here, data were paired between each patient and his/her

control to reflect the fact that each pair received the same visual stimulations. We used

spheres with a radius of 10 mm, a value which leads to a cortical surface in line with the one

reported for area hMT+ (see Kolster et al., 2010). It should be noted that the analyses were

also repeated with radius values of 5 and 15 mm to make sure that the choice of this parameter

did not influence our results.

2.9 Experimental protocol for psychophysical measurements: The psychophysical

measurements were performed separately from the fMRI recordings, and outside the MRI

scanner. During the experiment, participants sat in a chair whose height was adapted in order

to equalize the height of the eyes with the center of the screen. Their head was placed on a

head-support device clamped on top of a table and equipped with both chin and forehead

supports. The chair and head-support devices were positioned so as to ensure a fine

alignment between the participants' head and trunk axes. Participants had to keep this position

as constant as possible. Dot stimuli of 0.2° in size were presented on computer screens using

a large portion of the visual field (58.1° × 43.8° of visual angle in Toulouse and 48.6° x 30.7°

in Grenoble) to maximize coverage of peripheral vision. As for the fMRI recordings, during the

whole experiment, participants had to fixate monocularly (the other eye was patched) a red

cross. This cross was central for patients (who used their PRL) and at the position

corresponding to their patient fovea for age-matched controls.

We used the same random dot kinematogram stimuli as for the fMRI experiments (see the

'*Motion stimuli*' section), except that here, each dot had a limited lifetime of 200 ms (12 frames

at 60 Hz). At the end of its lifetime, it was randomly reassigned to a new spatial position. To

avoid a coherent flickering of the stimulus every 200 ms, each dot initial age was randomly

picked between 0 and 166 ms (11 frames) at the beginning of each trial. We chose to

manipulate lifetime to permit a direct comparison of the psychophysical results with those

obtained in a previous study using the same paradigm, but performed on other MD patients

(Guénot et al., 2022). It should be noted that we used unlimited lifetimes for the fMRI

recordings in order to maximize the BOLD responses of motion selective areas, and hence to

facilitate their localization.

A two-alternative forced-choice task (2-AFC) was used to estimate motion discrimination

thresholds. Stimuli were presented in blocks of 64 trials. Each block lasted about 3 min. Each

trial started with the presentation of the stimulus for 200 ms. Participants had to report whether

the motion of the stimulus was upward or downward. They were instructed to respond as

quickly as possible while maximizing their performances. Responses reported after 2 s were

considered as incorrect. After each trial, an auditory feedback indicated whether the chosen

direction was correct or not. During each block, we manipulated motion coherency (i.e., the

percentage of dots moving along the same direction while the other dots had random

directions) and estimated the thresholds corresponding to 80% of correct detection using an

adaptive Bayesian approach (QUEST, see below). Participants completed 4 blocks. The first

block was considered as a training and not included in the analyses. Breaks were included (3 minutes on average) between blocks to reduce fatigue. The whole experiment was completed in one session of about 20 minutes.

2.10 Robust estimation of motion coherence thresholds: Because it is complicated for MD patients to perform long psychophysical experiments, we used the QUEST adaptive procedure to obtain rapid, efficient and robust estimations of motion coherence thresholds (Guénot et al., 2022, 2023). QUEST is based on a Bayesian approach assuming that the psychophysics function underlying the participant performance follows a Weibull distribution. During a block, the estimated parameters of this function were updated after each trial on the basis of the participant's response. Coherence values corresponded to the current maximum likelihood estimate of the threshold. We fixed the maximum number of trials at 64 as it was previously shown that this value leads to robust thresholds in most circumstances (Watson & Pelli, 1983). We used an initial threshold value of 58 % based on previous recordings using the same stimuli in MD patients (Guénot et al., 2022).

3. Results

The aim of this study was to characterize the cortical network processing translational motion in maculopathy patients and to determine whether it undergoes cortical reorganizations. We recorded fMRI data in 7 patients (5 with AMD and 2 with Stargardt's disease) and in 7 age- and gender-matched controls using an experimental paradigm commonly employed to identify motion selective areas (Zeki et al., 1991; Watson et al., 1993; Tootell et al., 1995a; Huk & Heeger, 2002). For each participant, we also estimated the motion-direction discrimination thresholds using a two-alternative forced choice (2-AFC) task. In each control, central vision was masked using an artificial scotoma so that visual inputs matched those of his/her paired patient. In the following, we first detail the fMRI activation patterns observed in our MD patients

398   and their controls. Next, we describe the results of the psychophysical experiment and their

399   relationship with BOLD responses.

3.1 Motion-selective responses in the cortex of MD patients and their controls

Figure 3 shows the motion-selective activations (t-scores, p-value < 0.001, uncorrected)

obtained in one typical MD patient (MD3, upper panel) and her control (CTRL3, lower panel)

displayed on enlarged lateral views of their inflated individual cortical surfaces. Motion-

selective activations in the other participants are shown in Figure 4. Activations displayed on

medial views for all participants are shown on Figure 5. In all these figures, cortical nodes with

significantly stronger responses to translational motion than to static dots appear in hot colors

(yellow/red) while nodes with significantly stronger responses to the opposite contrast are

shown in cold (blue) colors. Data were thresholded at p-value < 10^{-3} (uncorrected).

Figure 3: Comparison between BOLD responses to the translational motion condition and to its control

condition (static dots) in one representative patient (MD3) and her age- and gender-matched control

(CTRL3). Activations are shown on inflated individual cortical surfaces. Gyri appear in pale gray while

sulci are shown in darker gray. Cortical nodes with significantly stronger responses to translational

motion appear in yellow/red colors while nodes with significantly stronger responses to static dots are

shown in blue. Data were thresholded at p-value < 10^{-3} (uncorrected). It should be noted that area V3A

in the right hemisphere of MD3 was detected using a threshold p-value of 10^{-2} and it is thus not visible

here. Colored lines provide the anatomical positions of the Inferior Occipital Sulcus (IOS, in olive green),

of the Lateral Occipital Sulcus (LOS, in green), of the Transverse Occipital Sulcus (TOS, in purple), of

the ventral part of the Anterior Occipital Sulcus (AOSv, in cyan), and of the Superior Temporal Sulcus

(STS, in red). The central panel illustrates visual stimulation in this case. During the fMRI recordings,

MD3 fixed the center of the screen (green cross) with her PRL while CTRL3 gazed at the position of

MD3's fovea (red cross). The dark disk is the artificial scotoma (see Figure 1 for more details on its

definition). The dotted lines (not shown during the recordings) specify the positions of the horizontal and

vertical meridians. We also provide the coherence thresholds (CoTh) estimated through psychophysical

measurements.

%%

In all participants, translational motion elicited significantly stronger responses in a cortical

network that is mainly localized within the dorsal visual pathway (i.e., in occipito-parietal areas)

and within early visual areas. It should be noted that this network is very restricted when

compared to the one obtained by contrasting responses to visual stimulation versus baseline

(ocular fixation) and which covers the majority of the posterior part of the brain and notably

includes areas along the ventral pathway (lateral and medial views of the corresponding

activations maps are provided in Supplementary Figures 2 and 3). In the following, we first

focus on responses within cortical regions that were consistently reported as motion-selective

in the literature. Then, we describe our results in early visual areas and in other parts of the

brain.

3.2 Activations in area hMT+: In all MD patients and their controls, we observed significantly stronger responses to translational motion than to static dots within (or posterior to) the ventral part of the anterior occipital sulcus (AOSv, in cyan in Figure 3), in between the inferior (IOS, in olive green) and the lateral (LOS, in green) occipital sulci (see Figures 3 and 4). The Talairach coordinates of the local maxima (t-scores) measured in this region are provided in Table 2. Averages values in the left and right hemispheres for patients ([-42±2; -71±2; 5±2] and [41±2; -66±2; 2±2]) and controls ([-42±2; -64±2; 4±3] and [42±2; -63±2; 3±2]) were similar. These anatomical locations and coordinates are in very good agreement with those reported in the literature for area hMT+ (e.g., [-41; -69; 2] and [41; -67; 2] in Watson et al., 1993 or [-47; -76; 2] and [44; -67; 0] in Dumoulin, 2000). To confirm that our data reflected motion-selective activations in this area, we used independently defined regions of interest (ROIs) centered on the Talairach coordinates provided by Watson et al., (1993) for hMT+ and tested whether the differences between the percentages of signal change (PSCs) measured during the translational motion condition and during its control (static dots) were significantly different from zeros across participants in this ROI (see the '*Methods*' section). This was the case for both groups (Wilcoxon sign-rank tests, p-value = 0.0156 in MD patients and p-value = 0.0078 in controls, see Figure 6-A). Importantly, we did not find significant differences between the differences in PSCs observed in this ROI for the two groups (Wilcoxon paired sign-rank tests, p-value = 0.1094). As an additional analysis, we also considered the activation extent within this ROI and found that they did not significantly differ between the two groups (see Supplementary figure 4). Altogether, these results suggest that both MD patients and their controls have motion-selective responses in area hMT+ and that these responses do not significantly differ between the two populations.

3.3 Activations in area V3A: In all MD patients and their controls, we observed significantly stronger responses to translational motion within the transverse occipital sulcus (in purple on Figure 3, see also Figure 4). For some participants, we had to decrease our threshold to a p-value of 10^{-2}, uncorrected) at similar Talairach coordinates between the two groups ([-18±2 -

81±2 16±3] and [21±3 -80±2 22±4] for patients and [-22±2 -80±1 25±3] and [23±1 -76±1 20±2]

for controls on average (see table 2). These anatomical locations and coordinates are in very

good agreement with those reported in the literature for area V3A (e.g., [±14; -84; 19] in Tootell

et al., 1997 or [±18; -78; 25] in Pitzalis et al., 2006). For both groups, we confirmed that our

data reflected motion-selective activations in this area from statistical analysis within an ROI

centered on the Talairach coordinates provided by Tootell et al. (1997) (Wilcoxon sign-rank

tests, p-value = 0.0078 in MD patients and p-value = 0.0234 in controls, see Figure 6-B and

above). We did not find significant differences between the differences of PSCs observed in

this ROI for the two groups (Wilcoxon paired sign-rank tests, p-value = 0.469), nor between

their activation extents (see Supplementary figure 4). Altogether, these results suggest that

both MD patients and their controls have motion-selective responses in area V3A and that

these responses do not significantly differ between the two populations.

3.4 Activations in area V6: In six MD patients and their controls (all participants except

MD3 and CTRL3), we observed significantly stronger responses to translational motion in the

upper part of the cuneus, within the posterior branch of the parieto-occipital sulcus (POS) at

similar Talairach coordinates between the two groups ([-12±2 -76±2 29±3] and [12±3 -73±3

31±3] for patients and [-14±2 -74±2 32±3] and [11±1 -75±1 35±3] for controls on average, see

Table 2). These anatomical locations and coordinates are in very good agreement with those

reported in the literature for area V6 (e.g., [±9; -78; 34] in Pitzalis et al., 2010). For both groups,

we confirmed that our data reflected selectivity to translational motion in this area from

statistical analysis within an ROI centered on the Talairach coordinates provided by Pitzalis et

al. (2010) (Wilcoxon sign-rank tests, p-value = 0.0156 in MD patients and p-value = 0.00781

in controls, see Figure 6-B and above). We did not find significant differences between the

differences of PSCs observed in this ROI for the two groups (Wilcoxon paired sign-rank tests,

p-value = 0.297), nor between their activation extents (see Supplementary figure 4).

Altogether, these results suggest that both MD patients and their controls have motion-

selective responses in area V6 and that these responses do not significantly differ between

the two populations.

3.5 Activations in early visual areas and relationship to retinotopic properties:

In all participants, we also observed motions selective activations below and above the most

anterior part of the calcarine sulcus (see Figure 5). These regions correspond to the portions

of primary visual areas (V1, V2 and V3) connected to peripheral vision. Their stronger

responses to translational motion are not surprising given that a vast proportion of their

neurons are motion-direction selective (Dumoulin et al., 2003). Here as well, we did not

observe different patterns of activations in statistical parametric maps between groups (Figure

5). It should be noted that for patients whose scotoma covers a larger surface in one visual

hemifield (because their PRL is lateralized) and for their controls (MD1/CTRL1, MD2/CTRL2,

MD3/CTRL3 and MD4/CTRL4), motion-selective activations were generally stronger in the

hemisphere ipsilateral to the scotoma (i.e., contralateral to the visual hemifield that receives

more visual inputs) which reflects the retinotopic properties of early visual areas (see e.g.,

Benson & Winawer, 2018). In V3A, hemispheric asymmetries were also observed for these

participants but they were less pronounced, reflecting the fact that neural receptive fields in

this area are bigger than in areas V1, V2 and V3 (e.g., Barton & Brewer, 2017 reported

population receptive fields with radiuses between 4 and 6° at eccentricities comparable to

those in our study). We did not observe hemispheric asymmetries in areas hMT+ and V6,

probably because these areas have very large receptive fields which overlap with the

ipsilateral visual hemifield (for hMT+, see Huk et al., 2002; for V6, see Pitzalis et al., 2010).

For hMT+, this is notably true in its hMST functional subdivision.

3.6 Additional activations: Beyond the activations reported above, we also observed

motion-selective responses in other parts of the brain but they were not as reliable and robust

across participants. In some patients and their controls (MD2/CTRL2, MD3/CTRL3,

MD4/CTRL4, STG1/STGCTRL1 and STG2/STGCTRL2, see also on the right hemisphere of

CTRL5), we notably observed motion-selective activations in the fundus of the superior

temporal sulcus (STS, in red on Figure 3) and also (but less frequently) on its lower bank (MD3

in the right hemisphere, MD4 in the right hemisphere, CTRL5 in the left hemisphere) or within

the middle temporal sulcus (MTS, CTRL2). These activations could correspond to the human

posterior temporal sulcus (pSTS) area, which is selective to biological motion (Grossman &

Blake, 2002) and which preferentially encodes peripheral vision (Barton & Brewer, 2017). The

associated Talairach coordinates are in good agreement with those reported for this area in

the literature (see supplementary Table 1) but an ROI-based analysis using a sphere centered

on the coordinates provided by Barton and Brewer (2017) did not confirm its implication

(Wilcoxon sign-rank tests, p-value = 0.109 in both MD patients and controls, see

Supplementary Figure 5).

In some participants (MD3, MD4, CTRL1 and CTRL2), additional motion-selective activations

were also observed in the parietal cortex and notably within the intra-parietal sulcus, where is

located the ventral intraparietal (VIP) area, which is motion selective (Bremmer et al., 2001).

Figure 4: Comparison between BOLD responses to the translational motion condition and to its control

condition (static dots) in all participants, except MD3 and her control (CTRL3), whose data are provided

on enlarged views in Figure 3. Significantly stronger responses to motion are shown on lateral views of

the cortical surfaces for MD patients (leftward panels) and their controls (rightward panels). See Figure

3 for the details of the legend.

%%%

		hMT+	V3A	V6
MD1	LH	-47; -71; 4	-17; -86; 13	-9; -74; 33
	RH	49; -66; -4	22; -84; 12	7; -77; 34
MD2	LH	-43; -66; 7	-22; -77; 18	-6; -80; 24
	RH	32; -66; 12	13; -78; 22	4; -78; 21
MD3	LH	-47; -66; 5	-16; -82; 14	X
	RH	38; -64; -2	29; -73; 32	X
MD4	LH	-43; -73; -2	-27; -72; 9	-17; -72; 23
	RH	49; -64; -5	29; -78; 27	20; -69; 24
MD5	LH	-39; -73; 13	-11; -77; 29	-13; -70; 32
	RH	40; -66; 7	18; -81; 33	19; -69; 34
STG1	LH	-33; -78; 0	-11; -87; 7	-12; -81; 38
	RH	37; -70; -1	11; -84; 8	6; -74; 34
STG2	LH	-42; -73; 8	-22; -83; 21	-12; -77; 21
	RH	43; -67; 5	24; -80; 19	18; -72; 38
CTRL1	LH	-44; -64; 15	-20; -79; 31	-16; -69; 34
	RH	49; -65; 11	24; -75; 21	13; -67; 45
CTRL2	LH	-42; -58; 7	-22; -73; 32	-19; -68; 46
	RH	38; -54; 6	21; -66; 28	10; -70; 45
CTRL3	LH	-36; -60; 3	-20; -77; 31	-15; -71; 31
	RH	38; -63; -3	26; -68; 24	14; -69; 27
CTRL4	LH	-40; -70; -4	-21; -79; 17	-13; -72; 31
	RH	48; -68; 3	19; -82; 23	7; -75; 29
CTRL5	LH	-40; -69; -7	-21; -79; 14	-12; -84; 15
	RH	38; -70; 0	29; -77; 19	9; -81; 27
STGCTRL1	LH	-47; -63; 13	-20; -89; 26	-6; -75; 34
	RH	45; -59; 2	23; -82; 15	12; -80; 33
STGCTRL2	LH	-47; -69; -2	-29; -87; 22	-18; -78; 30
	RH	35; -63; -1	18; -81; 10	10; -81; 36
Average patients	LH	-42±2; -71±2; 5±2	-18±2; -81±2; 16±3	-12±2; -76±2; 29±3
	RH	41±2; -66±2; 3±2	21±3; -80±3; 22±4	12±3; -73±3; 31±3
Average controls	LH	-42±2; -64±2; 4±3	-22±1; -80±2; 25±3	-14±2; -74±2; 32±3
	RH	42±2; -63±2; 3±2	23±1; ±-76±1; 20±2	11±1; -75±1; 35±3
Literature	LH	-41; -69; 2	-14; -84; 19	-9; -78; 34
	RH	41; -67; 2	14; -84; 19	9; -78; 34

Table 2: Talairach coordinates of the different motion-selective areas observed in our study (hMT+,

V3A and V6). Coordinates are provided for each MD patient and his/her control. The table also provides

average coordinates across each group as well as average coordinates reported in previous studies for

these 3 areas (Watson et al., 1993 for hMT+; Tootell et al., 1997 for V3A and Pitzalis et al., 2010 for

V6).

%%%

Figure 5: Comparison between BOLD responses to the translational motion condition and to its control condition (static dots) in all participants. Significantly stronger responses to motion are shown on medial views of the cortical surfaces for MD patients (leftward panels) and their controls (rightward panels). See Figure 3 for the details of the legend.

%%%

Figure 6: Motion-selective activations in the hMT+ (A), V3A (B) and V6 (C) ROIs. For each patient and

their age-matched control (connected by lines), colored dots provide the differences between the

average (across voxels and hemispheres) percentages of signal changes (PSCs) during the

translational motion and control (static dots) conditions. For each group, the boxplots provide the first

quartile, median, last quartile, and the extreme PSC difference. Stars (* or **) indicate that the PSC

difference is significantly different from zero (p-values < 0.05 or <0.01, respectively). Statistical analyses

based on paired Wilcoxon sign-rank tests did not reveal significant differences between activations in

the two groups (p-values > 0.05 for the three ROIs).

%%%

3.7 Motion-direction discrimination thresholds and their relationship with BOLD

responses

The motion direction discrimination thresholds estimated in each MD patient and their age-

matched control (see the 'Material and Methods' section) are provided in Figure 7-A.

Thresholds varied between 0.37 and 0.99 in patients (0.37 for MD3 whose fMRI data are also

shown on Figure 3; 0.67 on average) and between 0.21 and 0.98 in control participants (0.21

for CTRL3, see Figure 3; 0.58 on average). We did not find significant differences between

the thresholds measured in the two groups (Wilcoxon signed-rank test, p-value = 0.1698).

Altogether, these results confirm Guénot et al.'s (2022) results where motion direction

discrimination thresholds did not significantly differ between MD patients and their age-

matched controls under identical experimental conditions (see Figure 5 in Guénot et al., 2022).

Figure 7: Behavioral results and relationship with cortical responses in motion-selective ROIs. A)

Motion-direction discrimination thresholds estimated in the two groups. The boxplots show the different

values for the first quartile, median, last quartile, and the extreme data points. The cross represents the

means for both groups. Dots provide the individual data points of the distributions. It should be noted

that they were slightly offset horizontally to improve their visibility. B) Differences between the average

percentages of signal changes (PSCs) during the translational motion and control (static dots)

conditions in the hMT+ ROI as a function of coherence thresholds. Dots and triangles respectively

correspond to MD patients and controls. Solid and dotted lines provide the associated regressions lines.

Pearson correlation coefficients R are also shown (the associated p-values were respectively 0.086 for

patients and 0.4513 for controls). Data in the V3A and V6 ROIs are respectively provided in panels C

and D. The p-values associated with the Pearson correlation coefficients were 0.3043 and 0.0614 in

area V3A and 0.5689 and 0.5250 in area V6 for patients and control participants, respectively. The x-

axis was cropped at 0.2 for better visibility.

%%%

Next, we examine whether the performances measured in psychophysics are correlated with

fMRI activations in motion-selective areas. For each participant, panels B, C and D of Figure

7 show the differences between the averages percentages of signal changes (PSCs) during

the translational motion condition and during its control condition (static dots) in areas hMT+,

V3A and V6 as a function of motion-discrimination thresholds. For each group, we provide the

corresponding Pearson correlation coefficients. This measure does not assume normality of

the data and tests for linear correlation. Although stronger responses to motion stimuli were

generally associated with lower perceptual thresholds (Pearson coefficients were mostly

negative), we did not observe any significant correlations in the data (p-values > 0.5, see

details in the legend of the figure). In order to test whether other forms of correlations (e.g.,

non linear) existed in our data, we also estimated the Spearman rank coefficients (which test

whether the data are monotonically related and do not assume normality). Here as well, we

did not observe any significant correlations.

To complete these analyses, we also tested whether the size of the scotoma could influence

the percentages of signal changes measured in our ROIs but we did not observe significant

effects in both groups (see Supplementary figure 6). This is probably because receptive fields

in these areas are large enough to capture visual inputs outside the scotoma (see also the

'*Activations in early visual areas and relationship to retinotopic properties*' section).

3.8 Control analyses to verify the robustness of our ROI-based approach

All the ROIs-based analyses in our study used spheres of 10 mm. To make sure that this

choice did not impact our results, we reproduced these analyses with spheres of 5 and 15 mm

of radius and found the same patterns of results.

617   4. Discussion

The aim of this study was to characterize the cortical network processing translational

motion in maculopathy patients. We recorded fMRI data in 7 patients and in 7 age- and gender-

matched controls using an experimental paradigm based on RDKs classically employed to

identify motion selective areas (Zeki et al., 1991; Watson et al., 1993; Tootell et al., 1995a;

Huk et al., 2002). We also estimated motion-direction discrimination thresholds in each

participant. Stimuli in each control participant were masked using an artificial scotoma of the

same dimension as the scotoma of their paired patients, in order to equalize the visual inputs

received by the two populations.

A previous study on an animal model (Burnat et al., 2017) found that the induction of

a central retinal lesion in adult cats led to improved neuronal activity in the posteromedial

lateral suprasylvian area PMLS (the homologue of primate cortical area V5/MT, see Villeneuse

et al., 2006) and to a lesser degree in area 7 (which is also motion sensitive, see Pigarev &

Rodionova, 1998). Another study in humans suggested that functional connectivity between

MT and the portions of early visual areas that respond to peripheral vision is enhanced in

individuals with central vision loss (Fleming et al., 2024). We had thus hypothesized that the

cortical network processing motion could reorganize itself in MD patients in the presence of

the scotoma, either through an enhancement of the BOLD signal in motion selective areas or

through a modification of the areas involved and notably of their locations and/or extents. For

example, Shao et al. (2013) reported more extended activations in area V5/MT of a macaque

with juvenile MD. We found that the cortical networks activated by motion were actually similar

between the two populations. In both patients and controls, our protocol elicited significant and

robust responses in areas hMT+, V3A and V6, as well as within the portion of early visual

areas connected to peripheral vision. An ROI-based analysis confirmed the implication of

these regions and the associated Talairach coordinates are in good agreement with those

reported in the literature. Importantly, we did not observe significant motion-selective

activations outside of this network, which thus rules out the possibility that new motion areas

emerge in patients after the disease apparition. Finally, we did not find significant differences

in BOLD responses between patients and controls in hMT+, V3A and V6. Altogether, these

results do not demonstrate any significant large-scale modification of the motion-selective

network in MD patients in the presence of the scotoma. It is possible that the effects observed

in cats by Burnat et al. (2017) are not transferable to other mammal species. Alternatively,

maybe reorganizations exist in humans but fMRI is not an appropriate neuroimaging technique

to capture them since it only provides an indirect measure of cortical activity (see Calford et

al., 2005).

From fMRI recordings performed in macaques, Shao et al. (2013) found that population

receptive field (pRF) size in area V5/MT was significantly smaller in an animal with juvenile

MD than in control monkeys with a 10° artificial scotoma. It is not clear whether such a

shrinking of the pRF size in a cortical area (and notably in area hMT, localized within hMT+)

can lead to a modification of the fMRI responses in the context of our motion protocol. Given

our dot density (0.3945 dots per degree of visual angle) and the reported pRF size diminution

reported in V5/MT (~1.5°, see Figure 7-b in Shao et al., 2013), one might expect a reduction

of the BOLD signal. This is not what we observed in the hMT+ area (differences in PSCs were

on average lower for patients but this tendency was not significant) and also in the other motion

selective areas of our MD patients. To better understand whether the effects observed in

macaques are also present in humans, it will be important to reproduce retinotopic mapping

experiments in MD patients and to perform a pRF analysis. Although several research groups

demonstrated that such an approach is feasible in patients suffering from different kinds of

visual pathology (Carvalho et al., 2022; Papanikolaou et al., 2014; Silson et al., 2023), it

remains a challenge in maculopathy patients because their eye fixation is usually not stable

enough to obtain reliable signals.

While numerous neuroimaging studies on cortical reorganization in MD patients

focused on primary visual areas (see e.g., Morland, 2015 for a complete review), much less

investigated how activations in higher-level cortical areas are modified in maculopathy. Little

et al. (2008) found that age-related MD patients had stronger BOLD responses than control

participants in the prefrontal cortex and in the intraparietal sulcus during visually guided

saccade and smooth-pursuit tasks, two cognitive functions that rely on a form of motion

processing. Important modifications in brain activity were also observed in MD patients during

word recognition (Szlyk & Little, 2009) and scene processing (Ramanoël et al., 2018), maybe

because the visual field in control participants was not masked by an artificial scotoma. It was

suggested that enhanced fMRI activations measured in higher-level cortical areas in these

cases reflected a form of compensation for the decreased sensory functions. If we did not

observe significant modifications of brain activity in our case, it might be because motion

perception in MD patients is not significantly impaired in our task and no compensation is thus

needed.

We estimated motion direction discrimination thresholds in all our participants because

we wanted to replicate one previous result from Guénot et al. (2022) and also to determine

whether fMRI activations in some cortical areas were correlated with motion perception. We

found that discrimination thresholds for lateral motion do not significantly differ between

patients and controls under similar visual conditions. This confirms and further supports the

results of Guénot et al. (2022) which were obtained using the same protocol but with other

groups of patients and controls. While TMS studies suggested that hMT+ is causally involved

in motion perception (Chakraborty et al., 2021; McKeefry et al., 2008; Strong et al., 2017) and

fMRI activations in this area under different motion conditions were shown to correlate with

behavioral reports (Tootell et al., 1995), we did not observe significant correlations between

BOLD activations in our different motion-selective areas (notably hMT+, V3A and V6) and

perceptual thresholds. It should be noted that in our case, correlations were estimated across

MD patients (or across controls), while Tootell et al. (1995b) examined intra-individual

correlations across experimental conditions. The important inter-individual variability observed

in motion-discrimination thresholds and in brain activations in our data as well as our limited

number of participants might explain why our results differ (in addition to the differences of

experimental protocols).

A limitation of our study is the limited number of participants (7 patients and their

controls). It is very challenging to recruit MD patients for neuroimaging experiments and this

number can be explained by the inclusion criteria that we applied. We only recruited patients

with no long-term illnesses, with a stable PRL on at least one eye, with typical cognitive

functions and psychological states (as evaluated by the MMSE and BDI-II), and with no

contraindication for fMRI recordings. The present study is nonetheless in line with previous

neuroimaging works in maculopathy patients (e.g., 4 patients were scanned in Ramanoël et

al., 2018, 6 in Little et al., 2008, 8 in Szlyk & Little, 2009 and 6 in Masuda et al., 2021). Given

this small sample size, our study should rather be considered as a case study with matched

controls and might not have enough statistical power to exclude all forms of reorganizations

in MD patients. The very reliable activation maps that we obtained in all our patients and their

controls (with no significant differences between the response strengths and extent observed

within motion selective areas in the two populations) nonetheless suggest that the motion

processing network is preserved and does not undergo large-scale reorganization in MD

patients following the onset of the scotoma.

We did not perform whole brain analyses because there was an important variability

between the scotomas and PRL positions of our MD patients and hence between their visual

inputs and between the associated cortical activations. It makes it difficult to obtain reliable

results once fMRI responses across patients (or across controls) are projected onto a global

brain template. This problem is further amplified by the important variability that exists between

the position and extent of functional visual areas across individuals (see e.g., Kolster et al.,

2010 for the hMT+ cluster). We rather chose to analyze the data at the individual level and to

characterize the motion-selective network in each patient and their age-matched controls. We

performed group-level pairwise statistics using independently defined regions-of-interest

(ROIs), which permits us to get rid of the multiple comparisons problem (see e.g., Poldrack,

2007).

Data and code availability: MRI and fMRI data are provided at the following link:

These data have undergone spatial normalization, in compliance with the General Data

Protection Regulation (GDPR) under which this project is carried out. Data used for the

analyses reported in Figures 6 and 7 (differences of PSCs and motion direction discrimination

thresholds) are provided here:

https://osf.io/gpxhz/?view_only=43a1f1993d8742e6ba85ed4889b99a75

Fundings: This study was supported by a French grant from the Agence Nationale de la

Recherche (ANR-21-CE28-0021, ANR PRC ReViS-MD; awarded to C.P. and B.R.C.).

Competing interests: The authors have no competing interests or conflicts of interest

to disclose.

Ethics Statement: All participants gave their written informed consent and the protocol

was approved by the French Ethics Committee (CPP) of Sud Mediterranée IV (2022-00957-

36).

Acknowledgments: This work was performed on the Inserm/UPS UMR1214 Technical

Platform of Toulouse and the IRMaGe platform member of France Life Imaging network (grant

ANR-11-INBS-0006) for Grenoble. We thank Judith Eck for helping on fMRI analysis, and Ilia

Korjoukov and Louise Kauffmann for helping develop our MRI paradigm.

References

Aedo-Jury, F., Cottereau, B. R., Celebrini, S., & Séverac Cauquil, A. (2020). Antero-

Posterior vs. Lateral Vestibular Input Processing in Human Visual Cortex. *Frontiers in*

Integrative Neuroscience, *14*, 43. https://doi.org/10.3389/fnint.2020.00043

Audurier, P., Héjja-Brichard, Y., De Castro, V., Kohler, P. J., Norcia, A. M., Durand, J.-B., &

Cottereau, B. R. (2022). Symmetry Processing in the Macaque Visual Cortex.

Cerebral Cortex, *32*(10), 2277-2290. https://doi.org/10.1093/cercor/bhab358

Bach, M. (2007). The Freiburg Visual Acuity Test-variability unchanged by post-hoc re-

analysis. *Graefe's Archive for Clinical and Experimental Ophthalmology = Albrecht*

Von Graefes Archiv Fur Klinische Und Experimentelle Ophthalmologie, *245*(7),

965-971. https://doi.org/10.1007/s00417-006-0474-4

Baker, C. I. (2005). Reorganization of Visual Processing in Macular Degeneration. *Journal of*

Neuroscience, *25*(3), 614-618. https://doi.org/10.1523/JNEUROSCI.3476-04.2005

Barton, B., & Brewer, A. A. (2017). Visual Field Map Clusters in High-Order Visual

Processing : Organization of V3A/V3B and a New Cloverleaf Cluster in the Posterior

Superior Temporal Sulcus. *Frontiers in Integrative Neuroscience*, *11*.

https://doi.org/10.3389/fnint.2017.00004

Benson, N. C., & Winawer, J. (2018). Bayesian analysis of retinotopic maps. *eLife*, *7*,

775   e40224. https://doi.org/10.7554/eLife.40224

776 Bremmer, F., Schlack, A., Shah, N. J., Zafiris, O., Kubischik, M., Hoffmann, K.-P., Zilles, K.,

777   & Fink, G. R. (2001). Polymodal Motion Processing in Posterior Parietal and

778   Premotor Cortex. *Neuron*, *29*(1), 287‑296. https://doi.org/10.1016/S0896-

779   6273(01)00198-2

780 Burnat, K., Hu, T.-T., Kossut, M., Eysel, U. T., & Arckens, L. (2017). Plasticity Beyond V1 :

781   Reinforcement of Motion Perception upon Binocular Central Retinal Lesions in

782   Adulthood. *Journal of Neuroscience*, *37*(37), 8989‑8999.

783   https://doi.org/10.1523/JNEUROSCI.1231-17.2017

784 Calford, M. B., Chino, Y. M., Das, A., Eysel, U. T., Gilbert, C. D., Heinen, S. J., Kaas, J. H., &

785   Ullman, S. (2005). Rewiring the adult brain. *Nature*, *438*(7065), E3‑E3.

786   https://doi.org/10.1038/nature04359

787 Carvalho, J., Invernizzi, A., Martins, J., Renken, R. J., & Cornelissen, F. W. (2022). Local

788   neuroplasticity in adult glaucomatous visual cortex. *Scientific Reports*, *12*(1), 21981.

789   https://doi.org/10.1038/s41598-022-24709-1

790 Chakraborty, A., Tran, T. T., Silva, A. E., Giaschi, D., & Thompson, B. (2021). Continuous

791   theta burst TMS of area MT+ impairs attentive motion tracking. *The European*

792   *Journal of Neuroscience*, *54*(9), 7289‑7300. https://doi.org/10.1111/ejn.15480

793 Christen, M., & Abegg, M. (2017). The effect of magnification and contrast on reading

794   performance in different types of simulated low vision. *Journal of Eye Movement*

795   *Research*, *10*(2). https://doi.org/10.16910/jemr.10.2.5

796 Contemori, G., Maniglia, M., Guénot, J., Soler, V., Cherubini, M., Cottereau, B. R., & Trotter,

797   Y. (2024). tRNS boosts visual perceptual learning in participants with bilateral

798   macular degeneration. *Frontiers in Aging Neuroscience*, *16*, 1326435.

799   https://doi.org/10.3389/fnagi.2024.1326435

800 Dilks, D. D., Baker, C. I., Peli, E., & Kanwisher, N. (2009). Reorganization of visual

801   processing in macular degeneration is not specific to the « preferred retinal locus ».

802   *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience,*

29(9), 2768-2773. https://doi.org/10.1523/JNEUROSCI.5258-08.2009

Dumoulin, S. O. (2000). A New Anatomical Landmark for Reliable Identification of Human

Area V5/MT : A Quantitative Analysis of Sulcal Patterning. *Cerebral Cortex*, *10*(5),

454-463. https://doi.org/10.1093/cercor/10.5.454

Dumoulin, S. O., Baker, C. L., Jr, Hess, R. F., & Evans, A. C. (2003). Cortical Specialization

for Processing First- and Second-order Motion. *Cerebral Cortex, 13*(12), 1375-1385.

https://doi.org/10.1093/cercor/bhg085

Esteban, O., Birman, D., Schaer, M., Koyejo, O. O., Poldrack, R. A., & Gorgolewski, K. J.

(2017). MRIQC : Advancing the automatic prediction of image quality in MRI from

unseen sites. *PLOS ONE*, *12*(9), e0184661.

https://doi.org/10.1371/journal.pone.0184661

Fleming, L. L., Defenderfer, M. K., Demirayak, P., Stewart, P., Decarlo, D. K., & Visscher, K.

M. (2024). Impact of Deprivation and Preferential Usage on Functional Connectivity

Between Early Visual Cortex and Category-Selective Visual Regions. *Human Brain*

Mapping, *45*(17), e70064. https://doi.org/10.1002/hbm.70064

Fletcher, D. C., & Schuchard, R. A. (1997). Preferred Retinal Loci Relationship to Macular

Scotomas in a Low-vision Population. *Ophthalmology*, *104*(4), 632-638.

https://doi.org/10.1016/S0161-6420(97)30260-7

Goebel, R., Esposito, F., & Formisano, E. (2006). Analysis of functional image analysis

contest (FIAC) data with brainvoyager QX : From single-subject to cortically aligned

group general linear model analysis and self-organizing group independent

component analysis. *Human Brain Mapping*, *27*(5), 392-401.

https://doi.org/10.1002/hbm.20249

Graziano, M., Andersen, R., & Snowden, R. (1994). Tuning of MST neurons to spiral

motions. *The Journal of Neuroscience*, *14*(1), 54-67.

https://doi.org/10.1523/JNEUROSCI.14-01-00054.1994

Grossman, E. D., & Blake, R. (2002). Brain Areas Active during Visual Perception of

Biological Motion. *Neuron*, *35*(6), 1167-1175. https://doi.org/10.1016/S0896-

6273(02)00897-8

Guénot, J., Trotter, Y., Delaval, A., Baurès, R., Soler, V., & Cottereau, B. R. (2023).

Processing of translational, radial and rotational optic flow in older adults. *Scientific*

Reports, *13*(1), 15312. https://doi.org/10.1038/s41598-023-42479-2

Guénot, J., Trotter, Y., Fricker, P., Cherubini, M., Soler, V., & Cottereau, B. R. (2022). Optic

Flow Processing in Patients With Macular Degeneration. *Investigative*

Ophthalmology & Visual Science, *63*(12), 21. https://doi.org/10.1167/iovs.63.12.21

Gupta, A., Mesik, J., Engel, S. A., Smith, R., Schatza, M., Calabrèse, A., van Kuijk, F. J.,

Erdman, A. G., & Legge, G. E. (2018). Beneficial Effects of Spatial Remapping for

Reading With Simulated Central Field Loss. *Investigative Ophthalmology & Visual*

Science, *59*(2), 1105-1112. https://doi.org/10.1167/iovs.16-21404

Hassan, S. E., Lovie-Kitchin, J. E., & Woods, A. R. L. (2002). Vision and Mobility

Performance of Subjects with Age-Related Macular Degeneration: *Optometry and*

Vision Science, *79*(11), 697-707. https://doi.org/10.1097/00006324-200211000-

00007

Huk, A. C., & Heeger, D. J. (2002). Pattern-motion responses in human visual cortex. *Nature*

Neuroscience, *5*(1), 72-75. https://doi.org/10.1038/nn774

Kolster, H., Peeters, R., & Orban, G. A. (2010). The Retinotopic Organization of the Human

Middle Temporal Area MT/V5 and Its Cortical Neighbors. *Journal of Neuroscience*,

30(29), 9801-9820. https://doi.org/10.1523/JNEUROSCI.2069-10.2010

Kuyk, T., & Elliott, J. L. (1999). Visual factors and mobility in persons with age-related

macular degeneration. *Journal of Rehabilitation Research and Development*, *36*(4),

303-312.

Li, J. Q., Welchowski, T., Schmid, M., Mauschitz, M. M., Holz, F. G., & Finger, R. P. (2020).

Prevalence and incidence of age-related macular degeneration in Europe : A

systematic review and meta-analysis. *The British Journal of Ophthalmology*, *104*(8),

1077-1084. https://doi.org/10.1136/bjophthalmol-2019-314422

Little, D., Thulborn, K., & Szlyk, J. (2008). An FMRI study of saccadic and smooth-pursuit eye movement control in patients with age-related macular degeneration. *Investigative Ophthalmology & Visual Science*, *49*(4). https://doi.org/10.1167/iovs.07-0372

Maniglia, M., Soler, V., Cottereau, B., & Trotter, Y. (2018). Spontaneous and training-induced cortical plasticity in MD patients : Hints from lateral masking. *Scientific Reports*, *8*(1), 90. https://doi.org/10.1038/s41598-017-18261-6

Marron, J. A., & Bailey, I. L. (1982). Visual Factors and Orientation-Mobility Performance: *Optometry and Vision Science*, *59*(5), 413-426. https://doi.org/10.1097/00006324-198205000-00009

Masuda, Y., Dumoulin, S. O., Nakadomari, S., & Wandell, B. A. (2008). V1 projection zone signals in human macular degeneration depend on task, not stimulus. *Cerebral Cortex (New York, N.Y.: 1991)*, *18*(11), 2483-2493. https://doi.org/10.1093/cercor/bhm256

Masuda, Y., Takemura, H., Terao, M., Miyazaki, A., Ogawa, S., Horiguchi, H., Nakadomari, S., Matsumoto, K., Nakano, T., Wandell, B. A., & Amano, K. (2021). V1 Projection Zone Signals in Human Macular Degeneration Depend on Task Despite Absence of Visual Stimulus. *Current Biology*, *31*(2), 406-412.e3. https://doi.org/10.1016/j.cub.2020.10.034

McKeefry, D. J., Burton, M. P., Vakrou, C., Barrett, B. T., & Morland, A. B. (2008). Induced deficits in speed perception by transcranial magnetic stimulation of human cortical areas V5/MT+ and V3A. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *28*(27), 6848-6857. https://doi.org/10.1523/JNEUROSCI.1287-08.2008

Morland, A. B. (2015). Organization of the Central Visual Pathways Following Field Defects Arising from Congenital, Inherited, and Acquired Eye Disease. *Annual Review of*

Vision Science, *1*(1), 329-350. https://doi.org/10.1146/annurev-vision-082114-035600

Musel, B., Hera, R., Chokron, S., Alleysson, D., Chiquet, C., Romanet, J.-P., Guyader, N., & Peyrin, C. (2011). Residual abilities in age-related macular degeneration to process spatial frequencies during natural scene categorization. *Visual Neuroscience*, *28*(6), 529-541. https://doi.org/10.1017/S0952523811000435

Papanikolaou, A., Keliris, G. A., Papageorgiou, T. D., Shao, Y., Krapp, E., Papageorgiou, E., Stingl, K., Bruckmann, A., Schiefer, U., Logothetis, N. K., & Smirnakis, S. M. (2014). Population receptive field analysis of the primary visual cortex complements perimetry in patients with homonymous visual field defects. *Proceedings of the National Academy of Sciences*, *111*(16). https://doi.org/10.1073/pnas.1317074111

Peyrin, C., Ramanoël, S., Roux-Sibilon, A., Chokron, S., & Hera, R. (2017). Scene perception in age-related macular degeneration : Effect of spatial frequencies and contrast in residual vision. *Vision Research*, *130*, 36-47. https://doi.org/10.1016/j.visres.2016.11.004

Pigarev, I. N., & Rodionova, E. I. (1998). Two visual areas located in the middle suprasylvian gyrus (cytoarchitectonic field 7) of the cat's cortex. *Neuroscience*, *85*(3), 717-732. https://doi.org/10.1016/S0306-4522(97)00642-8

Pitzalis, S., Galletti, C., Huang, R.-S., Patria, F., Committeri, G., Galati, G., Fattori, P., & Sereno, M. I. (2006). Wide-Field Retinotopy Defines Human Cortical Visual Area V6. *The Journal of Neuroscience*, *26*(30), 7962-7973. https://doi.org/10.1523/JNEUROSCI.0178-06.2006

Pitzalis, S., Sereno, M. I., Committeri, G., Fattori, P., Galati, G., Patria, F., & Galletti, C. (2010). Human v6 : The medial motion area. *Cerebral Cortex (New York, N.Y.: 1991)*, *20*(2), 411-424. https://doi.org/10.1093/cercor/bhp112

Poldrack, R. A. (2007). Region of interest analysis for fMRI. *Social Cognitive and Affective Neuroscience*, *2*(1), 67-70. https://doi.org/10.1093/scan/nsm006

Priebe, N. J., Lisberger, S. G., & Movshon, J. A. (2006). Tuning for Spatiotemporal

Frequency and Speed in Directionally Selective Neurons of Macaque Striate Cortex.

The Journal of Neuroscience, *26*(11), 2941-2950.

https://doi.org/10.1523/JNEUROSCI.3936-05.2006

Ramanoël, S., Chokron, S., Hera, R., Kauffmann, L., Chiquet, C., Krainik, A., & Peyrin, C.

(2018). Age-related macular degeneration changes the processing of visual scenes

in the brain. *Visual Neuroscience*, *35*, E006.

https://doi.org/10.1017/S0952523817000372

Shao, Y., Keliris, G. A., Papanikolaou, A., Fischer, M. D., Zobor, D., Jägle, H., Logothetis, N.

K., & Smirnakis, S. M. (2013). Visual cortex organisation in a macaque monkey with

macular degeneration. *European Journal of Neuroscience*, *38*(10), 3456-3464.

https://doi.org/10.1111/ejn.12349

Silson, E. H., Baker, C. I., Aleman, T. S., Maguire, A. M., Bennett, J., & Ashtari, M. (2023).

Motion-selective areas V5/MT and MST appear resistant to deterioration in

choroideremia. *NeuroImage: Clinical*, *38*, 103384.

https://doi.org/10.1016/j.nicl.2023.103384

Strong, S. L., Silson, E. H., Gouws, A. D., Morland, A. B., & McKeefry, D. J. (2017). A Direct

Demonstration of Functional Differences between Subdivisions of Human V5/MT+.

Cerebral Cortex (New York, NY), *27*(1), 1-10. https://doi.org/10.1093/cercor/bhw362

Szlyk, J. P., & Little, D. M. (2009). An FMRI study of word-level recognition and processing

in patients with age-related macular degeneration. *Investigative Ophthalmology &*

Visual Science, *50*(9), 4487-4495. https://doi.org/10.1167/iovs.08-2258

Talairach, J., & Tournoux, P. (1988). *Co-planar Stereotaxic Atlas of the Human Brain : 3-*

dimensional Proportional System : an Approach to Cerebral Imaging. G. Thieme.

Tarita-Nistor, L., González, E. G., Markowitz, S. N., Lillakas, L., & Steinbach, M. J. (2008).

Increased Role of Peripheral Vision in Self-Induced Motion in Patients with Age-

Related Macular Degeneration. *Investigative Ophthalmology & Visual Science*, *49*(7),

3253-3258. https://doi.org/10.1167/iovs.07-1290

Tootell, R. B. H., Mendola, J. D., Hadjikhani, N. K., Ledden, P. J., Liu, Arthur. K., Reppas, J.

B., Sereno, M. I., & Dale, A. M. (1997). Functional Analysis of V3A and Related

Areas in Human Visual Cortex. *The Journal of Neuroscience*, *17*(18), 7060-7078.

https://doi.org/10.1523/JNEUROSCI.17-18-07060.1997

Tootell, R. B. H., Reppas, J. B., Dale, A. M., Look, R. B., Sereno, M. I., Malach, R., Brady, T.

J., & Rosen, B. R. (1995). Visual motion aftereffect in human cortical area MT

revealed by functional magnetic resonance imaging. *Nature*, *375*(6527), 139-141.

https://doi.org/10.1038/375139a0

Tootell, R., Reppas, J., Kwong, K., Malach, R., Born, R., Brady, T., Rosen, B., & Belliveau, J.

(1995). Functional analysis of human MT and related visual cortical areas using

magnetic resonance imaging. *The Journal of Neuroscience*, *15*(4), 3215-3230.

https://doi.org/10.1523/JNEUROSCI.15-04-03215.1995

Tran, T. H. C., Guyader, N., Guerin, A., Despretz, P., & Boucart, M. (2011). Figure Ground

Discrimination in Age-Related Macular Degeneration. *Investigative Opthalmology &*

Visual Science, *52*(3), 1655. https://doi.org/10.1167/iovs.10-6003

Villeneuve, M. Y., Ptito, M., & Casanova, C. (2006). Global motion integration in the postero-

medial part of the lateral suprasylvian cortex in the cat. *Experimental Brain Research*,

172(4), 485-497. https://doi.org/10.1007/s00221-006-0357-2

Watson, A. B., & Pelli, D. G. (1983). Quest : A Bayesian adaptive psychometric method.

Perception & Psychophysics, *33*(2), 113-120. https://doi.org/10.3758/BF03202828

Westeneng-van Haaften, S. C., Boon, C. J. F., Cremers, F. P. M., Hoefsloot, L. H., Den

Hollander, A. I., & Hoyng, C. B. (2012). Clinical and Genetic Characteristics of Late-

onset Stargardt's Disease. *Ophthalmology*, *119*(6), 1199-1210.

https://doi.org/10.1016/j.ophtha.2012.01.005

Wood, J. M., Lacherez, P. F., Black, A. A., Cole, M. H., Boon, M. Y., & Kerr, G. K. (2009).

Postural Stability and Gait among Older Adults with Age-Related Maculopathy.

Investigative Opthalmology & Visual Science, *50*(1), 482.

965   https://doi.org/10.1167/iovs.08-1942

966   Zeki, S., Watson, J. D., Lueck, C. J., Friston, K. J., Kennard, C., & Frackowiak, R. S. (1991).

967   A direct demonstration of functional specialization in human visual cortex. *The*

968   *Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *11*(3),

969   641-649. https://doi.org/10.1523/JNEUROSCI.11-03-00641.1991

Supplementary Materials

Supplementary Figure 1: Images of the eye fundus and estimation of the scotoma size (left panel) and PRL (middle panel) in three patients (MD2, MD5 and STG2). The right panels provide the artificial scotomas used in the corresponding control participants (CTRL2, CTRL5, STGCTRL2). See Figure 1 for more details.

%%%

Supplementary Figure 2: Comparison between BOLD responses to visual stimulation and to a blank

screen on the lateral cortical surfaces of all MD patients (left panels) and their respective controls (right

panels). See Figure 3 for the details of the legend.

993 ***Supplementary Figure 3:*** Comparison between BOLD responses to visual stimulation and to a blank

994 screen on the medial cortical surfaces of all MD patients (left panels) and their respective controls (right

995 panels). See Figure 3 for the details of the legend.

996 %%

		pSTS
MD1	LH	X
	RH	X
MD2	LH	X
	RH	51 ; -48 ; 10
MD3	LH	-47 ; -46 ; 2
	RH	53 ; -51 ; 5
MD4	LH	-47 ; -47 ; 9
	RH	56 ; -40 ; 5
MD5	LH	X
	RH	X
STG1	LH	-39 ; -49 ; 11
	RH	51 ; -43 ; 12
STG2	LH	-49 ; -50 ; 5
	RH	54 ; -41 ; 6
CTRL1	LH	X
	RH	X
CTRL2	LH	-51 ; -40 ; 3
	RH	52 ; -33 ; 5
CTRL3	LH	-46 ; -38 ; -1
	RH	43 ; -42 ; -1
CTRL4	LH	X
	RH	55 ; -35 ; 10
CTRL5	LH	X
	RH	44 ; -35 ; 1
STGCTRL1	LH	-45 ; -49 ; 10
	RH	53 ; -37 ; 11
STGCTRL2	LH	-47 ; -48 ; 5
	RH	40 ; -54 ; 12
Average patients	LH	-46±4 ; -48±8 ; 7±4
	RH	53±3 ; -45±3 ; 8±2
Average controls	LH	-47±2 ; -44±1 ; 4±2
	RH	48±1 ; -39±1 ; 6±1
Literature	LH	-56 ; -47 ; 8
	RH	52 ; -49 ; 15

998 ***Supplementary Table 1:*** Talairach coordinates of pSTS observed in patients and controls, the

999 calculated mean for both patients and controls, and the coordinates reported in Barton and Brewer

1000 (2017).

1001

1002

Supplementary Figure 4: Activation extents with the hMT+ (A), V3A (B) and V6 (C) ROIs. These

extents were computed as the percentage of voxels passing a threshold of t-score = 3 within each ROI.

For each group, the boxplots provide the first quartile, median, last quartile, and the extreme

percentages of activated voxels. ROIs were defined as spheres of 10 mm of radius centered on the

Talairach coordinates provided in the literature (see the '*Methods*' section) but similar results were

obtained when considering spheres of 5 and 15 mm of radius.

1009

Supplementary Figure 5: Motion-selective activations in the pSTS ROI. For each patient and their

age-matched control (connected by lines), colored dots provide the differences between the average

(across voxels and hemispheres) percentages of signal changes (PSCs) during the translational motion

and control (static dots) conditions. For each group, the boxplots provide the first quartile, median, last

quartile, and the extreme PSC difference. These differences are not significantly different from zero (p-

values = 0.109 in both MD patients and controls, Wilcoxon test). Statistical analyses based on paired

Wilcoxon sign-rank tests did not reveal significant differences between activations in the two groups (p-

values > 0.05).

1018

1019

Supplementary Figure 6: Differences between the average percentages of signal changes (PSCs)

during the translational motion and control (static dots) conditions in the hMT+ (A), V3A (B) and V6 (C)

ROIs as a function of the scotoma size, given by the radius of the associated disk (see the '*Methods*'

section). Dotted and solid lines respectively provide the regressions lines for MD patients and controls.

The corresponding Pearson correlation coefficients R are also shown. None of these correlations were

significant (p-values > 0.05). For patient MD5 (whose scotoma is elongated, see supplementary figure

1) and his control, we used the semi minor axis of the associated ellipse for the scotoma size in this

figure. We obtained similar results when using the semi major axis of the ellipse.

1028

1029