Stable states in an unstable landscape: microbial resistance at the front line of climate change

Authors: Dylan R. Cronin[1,2,3]†, Hannah Holland-Moritz[4,5]†, Derek A. Smith[3]†, Samuel T. N. Aroney[6]†, Suzanne B. Hodgkins[1], Mikayla Borton[7], Yueh-Fen Li[1], Ahmed Zayed[1,2], Kieran Healy[8], IsoGenie Field & Analytic Teams 2010-2017§, EMERGE Institute Coordinators§, Malak M. Tfaily[9,10], Patrick Crill[11], Carmody K. McCalley[12], Kelly Wrighton[7], Ruth K. Varner[11,13,14], Gene W. Tyson[6]‡, Ben J. Woodcroft[6]‡, Sarah C. Bagby[3]‡, Jessica Ernakovich[4,5]‡, Virginia I. Rich[1,15,16]‡*

† These lead authors contributed equally to this work.
‡ These senior authors contributed equally to this work.
§ Collective author designation: list of the authors and affiliations will be at the end of the manuscript.
* Corresponding author. Email: rich.270@osu.edu

Affiliations:
1. Department of Microbiology, The Ohio State University, Columbus, OH, USA
2. Center of Microbiome Science, The Ohio State University, Columbus, OH, USA
3. Department of Biology, Case Western Reserve University, Cleveland, OH, USA
4. Department of Natural Resources and the Environment, University of New Hampshire, Durham, NH, USA
5. Center for Soil Biogeochemistry and Microbial Ecology, University of New Hampshire, Durham, NH, USA
6. Centre for Microbiome Research, School of Biomedical Sciences, Queensland University of Technology (QUT), Translational Research Institute, Woolloongabba, QLD, Australia
7. Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO, USA
8. Department of Sociology, Duke University, Durham, NC, USA
9. Department of Environmental Science, The University of Arizona, Tucson, AZ, USA
10. Bio5 Institute, The University of Arizona, Tucson, AZ, USA
11. Department of Geological Sciences and Bolin Centre for Climate Research, Stockholm University, Stockholm, SE
12. Thomas H. Gosnell School of Life Sciences, Rochester Institute of Technology, Rochester, NY, USA
13. Department of Earth Sciences and Earth Systems Research Center, University of New Hampshire, Durham, NH, USA
14. Mathematics, Science & Engineering Education, University of New Hampshire, Durham, NH, USA
15. Translational Data Analytics Institute, The Ohio State University, Columbus, OH, USA
16. Byrd Polar Climate and Research Center, The Ohio State University, Columbus, OH, USA

Abstract

Microbiome responses to warming may amplify or ameliorate terrestrial carbon loss and thus are a critical unknown in predicting climate outcomes. Because the rapid thaw of permafrost peatlands makes a very large store of soil carbon available to microbial metabolism, understanding microbiome dynamics in these systems is particularly urgent. We quantified microbial warming response over seven years across three habitats in a thawing permafrost peatland, using large-scale multi-omics data. We integrated analyses of organisms (via taxonomy), functions (via metabolic pathways and proteins), and community organization (via network structure and ecological assembly) to deeply characterize response mechanisms. We consistently found a pattern of within-habitat microbiome stability, with virtually no signal of gradual change in the warming period studied. The resistance to change appeared bolstered by habitat-specific dispersal processes and community-level functional redundancy, particularly via versatile carbon generalists. Our findings also reveal key genome-inferred metabolic processes that underlie microbiome stability. Together, our results highlight the importance of understanding the limits of these stabilizing processes and suggest that future research should reorient towards critical habitat transitions.

Main Text

Introduction

Arctic permafrost peatlands, warming at a rate four times faster than the global average[1], contain globally significant reserves of soil organic carbon[2]. This carbon's fate and flux dynamics as carbon dioxide (CO_2) or methane (CH_4) will determine these systems' feedbacks to climate change[3–5]. Soil microbiota play a critical role in controlling these dynamics (reviewed in Jansson & Hofmockel, 2020), but the community structure, assembly, and metabolism of the microbiota are altered in turn by the cascading changes set off by rising Arctic temperatures[7–9]. Ecological factors such as functional stability and metabolic flexibility[10,11] reinforce habitat resistance and resilience to gradual change, but enough disturbance can push communities into non-equilibrium dynamics and trigger reassembly into a new community state[12–14]. The pressing need to understand how soil microbiomes respond to temperature-induced change has motivated numerous studies[15–23], but to date no clear pattern has emerged. Complicating interpretation, this body of work includes varying study designs and disturbance intensities (observation vs. manipulation[21]), timeframes (few- vs. many-year[18,22]), and environmental settings, in which weather, soil properties, depth, plant communities, and hydrology may outweigh climate warming impacts[15–17,19,20,23](see Supplementary note S1). Despite these challenges, understanding the dynamics of soil microbiome response to warming is vital for establishing how climate models should represent these critical ecosystems[24,25].

Model ecosystems create an opportunity for integrative study of these dynamics. Stordalen Mire is a discontinuous permafrost peatland in northern Sweden (68° 21' N, 19° 20' E; Supplementary note S2) that has been intensively studied over decades as its permafrost thaws. These studies have tracked the collapse of patches of dry elevated palsas to partially thawed bogs and fully thawed fens, with total fen extent doubling over the past 50 years[5]. Progression along this thaw gradient is marked by both abiotic and biotic shifts affecting redox

conditions (predominantly oxic in palsa, anoxic in fen), pH, hydrologic connectivity (dry palsa,
perched water tables in bog, permanently inundated fens), and plant, microbial, and viral
community compositions[5,7,9,26–28]. Importantly, these shifts across the thaw gradient induce
increasingly anaerobic microbial lineages and metabolisms, including methanogenesis[9,27],
driving substantial increases in net radiative forcing[5,29]. Because habitat conversion proceeds in
local patches, long-term observation of plant communities has provided many opportunities to
observe the conversion of the diverse forbs of the palsa to *Sphagnum*-dominated bog and
Eriophorum-dominated fen[5,29,30]. These repeated observations suggest that aboveground, at the
landscape scale, these habitat types behave as alternative stable states[31], local minima in a
stability landscape traversed by communities. Previous work has provided a detailed
understanding of microbial communities and processes in short periods within each
habitat[9,30,32,33], but the belowground temporal dynamics within each habitat have so far gone
unexplored.

Each of the alternative stable states observed aboveground, in the plant communities at
the surface of soil cores, could conceal any of three distinct belowground patterns. These
patterns can be visualized in terms of the classic "ball and cup" analogy[31] (Supplementary
Fig.1). First, a habitat's microbiome could be stable as long as its plant community is stable; a
time series of cores with plant communities belonging to one aboveground state would capture
only minor fluctuations belowground (Supplementary Fig.1(i)). Stability could arise either
because the community is insensitive to disturbance (resistance) or because it recovers rapidly
from disturbance (resilience)[10,11,15]. Second, if the microbiome states' minima are shallow and
aboveground-belowground coupling is loose, a microbiome could shift gradually towards the
next state in the canonical thaw progression (i.e., palsa to bog, bog to fen) or to a different
transitional state (Supplementary Fig.1(ii.a, ii.b)). Third, a microbiome could shift abruptly
between the belowground states associated with palsa, bog, and fen, either leading or lagging
aboveground shifts (Supplementary Fig.1(iii.a, iii.b)). A time series of cores with a given plant
community might capture this saltatory microbiome change; however, very long lags could make
the plant and microbial communities appear completely uncoupled, while with very short lags
the pattern would appear as within-habitat stability, with aboveground state consistently
predicting the belowground community. Critically, each of these patterns carries different
implications for how predictive modeling should represent a habitat's soil microbial community
and its activity, and for the type and resolution of data required to parameterize models. More
broadly, distinguishing between these patterns would reveal the structure of plant-soil feedbacks
in a rapidly changing ecosystem.

Here we examined microbiome temporal dynamics in Stordalen Mire's habitats through
three genome-resolved lenses: *organisms* (across taxon ranks ranging from phylum to strain),
functions (encoded metabolic pathways and proteins), and community *organization* (community
network structure and mechanisms of ecological assembly). Our depth-resolved dataset of 199
metagenomes (2.37 Tbp) and 66 metatranscriptomes (1.87 Tbp), from soil cores defined as
palsa, bog, or fen by the plant community observed at the surface, captures seven years of
progressive ~0.7°C/yr belowground warming for each habitat (2011-2017, Fig. 1A) at peak
growing season. This period included significant interannual fluctuations in air temperature and

water table depth (Extended Data Fig.1) and encompassed four of the planet's hottest years on
record[34]. Through each lens, in each habitat, at nearly every depth, we consistently observed
microbiome stability, closely coupled to the alternative stable states aboveground
(Supplementary Fig.1(i)). Together, our analyses suggest that this microbiome stability is driven
by (1) metabolic versatility within individual lineages, (2) functional redundancy between
lineages, and (3) stochastic dispersal in the waterlogged soils of bog and fen. Moreover, our
integrative approach suggests new hypotheses on nutrient-cycling pathways and partnerships in
each of Stordalen Mire's habitats.
142
143 ## **Microbiomes are resistant to increasing ground temperature**
144
Microbiome composition and function resisted change over time
To evaluate microbiome changes, we reconstructed 13,290 metagenome-assembled
genomes (MAGs) that represent 1,806 species-level taxa (i.e., clustered at 95% average
nucleotide identity, Fig. 1A). Together these MAGs captured 73% of the sequenced community
at genus-level [see Methods], which is high for natural systems. Read-mapping-based
abundances revealed that microbial diversity and composition were resistant to change over
seven years within habitat and depth (down to 39 cm) (Fig. 1A and Fig. 1B; Supplementary
Table1). Very little variation in community composition could be accounted for by sample year at
any level of resolution (strain: ≤3.0%, p = 0.03); genus: 1.6%, p = 0.03; phylum: 1.6%, p =
0.005) evaluated by distance-based redundancy analyses on single-copy marker genes (Fig. 2).
Similarly, in 92% of habitats and depths, genome-resolved community composition was not
explained by time (Fig. 1B and Fig. 2; Extended Data Fig. 2) and inter-annual fluctuation was
significant only at certain depths in the fen (for 0-9, 20-29, and 30-39 cm, respectively: R^2: 0.35,
0.62, 0.81; p-value: 0.036, 0.048, 0.048; Extended Data Fig. 3). Only the deeper fen (30-39 cm)
samples exhibited both significant inter-annual fluctuation (Extended Data Fig. 2, Extended Data
Fig. 3) and directional change ($R^2 = 0.50$, p = 0.012) in community composition with time (Fig.
2), and only 8 of 1,806 species were significantly correlated to time (Spearman rho ranging from
-0.7 to 0.91, p < 0.05; Extended Data Fig. 3) (see Supplementary note S3). The stability
observed across a range of taxonomic granularities suggests that the microbial community is
compositionally resistant over time.
165
Encoded microbiome functions were similarly consistent within habitats over time (Fig.
1B, Fig. 2). Beta diversity of community-wide protein clusters (a metric unconstrained by
functional annotation), as well as manually-curated genome-resolved pathways of carbon,
nitrogen, and sulfur metabolism did not significantly change over time within 92% of habitats
and depths (Fig. 2; Extended Data Fig. 4); the sole exception was 30-39 cm in the fen (R^2: 0.49-
0.55, PERMANOVA, p < 0.05 Fig. 1B; Fig. 2; Extended Data Fig. 3). At the coarsest level of
functional annotation (MAG-specific specialization), individual functional guilds of organisms
(e.g., methanogens, homoacetogens, fermenters, and macromolecule degraders) maintained
similar abundances through time. (Expressed microbiome function was not used for temporal
analysis due to discontinuity in the metatranscriptomic time series.) Thus, within the majority of
Stordalen Mire habitats and depths, we observed both compositional and functional resistance
to change over time across multiple levels of granularity.

178
Our third lens, microbial community organization, also revealed stability over time within
each habitat and depth. We used two distinct genome-resolved network approaches to
understand the influence of soil biogeochemical conditions (Weighted Genome Correlation
Network Analysis (WGCNA)) and inter-microbe interactions (multilayer temporal networks
(MTN)) on the organizational structure of the community with time. Habitat-specific WGCNA
modules showed no linear correlation of community composition to time (Pearson, $p<0.05$)
(Supplementary Table 2). MTN microbial co-occurrence networks exhibited variability in specific
microbial interactions over time (Fig 3A, Extended Data Fig. 5A-C; Jaccard similarity coefficient
of triangles), though the overall network structure remained more consistent at broader scales
(Fig. 21C, Fig. 3B, Extended Data Fig. 5A-C; degree distribution and degree correlation). This
suggests that broader community network patterns are resistant to change, even as individual
interactions shift temporally. To understand if subcommunities of microbes were resistant over
time, we asked how many years each MTN subcommunity persisted (Extended Data Fig. 5D).
Persistent subcommunities (occurring in ≥ four of the seven years) accounted for a conservative
estimate of 40% of the abundance-weighted community with time (depending on the interannual
relax rate; Extended data Fig. 5E); thus, while not all microbe-microbe interactions were stable
from year to year, persistent clusters contributed extensively to community composition.
196
Finally, we sought to understand whether the ecological mechanisms that reinforce
stability in each habitat's community organization were themselves changing over time. We
used beta-null modeling[35] to determine whether stochastic (dispersal and ecological drift) or
deterministic (abiotic or biotic selection) processes drove community structure in each habitat
(Extended Data Fig. 6). A change in community assembly mechanisms over time could indicate
a shift in the ecological pressures shaping the community (for example, increased selection due
to changing ground temperature). Instead, beta-null modeling showed that while each habitat
was governed by distinct assembly mechanisms (homogenizing selection in the palsa,
homogenizing dispersal in the water-logged habitats), the ratio between deterministic and
stochastic assembly processes within each habitat did not change with time (Mantel, $p > 0.05$)
(Fig. 1B and Fig. 3C).
208
Microbiome composition and function had weak susceptibility to temperature
Finally, we explored the possibility that organisms, functions, and organization of the
communities were being shaped by subtle or non-monotonic changes in abiotic drivers. Below
we focus on temperature. After accounting for covariation between temperature and depth
(strongest in palsa; Extended Data Fig. 1), genome-resolved community composition was
slightly, but significantly, linearly correlated to ground temperature in bog and fen but not palsa
(Fig. 2, Bray-Curtis dissimilarity: partial mantel rho = 0.152 and 0.24, $p < 0.001$, in bog and fen
respectively). However, when habitat and depth were included in non-linear generalized additive
models, no predictive relationship remained between community composition and temperature
($p = 0.4$; Supplementary note S4), suggesting that the effect of temperature was weak. Echoing
these results, while some WGCNA modules were significantly (Pearson, $p < 0.05$) correlated
with temperature, all but one of these modules were more strongly correlated with depth. Some
of these WGCNA modules were more strongly correlated to carbon cycling metrics (e.g.,

$\delta^{13}C_{CO2}$, $\delta^{13}C_{CH4}$, $\alpha_{CO2-CH4}$ (Extended Data Fig. 7), consistent with previous work at the
site[9,27,28,36,37], and other features (e.g., pH and total nitrogen). Ground temperature had no
significant impact on carbon, nitrogen, or sulfur pathways in the palsa or bog, and only on
xyloglucan degradation in the fen (Supplementary note S4). No community assembly
mechanisms were related to ground temperature or air temperature (day-of-sampling or prior
four weeks) (Mantel, $p > 0.05$), implying that even with lag effects, temperature was not a strong
selective or stochastic force on community formation. As with temporal change, microbiome
sensitivity to temperature was strongest in the fen but weak overall and overshadowed by other
abiotic factors (depth, pH, and nutrient availability).
231
Metabolic connections maintain stability
We next asked whether and how the activity of the microbiota within these communities
might contribute to reinforcing within-habitat stability. Microbiome capacity to degrade organic
matter depends not only on organic matter availability but also on access to other nutrients
including bioavailable nitrogen and sulfur. Access to nitrogen is a fundamental constraint on
microbial growth in northern peatlands, where nitrogen is frequently limiting[38,39]. Similarly,
despite small environmental pools of sulfur species[37,40], sulfur cycling can control organic matter
degradation rates, particularly in the bog[32]. Bottlenecks in nitrogen, sulfur, or carbon flow due to
warming-dependent changes in organic matter profiles, redox conditions, or enzyme kinetics[41]
might disturb the community, but access to alternative pathways or partners for substrate
processing could confer resistance to such pressure.
243
Chitin degradation and nitrogen fixation offer alternative pathways to nitrogen acquisition
When preferred low-molecular-weight organic nitrogen compounds[42,43] are not available,
nitrogen fixation and chitin degradation can provide alternative nitrogen sources. Nitrogen
fixation increases the bioavailable nitrogen pool, benefiting diazotrophs directly and the
community indirectly[44]. Up to 25% of the Stordalen Mire community encoded nitrogen fixation
(compared to ~0.3% in marine microbiomes[45]), and > 75% of encoding MAGs showed pathway
expression (Extended Data Fig. 8).
251
Pathways for degrading chitin ($(C_8H_{13}O_5N)_n$), the second most abundant biopolymer in
nature[46], were even more widespread. An insoluble compound whose breakdown requires
extracellular enzymes, chitin is targeted by many degradation pathways. At Stordalen Mire,
multiple chitin degradation pathways were ubiquitously encoded (abundance weighted median
73%, range 45%–94% (Extended Data Fig. 8)) and expressed (median 95% of encoding cells,
range 59–100% with one outlier sample at 5%). Furthermore, organisms containing these
functions were closely linked to key environmental outputs in WGCNA networks (as Variable
Importance in Projection (VIP) members of modules related to carbon and nitrogen cycling
(Extended Data Fig. 7)). In many aquatic and terrestrial environments, including peatlands,
extracellular chitin degradation products can facilitate commensal and cross-feeding
interactions[46–49], which may promote community stability[50] while buffering against fluctuations in
low-molecular-weight nitrogen availability. The ubiquity of both nitrogen fixation and chitin
degradation pathways across Stordalen Mire habitats may likewise confer resistance.

265
Cryptic sulfur cycling and degradation of sulfur-containing polysaccharides might minimize sulfur limitations

Cryptic sulfur cycling can profoundly influence system biogeochemistry[51]. We observed habitat-wide expression of dissimilatory sulfur oxidation and sulfate reduction (encoded pathway relative abundances ~25% and up to 10%, respectively (Fig. 4A; Extended Data Fig. 8), suggesting the active use of cryptic sulfur cycling at Stordalen Mire. Interestingly, although all microbes require sulfur for growth, we detected assimilatory sulfate reduction in only 25% (fen) to 50% (palsa, bog) of the population (Extended Data Fig. 8). Our analysis of carbohydrate degradation suggests a novel hypothesis: community members may meet some of their sulfur requirements through CAZyme degradation of sulfur-containing polysaccharides (Extended Data Fig. 8), known to be preferentially consumed from plant organic matter[37]. This novel linkage between the carbon and sulfur cycles in permafrost peatlands is analogous to some marine isolates' ability to degrade sulfur polysaccharides and thereby increase H_2, acetate, and sulfate concentrations[52,53]. Such a pathway is expected to release sulfur extracellularly, which could serve as a public good and partially alleviate sulfur limitation for other members of the community.

Carbon generalists are highly prevalent, and macromolecule degradation is widespread

We defined carbon generalists as MAGs encoding ≥ 3 alternative CAZyme pathways and ≥ 3 sugar degradation pathways (Supplementary note S5; Extended Data Fig. 9). These sorts of acquisition-oriented carbon generalists are typical of cold peatlands and other wetlands[11,54,55]. In Stordalen Mire, such MAGs constituted >80% of the community and were taxonomically diverse in each habitat (Fig. 4B) and at nearly all depths. The prevalence of these metabolic generalists produced community-level functional redundancy, with each CAZyme pathway encoded by a median of 531 carbon generalists, and each sugar degradation pathway by a median of 253. Macromolecule and sugar degradation dominated encoded and expressed pathways, particularly in the palsa and bog where >84% of the abundance-weighted community encoded and >50% expressed these pathways (Fig. 4A; Fig. 5A; Extended Data Fig. 4; Extended Data Fig. 9). Fermentation pathways were also highly abundant (47-93% cumulative relative abundance) (Fig. 4A; Extended Data Fig. 4). Collectively, functionally redundant carbon generalists should stabilize the flux of carbon through the system to downstream partners even as environmental conditions and carbon inputs shift.

The flexible community of carbon generalists supports methanogenesis in the bog and fen

Despite low abundances, methanogens have an outsized impact on ecosystem output, creating a pressing need to understand the mechanisms that support their activity. Consistent with previous reports[9], methanogenesis—primarily hydrogenotrophic—was encoded by a handful of MAGs (< 1%) in the bog and fen and was highly expressed there (Fig. 4A). Acetoclastic and methylotrophic methanogenesis were uncommon, with the former present only in the fen (Fig. 4A; Extended Data Fig. 4). We observed that several methanogens persisted throughout our study period (Fig. 5A&B; Supplementary note S6), even though the activity of these metabolic specialists depends on the supply of a narrow range of substrates. Diverse metabolic partnerships that reduce substrate limitation for methanogens have been observed to

309 stabilize aquatic microbial communities[56,57]. We asked whether bog and fen methanogens might
310 likewise partner with generalists and leverage these partners' metabolic flexibility to buffer thaw-
311 induced change.
312
Our network analyses allowed us to address this question in two ways. First, we
examined WGCNA relationships to the fractionation factor $\alpha_{CO2-CH4}$, an isotopic indicator of CH_4
metabolic pathways. The hydrogenotrophic methanogen $Ca.$ 'Methanoflorens' spp. has
previously been observed to be a strong predictor of $\alpha_{CO2-CH4}$ [9,27]. Strikingly, WGCNA not only
supported the correlation of $Ca.$ 'Methanoflorens' spp. with $\alpha_{CO2-CH4}$ but also identified thirteen
non-methanogens (including multiple $Acidimicrobiales$, $Terracidiphilus$ and $Ca.$ Acidiflorens) that
were highly predictive of (VIP > 1.5) and correlated to $\alpha_{CO2-CH4}$ (Spearman's rho, $p < 0.005$).
These correlations were comparable to or higher than that of $Ca.$ 'M. stordalenmirensis' itself
(Fig. 5A), suggesting that these predominantly carbon-generalist, nitrogen-flexible lineages may
be gatekeepers for methanogenic carbon flux. For example, several of the $Ca.$ 'Acidiflorens'
spp. may supply H_2 and formate to $Ca.$ 'M. stordalenmirensis'[9]. Second, we identified persistent
MTN associations between $Ca.$ 'M. stordalenmirensis' and with eight MAGs—all VIPs for $\alpha_{CO2-CH4}$,
$[CO_2]$, $\delta^{13}C_{CO2}$, and/or nitrogen content—from the same WGCNA module (Fig. 5B). These
MAGs included the Dormibacterota $Ca.$ 'Changshengia' spp., which was previously identified as
highly correlated with the fraction of carbon mineralized to CO_2 vs. CH_4 in the bog[9].
328
From close analysis of the metabolic capabilities of methanogen-associated lineages,
we propose a novel hypothesis that trading and acquisition of both nitrogen and carbon
underlies these relationships. We observed that, whereas methanogens with few network
connections tend to encode and express nitrogen fixation, highly connected methanogens
important to network structure (e.g., $Ca.$ 'M. stordalenmirensis') typically lacked recognizable
nitrogen-fixation or other nitrogen-linked decomposition pathways (Supplementary note S6;
Extended Data Fig. 10). In other systems, methanogenesis expression[58] and CO_2 and CH_4
production increase with addition of chitin, which methanogens do not themselves
degrade[48,59,60]. Here, all eight persistent methanogen partners (Fig. 5B) encode and express
chitin degradation, potentially mitigating nitrogen limitation for methanogens. Furthermore, we
identify a potential new linkage between methanogens and their partners through the reductive
glycine pathway. This energy-conserving and amine-recycling carbon-fixation pathway, capable
of producing methanogenesis substrates (pyruvate, acetate, and formate) and bioavailable
nitrogen (as ammonia)[61,62], is encoded and expressed by five of the eight methanogen partners.
These co-occurrence-based observations suggest that nitrogen may play a previously
unrecognized role in metabolic interactions supporting methanogen persistence and activity in
these ecosystems.
346
Habitat-specific microbiome resistance arises from functional redundancy and
dispersal mediated by hydrologic connectivity
Stordalen Mire is warming, causing rapid permafrost thaw and habitat conversion[5]
between visibly different alternative stable states aboveground. Our analysis of the soil
microbiome's organization, organisms, and functions in cores defined by their surface plant
communities suggests that belowground communities in palsa, bog, and fen are likewise stable

and tightly coupled to the aboveground state, to within the limits of our study's ability to detect lags. What ecological dynamics maintain the microbial communities' resistance to change? We hypothesize that functional redundancy, metabolic versatility, and homogenizing dispersal drive ecological resistance (Fig. 6).

Functional redundancy and functional similarity have been proposed to stabilize microbial communities via the "portfolio effect"[63–65]. We saw strong signals of both community-level functional redundancy and species-level metabolic versatility in the stable microbiomes of palsa, bog, and fen, each of which was dominated by diverse taxa encoding multiple alternative pathways for carbon and nitrogen acquisition (Fig. 4; Supplementary note S5; Extended Data Fig. 9). Studies of generalists have typically focused on habitat generalists rather than resource generalists[66] (Supplementary note S5), leaving the prevalence of versatile resource generalists in most habitats unknown. We argue that resource generalists merit attention as ecological stabilizers due to their potential for a species-level portfolio effect. That is, just as community function can be maintained by reciprocal changes in efficiency between two compensating species, so should a metabolically versatile cell be able to maintain metabolic activity by rebalancing flux between alternative pathways as conditions change. Increased access via mobile genetic elements to genes providing metabolic versatility would further backstop these stabilizing forces[67]. If redundant pathways within redundant populations stabilize communities, more narrowly distributed specialists[11] would be shielded from disturbance and the loss of taxa from critical functional groups.

Bog and fen microbiomes are likely further stabilized by homogenizing dispersal, i.e., dispersal high enough to result in a community that is more similar to a neighbor than would be expected by chance (Fig. 3C; Fig. 6; Extended Data Fig. 6). Dispersal dynamics are particularly important during disturbance, when dispersing organisms can determine ecosystem trajectory[68]. Local dispersal can promote community resistance and resilience by "rescuing" disturbed communities, preventing cross-site divergence after local population reductions or extinctions[10,14]. This is likely synergistic with functional redundancy and metabolic versatility, which improve the odds that dispersal will resupply organisms that fill key roles.

Conclusions

At Stordalen Mire, aboveground plant community reassembly makes habitat conversions detectable from space[69], but it is the belowground community that drives methane flux, and the dynamics of belowground change are unknown and not easy to remotely observe. Such knowledge gaps are a challenge in eco- and earth system climate models, which must calibrate the representation of microbial complexity[24] often in the absence of high-resolution datasets and parameter estimations[70,71]. The analyses presented here demonstrate that, despite warming and landscape-scale change[5], habitat-specific microbiome complexity can be distilled to essential functions that are stable during peak growing season on a near-decadal timescale.

Collectively, our findings offer new insights into processes underlying microbial community stability in critical permafrost ecosystems. Our integrated analysis over multiple levels of granularity for organismal ranks, metabolic functions, and community organization

revealed little to no significant temporal change within habitats over seven years of progressive
belowground warming. This analysis establishes a collection of metrics for assessing
community response to environmental disturbance. Multiple independent analyses suggest that
the stability of palsa, bog, and fen microbiomes under temperature-induced climate change
depends on functional redundancy within communities and dispersal-mediated resupply of
organisms encoding essential metabolic versatility.
403
To complement these insights into Stordalen Mire's microbiome stability landscape,
future sampling must target periods of habitat transition (Extended Data Fig. 1B), when
stabilizing capacity is exceeded. Further detailed study at spatial, temporal, and genetic
resolutions not captured in this study may reveal other mechanisms of microbial response to
changing conditions, as well as improving bounds on any lag between aboveground and
belowground stable-state changes. For instance, spring and fall observations capturing
changing phenology may illuminate an erosion of stability driven by aboveground/belowground
asynchronies. In addition, increased application of long-read sequencing may help resolve
strain-level functional differences, including those mediated by the activity of mobile genetic
elements. Despite these remaining unknowns, this work reveals the clear existence of stable
microbial communities from habitats in a rapidly warming unstable Arctic landscape.
415
416
417

References and Notes

1.　Rantanen, M. *et al.* The Arctic has warmed nearly four times faster than the globe since

1979. *Commun. Earth Environ.* **3**, 1–10 (2022).

2.　Schuur, E. A. G. *et al.* Permafrost and Climate Change: Carbon Cycle Feedbacks From the

Warming Arctic. *Annu. Rev. Environ. Resour.* **47**, 343–371 (2022).

3.　Hugelius, G. *et al.* Estimated stocks of circumpolar permafrost carbon with quantified

uncertainty ranges and identified data gaps. *Biogeosciences* **11**, 6573–6593 (2014).

4.　Kuhn, M. A. *et al.* Controls on Stable Methane Isotope Values in Northern Peatlands and

Potential Shifts in Values Under Permafrost Thaw Scenarios. *J. Geophys. Res.*

Biogeosciences **129**, e2023JG007837 (2024).

5.　Varner, R. K. *et al.* Permafrost thaw driven changes in hydrology and vegetation cover

increase trace gas emissions and climate forcing in Stordalen Mire from 1970 to 2014.

Philos. Trans. R. Soc. Math. Phys. Eng. Sci. **380**, 20210022 (2021).

6. Jansson, J. K. & Hofmockel, K. S. Soil microbiomes and climate change. *Nat. Rev.*

Microbiol. **18**, 35–46 (2020).

7. Hough, M. *et al.* Coupling plant litter quantity to a novel metric for litter quality explains C

storage changes in a thawing permafrost peatland. *Glob. Change Biol.* **28**, 950–968 (2022).

8. Persson, A., Hasan, A., Tang, J. & Pilesjö, P. Modelling Flow Routing in Permafrost

Landscapes with TWI : An Evaluation against Site-Specific Wetness Measurements. *Trans.*

GIS **16**, 701–713 (2012).

9. Woodcroft, B. J. *et al.* Genome-centric view of carbon processing in thawing permafrost.

Nature **560**, 49–54 (2018).

10. De Vries, F. & Shade, A. Controls on soil microbial community stability under climate

change. *Front. Microbiol.* **4**, (2013).

11. Weedon, J. T. *et al.* Compositional Stability of the Bacterial Community in a Climate-

Sensitive Sub-Arctic Peatland. *Front. Microbiol.* **8**, (2017).

12. Chang, C. & HilleRisLambers, J. Integrating succession and community assembly

perspectives. *F1000Research* **5**, 2294 (2016).

13. Fofana, A. *et al.* Mapping substrate use across a permafrost thaw gradient. *Soil Biol.*

Biochem. **175**, 108809 (2022).

14. Shade, A. *et al.* Fundamentals of Microbial Community Resistance and Resilience. *Front.*

Microbiol. **3**, (2012).

15. Curiel Yuste, J. *et al.* Strong functional stability of soil microbial communities under semiarid

Mediterranean conditions and subjected to long-term shifts in baseline precipitation. *Soil*

Biol. Biochem. **69**, 223–233 (2014).

16. DeAngelis, K. M. *et al.* Long-term forest soil warming alters microbial communities in

temperate forest soils. *Front. Microbiol.* **6**, (2015).

17. Dove, N. C. *et al.* Depth dependence of climatic controls on soil microbial community activity

and composition. *ISME Commun.* **1**, 1–11 (2021).

18. Feng, J. *et al.* Warming-induced permafrost thaw exacerbates tundra soil carbon

decomposition mediated by microbial community. *Microbiome* **8**, 3 (2020).

19. Gutknecht, J. L. M., Field, C. B. & Balser, T. C. Microbial communities and their responses

to simulated global change fluctuate greatly over multiple years. *Glob. Change Biol.* **18**,

2256–2269 (2012).

20. Monteux, S. *et al.* Long-term in situ permafrost thaw effects on bacterial communities and

potential aerobic respiration. *ISME J.* **12**, 2129–2141 (2018).

21. Wu, L. *et al.* Reduction of microbial diversity in grassland soil is driven by long-term climate

warming. *Nat. Microbiol.* **7**, 1054–1062 (2022).

22. Xue, K. *et al.* Tundra soil carbon is vulnerable to rapid microbial decomposition under

climate warming. *Nat. Clim. Change* **6**, 595–600 (2016).

23. Zhu, K., Chiariello, N. R., Tobeck, T., Fukami, T. & Field, C. B. Nonlinear, interacting

responses to climate limit grassland production under global change. *Proc. Natl. Acad. Sci.*

113, 10589–10594 (2016).

24. Kyker-Snowman, E. *et al.* Increasing the spatial and temporal impact of ecological research:

A roadmap for integrating a novel terrestrial process into an Earth system model. *Glob.*

Change Biol. **28**, 665–684 (2022).

25. Wieder, W. R., Bonan, G. B. & Allison, S. D. Global soil carbon projections are improved by

modelling microbial processes. *Nat. Clim. Change* **3**, 909–912 (2013).

26. Emerson, J. B. *et al.* Host-linked soil viral ecology along a permafrost thaw gradient. *Nat.*

Microbiol. **3**, 870–880 (2018).

27. McCalley, C. K. *et al.* Methane dynamics regulated by microbial community response to

permafrost thaw. *Nature* **514**, 478–481 (2014).

28. Mondav, R. *et al.* Microbial network, phylogenetic diversity and community membership in

the active layer across a permafrost thaw gradient. *Environ. Microbiol.* **19**, 3201–3218

(2017).

29. Christensen, T. R. *et al.* Thawing sub-arctic permafrost: Effects on vegetation and methane

emissions. *Geophys. Res. Lett.* **31**, (2004).

30. Hodgkins, S. B. *et al.* Changes in peat chemistry associated with permafrost thaw increase

greenhouse gas production. *Proc. Natl. Acad. Sci.* **111**, 5819–5824 (2014).

31. Beisner, B., Haydon, D. & Cuddington, K. Alternative stable states in ecology. *Front. Ecol.*

Environ. **1**, 376–382 (2003).

32. Freire-Zapata, V. *et al.* Microbiome–metabolite linkages drive greenhouse gas dynamics

over a permafrost thaw gradient. *Nat. Microbiol.* (2024) doi:10.1038/s41564-024-01800-z.

33. McGivern, B. B. *et al.* Microbial polyphenol metabolism is part of the thawing permafrost

carbon cycle. *Nat. Microbiol.* **9**, 1454–1466 (2024).

34. NOAA National Centers for Environmental Information. *NOAA National Centers for*

Environmental Information, Monthly Global Climate Report for Annual 2017.

https://www.ncei.noaa.gov/access/monitoring/monthly-report/global/201713 (2018).

35. Stegen, J. C. *et al.* Quantifying community assembly processes and identifying features that

impose them. *ISME J.* **7**, 2069–2079 (2013).

36. Wilson, R. M. *et al.* Soil metabolome response to whole-ecosystem warming at the Spruce

and Peatland Responses under Changing Environments experiment. *Proc. Natl. Acad. Sci.*

118, e2004192118 (2021).

37. Wilson, R. M. *et al.* Plant organic matter inputs exert a strong control on soil organic matter

decomposition in a thawing permafrost peatland. *Sci. Total Environ.* **820**, 152757 (2022).

38. Du, E. *et al.* Global patterns of terrestrial nitrogen and phosphorus limitation. *Nat. Geosci.*

13, 221–226 (2020).

39. Elser, J. J. *et al.* Global analysis of nitrogen and phosphorus limitation of primary producers

in freshwater, marine and terrestrial ecosystems. *Ecol. Lett.* **10**, 1135–1142 (2007).

40. Perryman, C. R. *et al.* Thaw Transitions and Redox Conditions Drive Methane Oxidation in a

Permafrost Peatland. *J. Geophys. Res. Biogeosciences* **125**, e2019JG005526 (2020).

41. Hodgkins, S. B. Changes in organic matter chemistry and methanogenesis due to permafrost thaw in a subarctic peatland. (The Florida State University, United States -- Florida, 2016).

42. Graham, E. B., Yang, F., Bell, S. & Hofmockel, K. S. High Genetic Potential for Proteolytic Decomposition in Northern Peatland Ecosystems. *Appl. Environ. Microbiol.* **85**, (2019).

43. Norman, J. S. & Friesen, M. L. Complex N acquisition by soil diazotrophs: how the ability to release exoenzymes affects N fixation by terrestrial free-living diazotrophs. *ISME J.* **11**, 315–326 (2017).

44. Lerch, B. A., Smith, D. A., Koffel, T., Bagby, S. C. & Abbott, K. C. How public can public goods be? Environmental context shapes the evolutionary ecology of partially private goods. *PLOS Comput. Biol.* **18**, e1010666 (2022).

45. Delmont, T. O. *et al.* Nitrogen-fixing populations of Planctomycetes and Proteobacteria are abundant in surface ocean metagenomes. *Nat. Microbiol.* **3**, 804–813 (2018).

46. Beier, S. & Bertilsson, S. Bacterial chitin degradation—mechanisms and ecophysiological strategies. *Front. Microbiol.* **4**, (2013).

47. Belova, S. E. *et al.* Hydrolytic Capabilities as a Key to Environmental Success: Chitinolytic and Cellulolytic Acidobacteria From Acidic Sub-arctic Soils and Boreal Peatlands. *Front. Microbiol.* **9**, (2018).

48. De Tender, C. *et al.* Peat substrate amended with chitin modulates the N-cycle, siderophore and chitinase responses in the lettuce rhizobiome. *Sci. Rep.* **9**, 9890 (2019).

49. Kaczmarek, M. B., Struszczyk-Swita, K., Li, X., Szczęsna-Antczak, M. & Daroch, M. Enzymatic Modifications of Chitin, Chitosan, and Chitooligosaccharides. *Front. Bioeng. Biotechnol.* **7**, (2019).

50. Lopes, W., Amor, D. R. & Gore, J. Cooperative growth in microbial communities is a driver of multistability. *Nat. Commun.* **15**, 4709 (2024).

51. Canfield, D. E. *et al.* A Cryptic Sulfur Cycle in Oxygen-Minimum–Zone Waters off the

Chilean Coast. *Science* **330**, 1375–1378 (2010).

52. Helbert, W. Marine Polysaccharide Sulfatases. *Front. Mar. Sci.* **4**, (2017).

53. Van Vliet, D. M. *et al.* Anaerobic Degradation of Sulfated Polysaccharides by Two Novel

Kiritimatiellales Strains Isolated From Black Sea Sediment. *Front. Microbiol.* **10**, 253 (2019).

54. Oliverio, A. M. *et al.* Rendering the metabolic wiring powering wetland soil methane

production. Preprint at https://doi.org/10.1101/2024.02.06.579222 (2024).

55. Tveit, A. T., Urich, T., Frenzel, P. & Svenning, M. M. Metabolic and trophic interactions

modulate methane production by Arctic peat microbiota in response to warming. *Proc. Natl.*

Acad. Sci. **112**, E2507–E2516 (2015).

56. Ciccarese, D. *et al.* Rare and localized events stabilize microbial community composition

and patterns of spatial self-organization in a fluctuating environment. *ISME J.* **16**, 1453–

1463 (2022).

57. Embree, M., Liu, J. K., Al-Bassam, M. M. & Zengler, K. Networks of energetic and metabolic

interactions define dynamics in microbial communities. *Proc. Natl. Acad. Sci.* **112**, 15450–

15455 (2015).

58. Ivanova, A. A., Wegner, C.-E., Kim, Y., Liesack, W. & Dedysh, S. N. Identification of

microbial populations driving biopolymer degradation in acidic peatlands by

metatranscriptomic analysis. *Mol. Ecol.* **25**, 4818–4835 (2016).

59. Kang, H., Freeman, C., Soon park, S. & Chun, J. N-Acetylglucosaminidase activities in

wetlands: a global survey. *Hydrobiologia* **532**, 103–110 (2005).

60. Wörner, S. & Pester, M. Microbial Succession of Anaerobic Chitin Degradation in

Freshwater Sediments. *Appl. Environ. Microbiol.* **85**, e00963-19 (2019).

61. Evans, P. N. *et al.* An evolving view of methane metabolism in the Archaea. *Nat. Rev.*

Microbiol. **17**, 219–232 (2019).

62. Sánchez-Andrea, I. *et al.* The reductive glycine pathway allows autotrophic growth of

Desulfovibrio desulfuricans. *Nat. Commun.* **11**, 5090 (2020).

63. Allison, S. D. & Martiny, J. B. H. Resistance, resilience, and redundancy in microbial

communities. *Proc. Natl. Acad. Sci.* **105**, 11512–11519 (2008).

64. Doak, D. F. *et al.* The Statistical Inevitability of Stability-Diversity Relationships in

Community Ecology. *Am. Nat.* **151**, 264–276 (1998).

65. Thibaut, L. M. & Connolly, S. R. Understanding diversity–stability relationships: towards a

unified model of portfolio effects. *Ecol. Lett.* **16**, 140–150 (2013).

66. Bell, T. H. & Bell, T. Many roads to bacterial generalism. *FEMS Microbiol. Ecol.* **97**, fiaa240

(2020).

67. Guo, J. *et al.* Mobile genetic elements that shape microbial diversity and functions in

thawing permafrost soils. Preprint at https://doi.org/10.1101/2025.02.12.637893 (2025).

68. HilleRisLambers, J., Adler, P. B., Harpole, W. S., Levine, J. M. & Mayfield, M. M. Rethinking

Community Assembly through the Lens of Coexistence Theory. *Annu. Rev. Ecol. Evol. Syst.*

43, 227–248 (2012).

69. Varner, R. K. *et al.* From Archaea to the atmosphere: remotely sensing Arctic methane.

Preprint at https://doi.org/10.1101/2025.02.13.638097 (2025).

70. Chang, K.-Y. *et al.* Methane Production Pathway Regulated Proximally by Substrate

Availability and Distally by Temperature in a High-Latitude Mire Complex. *J. Geophys. Res.*

Biogeosciences **124**, 3057–3074 (2019).

71. Wieder, W. R. *et al.* Explicitly representing soil microbial processes in Earth system models.

Glob. Biogeochem. Cycles **29**, 1782–1800 (2015).

Collective Author Designations

IsoGenie Field and Analytic Teams 2010-2017

Jeffrey P. Chanton[1], Darya Anderson[2], Gareth Trubl[3], Joachim Jansen[4], Moira Hough[5], Nicole Irwin-Raab[3], Rachel M. Wilson[1], Rhiannon Mondav[6], Robert Jones[3], Sky Dominguez[7], Apryl Perry[8]

1. Earth, Ocean, and Atmospheric Sciences, Florida State University, Tallahassee, FL, USA
2. Department of Soil, Water, and Environmental Sciences, The University of Arizona, Tucson, AZ, USA
3. Department of Microbiology, The Ohio State University, Columbus, OH, USA
4. Department of Ecology and Genetics, Uppsala University, Uppsala, SE
5. College of Forest Resources and Environmental Science, Michigan Technological University, Houghton, MI, USA
6. Centre for Environmental and Climate Science, Lund University, Lund, SE
7. Department of Ecology and Evolutionary Biology, The University of Arizona, Tucson, AZ, USA
8. Department of Earth Sciences and Earth Systems Research Center, University of New Hampshire, Durham, NH, USA

EMERGE Institute Coordinators

Jeffrey P. Chanton[1], Matthew B. Sullivan[2,3], Regis Ferriere[4], Scott R. Saleska[4], Rachel M. Wilson[1], Benjamin Bolduc[2], Eoin L. Brodie[5], Maria Florencia Fahnestock[6], Michael Ibba[7]

1. Earth, Ocean, and Atmospheric Sciences, Florida State University, Tallahassee, FL, USA
2. Department of Microbiology, The Ohio State University, Columbus, OH, USA
3. Center of Microbiome Science, The Ohio State University, Columbus, OH, USA
4. Department of Ecology and Evolutionary Biology, The University of Arizona, Tucson, AZ, USA
5. Earth and Environmental Sciences, Lawrence Berkeley National Laboratory, Berkeley, CA, USA
6. Department of Natural Resources and the Environment, University of New Hampshire, Durham, NH, USA
7. Schmid College of Science and Technology, Chapman University, Orange, CA, USA

Acknowledgments

Non-author contributions:
This study has been made possible by data provided by Abisko Scientific Research Station and the Swedish Infrastructure for Ecosystem Science (SITES) as well as the Swedish Polar

Research Secretariat. We thank the Swedish Polar Research Secretariat and SITES for the
support of the work done at the Abisko Scientific Research Station. A portion of this analysis
and non-monetary technical support was also provided through the U.S. Department of Energy
user facilities programs. Computational work was supported in part by an allocation of
computing time from the Ohio Supercomputer Center. Lastly, we thank contracted science artist
Elena Hartley for co-development of Figure 6.

Funding: We thank the many funding sources of this work, including:
● US National Science Foundation, Biology Integration Institutes Program, Award #
2022070 (VIR and RKV, with SH, MMT, GWT, BW, JE, SCB, and the EMERGE Institute
Coordinators)
● Genomic Science Program of the United States Department of Energy Office of
Biological and Environmental Research (BER), grants DE-SC0004632, DE-SC0010580
and DE-SC0016440 (SRS and VIR, with GWT, MMT). Associated with the third of these
was a BER Support Science award (Proposal #503530 DOI:
10.46936/10.25585/60001148) for nucleic acid sequencing for a portion of these
samples, conducted by the U.S. Department of Energy Joint Genome Institute
(https://ror.org/04xm1d337), a DOE Office of Science User Facility supported by the
Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-
05CH11231.
● US National Science Foundation MacroSystems Program, Award # NSF EF 1241037
(RV).
● US National Science Foundation Research for Undergraduates Program, the Northern
Ecosystems Research for Undergraduates, Award # NSF REU EAR 1063037 (RV)
● Nucleic acid sequencing and and dissolved organic matter fourier transform ion
cyclotron resonance mass spectrometry were partially supported by a Facilities
Integrating Collaborations for User Science (FICUS) awards, DOI:
10.46936/fics.proj.2016.49521/60006018 and 503547, which used resources at the DOE
Joint Genome Institute (https://ror.org/04xm1d337) and the Environmental Molecular
Sciences Laboratory (https://ror.org/04rc0xn13), which are DOE Office of Science User
Facilities. Both facilities are sponsored by the Office of Biological and Environmental
Research and operated under Contract Nos. DE-AC02-05CH11231 (JGI) and DE-AC05-
76RL01830 (EMSL).
● SITES is supported by the Swedish Research Council's grant 4.3-2021-00164

Author Contributions: DC, SA, DS, and HH-M all contributed equally to this work. Co-first
authors have the right to list their name first in their CV. In addition, VR, SB, GT, BW, and JE
are all co-senior authors and contributed equally to this work, order was paired with first author
order and all may list their name last on their CV. Following the CRediT model, authors made
the following contributions

Conceptualization - VR, CM, SH, RV, EMERGE Institute Coordinators, DC, HH-M, DS, SA, JE,
SB, BW, MT, PC
Data Curation - DC, HH-M, DS, SA, VR, BW, SH, FL, RV

Formal Analysis - DC, HH-M, DS, SA, CM, KW, MB, IsoGenie Field & Analytic Teams 2011-2017

Funding acquisition - VR, JE, SB, BW, GT, KW, SH, RV, EMERGE Institute Coordinators, MT, PC

Investigation - DC, HH-M, DS, SA, CM, SH, FL, RV, IsoGenie Field & Analytic Teams 2011-2017, PC

Methodology Development - DC, HH-M, DS, SA, VR, PC

Project Administration - DC, HH-M, DS, SA, VR, JE, SB, BW, GT, KW, SH, RV, EMERGE Institute Coordinators, PC

Resources Provision - VR, KW, SH, RV, EMERGE Institute Coordinators, PC

Software Programming - DC, HH-M, SA, KW, MB, SH

Supervision Oversight - VR, JE, KW, RV, MT

Validation Verification- SA, DC, HH-M

Visualization - DC, HH-M, DS, SA, VR, SB, KH

Writing - DC, HH-M, DS, SA, VR, JE, SB, BW, GT, CM, KW, MB, SH, FL, RV, IsoGenie Field & Analytic Teams 2011-2017, EMERGE Institute Coordinators, KH, MT, PC

Competing interests: Authors declare that they have no competing interests.

Data and materials availability:
Sample Metadata Sheet v1.0.0, containing field sample information and geochemical measurements are available on Zenodo at https://zenodo.org/records/12827096. Gas flux measurements are available on Zenodo at https://doi.org/10.5281/zenodo.13377109 and https://emerge-db.asc.ohio-state.edu/datasources/0138_DailyFlux_20122018. Temperature and water table summary for lagged effects analysis are available on Zenodo at https://doi.org/10.5281/zenodo.14189390. Gas flux summary for lagged effects analysis is available on Zenodo at https://doi.org/10.5281/zenodo.14834760. Water table depth measurements used for determining lagged effects are available on Zenodo at https://doi.org/10.5281/zenodo.10420396. ANS temperature measurements used in supplemental (Extended Data Fig.1) were recorded at the Abisko Scientific Research Station (ANS), Sveriges meteorologiska och hydrologiska institut (SMHI). (2024). Meteorological data from Abisko Observatory (Abisko: station 188800), daily mean air temperatures, 2011-2017. Retrieved [05/30/2024], from https://www.smhi.se/data/meteorologi/ladda-ner-meteorologiska-observationer#stationid=188800.

All metagenomes and metatranscriptomes used in this study are available in NCBI under the umbrella BioProject PRJNA888099. SRA accessions for the reads used for the interannual microbiome comparisons are listed in Supplementary Table 4, and those used to generate the MAGs are listed in the metadata table accompanying those MAGs (https://doi.org/10.5281/zenodo.13901514). Metagenome-assembled genomes (MAGs) are available in Zenodo at https://doi.org/10.5281/zenodo.13901514.

Figures

Figure 1: Stordalen Mire microbiomes resist temporal change. A) The three main thaw-

gradient habitats (palsas, bogs, and fens; colored throughout as brown, green and blue,

respectively) host microbiomes with distinct taxonomy (treemap with dereplicated MAGs as

rectangles, colored by phyla and sized proportional to relative abundance). Soil temperatures

increased significantly with time (lines indicate significant linear relationships, p<0.05, middle

panel). Microbiome beta diversity did not change significantly with time (bottom panels); each

point is one microbiome's average Bray-Curtis dissimilarity to those at the same depth for the

earliest baseline year (2011, or the first year for which at least two depth-matched microbiomes

were available). Black lines are the running dissimilarity average. Reference-year microbiomes'

within-year dissimilarities are indicated by gray dots, with their ranges and averages extended

as shaded gray bars and dashed lines, respectively. **B)** Habitat and depth structure microbiome

beta diversity data more strongly than year, observed in non-metric multidimensional scaling

(NMDS) ordinations of each sample's MAG-based taxonomic (top two panels) and functional

(lower two panels) composition, colored by year (purple shading) or depth (orange shading).

Habitats are denoted by shape (palsa, square; bog, circle; fen, triangle) and demarcated by

ellipses (dashed-line ellipses reflect each habitat's multivariate normal distribution, while shaded

ellipses are the multivariate t-distribution). Functions are cumulative pathway abundances: each

complete pathway observed in a MAG was weighted by that MAG's relative abundance, then all

occurrences are summed for each pathway.

Figure 2: Organisms, functions, and organizations exhibit temporal stability across nearly all Stordalen Mire habitats and depths examined. Granular heatmap of microbiome temporal change, in each habitat and depth. Change was assessed spanning organisms (from phylum to strains diversity metrics, both single copy marker genes and MAGs), functions (from pathways to protein clusters, both annotatable and non-annotatable), and organization (both network structure and community assembly metrics of stochasticity and determinism). The amount of variation explained by time (as Mantel R or R^2 depending on statistical test appropriate for each data type; see Methods) is indicated with the intensity of the shaded color; white asterisks denote significant correlations to time. Across all granularities surveyed only deeper fen (30-39 cm) showed significant change over time.

Figure 3: Microbiome organization is consistent within habitats over time. A) Multilayer

temporal networks (MTN) depict correlations between MAGs within annual networks for each

habitat. Each network layer corresponds to a specific year, with nodes representing MAGs.

Black edges represent significant (p <0.05) correlations (>0.3) between MAGs within the same

habitat and year. **B)** Pairwise interyear correlations (Spearman's rho) in network node degree

(x-axis, base year; colored points, comparison years) within each habitat. Top, palsa; middle,

bog; bottom, fen. Note that the fen's recovery to a near-stable state after 2013 suggests strong

resilience in this habitat. **C)** Assembly processes detected within each habitat (rows) by βNTI

and RCBC modeling in interyear comparisons. For each year, columns show pairwise

comparisons to all other years (self-comparisons are obscured by black bars). Stacked bar

755   colors denote the proportion of different assembly processes detected in each pairwise

756   comparison, revealing overall stability in assembly within habitats over time.

A

Elemental metabolism
- Carbon
- Nitrogen
- Sulfur

Encoded

Expressed

Palsa Bog Fen

B

Specialization
- Generalist
- Methanogen
- Fermenter
- Macromolecule degrader
- Methanotroph
- Homoacetogen
- Monomer degrader
- Other

Figure 4: Microbiome metabolisms differ in carbon, nitrogen, and sulfur cycling and

metabolic specialization across the thaw gradient. A) Aggregated over the study period, key

pathways in carbon, nitrogen, and sulfur metabolism (top center), metagenomic (middle row)

and metatranscriptomic (bottom row) data show differences in microbial investment in

methanotrophy and carbon fixation, nitrogen fixation, and sulfur redox cycling. **B)** Functional

composition of palsa, bog, and fen communities. Treemap rectangles are MAGs colored by

carbon metabolic specialization and scaled by relative abundance.

Figure 5: Microbiome metabolisms across the thaw gradient are interlinked by carbon

and nitrogen cycling. A) Carbon and nitrogen-related metapathways present in MAGs

belonging to the weighted genome correlation network analysis (WGCNA) module correlated

with the isotopic fractionation of carbon between carbon dioxide and methane carbon ($\alpha_{CO2-CH4}$)

in the bog. MAGs are colored by carbon metabolic specialization as in fig. 4 and ordered by VIP

score. **B)** Bog network structure of the persistent subcommunity of MAGs connected to *Ca.* 'M.

stordalenmirensis' in multi-layer temporal network (MTN) modeling. MAGs were included if they

maintained a connection to *Ca.* 'M. stordalenmirensis' for at least six years, had a VIP score >

1, and a correlation > 0.5 to at least one environmental variable. Edge color represents the

number of years MAGs were connected to *Ca.* 'M. stordalenmirensis'; node color indicates MAG

specialization, with colors as in Fig. 4B. We note that many other MAGs were connected to the

methanogen, but not persistently. Subplots show the Spearman correlation coefficient (rho) of

each MAGs to key environmental variables (teal) and their VIP value (pink).

Figure 6: Within-habitat communities resist change until increasing environmental pressures trigger transitions along the thaw gradient. Community composition fluctuates but does not change steadily with time in each habitat. Population density increases in tandem with water saturation [9]. Across the thaw gradient, ecological assembly shifts from predominantly deterministic processes in the palsa (community assembly, purple) to predominantly stochastic processes in the fen (orange), while the balance of functional specializations shifts from predominantly generalists (red) to a more metabolically diverse range of specialists. We hypothesize that functional redundancy, dispersal (bog, fen), and metabolic versatility (palsa, bog) drive within-habitat ecological resistance.