Stable states in an unstable landscape: microbial resistance at the front line of climate change

Authors: Dylan Cronin[1†], Hannah Holland-Moritz[2†], Derek Smith[3†], Samuel T. N. Aroney[4†], Suzanne Hodgkins[1], Mikayla Borton[5], Yueh-Fen Li[1], Kieran Healy[6], IsoGenie Field & Analytic Teams 2010-2017, EMERGE Institute Coordinators, Malak M. Tfaily[7], Patrick Crill[8], Carmody K. McCalley[9], Kelly Wrighton[5], Ruth Varner[2], Gene W. Tyson[4*], Ben Woodcroft[4*], Sarah C. Bagby[3*], Jessica Ernakovich[2*], Virginia I. Rich[1*].
† These authors contributed equally to this work, * Corresponding authors

[1]The Ohio State University, Columbus, OH; [2]University of New Hampshire, Durham, NH; [3]Case Western Reserve University, Cleveland, OH; [4]Queensland University of Technology, Brisbane, AU; [5]Colorado State University, Fort Collins, CO; [6]Duke University, Durham, NC; [7]University of Arizona, Tucson, AZ; [8]Stockholm University, Stockholm, SE; [9]Rochester Institute of Technology, Rochester, NY

One Sentence Summary

In a warming permafrost peatland, microbiome shifts are concerted and correspond to habitat change, rather than proceeding gradually with rising temperature, with multiple lines of evidence indicating that this microbiome ecological resistance is linked to functional redundancy and high dispersal.

Abstract

Microbiome responses to warming may amplify or ameliorate terrestrial carbon loss and thus are a critical unknown in predicting climate outcomes. We quantified microbial warming response over seven years in a thawing permafrost peatland, using large-scale multi-omics data. We integrated analyses of organisms (via taxonomy), functions (via metabolic pathways and proteins), and community organization (via network structure and ecological assembly) to deeply characterize response mechanisms. This revealed unexpected and comprehensive microbiome stability. This resistance to change appeared bolstered by habitat-specific dispersal and community-level functional redundancy, particularly via versatile carbon generalists. Our findings challenge widespread assumptions of significant soil microbiome alteration with climate change, and reveal key genome-inferred processes that underlie microbiome stability. This suggests that future research should reorient towards critical habitat transitions.

Main Text

Introduction

Arctic permafrost peatlands, warming at a rate four times faster than the global average (*1*), contain globally significant reserves of soil organic carbon (*2*). This carbon's fate and flux dynamics as carbon dioxide (CO_2) or methane (CH_4) will determine these systems' feedbacks to climate change (*3–5*). Soil microbiota play a critical role in controlling these dynamics (*6*). While rising Arctic temperatures trigger cascading changes that alter microbial community structure, assembly, and metabolism (*7–9*), underlying mechanisms of microbial change remain elusive (*10*). Resolving whether gradual or concerted change dominates, and at what scale, is vital for establishing how climate models should represent these critical ecosystems (*11*, *12*). Furthermore, modeling microbial responses to climate change remains a highly consequential, active area of research (*13*).

One prevailing assumption is that climate change will significantly alter belowground microbiome community and structure, yet previous studies reveal that microbial responses are dependent on disturbance intensity, timeframe, and ecological conditions. This variation in response could be driven by differences in study design and disturbance intensity (observation vs. manipulation (*14*)), timeframe (few- vs. many-year (*15*, *16*); see supplemental textS1), and environmental setting (with the impact of weather, soil properties and depth, plant communities, and hydrology potentially outweighing climate warming impacts (*17–22*)). These differences may mask a broad pattern of gradual microbiome response to climate change or alternatively, may indicate unexpected microbiome stability, with infrequent observations of concerted change. Stability may arise from community resistance or resilience (supplemental textS2), potentially driven by functional stability and metabolic flexibility (*23*, *24*). This stability, in turn, could create a pattern of concerted rather than gradual change, with potential for profoundly different outcomes in climate-critical systems.

A landscape in transition

Here we address the consequences of permafrost thaw on belowground microbiomes at Stordalen Mire, a discontinuous permafrost peatland in northern Sweden (68° 21' N, 19° 20' E; supplemental textS3) that has been intensively studied over decades. As permafrost thaws, dry elevated palsas collapse to form partially thawed bogs and then fully thawed fens, whose extent has doubled over the past 50 years (*5*). Thaw changes redox conditions from predominantly oxic to anoxic, significantly altering plant, microbial, and viral community compositions (*5*, *7*, *9*, *25–27*) and increasing anaerobic microbial lineages and metabolisms, including methanogenesis (*9*, *26*). While thaw-driven habitat conversion in Arctic peatlands is highly visible and its microbial impacts are well documented (*5*, *7*, *9*, *25–27*), it is unknown if rising temperatures also induce gradual within-habitat changes in microbiome composition, structure, and function or if microbiomes are broadly stable outside of concerted habitat transitions.

To address this gap, we quantified community change in Stordalen's habitats through three genome-resolved lenses: *organisms* (across taxon ranks ranging from phylum to strain), *functions* (encoded metabolic pathways and proteins), and community *organization* (community network structure and mechanisms of ecological assembly). Our dataset of 199 metagenomes

(2.37 Tbp) and 66 metatranscriptomes (1.87 Tbp) captures seven years of progressive
~0.7°C/yr belowground warming for each habitat (2011-2017, Fig1A) at peak growing season in
palsas, bogs, and fens (figS1). This period included significant interannual fluctuations in air
temperature and water table depth (figS2) and encompassed four of the planet's hottest years
(28). Given the strength of these study site fluctuations, due to widespread assumptions in the
literature we expected that warming would induce gradual year-over-year changes detectable
through all three lenses within each habitat. Contrary to our expectation, we consistently
observed within-habitat microbiome stability and resistance to climatic changes across taxon
ranks, metabolic functions, and community organization.
95
Microbiomes are resistant to increasing ground temperature
Microbiome composition and function resisted change over time
To evaluate microbiome changes, we reconstructed 13,290 metagenome-assembled
genomes (MAGs) that represent 1,806 species-level taxa (i.e., clustered at 95% average
nucleotide identity, Fig1A). Together these MAGs captured 73% of the community sequencing
reads at genus-level [see Methods], which is high for natural systems. Read-mapping-based
abundances revealed that microbial diversity and composition were resistant to change over
seven years within habitat and depth (down to 39 cm) (Fig1A and Fig1B; tableS1). Very little
variation in community composition could be accounted for by sample year at any level of
resolution (strain: ≤3.0%, p = 0.03; genus: 1.6%, p = 0.03; phylum: 1.6%, p = 0.005) evaluated
by distance-based redundancy analyses on single-copy marker genes (figS3; figS4). Similarly,
in 75% of habitats and depths, genome-resolved community composition was not explained by
time (p-value range: 0.22-0.79 Fig1B and Fig1C; figS5; figS6) and interannual fluctuation was
significant only at certain depths in the fen (for 0-9, 20-29, and 30-39 cm, respectively: R^2: 0.35,
0.62, 0.81; p-value: 0.036, 0.048, 0.048; figS7). Only the deeper fen (30-39 cm) samples
exhibited both significant non-directional inter-annual fluctuation (figS7) and directional change
(R^2 = 0.50, p = 0.012) in community composition with time (Fig1C), and only 8 of 1,806 species
were significantly correlated to time (Spearman rho ranging from -0.7 to 0.91, p < 0.05 figS8).
The stability observed across a range of taxonomic granularities suggests that the microbial
community is compositionally resistant over time and the rebounding of the fen community to its
original state after 2013 suggests resilience (see supplemental textS4).
117
Encoded microbiome functions were similarly consistent within habitats over time
(Fig1B&C). Beta diversity of community-wide protein clusters (a metric unconstrained by
functional annotation), as well as manually-curated genome-resolved pathways of carbon,
nitrogen, and sulfur metabolism did not significantly change over time within 92% of habitats
and depths (Fig1C; figS9); the sole exception was 30-39 cm in the fen (R^2: 0.49-0.55,
PERMANOVA, p < 0.05 Fig1B and Fig1C; figS9; figS10). At the finest level of functional
annotation (MAG-specific specialization), individual functional guilds of organisms (e.g.,
methanogens, homoacetogens, fermenters, and macromolecule degraders) maintained similar
abundances through time (figS11). (Expressed microbiome function was not used for temporal
analysis due to discontinuity in the metatranscriptomic time series.) Thus, within the majority of
Stordalen Mire habitats and depths, we observed both compositional and functional resistance
to change over time across multiple levels of granularity.

Our third lens, microbial community organization, was also stable over time within
habitat and depth. We used two distinct genome-resolved network approaches to understand
the influence of soil biogeochemical conditions (weighted genome correlation network analysis
(WGCNA)) and inter-microbe interactions (multilayer temporal networks (MTN)) on the
organizational structure of the community with time. Habitat-specific WGCNA modules showed
no linear correlation of community composition to time (Pearson, p<0.05) (table S3). MTN
microbial co-occurrence networks indicated variability in specific organisms over time, yet the
overall community network was resistant to change across different resolutions of network
structure (Fig1C, Fig2B, figS12) ranging from coarse (whole network degree distribution) to fine
(Jaccard similarity of triangles), suggesting that keystone taxa are stable (*29*). To understand if
subcommunities of microbes were resilient over time, we calculated the temporal persistence of
MTN subcommunities by categorizing each by the number of years it persisted (figS13).
Persistent subcommunities (occurring in ≥ four of the seven years) accounted for a conservative
estimate of 40% of the abundance-weighted community with time (depending on the interannual
relax rate; figS14) indicating that while not all microbe-microbe interactions were stable from
year to year, persistent clusters contributed extensively to community composition.

Finally, we sought to understand what ecological mechanisms may underlie the
apparent stability in community organization and to ask if they were changing over time. We
used beta-null modeling (*30*) to determine whether stochastic (dispersal and ecological drift) or
deterministic (abiotic or biotic selection) processes drove community structure (figS15; figS16).
A change in community assembly mechanisms over time could indicate a shift in the ecological
pressures shaping the community (for example, increased selection due to changing ground
temperature). Instead, beta-null modeling showed that while each habitat was governed by
distinct assembly mechanisms (homogenizing selection in the palsa, homogenizing dispersal in
the water-logged habitats), the ratio between deterministic and stochastic assembly processes
within each habitat did not change with time (Mantel, p > 0.05) (Fig1B and Fig2C).

Microbiome composition and function had weak susceptibility to temperature
Finally, we explored the possibility that organisms, functions, and organization of the
communities were being shaped by subtle or non-monotonic changes in abiotic drivers. Below
we focus on temperature; see figS17 for other abiotic factors. After accounting for covariation
between temperature and depth (strongest in palsa; figS18), genome-resolved community
composition was slightly, but significantly, linearly correlated to ground temperature in bog and
fen (Fig 1C, Bray-Curtis dissimilarity: partial mantel rho = 0.152 and 0.24, p < 0.001,
respectively; tableS2) but not palsa. However, non-linear generalized additive models revealed
that inclusion of habitat and depth removed any predictive relationship between community
composition and temperature (p = 0.4; supplemental textS5), suggesting that the effect of
temperature was weak. Ground temperature had no significant effect on carbon, nitrogen, or
sulfur pathways in the palsa or bog, and on only xyloglucan degradation in the fen
(supplemental textS5). While some WGCNA modules were significantly (Pearson, p < 0.05)
correlated with temperature, all but one of these modules (figS19) were more strongly correlated
with depth, echoing results from the models of community composition. Instead, WGCNA

modules were much more strongly correlated to carbon cycling metrics (e.g., $\delta^{13}C_{CO2}$, $\delta^{13}C_{CH4}$,

$\alpha_{CO2\text{-}CH4}$ (figS20; figS21)), consistent with previous work at the site (*9*, *26*, *27*, *31*, *32*), and other

features (e.g., pH and total nitrogen; figS22). No community assembly mechanisms were

related to ground temperature or air temperature (day-of-sampling or prior four weeks) (Mantel,

$p > 0.05$), implying that even with lag effects, temperature was not a strong selective or

stochastic force on community formation. As with temporal change, microbiome sensitivity to

temperature was strongest in the fen but weak overall and overshadowed by other abiotic

factors (depth, pH, and nutrient availability).

Metabolic connections maintain stability

Carbon generalists are highly prevalent, and macromolecule degradation is widespread

Microbiome capacity to degrade available organic matter determines the rate and form

of peat carbon loss to the atmosphere; therefore we sought to understand the prevalence and

ecological role of these organisms. Carbon generalists, which we defined as MAGs encoding ≥

3 alternative CAZyme pathways and ≥ 3 sugar degradation pathways (supplemental textS6;

figS23; figS24), were >80% of the community and taxonomically diverse across all three thaw

stages and nearly all depths. The prevalence of these metabolic generalists produced

community-level functional redundancy (figS37), with each CAZyme pathway encoded by a

median of 531 carbon generalists, and each sugar degradation pathway by a median of 253.

Macromolecule and sugar degradation dominated encoded and expressed pathways,

particularly in the palsa and bog where >84% of the abundance-weighted community encoded

and >50% expressed these pathways (Fig3A; Fig3C; figS9; figS25; figS26). Fermentation

pathways were also highly abundant (47-93% cumulative relative abundance) (Fig3A; figS9).

These sorts of acquisition-oriented carbon generalists are typical of cold peatlands and other

wetlands (*24*, *33*, *34*) and we hypothesize that these generalists confer resilience. As thaw

alters organic matter profiles and shifts redox conditions (*35*), functionally redundant carbon

generalists enable the community to continue exploiting changing carbon sources, yielding more

consistent downstream metabolites (e.g. for methanogenesis) even as environmental conditions

shift.

Chitin degradation and nitrogen fixation offer alternative pathways to nitrogen acquisition

Access to nitrogen is a fundamental constraint on microbial growth in northern peatlands

(figS27), where nitrogen is frequently limiting (*36*, *37*). We found organisms extensively used

two complementary nitrogen-acquisition pathways—nitrogen fixation and chitin degradation—

that may buffer a lack of preferred low-molecular-weight organic nitrogen compounds (*38*, *39*).

Nitrogen fixation increases the bioavailable nitrogen pool, benefiting diazotrophs directly and the

community indirectly (*40*). Up to 25% of the community encoded nitrogen fixation (compared to

~0.3% in marine microbiomes (*41*)), and > 75% of encoding MAGs showed pathway expression

(figS28).

Pathways for degrading chitin ($(C_8H_{13}O_5N)_n$), the second most abundant biopolymer in

nature (*42*), were even more widespread. Chitin is insoluble so mobilization of this nitrogen pool

requires extracellular enzymes and many degradation pathways exist (figS29). At Stordalen

Mire, multiple chitin degradation pathways were ubiquitously encoded (abundance weighted

median 73%, range 45%–94%) and expressed (median 95% of encoding cells, range 59–100%
with one outlier sample at 5% (figS29)). Furthermore, organisms containing these functions
were closely linked to key environmental outputs (as Variable Importance in Projection (VIP)
members of WGCNA modules related to carbon and nitrogen cycling (figS20; figS21; figS22)).
In many aquatic and terrestrial environments, including peatlands, extracellular chitin
degradation products can facilitate commensal and cross-feeding interactions (*42–45*), which
may promote community stability (*46*) while buffering against fluctuations in low-molecular-
weight nitrogen availability. The ubiquity of both nitrogen fixation and chitin degradation
pathways, may confer resilience to nutrient fluctuations in these habitats.
227
228   *Active sulfur cycling and sulfur-containing polysaccharide degradation might be a critical control*
229   *on organic matter degradation*
230   Despite small environmental pools of sulfur species (*32, 47*), sulfur cycling can control
organic matter degradation rates, particularly in the bog (*48*). We observed a range of sulfur
transformations across the thaw gradient (Fig3A). Habitat-wide expression of dissimilatory sulfur
oxidation and sulfate reduction (encoded pathway relative abundances ~25% and up to 10%,
respectively; figS30; figS31) suggest the presence of cryptic sulfur cycling, which can profoundly
influence system biogeochemistry (*49*). Interestingly, although all microbes require sulfur for
growth, we detected assimilatory sulfate reduction in only 25% (fen) to 50% (palsa, bog) of the
population (figS32). Our analysis of carbohydrate degradation suggests a novel hypothesis: that
community members may instead meet their sulfur requirements through CAZyme degradation
of sulfur-containing polysaccharides (figS33), known to be preferentially consumed from plant
organic matter (*32*). This novel linkage between the carbon and sulfur cycles in permafrost
peatlands is analogous to some marine isolates' ability to degrade sulfur polysaccharides and
thereby increase H_2, acetate, and sulfide concentrations (*50, 51*).
243
244   *The flexible community of carbon generalists supports methanogenesis in the bog and fen*
245   Despite being markedly less metabolically flexible than carbon generalists (supplemental
textS6), several methanogens persisted through seven years of thaw-induced change
(Fig3C&D; supplemental textS7). Consistent with previous reports (*9*), methanogenesis—
primarily hydrogenotrophic—was encoded by a handful of MAGs (< 1%) in the bog and fen and
was highly expressed there (Fig3A). Acetoclastic and methylotrophic methanogenesis were
uncommon, with the former present only in the fen (Fig3A; figS25). Despite low abundances,
methanogens have an outsized impact on ecosystem output. For example, the
hydrogenotrophic methanogen *Ca.* 'Methanoflorens' spp. is a strong predictor of $\alpha_{CO2-CH4}$ (*9, 26*),
which is an isotopic indicator of CH_4 metabolic pathways. Strikingly, WGCNA not only confirmed
this relationship but also identified thirteen non-methanogens (including multiple
Acidimicrobiales, *Terracidiphilus* and *Ca.* Acidiflorens) that were highly predictive of (VIP > 1.5)
and correlated to $\alpha_{CO2-CH4}$ (Spearman's rho, p < 0.005). These correlations were comparable to
or higher than that of *Ca.* 'M. stordalenmirensis' itself (Fig3C), suggesting that these
predominantly carbon-generalist, nitrogen-flexible lineages may be gatekeepers for
methanogenic carbon flux. For example, several of the *Ca*. 'Acidiflorens' spp. may supply H_2
and formate to *Ca.* 'M. stordalenmirensis' (*9*). Further, the MTN analysis showed *Ca.* 'M.
stordalenmirensis' is persistently associated with eight MAGs—all VIPs for [CO_2], $\delta^{13}C_{CO2}$, α_{CO2-}

$_{CH4}$ and/or nitrogen content—from the same WGCNA module (Fig3D). These MAGs included
the Dormibacterota *Ca*. 'Changshengia' spp., which was previously identified as highly
correlated with the fraction of carbon mineralized to CO_2 vs. CH_4 in the bog (*9*). These diverse
metabolic partnerships may increase stability by reducing substrate limitation for methanogens,
as was seen in aquatic microbial communities (*52, 53*).
267
From close analysis of the metabolic capabilities of methanogen-associated lineages,
we propose a novel hypothesis that trading and acquisition of both nitrogen and carbon
underlies these relationships. We observed that highly connected methanogens important to
network structure (e.g., *Ca*. 'M. stordalenmirensis') typically lacked recognizable nitrogen-
fixation or other nitrogen-linked decomposition pathways (supplemental textS7). Conversely,
methanogens encoding and expressing nitrogen-fixation have fewer connections and appear to
be more independent (see supplemental textS7; figS34; figS35; figS36). In other systems,
methanogenesis expression (*54*) and CO_2 and CH_4 production increase with addition of chitin,
which methanogens do not themselves degrade (*44, 55, 56*). Here, all eight persistent
methanogen partners (Fig3D) encode and express chitin degradation. Furthermore, we identify
a potential new linkage between methanogens and their partners through the reductive glycine
pathway. This energy-conserving and amine-recycling carbon-fixation pathway, capable of
producing methanogenesis substrates (pyruvate, acetate, and formate) and bioavailable
nitrogen (as ammonia) (*57, 58*), is encoded and expressed by five of the eight methanogen
partners. These observations suggest a previously unrecognized role for nitrogen in metabolic
interactions supporting methanogen persistence and activity in these ecosystems.
284
Habitat-specific microbiome resistance arises from functional redundancy and
dispersal mediated by hydrologic connectivity
Stordalen Mire is warming, causing rapid permafrost thaw and habitat conversion (*5*), yet
we observed within-habitat stability in microbiomes' organization, organisms, and functions.
What ecological dynamics maintain the community's resistance and resilience to change
(textS2)? We hypothesize that functional redundancy and homogenizing dispersal drive
ecological resistance (Fig4).
292
Functional redundancy and functional similarity have been proposed to stabilize
microbial communities (*59*). We saw strong signals of functional redundancy at both the
community and the individual level: communities across the thaw gradient were dominated by
diverse taxa encoding multiple alternative pathways for carbon and nitrogen acquisition (Fig3C;
supplemental textS6; figS37; figS38; figS39). It is unclear if such abundance of versatile
resource generalists is unusual or typical (*60*) (supplemental textS6). We argue that a "portfolio
effect" (*61, 62*) should operate at the individual as well as community level to smooth ecosystem
outputs: under changing conditions, just as one lineage's decrease in efficiency may be
compensated for by another's increase, maintaining community function, so can a metabolically
versatile cell (i.e. with a broad portfolio of carbon/nitrogen-handling pathways) redirect metabolic
flux under different conditions to maintain activity. Increased access via mobile genetic elements
to genes providing metabolic versatility would further backstop these stabilizing forces (see
companion paper by Guo et al.). If redundant pathways within redundant populations stabilize

communities, more narrowly distributed specialists (*24*) would be shielded from disturbance and
the loss of taxa from critical functional groups.

Bog and fen microbiomes are likely further stabilized by homogenizing dispersal, i.e.,
dispersal high enough to result in a community that is more similar to a neighbor than would be
expected by chance (Fig2c; Fig4; figS15; figS16). Dispersal dynamics are particularly important
during disturbance, when dispersing organisms can determine ecosystem trajectory (*63*). Local
dispersal can promote community resistance and resilience by "rescuing" disturbed
communities, preventing cross-site divergence after local population reductions or extinctions
(*23*, *64*). This is likely synergistic with functional redundancy and metabolic versatility, which
improve the odds that dispersal will resupply organisms that fill key roles.

Conclusions

Ecological factors reinforce habitat resistance and resilience to gradual change
(supplemental textS2), but enough disturbance can push communities into non-equilibrium
dynamics and reassembly into a new community state (*64–66*). At Stordalen Mire, the surficial
plant community reassembly makes habitat conversions detectable from space (Varner et al,
co-submitted), but the dynamics of belowground change are unknown and not easy to remotely
observe. Such knowledge gaps are a challenge in eco- and earth system climate models, which
must calibrate the representation of microbial complexity (*11*) often in the absence of high-
resolution datasets and parameter estimations (*13*, *67*). The analyses presented here
demonstrate that, despite warming and landscape-scale change (*5*), habitat-specific microbiome
complexity can be distilled to essential functions that are stable during peak growing season on
a near-decadal timescale—supporting their representation in models.

Collectively, our findings challenge prevailing assumptions about microbial responses to
climate change in critical permafrost ecosystems and offer new insights into processes
underlying community stability. Our novel, integrated analysis over multiple levels of granularity
for organismal ranks, metabolic functions, and community organization revealed little to no
significant temporal change over seven years of progressive belowground warming. This
analysis establishes a collection of metrics for assessing community response to environmental
disturbance and a new standard of evidence that should be met before soil microbiome change
is invoked. It also highlights that shorter-term or large disturbance experiments may incorrectly
interpret fluctuating microbiome composition as longer-term temporal change. Critically,
belowground microbial communities might be more resistant and resilient to temperature-
induced climate change than previously thought, likely due to synergy between functional
redundancy within communities and dispersal-mediated resupply of organisms encoding
essential metabolic versatility.

Given that the three habitats' microbiomes were stable and highly distinct over time—
and across the landscape (Varner et al, co-submitted)—future sampling must target periods of
habitat transition (figS2) when stabilizing capacity is exceeded. Further detailed study at spatial,
temporal, and genetic resolutions not captured in this study may reveal other mechanisms of
microbial response to changing conditions. For instance, spring and fall observations capturing

changing phenology may illuminate an erosion of stability driven by aboveground/belowground asynchronies. In addition, increased application of long-read sequencing may help resolve strain-level functional differences, including those mediated by the activity of mobile genetic elements. Despite these remaining unknowns, this work reveals the clear existence of stable habitats in a rapidly warming unstable Arctic landscape.

References and Notes

1. M. Rantanen, A. Y. Karpechko, A. Lipponen, K. Nordling, O. Hyvärinen, K. Ruosteenoja, T. Vihma, A. Laaksonen, The Arctic has warmed nearly four times faster than the globe since 1979. *Commun. Earth Environ.* **3**, 1–10 (2022).

2. E. A. G. Schuur, B. W. Abbott, R. Commane, J. Ernakovich, E. Euskirchen, G. Hugelius, G. Grosse, M. Jones, C. Koven, V. Leshyk, D. Lawrence, M. M. Loranty, M. Mauritz, D. Olefeldt, S. Natali, H. Rodenhizer, V. Salmon, C. Schädel, J. Strauss, C. Treat, M. Turetsky, Permafrost and Climate Change: Carbon Cycle Feedbacks From the Warming Arctic. *Annu. Rev. Environ. Resour.* **47**, 343–371 (2022).

3. G. Hugelius, J. Strauss, S. Zubrzycki, J. W. Harden, E. a. G. Schuur, C.-L. Ping, L. Schirrmeister, G. Grosse, G. J. Michaelson, C. D. Koven, J. A. O'Donnell, B. Elberling, U. Mishra, P. Camill, Z. Yu, J. Palmtag, P. Kuhry, Estimated stocks of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps. *Biogeosciences* **11**, 6573–6593 (2014).

4. M. A. Kuhn, R. K. Varner, C. K. McCalley, C. R. Perryman, M. Aurela, S. A. Burke, J. P. Chanton, P. M. Crill, J. DelGreco, J. Deng, L. Heffernan, C. Herrick, S. B. Hodgkins, C. P. Jones, S. Juutinen, E. S. Kane, L. J. Lamit, T. Larmola, E. Lilleskov, D. Olefeldt, M. W. Palace, V. I. Rich, C. Schulze, J. H. Shorter, F. B. Sullivan, O. Sonnentag, M. R. Turetsky, M. P. Waldrop, Controls on Stable Methane Isotope Values in Northern Peatlands and Potential Shifts in Values Under Permafrost Thaw Scenarios. *J. Geophys. Res. Biogeosciences* **129**, e2023JG007837 (2024).

5. R. K. Varner, P. M. Crill, S. Frolking, C. K. McCalley, S. A. Burke, J. P. Chanton, M. E. Holmes, Isogenie Project Coordinators, S. Saleska, M. W. Palace, Permafrost thaw driven changes in hydrology and vegetation cover increase trace gas emissions and climate forcing in Stordalen Mire from 1970 to 2014. *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.* **380**, 20210022 (2021).

6. J. K. Jansson, K. S. Hofmockel, Soil microbiomes and climate change. *Nat. Rev. Microbiol.* **18**, 35–46 (2020).

7. M. Hough, S. McCabe, S. R. Vining, E. Pickering Pedersen, R. M. Wilson, R. Lawrence, K.-Y. Chang, G. Bohrer, T. I. Coordinators, W. J. Riley, P. M. Crill, R. K. Varner, S. J. Blazewicz, E. Dorrepaal, M. M. Tfaily, S. R. Saleska, V. I. Rich, Coupling plant litter quantity to a novel metric for litter quality explains C storage changes in a thawing permafrost peatland. *Glob. Change Biol.* **28**, 950–968 (2022).

8.   A. Persson, A. Hasan, J. Tang, P. Pilesjö, Modelling Flow Routing in Permafrost
Landscapes with TWI: An Evaluation against Site-Specific Wetness Measurements. *Trans.*
GIS **16**, 701–713 (2012).

9.   B. J. Woodcroft, C. M. Singleton, J. A. Boyd, P. N. Evans, J. B. Emerson, A. A. F. Zayed,
R. D. Hoelzle, T. O. Lamberton, C. K. McCalley, S. B. Hodgkins, R. M. Wilson, S. O.
Purvine, C. D. Nicora, C. Li, S. Frolking, J. P. Chanton, P. M. Crill, S. R. Saleska, V. I.
Rich, G. W. Tyson, Genome-centric view of carbon processing in thawing permafrost.
Nature **560**, 49–54 (2018).

10. J. G. Ernakovich, R. A. Barbato, V. I. Rich, C. Schädel, R. E. Hewitt, S. J. Doherty, E. D.
Whalen, B. W. Abbott, J. Barta, C. Biasi, C. L. Chabot, J. Hultman, C. Knoblauch, M. C. Y.
L. Vetter, M. Leewis, S. Liebner, R. Mackelprang, T. C. Onstott, A. Richter, U. M. E.
Schütte, H. M. P. Siljanen, N. Taş, I. Timling, T. A. Vishnivetskaya, M. P. Waldrop, M.
Winkel, Microbiome assembly in thawing permafrost and its feedbacks to climate. *Glob.*
Change Biol., 5007–5026 (2022).

11. E. Kyker-Snowman, D. L. Lombardozzi, G. B. Bonan, S. J. Cheng, J. S. Dukes, S. D. Frey,
E. M. Jacobs, R. McNellis, J. M. Rady, N. G. Smith, R. Q. Thomas, W. R. Wieder, A. S.
Grandy, Increasing the spatial and temporal impact of ecological research: A roadmap for
integrating a novel terrestrial process into an Earth system model. *Glob. Change Biol.* **28**,
665–684 (2022).

12. W. R. Wieder, G. B. Bonan, S. D. Allison, Global soil carbon projections are improved by
modelling microbial processes. *Nat. Clim. Change* **3**, 909–912 (2013).

13. W. R. Wieder, S. D. Allison, E. A. Davidson, K. Georgiou, O. Hararuk, Y. He, F. Hopkins,
Y. Luo, M. J. Smith, B. Sulman, K. Todd-Brown, Y. Wang, J. Xia, X. Xu, Explicitly
representing soil microbial processes in Earth system models. *Glob. Biogeochem. Cycles*
29, 1782–1800 (2015).

14. L. Wu, Y. Zhang, X. Guo, D. Ning, X. Zhou, J. Feng, M. M. Yuan, S. Liu, J. Guo, Z. Gao, J.
Ma, J. Kuang, S. Jian, S. Han, Z. Yang, Y. Ouyang, Y. Fu, N. Xiao, X. Liu, L. Wu, A. Zhou,
Y. Yang, J. M. Tiedje, J. Zhou, Reduction of microbial diversity in grassland soil is driven
by long-term climate warming. *Nat. Microbiol.* **7**, 1054–1062 (2022).

15. J. Feng, C. Wang, J. Lei, Y. Yang, Q. Yan, X. Zhou, X. Tao, D. Ning, M. M. Yuan, Y. Qin,
Z. J. Shi, X. Guo, Z. He, J. D. Van Nostrand, L. Wu, R. G. Bracho-Garillo, C. R. Penton, J.
R. Cole, K. T. Konstantinidis, Y. Luo, E. A. G. Schuur, J. M. Tiedje, J. Zhou, Warming-
induced permafrost thaw exacerbates tundra soil carbon decomposition mediated by
microbial community. *Microbiome* **8**, 3 (2020).

16. K. Xue, M. M. Yuan, Z. J. Shi, Y. Qin, Y. Deng, L. Cheng, L. Wu, Z. He, J. D. Van
Nostrand, R. Bracho, S. Natali, E. A. G. Schuur, C. Luo, K. T. Konstantinidis, Q. Wang, J.
R. Cole, J. M. Tiedje, Y. Luo, J. Zhou, Tundra soil carbon is vulnerable to rapid microbial
decomposition under climate warming. *Nat. Clim. Change* **6**, 595–600 (2016).

17. J. Curiel Yuste, A. J. Fernandez-Gonzalez, M. Fernandez-Lopez, R. Ogaya, J. Penuelas,
J. Sardans, F. Lloret, Strong functional stability of soil microbial communities under
semiarid Mediterranean conditions and subjected to long-term shifts in baseline
precipitation. *Soil Biol. Biochem.* **69**, 223–233 (2014).

18. K. M. DeAngelis, G. Pold, B. D. Topçuoğlu, L. T. A. van Diepen, R. M. Varney, J. L.
Blanchard, J. Melillo, S. D. Frey, Long-term forest soil warming alters microbial
communities in temperate forest soils. *Front. Microbiol.* **6** (2015).

19. N. C. Dove, M. E. Barnes, K. Moreland, R. C. Graham, A. A. Berhe, S. C. Hart, Depth
dependence of climatic controls on soil microbial community activity and composition.
ISME Commun. **1**, 1–11 (2021).

20. J. L. M. Gutknecht, C. B. Field, T. C. Balser, Microbial communities and their responses to
simulated global change fluctuate greatly over multiple years. *Glob. Change Biol.* **18**,
2256–2269 (2012).

21. S. Monteux, J. T. Weedon, G. Blume-Werry, K. Gavazov, V. E. J. Jassey, M. Johansson,
F. Keuper, C. Olid, E. Dorrepaal, Long-term in situ permafrost thaw effects on bacterial
communities and potential aerobic respiration. *ISME J.* **12**, 2129–2141 (2018).

22. K. Zhu, N. R. Chiariello, T. Tobeck, T. Fukami, C. B. Field, Nonlinear, interacting
responses to climate limit grassland production under global change. *Proc. Natl. Acad. Sci.*
113, 10589–10594 (2016).

23. F. De Vries, A. Shade, Controls on soil microbial community stability under climate change.
Front. Microbiol. **4** (2013).

24. J. T. Weedon, G. A. Kowalchuk, R. Aerts, S. Freriks, W. F. M. Röling, P. M. van Bodegom,
Compositional Stability of the Bacterial Community in a Climate-Sensitive Sub-Arctic
Peatland. *Front. Microbiol.* **8** (2017).

25. J. B. Emerson, S. Roux, J. R. Brum, B. Bolduc, B. J. Woodcroft, H. B. Jang, C. M.
Singleton, L. M. Solden, A. E. Naas, J. A. Boyd, S. B. Hodgkins, R. M. Wilson, G. Trubl, C.
Li, S. Frolking, P. B. Pope, K. C. Wrighton, P. M. Crill, J. P. Chanton, S. R. Saleska, G. W.
Tyson, V. I. Rich, M. B. Sullivan, Host-linked soil viral ecology along a permafrost thaw
gradient. *Nat. Microbiol.* **3**, 870–880 (2018).

26. C. K. McCalley, B. J. Woodcroft, S. B. Hodgkins, R. A. Wehr, E.-H. Kim, R. Mondav, P. M.
Crill, J. P. Chanton, V. I. Rich, G. W. Tyson, S. R. Saleska, Methane dynamics regulated
by microbial community response to permafrost thaw. *Nature* **514**, 478–481 (2014).

27. R. Mondav, C. K. McCalley, S. B. Hodgkins, S. Frolking, S. R. Saleska, V. I. Rich, J. P.
Chanton, P. M. Crill, Microbial network, phylogenetic diversity and community membership
in the active layer across a permafrost thaw gradient. *Environ. Microbiol.* **19**, 3201–3218
(2017).

28. NOAA National Centers for Environmental Information, "NOAA National Centers for
Environmental Information, Monthly Global Climate Report for Annual 2017" (Climate
Monitoring Branch, DOC/NOAA/NESDIS/NCDC/, National Climatic Data Center, NESDIS,
NOAA, U.S. Department of Commerce, 2018);
https://www.ncei.noaa.gov/access/monitoring/monthly-report/global/201713.

29. S. Banerjee, K. Schlaeppi, M. G. A. van der Heijden, Keystone taxa as drivers of
microbiome structure and functioning. *Nat. Rev. Microbiol.* **16**, 567–576 (2018).

30. J. C. Stegen, X. Lin, J. K. Fredrickson, X. Chen, D. W. Kennedy, C. J. Murray, M. L.
Rockhold, A. Konopka, Quantifying community assembly processes and identifying
features that impose them. *ISME J.* **7**, 2069–2079 (2013).

31. R. M. Wilson, M. M. Tfaily, M. Kolton, E. R. Johnston, C. Petro, C. A. Zalman, P. J.
Hanson, H. M. Heyman, J. E. Kyle, D. W. Hoyt, E. K. Eder, S. O. Purvine, R. K. Kolka, S.
D. Sebestyen, N. A. Griffiths, C. W. Schadt, J. K. Keller, S. D. Bridgham, J. P. Chanton, J.
E. Kostka, Soil metabolome response to whole-ecosystem warming at the Spruce and
Peatland Responses under Changing Environments experiment. *Proc. Natl. Acad. Sci.*
118, e2004192118 (2021).

32. R. M. Wilson, M. A. Hough, B. A. Verbeke, S. B. Hodgkins, G. Tyson, M. B. Sullivan, E.
Brodie, W. J. Riley, B. Woodcroft, C. McCalley, S. C. Dominguez, P. M. Crill, R. K. Varner,
S. Frolking, W. T. Cooper, J. P. Chanton, S. D. Saleska, V. I. Rich, M. M. Tfaily, Plant
organic matter inputs exert a strong control on soil organic matter decomposition in a
thawing permafrost peatland. *Sci. Total Environ.* **820**, 152757 (2022).

33. A. M. Oliverio, A. B. Narrowe, J. A. Villa, C. Rinke, D. W. Hoyt, P. Liu, B. B. McGivern, E.
K. Bechtold, J. B. Ellenbogen, R. A. Daly, G. J. Smith, J. C. Angle, R. M. Flynn, A. P.
Freiburger, K. B. Louie, B. Stemple, T. R. Northen, C. Henry, C. S. Miller, T. H. Morin, G.
Bohrer, M. A. Borton, K. C. Wrighton, Rendering the metabolic wiring powering wetland
soil methane production. [Preprint] (2024). https://doi.org/10.1101/2024.02.06.579222.

34. A. T. Tveit, T. Urich, P. Frenzel, M. M. Svenning, Metabolic and trophic interactions
modulate methane production by Arctic peat microbiota in response to warming. *Proc.*
Natl. Acad. Sci. **112**, E2507–E2516 (2015).

35. S. B. Hodgkins, "Changes in organic matter chemistry and methanogenesis due to
permafrost thaw in a subarctic peatland," thesis, The Florida State University, United
States -- Florida (2016).

36. E. Du, C. Terrer, A. F. A. Pellegrini, A. Ahlström, C. J. van Lissa, X. Zhao, N. Xia, X. Wu,
R. B. Jackson, Global patterns of terrestrial nitrogen and phosphorus limitation. *Nat.*
Geosci. **13**, 221–226 (2020).

37. J. J. Elser, M. E. S. Bracken, E. E. Cleland, D. S. Gruner, W. S. Harpole, H. Hillebrand, J.
T. Ngai, E. W. Seabloom, J. B. Shurin, J. E. Smith, Global analysis of nitrogen and
phosphorus limitation of primary producers in freshwater, marine and terrestrial
ecosystems. *Ecol. Lett.* **10**, 1135–1142 (2007).

38. E. B. Graham, F. Yang, S. Bell, K. S. Hofmockel, High Genetic Potential for Proteolytic
Decomposition in Northern Peatland Ecosystems. *Appl. Environ. Microbiol.* **85** (2019).

39. J. S. Norman, M. L. Friesen, Complex N acquisition by soil diazotrophs: how the ability to
release exoenzymes affects N fixation by terrestrial free-living diazotrophs. *ISME J.* **11**,
315–326 (2017).

40. B. A. Lerch, D. A. Smith, T. Koffel, S. C. Bagby, K. C. Abbott, How public can public goods
be? Environmental context shapes the evolutionary ecology of partially private goods.
PLOS Comput. Biol. **18**, e1010666 (2022).

41. T. O. Delmont, C. Quince, A. Shaiber, Ö. C. Esen, S. T. Lee, M. S. Rappé, S. L. McLellan,
S. Lücker, A. M. Eren, Nitrogen-fixing populations of Planctomycetes and Proteobacteria
are abundant in surface ocean metagenomes. *Nat. Microbiol.* **3**, 804–813 (2018).

42. S. Beier, S. Bertilsson, Bacterial chitin degradation—mechanisms and ecophysiological
strategies. *Front. Microbiol.* **4** (2013).

43. S. E. Belova, N. V. Ravin, T. A. Pankratov, A. L. Rakitin, A. A. Ivanova, A. V. Beletsky, A.
V. Mardanov, J. S. Sinninghe Damsté, S. N. Dedysh, Hydrolytic Capabilities as a Key to
Environmental Success: Chitinolytic and Cellulolytic Acidobacteria From Acidic Sub-arctic
Soils and Boreal Peatlands. *Front. Microbiol.* **9** (2018).

44. C. De Tender, B. Mesuere, F. Van der Jeugt, A. Haegeman, T. Ruttink, B. Vandecasteele,
P. Dawyndt, J. Debode, E. E. Kuramae, Peat substrate amended with chitin modulates the
N-cycle, siderophore and chitinase responses in the lettuce rhizobiome. *Sci. Rep.* **9**, 9890
(2019).

45. M. B. Kaczmarek, K. Struszczyk-Swita, X. Li, M. Szczęsna-Antczak, M. Daroch, Enzymatic
Modifications of Chitin, Chitosan, and Chitooligosaccharides. *Front. Bioeng. Biotechnol.* **7**
(2019).

46. W. Lopes, D. R. Amor, J. Gore, Cooperative growth in microbial communities is a driver of
multistability. *Nat. Commun.* **15**, 4709 (2024).

47. C. R. Perryman, C. K. McCalley, A. Malhotra, M. F. Fahnestock, N. N. Kashi, J. G. Bryce,
R. Giesler, R. K. Varner, Thaw Transitions and Redox Conditions Drive Methane Oxidation
in a Permafrost Peatland. *J. Geophys. Res. Biogeosciences* **125**, e2019JG005526 (2020).

48. V. Freire-Zapata, H. Holland-Moritz, D. R. Cronin, S. Aroney, D. A. Smith, R. M. Wilson, J.
G. Ernakovich, B. J. Woodcroft, S. C. Bagby, EMERGE 2012 Field Team, S. B. Hodgkins,
R. Mondav, EMERGE Biology Integration Coordinators, J. E. Cross, M. F. Fahnestock, R.
Ferriere, M. Ibba, S. R. Saleska, R. K. Varner, A. A. Zayed, V. I. Rich, M. B. Sullivan, J. C.
Stegen, M. M. Tfaily, Microbiome–metabolite linkages drive greenhouse gas dynamics
over a permafrost thaw gradient. *Nat. Microbiol.*, doi: 10.1038/s41564-024-01800-z (2024).

49. D. E. Canfield, F. J. Stewart, B. Thamdrup, L. De Brabandere, T. Dalsgaard, E. F. Delong,
N. P. Revsbech, O. Ulloa, A Cryptic Sulfur Cycle in Oxygen-Minimum–Zone Waters off the
Chilean Coast. *Science* **330**, 1375–1378 (2010).

50. W. Helbert, Marine Polysaccharide Sulfatases. *Front. Mar. Sci.* **4** (2017).

51. D. M. Van Vliet, S. Palakawong Na Ayudthaya, S. Diop, L. Villanueva, A. J. M. Stams, I.
Sánchez-Andrea, Anaerobic Degradation of Sulfated Polysaccharides by Two Novel
Kiritimatiellales Strains Isolated From Black Sea Sediment. *Front. Microbiol.* **10**, 253
(2019).

52. D. Ciccarese, G. Micali, B. Borer, C. Ruan, D. Or, D. R. Johnson, Rare and localized
events stabilize microbial community composition and patterns of spatial self-organization
in a fluctuating environment. *ISME J.* **16**, 1453–1463 (2022).

53. M. Embree, J. K. Liu, M. M. Al-Bassam, K. Zengler, Networks of energetic and metabolic

interactions define dynamics in microbial communities. *Proc. Natl. Acad. Sci.* **112**, 15450–
15455 (2015).

54.   A. A. Ivanova, C.-E. Wegner, Y. Kim, W. Liesack, S. N. Dedysh, Identification of microbial
populations driving biopolymer degradation in acidic peatlands by metatranscriptomic
analysis. *Mol. Ecol.* **25**, 4818–4835 (2016).

55.   H. Kang, C. Freeman, S. Soon park, J. Chun, N-Acetylglucosaminidase activities in
wetlands: a global survey. *Hydrobiologia* **532**, 103–110 (2005).

56.   S. Wörner, M. Pester, Microbial Succession of Anaerobic Chitin Degradation in Freshwater
Sediments. *Appl. Environ. Microbiol.* **85**, e00963-19 (2019).

57.   P. N. Evans, J. A. Boyd, A. O. Leu, B. J. Woodcroft, D. H. Parks, P. Hugenholtz, G. W.
Tyson, An evolving view of methane metabolism in the Archaea. *Nat. Rev. Microbiol.* **17**,
219–232 (2019).

58.   I. Sánchez-Andrea, I. A. Guedes, B. Hornung, S. Boeren, C. E. Lawson, D. Z. Sousa, A.
Bar-Even, N. J. Claassens, A. J. M. Stams, The reductive glycine pathway allows
autotrophic growth of Desulfovibrio desulfuricans. *Nat. Commun.* **11**, 5090 (2020).

59.   S. D. Allison, J. B. H. Martiny, Resistance, resilience, and redundancy in microbial
communities. *Proc. Natl. Acad. Sci.* **105**, 11512–11519 (2008).

60.   T. H. Bell, T. Bell, Many roads to bacterial generalism. *FEMS Microbiol. Ecol.* **97**, fiaa240
(2020).

61.   D. F. Doak, D. Bigger, E. K. Harding, M. A. Marvier, R. E. O'Malley, D. Thomson, The
Statistical Inevitability of Stability-Diversity Relationships in Community Ecology. *Am. Nat.*
151, 264–276 (1998).

62.   L. M. Thibaut, S. R. Connolly, Understanding diversity–stability relationships: towards a
unified model of portfolio effects. *Ecol. Lett.* **16**, 140–150 (2013).

63.   J. HilleRisLambers, P. B. Adler, W. S. Harpole, J. M. Levine, M. M. Mayfield, Rethinking
Community Assembly through the Lens of Coexistence Theory. *Annu. Rev. Ecol. Evol.*
Syst. **43**, 227–248 (2012).

64.   A. Shade, H. Peter, S. Allison, D. Baho, M. Berga, H. Buergmann, D. Huber, S.
Langenheder, J. Lennon, J. Martiny, K. Matulich, T. Schmidt, J. Handelsman,
Fundamentals of Microbial Community Resistance and Resilience. *Front. Microbiol.* **3**
(2012).

65.   C. Chang, J. HilleRisLambers, Integrating succession and community assembly
perspectives. *F1000Research* **5**, 2294 (2016).

66.   A. Fofana, D. Anderson, C. K. McCalley, S. Hodgkins, R. M. Wilson, D. Cronin, N. Raab,
M. Torabi, R. K. Varner, P. Crill, S. R. Saleska, J. P. Chanton, M. M. Tfaily, V. I. Rich,
Mapping substrate use across a permafrost thaw gradient. *Soil Biol. Biochem.* **175**, 108809
(2022).

67.   K.-Y. Chang, W. J. Riley, E. L. Brodie, C. K. McCalley, P. M. Crill, R. F. Grant, Methane
Production Pathway Regulated Proximally by Substrate Availability and Distally by
Temperature in a High-Latitude Mire Complex. *J. Geophys. Res. Biogeosciences* **124**,
3057–3074 (2019).

Acknowledgments

Non-author contributions:

This study has been made possible by data provided by Abisko Scientific Research Station and the Swedish Infrastructure for Ecosystem Science (SITES) as well as the Swedish Polar Research Secretariat. We thank the Swedish Polar Research Secretariat and SITES for the support of the work done at the Abisko Scientific Research Station. A portion of this analysis and non-monetary technical support was also provided through the U.S. Department of Energy user facilities programs. Computational work was supported in part by an allocation of computing time from the Ohio Supercomputer Center. Lastly, we thank contracted science artist Elena Hartley for co-development of Figure 4.

Funding: We thank the many funding sources of this work, including:

- US National Science Foundation, Biology Integration Institutes Program, Award # 2022070 (VIR and RKV, with SH, MMT, GWT, BW, JE, SCB, and the EMERGE Institute Coordinators)
- Genomic Science Program of the United States Department of Energy Office of Biological and Environmental Research (BER), grants DE-SC0004632, DE-SC0010580 and DE-SC0016440 (SRS and VIR, with GWT, MMT). Associated with the third of these was a BER Support Science award (Proposal #503530 DOI: 10.46936/10.25585/60001148) for nucleic acid sequencing for a portion of these samples, conducted by the U.S. Department of Energy Joint Genome Institute (https://ror.org/04xm1d337), a DOE Office of Science User Facility supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231.
- US National Science Foundation MacroSystems Program, Award # NSF EF 1241037 (RV).
- US National Science Foundation Research for Undergraduates Program, the Northern Ecosystems Research for Undergraduates, Award # NSF REU EAR 1063037 (RV)
- Nucleic acid sequencing and and dissolved organic matter fourier transform ion cyclotron resonance mass spectrometry were partially supported by a Facilities Integrating Collaborations for User Science (FICUS) awards, DOI: 10.46936/fics.proj.2016.49521/60006018 and 503547, which used resources at the DOE Joint Genome Institute (https://ror.org/04xm1d337) and the Environmental Molecular Sciences Laboratory (https://ror.org/04rc0xn13), which are DOE Office of Science User Facilities. Both facilities are sponsored by the Office of Biological and Environmental Research and operated under Contract Nos. DE-AC02-05CH11231 (JGI) and DE-AC05-76RL01830 (EMSL).
- SITES is supported by the Swedish Research Council's grant 4.3-2021-00164

Author Contributions: DC, SA, DS, and HH-M all contributed equally to this work. Co-first authors have the right to list their name first in their CV. In addition, VR, SB, GT, BW, and JE are all co-senior authors and contributed equally to this work, order was paired with first author order and all may list their name last on their CV. Following the CRediT model, authors made the following contributions

660
Conceptualization - VR, CM, SH, RV, EMERGE Institute Coordinators, DC, HH-M, DS, SA, JE,
SB, BW, MT, PC
Data Curation - DC, HH-M, DS, SA, VR, BW, SH, FL, RV
Formal Analysis - DC, HH-M, DS, SA, CM, KW, MB, IsoGenie Field & Analytic Teams 2011-
2017
Funding acquisition - VR, JE, SB, BW, GT, KW, SH, RV, EMERGE Institute Coordinators, MT,
PC
Investigation - DC, HH-M, DS, SA, CM, SH, FL, RV, IsoGenie Field & Analytic Teams 2011-
2017, PC
Methodology Development - DC, HH-M, DS, SA, VR, PC
Project Administration - DC, HH-M, DS, SA, VR, JE, SB, BW, GT, KW, SH, RV, EMERGE
Institute Coordinators, PC
Resources Provision - VR, KW, SH, RV, EMERGE Institute Coordinators, PC
Software Programming - DC, HH-M, SA, KW, MB, SH
Supervision Oversight - VR, JE, KW, RV, MT
Validation Verification- SA, DC, HH-M
Visualization - DC, HH-M, DS, SA, VR, SB, KH
Writing - DC, HH-M, DS, SA, VR, JE, SB, BW, GT, CM, KW, MB, SH, FL, RV, IsoGenie Field &
Analytic Teams 2011-2017, EMERGE Institute Coordinators, KH, MT, PC
680
681
682   **Competing interests:** Authors declare that they have no competing interests.
683
Data and materials availability:
Sample Metadata Sheet v1.0.0, containing field sample information and geochemical
measurements are available on Zenodo at https://zenodo.org/records/12827096. Gas flux
measurements are available on Zenodo at https://doi.org/10.5281/zenodo.13377109 and
https://emerge-db.asc.ohio-state.edu/datasources/0138_DailyFlux_20122018. Temperature and
water table summary for lagged effects analysis are available on Zenodo at
https://doi.org/10.5281/zenodo.14189390. Gas flux summary for lagged effects analysis is
available on Zenodo at https://doi.org/10.5281/zenodo.14834760. Water table depth
measurements used for determining lagged effects are available on Zenodo at
https://doi.org/10.5281/zenodo.10420396. ANS temperature measurements used in
supplemental (figS2) were recorded at the Abisko Scientific Research Station (ANS), Sveriges
meteorologiska och hydrologiska institut (SMHI). (2024). Meteorological data from Abisko
Observatory (Abisko: station 188800), daily mean air temperatures, 2011-2017. Retrieved
[05/30/2024], from https://www.smhi.se/data/meteorologi/ladda-ner-meteorologiska-
observationer#stationid=188800.
699
All metagenomes and metatranscriptomes used in this study are available in NCBI under the
umbrella BioProject PRJNA888099. SRA accessions for the reads used for the interannual
microbiome comparisons are listed in SuppTable_temporal-paper_reads, and those used to
generate the MAGs are listed in the metadata table accompanying those MAGs

(https://doi.org/10.5281/zenodo.13901514). Metagenome-assembled genomes (MAGs) are
available in Zenodo at https://doi.org/10.5281/zenodo.13901514.

Figures

Figure 1: Stordalen Mire microbiomes resist temporal change. A) The three main thaw-

gradient habitats (palsas, bogs, and fens; colored throughout as brown, green and blue,

respectively) host microbiomes with distinct taxonomy (treemap with dereplicated MAGs as

rectangles, colored by phyla and sized proportional to relative abundance). Microbiome beta

diversity did not change significantly with time (middle panels); each point is one microbiome's

average Bray-Curtis dissimilarity to those at the same depth for the earliest baseline year (2011,

or the first year for which at least two depth-matched microbiomes were available). Black lines

are the running dissimilarity average. Reference-year microbiomes' within-year dissimilarities

are indicated by gray dots, with their ranges and averages extended as shaded gray bars and

dashed lines, respectively. Soil temperatures increased significantly with time (lines indicate

significant linear relationships, p<0.05, lowest panel). **B)** Habitat and depth structure

microbiome beta diversity data more strongly than year, observed in non-metric

multidimensional scaling (NMDS) ordinations of each sample's MAG-based taxonomic (left

panels) and functional (right panels) composition, colored by year (upper panel) or depth (lower

panel). Habitats are denoted by shape (palsa, square; bog, circle; fen, triangle) and demarcated

by ellipses (dashed-line ellipses reflect each habitat's multivariate normal distribution, while

shaded ellipses are the multivariate t-distribution). Functions are cumulative pathway

abundances: each complete pathway observed in a MAG was weighted by that MAG's relative

abundance, then all occurrences are summed for each pathway. **C)** Granular heatmap of

microbiome temporal change, in each habitat and depth. Change was assessed spanning

organisms (from phylum to strains diversity metrics, both single copy marker genes and MAGs),

functions (from pathways to protein clusters, both annotatable and non-annotatable), and

organization (both network structure and community assembly metrics of stochasticity and

determinism). The amount of variation explained by time (as Mantel R or R^2 depending on

statistical test appropriate for each data type; see Methods) is indicated with the intensity of the

shaded color; white asterisks denote significant correlations to time. Across all granularities

surveyed only deeper fen (30-39 cm) showed significant change over time.

Figure 2: Microbiome organization is consistent within habitats over time. A) Multilayer

temporal networks depict correlations between MAGs within annual networks for each habitat.

Each network layer corresponds to a specific year, with nodes representing MAGs. Black edges

represent significant (p <0.05) correlations (>0.3) between MAGs within the same habitat and

year, while node size reflects node degree (the number of connections for that node) within its

respective layer (year). **B)** Pairwise interyear correlations (Spearman's rho) in network node

degree (x-axis, base year; colored points, comparison years) within each habitat. Note that the

fen's recovery to a near-stable state after 2013 suggests strong resilience in this habitat. **C)**

Assembly processes detected within each habitat (rows) by βNTI and RCBC modeling in

interyear comparisons. For each year, columns show pairwise comparisons to all other years

(self-comparisons are obscured by black bars). Stacked bar colors denote the proportion of

different assembly processes detected in each pairwise comparison, revealing overall stability in

assembly within habitats over time.

Figure 3: Microbiome metabolisms across the thaw gradient are interlinked by carbon,

nitrogen, and sulfur cycling. A) Aggregated over the study period, key pathways in carbon,

nitrogen, and sulfur metabolism (top center), metagenomic (middle row) and metatranscriptomic

(bottom row) data show differences in microbial investment in methanotrophy and carbon

fixation, nitrogen fixation, and sulfur redox cycling. **B)** Functional composition of palsa, bog, and

fen communities. Treemap rectangles are MAGs colored by carbon metabolic specialization and

scaled by relative abundance. **C)** Metabolic capabilities of MAGs belonging to the weighted

genome correlation network analysis (WGCNA) module correlated with the isotopic fractionation

of carbon between carbon dioxide and methane carbon ($\alpha_{CO2\text{-}CH4}$) in the bog. Carbon and

nitrogen-related metapathways present in these MAGs. MAGs are colored by carbon metabolic

specialization (see panel B) and ordered by VIP score. **D)** Bog network structure of the

persistent subcommunity of MAGs connected to *Ca.* 'M. stordalenmirensis' in multi-layer

temporal network (MTN) modeling. MAGs were included if they maintained a connection to *Ca.*

'M. stordalenmirensis' for at least six years and had a VIP score > 1 and a correlation > 0.5 in at

least one metric. MAGs are colored by carbon metabolic specialization (see panel B). Edge

color represents the number of years MAGs were connected to *Ca.* 'M. stordalenmirensis'.

Subplots show the Spearman correlation coefficient (rho) of each MAG to key environmental

variables (teal) and their VIP value (pink).

Figure 4: Within-habitat communities resist change until increasing environmental pressures trigger a concerted transition along the thaw gradient. Community composition fluctuates but does not change steadily with time in each habitat. Population density increases in tandem with water saturation (*9*). Across the thaw gradient, ecological assembly shifts from predominantly deterministic processes in the palsa (community assembly, purple) to predominantly stochastic processes in the fen (orange), while the balance of functional specializations shifts from predominantly generalists (red) to a more metabolically diverse range of specialists. We hypothesize that functional redundancy, dispersal (bog, fen), and metabolic versatility (palsa, bog) drive within-habitat ecological resistance.

Supplemental Text

Supplemental Note 1: In-depth literature review on microbial community responses to
warming and climate forcings
Supplemental Note 2: Literature-based definitions of resistance, resilience and stability
Supplemental Note 3: Site description of the model permafrost peatland ecosystem,
Stordalen Mire
Supplemental Note 4: The peat of the fen at deeper depths are more variable, which
might explain why this is the sole depth and habitat that shows directional microbiome
change with time rather than just fluctuation.
Supplemental Note 5: Further analyses on stability with time and temperature
Supplemental Note 6: Carbon generalist genomes are significantly larger in size than
specialists, which supports ecological literature, linking genome size to metabolic flexibility and
functional redundancy
Supplemental Note 7: Methanogens have simplistic and specialized metabolic calls
oftentimes with limited to no recognized nitrogen capabilities, which leads to a larger number of
associations with other organisms.
Supplemental Note 8: Etymology.
875

Supplemental Figures
876

Supplemental Text

In order to observe significant change in the soil microbial community or structure, some studies have required large pressed and/or pulsed disturbances with experimental climate forcings (e.g. increased temperature, precipitation, N addition, etc.) (e.g. ref (*1*) and/or shorter time scales (<3 years) of observation (*2, 3*), however there are inconsistencies across the literature and longer-term (>5 years) studies. In a 12-year grassland study, it took 8 years for the effects of enhanced CO_2 exposure to significantly alter soil respiration rates and community function. These 8 years of data were not reported on in the main text of that study due in part to no functional genes involved in C degradation or N cycling being significantly correlated with soil CO_2 efflux rates, however the second phase of the experiment (years 9 to 12) showed significant correlations and were highlighted (*4*). In anaerobic microcosms from the Spruce and Peatland Responses Under Changing Environments (SPRUCE) site incubated at 6°C and 15°C going down to 2 m over 70 days, temperature did not significantly alter the relative abundance of the predominant taxa, alpha diversity metrics, or bacterial and archaeal 16S rRNA gene copies (*5*). At the Harvard Forest Long-Term Ecological Research (LTER) project, where soils were warmed 5°C above ambient for 5, 8, and 20 years, only at the 20-year site in the organic and mineral soil horizons had the community structure and Shannon H diversity been significantly altered (*6*).

Natural variation and variability in environmental conditions and the microbial community can outweigh observed perturbations due to climate change. This suggests that communities may maintain some baseline amount of natural functional stability and metabolic flexibility that can be on par with the magnitudes required to respond to climate-change induced perturbation. Soil depth and pH can be two of the largest effectors on microbial community diversity and structure and are often more significant effectors than climate change forcings (*7–9*). Seasonal variation in sampling date, attributed to short-term variations in environmental parameters, was more important than 10 years of shifted baseline precipitation in a Mediterranean climate, which led to the conclusion that the functional stability of these particular communities is key (*10*). There were inconsistent and non-unidirectional yearly temporal trends in the community response in the Jasper Ridge Global Change Experiment (JRGCE), a 17-year multifactorial [warming, added rainfall, elevated CO_2 and additional nitrogen] experiment. The natural inter-annual variability of the microbial community was more significant than any treatment effect (*11, 12*), which might suggest this community possessed the metabolic flexibility to deal with even large pressed climate forcings.

Yet, individual metabolic processes and pathways, particularly in the case of nitrogen fixation and methanogenesis, have shown significant response to climate change-associated effects, even when the overall community structure or diversity remained unchanged. At JRGCE, individual metabolic processes were linked to specific forcings, for example enzymes related to carbon degradation (starch, hemicellulose, chitin, pectinase, and aromatics) showed a response to elevated CO_2, however the direction and magnitude of these responses was

variable and carbon fixation related genes remained unchanged (*13*). Across study sites, N-
fixation and methanogenesis (measured via abundance of *nifH* and *mcrA*, respectively) are two
of the more commonly reported pathways that show significant change with increased heating
or other climate-change environmental factors (*3, 7, 14*). However, this pattern is not consistent
across all sites, since at the SPRUCE site in a 5-year study where deep peat experienced
intense heating (up to +9°C over ambient), neither a significant change in methanogen-related
genes at 40-50 cm nor a significant change in carbon substrate preference were observed (*15,
16*).
951
The consequences of pulsed vs. pressed disturbances and the sensitivity and response
of the microbial community to those disturbances varies and is seen over timeframes that can
range from hours to multiple decades, due in part to the resistance and resiliency of the
microbial community (*17, 18*). The diversity and functional stability of the peatlands may arise
from strong filtering by the environment (low pH, oligotrophy, metabolite, and plant exudate
concentrations) which promotes a community that is resistant, or resilient, to change (*19, 20*). At
Abisko in northern Sweden, a temperature experiment, raising growing period ambient
conditions by ~1°C for more than 10 years, did cause alterations in the phylogenetic
composition of the bacterial community despite documented changes in gas fluxes (*20*). In a
similarly established warming-focused decadal experiment nearby Abisko at Storflaket, while
the microbial community composition was dependent on depth and decadal heating, the overall
community structure was not different in the thawed layers, and the active layer community
structure was not significantly altered between treatments (*21*). Given these observations, we
advocate a change in focus: from examining whether microbial function can be directly
associated with modifications of community composition associated with climate change
forcings to exploring under what specific contexts functional alterations agree with compositional
alterations (in accordance with refs (*20, 22*)).
969
***Supplemental Note 2**: Literature-based definitions of resistance, resilience and stability*
Resistance is the insensitivity of a community to a disturbance or stressor, or the ability
to withstand change (*17, 23, 24*). Resilience is the rate at which a community is able to return to
its original state after disturbance (*17, 23, 24*). Both resistance and resilience must address the
natural variability that exists in a community's mean state. This is important as stability, defined
as little or no difference between a community's average state before and after disturbance, has
factors of both resistance and resilience and can be assessed through the persistence of
microbial taxa and metabolic functions with time (*18, 25*). It is worth noting that one aspect of
ecological stability depends on the community's ability to coexist in various stable states (*18,
26*).
980
***Supplemental Note 3**: Site description of the model permafrost peatland ecosystem,*
Stordalen Mire
Stordalen Mire, in arctic Sweden, located at 68°20'N 19°03'E, is a permafrost peatland
complex underlain by discontinuous permafrost. The presence or absence of permafrost
determines microtopography and hydrology, creating three thaw stages, palsa, bog and fen
(Fig. 4), with distinct biogeochemistry and plant communities. These three distinct sub-habitats

have been utilized for more than 60 years for plant ecology studies ranging from the Swedish
IBP/PT tundra biome project 1968-1974 to more recent dynamic landscape studies of climate
sensitive arctic atmosphere-biosphere interactions (*27–29*).

Palsa are common in the zone of discontinuous permafrost. The largest central area of
Stordalen Mire is a palsa underlain and elevated 1-2 m above the surrounding landscape and
ponds by continuous ice-rich permafrost (*30*). Palsa is underlain by continuous ice-rich
permafrost with a shallow seasonal active layer (20-50 cm) and unsaturated peats. It is
dominated by cryptogams and low form ericoid shrubs such as those from the genera
Andromeda, *Vaccinium*, and *Rubus*. Permafrost becomes unstable at ambient annual air
temperatures above 0°C. The mean annual air temperatures (MAATs) at the Abisko Research
Station (ANS), ca. 10 km west of Stordalen Mire, have mostly been above 0°C since ca.1995
(*31*) and averaged +0.7°C for the duration of this study, 2011-2017 (SMHI 2024). The Stordalen
Mire palsa is maintained by strong bidirectional winds (*32*) which keeps the snow cover minimal
and allows deep penetration of the winter frost. Wind will also drift snow into surface
depressions and the palsa edges, which change the thermal structure, increasing the heat
capacities of the now wet peat causing the underlying ice-rich permafrost to thaw, further
collapsing the surface and changing the plant community (*33*).

Bogs are found in the intermediate state of the thaw gradient if the permafrost is not
entirely lost. Bogs remain perched above the local water table. The bogs are ombrotrophic
collecting water from limited runoff from surrounding palsa, snow melt and rain, and are
dominated by *Sphagnum* spp.. Bogs are also intermediate between the palsa and fen sites in
their productivity and carbon gas exchange rates (*34*). The seasonal active layer in the bog is
deeper (50-90 cm) than that of the palsa and the water table can vary substantially during the
year but the bog sites remain wet with the deeper peats above the permafrost remaining
saturated.

Fens are the lowest and wettest areas in Stordalen Mire and lack discernible underlying
permafrost. They are connected to the regional water table flowing from the surrounding
watershed and, more immediately from shallow post-glacial lakes to the east and south of the
Mire with a slow flow towards Lake Tornetrask to the north. The water table is usually at or
above the fen surface (*35*): The persistent supply of relatively nutrient rich lake and groundwater
supports productive communities of emergent macrophytes such as *Equisetum* spp. and
graminoids such as *Eriophorum angustifolium* and *Carex rostrata* that grow to heights >1 m in a
single growth season

***Supplemental Note 4**: The peat of the fen at deeper depths are more variable, which*
might explain why this is the sole depth and habitat that shows directional microbiome
change with time rather than just fluctuation.

The deeper fen peats are different and more prone to variability because of more, fresh,
annual inputs of belowground litter due to macrophyte productivity combined with higher rates of
lateral hydrologic transport (lower porewater residence times) because of their local water table
connectivity. As illustrated in Supplemental Note 3, the habitat-specific plant communities have

different productivities (*34*) that are reflected in and are integrally linked to the microbial
communities in the peat below. There are distinct microbial communities and specific
metabolisms in the peats coincident with the obvious surface plant communities (*36*).
Aboveground productivity of these plant communities is affected by abiotic factors such as
temperature, water, and belowground litter inputs (*37*) which in turn will likely affect
belowground microbial and plant activity and variability. Palsa and bog peats are likely to have
more stable conditions in the deeper peat layers. Both of those habitats have lower productivity
of recalcitrant organic matter and restricted seasonal active layers. Palsa has the shallowest
active layer with colder drained peats and woody litter inputs. *Sphagnum* dominated bogs grow
from the top with less variability in belowground litter input into recalcitrant *Sphagnum* peat.
Because bogs are perched there is minimal or no water flow in the deeper, more decomposed
catotelm with longer porewater residence times. Fens, on the other hand, are more productive
with high water conductivity. The roots of the graminoids in the fen will extend through the peat
profile to the basement depth providing substantial amounts of seasonally fresh belowground
litter input.

The MAAT and precipitation at ANS (SMHI 2024) and the CO_2/CH_4 flux rates (*34*)
measured in the Stordalen Mire during 2013 were not anomalous for the study period. Two
events related to the hydrology and temperatures occurred during the 2012 to 2013 winter that
may particularly have affected the deep fen peat communities (Figs 1a and 2a). Pascual and
Johansson (*38*), using meteorological data from ANS, calculated a larger number of melt days
and positive degree days at the Storflaket (7 km west of Stordalen). Thaw days may have
affected the input of the quantity and quality of fresh belowground litter. It was also noted that
the post-glacial ponds lost their ice cover earlier in the spring of 2013 (*39*) which would have
also provided more fen waterflow earlier in the season affecting productivity. Both processes
would have provided more and earlier substrate and water supply potentially affecting the
microbial community structure specifically in the fens.

***Supplemental Note 5**: There is little compelling evidence non-linear responses to time or*
temperature for microbial community composition, function, or assembly.
The data shows little to no evidence of a linear relationship with time or temperature for the
microbial community composition, function, or assembly. There is still a possibility of a
significant non-linear relationship between community composition and time or temperature, or
individual metabolic pathways or organisms that show relationships to temperature. At the
community scale non-linearity may occur since temperature or time could have non-linear
dynamics that create inflection points or other complex structures in the data. Furthermore,
individual pathways or organisms may show more sensitivity than the whole community.
Therefore, we performed several tests for non-linear responses of the community and taxa to
time and temperature.
To investigate non-linear relationships between the microbial community and time or
temperature, we ran generalized additive models (GAMs) to better understand if complex non-
linear dynamics of the microbial community (represented by the first two principle coordinate
axes) could predict year or temperature. To mirror our tests for linear relationships we ran the
models both with and without the effects of habitat and depth. We found no evidence that
microbial composition had a significant non-linear relationship to either time or temperature.

Microbial community composition was a poor, non-significant predictor of year both when
differences due to depth and habitat were not included in the model (GAM; $R^2 = 6.5 \times 10^{-3}$,
microbial smooth term p-value = 0.4, deviance explained = 2.02%) and when they were
included as covariates (GAM $R^2 = 5.6 \times 10^{-2}$, microbial smooth term p-value = 0.142, deviance
explained = 12.3%). For temperature, microbial community was a good predictor of ground
temperature only when the model did not account for habitat and depth (GAM $R^2 = 0.34$,
microbial smooth term p-value = 1.64×10^{-16}, deviance explained = 40.5%). When habitat and
depth were included in the model, the explanatory power of the model increased (GAM $R^2 = 0.6$,
deviance explained = 63.7%, microbial smooth term p-value = 0.38), but the significance of the
terms containing the microbial community principle coordinate axes were rendered non-
significant. Taken together these results support our findings that microbial community
composition was remarkably insensitive over time and to fluctuations in temperature.
To investigate the sensitivity of particular organisms and pathways to temperature we
first chose organisms and pathways of interest to minimize the chance of inflating error rates
due to multiple comparisons. In the case of organisms, we selected those that were present in
WGCNA modules that showed >0.5 correlation coefficients with ground temperature and further
selected only organisms within those modules that had the highest variable importance in
projection (VIP) score (VIP > 1.5). We reasoned that these organisms would be the most likely
to exhibit a strong population-level response to temperature change in our dataset. We then ran
linear models to the relative importance of ground temperature and depth (as well as their
interaction) on the centered log-ratio (CLR)-transformed abundance of each organism within a
particular habitat (model formulas took the form: $GenomeAbundance_{ij}$ ~ $SoilTemperature_j$ +
$Depth_j$ + $SoilTemperature_j{:}Depth_j$, where $GenomeAbundance_{ij}$ is the CLR-transformed genome
abundance of taxon i in habitat j). In some cases, $GenomeAbundance_{ij}$ was log-transformed to
meet the assumptions of linear regression. Of the 109 organisms tested, only 14 showed any
significant effect of temperature (linear regression, R^2 range: 0.11-0.57, $p<0.05$), and in all 14
cases, temperature had a minor negative effect on abundance (standardized effects size ranged
from -0.079 to -0.036).
In the case of metabolic pathways, sixteen pathways of the 48 pathways analyzed,
selected for their central contributions to carbon, nitrogen, and sulfur cycling in Stordalen Mire,
were chosen for temperature analysis. We reasoned that temperature sensitivity in these
sixteen critical pathways were most likely to have downstream effects in the community-level
metabolism due to their centrality to key processes such as methanogenesis, or nitrogen and
sulfur acquisition. The temperature sensitivity of pathways was analyzed similarly as for
organisms, substituting cumulative weighted pathway abundance for genome abundance. Of
the sixteen pathways only CAZymes for xyloglucan degradation in the fen showed any
significant effect of temperature ($R^2 = 0.57$, temperature effect p-value < 0.05, standardized
temperature effect size = -0.097). Collectively, these results support our findings that even at
the granular level of individual organisms and pathways, the microbes and metabolisms in these
communities are resistant to fluctuations in temperature over the 7 years, and that differences in
habitat and depth have a larger effect on community variation in abundance of organisms and
pathways than temperature.
1112

Supplemental Note 6*: **Carbon generalist genomes are significantly larger in size than**
specialists, which supports ecological literature, linking genome size to metabolic
flexibility and functional redundancy.

To date, research on microbial generalism has focused more often on "habitat
generalism" than on the metabolic generalism we investigate here (*40*, *41*). In these studies, the
breadth of an organism's niche has typically been inferred through measurements of abundance
across different habitats (*42*, *43*). Organisms occupying narrow or broad fundamental niches
have then been taken to be specialists or generalists, respectively, with functional inferences
occurring only downstream of this classification (*40*, *44*, *45*). Because generalism is expected to
come at a cost (e.g., lower catalytic efficiency; higher maintenance costs for larger genomes;
competitive exclusion), specialists have been hypothesized to dominate in many habitats (*46*,
47). Alternatively, ecological theory suggests that broader niches should allow generalists to
maintain more stable populations, while narrow-niche specialists should be less able to
withstand environmental variation. Ready access to a broad pan-genome has been
hypothesized to support the production of metabolic generalists (*48*), which should then
succeed under selection in dynamic environments (*45*). It is not clear what aspects of
environmental variability at Stordalen Mire drive the abundance of metabolic generalists.
Furthermore, depending on the strength and ordering of substrate preferences in different
organisms, apparent metabolic generalism may effectively be specialization in disguise (*48*).
1132
To define specialists and generalists on the basis of function rather than habitat range,
we first annotated genomes for metabolic functioning and then grouped by specialization (see
Methods). We did not consider genome size in our classification of metabolic specialists and
generalists. We then hypothesized, in accordance with genome streamlining and ecological
inference (*49–51*), that generalist organisms encoding more metabolic pathways would have
larger genomes than methanogens and other metabolic specialists (figS23). Consistent with this
prediction, our carbon generalists had significantly larger genomes (mean 5.00 ± 1.63 Mb,
Tukey test, $p < 7.85$ x10^{-12}) than methanogens (mean 2.30 ± 0.45 Mb) and most other carbon
specialists (Tukey test, p-values range from <7.78 x10^{-12} to <1.48 x10^{-4}). Among specialists,
homoacetogen genome sizes (mean 4.06 ± 1.70 Mb) were indistinguishable from those of
methanotrophs or fermenters (mean 3.50 ± 1.28 Mb) but significantly larger than the
macromolecule degraders (mean 3.36 ± 1.36 Mb, Tukey test, $p < 0.039$). Monomer degraders
(mean 2.75 ± 0.97 Mb) were indistinguishable in genome size from both these larger specialists
and from methanogens. Methanotrophs were the sole exception (mean 4.25 ± 0.64 Mb, Tukey
test, not significant) to the finding that generalists had larger genomes than specialists. The
methanotrophs identified here are aerobic, which might partially explain why their genomes are
larger than those of other specialists (*52*); moreover, we found that these methanotrophs
typically encoded chemotaxis, various CAZyme and carbon fermentation pathways, and
nitrogen, sulfur and phosphate metabolic capabilities. Collectively, this pattern of genome size
differentiation between generalists and specialists across metabolism types provides confidence
in our pathway annotation-based calls of carbon generalists and specialists.
1154
Functional generalism, observed within individual organisms, can lead to functional
redundancy in the community (*41*, *43*, *53*). Functional redundancy, defined here and elsewhere

as the ability of taxonomically distinct organisms to encode the same dissimilatory metabolic
capabilities (*44*), is a common observation in microbial soil systems across global latitudes (*54*).
Consistent with the observations throughout this work, functional redundancy that results in
taxonomic differences in intermediate reactions of the metabolic network is unlikely to affect the
endpoints of dissimilatory metabolic pathways (*44*), e.g., methane production in anaerobic
digesters (*55*).
1163
1164   ***Supplemental Note 7: Methanogens have simplistic and specialized metabolic calls often***
1165   ***with limited to no recognized nitrogen capabilities, which leads to a larger number of***
1166   ***associations with other organisms.***
1167   The methanogens with a high importance to the overall structure and number of
connections in WGCNA and MTN networks, respectively, were highly specialized with few
annotated metabolisms. Hydrogenotrophic methanogens, dominant in the bog, were quite
specialized with simplified carbon and nitrogen metabolic calls. There was greater versatility and
diversity in methanogenic metabolisms in the fen with more variations of carbon fixation and
degradation pathways, and more types of methanogenesis. Two of the most crucial organisms
in the bog and fen networks, the hydrogenotrophic methanogens *Ca*. 'M. stordalenmirensis'
(bog) and *Ca*. 'M. crilli' (fen), lacked recognizable nitrogen metabolism calls (figS34).
Methanogens capable of nitrogen fixation had fewer connections in the MTN and lesser
influence (via degree centrality and betweenness metrics) in WGCNA networks (figS34; figS35;
figS36). These observations, along with those discussed in association with Fig 3c&d of the
main text, led us to propose the hypothesis that if a methanogen has nitrogen-fixation or other
nitrogen-linked decomposition pathways it will have fewer temporal connections since it is less
likely to need additional nitrogen inputs from other members of the community.
1181
1182   To further assess this hypothesis we examined whether we could identify other genes
and functions, not directly addressed in our metabolic annotations, that could differentiate the
methanogens with more connections from those with fewer connections (See Methods section
on Genome Annotation for more details). Those methanogens that tended to be more
independent with fewer associations to other organisms had five different functions that were
either significantly enriched (adjusted q-value ≤ 0.05) or nearly significant (q-value < 0.055).
Four of the functions (K02586, K02587, K02591, K02592) pertained to nitrogenase and nitrogen
fixation, and one (COG1048) was involved in lysine biosynthesis, a critical amino acid. That a
group of nitrogen-fixing methanogens were enriched in genes that pertain to nitrogen cycling
and fixation, when compared to non-nitrogen fixing methanogens, was not unexpected but was
consistent with our hypothesis. The only nearly significant (q value < 0.054) functions that were
enriched in the methanogenic group with more connections and associates with time ("less
independent" methanogens) were COG4656 and COG1035 which encode Na+ translocating
ferredoxin: NADH oxidoreductase (Rnf). This Na^+ pump and Rnf complex have been found to
play important roles in energy conservation, ATP generation and electron shuttling (*56*, *57*).
Perhaps, the highly reducing conditions of the anoxic environment and reliance on reverse
electron transport could be beneficial to those organisms, but the precise reason why an energy
conservation function would be enriched in an apparent more collaborative group of
methanogens is not entirely clear. Collectively, these results support the hypothesis and trends

seen in the metabolic, MTN and WGCNA analysis that if a methanogen isn't clearly limited in
nitrogen capabilities they have fewer associations and alternatively when there isn't a known
nitrogen pathway those methanogens likely need external sources of nitrogen from surrounding
community members.
1205
1206 ***Supplemental Note 8: Etymology.***
1207 'Acidi' from Latin noun neuter acidum an acid, adj. acidus, sour.
1208
Description of 'Candidatus Acididegradans cibantis' sp. nov.
Classification and differentiation based on average nucleotide identity (ANI). 'degradans' Latin
verb neuter present active participle: degrade. 'Cibantis' Latin verb neuter present active
participle: to feed. This organism is of the order Acidimicrobiales in the Actinobacteriota phylum,
was classified as a carbon generalist and proposed to be providing key substrates to other
organisms, including methanogens.
1215
Description of 'Candidatus Acididegradans virginiarichiae' sp. nov.
Classification and differentiation based on ANI. 'degradans' Latin verb neuter present active
participle: degrade. 'virginiarichiae', New Latin noun feminine, named after Virginia Rich, the
Ohio State University, USA, in recognition of her pioneering efforts on deciphering the controls
that climate change has from scales of genes to ecosystems in thawing permafrost. This
organism is of the order Acidimicrobiales in the Actinobacteriota phylum, and was classified as a
carbon generalist.
1223
1224 Description of 'Candidatus Acidimolecula latum' sp. nov.
1225 Classification and differentiation based on topology and ANI. 'molecula' Latin noun feminine:
1226 molecule. 'latum' Latin adjective neuter: broad. This organism is of the genus Terracidiphilus in
1227 the Acidobacteriota phylum, was classified as a carbon generalist and possessed a broad range
1228 of the carbon metabolisms considered.
1229
1230 Description of 'Candidatus Acidimolecula amplum' sp. nov.
1231 Classification and differentiation based on ANI. 'molecula' Latin noun feminine: molecule.
1232 'amplum' Latin adjective neuter: large. This organism is of the genus Terracidiphilus in the
1233 Acidobacteriota phylum and was classified as a carbon macro-molecule degrader.
1234
1235 Description of 'Candidatus Acidiacetogenum stordalenmirensis' sp. nov.
1236 Classification and differentiation based on ANI. 'acetogenum' New Latin noun neuter: acetogen.
1237 'stordalenmirensis' New Latin adjective masculine, of or belonging to Stordalen Mire, Sweden
1238 from where the species was characterized. This organism belongs to the family
1239 Thermacetogeniaceae in the Firmicutes phylum and was classified as a homoacetogen.
1240
Description of 'Candidatus Abiskobacterium ruthvarnerae' sp. nov.
Classification and differentiation based on ANI. 'Abiskobacterium' New Latin noun neuter:
bacterium from Abisko, Sweden from the surrounding area where the species was
characterized. 'ruthvarnerae' New Latin noun feminine, named after Ruth Varner, University of

New Hampshire, USA in recognition of her pioneering efforts on deciphering the controls that
climate change has from scales of genes to ecosystems in thawing permafrost. This organism
belongs to the family of Bryobacteraceae, was classified as a carbon generalist and possessed
nearly all of the carbon metabolisms considered.
1249
Description of 'Candidatus Acidiflorens viviennsis' and 'Candidatus Acidiflorens anabellensis',
sp. nov.
Classification and differentiation based on ANI and topology, and ANI, respectively.
'Acidiflorens', New Latin an organism that blooms in acid (*58*). 'viviennsis' and 'anabellensis'
New Latin adjective feminine, of or belonging to Vivien and Annabelle, respectively, the mothers
of Virginia Rich, the Ohio State University, USA and Ruth Varner, University of New Hampshire,
USA, respectively, whom they wish to be honored.
1257
1258 Description of 'Candidatus Acidiaspiraculum latum' sp. nov.
1259 Classification and differentiation based on topology and ANI. 'a' Latin preposition: away from.
1260 'spiraculum' Latin noun neuter: air. ''latum' Latin adjective neuter: broad. This organism belongs
1261 to Verrucomicrobiota, found to be important in the anoxic fen habitat and was classified as a
1262 carbon generalist and possessed a broad range of the carbon metabolisms considered.
1263
1264 Description of 'Candidatus Acidiaspiraculum fermentum' sp. nov.
1265 Classification and differentiation based on topology and ANI. 'a' Latin preposition: away from.
1266 'spiraculum' Latin noun neuter: air. ''fermentum' Latin noun neuter: fermentation. This organism
1267 belongs to Verrucomicrobiota, found to be important in the anoxic fen habitat and was classified
1268 as a fermenter.
1269
1270

1271

1272

1273

1274

1275

1276

1277

1278

1279

1280

Supplemental Figures

1282

Supplemental Figure 1: Overview of samples included in this study. Each point represents the average depth of a sample (cm) with error bars showing the minimum and maximum depth of the sample. Blue lines represent water table depth where appropriate, black lines show active layer depth, black lines with x's show the maximum depth of the active-layer probe (where the permafrost was not reached). We note that in 2012, a deeper probe was used but broke during sampling of the first bog core, after which a standard 1 m probe was used.

A)

B)

Supplemental Figure 2: A) The trends of mean annual temperature at Stordalen Mire
(as measured by the Abisko weather station SMHI) have been increasing for over fifty
years (light grey, C-losses and associated habitat conversion discussed in ref _29_),
despite large inter-annual fluctuations. Our study captures the recent period from 2011-
2017 (dark grey). **B)** There are important seasonal differences in the temperature
change over the last fifty years. The summer, which represents the time of peak
emissions, is one of the more stable times of year, while the winters and shoulder
seasons have been experiencing the most rapid increases in temperature. Panels are
presented in the following order: winter, spring, summer, fall, all seasons.

1332

Supplemental Figure 3: There are no significant relationships between SingleM based
alpha diversity metrics and time as a continuous variable. Alpha diversity, as measured
as richness **(A)**, Pielou's evenness **(B)**, and Shannon diversity **(C)** for SingleM results
rarefied to 1000 sequence hits per marker gene and averaged across marker genes
with linear model significance testing against time (in years) as a continuous variable.

SingleM based Alpha Diversity for each habitat by depth against continuous time (as year)

Habitat	Depth	Alpha	R^2	p-value
Fen	10-19	diversity	0.52	0.043
Fen	10-19	richness	0.57	0.031
Fen	30-39	evenness	0.58	0.017
Fen	30-39	diversity	0.58	0.017
Fen	30-39	richness	0.54	0.023

Supplemental Figure 4: The only significant relationships between SingleM based alpha diversity metrics and time as a continuous variable occurred in the fen (10-19 cm and 30-39 cm). Alpha diversity (measured as richness, Pielou's evenness, and Shannon diversity) for SingleM results rarefied to 1000 sequence hits per marker gene and averaged across marker genes by depth across habitats with linear model significance testing against time (in years) as a continuous variable.

1391

<u>Supplemental Figure 5:</u> Divergence of each sample from the median community
profile in each habitat. Points within each boxplot are jittered to represent individual
samples, with the Bray-Curtis distance (determined from MAG abundances) indicating
the divergence of each sample from the median profile. The data is plotted in two
panels: **(a)** samples are colored by depth, and **(b)** samples are colored by year. This
allows for comparison of divergence patterns across different depths and years within
each habitat.

1419
<u>Supplemental Figure 6:</u> Comparing pairwise sample distances within year and
between years for each habitat (palsa, bog, and fen). Points represent individual
pairwise comparisons between samples, with the year differential between the two
samples indicated by a gradient color scale. The Wilcoxon ranked sum test p value is
displayed on the plots, comparing within-year versus between-year groups. This
provides statistical insights into the temporal variability of samples within each habitat
type.

1446

<u>Supplemental Figure 7:</u> Fluctuation over time was assessed by using time as an unordered categorical variable. Heatmap visualization of microbiome temporal fluctuation at different levels of granularity, spanning organisms (from phylum to strains to diversity metrics, via single copy marker genes and MAGs), functions (from pathways to protein clusters, both annotatable and non-annotatable), and organization (community assembly metrics of stochasticity and determinism), in each habitat and depth. The amount of variation explained by time (as Mantel R or R^2 depending on statistical test appropriate for each data type; see Methods) is indicated with the intensity of the shaded color; white asterisks denote significant correlations to time. Different statistical tests were used to test structure with time depending on the data type (e.g. multivariate vs. single variate data). Change in community phylogenetic and stochastic assembly data was tested with a Mantel test. Change in organisms and functions metrics was tested with a redundancy analysis with time as an explanatory factor.

1476

Supplemental Figure 8:

8 MAGs from the fen at 30-39 cm depth showed both significant differential centered log-ratio
(CLR) abundances across years and significant correlations to time. Lines of best fit are shown
in blue with standard error indicated in gray. Spearman rho and false discovery rate (FDR)
adjusted p-values for each test are displayed in the top left-hand corner of each subplot. Each
organism is designated at the top of the plot by its phylum and unique genome identifier.

1498
1499 **Supplemental Figure 9:** Encoded cumulative relative abundance of meta-pathways
1500 with depth and time across habitats. Each point represents the cumulative relative
1501 abundance of all species that encode a pathway within the given meta-pathway.

1518
Supplemental Figure 10: Metabolic pathways that significantly change over time. Pathways with
changing cumulative abundance across years in metagenomes (top), in 30-39cm Fen,
representing 10/13 of the significantly changing pathways; and metatranscriptomes (bottom), in
the 10-19cm Fen, representing 34/132 of the significantly changing pathways. Left column
represents cumulative abundances over time for significantly changing pathways, right column
represents year-to-year log2 fold-change. Error bars represent standard error. All pathways
were significant p<0.05 in Likelihood-ratio test of beta-regression (logit) with and without Year
parameter.

1546

1547 **<u>Supplemental Figure 11:</u>** Relative abundance of each group of carbon specializations
1548 across habitats, depths and years and averaged across samples. "NA" represents
1549 genomes that were unable to be assigned to a specialization.

1550

1551

1552

1553

1554

1555

1556

1557

1558

1559

1560

1561

1562

1563

1564

1565

1566

1567

1568

1569

1570

1571

1572

1573

Supplemental Figure 12: Temporal comparison of network structures across habitats.

This figure examines year-to-year variations in network metrics for the three habitats: Palsa, Bog, and Fen. Multilayer networks (MTN) were constructed for each year and habitat, and metrics were calculated to compare network structure over time. Pairwise comparisons are shown using boxplots, with individual data points jittered and color-coded to indicate the comparator year. Statistical significance is indicated above each boxplot through a Kruskal-Wallis test. (A) Dissimilarity index of degree distributions. The dissimilarity in degree distributions between years within each habitat. The degree distribution reflects the frequency of nodes having a specific number of connections. A higher dissimilarity index indicates greater differences in connectivity patterns between years. (B) Degree correlation (ρ\rho$\rho$). Pairwise correlations of node connectivity patterns across years. (C) Jaccard similarity coefficient of triangles. Pairwise comparisons of triangle connectivity (three connected nodes) between years. The coefficient measures the proportion of shared triangles relative to the total number of unique triangles across the years, indicating the stability or change in triadic structures over time.

1598
Supplemental Figure 13: This figure shows upset plots of clustered modules,
generated by selecting the highest possible likelihood of clustering across layers, based
on varying the interlayer edge weights. The y-axis represents the number of modules
present in the years indicated by the bolded points below the x-axis. The values below
the x-axis indicate the year where a module was present (1-7 is 2011-2017). Generally,
the plots reveal higher levels of module persistence in the Palsa and Bog compared to
the Fen, where ephemeral modules are more prevalent.

1631

<u>Supplemental Figure 14:</u> Since we have no direct measurement of interlayer edge

weight (which roughly corresponds how strongly one year's network influences the

subsequent year's community), we tested a variety of interlayer edge weights (ranging

from 0 to 1 with 0 being the least relationship between years) to assess the impact on

the clustering of modules. The top row of panels shows how many modules formed

across the range of relax rates (the change in interlayer edge weight). The middle row

shows the distribution of the persistence of modules (the number of layers where a

module is present) at various relax rates. The bottom row shows the average relative

abundance of organisms within each kind of module. Modules are grouped into three

categories: ephemeral (present in 1 layer), episodic (present in 2–3 layers), and

persistent (present in ≥ 4 layers). At higher relax rates, fewer modules are detected but

have higher persistence over time. Compared to the bog and palsa, the fen shows a

higher abundance of episodic and ephemeral modules across all relax rates.

1645

1646

1647

1648

1649

1650

1651

1652

1653

1654

1655

1656

1657

Supplemental Figure 15: The proportion of assembly processes in each habitat. To control for depth, cross-depth sample-to-sample comparisons have been filtered out. The proportion of sample-to-sample comparisons within each habitat whose assembly processes were characterized as homogenous selection, heterogenous selection, homogenizing dispersal, dispersal limitation and drift, or pure drift are indicated on the y-axis. While each depth interval is indicated on the x-axis. Purple-tinted colors are indicative of deterministic assembly processes, orange-tinted colors are indicative of stochastic assembly processes.

1691

Supplemental Figure 16: The proportion of assembly processes in each habitat as a function of depth (in 10-cm intervals). The proportion of sample-to-sample comparisons whose assembly processes were characterized as homogenous selection, heterogenous selection, homogenizing dispersal, dispersal limitation and drift, or pure drift are indicated on the y-axis. Each depth interval is indicated on the x-axis. Each row of plots represents a different habitat across the thaw gradient. Hydrologic connectivity as measured by time below the water table during the growing season prior to sampling increases with thaw (bogs are sometimes waterlogged and fens are nearly always waterlogged) and time below the water table increases with depth, particularly in the bog.

1715
Supplemental Figure 17: Mantel correlograms showing the Pearson correlation
between between-MAG environmental optima and between-MAG phylogenetic distance
for a number of environmental gradients at the site. Significant correlations (P < 0.05,
filled points) indicate that a phylogenetic signal in MAG ecological niche for the
associated distance class. Significant positive correlations indicate ecological similarity
among MAGs is higher than expected for the associated distance class. Gradients
summarized from left to right and top to bottom: all - composite of all site environmental
data distances summarized by gower distance, alphaC - fractionation coefficient,
d15N_peat - delta 15N value of peat N, DepthAvg_neg - the depth optima of each MAG,
N.percent - the % N in peat (by weight), Nitrate.uM - Nitrate concentrations (uM),
pH_peat - pH of peat, samp_dist_WT - typical distance of a MAG from the water table,
Sulfate.uM - sulfate concentration (uM), T_air_deg_C - air temperature, T_soil_deg_C -
soil temperature, TN.mM - total nitrogen concentration (organic and inorganic).

1743

Supplemental Figure 18: Depth-dependent temperature decreases for all habitats across years. We ran several different mixed effects models to understand the effect of year on the relationship between ground temperature and depth, we tested year as both a random slope and a random intercept, as well as running the model without year as a random effect. The best model (as determined by Akaike and Bayesian information criterion, AIC and BIC, respectively) was one in which year was a random intercept, indicating that while the surface temperature does vary from year to year, the relationship (slope) between temperature and depth is fairly consistent within habitat from year to year. Soil temperature (y-axis) was most strongly related to depth (x-axis) in the palsa (slope = -0.352 +/- 0.024 deg/cm), and more weakly related in the bog and fen (slope = -0.25 +/- 0.02 and -0.11 +/- 0.02 deg/cm, respectively). Here, "grand slope" and "grand intercept" represent the best fit pooled mean estimate of the relationship between temperature and depth (slope), and average temperature at the surface (intercept) over the seven years. sd_Year shows the variation among years around that mean. Year-to-year differences in surface temperature were most variable in the bog (sd_Year = 3.39), and least variable in the Palsa (sd_Year = 1.22).

	Not Depth Corrected		Depth-Corrected (partial mantel)	
Habitat	Rho	Significance	Rho	Significance
All	0.126	0.001	0.106	0.001
Palsa	0.314	0.001	-0.129	0.978 (NS)
Bog	0.269	0.001	0.152	0.003
Fen	0.24	0.004	0.24	0.005

1776
1777 **Supplemental Table 2**: Mantel tests of the relationship between community composition
1778 and temperature. Ground temperatures show low-to-moderate positive relationships with
1779 community composition (rho ranges 0.106-0.31). When depth-corrected (via partial
1780 Mantel tests) the relationship weakens but is mostly still significant except in palsa
1781 where the depth-temperature relationship is strongest.

1811
Supplemental Figure 19: A single WGCNA module showed a relationship to
temperature that was greater than its relationship to depth. The 'brown' module from the
fen samples was strongly correlated to the 2-week average air temperature leading to
sample collection date. Each data point represents one MAG within the brown module,
where the x-axis shows the correlation between the MAG shown and the representative
eigenvector of the module. Additionally, each MAG is also correlated to the air
temperature on the y-axis.

1844

<u>Supplemental Figure 20:</u> (a) The WGCNA module containing the MAGs from all

habitats most highly correlated to $\delta^{13}C_{CO2}$ of porewater (y-axis) over seven years of data

with the relationship of the MAG to the module representative (x-axis), where stronger

relationships between the environmental variable and the module representative

eigenvector indicate a robust relationship. **(b)** Measured vs predicted values of $\delta^{13}C_{CO2}$

porewater values using partial least squares regression with leave-one-out cross

validation.

1852

1853

1854

1855

1856

1857

1858

1859

1860

1861

1862

1863

1864

1865

1866

1867

1868

1869

1870

1871

1872

1873

Supplemental Figure 21: (a) The WGCNA module containing the MAGs from all

habitats most highly correlated to $\alpha_{CO2\text{-}CH4}$ of porewater (y-axis) over seven years of

data with the relationship of the MAG to the module representative eigenvector (x-axis),

where stronger relationships between the environmental variable and module

representative indicate a robust relationship. **(b)** Measured vs predicted values of $\alpha_{CO2\text{-}}$

$_{CH4}$ of porewater values using partial least squares regression with leave-one-out cross

validation.

1881

1882

1883

1884

1885

1886

1887

1888

1889

1890

1891

1892

1893

1894

1895

1896

1897

1898

1899

1900

1901

1902

1903

1904 **Supplemental Figure 22: (a)** The WGCNA module containing the MAGs from all
1905 habitats most highly correlated to total nitrogen concentration (TN.mM) of porewater (y-
1906 axis) over seven years of data with the relationship of the MAG to the module
1907 representative (x-axis), where stronger relationships between the environmental
1908 variable and module representative indicate a robust relationship. **(b)** Measured vs
1909 predicted values of TN.mM porewater values using partial least squares regression with
1910 leave-one-out cross validation.

1932
1933 **<u>Supplemental Figure 23:</u>** Genome size for each of the different carbon specializations.
1934 Carbon generalists had significantly larger genomes (mean 5.00 ± 1.63 Mb, Tukey test,
1935 $p < 7.85 \times 10^{-12}$) than methanogens (mean 2.30 ± 0.45 Mb) and most other carbon
1936 specialists (Tukey test, p-values range from <7.78 $\times 10^{-12}$ to <1.48 $\times 10^{-4}$).

Decision tree

Per MAG/Per pathway

1. % of total pathway

<60%

≥60%

2. % of reactions present

<70%

≥70%

3. Signature gene/s present

False

True

MAG does not
have pathway

MAG has pathway

1. 60% cutoff chosen by inflection point
2. 70% cutoff chosen to require all reactions to be present for 1/2/3-reaction pathways
3. Signature genes used to distinguish pathways with high overlap (e.g. Wood-Ljungdahl with hydrogenotrophic methanogenesis). In addition, signature gene within CAZy context means gene unique to that pathway.

1959
1960
1961 **Supplemental Figure 24:** A conservative decision tree for positive functional pathway
1962 calls: genomes had to contain ≥60% of the total genes in a pathway, ≥70% of the total
1963 reactions in a pathway, and the 'signature' gene(s) of a pathway to be positively
1964 identified. The 70% cutoff was chosen to require all reactions to be present for
1965 pathways with 3 or fewer reactions, and inclusion of signature genes helped distinguish
1966 between pathways with high overlap (e.g. Wood-Ljungdahl and hydrogenotrophic
1967 methanogenesis). In most cases, signature genes are those typically used in the
1968 literature to identify a pathway's presence (e.g. *acsBCD* for Wood-Ljungdahl). See
1969 Methods section on Genome Annotation for more details.

Supplemental Figure 25: Metabolic pathway expression, determined as pathway-
based averaged transcripts per million (tpm), across habitats. Transcripts were
averaged across reaction proteins and reactions, and summed across parallel
reactions. Each bar represents the mean ± SD of pathway-averaged expression,
summed across genomes and averaged across samples.

Carbohydrate Associated Enzymes (CAZymes)

Supplemental Figure 26: The abundance and taxonomy of the MAGs across habitats that encode xylan **(a)**, starch **(b)**, and amorphous cellulose **(c)** CAZymes: box plots of encoded cumulative relative abundance over 7 years for each habitat with depth lumping of 0-9, 10-19, 20-29 and 30-39cm. Black stars represent significant differences between habitat and specific colored stars represent significant differences between depths within that habitat (p<0.05, Likelihood-ratio test of beta-regression (logit) with and without Habitat parameter). Treemap taxonomy where rectangles represent dereplicated MAGs, colored according to phylum and sized proportional to relative abundance.

Nitrate reduction

2051

2052 **<u>Supplemental Figure 27:</u>** Dissimilatory **(a)** and assimilatory **(b)** nitrate reduction: box
2053 plots of encoded cumulative relative abundance over 7 years for each habitat with depth
2054 lumping of 0-9, 10-19, 20-29 and 30-39cm. Black stars represent significant difference
2055 between habitat, and specific colored stars represent significant difference between
2056 depths within that habitat (p<0.05, Likelihood-ratio test of beta-regression (logit) with
2057 and without Habitat parameter). Treemap taxonomy where rectangles represent
2058 dereplicated MAGs, colored according to phylum and sized proportional to relative
2059 abundance.

Nitrogen fixation

2077

Supplemental Figure 28: The abundance and taxonomy of the MAGs that encode and transcribe nitrogen fixation across habitats. **a)** Box plots of encoded cumulative relative abundance over 7 years for each habitat with depth lumping of 0-9, 10-19, 20-29 and 30-39cm. Black stars represent significant differences between habitat and specific colored stars represent significant differences between depths within that habitat (p<0.05, Likelihood-ratio test of beta-regression (logit) with and without Habitat parameter). Treemap taxonomy where rectangles represent dereplicated MAGs, colored according to phylum and sized proportional to relative abundance. **b)** Taxonomic expression, averaged over specific metabolic pathway, across habitats and with depth.

Chitin related

2103
2104

Supplemental Figure 29: (a) Various possible degradation pathways of chitin as
determined via KEGG and CAZy.org databases. Overall positive pathway call for chitin
(b), specific glycoside hydrolase (GH) 18 exo-chitinase (non-reducing end), *chiB*, EC:
3.2.1.200 **(d)**, and carbohydrate esterase (CE)4 **(e)**: box plots of encoded cumulative
relative abundance over 7 years for each habitat with depth lumping of 0-9, 10-19, 20-
29 and 30-39cm. Black stars represent significant differences between habitat and
specific colored stars represent significant differences between depths within that
habitat (p<0.05, Likelihood-ratio test of beta-regression (logit) with and without Habitat
parameter). Treemap taxonomy where rectangles represent dereplicated MAGs,
colored according to phylum and sized proportional to relative abundance. **(c)**
Proportion of cells transcribing positive pathway call for chitin **(b)**, as determined by
dividing the number of MAGs encoding the pathway that had expression by the total
number of MAGs encoding the specific pathway across habitats and with depth.

Dissimilatory sulfur oxidation

2129

Supplemental Figure 30: The abundance and taxonomy of the MAGs that encode and

transcribe dissimilatory sulfur oxidation across habitats. **a)** Box plots of encoded

cumulative relative abundance over 7 years for each habitat with depth lumping of 0-9,

10-19, 20-29 and 30-39cm. Black stars represent significant differences between habitat

and specific colored stars represent significant differences between depths within that

habitat ($p<0.05$, Likelihood-ratio test of beta-regression (logit) with and without Habitat

parameter). **b)** Treemap taxonomy where rectangles represent dereplicated MAGs,

colored according to phylum and sized proportional to relative abundance. **c)**

Taxonomic expression, averaged over specific metabolic pathway, across habitats and

with depth.

2140

2141

2142

2143

2144

2145

2146

2147

2148

2149

2150

2151

2152

2153

2154

Dissimilatory sulfate reduction

2155

Supplemental Figure 31: The abundance and taxonomy of the MAGs that encode and
transcribe dissimilatory sulfate reduction across habitats. **a)** Box plots of encoded
cumulative relative abundance over 7 years for each habitat with depth lumping of 0-9,
10-19, 20-29 and 30-39cm. Black stars represent significant differences between habitat
specific colored stars represent significant differences between depths within that
habitat (p<0.05, Likelihood-ratio test of beta-regression (logit) with and without Habitat
parameter). **b)** Treemap taxonomy where rectangles represent dereplicated MAGs,
colored according to phylum and sized proportional to relative abundance. **c)**
Taxonomic expression, averaged over specific metabolic pathway, across habitats and
with depth.

Assimilatory sulfate reduction

2181

Supplemental Figure 32: The abundance and taxonomy of the MAGs that encode and transcribe assimilatory sulfate reduction across habitats. **a)** Box plots of encoded cumulative relative abundance over 7 years for each habitat with depth lumping of 0-9, 10-19, 20-29 and 30-39cm. Black stars represent significant differences between habitat and specific colored stars represent significant differences between depths within that habitat (p<0.05, Likelihood-ratio test of beta-regression (logit) with and without Habitat parameter). **b)** Treemap taxonomy where rectangles represent dereplicated MAGs, colored according to phylum and sized proportional to relative abundance. **c)** Taxonomic expression, averaged over specific metabolic pathway, across habitats and with depth.

Sulfur-containing Polysaccharide CAZymes

2207

Supplemental Figure 33: The abundance and taxonomy of the MAGs that encode and
transcribe the genes necessary to degrade sulfur-containing polysaccharides across
habitats. a) Box plots of encoded cumulative relative abundance over 7 years for each
habitat with depth lumping of 0-9, 10-19, 20-29 and 30-39cm. Black stars represent
significant differences between habitat and specific colored stars represent significant
differences between depths within that habitat (p<0.05, Likelihood-ratio test of beta-
regression (logit) with and without Habitat parameter). b) Treemap taxonomy where
rectangles represent dereplicated MAGs, colored according to phylum and sized
proportional to relative abundance. c) Taxonomic expression, averaged over specific
metabolic pathway, across habitats and with depth.

2233
2234 **Supplemental Figure 34:** The average degree centrality of methanogens, and their
2235 carbon and nitrogen related metabolic pathway calls from genome annotations for the
2236 **(a)** bog and **(b)** fen habitats.

Supplemental Figure 35: The number of yearly connections in the bog habitat, derived from MTN, that exist between methanogens and other community members. The x-axis ("number of layers") is the number of years that specific neighbors persist (significant correlation exists). The y-axis (count) is the number of lineages positively correlated to each methanogen for each number of years.

Supplemental Figure 36: The number of yearly connections in the fen habitat, derived from MTN, that exist between methanogens and other community members. The x-axis ("number of layers") is the number of years that specific neighbors persist (significant correlation exists). The y-axis (count) is the number of lineages positively correlated to each methanogen for each number of years.

2320

<u>Supplemental Figure 37:</u> The specific genes that encode the carbon, nitrogen and
sulfur pathways annotated in this work show functional redundancy. The difference in
community composition changes (as measured by the MAG-resolved community Bray-
Curtis distance) more rapidly than changes in genes encoding annotated functional
pathways ("Bray-Curtis Distance - Pathways"). This means that as new MAGs are
added to the community, new pathways accumulate more slowly; stated differently, as
every new MAG is added to the community, it is more likely to have the same pathways
as are already observed.

2353

<u>Supplemental Figure 38:</u> Temporal variation in carbon specialization of MAGs across

habitats and depths. Violin plots display the coefficient of variation (CV) over time for

approximately 1,800 dereplicated MAGs (metagenome-assembled genomes),

categorized by metabolic specialization across three habitats and four depth intervals.

Each panel represents a depth interval, with the y-axis showing the CV and each data

point corresponding to an individual MAG. The x-axis and colors indicate the metabolic

specialization categories, allowing for comparisons of CV across specializations within

and between habitats. Temporal variation in specialization suggests overall habitat

stability but also reveals potential inconsistencies with traditional generalist-specialist

definitions, particularly in the context of carbon metabolism.

2364

2365

2366

2367

2368

2369

2370

2371

2372

2373 **Supplemental Figure 39:** Phylogenetic tree of the dereplicated ~1,800 MAGs across
2374 habitat, depth interval, shaded by habitat and the coefficient of variation, color coded by
2375 carbon specialization.

Supplemental References

2385

1.   L. Wu, Y. Zhang, X. Guo, D. Ning, X. Zhou, J. Feng, M. M. Yuan, S. Liu, J. Guo, Z. Gao, J.
Ma, J. Kuang, S. Jian, S. Han, Z. Yang, Y. Ouyang, Y. Fu, N. Xiao, X. Liu, L. Wu, A. Zhou,
Y. Yang, J. M. Tiedje, J. Zhou, Reduction of microbial diversity in grassland soil is driven
by long-term climate warming. *Nat. Microbiol.* **7**, 1054–1062 (2022).

2.   K. Xue, M. M. Yuan, Z. J. Shi, Y. Qin, Y. Deng, L. Cheng, L. Wu, Z. He, J. D. Van
Nostrand, R. Bracho, S. Natali, E. A. G. Schuur, C. Luo, K. T. Konstantinidis, Q. Wang, J.
R. Cole, J. M. Tiedje, Y. Luo, J. Zhou, Tundra soil carbon is vulnerable to rapid microbial
decomposition under climate warming. *Nat. Clim. Change* **6**, 595–600 (2016).

3.   J. Feng, C. Wang, J. Lei, Y. Yang, Q. Yan, X. Zhou, X. Tao, D. Ning, M. M. Yuan, Y. Qin,
Z. J. Shi, X. Guo, Z. He, J. D. Van Nostrand, L. Wu, R. G. Bracho-Garillo, C. R. Penton, J.
R. Cole, K. T. Konstantinidis, Y. Luo, E. A. G. Schuur, J. M. Tiedje, J. Zhou, Warming-
induced permafrost thaw exacerbates tundra soil carbon decomposition mediated by
microbial community. *Microbiome* **8**, 3 (2020).

4.   Q. Gao, G. Wang, K. Xue, Y. Yang, J. Xie, H. Yu, S. Bai, F. Liu, Z. He, D. Ning, S. E.
Hobbie, P. B. Reich, J. Zhou, Stimulation of soil respiration by elevated CO2 is enhanced
under nitrogen limitation in a decade-long grassland study. *Proc. Natl. Acad. Sci.* **117**,
33317–33324 (2020).

5.   L. A. Kluber, E. R. Johnston, S. A. Allen, J. N. Hendershot, P. J. Hanson, C. W. Schadt,
Constraints on microbial communities, decomposition and methane production in deep
peat deposits. *PLOS ONE* **15**, e0223744 (2020).

6.   K. M. DeAngelis, G. Pold, B. D. Topçuoğlu, L. T. A. van Diepen, R. M. Varney, J. L.
Blanchard, J. Melillo, S. D. Frey, Long-term forest soil warming alters microbial
communities in temperate forest soils. *Front. Microbiol.* **6** (2015).

7.   J. Feng, C. R. Penton, Z. He, J. D. Van Nostrand, M. M. Yuan, L. Wu, C. Wang, Y. Qin, Z.
J. Shi, X. Guo, E. A. G. Schuur, Y. Luo, R. Bracho, K. T. Konstantinidis, J. R. Cole, J. M.
Tiedje, Y. Yang, J. Zhou, Long-Term Warming in Alaska Enlarges the Diazotrophic
Community in Deep Soils. *mBio* **10**, e02521-18 (2019).

8.   J. Seward, M. A. Carson, L. J. Lamit, N. Basiliko, J. B. Yavitt, E. Lilleskov, C. W. Schadt, D.
S. Smith, J. Mclaughlin, N. Mykytczuk, S. Willims-Johnson, N. Roulet, T. Moore, L. Harris,
S. Bräuer, Peatland Microbial Community Composition Is Driven by a Natural Climate
Gradient. *Microb. Ecol.* **80**, 593–602 (2020).

9.   N. C. Dove, M. E. Barnes, K. Moreland, R. C. Graham, A. A. Berhe, S. C. Hart, Depth
dependence of climatic controls on soil microbial community activity and composition.
ISME Commun. **1**, 1–11 (2021).

10. J. Curiel Yuste, A. J. Fernandez-Gonzalez, M. Fernandez-Lopez, R. Ogaya, J. Penuelas,
J. Sardans, F. Lloret, Strong functional stability of soil microbial communities under
semiarid Mediterranean conditions and subjected to long-term shifts in baseline
precipitation. *Soil Biol. Biochem.* **69**, 223–233 (2014).

11. J. L. M. Gutknecht, C. B. Field, T. C. Balser, Microbial communities and their responses to

simulated global change fluctuate greatly over multiple years. *Glob. Change Biol.* **18**,
2256–2269 (2012).

12. K. Zhu, N. R. Chiariello, T. Tobeck, T. Fukami, C. B. Field, Nonlinear, interacting
responses to climate limit grassland production under global change. *Proc. Natl. Acad. Sci.*
113, 10589–10594 (2016).

13. S. Yang, Q. Zheng, M. Yuan, Z. Shi, N. R. Chiariello, K. M. Docherty, S. Dong, C. B. Field,
Y. Gu, J. Gutknecht, B. A. Hungate, X. Le Roux, X. Ma, A. Niboyet, T. Yuan, J. Zhou, Y.
Yang, Long-term elevated CO_2 shifts composition of soil microbial communities in a
Californian annual grassland, reducing growth and N utilization potentials. *Sci. Total*
Environ. **652**, 1474–1481 (2019).

14. A. A. Carrell, M. Kolton, J. B. Glass, D. A. Pelletier, M. J. Warren, J. E. Kostka, C. M.
Iversen, P. J. Hanson, D. J. Weston, Experimental warming alters the community
composition, diversity, and N2 fixation activity of peat moss (Sphagnum fallax)
microbiomes. *Glob. Change Biol.* **25**, 2993–3004 (2019).

15. R. M. Wilson, M. M. Tfaily, M. Kolton, E. R. Johnston, C. Petro, C. A. Zalman, P. J.
Hanson, H. M. Heyman, J. E. Kyle, D. W. Hoyt, E. K. Eder, S. O. Purvine, R. K. Kolka, S.
D. Sebestyen, N. A. Griffiths, C. W. Schadt, J. K. Keller, S. D. Bridgham, J. P. Chanton, J.
E. Kostka, Soil metabolome response to whole-ecosystem warming at the Spruce and
Peatland Responses under Changing Environments experiment. *Proc. Natl. Acad. Sci.*
118, e2004192118 (2021).

16. M. R. Baysinger, R. M. Wilson, P. J. Hanson, J. E. Kostka, J. P. Chanton, Compositional
stability of peat in ecosystem-scale warming mesocosms. *PLOS ONE* **17**, e0263994
(2022).

17. S. D. Allison, J. B. H. Martiny, Resistance, resilience, and redundancy in microbial
communities. *Proc. Natl. Acad. Sci.* **105**, 11512–11519 (2008).

18. A. Shade, H. Peter, S. Allison, D. Baho, M. Berga, H. Buergmann, D. Huber, S.
Langenheder, J. Lennon, J. Martiny, K. Matulich, T. Schmidt, J. Handelsman,
Fundamentals of Microbial Community Resistance and Resilience. *Front. Microbiol.* **3**
(2012).

19. R. Andersen, S. J. Chapman, R. R. E. Artz, Microbial communities in natural and disturbed
peatlands: A review. *Soil Biol. Biochem.* **57**, 979–994 (2013).

20. J. T. Weedon, G. A. Kowalchuk, R. Aerts, S. Freriks, W. F. M. Röling, P. M. van Bodegom,
Compositional Stability of the Bacterial Community in a Climate-Sensitive Sub-Arctic
Peatland. *Front. Microbiol.* **8** (2017).

21. S. Monteux, J. T. Weedon, G. Blume-Werry, K. Gavazov, V. E. J. Jassey, M. Johansson,
F. Keuper, C. Olid, E. Dorrepaal, Long-term in situ permafrost thaw effects on bacterial
communities and potential aerobic respiration. *ISME J.* **12**, 2129–2141 (2018).

22. F. De Vries, A. Shade, Controls on soil microbial community stability under climate change.
Front. Microbiol. **4** (2013).

23.   S. Pimm, The complexity and stability of ecosystems. *Nature* **307**, 321–326 (1984).

24.   J. W. Sorensen, A. Shade, Dormancy dynamics and dispersal contribute to soil
microbiome resilience. doi: https://doi.org/10.1098/rstb.2019.0255 (2020).

25.   B. Worm, J. E. Duffy, Biodiversity, productivity and stability in real food webs. *Trends Ecol.*
Evol. **18**, 628–632 (2003).

26.   C. Chang, J. HilleRisLambers, Integrating succession and community assembly
perspectives. *F1000Research* **5**, 2294 (2016).

27.   M. Sonesson, S. Jonsson, T. Rosswall, B. E. Rydén, The Swedish IBP/PT Tundra Biome
Project Objectives-Planning-Site. *Ecol. Bull.*, 7–25 (1980).

28.   T. Johansson, N. Malmer, P. M. Crill, T. Friborg, J. H. Åkerman, M. Mastepanov, T. R.
Christensen, Decadal vegetation changes in a northern peatland, greenhouse gas fluxes
and net radiative forcing. *Glob. Change Biol.* **12**, 2352–2369 (2006).

29.   R. K. Varner, P. M. Crill, S. Frolking, C. K. McCalley, S. A. Burke, J. P. Chanton, M. E.
Holmes, null null, S. Saleska, M. W. Palace, Permafrost thaw driven changes in hydrology
and vegetation cover increase trace gas emissions and climate forcing in Stordalen Mire
from 1970 to 2014. *Philos. Trans. R. Soc. Math. Phys. Eng. Sci.* **380**, 20210022 (2021).

30.   K. Kujala, M. Seppälä, T. Holappa, Physical properties of peat and palsa formation. *Cold*
Reg. Sci. Technol. **52**, 408–414 (2008).

31.   T. V. Callaghan, F. Bergholm, T. R. Christensen, C. Jonasson, U. Kokfelt, M. Johansson, A
new climate era in the sub-Arctic: Accelerating climate changes and multiple impacts.
Geophys. Res. Lett. **37** (2010).

32.   P. Łakomiec, J. Holst, T. Friborg, P. Crill, N. Rakos, N. Kljun, P.-O. Olsson, L. Eklundh, A.
Persson, J. Rinne, Field-scale CH_4 emission at a subarctic mire with heterogeneous
permafrost thaw status. *Biogeosciences* **18**, 5811–5830 (2021).

33.   M. Johansson, T. V. Callaghan, J. Bosiö, H. J. Åkerman, M. Jackowicz-Korczynski, T. R.
Christensen, Rapid responses of permafrost and vegetation to experimentally increased
snow cover in sub-arctic Sweden. *Environ. Res. Lett.* **8**, 035025 (2013).

34.   M. E. Holmes, P. M. Crill, W. C. Burnett, C. K. McCalley, R. M. Wilson, S. Frolking, K. -Y.
Chang, W. J. Riley, R. K. Varner, S. B. Hodgkins, IsoGenie Project Coordinators, IsoGenie
Field Team, A. P. McNichol, S. R. Saleska, V. I. Rich, J. P. Chanton, Carbon
Accumulation, Flux, and Fate in Stordalen Mire, a Permafrost Peatland in Transition. *Glob.*
Biogeochem. Cycles **36**, e2021GB007113 (2022).

35.   D. Olefeldt, N. T. Roulet, Effects of permafrost and hydrology on the composition and
transport of dissolved organic carbon in a subarctic peatland complex. *J. Geophys. Res.*
Biogeosciences **117** (2012).

36.   M. Hough, A. McClure, B. Bolduc, E. Dorrepaal, S. Saleska, V. Klepac-Ceraj, V. Rich,
Biotic and Environmental Drivers of Plant Microbiomes Across a Permafrost Thaw
Gradient. *Front. Microbiol.* **11**, 796 (2020).

37. S. Frolking, N. T. Roulet, T. R. Moore, P. J. H. Richard, M. Lavoie, S. D. Muller, Modeling
Northern Peatland Decomposition and Peat Accumulation. *Ecosystems* **4**, 479–498 (2001).

38. D. Pascual, M. Johansson, Increasing impacts of extreme winter warming events on
permafrost. *Weather Clim. Extrem.* **36**, 100450 (2022).

39. B. F. Thornton, M. Wik, P. M. Crill, Climate-forced changes in available energy and
methane bubbling from subarctic lakes. *Geophys. Res. Lett.* **42**, 1936–1942 (2015).

40. T. H. Bell, T. Bell, Many roads to bacterial generalism. *FEMS Microbiol. Ecol.* **97**, fiaa240
(2020).

41. L. A. Malard, A. Guisan, Into the microbial niche. *Trends Ecol. Evol.* **38**, 936–945 (2023).

42. N. Fierer, M. A. Bradford, R. B. Jackson, TOWARD AN ECOLOGICAL CLASSIFICATION
OF SOIL BACTERIA. *Ecology* **88**, 1354–1364 (2007).

43. A. Barberán, K. S. Ramirez, J. W. Leff, M. A. Bradford, D. H. Wall, N. Fierer, Why are
some microbes more ubiquitous than others? Predicting the habitat breadth of soil
bacteria. *Ecol. Lett.* **17**, 794–802 (2014).

44. S. Louca, M. F. Polz, F. Mazel, M. B. N. Albright, J. A. Huber, M. I. O'Connor, M.
Ackermann, A. S. Hahn, D. S. Srivastava, S. A. Crowe, M. Doebeli, L. W. Parfrey, Function
and functional redundancy in microbial systems. *Nat. Ecol. Evol.* **2**, 936–943 (2018).

45. Y.-J. Chen, P. M. Leung, J. L. Wood, S. K. Bay, P. Hugenholtz, A. J. Kessler, G. Shelley,
D. W. Waite, A. E. Franks, P. L. M. Cook, C. Greening, Metabolic flexibility allows bacterial
habitat generalists to become dominant in a frequently disturbed ecosystem. *ISME J.* **15**,
2986–3004 (2021).

46. M. Mariadassou, S. Pichon, D. Ebert, Microbial ecosystems are dominated by specialist
taxa. *Ecol. Lett.* **18**, 974–982 (2015).

47. E. E. L. Muller, Determining Microbial Niche Breadth in the Environment for Better
Ecosystem Fate Predictions. *mSystems* **4**, e00080-19 (2019).

48. D. Bajic, A. Sanchez, The ecology and evolution of microbial metabolic strategies. *Curr.*
Opin. Biotechnol. **62**, 123–128 (2020).

49. K. T. Konstantinidis, J. M. Tiedje, Trends between gene content and genome size in
prokaryotic species with larger genomes. *Proc. Natl. Acad. Sci.* **101**, 3160–3165 (2004).

50. B. Guieysse, S. Wuertz, Metabolically versatile large-genome prokaryotes. *Curr. Opin.*
Biotechnol. **23**, 467–473 (2012).

51. S. J. Giovannoni, J. Cameron Thrash, B. Temperton, Implications of streamlining theory for
microbial ecology. *ISME J.* **8**, 1553–1565 (2014).

52. D. A. Nielsen, N. Fierer, J. L. Geoghegan, M. R. Gillings, V. Gumerov, J. S. Madin, L.
Moore, I. T. Paulsen, T. B. K. Reddy, S. G. Tetu, M. Westoby, Aerobic bacteria and
archaea tend to have larger and more versatile genomes. *Oikos* **130**, 501–511 (2021).

53.   D. R. Johnson, F. Goldschmidt, E. E. Lilja, M. Ackermann, Metabolic specialization and the
assembly of microbial communities. *ISME J.* **6**, 1985–1991 (2012).

54.   H. Chen, K. Ma, C. Lu, Q. Fu, Y. Qiu, J. Zhao, Y. Huang, Y. Yang, C. W. Schadt, H. Chen,
Functional Redundancy in Soil Microbial Community Based on Metagenomics Across the
Globe. *Front. Microbiol.* **13**, 878978 (2022).

55.   I. Vanwonterghem, P. D. Jensen, K. Rabaey, G. W. Tyson, Genome-centric resolution of
microbial diversity, metabolism and interactions in anaerobic digestion. *Environ. Microbiol.*
18, 3144–3158 (2016).

56.   R. K. Thauer, A.-K. Kaster, H. Seedorf, W. Buckel, R. Hedderich, Methanogenic archaea:
ecologically relevant differences in energy conservation. *Nat. Rev. Microbiol.* **6**, 579–591
(2008).

57.   D. Gupta, K. Chen, S. J. Elliott, D. D. Nayak, MmcA is an electron conduit that facilitates
both intracellular and extracellular electron transport in Methanosarcina acetivorans. *Nat.*
Commun. **15**, 3300 (2024).

58.   B. J. Woodcroft, C. M. Singleton, J. A. Boyd, P. N. Evans, J. B. Emerson, A. A. F. Zayed,
R. D. Hoelzle, T. O. Lamberton, C. K. McCalley, S. B. Hodgkins, R. M. Wilson, S. O.
Purvine, C. D. Nicora, C. Li, S. Frolking, J. P. Chanton, P. M. Crill, S. R. Saleska, V. I.
Rich, G. W. Tyson, Genome-centric view of carbon processing in thawing permafrost.
Nature **560**, 49–54 (2018).

2557

2558