1 **CRISPR targeting of H3K4me3 activates gene expression and unlocks**
2 **centromeric crossover recombination in Arabidopsis**
3

4 Jenia Binenbaum[1,*], Vanda Adamkova[1,*], Hannah Fryer[1], Linhao Xu[1], Nicola
5 Gorringe[1], Piotr Włodzimierz[1,3], Robin Burns[1], Ashot Papikian[2,4], Steven E.
6 Jacobsen[2,5], Ian Henderson[1], C. Jake Harris[1,#]
7

[1] Department of Plant Sciences, University of Cambridge, Cambridge CB2 3EA, UK.
[2] Department of Molecular, Cell and Developmental Biology, University of California,
Los Angeles, CA, USA.
[3] Current Address: Institute of Biochemistry and Biophysics, Polish Academy of
Sciences, Warsaw, Poland.
[4] Current Address: Plant Molecular and Cellular Biology Laboratory, Salk Institute for
Biological Studies, La Jolla, CA 92037, USA.
[5] Howard Hughes Medical Institute, University of California, Los Angeles, CA, USA.
* These authors contributed equally.
[#] For correspondance: cjh92@cam.ac.uk

Abstract:

H3K4me3 is a fundamental and highly conserved chromatin mark across eukaryotes, playing a central role in many genome-related processes, including transcription, maintenance of cell identity, DNA damage repair, and meiotic recombination. However, identifying the causal function of H3K4me3 in these diverse pathways remains a challenge, and we lack the tools to manipulate it for agricultural benefit. Here we use the CRISPR-based SunTag system to direct H3K4me3 methyltransferases in the model plant, *Arabidopsis thaliana*. Targeting of SunTag-SDG2 activates the expression of the endogenous reporter gene, *FWA*. We show that SunTag-SDG2 can be employed to increase pathogen resistance by targeting the H3K4me3-dependent disease resistance gene, *SNC1*. Meiotic crossover recombination rates impose a limit on the speed with which new traits can be transferred to elite crop varieties. We demonstrate that targeting of SunTag-SDG2 to low recombining centromeric regions can significantly stimulate crossover formation. Finally, we reveal that the effect is not specific to SDG2 and is likely dependent on the H3K4me3 mark itself, as the orthogonal mammalian-derived H3K4me3 methyltransferase, PRDM9, produces a similar effect on gene expression with reduced off-target potential. Overall, our study supports an instructive role for H3K4me3 in transcription and meiotic recombination and opens the door to precise modulation of important agricultural traits.

Introduction:

Re-programming of transcriptional landscapes in response to developmental and environmental cues critically underlies a plant's ability to adapt and optimise their performance. The potential to re-wire and fine-tune expression of endogenous genes therefore holds great promise for agricultural improvement. Chromatin modifications associated with gene expression play an essential role in this process (Lloyd and Lister, 2021).

Trimethylation of histone 3 lysine 4 (H3K4me3) is a well-studied and highly conserved chromatin mark. It's enrichment around the transcriptional start site of expressed genes is observed across eukaryotes (Schwaiger et al., 2014), and is thought to assist in transcriptional maintenance and memory of cell identity (Wang and Helin, 2024). Despite the ubiquity of H3K4me3 and its positive correlation with gene expression, whether it plays an instructive role in transcription remains a matter of debate, due in part to the limited evidence of transcriptional impact when H3K4me3 levels are perturbed (Francisco et al., 2023; Howe et al., 2017). In animals, TAF3 directly binds to H3K4me3 and recruits basal transcription factor machinery, providing a plausible mechanism to stimulate transcription (Lauberth et al., 2013; Vermeulen et al., 2007). Recent work in embryonic stem cells shows that H3K4me3 plays a role in RNA Polymerase II pause-release and elongation, rather than transcriptional initiation (H. Wang et al., 2023). In plants, there are no TAF3 homologues and whether similar mechanisms exist to promote transcription downstream of H3K4me3 remains unknown.

H3K4me3 also plays a key role in several other genome-related processes. For instance, meiotic recombination, which allows for genetic exchange between parental chromosomes (Borde et al., 2009). This process is essential for breeders to transfer traits from wild relatives into elite cultivars. However, the positioning of meiotic recombination events is non-uniform across the genome, with gene-rich chromosomal arms having relatively higher rates of recombination, and centromeric repeat-rich regions being more recombination supressed (Rowan et al., 2019; Tock et al., 2021). Loci encoding desirable traits that reside within these low-recombining regions of the genome are largely intractable to selection by traditional breeding approaches.

Further, in mammals, the H3K4me3 SET domain methyltransferase containing
protein, PRDM9, helps to define active sites of crossover recombination (Powers et
al., 2016). While H3K4me3 levels are positively correlated with recombination rates in
plants (Choi et al., 2013; Rowan et al., 2019), whether deposition of H3K4me3 is
sufficient to promote crossover formation is entirely unknown.

One way to reveal the causal role of an epigenetic modification is to directly deposit
the mark at targeted genomic locations (Dubois and Roudier, 2021; Harris et al., 2023;
Policarpi and Hackett, 2021; Villiger et al., 2024; Wang and Yamaguchi, 2024).
Epigenome editing tools, such as the CRISPR-based SunTag system (Tanenbaum et
al., 2014), have proven to be highly efficient at targeting DNA methylation addition,
removal and histone modification (Gallego-Bartolomé et al., 2018; Papikian et al.,
2019; Pflueger et al., 2018; Policarpi et al., 2024; M. Wang et al., 2023), although their
utility for modulating agronomically useful traits is only just beginning to be realised
(Selma and Orzáez, 2021), and is limited by the range of possible marks that can be
directed (Lloyd and Lister, 2021). In the model plant *Arabidopsis thaliana*, SDG2 is the
main histone methyltransferase responsible for deposition of H3K4me3 (Guo et al.,
2010). Here, we describe CRISPR directed H3K4me3 deposition in plants and
demonstrate its utility in a range of scenarios. We translationally fuse the SDG2
methyltransferase domain to the scFv single chain antibody, which can be recruited to
the SunTag epitope chain connected to dead Cas9 (dCas9). We show that SunTag-
SDG2 mediated H3K4me3 deposition can activate the expression of endogenous
genes. Further, by targeting a disease resistance gene, we generate plants with
enhance resistance capabilities. Next, we show that SunTag-SDG2 can unlock meiotic
crossover recombination across centromeric regions. Finally, we incorporate the
methyltransferase domain from PRDM9 into SunTag – which was recently shown to
instruct transcription in a mammalian context (Policarpi et al., 2024) – showing that it
is also highly efficient for transcriptional activation in plants, and displays reduced off-
target potential. These results demonstrate the power and versatility of a functional
H3K4me3 targeting system for modulation of critical genome templated processes,
such as transcriptional enhancement and meiotic recombination.

Results:

SunTag SDG2 mediates transcriptional activation of FWA

The SunTag system we chose is composed of three components (Gardiner et al., 2022; Tanenbaum et al., 2014) (Figure 1A): 1) A catalytically deactivated (nuclease deficient) Cas9 (dCas9) translationally fused to a tail with 10x copies of the GCN4 epitope, each separated by a 22 amino acid flexible linker. 2) An effector module that encodes a single chain fragment variable (scFv) antibody that recognises GCN4, super folder GFP (sfGFP), and the effector of interest (e.g. enzymatic modifier / recruitment scaffold). 3) *U6* promoter driven CRISPR guide RNA. When the three components are expressed within the same cell, the guide RNA recruits the dCas9-10xGCN4 to the target locus of interest, while the GCN4 single chain antibody containing effector module binds to the dCas9-10xGCN4 epitope tail. The system allows for conformational flexibility and the potential for high stochiometric concentration of the effector at the locus of interest.

We inserted the C-terminal coding sequence of the H3K4 methyltransferase, SDG2 (amino acids 1571-2335), into the SunTag system (Figure 1B, Supp. Fig. S1). As a control, we also generated a version of SDG2 with an amino acid change (Y1903F) predicted to abolish catalytic activity (Supp. Fig. S2) (Du et al., 2014; Guo et al., 2010). Hereafter, we refer to these as SDG2 and dSDG2 (for catalytically deactivated). We targeted the SunTag system to the promoter of the epigenetically repressed *FWA* gene (Soppe et al., 2000), using the previously published CRISPR guide RNA 4, which is complementary to two tandem repeat regions directly upstream of the *FWA* transcription start site (TSS) (Gallego-Bartolomé et al., 2018). These were transformed into the *rdr6* background, as previous results have shown that SunTag is less prone to silencing and achieves higher gene expression (M. Wang et al., 2023)*.* We observed that SunTag:SDG2:FWA_g4 activated *FWA* mRNA expression, while the SunTag:dSDG2:FWA_g4 and no guide (SunTag:SDG2:No_g) RNA controls did not (Fig 1C, Supp. Fig. 1C). Importantly, effector module expression was detected in all constructs, indicating that both guide RNA targeting and catalytic activity are required for *FWA* activation (Sup Fig 3A). To validate the presence of the SunTag and test for the deposition of H3K4me3, we performed ChIP qPCR and observed significant

enrichment both for the presence of SunTag and for H3K4me3 accumulation at the
target site in these transgenic lines (Figure 1D and Sup Fig 3B), confirming that
SunTag-SDG2 binds the *FWA* locus and deposits H3K4me3. To investigate
SunTag:SDG2:FWA_g4 binding genome-wide, we performed high-throughput
sequencing (anti-HA ChIP-seq), which confirmed the previously reported high
specificity of SunTag targeting (Papikian et al., 2019), with the *FWA* target locus as
the top binding target (Figure 1E, Supp. Fig. 4). Only 3 conserved binding sites were
identified in the SunTag:SDG2:FWA_g4 lines; *FWA* and two off-target sites previously
identified as having sequence-similarity to FWA_g4 (Papikian et al., 2019).
Furthermore, transcriptome analyses confirmed *FWA* as the highest fold change
differentially expressed upregulated gene (Figure 1F). Together, these data indicate
that deposition of H3K4me3 by SunTag:SDG2 is sufficient to overcome epigenetic
silencing and drive transcriptional activation in a locus-specific manner.

Figure 1

Figure 1: SunTag:SDG2 activates *FWA* mRNA expression.

A) Schematic showing the SunTag system for modulation of chromatin (Gardiner et al., 2022; Papikian et al., 2019). B) Upper panel shows SDG2 with catalytic SET domain (drawn approximately to scale). Lower left depicts predicted alignment error (PAE) plot from AlphaFold3 (Abramson et al., 2024) prediction of the SDG2 coding sequence with a box drawn around the region cloned into SunTag (amino acids 1571-2335). Lower right depicts the AlphaFold3 model for $SDG2_{1571-2335}$, with the position of the amino acid change (Y1903F) for dSDG2 indicated (shown in red). C) RT-qPCR for *FWA* expression from the genotypes indicated. Error bars represent SEM from three biological replicates. *** indicates p-value

<0.001 (students T-test). D) ChIP qPCR for the presence of SunTag (left panel) and H3K4me3

enrichment (right panel) over the TSS of *FWA*. The control region (AT5G65130) used for

normalisation has high H3K4me3. Error bars represent SEM from 2 biological replicates. E)

Genome browser image showing the enrichment of two independent SunTag:SDG2:FWA_g4

lines (2 biological replicates each, bio replicate tracks are overlaid) at the *FWA* TSS. F) MA

plot comparing the transcriptome of SunTag:SDG2:FWA_g4 as compared to the non-

transformed control (*rdr6*). Differentially expressed genes are show in in red (FDR < 0.01),

with *FWA* labelled and enlarged for visibility.

SNC1 mediated modulation of disease resistance

Given that SunTag:SDG2 can modulate chromatin and activate gene expression, we

were interested use the tool to modify an important plant trait. Gene expression during

pathogen attack is a key factor in determining disease outcome. *SNC1* is disease

resistance gene, encoding a nucleotide-binding leucine-rich repeat protein (M. Y.

Wang et al., 2023) that sits within the partially epigenetically repressed *RPP5* gene

cluster (Yi and Richards, 2009). *SNC1* expression activity is modulated by H3K4me3

(Yang et al., 2023), and its level of expression is positively correlated with basal

resistance (Yang et al., 2023; Yi and Richards, 2009). Therefore, *SNC1* is an attractive

target for proof-of-principle modulation of disease resistance by epigenome

engineering.

We designed a guide RNA to target the promoter region of *SNC1* directly upstream of

the TSS and inserted it into our SunTag:SDG2 and SunTag:dSDG2 constructs (Figure

2A). As high pathogen resistance is generally associated with dwarfed plants, we

inspected the lines for growth and developmental phenotypes. In our *SNC1* targeting

SunTag:SDG2:SNC1_g4, but not our SunTag:dSDG2:SNC1_g4 controls, we

observed small, dwarfed phenotype with reduced rosette diameter (Figure 2AB).

These plant phenotypes resembled those of our positive control *bal* lines, which

overexpress *SNC1*, resulting in the small dwarfed plant phenotype and high basal

disease resistance (Yi and Richards, 2009). As the phenotype is consistent with

upregulation of *SNC1* by SunTag, we confirmed the increased expression and

successful targeting of SunTag:SDG2 to the *SNC1* locus (Figure 2CD).

To assess whether these lines show increased pathogen resistance, we challenged
the plants with the generalist plant pathogen, *Pseudomonas syringae* pathovar *tomato*
strain DC3000 (*Pst*). The recently described bioluminescent *Pst* (*Pst::LUX*) was used
to allow for non-destructive quantification of pathogen colonisation (Furci et al., 2021).
We developed a high-throughput assay system, inoculating 7-day-old seedlings with
Pst::LUX and imaging to quantify the levels of *Pst::LUX* at 24 hour intervals over the
course of infection. We challenged a range of control genotypes that are known to be
Pst hyper-resistant (*bal*), constitutively primed (*edr1*) and susceptible (*NahG*),
recapitulating the expected pattern of *Pst::LUX* colonisation dynamics in these
backgrounds (Supp. Fig. S5) (Furci et al., 2021). Next, we challenged our *SNC1*
targeting lines with *Pst::LUX*, finding that SunTag:SDG2:SNC1_g4 lines displayed
hyper resistance that was nearly identical in magnitude to that of the *SNC1*
overexpressing positive control, *bal* (Figure 2E). The results demonstrate that
epigenome-engineering of a single defense gene, *SNC1*, is sufficient to generate
plants with improved disease resistance phenotypes.

Figure 2

Figure 2: SunTag:SDG2 targeting to *SNC1* enhances resistance to *P. syringae*.

A) Upper, schematic of the *SNC1* gene, with the red line indicating the *SNC1* gRNA target site, drawn to scale. Lower, representative images of 3-week old plants from the genotypes indicated. B) Rosette size quantification. Each dot represents an individual plant. Error bars represent SEM. Different letters indicate significant difference by ANOVA with post-hoc Tukey HSD. C) ChIP qPCR for the presence of SunTag at SNC1. D) RT-qPCR for *SNC1* expression. Error bars represent SEM. * indicates p-value < 0.05. E) Pst::LUX assay for colonisation

216 quantification at 3 days post inoculation. Different letters indicate significant difference by
217 ANOVA with post-hoc Tukey HSD.

219 Centromeric targeting of SDG2 drives increased meiotic crossover
220 recombination

Beyond transcription, H3K4me3 is associated with many other fundamental genome
related processes including meiotic recombination. In plants, meiotic crossover
recombination rates broadly correlate with epigenetic territories and are non-uniform
across the genome (Rowan et al., 2019; Tock et al., 2021). In *A. thaliana*,
recombination occurs at a relatively high frequency in euchromatic arms, it is
significantly reduced over the more heterochromatic pericentromeric regions, and
drops to virtually zero over the centromere (Naish et al., 2021; Yelina et al., 2012).
Crossover hotspots are often found to be high at gene promoters where H3K4me3 is
enriched (Choi et al., 2013). In previous work, we showed that gene-associated
crossover recombination hotspots can be supressed by small RNA directed DNA
methylation (Yelina et al., 2015). Therefore, we hypothesised that recombination within
the centromeric regions might be increased by SunTag mediated epigenetic activation
via deposition of H3K4me3.

The Arabidopsis centromeres are composed of highly repetitive *CEN178* satellite
repeat arrays (Naish et al., 2021; Wlodzimierz et al., 2023). We designed a guide RNA
that perfectly matches to 250 individual *CEN178* repeats within the centromeric region
of chromosome 3 (LRCen3_g), and inserted this guide into the SunTag:SDG2 system.
As centromere-proximal crossover recombination events are relatively rare, to
quantitatively assess the impact on recombination rate over centromere 3, we crossed
(non-transgenic) Col-0 plants with *CTL3.9* (Wu et al., 2015), which is a transgenic a
line that encodes red and green fluorescent markers flanking *CEN3*. The red and
green T-DNA insertions span a region of approximately 9Mb (positions 9.7Mb and
18.8Mb, respectively, on the Col-CEN assembly) of chromosome 3, while the
centromeric satellite arrays are located from 13.6-15.7Mb (Figure 3A). Meiotic
recombination events that occur within this region result in progeny inheriting either
red or green fluorescent markers, the frequency of which can be measured by
automated seed imaging (Kbiri et al., 2022). We transformed *CTL3.9* F_1 double

hemizygous plants with the SunTag:SDG2:LRCen3_g construct (Figure 3AB). The F_2

plants represented individual transformants of the SunTag (T_1 for SunTag), and so F_3

seeds from individual F_2 plants were imaged to measure the rate of recombination, in

centiMorgans, over the centromere 3 spanning region of *CTL3.9* (Fernandes et al.,

2024).

Remarkably, the SunTag:SDG2:LRCen3_g containing lines showed significantly

elevated crossover recombination within the *CTL3.9* interval, which was not observed

in the no guide RNA controls (ANOVA with post-hoc Tukey HSD cutoff <0.05) (Figure

3C, Supp. Fig. 6). Some individual transgenic plants displayed a substantial increase

in *CTL3.9* crossover recombination rate, over 50% higher as compared the average

in non-transgenics, across this megabase-scale centromere-spanning region.

Progeny from the most highly recombining line that also showed Mendelian

inheritance of the *CTL3.9* T-DNAs was sown to assess whether the effect could persist

in the subsequent (F_4) generation. The plants retained a significantly elevated

crossover frequency across *CTL3.9* in the next generation, suggesting that the

SunTag:SDG2:LRCen3_g effect is relatively stable in the presence of the transgene

(Figure 3C).

To examine the distribution of SunTag:SDG2:LRCen3_g in the high recombination

lines (P4), we performed ChIP-seq (anti-HA) comparing progeny from sibling plants

that either have (+SunTag) or do not have (-SunTag) the SunTag machinery, as well

as non-transgenic (Col-0) controls. Chromosome-wide plots at 100 kb resolution

showed that SunTag:SDG2:LRCen3_g is enriched over the entire centromeric region

of chromosome 3 (Figure 3D). SunTag was also enriched over the other four *A.*

thaliana centromeres (Supp. Fig. 7). As these regions are highly repetitive, we

reasoned that the LRCen3 guide RNA might additionally bind the other centromeres

due to mismatch tolerance. Indeed, while the LRCen3 guide RNA aligns perfectly to

250 loci within the centromere of chromosome 3, by allowing a single mismatch the

guide is predicted to bind regions in all 5 centromeres, which is consistent with the

ChIP-seq data observed (Supp. Fig. 8). Having confirmed that SunTag is enriched

within the centromeric regions in these lines, we were next interested to examine

levels of H3K4me3. ChIP-seq of H3K4me3 showed that levels were broadly increased

throughout the genome (Figure 3D). This increase was stronger in centromeric regions

as compared to euchromatic arms, as expected, and was most prominent over the pericentromere. As the markers on *CTL3.9* span the centromere and adjacent pericentromeric regions (Fernandes et al., 2024) (Figure 3D), this is consistent with the significantly elevated crossover recombination rate observed. Overall, the results demonstrate that centromerically targeted SunTag-SDG2 can elevate the meiotic crossover potential of these typically recombination-suppressed regions.

Figure 3

Figure 3: Centromeric targeting of SunTag:SDG2 elevates meiotic crossover recombination rate.

A) Region of chromosome 3, showing CENH3 ChIP-seq enrichment (Wlodzimierz et al., 2023) with *CTL3.9* (Wu et al., 2015) red / green T-DNA marker positions and LRCen3 guide RNA binding sites indicated. B) Left panel, representative image of red and green fluorescing seeds from a *CTL3.9* double homozygous line. Right panel, schematic of showing how crossovers are identified within *CTL3.9*. C) Boxplot showing crossover recombination frequency over *CTL3.9* in centiMorgans. Control data is from Col-0 (non-transgenic) crossed to *CTL3.9*.

Different letters indicate significant difference by ANOVA with post-hoc Tukey HSD. The red dot indicates the seed set line taken to the next generation (F_4). D) Chromosome-wide plots of chromosome 3. Upper and middle panels show enrichment of H3K4me3 and SunTag (anti-HA) by ChIP-seq in sibling lines with / without SunTag (+/-). Enrichment is calculated as log_2 fold-change over non-transgenic (Col-0) controls in 100 kb windows. The lower panel depicts an LRCen3 guide RNA binding site density over chromosome 3. *CTL3.9* marker positions are indicated, and centromeric regions are shown in grey. All data are mapped the Col-CEN genome assembly (Naish et al., 2021).

Mammalian PRDM9 drives efficient transcriptional activation in Arabidopsis

As the enrichment of H3K4me3 in SunTag:SDG2:LRCen3_g was not exclusive to the guide RNA targeted centromeric regions, we reasoned that the SunTag system may afford some level of off-target activity due to non-guide RNA directed ectopic interactions. Consistent with this observation, we also noticed that our SunTag:SDG2 no guide RNA controls frequently exhibited pleotropic developmental phenotypes (Supp. Fig. S9). To quantify global levels of H3K4me3, we performed bulk histone westerns, comparing no guide RNA lines to successful *FWA* expression-activating guide RNA containing lines. Importantly, the SDG2 effector itself is expressed to similar levels as in the *FWA* targeting lines (Supp. Fig. S3). The SunTag:SDG2 no guide lines all showed significant (3-4 fold) increases in global levels of H3K4me3, as compared to the *FWA* guide containing lines (Supp. Fig. 10A). This indicates that the presence of the guide RNA may enhance the targeting specificity of the SunTag complex by reducing the frequency of non-specific interactions that result in off-target deposition of H3K4me3. We also noticed that a small number of SunTag:dSDG2:FWA_g4 lines could activate *FWA*, but this effect was limited to lines where the dSDG2 effector was massively overexpressed (Supp. Fig. 10B). This suggests that either dSDG2 retains some residual catalytic activity, or that it can recruit endogenous machinery to initiate transcription.

In order to reduce off-target activity and the potential for co-recruitment of endogenous complexes, we replaced SDG2 with an orthogonal effector, PRDM9. PRDM9 is a mammalian derived H3K4me3 methyltransferase, and Policarpi et al (2024) recently showed that the catalytic domain could be incorporated into the SunTag system for

targeting of H3K4me3 mammalian cells. Therefore, we inserted the PRDM9 methyltransferase domain, as well as the catalytically deactivated version, into our SunTag system and targeted *FWA* (Figure 4A). Strikingly, SunTag:PRDM9:FWA_g4 was highly effective at activating *FWA* mRNA expression (Figure 4B), (23/32 in the T_1 generation, 72%), while none of the dPRDM9 (0/16) and no-guide RNA (0/19) versions were able to do so (Supp. Fig. 11). Importantly, we could detect robust expression of the effector in all constructs (Supp. Fig. 11), and ChIP qPCR confirmed the presence of SunTag and the enrichment of H3K4me3 at *FWA* (Fig 4C). At the transcriptome level, the effect of SunTag:PRDM9:FWA_g4 was highly specific, with only 19 genes identified as differentially expressed (15 up, 4 down, FDR<0.01), and with *FWA* itself having the highest fold change among them (>158 fold change over *rdr6*) (Figure 4D). Unlike the no-guide SDG2 lines, we did not observe any developmental defects in the PRDM9 no-guides, and correspondingly, we observed only minor transcriptional defects when comparing the no-guide lines to non-transformed *rdr6* controls (Supp. Fig. S12). Finally, we performed ChIP-seq in these lines, confirming *FWA* as the most significant and enriched peak for SunTag binding and revealing deposition of H3K4me3 at the transcriptional start site of *FWA* (Fig 4E). Overall, the results demonstrate that PRDM9 is highly effective for deposition of H3K4me3 in plant as well as mammalian SunTag systems. Furthermore, as PRDM9 is not likely to have endogenous partners in the plant nucleus, the results provide strong evidence in favour of a direct role for H3K4me3 in transcriptional stimulation.

Figure 4

354
355
Figure 4: SunTag PRDM9 is sufficient for activation of *FWA*.
A) Upper panel shows PRDM9 with catalytic SET domain (drawn approximately to scale).
Lower left depicts predicted alignment error (PAE) plot from AlphaFold3 prediction of the
PRDM9 coding sequence with a box drawn around the region cloned into SunTag (amino
acids 110-417). Lower right depicts the AlphaFold3 model for PRDM9$_{110-417}$, with the amino
acid change for dPRDM9 indicated (position shown in red). B) RT-qPCR for *FWA* (upper
panel) and the effector module (sfGFP, lower panel) for the genotypes indicated. Two

independent T$_3$ lines are used per genotype. Error bars represent SEM. C) ChIP qPCR for the presence of SunTag (left panel) and H3K4me3 enrichment (right panel) over the TSS of *FWA*. Error bars represent SEM from 2 biological replicates. D) MA plot comparing the transcriptome of SunTag:PRDM9:FWA_g4 as compared to the non-transformed control (*rdr6*). Differentially expressed genes are show in in red (FDR < 0.01), with *FWA* labelled and enlarged for visibility. E) Genome browser image showing efficient SunTag targeting and H3K4me3 enrichment at *FWA*. Biological replicate tracks from independent lines are overlaid.

Discussion:

Here we show that SunTag mediated targeting of H3K4me3 facilitates control of gene expression states, can allow for improved pathogen resistance capabilities, and can drive targeted increases in centromere-proximal recombination. From a fundamental biological perspective, our data provide strong support in favour of an instructive role for H3K4me3 in transcription, as we show that site-specific deposition of H3K4me3 results in activation of mRNA expression. As the effect is dependent on the presence of a functional catalytic domain, and is observed using both native and non-native methyltransferases, co-recruitment of endogenous complexes can be likely ruled out as an explanatory factor in driving this transcriptional enhancement. Furthermore, we reveal H3K4me3 targeting as a novel strategy for stimulating crossover formation in low-recombining, repeat-rich regions of the genome. The results open the door to non-GM approaches for precise genome engineering and rational design of agriculturally relevant traits.

Materials & Methods

Plant materials and growth conditions:

All *Arabidopsis thaliana* lines are in a Col-0 ecotype background. Plants grown on plates were sown on to 1/2x Murashige-Skoog (MS) medium, 1% sucrose and 0.8% agar (pH 5.7), stratified for 2 d at 4 °C in the dark, then transferred to growth chambers (Percival CU-41L4D) at 21 °C under long day light conditions (16 hour light/8 hour dark). Plants grown on soil were grown in F2 soil at 20°C under long day light conditions (16 hour light/8 hour dark). Lines: *rdr6-15* (SAIL_617_H07), and *fwa* lines were previously described (M. Wang et al., 2023). NahG and *edr1* were kindly provided by Prof. Juriaan Ton, previously described in (Furci et al., 2021). *bal* lines were previously described in (Yi and Richards, 2009). The CTL 3.9 line was previously described in (Fernandes et al., 2024; Wu et al., 2015). All transgenic plants were generated by *Agrobacterium tumefaciens* strain AGL1 using floral dip (Clough and Bent, 1998).

Plasmid construction:

To generate the SunTag:SDG2:FWA_g4 construct: the CDS of the catalytic domain of SDG2 (from amino acids 1571-2335) was amplified and coned into into the BsiWI linearised SunTag construct (previously described (Papikian et al., 2019)), using In-Fusion (Takara). To generate dSDG2: the conserved amino acid predicted to abrogate catalytic activity when mutated (Y1903F) was identified using ConSurf (Ashkenazy et al., 2016). Overlapping PCR was used to generate a version of SDG2 containing the desired mutation (see Supplementary Table 1), and the amplified PCR product was cloned into a BsiWI linearized SunTag-SDG2 construct by In-Fusion (Takara). To generate SunTag:PRDM9, the catalytic domain of mouse PRDM9 (110-417aa) (described in Policarpi et al 2024) was synthesised (TWIST Bioscience) and cloned into the SunTag using the same BsiWI linearization and In-Fusion method described above. Subsequently, overlap PCR was used to generate a version of PRDM9 with the G282A mutation (see Supplementary Table 1). The SNC1, LRCen3, and the no guides controls, were cloned using the protocol described in (Ghoshal and Gardiner, 2021) (see Supplementary Table 1).

Chromatin Immunoprecipitation:

The chromatin immunoprecipitation experiments were performed as previously
described (Harris et al., 2024). 2g of 14-day-old seedlings were collected per biological
replicate. For SunTag-SDG2 FWA and SunTag-SDG2 SNC1 input material are pooled
T_2 plants. For the SunTag:PRDM9:FWA_g4 in *rdr6*, 2 independent T_3 lines expressing
FWA, 2 independent T_3 lines NoG were used alongside *rdr6* controls. 7 µl of anti-HA
(3F10, Merck) and 5 µl of anti-H3K4me3 (ab8580, abcam) were added to up to 2 ml
of sheared chromatin per sample for pulldown. The ChIP purified DNA was directly
used for ChIP-qPCR and ChIP-seq library preparation. Primer information is listed in
Suppletory Table 1. Values were normalised to Input $(2^{-(\text{ChIP Ct} - \text{Input Ct})}*100)$. ChIP-seq
libraries were generated using NuGen Ovation Ultra Low System V2 kits according to
manufacturer's instructions and were sequenced on an Illumina NovaSeq X
instrument with 150bp paired-end reads.
432
<u>Crossover measurement:</u>
For the recombination rate experiment, SunTag:SDG2:LRCen3_g/No_g plamids were
transformed into F_1 plants derived from a cross between Col-0 and *CTL3.9* (double
homozygous for the seed coat fluorescent markers) (Wu et al., 2015). 70-90 individual
T_1 transgenic plants were recovered per construct after selection on hygromycin (F_2
generation). Seeds derived from individual T_1s (F_3) were imaged and scored for
recombination rate using the *SeedScoring* CellProfiler pipeline, previously described
(Fernandes et al., 2024; Kbiri et al., 2022) with approximately 1,000 seeds per
biological sample. Note, seed sets were pre-screened to remove lines in which either
of the fluorescent markers had become homozygous (presence / absence) in the F_2
generation. For the F_4 data, seeds derived from the most highly recombining and non-
distorted line (P4) were sown (F_3), grown for a generation and seeds from individual
plants were collected (F_4) for recombination rate analysis as above, again filtering
seed sets from which either of the fluorescent markers had become homozygous
(presence / absence) in the previous generation.
448
<u>Western blots:</u>
Western blot were performed as previously described (Ichino et al., 2021) with minor
modifications described below. For detection of the large dCAS9-10xGCN4 and SDG2
effector components as shown in Fig S1, single leaf punch (1cm diameter) samples
from individual plants were used as input. Tissue was ground in a Tissuelyser II

(Retsch) and resuspended in 100 ul of a 60:40 ratio of 2XSDS:8M Urea, mixed, and

boiled at 95 degrees C for 5 minutes. Samples were centrifuged at >10,000xg to

remove debris and loaded into a 3-8% tris acetate gel using MOPS running buffer.

After electrophoresis, protein was transferred from the gel to a methanol activated

PVDF membrane in Tobin Buffer (25mM Tris, 192mM Glycine, 20% methanol, 0.035%

SDS). Membranes were blocked in 5% milk (w/v) in PBST (Phosphate Buffered Saline,

0.1% Tween20) and then incubated with HRP conjugated anti-HA at a 1:3,000 dilution

in block. The blot was imaged by ECL detection.

For the western blots used to detect levels of H3K4me3, 100 mg of 7-day-old seedlings

were frozen in liquid nitrogen and ground using a Tissuelyser II (Retsch). Total

histones were extracted using the The EpiQuik™ Total Histone Extraction Kit

(EpigenTek) according to the manufacturers instructions. 5 µl of sample for H3 and 20

µl of sample for H3K4me3 were loaded onto a NuPage 4-12% BIS-Tris gel (Invitrogen)

in NuPage MOPS SDS running buffer (Invitrogen). Transfer from gel to PVDF

membrane was performed using an iBlot 3 machine with the dedicated transfer stack

(Invitrogen). Following transfer, membrane was blocked in 5% milk (w/v) in TBST (Tris

Buffered Saline, 0.1% Tween20) and shaken in room temp for 1 hour. Blocking buffer

was discarded and membranes were incubated in primary antibody (anti-H3, ab1791

and anti-H3K4me3, ab8580, Abcam) in blocking buffer (1:1,000) shaking overnight in

4C. Following morning membrane was rinsed in TBST 5 times for 5 minutes in room

temp. Secondary antibody was added for 1 hour (Goat anti-mouse HRP 1:10,000 in

blocking buffer, ab6789, Abcam). Membrane was washed again 5 times for 5 minutes.

Detection was performed using a SuperSignal West Pico chemiluminescent substrate

(#34080, Thermo) using an ImageQuant 800 (Amersham).

<u>Fluorescent imaging:</u>

Roots of 7-day-old plants grown in vertical 1/2x MS solid media plates were imaged

using a Leica DM6000B epifluorescent microscope.

<u>Rosette size surface area measurement:</u>

3-week-old plants grown on soil were imaged using a Google Pixel 7 mounted on a

tripod. Images were then analyzed using Fiji.

Pseudomonas syringae colonisation assay:

Infection assays were performed with *P. syringae* pathovar *tomato* strain DC3000 (*Pst*) strain expressing a stable chromosomal insertion of the *lux- CDABE* operon from *Photorhabdus luminescens* (*Pst::LUX*) (Fan et al. 2008), using a slightly modified version of a previously published protocol (Furci et al. 2021). Plants were grown on 1/2X MS solid media supplemented with 200 µg/ml Timentin in 96-well plates (Falcon) for 7 days. Prior inoculation, *Pst::LUX* were grown in King's B media containing 50 µg/ml Kanamycin and 50 µg/ml Rifampicin overnight at 28 °C. Bacteria were pelleted by centrifugation, washed in 10 mM MgSO4 and finally re-suspended to an OD600 of 0.2 in 10 mM MgSO4 containing 0.015% v/v Silwet-L77. 7 days-old seedlings were sprayed with bacteria solution and 96-well plates were sealed with parafilm to maintain 100% relative humidity. Seedlings were imaged at 1-4 days post infection (dpi). Prior imaging using (ImageQuant 800, Amersham), plates were left in the dark for 2 mins. Bacterial bioluminescence images were acquired with 4 mins exposure time and bright field images were taken using OD settings of the ImageQuant 800. Image-based quantification of bioluminescence was carried out in Fiji. For each well, the well outline was obtained from bright field images and added to ROI Manager. These ROIs were then transposed onto the bioluminescence images. The bioluminescence intensity from infected seedlings was obtained using the Fiji functions Analyse, Measure and Mean value.

RT-qPCR:

RNA was extracted from indicated plant material using TRIzol reagent (Invitrogen) and the Direct-Zol RNA MiniPrep (Zymo) kit, including in-column DNase treatment following the manufacturer's instructions. cDNA was synthesized using SuperScript IV (Invitrogen). qRT-PCRs were performed using a Luna Universal qPCR Master Mix (NEB) and a CFX connect Real-time PCR detection system (Bio-rad). For SunTag-SDG2 targeting FWA and SNC1, RNA was extracted from 14-day-old whole seedlings grown on 1/2x MS plates. SNC1 targeted lines were treated with Pst::LUX 4H prior to harvesting. For SunTag:PRDM9 qRT-PCR, RNA was extracted from 4 weeks-old leaf tissue of all T_1 lines and 2 independent T_3 lines per construct. 450 ng of total RNA was used for cDNA synthesis. Primers are listed in Supplementary Table 1.

Quant-Seq Library construction:

For Suntag-SDG2 QuantSeq, 350 ng of total RNA from ~10 pooled T_2 plants were

used as input material. For Suntag:PRDM9 Quant-seq, 350 ng of total RNA from 3 T_1

SunTag:PRDM9:FWA_g4 plants expressing FWA, 3 T_1 no guide SunTag:PRDM9

lines and *rdr6* and plants was used for library preparation. Libraries were prepared

using QuantSeq 3 'mRNA-Seq Library Prep Kit FWD (Lexogen) according to

manufacturer's instructions and sequenced on Illumina NovaSeq 6000 PE150

instrument.

Bioinformatic Analysis:

ChIP-seq analysis:

The ChIP-seq analysis pipeline was performed as previously described with minor

modifications. Bowtie2 (version 2.5.0) was used to map the PE150 bp read data in

fastq format to the TAIR10 genome (--no-unal), and were converted to bam format

using Samtools (version 1.10). Reads were de-duplicated using the samtools fixmate

and markdup commands (see https://github.com/C-Jake-Harris/Binenbaum-SunTag-

H3K4me3 for more details). Tracks were generated in DeepTools (version 3.5.1) using

bamCoverage (--normalizeUsing RPGC, --effectiveGenomeSize 135000000 --binSize

10) with multicopy regions blacklisted (--blackListFileName) using the regions

identified in (Klasfeld et al., 2022). For analysis of reads mapping to the centromere,

a modified version of the previously described pipeline (Wlodzimierz et al., 2023) was

used. Briefly, PE150 bp read data in fastq format were mapped to the Col-CEN

genome (Naish et al., 2021) using Bowtie2 (version 2.2.5) with (--very-sensitive -k 200

--no-unal --no-discordant) and were filtered for primary alignments using samtools

view (-F 256 -q 5) prior to deduplication and track generation, as above. Peaks were

called using MACS2 (version 2.2.9.1) with default parameters. Correlation plots were

generated in DeepTools using MultiBamSummary (--binSize 25) and PlotCorrelation

(-c pearson --removeOutliers --plotNumbers -p heatmap).

QuantSeq Analysis:

In accordance with the manufacturer's (Lexogen) instructions, only read (*1.fq.gz)

from PE150 bp data was used for downstream analysis. Briefly, reads were trimmed

using cutadapt (version 1.18) and were mapped to the genome using STAR (version

2.7.10b) to the TAIR10 genome (--quantMode GeneCounts --alignIntronMax 10000 --

outSAMmultNmax 20). Gene counts were used as input for DESeq2 analysis in R.

Differentially expressed genes were defined using an adjusted p-value (Bonferroni

method) cutoff of <0.01.

<u>gRNA binding analysis:</u>

For the LRCen3 binding and mismatch analysis, the LRCen3 guide RNA sequence

(5'-AGGCTTACAAGATTGGGTTG-3'), was mapped to the Col-CEN genome using

bowtie2 (version 2.2.5). The following options were used for no mismatches (-f -a --

end-to-end --np 0 --score-min L,0,0) and for 1 mismatch (-f -a --end-to-end --score-

min L,0,-1). Mapped reads were sorted and converted to bam format using Samtools

(version 1.10), and subsequently converted to .bed format using the bamtobed

function in bedtools (2.20.1). Bed files were used for downstream analysis in R to

generate chromosome-wide density plots over 100kb regions.

Data availability:

The high-throughput sequencing data generated has been deposited to NCBI

(GSEXXXXXX). Previously published datasets: CENH3 (Wlodzimierz et al., 2023),

H3K4me3 (Xu et al., 2024). Custom scripts used are available on github

(https://github.com/C-Jake-Harris/Binenbaum-SunTag-H3K4me3).

Acknowledgements

We would like to thank Jurriaan Ton's group at the University of Sheffield for their

generous gift of the *Pst::LUX* strain, and for their support in helping to establish the

infection and quantification protocol in the Harris group. **Funding:** C.J.H. is supported

by a Royal Society University Research Fellowship (URF\R1\201016) and an ERC

Starting Grant (TransPlantMemory), funded by UKRI EPSRC (EP/X025306/1). J.B. is

supported by a Marie Curie postdoctoral fellowship, funded by UKRI (EP/Z001749/1).

V.A. is supported by a 4-year BBSRC-funded doctoral studentship. I.R.H is supported

by an ERC Advanced Grant (EvoPanCen). N.G. is supported by a BBSRC-UKRI

doctoral studentship. S.E.J. is an investigator of the Howard Hughes Medical Institute.

Author contributions:

C.J.H., I.R.H. and S.E.J. conceived, designed and supervised the research. C.J.H.

performed experiments, NGS bioinformatic analysis, and assembled the manuscript.

J.B. performed the majority of the experiments relating to SunTag:SDG2. V.A.

performed the experiments relating to SunTag:PRDM9. H.F. performed the majority of
experiments and analysis relating to meiotic recombination. L.X. performed the
SunTag:SDG2 centromere-targeting ChIP-seq. N.G. provided the Col-0 x *CTL3.9* F$_1$
seeds and guidance on the analysis of recombination frequency. P.W. and R.B.
contributed to the design of the centromere targeting guide RNA. A.P. cloned SDG2
into SunTag.
596

References

Abramson J, Adler J, Dunger J, Evans R, Green T, Pritzel A, Ronneberger O, Willmore
L, Ballard AJ, Bambrick J, Bodenstein SW, Evans DA, Hung CC, O'Neill M,
Reiman D, Tunyasuvunakool K, Wu Z, Žemgulytė A, Arvaniti E, Beattie C, Bertolli
O, Bridgland A, Cherepanov A, Congreve M, Cowen-Rivers AI, Cowie A, Figurnov
M, Fuchs FB, Gladman H, Jain R, Khan YA, Low CMR, Perlin K, Potapenko A,
Savy P, Singh S, Stecula A, Thillaisundaram A, Tong C, Yakneen S, Zhong ED,
Zielinski M, Žídek A, Bapst V, Kohli P, Jaderberg M, Hassabis D, Jumper JM.
2024. Accurate structure prediction of biomolecular interactions with AlphaFold 3.
Nature **630**:493–500. doi:10.1038/s41586-024-07487-w
Ashkenazy H, Abadi S, Martz E, Chay O, Mayrose I, Pupko T, Ben-Tal N. 2016.
ConSurf 2016: an improved methodology to estimate and visualize evolutionary
conservation in macromolecules. *Nucleic Acids Res* **44**:W344–W350.
doi:10.1093/NAR/GKW408
Borde V, Robine N, Lin W, Bonfils S, Géli V, Nicolas A. 2009. Histone H3 lysine 4
trimethylation marks meiotic recombination initiation sites. *EMBO J* **28**:99–111.
doi:10.1038/emboj.2008.257
Choi K, Zhao X, Kelly KA, Venn O, Higgins JD, Yelina NE, Hardcastle TJ, Ziolkowski
PA, Copenhaver GP, Franklin FCH, Mcvean G, Henderson IR. 2013. Arabidopsis
meiotic crossover hot spots overlap with H2A.Z nucleosomes at gene promoters.
Nat Genet **45**:1327–1338. doi:10.1038/ng.2766
Clough SJ, Bent AF. 1998. Floral dip: A simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. *Plant J* **16**:735–743. doi:10.1046/j.1365-
313X.1998.00343.x
Du J, Johnson LM, Groth M, Feng S, Hale CJ, Li S, Vashisht AA, Gallego-Bartolome
J, Wohlschlegel JA, Patel DJ, Jacobsen SE. 2014. Mechanism of DNA
methylation-directed histone methylation by KRYPTONITE. *Mol Cell* **55**:495–504.

doi:10.1016/j.molcel.2014.06.009

Dubois A, Roudier F. 2021. Deciphering Plant Chromatin Regulation via
CRISPR/dCas9-Based Epigenome Engineering. *Epigenomes* **5**:17.
doi:10.3390/epigenomes5030017

Fernandes JB, Naish M, Lian Q, Burns R, Tock AJ, Rabanal FA, Wlodzimierz P,
Habring A, Nicholas RE, Weigel D, Mercier R, Henderson IR. 2024. Structural
variation and DNA methylation shape the centromere-proximal meiotic crossover
landscape in Arabidopsis. *Genome Biol* **25**:1–31. doi:10.1186/s13059-024-
03163-4

Francisco M, Id P, Id PS. 2023. Histone methyltransferase activity affects metabolism
in human cells independently of transcriptional regulation.
doi:10.1371/journal.pbio.3002354

Furci L, Pardo DP, Ton J. 2021. A rapid and non - destructive method for spatial –
temporal quantification of colonization by Pseudomonas syringae pv . tomato
DC3000 in Arabidopsis and tomato. *Plant Methods* 1–8. doi:10.1186/s13007-021-
00826-2

Gallego-Bartolomé J, Gardiner J, Liu W, Papikian A, Ghoshal B, Kuo HY, Zhao JM-C,
Segal DJ, Jacobsen SE. 2018. Targeted DNA demethylation of the *Arabidopsis*
genome using the human TET1 catalytic domain. *Proc Natl Acad Sci*
115:201716945. doi:10.1073/pnas.1716945115

Gardiner J, Ghoshal B, Wang M, Jacobsen SE. 2022. CRISPR–Cas-mediated
transcriptional control and epi-mutagenesis. *Plant Physiol* **188**:1811–1824.
doi:10.1093/plphys/kiac033

Ghoshal B, Gardiner J. 2021. CRISPR-dCas9-Based Targeted Manipulation of DNA
Methylation in Plants. *Springer Protoc Handbooks* **2**:57–71.
doi:https://doi.org/10.1007/978-1-0716-1657-4_5

Guo L, Yu Y, Law JA, Zhang X. 2010. Set domain group2 is the major histone H3
lysine 4 trimethyltransferase in Arabidopsis. *Proc Natl Acad Sci U S A* **107**:18557–
18562. doi:10.1073/pnas.1010478107

Harris CJ, Amtmann A, Ton J. 2023. Epigenetic processes in plant stress priming :
Open questions and new approaches. *Curr Opin Plant Biol* **75**:102432.
doi:10.1016/j.pbi.2023.102432

Harris CJ, Zhong Z, Ichino L, Feng S, Jacobsen SE. 2024. H1 restricts euchromatin-
associated methylation pathways from heterochromatic encroachment. *Elife* 1–

17.

Howe FS, Fischl H, Murray SC, Mellor J. 2017. Is H3K4me3 instructive for transcription
activation? *BioEssays* **39**:1–12. doi:10.1002/bies.201600095

Ichino L, Boone BA, Strauskulage L, Harris CJ, Kaur G, Gladstone MA, Tan M, Feng
S, Jami-alahmadi Y, Duttke SH. 2021. MBD5 and MBD6 couple DNA methylation
to gene silencing through the J-domain protein SILENZIO. *Science (80-)*
1439:1434–1439.

Kbiri N, Dluzewska J, Henderson IR, Ziolkowski PA. 2022. Quantifying Meiotic
Crossover Recombination in Arabidopsis Lines Expressing Fluorescent
Reporters in Seeds Using SeedScoring Pipeline for CellProfiler In: Lambing C,
editor. Plant Gametogenesis. Methods in Molecular Biology. New York, NY:
Springer US. pp. 121–134. doi:10.1007/978-1-0716-2253-7_10

Klasfeld S, Roulé T, Wagner D. 2022. Greenscreen: A simple method to remove
artifactual signals and enrich for true peaks in genomic datasets including ChIP-
seq data. *Plant Cell* **34**:4795–4815. doi:10.1093/plcell/koac282

Lauberth SM, Nakayama T, Wu X, Ferris AL, Tang Z, Hughes SH, Roeder RG. 2013.
H3K4me3 interactions with TAF3 regulate preinitiation complex assembly and
selective gene activation. *Cell* **152**:1021–1036. doi:10.1016/j.cell.2013.01.052

Lloyd JPB, Lister R. 2021. Epigenome plasticity in plants. *Nat Rev Genet* **0123456789**.
doi:10.1038/s41576-021-00407-y

Naish M, Alonge M, Wlodzimierz P, Tock AJ, Abramson BW, Schmücker A,
Mandáková T, Jamge B, Lambing C, Kuo P, Yelina N, Hartwick N, Colt K, Smith
LM, Ton J, Kakutani T, Martienssen RA, Schneeberger K, Lysak MA, Berger F,
Bousios A, Michael TP, Schatz MC, Henderson IR. 2021. The genetic and
epigenetic landscape of the Arabidopsis centromeres. *Science (80-)* **374**.
doi:10.1126/science.abi7489

Papikian A, Liu W, Gallego-bartolomé J, Jacobsen SE. 2019. Site-specific
manipulation of Arabidopsis loci using CRISPR-Cas9 SunTag systems. *Nat*
Commun **10**. doi:10.1038/s41467-019-08736-7

Pflueger C, Tan D, Swain T, Nguyen T, Pflueger J, Nefzger C, Polo JM, Ford E, Lister
R. 2018. A modular dCas9-SunTag DNMT3A epigenome editing system
overcomes pervasive off-target activity of direct fusion dCas9-DNMT3A
constructs. *Genome Res* **28**:1193–1206. doi:10.1101/gr.233049.117

Policarpi C, Hackett JA. 2021. Epigenetic editing : Dissecting chromatin function.

BioEssays 1–16.

Policarpi C, Munafò M, Tsagkris S, Carlini V, Hackett JA. 2024. Systematic epigenome
editing captures the context-dependent instructive function of chromatin
modifications. *Nat Genet* **56**:1168–1180. doi:10.1038/s41588-024-01706-w

Powers NR, Parvanov ED, Baker CL, Walker M, Petkov PM, Paigen K. 2016. The
Meiotic Recombination Activator PRDM9 Trimethylates Both H3K36 and H3K4 at
Recombination Hotspots In Vivo. *PLoS Genet* **12**:1–24.
doi:10.1371/journal.pgen.1006146

Rowan BA, Heavens D, Feuerborn TR, Tock AJ, Henderson IR, Weigel D. 2019. An
Ultra High-Density Arabidopsis thaliana Crossover. *Genetics* **213**:771–787.

Schwaiger M, Schönauer A, Rendeiro AF, Pribitzer C, Schauer A, Gilles AF, Schinko
JB, Renfer E, Fredman D, Technau U. 2014. Evolutionary conservation of the
eumetazoan gene regulatory landscape. *Genome Res* **24**:639–650.
doi:10.1101/gr.162529.113

Selma S, Orzáez D. 2021. Perspectives for epigenetic editing in crops. *Transgenic*
Res **30**:381–400. doi:10.1007/s11248-021-00252-z

Soppe WJ, Jacobsen SE, Alonso-Blanco C, Jackson JP, Kakutani T, Koornneef M,
Peeters a J. 2000. The late flowering phenotype of fwa mutants is caused by
gain-of-function epigenetic alleles of a homeodomain gene. *Mol Cell* **6**:791–802.

Tanenbaum ME, Gilbert L a, Qi LS, Weissman JS, Vale RD. 2014. A versatile protein
tagging system for signal amplification in single molecule imaging and gene
regulation. *Cell* **159**:635–646. doi:10.1016/j.cell.2014.09.039

Tock AJ, Holland DM, Jiang W, Osman K, Sanchez-Moran E, Higgins JD, Edwards
KJ, Uauy C, Franklin FCH, Henderson IR. 2021. Crossover-active regions of the
wheat genome are distinguished by DMC1, the chromosome axis, H3K27me3,
and signatures of adaptation. *Genome Res* **31**:1614–1628.
doi:10.1101/gr.273672.120

Vermeulen M, Mulder KW, Denissov S, Pijnappel WWMP, van Schaik FMA, Varier
RA, Baltissen MPA, Stunnenberg HG, Mann M, Timmers HTM. 2007. Selective
Anchoring of TFIID to Nucleosomes by Trimethylation of Histone H3 Lysine 4.
Cell **131**:58–69. doi:10.1016/j.cell.2007.08.016

Villiger L, Joung J, Koblan L, Weissman J, Abudayyeh OO, Gootenberg JS. 2024.
CRISPR technologies for genome, epigenome and transcriptome editing. *Nat Rev*
Mol Cell Biol. doi:10.1038/s41580-023-00697-6

Wang H, Fan Z, Shliaha P V., Miele M, Hendrickson RC, Jiang X, Helin K. 2023.
727   H3K4me3 regulates RNA polymerase II promoter-proximal pause-release.
728   *Nature* **615**:339–348. doi:10.1038/s41586-023-05780-8

729 Wang H, Helin K. 2024. Roles of H3K4 methylation in biology and disease. *Trends*
730   *Cell Biol* **xx**:1–14. doi:10.1016/j.tcb.2024.06.001

731 Wang M, Zhong Z, Gallego-bartolomé J, Li Z, Feng S, Kuo HY, Kan RL, Lam H, Richey
732   JC, Tang L, Zhou J, Liu M, Jami-Alahmadi Y, Wohlschlegel JA, Jacobsen SE.
733   2023. A gene silencing screen uncovers diverse tools for targeted gene
734   repression in Arabidopsis . *Nat Plants*. doi:10.1038/s41477-023-01362-8

735 Wang MY, Chen J Bin, Wu R, Guo HL, Chen Y, Li ZJ, Wei LY, Liu C, He SF, Du M
736   Da, Guo Y long, Peng YL, Jones JDG, Weigel D, Huang JH, Zhu WS. 2023. The
737   plant immune receptor SNC1 monitors helper NLRs targeted by a bacterial
738   effector. *Cell Host Microbe* **31**:1792-1803.e7. doi:10.1016/j.chom.2023.10.006

739 Wang X, Yamaguchi N. 2024. Cause or effect: Probing the roles of epigenetics in plant
740   development and environmental responses. *Curr Opin Plant Biol* **81**:102569.
741   doi:10.1016/j.pbi.2024.102569

742 Wlodzimierz P, Rabanal FA, Burns R, Naish M, Primetis E, Scott A, Mandáková T,
743   Gorringe N, Tock AJ, Holland D, Fritschi K, Habring A, Lanz C, Patel C, Schlegel
744   T, Collenberg M, Mielke M, Nordborg M, Roux F, Shirsekar G, Alonso-Blanco C,
745   Lysak MA, Novikova PY, Bousios A, Weigel D, Henderson IR. 2023. Cycles of
746   satellite and transposon evolution in Arabidopsis centromeres. *Nature* **618**:557–
747   565. doi:10.1038/s41586-023-06062-z

748 Wu G, Rossidivito G, Hu T, Berlyand Y, Poethig RS. 2015. Traffic lines: New tools for
749   genetic analysis in Arabidopsis thaliana. *Genetics* **200**:35–45.
750   doi:10.1534/genetics.114.173435

751 Xu L, Wang Y, Li X, Hu Q, Adamkova V, Xu J, Harris CJ, Ausin I. 2024. The H3K4me3
752   binding ALFIN-LIKE proteins recruit SWR1 for gene-body deposition of H2A.Z.
753   *bioRxiv*. doi:https://doi.org/10.1101/2024.09.12.612642

754 Yang L, Wang Z, Hua J. 2023. Multiple chromatin-associated modules regulate
755   expression of an intracellular immune receptor gene in Arabidopsis. *New Phytol*
756   **237**:2284–2297. doi:10.1111/nph.18672

757 Yelina NE, Choi K, Chelysheva L, Macaulay M, de Snoo B, Wijnker E, Miller N,
758   Drouaud J, Grelon M, Copenhaver GP, Mezard C, Kelly K a, Henderson IR. 2012.
759   Epigenetic remodeling of meiotic crossover frequency in Arabidopsis thaliana

DNA methyltransferase mutants. *PLoS Genet* **8**:e1002844.
doi:10.1371/journal.pgen.1002844

Yelina NE, Lambing C, Hardcastle TJ, Zhao X, Santos B, Henderson IR. 2015. DNA
methylation epigenetically silences crossover hot spots and controls
chromosomal domains of meiotic recombination in Arabidopsis. *Genes Dev*
29:2183–2202. doi:10.1101/gad.270876.115

Yi H, Richards EJ. 2009. Gene duplication and hypermutation of the pathogen
Resistance gene SNC1 in the arabidopsis bal variant. *Genetics* **183**:1227–1234.
doi:10.1534/genetics.109.105569

Figure Legends

Figure 1: SunTag:SDG2 activates *FWA* mRNA expression.

A) Schematic showing the SunTag system for modulation of chromatin (Gardiner et al., 2022; Papikian et al., 2019). B) Upper panel shows SDG2 with catalytic SET domain (drawn approximately to scale). Lower left depicts predicted alignment error (PAE) plot from AlphaFold3 (Abramson et al., 2024) prediction of the SDG2 coding sequence with a box drawn around the region cloned into SunTag (amino acids 1571-2335). Lower right depicts the AlphaFold3 model for $SDG2_{1571-2335}$, with the position of the amino acid change (Y1903F) for dSDG2 indicated (shown in red). C) RT-qPCR for *FWA* expression from the genotypes indicated. Error bars represent SEM from three biological replicates. *** indicates p-value <0.001 (students T-test). D) ChIP qPCR for the presence of SunTag (left panel) and H3K4me3 enrichment (right panel) over the TSS of *FWA*. The control region (AT5G65130) used for normalisation has high H3K4me3. Error bars represent SEM from 2 biological replicates. E) Genome browser image showing the enrichment of two independent SunTag:SDG2:FWA_g4 lines (2 biological replicates each, bio replicate tracks are overlaid) at the *FWA* TSS. F) MA plot comparing the transcriptome of SunTag:SDG2:FWA_g4 as compared to the non-transformed control (*rdr6*). Differentially expressed genes are show in in red (FDR < 0.01), with *FWA* labelled and enlarged for visibility.

Figure 2: SunTag:SDG2 targeting to *SNC1* enhances resistance to *P. syringae*.

A) Upper, schematic of the *SNC1* gene, with the red line indicating the *SNC1* gRNA target site, drawn to scale. Lower, representative images of 3-week old plants from the genotypes indicated. B) Rosette size quantification. Each dot represents an individual plant. Error bars represent SEM. Different letters indicate significant difference by ANOVA with post-hoc Tukey HSD. C) ChIP qPCR for the presence of SunTag at SNC1. D) RT-qPCR for *SNC1* expression. Error bars represent SEM. * indicates p-value < 0.05. E) Pst::LUX assay for colonisation quantification at 3 days post inoculation. Different letters indicate significant difference by ANOVA with post-hoc Tukey HSD.

Figure 3: Centromeric targeting of SunTag:SDG2 elevates meiotic crossover recombination rate.

A) Region of chromosome 3, showing CENH3 ChIP-seq enrichment (Wlodzimierz et al., 2023) with *CTL3.9* (Wu et al., 2015) red / green T-DNA marker positions and LRCen3 guide RNA binding sites indicated. B) Left panel, representative image of red and green fluorescing seeds from a *CTL3.9* double homozygous line. Right panel, schematic of showing how crossovers are identified within *CTL3.9*. C) Boxplot showing crossover recombination frequency over *CTL3.9* in centiMorgans. Control data is from Col-0 (non-transgenic) crossed to *CTL3.9*. Different letters indicate significant difference by ANOVA with post-hoc Tukey HSD. The red dot indicates the seed set line taken to the next generation (F_4). D) Chromosome-wide plots of chromosome 3. Upper and middle panels show enrichment of H3K4me3 and SunTag (anti-HA) by ChIP-seq in sibling lines with / without SunTag (+/-). Enrichment is calculated as $\log_2$ fold-change over non-transgenic (Col-0) controls in 100 kb windows. The lower panel depicts an LRCen3 guide RNA binding site density over chromosome 3. *CTL3.9* marker positions are indicated, and centromeric regions are shown in grey. All data are mapped the Col-CEN genome assembly (Naish et al., 2021).

Figure 4: SunTag PRDM9 is sufficient for activation of *FWA*.

A) Upper panel shows PRDM9 with catalytic SET domain (drawn approximately to scale). Lower left depicts predicted alignment error (PAE) plot from AlphaFold3 prediction of the PRDM9 coding sequence with a box drawn around the region cloned into SunTag (amino acids 110-417). Lower right depicts the AlphaFold3 model for $PRDM9_{110-417}$, with the amino acid change for dPRDM9 indicated (position shown in red). B) RT-qPCR for *FWA* (upper panel) and the effector module (sfGFP, lower panel) for the genotypes indicated. Two independent T_3 lines are used per genotype. Error bars represent SEM. C) ChIP qPCR for the presence of SunTag (left panel) and H3K4me3 enrichment (right panel) over the TSS of *FWA*. Error bars represent SEM from 2 biological replicates. D) MA plot comparing the transcriptome of SunTag:PRDM9:FWA_g4 as compared to the non-transformed control (*rdr6*). Differentially expressed genes are show in in red (FDR < 0.01), with *FWA* labelled and enlarged for visibility. E) Genome browser image showing efficient SunTag targeting and H3K4me3 enrichment at *FWA*. Biological replicate tracks from independent lines are overlaid.

Supplementary Figure 1: A) epifluorescence image of rootstocks from 2-week old
representative plants showing nuclear localisation of sfGFP from SunTag lines. B)
Western blot for the dCas9-10xGCN4 and effector module in SunTag lines showing
activation of FWA expression by RT-qPCR show in C).

Supplementary Figure 2: A) Amino acid alignment between SDG2 and two other well
characterised A. thaliana hisone methyltransferases, SUVH4 (H3K9me2) and CLF
(H3L27me3). The Y at amino acid position 1903 of SDG2 that was mutated to F in the
SunTag:dSDG2 constructs corresponds to the Y593F in SUVH4(/KYP) that has
previously been shown to abolish methyltransferase catalytic activity (Du et al., 2014)
(position indicated in green rectangle). B) CONSURF analysis (Ashkenazy et al.,
2016) on the SDG2 predicted SET domain (amino acids 1761-1911) showing the
maximum level of conservation for $SDG2_{Y1903}$ (position indicated in green rectangle).

Supplementary Figure 3: A) qRT-PCR for the expression of FWA (upper panel) and
sfGFP (lower panel) with individual biological replicates shown. B) ChIP qPCR with
two independent FWA targeting lines.

Supplementary Figure 4: A) Venn diagram showing the overlap between the peaks
called from HA ChIP-seq of two independent SunTag:SDG2 FWA targeting lines. B)
Genome browser images for the overlapping peak regions, the top most significant
peak is FWA, while the other two are off-target regions with sequence similarity to the
FWA binding site that were previously identified (Papikian et al., 2019).

Supplementary Figure 5: Colonisation dynamics of Pst::LUX in the genotypes
indicated. Error bars represent SEM.

Supplementary Figure 6: A) Recombination rate over *CTL3.9* of the F_3 lines (as
shown in main Figure 3). Blue arrow (also shown in B) points the P4 seed set taken to
the next generation (also see main Figure 3C, red dot). B) Distortion ratios for the red
/ green T-DNAs from *CTL3.9* from the F_3's. C) Recombination rate over *CTL3.9* (F_3's)
including all scorable individuals including no guide lines. D) Recombination rate over
CTL3.9 (F_3's) filtered for individuals with no distortion (i.e. 3:1).

Supplementary Figure 7: A-B) ChIP replicate comparisons by Pearsons correlation
clustering using 25bp bin resolution. C-D) ChIPseq data chromosomal-wide plots as
shown in Figure 3, over all chromosomes from sibling lines with or without
SunTag:SDG2:LRCen3 (+/-). Enrichment is calculated as log2 fold change over non-
transgenic (Col-0) controls in 100kb windows. *CTL3.9* marker positions are indicated,
and centromeric regions are shown in grey.
876
877 **Supplementary Figure 8:** A) LRCen3 guide RNA binding site density plot with 0
878 mismatches or B) with 1 mismatch tolerance.
879
880 **Supplementary Figure 9:** Representative images from 3-week old plants.
881
882 **Supplementary Figure 10:** A) Upper panel, western blot showing bulk levels of
883 H3K4me3 normalised to H3 in the genotypes indicated. Lower panel, ratio of
884 H3K4me3/H3. B) RT-qPCR for expression of FWA (upper) and sfGFP (middle) and
885 ratio (lower) in the genotypes indicated.
886
887 **Supplementary Figure 11:** A) RT-qPCR of FWA expression (upper) and sfGFP
888 expression (lower) for individual T_1 plants from the genotypes indicated. B) summary
889 table showing the number of FWA activating T_1 lines per construct.
890
891 **Supplementary Figure 12:** A) MA plot for transcriptome of SunTag:SDG2:No_guide
892 versus the non-transformed control (*rdr6*). B) MA plot for transcriptome of
893 SunTag:PDRM9:No_guide versus the non-transformed control (*rdr6*). Differentially
894 expressed genes are depicted in red (FDR < 0.1). Note that FWA is not identified as
895 differentially expressed in either comparison.
896
897 **Supplementary Table 1:** Oligo List.

SunTag cloning	
SDG2_BsiWI_Fw	GAGGTCGGACCGGTCGTACG
SDG2_mutation_Rev	ATTAAAATCGAAAGTTATCTCCTCGCC
SDG2_mutation_Fw	GATTTTAATTCTGTAACTGAGAGTAAG
SDG2_BsiWI_Rev	CGAGATCCTCCTCCCGTACG
Universal F1	GTCATCTATGTTACGAAGCTTTCGTTGAACAACGG
Universal R2	TCGTGCTCCACCATGGTACCAAAAAAAGCACCGACTCGGT GC

pFWA_gRNA4_F2	acggaaagatgtatgggcttGTTTTAGAGCTAGAAATAGC
pFWA_gRNA4_R1	aagcccatacatctttccgtCAATCACTACTTCGACTCTA
pSNC1_gRNA4_F2	ctatgcaagtcttctaagaaGTTTTAGAGCTAGAAATAGC
pSNC1_gRNA4_R1	ttcttagaagacttgcatagCAATCACTACTTCGACTCTA
LRCen3_g_F2	aggcttacaagattgggttgGTTTTAGAGCTAGAAATAGC
LRCen3_g_R1	caacccaatcttgtaagcctCAATCACTACTTCGACTCTA
PRDM9 cloning into SunTag	
RP_PRDM9_G282A_3'	cgagatcctcctcccgtacgtcaaa
FP_PRDM9_G282A_5'	GAGGTCGGACCGGTCGTACGTATC
RP_PRDM9_G282A_middle	CTCTGTGATCTGAGCCTCATAGGGGCCA
FP_PRDM9_G282A_middle	TGGCCCCTATGAGGCTCAGATCACAGAG
ChIP-qPCRs	
High_H3K4me3_AT5G65130_F	CCAGACATTGTTCGTCAAGGACAC
High_H3K4me3_AT5G65130_R	CAACGTTATATCCGACTCAGGAGATG
Low_H3K4me3_AT2G40004up_F	CAGCTACAAGTTCAACTAATCATGC
Low_H3K4me3_AT2G40004up_R	CTTTAATTGCCCGTCATTACGCTAC
pFWA_ChIP_Fw	AATATCAGATCTTGCGCCGCTC
pFWA_ChIP_Rev	GAAAACAGACAAATCGGGAACC
pSNC1_ChIP_Fw1	ATGGCCTTATCTTGTTAGG
pSNC1_ChIP_Fw2	GAGTAGAAGCTTTCCAAGG
RT-qPCRs	
sfGFP_qPCR_Fw	CAAAGATGACGGGACCTACAA
sfGFP_qPCR_Rev	GTACTCGAGTTTGTGTCCAAGA
FWA_qPCR_Fw	TTAGATCCAAAGGAGTATCAAAG
FWA_qPCR_Rev	CTTTGGTACCAGCGGAGA
SNC1_qPCR_Fw	TTTGAAGAGACATGCAAGGC
SNC1_qPCR_Rev	TCGGCAAGCTCTTCAATCATGG

Figure 1

a

dCas9 + gRNA

— effector
— sfGFP
— scFv
— GCN4

Gene

b

SDG2 (1-2335)   C-term (1571-2335)

SET

Residue position

PAE (Å)
30
20
10

Residue position
0 500 1000 1500 2000

SDG2^Y1903F

ITFDYNSVT
↓
ITFDFNSVT

c

RT-qPCR
FWA expression

FWA (IPP2 normalised)

0.6
0.4
0.2
0.0

***

rdr6 +

non-transformed
SunTag:SDG2:FWA_g4 #12
SunTag:**dSDG2**:FWA_g4 #2
SunTag:SDG2:**No_g** #13–5

d

ChIP-qPCR

HA

SunTag (anti-HA) (% input)

0.5
0.4
0.3
0.2
0.1
0.0

rdr6 +

non-transformed
SunTag:SDG2:FWA_g4 #12

H3K4me3

H3K4me3 (FWA / control region)

0.3
0.2
0.1
0.0

rdr6 +

non-transformed
SunTag:SDG2:FWA_g4 #12

f

Transcriptome
SunTag:SDG2:FWA_g4 vs. non-transformed

Log_2(SunTag / N.T.)

-5

FWA

Significance (adj. p < 0.01)
No
Yes

-5   0   5   10   15

Log_2(Average Expression)

e

15 kb

13,032 kb   13,034 kb   13,036 kb   13,038 kb   13,040 kb   13,042 kb   13,044 kb   13,04

H3K4me3 (Xu et al. 2024)   [0 - 11]

rdr6   [0 - 73]

anti-HA ChIP-seq

ST:SDG2:FWA_g4 line #10 (in rdr6)   [0 - 73]

ST:SDG2:FWA_g4 line #12 (in rdr6)   [0 - 73]

FWA g4 binding sites   FWA_g4

AT4G25515   AT4G25520   *FWA* (AT4G25530)   AT4G25540

Figure 2

Figure 3

Figure 4

S1

a

WT | SunTag-SDG2

100 μm

b

SunTag-SDG2 FWA g4

no SunTag control

rep 1 | rep 2 | rep 3 | rep 1 | rep 2 | rep 3

dCas9 module:
dCas9-HA-3xNLS-10xGCN4
(predicted size = ~203 kDa)

effector module:
scFv-HA-sfGFP-HA-2xNLS-SDG2
(predicted size = ~145 kDa)

RuBisCo

c

FWA relative expression (normalised to IPP2)

1.2

0.8

0.4

rep 1 | rep 2 | rep 3 | rep 1 | rep 2 | rep 3

SunTag-SDG2 FWA g4

no SunTag control

S2

a   SDG2 alignment over SET domain:

```
SDG2   1858 ANYASRICHSCRPNCEAKVTAVDG----HYQIGIYSVRAIEYGEEITFDYNSVTESKEEY   1913
SUVH4   544 GNFARFINHSCEPNLFVQCVLSSHQDIRLARVVLFAADNISPMQELTYDYGYALDSVHGP    603
CLF     821 GDKLKFANHSPEPNCYAKVIMVAG----DHRVGIFAKERILAGEELFYDYRYEPDRAPAW    876
```

b   CONSURF analysis:

The conservation scale:

? 1 2 3 4 5 6 7 8 9
Variable   Average   Conserved

e – An exposed residue according to the neural network algorithm.

b – A buried residue according to the neural network algorithm.

f – A predicted functional residue (highly conserved and exposed).

s – A predicted structural residue (highly conserved and buried).

S3

S4

a

ST-SDG2 FWA #10 peaks — 14 — 3 — 1 — ST-SDG2 FWA #12 peaks

b

FWA (target)

Off-target 1

Off-target 2

S5

Pst::LUX colonisation dynamics on control genotypes

S6

a   Recombination rate of SunTag:SDG2:LRCen3_g independent transformants over CTL3.9

b

Green to non-green distortion

Red to non-red distortion

Recombination frequency over CTL3.9 with different SunTag-SDG2 guide RNAs

c   all scored individuals

d   filtered for distortion

S7

a

anti-H3K4me3 bam comparison

b

anti-HA comparison bam comparison

c

log$_2$(SunTag/Col-0) H3K4me3 over all chromosomes

- H3K4me3_B12overCol_log2_100bp.bg (+ SunTag)
- H3K4me3_B5overCol_log2_100bp.bg (- SunTag)

H3K4me3 in SunTag/WT across Chromosomes 1–5 (100kb bins)

d

log$_2$(SunTag/Col-0) HA over all chromosomes

- HA_B12overCol_log2_100bp.bg (+ SunTag)
- HA_B5overCol_log2_100bp. (- SunTag)

HA in SunTag/WT across Chromosomes 1–5 (100kb bins)

S8

a

Density plot of LRCen3 gRNA binding sites
allowing 0 mismatch across the genome

b

Density plot of LRCen3 gRNA binding sites
allowing 1 mismatch across the genome

a

SunTag-SDG2 no guide causes bulk
increases in H3K4me3

b

Overexpression of SunTag-SDG2**dcat**
effector can activate FWA

a

b

Genotype	T1's screened (no.)	FWA expressing (no.)	FWA expressing (%)
SunTag:PRDM9:FWA_g4	32	23	72%
SunTag:dPRDM9:FWA_g4	19	0	0%
SunTag:PRDM9:No_g	16	0	0%

a

SunTag:SDG2:NoGuide vs. *rdr6*

b

SunTag:PRDM9:NoGuide vs. *rdr6*

*Note that FWA is not a differentially expressed gene in either comparisons