CRISPR targeting of H3K4me3 activates gene expression and unlocks centromere-proximal crossover recombination in Arabidopsis

Jenia Binenbaum[1,*], Vanda Adamkova[1,*], Hannah Fryer[1], Linhao Xu[1], Nicola Gorringe[1], Piotr Włodzimierz[1,2], Robin Burns[1], Ashot Papikian[3,4], Steven E. Jacobsen[3,5], Ian R. Henderson[1], C. Jake Harris[1,#]

[1] Department of Plant Sciences, University of Cambridge, Cambridge CB2 3EA, UK.

[2] Current Address: Institute of Biochemistry and Biophysics, Polish Academy of Sciences, Warsaw, Poland.

[3] Department of Molecular, Cell and Developmental Biology, University of California, Los Angeles, CA, USA.

[4] Current Address: Plant Molecular and Cellular Biology Laboratory, Salk Institute for Biological Studies, La Jolla, CA 92037, USA.

[5] Howard Hughes Medical Institute, University of California, Los Angeles, CA, USA.

* These authors contributed equally.

[#] For correspondence: cjh92@cam.ac.uk

ABSTRACT

H3K4me3 is a fundamental and highly conserved chromatin mark across eukaryotes,

playing a central role in many genome-related processes, including transcription,

maintenance of cell identity, DNA damage repair, and meiotic recombination. However,

identifying the causal function of H3K4me3 in these diverse pathways remains a

challenge, and we lack the tools to manipulate it for agricultural benefit. Here we use

the CRISPR-based SunTag system to direct H3K4me3 methyltransferases in the

model plant, *Arabidopsis thaliana*. Targeting of SunTag-SDG2 activates the

expression of the endogenous reporter gene, *FWA*. We show that SunTag-SDG2 can

be employed to increase pathogen resistance by targeting the H3K4me3-dependent

disease resistance gene, *SNC1*. Meiotic crossover recombination rates impose a limit

on the speed with which new traits can be transferred to elite crop varieties. We

demonstrate that targeting of SunTag-SDG2 to low recombining centromeric regions

can significantly stimulate proximal crossover formation. Finally, we reveal that the

effect is not specific to SDG2 and is likely dependent on the H3K4me3 mark itself, as

the orthogonal mammalian-derived H3K4me3 methyltransferase, PRDM9, produces

a similar effect on gene expression with reduced off-target potential. Overall, our study

supports an instructive role for H3K4me3 in transcription and meiotic recombination

and opens the door to precise modulation of important agricultural traits.

INTRODUCTION

Re-programming of transcriptional landscapes in response to developmental and environmental cues critically underlies a plant's ability to adapt and optimise its performance. The potential to re-wire and fine-tune expression of endogenous genes therefore holds great promise for agricultural improvement. Chromatin modifications associated with gene expression play an essential role in this process [1].

Trimethylation of histone 3 lysine 4 (H3K4me3) is a well-studied and highly conserved chromatin mark. H3K4me3 enrichment around the transcriptional start site of expressed genes is observed across eukaryotes [2], and is thought to assist in transcriptional maintenance and memory of cell identity [3]. Despite the ubiquity of H3K4me3 and its positive correlation with gene expression, whether this chromatin mark plays an instructive role in transcription remains a matter of debate, due in part to the limited evidence of transcriptional impact when H3K4me3 levels are perturbed [4,5]. In animals, TAF3 directly binds to H3K4me3 and recruits basal transcription factor machinery, providing a plausible mechanism to stimulate transcription [6,7]. Recent work in embryonic stem cells shows that H3K4me3 plays a role in RNA Polymerase II pause-release and elongation, rather than transcriptional initiation [8]. In plants, there are no TAF3 homologues and whether similar mechanisms exist to promote transcription downstream of H3K4me3 remains unknown [9-13].

H3K4me3 also plays a key role in several other genome-related processes. For instance, meiotic recombination, which allows for genetic exchange between parental chromosomes [14]. This process is essential for breeders to transfer traits from wild relatives into elite cultivars [15,16]. However, the positioning of meiotic recombination events is non-uniform across the genome, with gene-rich chromosomal arms having relatively higher rates of crossover recombination, and centromeric repeat-rich regions being more recombination-suppressed [17,18]. Loci encoding desirable traits that reside within these low-recombining regions of the genome can show linkage drag with deleterious variants during traditional breeding approaches, which limits genetic gains [15,16]. In mammals, the H3K4me3 SET domain methyltransferase containing protein, PRDM9, helps to define active sites of crossover recombination [19]. While H3K4me3 levels are positively correlated with recombination rates in plants [17,20], whether

deposition of H3K4me3 is sufficient to promote crossover formation is entirely
unknown.

One way to reveal the causal role of an epigenetic modification is to directly deposit
the mark at targeted genomic locations [21–25]. Epigenome editing tools, such as the
CRISPR-based SunTag system [26], have proven to be highly efficient at targeting DNA
methylation addition, removal and histone modification [23,27–32], although their utility for
modulating agronomically useful traits is only just beginning to be realised [33], and is
limited by the range of possible marks that can be directed [1]. In the model plant
Arabidopsis thaliana, SDG2 is the main histone methyltransferase responsible for
deposition of H3K4me3 [34]. Here, we describe CRISPR directed H3K4me3 deposition
in plants and demonstrate its utility in a range of scenarios. We translationally fuse the
SDG2 methyltransferase domain to the scFv single chain antibody, which can be
recruited to the SunTag epitope chain connected to a dead Cas9 (dCas9). We show
that SunTag-SDG2 mediated H3K4me3 deposition can activate the transcription of
endogenous genes. Further, by targeting a disease resistance gene, we generate
plants with enhanced resistance capabilities. Next, we show that SunTag:SDG2 can
unlock meiotic crossover recombination in centromere-proximal regions. Finally, we
incorporate the methyltransferase domain from PRDM9 into SunTag - which was
recently shown to instruct transcription in a mammalian context [31] - showing that it is
also highly efficient for transcriptional activation in plants, and displays reduced off-
target potential. These results demonstrate the power and versatility of a functional
H3K4me3 targeting system for modulation of critical genome templated processes,
such as transcriptional enhancement and meiotic recombination.

 RESULTS

SunTag SDG2 mediates transcriptional activation of *FWA*

The SunTag system we chose is composed of three components [26,35] (Fig. 1A): 1) A

catalytically deactivated (nuclease deficient) Cas9 (dCas9) translationally fused to a

tail with 10x copies of the GCN4 epitope, each separated by a 22 amino acid flexible

linker. 2) An effector module that encodes a single chain fragment variable (scFv)

antibody that recognises GCN4, super folder GFP (sfGFP), and the effector of interest

(e.g. enzymatic modifier / recruitment scaffold). 3) *U6* promoter driven CRISPR guide

RNA. When the three components are expressed within the same cell, the guide RNA

recruits the dCas9-10xGCN4 to the target locus of interest, while the GCN4 single

chain antibody containing effector module binds to the dCas9-10xGCN4 epitope tail.

The system allows for conformational flexibility and the potential for high stochiometric

concentration of the effector at the locus of interest.

We inserted the C-terminal coding sequence of the H3K4 methyltransferase, SDG2

(amino acids 1571-2335), into the SunTag system (Fig. 1B, Supp. Fig. S1). As a

control, we also generated a version of SDG2 with an amino acid change (Y1903F)

predicted to abolish catalytic activity (Supp. Fig. S2) [34,36]. Hereafter, we refer to these

as SDG2 and dSDG2 (for catalytically deactivated). We targeted the SunTag system

to the promoter of the epigenetically repressed *FWA* gene [37], using the previously

published CRISPR guide RNA 4, which is complementary to two tandem repeat

regions directly upstream of the *FWA* transcription start site (TSS) [29]. These were

transformed into the *rdr6* background, as previous results have shown that SunTag is

less prone to silencing and achieves higher transgene expression [27]. We observed

that SunTag:SDG2:FWA_g4 activated *FWA* mRNA expression, while the

SunTag:dSDG2:FWA_g4 and no guide RNA (SunTag:SDG2:No_g) controls did not

(Fig 1C, Supp. Fig. 1C). Importantly, effector module expression was detected in all

constructs, indicating that both guide RNA targeting and catalytic activity are required

for *FWA* activation (Supp. Fig 3A). To validate the presence of the SunTag and test for

the deposition of H3K4me3, we performed ChIP qPCR and observed significant

enrichment both for the presence of SunTag and for H3K4me3 accumulation at the

target site in these transgenic lines (Fig. 1D and Sup Fig 3B), confirming that SunTag-

SDG2 binds the *FWA* locus and deposits H3K4me3. To investigate
SunTag:SDG2:FWA_g4 binding genome-wide, we performed high-throughput
sequencing (anti-HA ChIP-seq; the HA-tag is present on both the dCas9 and effector
fusion modules), which confirmed the previously reported high specificity of SunTag
targeting [28], with the *FWA* target locus as the most enriched binding target (Fig. 1E,
Supp. Fig. 4). Only 3 conserved binding sites were identified in the
SunTag:SDG2:FWA_g4 lines; *FWA* and two off-target sites previously identified as
having sequence-similarity to FWA_g4 [28]. Furthermore, transcriptome analyses
confirmed *FWA* as the highest fold change differentially upregulated gene (Fig. 1F).
Together, these data indicate that deposition of H3K4me3 by SunTag:SDG2 is
sufficient to overcome epigenetic silencing and drive transcriptional activation in a
locus-specific manner.
142
Next, to examine the genome-wide effect on H3K4me3 by SunTag-SDG2 we
performed ChIP-seq analysis, confirming enrichment at the *FWA* locus and canonical
genic H3K4me3 patterning (Fig 1E, Supp. Fig. 5). In addition to a modest peak at the
+1 nucleosome at *FWA*, the H3K4me3 signal extended into the gene body. Performing
metaplots over genes, transcriptional start sites (TSS's), and endogenous H3K4me3
peaks, we observed that SunTag:SDG2 lines generally display a moderate increase
of H3K4me3 over the 3' end of gene bodies (Fig. 1G). Correspondingly, we observed
reduced H3K4me3 signal at endogenous peaks and TSS regions (Fig. 1G). As the
H3K4me3 peak at *FWA* in our SunTag:SDG2:FWA_g4 lines was not as strong as that
observed for *fwa* in the epiallelic state [27,38], we asked whether the effect could persist
in the next generation. Analysis of T_2 seedlings from two independent lines
segregating for the presence of SunTag:SDG2:FWA_g4 showed that *FWA* activation
was entirely dependent on the presence of the transgene (Supp. Fig. 6). Together
these data indicate that while SunTag:SDG2:FWA_g4 can site-specifically activate
FWA expression, it is insufficient to overcome the endogenous silencing machinery as
required to switch *FWA* into the stably active *fwa* epiallelic state [32,39].

Figure 1: SunTag:SDG2 activates *FWA* mRNA expression.

A) Schematic showing the SunTag system for modulation of chromatin [28,35]. Created in BioRender. Harris, J. (2025) https://BioRender.com/mqs3hbr. B) Upper panel shows SDG2

with catalytic SET domain (drawn approximately to scale). Lower left depicts predicted alignment error (PAE) plot from AlphaFold3 [75] prediction of the SDG2 coding sequence with a box drawn around the region cloned into SunTag (amino acids 1571-2335). Lower right depicts the AlphaFold3 model for $SDG2_{1571-2335}$, with the position of the amino acid change (Y1903F) for dSDG2 indicated (shown in red). C) RT-qPCR for *FWA* expression from the genotypes indicated. Error bars represent SEM from three biological replicates (a pool of 10 seedlings) from the lines (#) and genotypes indicated. *** indicates *p* <0.005 (ANOVA with post-hoc Tukey HSD. D) ChIP qPCR for the presence of SunTag (left panel) and H3K4me3 enrichment (right panel) over the TSS of *FWA*. The control region (AT5G65130) used for normalisation has high H3K4me3. E) Genome browser images showing ChIP-seq enrichment of SunTag (anti-HA) and H3K4me3 levels of two independent SunTag:SDG2:FWA_g4 lines (2 biological replicates each, bio replicate tracks are overlaid) at the *FWA* TSS (left panel) and at a random genic control region. F) MA plot comparing the transcriptome of SunTag:SDG2:FWA_g4 as compared to the non-transformed control (*rdr6*). Differentially expressed genes were defined using an adjusted p-value (Bonferroni method), shown in red (FDR < 0.01), with *FWA* labelled and enlarged for visibility. G) H3K4me3 ChIP-seq metaplots over all genes, H3K4me3 endogenous peaks, and TSS regions.

SNC1 mediated modulation of disease resistance

Given that SunTag:SDG2 can modulate chromatin and activate *FWA* gene expression, we were interested use the tool to modify an important plant trait. Gene expression during pathogen attack is a key factor in determining disease outcome. *SNC1* is disease resistance gene, encoding a nucleotide-binding leucine-rich repeat protein [40] that is located within the partially epigenetically repressed *RPP5* gene cluster [41–43]. *SNC1* expression activity is modulated by H3K4me3 [44], and its level of expression is positively correlated with basal resistance [41,44]. Therefore, *SNC1* is an attractive target for proof-of-principle modulation of disease resistance by epigenome engineering.

We designed guide RNAs to target the promoter region of *SNC1* directly upstream of the TSS and inserted it into our SunTag:SDG2 and SunTag:dSDG2 constructs (Fig. 2A, Supp. Fig 7). As high pathogen resistance is generally associated with dwarfed plants, we inspected the lines for growth and developmental phenotypes. In our *SNC1* targeting SunTag:SDG2, but not our SunTag:dSDG2 control, we observed small, dwarfed phenotype with reduced rosette surface area (Fig. 2AB, Supp. Fig 7). These

198 plant phenotypes resembled those of our positive control *bal* lines, which overexpress
199 *SNC1*, resulting in the small dwarfed plant phenotype and high basal disease
200 resistance [41]. As the phenotype is consistent with upregulation of *SNC1* by SunTag,
201 we confirmed the increased expression and successful targeting of SunTag:SDG2 to
202 the *SNC1* locus (Fig. 2CD, Supp. Fig 7).
203
To assess whether these lines show increased pathogen resistance, we challenged
the plants with the generalist plant pathogen, *Pseudomonas syringae* pathovar *tomato*
strain DC3000 (*Pst*). The recently described bioluminescent *Pst* (*Pst::LUX*) was used
to allow for non-destructive quantification of pathogen colonisation [45]. We developed
a high-throughput assay system, inoculating 7-day-old seedlings with *Pst::LUX* and
imaging to quantify the levels of *Pst::LUX* at 24 hour intervals over the course of
infection. We challenged a range of control genotypes that are known to be *Pst* hyper-
resistant (*bal*), constitutively primed (*edr1*), and susceptible (*NahG*), recapitulating the
expected pattern of *Pst::LUX* colonisation dynamics in these backgrounds (Supp. Fig.
S8) [45]. Next, we challenged our *SNC1* targeting lines with *Pst::LUX*, finding that they
displayed hyper-resistance that was nearly identical in magnitude to that of the *SNC1*
overexpressing positive control, *bal* (Fig. 2E, Supp. Fig. 7). The results demonstrate
that epigenome-engineering of a single defense gene, *SNC1*, is sufficient to generate
plants with improved disease resistance phenotypes in Arabidopsis.
218

Figure 2: SunTag:SDG2 targeting to *SNC1* enhances resistance to *P. syringae*.
A) Upper, schematic of the *SNC1* gene, with the red line indicating the *SNC1* gRNA target
site, drawn to scale. Lower, representative images of 3-week old plants from the genotypes
indicated. Scale bar = 1cm. B) Rosette size quantification. Each dot represents an individual
plant. Error bars represent SEM. Different letters indicate significant difference by ANOVA with
post-hoc Tukey HSD, $p < 0.05$. n = 14 for SunTag:dSDG2:SNC1_g4 #1, n = 15 for the rest. C)
ChIP qPCR for the presence of SunTag at SNC1. D) RT-qPCR for *SNC1* expression. Error

bars represent SEM. * indicates p-value = 0.045 by student's *t*-test. n = 4 biological replicates

(each a pool of 10 seedlings). E) *Pst::LUX* assay for colonisation quantification at 3 days post

inoculation. Different letters indicate significant difference by ANOVA with post-hoc Tukey

HSD, *p* < 0.05. Boxplots show the median, the interquartile range, whiskers extending to 1.5×

the interquartile range, and individual data points plotted as dots. n = 96 wells, containing 3

seedlings each.

Centromeric targeting of SDG2 drives increased meiotic crossover

recombination

Beyond transcription, H3K4me3 is associated with many other fundamental genome

related processes including meiotic recombination. In plants, meiotic crossover

recombination rates broadly correlate with epigenetic territories and are non-uniform

across the genome [17,18]. In *A. thaliana*, recombination occurs at a relatively high

frequency in euchromatic arms, whereas it is significantly reduced over the

heterochromatic pericentromeric and centromeric [46–48]. Crossover hotspots are often

found to be high at gene promoters where H3K4me3 is enriched [20]. In previous work,

we showed that gene-associated crossover recombination hotspots can be supressed

by small RNA directed DNA methylation [49]. Therefore, we hypothesised that

recombination within the centromeric regions might be increased by SunTag mediated

epigenetic activation via deposition of H3K4me3.

The Arabidopsis centromeres are composed of highly repetitive *CEN178* satellite

repeat arrays [47,50]. We designed a guide RNA that perfectly matches to 241 target sites

within the *CEN178* repeats within the centromeric region of chromosome 3

(LRCen3_g), and inserted this guide into the SunTag:SDG2 system. As centromere-

proximal crossover recombination events are relatively rare, to quantitatively assess

the impact on recombination rate over centromere 3, we crossed (non-transgenic) Col-

0 plants with the *CTL3.9* crossover reporter [51]. *CTL3.9* is a transgenic a line that

encodes linked T-DNAs that encode red and green fluorescent markers expressed

from a seed-specific promoter, that flank *CEN3* [48]. The red and green T-DNA insertions

span a region of approximately 8.91 Mb in the Col-CEN assembly of chromosome 3,

which contains the centromeric satellite arrays in addition to flanking pericentromeric

heterochromatin (Fig. 3A) [48]. Meiotic crossover recombination events that occur within

this region result in progeny inheriting either red or green fluorescent markers, the
frequency of which can be measured by automated seed imaging [52]. We transformed
CTL3.9 F_1 double hemizygous plants with the SunTag:SDG2:LRCen3_g construct
(Fig. 3AB). The F_2 plants represented individual transformants of the SunTag (T_1 for
SunTag), and so F_3 seeds from individual F_2 plants were imaged to measure the rate
of crossover recombination, in centiMorgans, over the centromere 3 spanning region
of *CTL3.9* [48].

Remarkably, the SunTag:SDG2:LRCen3_g containing lines showed significantly
elevated crossover recombination within the *CTL3.9* interval as compared to the non-
SunTag control lines (ANOVA with post-hoc Tukey HSD cutoff <0.05) (Fig. 3C, Supp.
Fig. 9). Some individual transgenic plants displayed a substantial increase in *CTL3.9*
crossover recombination rate, with over 50% higher frequency, as compared to the
average in non-transgenics, across this megabase-scale centromere-spanning region
(Fig. 3C). SunTag:SDG2:LRCen3_g progeny from the most highly recombining line
that also showed Mendelian inheritance of the *CTL3.9* T-DNAs was sown to assess
whether the effect could persist in the subsequent (F_4) generation. The plants retained
a significantly elevated crossover frequency across *CTL3.9* in the next generation,
suggesting that the SunTag:SDG2:LRCen3_g effect is relatively stable in the presence
of the transgene (Fig. 3C).

To examine the distribution of SunTag:SDG2:LRCen3_g in the high recombination
lines (P4, Supp. Fig. 9), we performed ChIP-seq (anti-HA) comparing progeny from
sibling plants that either inherited (+SunTag), or did not inherit (-SunTag) the SunTag
machinery, as well as non-transgenic (Col-0) controls. Chromosome-wide plots at 100
kb resolution showed that SunTag:SDG2 is enriched over the entire centromeric
region of chromosome 3 (Fig. 3D). SunTag was also enriched over the other four *A.*
thaliana centromeres (Supp. Fig. 10). As these regions are highly repetitive, we
reasoned that the LRCen3 guide RNA might additionally bind the other centromeres
due to mismatch tolerance. Indeed, while the LRCen3 guide RNA aligns perfectly to
250 loci within the centromere of chromosome 3, by allowing a single mismatch the
guide is predicted to bind to target sites in all 5 centromeres (Supp. Fig. 11), which is
consistent with the ChIP-seq data observed. Having confirmed that SunTag is
enriched within the centromeric regions in these lines, we next examined levels of

H3K4me3. Inspecting centromeric regions, we observed many instances of novel or
elevated H3K4me3 peaks in the LRCen3 targeted lines (Supp. Fig. 12). Quantifying
H3K4me3 levels at these peaks confirmed the enrichment of H3K4me3 levels over
the centromere region and further extending into the neighbouring pericentromeres
(Fig. 3D, Supp. Fig. 13). As the markers on *CTL3.9* span the centromere and adjacent
pericentromeric regions [48] (Fig. 3D), this is consistent with the significantly elevated
crossover recombination rate observed. Overall, the results demonstrate that
centromerically targeted SunTag-SDG2 can elevate the meiotic crossover potential of
these typically recombination-suppressed regions.

304
305
306 **Figure 3: Centromeric targeting of SunTag:SDG2 elevates meiotic crossover**
307 **recombination rate.**
308 A) Region of chromosome 3, showing CENH3 ChIP-seq enrichment [50] with *CTL3.9* [51] red /
309 green T-DNA marker positions and LRCen3 guide RNA binding sites indicated. B) Left panel,

representative image of red and green fluorescing seeds from a *CTL3.9* double homozygous line. Right panel, schematic of showing how crossovers are identified within *CTL3.9*. C) Boxplot showing crossover recombination frequency over *CTL3.9* in centiMorgans. Control data is from Col-0 (non-transgenic) crossed to *CTL3.9*. Different letters indicate significant difference by ANOVA with post-hoc Tukey HSD ($p<0.05$). Control: n = 12, LRCen3 F3: n = 19, No guide: n = 6, LRCen3 F4: n = 14. Boxplots show the median, the interquartile range, whiskers extending to 1.5× the interquartile range, and individual data points plotted as dots. The red dot indicates the seed set line taken to the next generation (F_4). D) Chromosome-wide plots of chromosome 3. Upper panels shows enrichment of SunTag (anti-HA) by ChIP-seq in sibling lines with / without SunTag (+/-). Enrichment is calculated as $\log_2$ fold-change over non-transgenic (Col-0) controls in 100 kb windows. Middle panel shows H3K4me3 enrichment over H3K4me3 peaks ($\log_2$ fold-change over non-transgenic (Col-0) controls). The lower panel depicts an LRCen3 guide RNA binding site density over chromosome 3. *CTL3.9* marker positions are indicated, and centromeric regions are shown in grey. All data are mapped the Col-CEN genome assembly [47].

Mammalian PRDM9 drives efficient transcriptional activation in Arabidopsis

As the enrichment of H3K4me3 in SunTag:SDG2:LRCen3_g was not exclusive to the guide RNA targeted centromeric regions, we reasoned that the SunTag system may afford some level of off-target activity due to non-guide RNA directed ectopic interactions. Consistent with this observation, we also noticed that our SunTag:SDG2 no guide RNA controls frequently exhibited pleotropic developmental phenotypes (Supp. Fig. 14). To quantify global levels of H3K4me3, we performed bulk histone westerns, comparing no guide RNA lines to *FWA* expression-activating guide RNA containing lines. Importantly, the SDG2 effector itself is expressed to similar levels as in the *FWA* targeting lines (Supp. Fig. S3a). The SunTag:SDG2 no guide lines all showed significant (3-4 fold) increases in global levels of H3K4me3, as compared to the *FWA* guide containing lines (Supp. Fig. 15A). This indicates that the presence of the guide RNA may enhance the targeting specificity of the SunTag complex by reducing the frequency of non-specific interactions that result in off-target deposition of H3K4me3. We also observed that a small number of SunTag:dSDG2:FWA_g4 lines could activate *FWA* expression, but this effect was limited to lines where the dSDG2 effector was highly overexpressed (Supp. Fig. 15B). To systematically examine this effect, we screened T_1 plants finding that 6% (1/16) of the *FWA* targeted dSDG2, and

14% (2/16) of the no guide SDG2 lines could activate *FWA* expression (Supp. Fig. 16). This underscores the potential off-target effects from SunTag:SDG2, and suggests that dSDG2 either retains some residual catalytic activity, or that it can recruit endogenous machinery to initiate transcription independently of H3K4me3 deposition.

In order to reduce off-target activity and the potential for co-recruitment of endogenous complexes, we replaced SDG2 with an orthogonal effector, PRDM9. PRDM9 is a mammalian H3K4me3 methyltransferase, and Policarpi *et al.,* (2024) recently showed that the catalytic domain could be incorporated into the SunTag system for targeting of H3K4me3 in mammalian cells. Therefore, we inserted the PRDM9 methyltransferase domain, as well as a catalytically deactivated version, into our SunTag system and targeted *FWA* (Fig. 4A). Strikingly, SunTag:PRDM9:FWA_g4 was highly effective at activating *FWA* mRNA expression (Fig. 4B). In the T_1 generation, 72% (23/32) of SunTag:PRDM9:FWA_g4 plants could activate FWA, while no of plants with the dPRDM9 (0/16), or no-guide RNA (0/19) constructs were able to do so (Supp. Fig. 17). Importantly, we could detect robust expression of the effector from all constructs (Supp. Fig. 11). ChIP qPCR confirmed the presence of SunTag and the enrichment of H3K4me3 at *FWA* (Fig 4C). At the transcriptome level, the effect of SunTag:PRDM9:FWA_g4 was highly specific, with only 19 genes identified as differentially expressed (15 up, 4 down, FDR<0.01), and with *FWA* itself having the highest fold change amongst them (>158 fold change over *rdr6*) (Fig. 4D). Unlike the no-guide SDG2 transgenic lines, bulk histone westerns confirmed no H3K4me3 increases in the PRDM9 no-guide lines (Supp. Fig. 18). Correspondingly we did not observe any developmental defects and saw only minor transcriptional defects when comparing the no-guide lines to non-transformed *rdr6* controls (Supp. Fig. 19). Finally, we performed ChIP-seq in these lines, confirming *FWA* as the most significant and enriched peak for SunTag binding and revealing deposition of H3K4me3 at the transcriptional start site of *FWA* (Fig. 4E).

At the genome-wide level, while SunTag:PRDM9:FWA_g4 lines also displayed modest decrease in H3K4me3 signal at endogenous peaks with a corresponding increase over 3′ gene body regions (Fig. 4F, Supp. Fig. 20), these effects were less pronounced than SunTag:SDG2 (Fig. 1G), again suggesting that the PRDM9 effector fusion has reduced off-target potential. Finally, we performed the *CTL3.9* meiotic recombination

assay, finding that PRDM9 was similarly effective at increasing crossover frequency
over the centromeric *CTL3.9* interval (Fig. 4G, Supp. Fig. 21). Overall, our results
demonstrate that PRDM9 is highly effective for deposition of H3K4me3 in plant, as
well as mammalian SunTag systems. Furthermore, as PRDM9 is not likely to have
endogenous partners in the plant nucleus, the results provide strong evidence in
favour of a direct role for H3K4me3 in transcriptional stimulation.

Figure 4: SunTag PRDM9 is sufficient for activation of *FWA*.

A) Upper panel shows PRDM9 with catalytic SET domain (drawn approximately to scale). Lower left depicts predicted alignment error (PAE) plot from AlphaFold3 prediction of the PRDM9 coding sequence with a box drawn around the region cloned into SunTag (amino acids 110-417). Lower right depicts the AlphaFold3 model for PRDM9$_{110-417}$, with the amino acid change for dPRDM9 indicated (position shown in red). B) RT-qPCR for *FWA* (upper panel) and the effector module (sfGFP, lower panel) for the genotypes indicated. Dots represent individual plants, with two independent T$_3$ lines used per construct, 3 biological replicates per construct. Error bars represent SEM. C) ChIP qPCR for the presence of SunTag (left panel) and H3K4me3 enrichment (right panel) over the TSS of *FWA*. D) MA plot comparing the transcriptome of SunTag:PRDM9:FWA_g4 as compared to the non-transformed control (*rdr6*). Differentially expressed genes were defined using an adjusted p-value (Bonferroni method), shown in red (FDR < 0.01), with FWA labelled and enlarged for visibility. E) Genome browser image showing efficient SunTag targeting and H3K4me3 enrichment at *FWA* and at a random genic control region. Biological replicate tracks from independent lines are overlaid. F) H3K4me3 ChIP-seq metaplots over all genes, H3K4me3 endogenous peaks, and TSS regions. G) Boxplot showing crossover recombination frequency over *CTL3.9* in centiMorgans (*p*<0.001, two-sample, two-sided *t*-test). Boxplots show the median, the interquartile range, whiskers extending to 1.5× the interquartile range, and individual data points plotted as dots. n = 24 for control, n = 17 for Suntag:PRDM9 LRCen3.

DISCUSSION

Our results demonstrate that targeted deposition of H3K4me3 using the SunTag system enables transcriptional activation, enhanced pathogen resistance, and increased crossover recombination over megabase-scale centromere-proximal regions. These findings support an instructive role for H3K4me3 in transcriptional control and open new avenues for precise, potentially non-GM crop improvement through epigenome engineering. Because the effect depends on the presence of a functional catalytic domain, and is observed using both native and non-native methyltransferases, it is unlikely to result from co-recruitment of endogenous complexes, strengthening the argument that H3K4me3 itself can act as a causal regulatory mark.

In a related study [38] showed that full activation of *FWA* by recruitment of SDG2

required active removal of DNA methylation by ROS1 and related proteins. Whether

H3K4me3 directly recruits ROS1 machinery is currently unknown. However, the

finding that PRDM9 also causes potent transcriptional activation of *FWA* suggests that

ROS1 recruitment is downstream of H3K4me3, rather than being co-recruited via

COMPASS complex components [9,10,53]. Removal of DNA methylation could also help

explain the increased centromeric crossover recombination observed over the *CTL3.9*

interval, consistent with prior work showing elevated recombination rates in non-CG

DNA methylation mutants in similar interval [48,54]. However, in mammalian systems,

H3K4me3 has been shown to stimulate transcription by promoting Pol II pause release

[8]. Similarly, using zinc finger-fused SDG2 in *A. thaliana*, the Wang et al 2025 study

found many instances of ectopic H3K4me3 driving transcriptional stimulation in the

absence of pre-existing or DNA methylation removal. Thus, DNA methylation-

independent mechanisms must also play a role H3K4me3 mediated transcriptional

activation.

In our system, SunTag:SDG2 and PRDM9 targeting of *FWA* with a single guide caused

strong upregulation at this locus, yet we did not observe any plants that switched to

late-flowering as previously reported when *FWA* transitions to an active *fwa* epistate.

The Wang et al, 2025 study only observed flowering delays when two guides were

used to target the *FWA* promoter. We cannot exclude the possibility that *FWA* failed

to reach a critical expression threshold in our experiments, or that expression was

developmentally mis-timed or mis-localised. It is also possible that in the single guide

system the H3K4me3 deposition induces more non-functional transcripts, consistent

with chromatin's roles in co-transcriptional regulation [55,56]. The lack of a flowering time

phenotype also aligns with the absence of transgenerational effects, suggesting that

the targeting did not remove sufficient DNA methylation to establish heritable gene

FWA expression. This is further supported by the modest H3K4me3 peak observed at

FWA in our targeted lines, compared to lines that have fully switched to late flowering

[38].

Both mammalian PRDM9 and the endogenous SDG2 catalytic domains were able to

transcriptionally activate *FWA* and elevate crossover recombination rates over a

recombination-suppressed centromere spanning interval. Seminal work in *Drosophila*

melanogaster and showed that while loss of the H3K4 monomethyltransferases, *Trr*,
was lethal, the corresponding catalytic-null caused only minor developmental defects
[57]. This underscores the importance of using catalytic mutations and orthogonal
effectors to separate the effects from histone marks themselves, over the histone
modifying enzymes and associated complexes. It also highlights the utility of modular
targeting platforms that can decouple chromatin modifications from native pathways.
Continued development of such tools will be crucial for both basic discovery and
applied crop engineering [23,25,27,28,58–66].

We observed off-target effects in both SunTag:PRDM9 and SunTag:SDG2 transgenic
lines, manifested as increased non-specific H3K4me3 background signal and reduced
occupancy at endogenous locations (Fig. 1G, Fig. 4F, Supp. Fig. 13). This off-target
signal is inherently difficult to diagnose, as it is virtually indistinguishable from a ChIP-
seq with reduced signal-to-noise. However, as we observed the same trend in multiple
independent experiments, with the SunTag lines always showing this same pattern as
compared to the non-transgenics (Fig. 1G, Fig 4F, Supp. Fig. 13A), we conclude that
the effect is due to SunTag off-targeting, rather than ChIP signal variability. Notably,
these effects were weaker in PRDM9 lines, supporting the idea that orthogonality
reduces off-target potential. Unlike SDG2, PRDM9 lines without guide RNAs did not
show elevated bulk H3K4me3 levels (Supp. Fig. 15 vs. 18). The strong developmental
phenotypes in the SDG2 no-guide lines (Supp. Fig. S14) may result from
transcriptional misregulation at ectopically modified loci, or could be caused by
transcription-independent metabolic defects [5]. Interestingly, these phenotypes were
largely absent in the SDG2 guide RNA targeted lines, suggesting that the guide RNAs
reduce this off-target activity. With the guided lines, it is worth reiterating that the
SunTag on-target activity is highly specific for both SDG2 and PRDM9, with *FWA* as
the most significantly enriched SunTag peak by ChIPseq, and is the most upregulated
transcript with few other DEGs observed (Fig. 1F and Fig. 4D). However, off-target
effects and metabolic impacts are important considerations when designing these
systems for use in crops, which possess larger genomes and more complex regulatory
architectures than *A. thaliana* [67]. Notably, SunTag has already been successfully
deployed in rice for precise removal of DNA methylation in the context of cold stress
adaptation [60], demonstrating the feasibility and potential of these epigenome
engineering approaches in agricultural settings.

In summary, our work demonstrates that targeted deposition of H3K4me3 is a powerful

tool for modulating transcription and meiotic crossover recombination in plants. By

deploying both endogenous and orthogonal methyltransferases, we reveal core

insights into H3K4me3 function and highlight the potential of this platform for

translational application. Future optimisation of guide design, effector choice, and off-

target mitigation will be important for enabling both mechanistic discovery and crop

engineering through epigenome editing.

METHODS:

Plant materials and growth conditions

All *Arabidopsis thaliana* lines are in a Col-0 ecotype background. Plants grown on plates were sown on to 1/2x Murashige-Skoog (MS) medium, 1% sucrose and 0.8% agar (pH 5.7), stratified for 2 d at 4 °C in the dark, then transferred to growth chambers (Percival CU-41L4D) at 21 °C under long day light conditions (16 hour light/8 hour dark). Plants grown on soil were grown in F2 soil at 20°C under long day light conditions (16 hour light/8 hour dark). Lines: *rdr6-15* (SAIL_617_H07), and *fwa* lines were previously described [27]. NahG and *edr1* were kindly provided by Prof. Juriaan Ton, previously described in [45]. *bal* lines were previously described in [41]. The CTL 3.9 line was previously described in [48,51]. All transgenic plants were generated by *Agrobacterium tumefaciens* strain AGL1 using floral dip [68].

Plasmid construction

To generate the SunTag:SDG2:FWA_g4 construct: the CDS of the catalytic domain of SDG2 (from amino acids 1571-2335) was amplified and coned into into the BsiWI linearised SunTag construct (previously described [28]), using In-Fusion (Takara). To generate dSDG2: the conserved amino acid predicted to abrogate catalytic activity when mutated (Y1903F) was identified using ConSurf [69]. Overlapping PCR was used to generate a version of SDG2 containing the desired mutation (see Supplementary Table 1), and the amplified PCR product was cloned into a BsiWI linearized SunTag-SDG2 construct by In-Fusion (Takara). To generate SunTag:PRDM9, the catalytic domain of mouse PRDM9 (110-417aa) (described in Policarpi et al 2024) was synthesised (TWIST Bioscience) and cloned into the SunTag using the same BsiWI linearization and In-Fusion method described above. Subsequently, overlap PCR was used to generate a version of PRDM9 with the G282A mutation (see Supplementary Table 1). The SNC1 (guides were chosen based on the predicted binding proximity to the TSS), LRCen3, and the no guides controls, were cloned using the protocol described in [70] (see Supplementary Table 1).

Chromatin Immunoprecipitation

The chromatin immunoprecipitation experiments were performed as previously described [71]. 2g of 14-day-old seedlings were collected per biological replicate. For

SunTag-SDG2 FWA and SunTag-SDG2 SNC1 input material are pooled T_2 plants. For

the SunTag:PRDM9:FWA_g4 in *rdr6*, 2 independent T_3 lines expressing FWA, 2

independent T_3 lines NoG were used alongside *rdr6* controls. 7 µl of anti-HA (3F10,

Merck) and 5 µl of anti-H3K4me3 (ab8580, abcam) were added to up to 2 ml of

sheared chromatin per sample for pulldown. The ChIP purified DNA was directly used

for ChIP-qPCR and ChIP-seq library preparation. Primer information is listed in

Suppletory Table 1. Values were normalised to Input ($2^{-(ChIP\ Ct - Input\ Ct)}*100$). ChIP-seq

libraries were generated using NuGen Ovation Ultra Low System V2 kits according to

manufacturer's instructions and were sequenced on an Illumina NovaSeq X

instrument with 150bp paired-end reads.

Crossover measurement

For the recombination rate experiment, SunTag plasmids were transformed into F_1

plants derived from a cross between the traffic line *CTL3.9* (containing insertions of

NAP:eGFP and NAP:dsRED) and Col-0 (Wu et al., 2015). Note that as the red and

green T-DNA insertions are separated by nearly 9-Mb, disruption from the T-DNAs

insertions (5-Kb and 12-Kb, respectively) are likely to be negligible at this scale [48]. To

help reduce the impact of transgene silencing in the absence of the *rdr6* background

(due to the impracticality of generating homozygous *rdr6* in the double heterozygous

CTL3.9 F1 background) and to ensure a sufficient number of scorable lines, we aimed

to screen 70-100 T_1's per constructs. T_1 transgenic plants (F_2 plants) were recovered

on hygromycin selection plates. Seeds harvested from T_1 (F_3) plants were first pre-

screened using a Leica DFC310 FX dissecting microscope with ultraviolet filters. Only

seeds which contained both red and green markers were cleaned to remove plant

debris and included in the analysis pipeline. The seed monolayer containing between

1,000 and 2,500 seeds was captured with a Leica DFC310 FX dissecting microscope,

first using brightfield followed by UV through a dsRED filter and then UV through a

GFP filter. Images were analysed using a modified CellProfiler pipeline (version 4.2.5)

described in [48,52,72]. For the F_4 data, seeds derived from the most highly recombining

and non-distorted line (P4) were sown (F_3), grown for a generation and seeds from

individual plants were collected (F_4) for recombination rate analysis as above, again

filtering seed sets from which either of the fluorescent markers had become

homozygous (presence / absence) in the previous generation.

Western blots

Western blot were performed as previously described [73] with minor modifications described below. For detection of the large dCAS9-10xGCN4 and SDG2 effector components as shown in Fig S1, single leaf punch (1cm diameter) samples from individual plants were used as input. Tissue was ground in a Tissuelyser II (Retsch) and resuspended in 100 ul of a 60:40 ratio of 2XSDS:8M Urea, mixed, and boiled at 95 degrees C for 5 minutes. Samples were centrifuged at >10,000xg to remove debris and loaded into a 3-8% tris acetate gel using MOPS running buffer. After electrophoresis, protein was transferred from the gel to a methanol activated PVDF membrane in Tobin Buffer (25mM Tris, 192mM Glycine, 20% methanol, 0.035% SDS). Membranes were blocked in 5% milk (w/v) in PBST (Phosphate Buffered Saline, 0.1% Tween20) and then incubated with HRP conjugated anti-HA at a 1:3,000 dilution in block. The blot was imaged by ECL detection.

For the western blots used to detect levels of H3K4me3, 100 mg of 7-day-old seedlings were frozen in liquid nitrogen and ground using a Tissuelyser II (Retsch). Total histones were extracted using the The EpiQuik™ Total Histone Extraction Kit (EpigenTek) according to the manufacturer's instructions. 5 µl of sample for H3 and 20 µl of sample for H3K4me3 were loaded onto a NuPage 4-12% BIS-Tris gel (Invitrogen) in NuPage MOPS SDS running buffer (Invitrogen). Transfer from gel to PVDF membrane was performed using an iBlot 3 machine with the dedicated transfer stack (Invitrogen). Following transfer, membrane was blocked in 5% milk (w/v) in TBST (Tris Buffered Saline, 0.1% Tween20) and shaken in room temp for 1 hour. Blocking buffer was discarded and membranes were incubated in primary antibody (anti-H3, ab1791 and anti-H3K4me3, ab8580, Abcam) in blocking buffer (1:1,000) shaking overnight in 4C. Following morning membrane was rinsed in TBST 5 times for 5 minutes in room temp. Secondary antibody was added for 1 hour (Goat anti-rabbit HRP, G-21234, Thermo). Membrane was washed again 5 times for 5 minutes. Detection was performed using a SuperSignal West Pico chemiluminescent substrate (#34080, Thermo) using an ImageQuant 800 (Amersham).

Fluorescent imaging

Roots of 7-day-old plants grown in vertical 1/2x MS solid media plates were imaged using a Leica DM6000B epifluorescent microscope.

Rosette size surface area measurement

3-week-old plants grown on soil were imaged using a Google Pixel 7 mounted on a
tripod. Images were then analyzed using Fiji.

Pseudomonas syringae colonisation assay

Infection assays were performed with *P. syringae* pathovar *tomato* strain DC3000
(*Pst*) strain expressing a stable chromosomal insertion of the *lux- CDABE* operon from
Photorhabdus luminescens (*Pst::LUX*) (Fan et al. 2008), using a slightly modified
version of a previously published protocol (Furci et al. 2021). Plants were grown on
1/2X MS solid media supplemented with 200 µg/ml Timentin in 96-well plates (Falcon)
for 7 days. Prior inoculation, *Pst::LUX* were grown in King's B media containing 50
µg/ml Kanamycin and 50 µg/ml Rifampicin overnight at 28 °C. Bacteria were pelleted
by centrifugation, washed in 10 mM MgSO4 and finally re-suspended to an OD600 of
0.2 in 10 mM MgSO4 containing 0.015% v/v Silwet-L77. 7 days-old seedlings were
sprayed with bacteria solution and 96-well plates were sealed with parafilm to maintain
100% relative humidity. Seedlings were imaged at 1-4 days post infection (dpi). Prior
imaging using (ImageQuant 800, Amersham), plates were left in the dark for 2 mins.
Bacterial bioluminescence images were acquired with 4 mins exposure time and bright
field images were taken using OD settings of the ImageQuant 800. Image-based
quantification of bioluminescence was carried out in Fiji. For each well, the well outline
was obtained from bright field images and added to ROI Manager. These ROIs were
then transposed onto the bioluminescence images. The bioluminescence intensity
from infected seedlings was obtained using the Fiji functions Analyse, Measure and
Mean value.

RT-qPCR

RNA was extracted from indicated plant material using TRIzol reagent (Invitrogen) and
the Direct-Zol RNA MiniPrep (Zymo) kit, including in-column DNase treatment
following the manufacturer's instructions. cDNA was synthesized using SuperScript IV
(Invitrogen). qRT-PCRs were performed using a Luna Universal qPCR Master Mix
(NEB) and a CFX connect Real-time PCR detection system (Bio-rad). For SunTag-
SDG2 targeting FWA and SNC1, RNA was extracted from 14-day-old whole seedlings
grown on 1/2x MS plates. SNC1 targeted lines were treated with *Pst::LUX* 4 hours

prior to harvesting. For SunTag:PRDM9 qRT-PCR, RNA was extracted from 4 weeks-

old leaf tissue of all T_1 lines and 2 independent T_3 lines per construct. 450 ng of total

RNA was used for cDNA synthesis. Primers are listed in Supplementary Table 1.

Quant-Seq Library construction

For Suntag-SDG2 QuantSeq, 350 ng of total RNA from ~10 pooled T_2 plants were

used as input material. For Suntag:PRDM9 Quant-seq, 350 ng of total RNA from 3 T_1

SunTag:PRDM9:FWA_g4 plants expressing FWA, 3 T_1 no guide SunTag:PRDM9

lines and *rdr6* and plants was used for library preparation. Libraries were prepared

using QuantSeq 3 'mRNA-Seq Library Prep Kit FWD (Lexogen) according to

manufacturer's instructions and sequenced on Illumina NovaSeq 6000 PE150

instrument.

Bioinformatic Analysis

ChIP-seq analysis

The ChIP-seq analysis pipeline was performed as previously described with minor

modifications. Bowtie2 (version 2.5.0) was used to map the PE150 bp read data in

fastq format to the TAIR10 genome (--no-unal), and were converted to bam format

using Samtools (version 1.10). Reads were de-duplicated using the samtools fixmate

and markdup commands (see https://github.com/C-Jake-Harris/Binenbaum-SunTag-

H3K4me3 for more details). Tracks were generated in DeepTools (version 3.5.1) using

bamCoverage (--normalizeUsing RPGC, --effectiveGenomeSize 135000000 --binSize

10) with multicopy regions blacklisted (--blackListFileName) using the regions

identified in [74]. For analysis of reads mapping to the centromere, a modified version of

the previously described pipeline [50] was used. Briefly, PE150 bp read data in fastq

format were mapped to the Col-CEN genome [47] using Bowtie2 (version 2.2.5) with (--

very-sensitive -k 200 --no-unal --no-discordant) and were filtered for primary

alignments using samtools view (-F 256 -q 5) prior to deduplication and track

generation, as above. Peaks were called using MACS2 (version 2.2.9.1) with default

parameters. multiBigwigSummary (BED-file --outRawCounts) was used to gather

normalised read enrichment over called peaks. Correlation plots were generated in

DeepTools using MultiBamSummary (--binSize 25) and PlotCorrelation (-c pearson --

removeOutliers --plotNumbers -p heatmap).

QuantSeq Analysis

In accordance with the manufacturer's (Lexogen) instructions, only read (*1.fq.gz)

from PE150 bp data was used for downstream analysis. Briefly, reads were trimmed

using cutadapt (version 1.18) and were mapped to the genome using STAR (version

2.7.10b) to the TAIR10 genome (--quantMode GeneCounts --alignIntronMax 10000 --

outSAMmultNmax 20). Gene counts were used as input for DESeq2 analysis in R.

Differentially expressed genes were defined using an adjusted p-value (Bonferroni

method) cutoff of <0.01.

gRNA binding analysis

For the LRCen3 binding and mismatch analysis, the LRCen3 guide RNA sequence

(5'-AGGCTTACAAGATTGGGTTG-3'), was mapped to the Col-CEN genome using

bowtie2 (version 2.2.5). The following options were used for no mismatches (-f -a --

end-to-end --np 0 --score-min L,0,0) and for 1 mismatch (-f -a --end-to-end --score-

min L,0,-1). Mapped reads were sorted and converted to bam format using Samtools

(version 1.10), and subsequently converted to .bed format using the bamtobed

function in bedtools (2.20.1). Bed files were used for downstream analysis in R to

generate chromosome-wide density plots over 100kb regions.

Statistics and reproducibility

Statistical significance was determined by using unpaired two-tailed Student's t-test

for two-group comparisons and ANOVA with post-hoc Tukey HSD for multiple

comparisons. Asterisks indicate significant differences (*$P < 0.05$, **$P < 0.01$,

***$P < 0.001$). Different lowercase letters indicate significant differences ($P < 0.05$). No

statistical method was used to predetermine sample size, no data were excluded from

the analyses, the experiments were not randomized, and the investigators were not

blinded to allocation during experiments and outcome assessment.

DATA AVAILABILITY:

The high-throughput sequencing data generated has been deposited to NCBI

(GSE288686 [https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE288686].

Previously published datasets: CENH3 ChIP-seq and input are on ArrayExpress (E-

MTAB-11974, [https://www.ebi.ac.uk/biostudies/arrayexpress/studies/E-MTAB-

11974])[47,50]. Custom scripts used are available on GitHub (https://github.com/C-Jake-

Harris/Binenbaum-SunTag-H3K4me3). Source Data are provided with this paper.

REFERENCES:

1. Lloyd, J. P. B. & Lister, R. Epigenome plasticity in plants. *Nat. Rev. Genet.* **23**, 55–68 (2022).

2. Schwaiger, M. *et al.* Evolutionary conservation of the eumetazoan gene regulatory landscape. *Genome Res.* **24**, 639–650 (2014).

3. Wang, H. & Helin, K. Roles of H3K4 methylation in biology and disease. *Trends Cell Biol.* **xx**, 1–14 (2024).

4. Howe, F. S., Fischl, H., Murray, S. C. & Mellor, J. Is H3K4me3 instructive for transcription activation? *BioEssays* **39**, 1–12 (2017).

5. Perez, M. F. & Sarkies, P. Histone methyltransferase activity affects metabolism in human cells independently of transcriptional regulation. *PLoS Biol.* **21**, 1–31 (2023).

6. Vermeulen, M. *et al.* Selective Anchoring of TFIID to Nucleosomes by Trimethylation of Histone H3 Lysine 4. *Cell* **131**, 58–69 (2007).

7. Lauberth, S. M. *et al.* H3K4me3 interactions with TAF3 regulate preinitiation complex assembly and selective gene activation. *Cell* **152**, 1021–1036 (2013).

8. Wang, H. *et al.* H3K4me3 regulates RNA polymerase II promoter-proximal pause-release. *Nature* **615**, 339–348 (2023).

9. Xie, S. S. *et al.* JMJ28 guides sequence-specific targeting of ATX1/2-containing COMPASS-like complex in Arabidopsis. *Cell Rep.* **42**, 112163 (2023).

10. Shang, J. Y. *et al.* COMPASS functions as a module of the INO80 chromatin remodeling complex to mediate histone H3K4 methylation in Arabidopsis. *Plant Cell* **33**, 3250–3271 (2021).

11. Xue, M. *et al.* Single amino acid mutations in histone H3.3 illuminate the functional significance of H3K4 methylation in plants. *Nat. Commun.* **16**, (2025).

12. Xiao, J., Lee, U. & Wagner, D. ScienceDirect Tug of war : adding and removing histone lysine methylation in Arabidopsis. *Curr. Opin. Plant Biol.* **34**, 41–53 (2016).

13. Xu, L. *et al.* H3K4me3 binding ALFIN - LIKE proteins recruit SWR1 for gene - body deposition of H2A . Z. *Genome Biol.* (2025). doi:10.1186/s13059-025-03605-7

14. Borde, V. *et al.* Histone H3 lysine 4 trimethylation marks meiotic recombination initiation sites. *EMBO J.* **28**, 99–111 (2009).

15. Fayos, I. *et al.* Manipulation of Meiotic Recombination to Hasten Crop Improvement. *Biology (Basel).* **11**, 1–15 (2022).

16. Taagen, E., Bogdanove, A. J. & Sorrells, M. E. Counting on Crossovers: Controlled Recombination for Plant Breeding. *Trends Plant Sci.* **25**, 455–465 (2020).

17. Rowan, B. A. *et al.* An Ultra High-Density Arabidopsis thaliana Crossover. *Genetics* **213**, 771–787 (2019).

18. Tock, A. J. *et al.* Crossover-active regions of the wheat genome are distinguished by DMC1, the chromosome axis, H3K27me3, and signatures of adaptation. *Genome Res.* **31**, 1614–1628 (2021).

19. Powers, N. R. *et al.* The Meiotic Recombination Activator PRDM9 Trimethylates Both H3K36 and H3K4 at Recombination Hotspots In Vivo. *PLoS Genet.* **12**, 1–24 (2016).

20. Choi, K. *et al.* Arabidopsis meiotic crossover hot spots overlap with H2A.Z nucleosomes at gene promoters. *Nat. Genet.* **45**, 1327–1338 (2013).

21. Policarpi, C. & Hackett, J. A. Epigenetic editing : Dissecting chromatin function. *BioEssays* 1–16 (2021).

22. Villiger, L. *et al.* CRISPR technologies for genome, epigenome and transcriptome editing. *Nat. Rev. Mol. Cell Biol.* (2024). doi:10.1038/s41580-023-00697-6

23. Wang, X. & Yamaguchi, N. Cause or effect: Probing the roles of epigenetics in plant development and environmental responses. *Curr. Opin. Plant Biol.* **81**, 102569 (2024).

24. Dubois, A. & Roudier, F. Deciphering Plant Chromatin Regulation via CRISPR/dCas9-Based Epigenome Engineering. *Epigenomes* **5**, 17 (2021).

25. Harris, C. J., Amtmann, A. & Ton, J. Epigenetic processes in plant stress priming : Open questions and new approaches. *Curr. Opin. Plant Biol.* **75**, 102432 (2023).

26. Tanenbaum, M. E., Gilbert, L. a, Qi, L. S., Weissman, J. S. & Vale, R. D. A versatile protein tagging system for signal amplification in single molecule imaging and gene regulation. *Cell* **159**, 635–646 (2014).

27. Wang, M. *et al.* A gene silencing screen uncovers diverse tools for targeted gene repression in Arabidopsis . *Nat. Plants* (2023). doi:10.1038/s41477-023-01362-8

28.   Papikian, A., Liu, W., Gallego-bartolomé, J. & Jacobsen, S. E. Site-specific
manipulation of Arabidopsis loci using CRISPR-Cas9 SunTag systems. *Nat.*
Commun. **10**, (2019).

29.   Gallego-Bartolomé, J. *et al.* Targeted DNA demethylation of the *Arabidopsis*
genome using the human TET1 catalytic domain. *Proc. Natl. Acad. Sci.* **115**,
201716945 (2018).

30.   Pflueger, C. *et al.* A modular dCas9-SunTag DNMT3A epigenome editing
system overcomes pervasive off-target activity of direct fusion dCas9-DNMT3A
constructs. *Genome Res.* **28**, 1193–1206 (2018).

31.   Policarpi, C., Munafò, M., Tsagkris, S., Carlini, V. & Hackett, J. A. Systematic
epigenome editing captures the context-dependent instructive function of
chromatin modifications. *Nat. Genet.* **56**, 1168–1180 (2024).

32.   Cheng, Y., Zhou, Y. & Wang, M. Targeted gene regulation through epigenome
editing in plants. *Curr. Opin. Plant Biol.* **80**, 102552 (2024).

33.   Selma, S. & Orzáez, D. Perspectives for epigenetic editing in crops. *Transgenic*
Res. **30**, 381–400 (2021).

34.   Guo, L., Yu, Y., Law, J. A. & Zhang, X. Set domain group2 is the major histone
H3 lysine 4 trimethyltransferase in Arabidopsis. *Proc. Natl. Acad. Sci. U. S. A.*
107, 18557–18562 (2010).

35.   Gardiner, J., Ghoshal, B., Wang, M. & Jacobsen, S. E. CRISPR–Cas-mediated
transcriptional control and epi-mutagenesis. *Plant Physiol.* **188**, 1811–1824
(2022).

36.   Du, J. *et al.* Mechanism of DNA methylation-directed histone methylation by
KRYPTONITE. *Mol. Cell* **55**, 495–504 (2014).

37.   Soppe, W. J. *et al.* The late flowering phenotype of fwa mutants is caused by
gain-of-function epigenetic alleles of a homeodomain gene. *Mol. Cell* **6**, 791–
802 (2000).

38.   Wang, M. *et al.* Histone H3 lysine 4 methylation recruits DNA demethylases to
enforce gene expression in Arabidopsis. *Nat. Plants* **11**, 206–217 (2025).

39.   Johnson, L. M. *et al.* SRA- and SET-domain-containing proteins link RNA
polymerase V occupancy to DNA methylation. *Nature* **507**, 124–128 (2014).

40.   Wang, M. Y. *et al.* The plant immune receptor SNC1 monitors helper NLRs
targeted by a bacterial effector. *Cell Host Microbe* **31**, 1792-1803.e7 (2023).

41.   Yi, H. & Richards, E. J. Gene duplication and hypermutation of the pathogen

Resistance gene SNC1 in the arabidopsis bal variant. *Genetics* **183**, 1227–1234 (2009).

42. Yi, H. & Richards, E. J. A cluster of disease resistance genes in Arabidopsis is coordinately regulated by transcriptional activation and RNA silencing. *Plant Cell* **19**, 2929–39 (2007).

43. Stokes, T. L., Kunkel, B. N. & Richards, E. J. Epigenetic variation in Arabidopsis disease resistance. *Genes Dev.* **16**, 171–182 (2002).

44. Yang, L., Wang, Z. & Hua, J. Multiple chromatin-associated modules regulate expression of an intracellular immune receptor gene in Arabidopsis. *New Phytol.* **237**, 2284–2297 (2023).

45. Furci, L., Pardo, D. P. & Ton, J. A rapid and non - destructive method for spatial – temporal quantification of colonization by Pseudomonas syringae pv . tomato DC3000 in Arabidopsis and tomato. *Plant Methods* 1–8 (2021). doi:10.1186/s13007-021-00826-2

46. Yelina, N. E. *et al.* Epigenetic remodeling of meiotic crossover frequency in Arabidopsis thaliana DNA methyltransferase mutants. *PLoS Genet.* **8**, e1002844 (2012).

47. Naish, M. *et al.* The genetic and epigenetic landscape of the Arabidopsis centromeres. *Science (80-.).* **374**, (2021).

48. Fernandes, J. B. *et al.* Structural variation and DNA methylation shape the centromere-proximal meiotic crossover landscape in Arabidopsis. *Genome Biol.* **25**, 1–31 (2024).

49. Yelina, N. E. *et al.* DNA methylation epigenetically silences crossover hot spots and controls chromosomal domains of meiotic recombination in Arabidopsis. *Genes Dev.* **29**, 2183–2202 (2015).

50. Wlodzimierz, P. *et al.* Cycles of satellite and transposon evolution in Arabidopsis centromeres. *Nature* **618**, 557–565 (2023).

51. Wu, G., Rossidivito, G., Hu, T., Berlyand, Y. & Poethig, R. S. Traffic lines: New tools for genetic analysis in Arabidopsis thaliana. *Genetics* **200**, 35–45 (2015).

52. Kbiri, N., Dluzewska, J., Henderson, I. R. & Ziolkowski, P. A. Quantifying Meiotic Crossover Recombination in Arabidopsis Lines Expressing Fluorescent Reporters in Seeds Using SeedScoring Pipeline for CellProfiler. in *Plant Gametogenesis. Methods in Molecular Biology* (ed. Lambing, C.) 121–134 (Springer US, 2022). doi:10.1007/978-1-0716-2253-7_10

53.   Wang, Y. *et al.* The Arabidopsis DREAM complex antagonizes WDR5A to
modulate histone H3K4me2/3 deposition for a subset of genome repression.
Proc. Natl. Acad. Sci. U. S. A. **119**, 1–9 (2022).

54.   Underwood, C. J. *et al.* Epigenetic activation of meiotic recombination near
Arabidopsis thaliana centromeres via loss of H3K9me2 and non-CG DNA
methylation. *Genome Res.* **28**, 519–531 (2018).

55.   Leng, X., Thomas, Q., Rasmussen, S. H. & Marquardt, S. A
G(enomic)P(ositioning)S(ystem) for Plant RNAPII Transcription. *Trends Plant*
Sci. **25**, 744–764 (2020).

56.   Marquardt, S., Petrillo, E. & Manavella, P. A. Cotranscriptional RNA processing
and modification in plants. *Plant Cell* **35**, 1654–1670 (2023).

57.   Rickels, R. *et al.* Histone H3K4 monomethylation catalyzed by Trr and
mammalian COMPASS-like proteins at enhancers is dispensable for
development and viability. *Nat. Genet.* **49**, 1647–1653 (2017).

58.   Fal, K., El Khoury, S., Le Masson, M., Berr, A. & Carles, C. C. CRISPR/dCas9-
targeted H3K27me3 demethylation at the CUC3 boundary gene triggers ectopic
transcription and impacts plant development. *iScience* **28**, 112475 (2025).

59.   Cuimei Zhang, Yajun Tang, Shanjie Tang, Lei Chen, Haidi Yuan, Yujun Xu,
Yangyan Zhou, Shuaibin Zhang, Jianli Wang, Hongyu Wen, Wenbo Jiang,
Yongzhen Pang, Xian Deng, Xiaofeng Cao, Junhui Zhou, Xianwei Song, Q. L.
P. An Inducible CRISPR-activation tool for accelerated plant regeneration
Cuimei. *Plant Commun.* (2024).

60.   Song, X. *et al.* Inheritance of acquired adaptive cold tolerance in rice through
DNA methylation. *Cell* 1–12 (2025). doi:10.1016/j.cell.2025.04.036

61.   Oberkofler, V. & Bäurle, I. Inducible epigenome editing probes for the role of
histone H3K4 methylation in Arabidopsis heat stress memory. *Plant Physiol.*
703–714 (2022). doi:10.1093/plphys/kiac113

62.   Casas-Mollano, J. A., Zinselmeier, M. H., Sychla, A. & Smanski, M. J. Efficient
gene activation in plants by the MoonTag programmable transcriptional
activator. *Nucleic Acids Res.* **51**, 7083–7093 (2023).

63.   Pan, C. *et al.* CRISPR–Act3.0 for highly efficient multiplexed gene activation in
plants. *Nat. Plants* **7**, 942–953 (2021).

64.   de Melo, B. P. *et al.* Transcriptional modulation of AREB-1 by CRISPRa
improves plant physiological performance under severe water deficit. *Sci. Rep.*

10, 1–10 (2020).

65.   Boone, B. A. *et al.* ACD15, ACD21, and SLN regulate the accumulation and
870   mobility of MBD6 to silence genes and transposable elements. *Sci. Adv.* **9**, 1–
871   14 (2023).

872   66.   Baldini, A., Battaglia, F. & Perrella, G. The generation of novel epialleles in
873   plants: the prospective behind re-shaping the epigenome. *Front. Plant Sci.* **16**,
874   (2025).

875   67.   Lu, Z. *et al.* The prevalence, evolution and chromatin signatures of plant
876   regulatory elements. *Nat. Plants* **5**, 1250–1259 (2019).

877   68.   Clough, S. J. & Bent, A. F. Floral dip: A simplified method for Agrobacterium-
878   mediated transformation of Arabidopsis thaliana. *Plant J.* **16**, 735–743 (1998).

879   69.   Ashkenazy, H. *et al.* ConSurf 2016: an improved methodology to estimate and
880   visualize evolutionary conservation in macromolecules. *Nucleic Acids Res.* **44**,
881   W344–W350 (2016).

882   70.   Ghoshal, B. & Gardiner, J. CRISPR-dCas9-Based Targeted Manipulation of
883   DNA Methylation in Plants. *Springer Protoc. Handbooks* **2**, 57–71 (2021).

884   71.   Harris, C. J., Zhong, Z., Ichino, L., Feng, S. & Jacobsen, S. E. H1 restricts
885   euchromatin-associated   methylation   pathways   from   heterochromatic
886   encroachment. *Elife* 1–17 (2024).

887   72.   Gorringe, N., Topp, S., Burns, R., Yamaguchi, S. & Fernando, A. Natural
888   variation modifies centromere-proximal meiotic crossover frequency and
889   segregation distortion in Arabidopsis thaliana. *bioRxiv* 1–69 (2025).

890   73.   Ichino, L. *et al.* MBD5 and MBD6 couple DNA methylation to gene silencing
891   through the J-domain protein SILENZIO. *Science (80-.).* **1439**, 1434–1439
892   (2021).

893   74.   Klasfeld, S., Roulé, T. & Wagner, D. Greenscreen: A simple method to remove
894   artifactual signals and enrich for true peaks in genomic datasets including ChIP-
895   seq data. *Plant Cell* **34**, 4795–4815 (2022).

896   75.   Abramson, J. *et al.* Accurate structure prediction of biomolecular interactions
897   with AlphaFold 3. *Nature* **630**, 493–500 (2024).

ACKNOWLEDGEMENTS

We would like to thank Jurriaan Ton's group at the University of Sheffield for their generous gift of the *Pst::LUX* strain, and for their support in helping to establish the infection and quantification protocol in the Harris group. **Funding:** C.J.H. is supported by a Royal Society University Research Fellowship (URF\R1\201016) and an ERC Starting Grant (TransPlantMemory), funded by UKRI EPSRC (EP/X025306/1). J.B. is supported by a Marie Curie postdoctoral fellowship, funded by UKRI (EP/Z001749/1). V.A. is supported by a 4-year BBSRC-funded doctoral studentship. I.R.H is supported by UKRI/BBSRC grants BB/Y009487/1 and BB/V003984/1, and an ERC Advanced Grant (EvoPanCen). N.G. is supported by a BBSRC-UKRI doctoral studentship. S.E.J. is an investigator of the Howard Hughes Medical Institute.

AUTHOR CONTRIBUTIONS

C.J.H., I.R.H. and S.E.J. conceived, designed and supervised the research. C.J.H. performed experiments, NGS bioinformatic analysis, and assembled the manuscript. J.B. performed the majority of the experiments relating to SunTag:SDG2. V.A. performed the experiments relating to SunTag:PRDM9. H.F. performed the majority of experiments and analysis relating to meiotic recombination. L.X. performed the SunTag:SDG2 centromere-targeting ChIP-seq. N.G. provided the Col-0 x *CTL3.9* F_1 seeds and guidance on the analysis of recombination frequency. P.W. and R.B. contributed to the design of the centromere targeting guide RNA. A.P. cloned SDG2 into SunTag.

COMPETING INTERESTS

The authors declare no competing interests.