Synergistic *MAPT* mutations as a platform to uncover modifiers of tau pathogenesis

Miles R. Bryan III[1,2], Michael F. Almeida[2,3,4], Kyle Pellegrino[1,4], Carli K. Opland[1], Ethan Paulakonis, J.[1] McGillion-Moore[1], Hanna Trzeciakiewicz[1], Diamond King[1], Xu Tian[1], Jui-Heng Tseng[1,2,5], Jonathan C. Schisler[3,4], Nicholas G. Brown[4], Ben A. Bahr[2,6], Todd J. Cohen[1,2*]

[1]Neurology, University of North Carolina- Chapel Hill, NC
[2]Duke-University of North Carolina Alzheimer's Disease Research Center (ADRC), 2424 Erwin Road, Durham, North Carolina 27705
[3]McAllister Heart Institute and the Department of Pharmacology, 111 Mason Farm Road, University of North Carolina - Chapel Hill, Chapel Hill, North Carolina 27514 U.S.A.
[4]Pharmacology, University of North Carolina- Chapel Hill, NC
[5]School of Life Sciences, Arizona State University- Tempe, AZ
[6]Biotechnology Research and Training Center, University of North Carolina- Pembroke, NC

From the Department of Neurology, UNC Neuroscience Center, University of North Carolina, Chapel Hill North Carolina, 27599

*To whom correspondence should be addressed: Todd J. Cohen: Department of Neurology, University of North Carolina at Chapel Hill, Chapel Hill NC 27599
toddcohen@neurology.unc.edu; Tel. (919) 966-3759; Fax. (919) 843-4576

Keywords*: aggregation, Alzheimer's disease, FTDP-17, MAPT, posttranslational modification (PTM), heat-shock proteins, Hsp70, organotypic brain slices, neurodegeneration, tau, tauopathy

Abstract

The natively unfolded tau (MAPT) protein is extremely soluble, which poses challenges when

modeling neurofibrillary tangle (NFT) pathology in Alzheimer's disease (AD). To overcome this

hurdle, we combined P301L and S320F mutations (PL-SF) to generate a rapid and reliable tau

pathology platform to expedite the discovery of factors that modify tau aggregation. Using this

model, we evaluated heat-shock proteins (Hsp), which have been linked to tau pathology, but

whose role in AD remains enigmatic and controversial. In primary neurons, expression of Hsp70,

but not Hsc70 or Hsp90, exacerbated tau aggregation. Conversely, lowering Hsp70 or employing

a chaperone-deficient tau mutant (PL-SF-4Δ) reduced tau phosphorylation and abrogated tau

aggregation, highlighting Hsp70 as a key driver of tau aggregation. Hsp70 foci clustered within

and surrounding neuritic plaques and NFTs in post-mortem AD brain. Functionally, mature aggregate-bearing neurons showed deficits in neuronal firing and network communication, which were restored by chaperone-binding deficient tau variants that abrogated tau pathology. This study provides a powerful cell-intrinsic model for accelerated tau aggregation, which can be harnessed to identify potent regulators of tau aggregation as promising therapeutic targets.

Introduction

Aggregation of the microtubule-associated protein tau (MAPT) and the formation of insoluble tau inclusion pathology are hallmark features of Alzheimer's disease and related neurodegenerative disorders. In contrast to AD, in which tau pathology is comprised of normal wild-type tau proteins, pathogenic disease-causing mutations in *MAPT* are implicated in familial frontotemporal dementia linked to chromosome 17 (FTDP-17). Over 50 different pathogenic *MAPT* missense, silent, and intronic mutations have been identified[1]. In many instances, missense mutations generate loss and gain of function phenotypes including altered microtubule (MT) binding and enhanced tau aggregation propensity, which are implicated in prion-like tau spreading along defined spatial and temporal brain networks.

While familial missense mutations at residue P301 (e.g., P301L and P301S) have been well characterized as aggregate-prone and MT-binding deficient, less is known about nearby mutations within the MT-binding repeat domain (MTBR). Our recent study characterized a panel of FTDP-17 tau mutations spanning this domain and found that the cellular surveillance machinery composed of HDAC6 and associated chaperones (e.g. Hsp70) shows increased binding to two specific mutations within the R3 domain (P301L and S320F), but not others, suggesting these two

mutants are conformationally distinct and are more effectively recognized by the quality control

machinery that targets tau for degradation[2].

In separate studies, the P301L and S320F mutations were combined into the same full-

length tau molecule (termed PL-SF), which generated a surprisingly aggregate-prone

conformation leading to an accelerated AD-like tangle pathology, neuroinflammation, and

memory impairments in mice and rhesus macaques[3–7]. Thus, PL-SF could potentially be leveraged

as an intrinsic, cell-autonomous, and seeding-independent model of full-length tau aggregation,

which would otherwise be challenging to achieve with full-length tau. In fact, many widely used

tau biosensor models employ artificial, truncated tau mutants comprising only the tau repeat

domain, a fragment with questionable physiological relevance to neurons[8]. Once fully

characterized, the PL-SF model offers several advantages, including serving as a rapid model of

mature tau aggregation and as a cornerstone for high-throughput discovery of tau modifiers.

The extent to which tau-bound chaperones such as inducible Hsp70, constitutive Hsc70,

and related Hsp90 modulate tau function remains unclear. This debate has obvious therapeutic

implications since Hsp inhibitors are in clinical development and have been evaluated in

preclinical models of AD[9–14]. However, prior studies have yielded quite contradictory and even

opposing results, since on the one hand, Hsp70 facilitates the MC1-positive tau conformation[15],

while other studies suggest Hsp70 and Hsc70 mediate tau degradation via autophagy and the

ubiquitin-proteasome system (UPS)[16,17]. One possible explanation for these discrepancies is the

inability to generate mature tau pathology as observed in human AD brains. Adding additional

complexity, tau interacts with dozens of chaperone adaptors and cofactors that may promote or

suppress chaperone function directed towards tau[18–21]. Here, we characterized a tau PL-SF model

of accelerated tau pathology, which allowed us to interrogate the role of Hsp70 and other

chaperones as bona fide modifiers of tau-mediated toxicity and aggregation.

Results

Synergistic familial *MAPT* mutations drive tau phosphorylation and aggregation

We first evaluated tau aggregation in PL-SF expressing immortalized QBI-293 cells. Cells

transfected with human T40 (2N4R) wild-type, P301L, S320F, or PL-SF were monitored for

RIPA-soluble expression of phosphorylated tau (AT8 and AT100) and cleaved tau (TauC3), two

well-established markers of tau pathogenesis. Phosphorylated tau accumulated in PL-SF

expressing cells, more so than WT or either single mutant (**Figure 1 A, B**). Confirming this finding,

total tau immunoblotting revealed up-shifted tau protein bands in SF and PL-SF, indicative of tau

hyperphosphorylation. Surprisingly, the single PL mutant appeared hypo-phosphorylated

compared to WT or SF mutations. Immunocytochemistry confirmed these findings and revealed

Thioflavin S (ThioS) positive SF and PL-SF aggregates, consistent with enhanced tau aggregation

(**Supplemental Figure 1, Figure 1C**). Similar to phosphorylated tau, cleaved tau also

preferentially accumulated in SF or PL-SF expressing cells.

Multiple familial FTD-associated mutations occur at residue P301 (P301L, P301S, P301T)

and S320 (S320F, S320Y). We evaluated a panel of other combination mutants by immunoblotting

and immunocytochemistry and found that all double mutants formed phosphorylated, cleaved, and

insoluble tau species compared to individual single tau mutants (**Supplemental Figure 2**). The

most consistently insoluble tau species was PL-SF, which we selected for follow-up analysis.

A prior study found that deletion of four key residues in tau (I277/I278 and I308/V309) can abrogate Hsp70 binding[22]. Since the role of Hsp70 in tau pathology is unclear, we generated a PL-SF mutant in which these four residues were depleted, termed PL-SF 4Δ. In transfected QBI-293 cells, PL-SF 4Δ was nearly completely dephosphorylated and accumulated negligible amounts of cleaved tau. PL-SF 4Δ was also downshifted compared to PL-SF (**Figure 1D, E**) and showed significantly less phosphorylated (AT8), cleaved (TauC3), and aggregated (ThioS) tau (**Supplemental Figure 3, Figure 1F**). By co-immunoprecipitation assays, we confirmed that PL-SF 4Δ abrogated the majority of Hsc70 and Hsp70 binding to tau. We did not observe detectable and specific binding to endogenous Hsp90 in our model (**Figure 1G**).

We also assessed the impact of tau mutants on acetylated-tubulin (ac-tubulin), a readout for microtubule (MT) stability and dynamics. Wild-type tau increased ac-tubulin, as expected, while P301L or S320F were less effective (~30-40%). Consistent with a near-complete loss of MT regulatory function, PL-SF reduced ac-tubulin levels down to the levels of non-transfected cells. Surprisingly, PL-SF 4Δ partly restored the tubulin interactions (**Figure 1H, I**).

To confirm that PL-SF has increased intrinsic propensity to self-aggregate, as opposed to being influenced by other cellular factors, we performed in vitro Thioflavin T (ThioT) assays using recombinant purified tau proteins to assess aggregation kinetics in the presence of heparin as an inducing agent. Indeed, PL-SF tau rapidly aggregated within 1 hr post-induction, at which point the formation of recombinant tau fibrils plateaued, while PL-SF 4Δ was completely resistant to aggregation, even though similar tau concentrations were present (**Figure 1J**).

Impaired chaperone binding alleviates tau pathology in neurons

To validate our findings in neurons, we developed mutant tau lentiviral constructs suitable

for high-efficiency transduction into mouse primary cortical neurons.   Transduced primary

neurons were evaluated with a panel of tau mutants, including PL-SF. Immunoblotting of neuronal

lysates showed that PL-SF tau was again up-shifted and accumulated more phosphorylated tau

(**Figure 2A, B).** Unlike immortalized cell lines, neurons did not show appreciable levels of cleaved

tau (TauC3). Only PL-SF accumulated in the insoluble fraction (**Figure 2C**). PL-SF 4Δ showed a

marked reduction in tau phosphorylation and aggregation, suggesting a similar phenotype in

neurons as well. In fact, deletion of just two of the Hsp-binding residues (ΔI277/I278 or

ΔI308/V309) abrogated tau phosphorylation and aggregation, with residues I308/V309 showing

the most dominant effect (**Figure 2A-C**). Introduction of β-sheet disrupting proline mutations has

been shown to abrogate tau aggregation[23–27]. Indeed, by mutating residues I277 and I308 to

prolines (PL-SF I277P/I308P), we were able to suppress tau aggregation to similar levels as seen

in PL-SF 4Δ (**Supplemental Figure 4**).

By performing microscopy on transduced neurons, we found that PL-SF accumulated AT8,

MC1, and ThioS-positive tau inclusions (**Figure 2D, E**). Both MC1 (pre-tangle confirmational

tau) and AT8 immunoreactive tau were amorphous and globular and accumulated as large

varicosities, or swellings, within neurites. The large globular tau aggregates colocalized with

ThioS, supporting tangle-like pathology (**Figure 2F, G**). By transducing neuronal cultures with

GFP-tagged PL-SF and labeling neurons with a mouse-specific tau antibody (monoclonal T49),

we observed recruitment of endogenous mouse tau to PL-SF inclusions, implying that no species

barrier exists between human PL-SF aggregates and mouse tau (**Figure 2H**). PL-SF behaved

similarly in tau knockout neurons, and therefore, endogenous tau is not required for the

accumulation of PL-SF pathology in neurons (**Supplemental Figure 5**).

Hsp70 promotes tau hyperphosphorylation and aggregation

To determine whether Hsp70 is a more potent regulator of tau pathology compared to other tau-associated heat shock proteins, PL-SF expressing neurons were co-transduced with either HSP70, HSC70, or HSP90. Only Hsp70, but not Hsc70 or Hsp90, exacerbated tau phosphorylation (AT8), which was supported by an appreciable upshift in tau's molecular weight (**Figure 3A, B**) and a trend towards increased levels of total insoluble tau (**Figure 3C, D**). Hsp70 significantly increased AT8, MC1, and ThioS immunoreactivity (**Figure 3E, F**). The effect of Hsp70 appears remarkably specific to the aggregate-prone PL-SF tau, since the P301L single mutant was unaffected **(Supplemental Figure 6).**

Next, we depleted HSP70 in PL-SF transduced neurons. We achieved ~ 50% Hsp70 knockdown using lentiviral shRNA transduction, which did not affect HSC70 expression. Strikingly, HSP70 knockdown led to ~70% decrease in phosphorylated tau (**Figure 4 A, B**) and ~50% decrease in total insoluble tau (**Figure 4C, D**). We note that a ~50% decrease in soluble tau was also observed, suggesting that Hsp70 depletion enhances the turnover of steady-state tau levels (**Figure 4A, B**). The impact of Hsp70 on tau also appears to be quite specific since knockdown of HSP90 did not alter total or phosphorylated tau (**Supplementary Figure 7**). Overall, our findings highlight Hsp70 as a primary regulator of aggregate-prone tau species in this accelerated PL-SF tau model.

Cell-autonomous aggregated tau alters neuronal activity and network synchrony

Given the extensive tau pathology observed in PL-SF transduced neurons, we evaluated any deficits in neuronal activity and connectivity. Electrically active neurons were transduced at DIV18 following an initial multi-electrode array (MEA) recording. Daily recordings were taken

for 53 days, and in parallel, neurons were immunolabeled at DIV29 and DIV62 to assess the extent

of tau pathology.

We evaluated the mean firing rate (a measure of overall activity), the number of network

bursts (a measure of neuronal connectivity), and the area under the normalized crossed-correlation

(a readout for neuronal synchrony). After 15-20 days post-lentiviral transduction, PL-SF neurons

showed a decline in all three metrics, suggesting a loss in neuronal activity and reduced

synchronous firing across neuronal networks. The deficits worsened over time, but some neurons

continued to fire even past DIV50 (**Figure 5A-C**). Raster plots at DIV54 highlight reduced

network firing across electrodes and elongated network firing time (**Figure 5D-F**). PL-SF 4Δ did

not exhibit any deficits in any of the activity metrics and showed similar firing patterns as

compared to wild-type control neurons (**Figure 5A-C**).

The impact of aggregate-prone tau mutants on neuronal firing could represent a direct

effect of tau at the synapse. Therefore, we assessed whether PL-SF accumulated at the post-

synaptic density (PSD), a re-localization thought to coincide with synaptic dysfunction. PSD

fractions were biochemically isolated from transduced neurons at DIV18 and probed by

immunoblotting for phosphorylated and total tau. PL-SF accumulated in the PSD fraction, which

was not observed with PL-SF 4Δ or any single tau mutants (e.g., P301L) (**Figure 5G**). We

conclude that mature PL-SF tau aggregates accumulate at the PSD and drive synaptic dysfunction.

By abrogating Hsp binding, PL-SF 4Δ bypasses these defects.

PL-SF expressing neurons were aged in culture to determine tau phenotypes over time.

Indeed, aged neurons accumulated dense tau inclusions comprised of phosphorylated, MC1, and

ThioS-positive tau at DIV29, which became even more intensely labeled by DIV62. All indicators

of tau pathology were absent in PL-SF 4Δ expressing neurons, even at late time points (**Figure**

5H). We also note that Hsp70 began to colocalize with aggregated tau in aged neurons (**Figure**

5I), supporting aggregation-dependent recruitment of Hsp70.

Cell-autonomous tau pathology in organotypic brain slice cultures

To validate our findings in brain culture systems comprised of both neurons and glia, we

transduced adult rat organotypic hippocampal brain slices ex vivo with either GFP alone, wild-

type tau-GFP, PL-SF-GFP, or PL-SF 4Δ-GFP at DIV1 and imaged brain slices after nine weeks in

culture **(Figure 6A, B)**. Similar to cultured neurons, PL-SF accumulated dense AT8 and MC1-

positive inclusions and distinct neuritic puncta, or foci. Most tau pathology preferentially

accumulated in the dentate gyrus (DG) and extended through the hippocampal CA1 and CA3

regions. In contrast, PL-SF 4Δ exhibited diffuse, non-aggregated neuronal labeling in the DG and

showed far lower levels of pathogenic tau markers, such as MC1 and AT8 **(Figure 6C, D)**. These

data suggest that, in a multicellular model with complete hippocampal architecture, abrogating

tau-Hsp binding can confer neuroprotection.

Hsp70 is upregulated in human AD brain and intercalates within tau aggregates.

Finally, we evaluated Hsp70 levels and localization in AD human post-mortem brain.

Biochemical fractionation of control and AD brain lysates showed elevated Hsp70 expression in

both soluble (hi-salt extracted) and insoluble (urea extracted) whole cortex AD brain homogenates

(Figure 7A,B). Hsc70 was only upregulated in the insoluble fraction of AD brains (**Figure 7C,D**).

We then performed immunohistochemistry on AD hippocampus and found that the vast majority

of AT8-positive tau aggregates (plaques and tangles) were labeled by Hsp70-positive foci (**Figure**

7E, Supplemental Figure 8). Interestingly, the Hsp70 foci formed discrete puncta that appeared

to intercalate within the AT8-positive tau neuritic plaques and neurofibrillary tangles (NFTs). This

data further supports a prominent role for Hsp70 in modulating tau accumulation in human AD

brain.

Discussion

Here, we developed and characterized a cell-autonomous model of tau aggregation based

on the synergistic aggregation of tau missense mutations at residues P301 and S320. Given the

rapid and robust tau aggregation observed within days (in primary neurons), the PL-SF model

readily lends itself to identifying factors that modulate tau aggregation. We leveraged this model

to clarify the longstanding debate on the role of Hsp70 as a tau modifier. The PL-SF model could

address the stepwise development of tau pathology and provide a more reliable and sensitive

system for identifying subtle regulators of tau pathogenesis.

Synergistic tau mutants as a model of mature tauopathy

Prior studies have suggested that combining tau mutations at P301 and S320 (e.g. PL-SF)

can lead to accelerated tau aggregation[3–6,28], leading to the development of new cell culture and

mammalian models that better recapitulate tau pathology. For example, the S320F mutation was

introduced in P301S/S305N knock-in mice, resulting in enhanced tau pathology[29]. Recently,

AAV-mediated delivery of PL-SF tau to rhesus macaques was able to induce rapid and progressive

tau pathology, demonstrating its utility as an efficient primate model for studying AD[30] . Here,

we analyzed the biochemical and histological properties of PL-SF in cells, cortical neurons, and

organotypic brain slice cultures. Strikingly, PL-SF tau inclusions appear quite similar to those

detected in post-mortem AD brain (**Figures 1, 2, 6, 7**). Future cryo-EM studies may be needed to

better define their ultrastructural properties[31–34]. However, if PL-SF does indeed form AD-like

filaments, it may be a preferred model over other single tau variants (P301S and P301L) that lack

mature amyloid structure in culture and mouse models[35,36]. The PL-SF tau variant also

accumulated at the post-synaptic density (PSD), leading to functional deficits in neuronal firing

and connectivity (**Figure 5 A-H**), which supports its synaptic toxicity. This rapidly maturing

pathology likely allows for the rapid occurrence of sensitized functional defects at the synapse and

may also affect other cellular processes.

Exactly how these two mutations (P301L or S320F) synergize to promote tau pathology

and drive tau-mediated toxicity is unclear. But we can envision several scenarios. First, P301L and

P301S have been well-characterized for disrupting MT dynamics and promoting tau aggregation;

however, less is known about S320F, which is linked to Pick's disease. Recently, S320F was found

to stabilize a hydrophobic pocket of tau, exposing the ^{306}VQIVYK^{311} amyloid motif and thereby

promoting aggregation[37]. The combination of these two mutations likely amplifies their proclivity

towards β-sheet self-assembly. To further test this, we generated PL-SF I277P/I308P containing

two amyloid-breaking proline mutations within the MTBR. Indeed, insertion of β-breaking

mutations suppressed tau aggregation in primary neurons (**Supplemental Figure 4**), in line with

previous in vitro studies[23–27].

Second, by extension of its aggregation propensity, PL-SF tau may also be more highly

decorated with pathological PTMs than either single tau mutant, driving conformational changes

(i.e., MC1). In this study, we only evaluated well-known tau phosphorylation epitopes (AT8,

AT100, AT180), but future mass spectrometry approaches could determine the temporal

emergence of global tau PTMs (**Figure 2A, B**). For instance, increased PTM complexity could

create a feed-forward cycle leading to further loss of tau-MT binding. One can also envision PTM-

induced loss of tau binding to quality control machinery, which typically acts in a compensatory

manner to prevent aberrant tau[2]. One example is the surveillance factor HDAC6, which binds and

maintains tau in the deacetylated state[2]. Thus, PL-SF could overburden the cellular proteostasis

capacity, as suggested for human AD[38–40].

Third, P301L and S320F mutations are also known to dissociate tau from microtubules,

which would shift and favor their equilibrium, further concentrating tau pools in the cytosol and

facilitating their aggregation. PL-SF drastically reduced acetylated-tubulin levels back down to

baseline (i.e., non-tau expressing cells) (**Figure 1 H, I**), suggesting PL-SF is almost completely

disassociated from microtubules and therefore more readily primed for aggregation.

Lastly, once formed, we found that neuronal PL-SF tau can recruit endogenous mouse tau

(detected by T49 antibody) into tau inclusions (**Figure 2H**), consistent with a prior study showing

that human PL-SF-expressing mice (SPAM mice) recruited endogenous mouse tau[6,28]. PL-SF tau,

therefore, may be more highly seeding competent, in a species-independent manner, which could

be leveraged to model tau spreading since it poses several advantages over the use of exogenously

added, brain-derived AD seeds (e.g., large-scale production, reliability and reproducibility, and

fully intact N-terminal domains). We reiterate that further ultrastructural studies are needed to

determine whether seeding-competent PL-SF resembles human AD brain-derived tau.

The role of Hsp70 in tau pathology

We added clarity to the long-standing question of whether Hsp70 modulates tau

aggregation, as the prior literature is inconsistent and often highlights contradictory models for tau

triage via coordinated Hsp binding. Some studies have found that activation or overexpression of

Hsp70 can suppress tau aggregation in vitro, which would intuitively suggest a classical Hsp70

chaperone activity as a suppressor of tau aggregation[16,17,41,42]. Adding additional complexity,

while Hsp70 can disassemble tau filaments isolated from human AD brain, this chaperone activity

can release seeding-competent tau species, suggesting any protection afforded by Hsp70 may

depend on the nature of the chaperone-liberated tau species[43]. In contrast, other studies have shown

that Hsp70 depletion or small molecule inhibition (YM-01, JG-23/48) may drive tau

clearance[11,13,44–48]. We sought to reconcile these differences in the context of our PL-SF cellular

model of tau aggregation.

We initially found that Hsp70 was tightly associated with PL-SF tau aggregates in vitro

and in cell culture, a finding consistent with a previous study[49] (**Figure 5I**). HSP70 overexpression

was associated with PL-SF phosphorylation and aggregation, while HSP70 depletion reduced total

tau levels, including hyperphosphorylated and aggregated tau species (**Figure 3A-D, Figure 4A-**

D). Furthermore, abrogating Hsp70 binding using an Hsp70-binding deficient mutant (PL-SF-4Δ)

was also sufficient to suppress tau aggregation (**Figure 1, 2, 5, 6**). Further supporting a role for

Hsp70 in human AD pathology, we found that Hsp70 is upregulated in AD hippocampus and

heavily intercalated within neuritic plaques and NFTs (**Figure 7**). The Hsp70 localization pattern

was highly consistent in human AD hippocampus (**Supplemental Figure 8**) and illustrates a

unique tau-Hsp70 pattern of co-pathology[50–52]. The lack of complete colocalization suggests that

Hsp70 recruitment to tau occurs in a time-dependent manner. At least in a mature tau aggregation

model, these data are consistent with Hsp70 stabilizing a misfolded tau conformation that

potentially evades proper triage and/or degradation.

In addition to Hsp70, tau is also an Hsp90 substrate, which competes for binding sites

within tau's MTBR and is thought to promote CHIP-mediated tau degradation and clearance[21,53–]

[59]. It is plausible that aggregated tau may form as a consequence of an imbalance in the ratio of

Hsp70 vs. Hsp90 binding, which, if shifted towards Hsp70, could radically shift tau processing

away from degradation towards stabilization of aggregated tau intermediates. In this scenario,

Hsp70 depletion or inhibition would facilitate proteasome or autophagic targeting (degradation)

by shifting the balance away from tau refolding. Future proteomics studies will be needed to

determine whether distinct tau triage complexes emerge temporally in response to tau aggregation.

For example, identifying the PL-SF interactome over time may reveal distinct triage stages in

which Hsp70 has shifted from pro-folding to pro-aggregation[60].

We did not observe any effect of Hsc70 on tau aggregation. Though highly homologous,

Hsc70 is constitutively expressed at ~30-fold higher levels in the brain compared to the inducible

Hsp70 isoform[61]. Hsp70, but not Hsc70, overexpression increased the number of MC1-positive

neurons by nearly 3-fold (**Figure 3E, F**). Since Hsp70 is more sensitive to redox signaling (via

cysteine residue reactivity), its activity towards tau may be more tunable under stress conditions[62].

Prior work has shown that the functional outcome of Hsp70-tau interactions is highly

dependent on the specificity and fine-tuning provided by co-chaperones that regulate Hsc70[19],

Hsp90[63–65], CHIP[61], as well as small heat shock proteins[66–68]. Thus, we suspect that many

variables could dictate how effectively and efficiently Hsp70 targets tau, including: 1) the extent

of MT-dissociated aggregated tau present, 2) the relative balance and abundance of chaperones,

co-chaperones, and cofactors that target and triage tau, and 3) the notion that disease onset and

disease progression (either acute or chronic) may involve distinct signaling pathways or

compensatory responses.

Taken together, our study supports the notion that depletion or activity-modifying

approaches targeting Hsp70 (or its cofactors and co-chaperones) may be neuroprotective in the

context of the AD brain. While brain-specific HSP70 knockout mice are not available (to the best

of our knowledge), brain-permeable Hsp70 inhibitors may have therapeutic utility, although broad

Hsp70 inhibition could have detrimental effects on overall brain proteostasis. Therapeutic

strategies aimed at "precision proteostasis" could selectively modulate chaperone–client

interactions to favor the clearance of pathogenic tau without compromising global protein

homeostasis and without directly targeting tau itself.

Materials and Methods

Plasmid and lentivirus cloning

To generate the pUltra-T40 WT, PL, PL-SF, SF, PL/SF-2Δ (277-278), PL/SF-2Δ (308-309),

PL/SF-4Δ, HSC70-flag, HSP70-flag, and HSP90-flag constructs, the cDNAs were amplified

through PCR and inserted into the pUltra vector using AgeI and SalI restriction endonucleases to

replace eGFP in pUltra. pUltra-XeGFP was made by removing eGFP using AgeI and BspEI,

creating a blunt end with Klenow before circularizing. pUltra-T40 WT (eGFP), PL/SF (eGFP),

and PL/SF-4Δ (eGFP) were created by replacing TDP43 in pUltra TDP43 WT (eGFP) using BspEI

and SalI. To generate PL/SF/2IP, two consecutive Q5 site-directed mutagenesis were performed

on PL/SF(eGFP). pUltra was ordered from Addgene (Plasmid #24129). pUltra was a gift from

Malcolm Moore (Addgene plasmid # 24129; RRID:Addgene_24129). shRNA plasmids against

HSP70   (TRCN0000098597,   TRCN0000098598)   and   HSP90   (TRCN0000008491,

TRCN0000008494) were ordered from the UNC Lenti-shRNA Core Facility. Primers used for

vector construction are in Table 1.

QBI-293 cell culture

QBI-293 (HEK293A) cells (ThermoFisher, #R70507) are commercially available and maintained

according to standard protocols. QBI-293 cells were cultured in complete DMEM media (10%

fetal bovine serum, 1X L-glutamine, and 1X penicillin/streptomycin). QBI-293 cells were

transfected with FuGENE 6 transfection reagent (Promega) to express 2N4R tau constructs for 48

hr. Following transfection, cells were lysed and collected in 1X RIPA buffer [50 mM Tris pH 8.0,

150 mM NaCl, 5 mM EDTA, 0.5% sodium deoxycholate, 1% Igepal CA-630, 0.1% SDS]

supplemented with deacetylase, phosphatase and protease inhibitors: 1 mM phenylmethylsulfonyl

fluoride (PMSF), 10 mM Nicotinamide (NCA), 1 mM sodium orthovanadate (Na3VO4), 1 mM

sodium fluoride (NaF), 0.5 M β-glycerophosphate, 2 μM Trichostatin A (TSA), and protease

inhibitor cocktail composed of L-1-tosylamide-2-phenylethylchloromethyl, N-tosyl-L-lysine

chloromethyl ketone, leupeptin, pepstatin, and soybean trypsin inhibitor, each at 1 μg/ml.

Primary cortical neuron cell culture

Mouse primary cortical culture was performed in accordance with the University of North Carolina

(UNC) Institutional Animal Care and Use Committee (UNC IACUC protocol 24-190). Mouse

cortical neurons were dissected and isolated from C57Bl/6 mice (Charles River) at embryonic day

15-16 following lethal isoflurane anesthesia. Following anesthesia, embryos were removed from

the pregnant abdominal cavity, washed well with ice-cold 70% ethanol and placed in cold Hanks-

buffered balanced salt solution (HBSS). Once rinsed, these embryos were moved to a 10cm dish

for dissection. The embryos were then removed from the placental membrane and brains were

isolated. The cortex was dissected then moved to Hibernate-E solution (BrainBits) with B27

(Gibco) and Glutamax (ThermoFisher). The cerebral cortices were minced thoroughly with

forceps and incubated for 30 minutes at 37 °C in HBSS with 20 U/mL papain (Worthington

Biomedical), 1mM EDTA, 0.2 mg/mL L-Cysteine, and 5U/mL DNAse (Promega). Cortical tissue

was further dissociated in plating media (BrainPhys media (Stemcell), 5% fetal bovine serum, 1X

B27 (Gibco), 1X penicillin/streptomycin (Gibco), and 1X Glutamax by triturating the sample using

a P1000 micropipette. This suspension was centrifuged for 5 minutes at 1.5 rcf to pellet the cells.

The pellet was then resuspended in plating media and passed through a 40 μm cell strainer. A

Countess 3 automated cell counter (Invitrogen) was used to count the cells. Cells were plated in

plating media at 600,000 cells/well in 6-well plates or 30,000 cells/well in 96-well plates. The next

day (DIV1), plating media was completely aspirated and replaced with fresh neuronal media

(BrainPhys media (Stemcell), B27 (Gibco), 1X penicillin/streptomycin (Gibco), and 1X

Glutamax).

364   Lentiviral production

Lentiviral production was performed by co-transfecting 37.5 mg lentivirus plasmid with 25 mg

psPAX2, 12.5 mg VSVG, and 6.25 mg REV plasmids per 15 cm culture plates of Lenti-X 293 T

cells (Takara). For each lentiviral construct, three 15 cm plates were used for production. Roughly

48 hrs after the transfection, cell culture media was collected and subsequently centrifuged at

2000 rcf for 10 minutes. Lentiviral particles were purified using a double-sucrose gradient method

as follows: supernatants were loaded onto a 70–60% to 30–20% sucrose gradient and centrifuged

at 70,000 rcf for 2 hr at 17 °C. The 30–60% fraction containing the viral particles was retrieved,

resuspended in PBS, filtered with a 0.45 mm filter flask, loaded onto a 20% sucrose solution, and

centrifuged a second time at 70,000 rcf for 2 hr at 17 °C. The supernatants were carefully

discarded, and the viral particles in the pellet were resuspended in PBS, aliquoted and stored at

− 80 °C.

Lentiviral transductions

Primary neurons were transduced with ~3.0×10^{12} viral particles per mL at DIV4 for most experiments (viruses were added at DIV18 for MEA experiments). If neurons were co-transduced with two viruses, both were added sequentially (each at ~3.0×10^{12} viral particles per mL). For transductions, viruses were added directly to the media in each well (1 µl virus per 2 mL media), followed by gentle rocking by hand to distribute viral particles. The following day, virus-containing media was completely aspirated and replaced with half conditioned media (neuronal media from the same neuron culture) and half fresh neuronal media (BrainPhys (StemCell) with B27 (Gibco), Pen-strep (Gibco), and Glutamax (Gibco)). Every two-three days, half of the neuronal media was discarded and replenished with fresh neuronal media until neurons were collected. Most neurons were collected at DIV10 (7 days after transduction), unless otherwise noted.

Immunoblotting and biochemical analysis

RIPA soluble lysates (250 µl per well of a 6-well plate) collected from QBI-293 and primary cortical neuron were sonicated 20 times on ice, then centrifuged at 15,000 rcf for 20 minutes at 4 °C. The supernatant was removed from the pelleted material, leaving behind a RIPA insoluble pellet. The RIPA insoluble pellet was washed with 250 µl of RIPA lysis buffer with protease, phosphatase, and deacetylase inhibitors and centrifuged at 15,000 rcf for 20 minutes at 4 °C. This supernatant was removed and discarded and 75 µl of 1% SDS was added to each pellet then sonicated 20 times at room temperature. These samples were then centrifuged at 15,000 rcf at room temperature. The supernatant from these tubes was removed and stored in separate tubes as the

SDS-soluble fraction. The RIPA and SDS-soluble supernatants were then mixed with 6X loading

dye with DTT and boiled at 98 °C for 10 minutes in a PCR thermocycler (BioRad). Samples were

run and resolved via SDS-PAGE on 4–20% gradient gels (BioRad). Gels were transferred to iBlot

2 Midi nitrocellulose membranes using the iBlot 2 (Invitrogen) following the standard P0, 7-

minute settings. Following transfer, membranes were blocked with 2% milk in 1X TBS for 30

minutes and then incubated with primary antibody overnight at 4 °C on a rocker. The next day,

membranes were incubated with HRP-conjugated anti-mouse or anti-rabbit antibodies (1:1000)

for 1-2 hr at room temperature. Blots were developed with ECL Western Blotting substrate

(Cytiva, Amsterdam) and imaged on a GE ImageQuant LAS4000.

Immunocytochemistry

Immunocytochemistry experiments were performed using QBI-293 cells or cortical primary

neurons plated on poly-D-lysine (PDL)-coated 96-well plates. Following either plasmid

transfection (48 hr) or lentiviral transduction (7 days unless otherwise stated), cells were washed

once in PBS, immediately fixed in 4% paraformaldehyde for 15 minutes, then washed once more

with PBS. Following fixation, cells were then permeabilized with permeabilization buffer (0.2%

Triton X-100 (Sigma) in PBS) for 20 minutes then blocking buffer (4% normal goat serum in PBS

with 0.05% Tween 20) for 1 hr. Primary antibodies were added in blocking buffer and incubated

overnight on a rocker at 4 °C. The following day, cells were washed 3X in PBS with 0.05% tween

20 and incubated with Alexa 488, Alexa 594, or Alexa 647 conjugated secondary antibodies at

1:1000 and DAPI for 1 hr. Cells were then washed 3X in PBS with 0.05% Tween 20. Cells were

then imaged using an EVOS M7000 automated microscope with EVOS 20X objective (Invitrogen

AMEP4982) and 40X objectives (Invitrogen AMEP4754) and DAPI, GFP, Texas Red, and Cy5

light cubes. Note: Some representative images were artificially brightened or contrasted in

Microsoft PowerPoint or Celleste 5.0 and 6.0 software (Invitrogen) to aid visualization. Any panels

assessing the same set of conditions were brightened to identical levels. Images were not

manipulated prior to quantification.

Antibodies

The following primary antibodies were used in this study: AT8 (ThermoFisher Scientific-

MN1020), AT100 (ThermoFisher Scientific MN1060), AT180 (ThermoFisher Scientific

MN1040), TauC3 (ThermoFisher Scientific-AHB0061), T49 mouse tau (Millipore- MABN827),

Tau12 (Millipore, MAB2241, 1:1000), Tau5 (AHB0042), Hsp70 used for immunoblotting (Enzo-

ADI-SPA-810-D), Hsp70 used for immunoblotting and immunocytochemistry (Cell Signaling

Technologies- 4872), Hsc70 1B5 (ADI-SPA-815-F), Hsp90 16F1 (Enzo- adi-spa-835), Total tau

rabbit polyclonal- formerly K9JA (DAKO, A0024 1:2000), GAPDH 6C5 (Millipore- MAB374),

ac-tubulin (Sigma T7451), alpha tubulin (Sigma, T9026), Anti-FLAG M2 (Sigma, F1804, 1:2000),

GFAP (Dako, 20334). The MC1 antibody was kindly shared by Dr. Peter Davies. Quantification

of immunoblotting was measured by ImageStudio Lite Version 5.2.

Co-immunoprecipitation reactions

All co-immunoprecipitations were performed in QBI-293 cells transfected with plasmids

expressing 2N4R wild-type, P301L, P301L/S320F, and P301L/S320F 4Δ for 48 hrs prior to

collection. Each well of a 6-well plate was used as separate conditions. Cells were collected in 250

µl RIPA buffer with protease, phosphatase, and deacetylase inhibitors prior to co-

immunoprecipitation. Lysates were immediately rotated on a wheel for 30 minutes at 20 rpm at 4

°C. Lysates were then centrifuged at 15,000 rcf for 15 minutes at 4 °C. Supernatants were removed

and 40 µl were kept as input. The remaining supernatant was pre-cleared with 50 µl of Protein

A/G PLUS-Agarose beads (Santa Cruz, sc2003) for 30 minutes at 4 °C. Lysates were centrifuged

for 1 minute at 1,500 rcf, and the supernatant was removed. The volume of each reaction was

brought to 500 µl of RIPA buffer with inhibitors. For each reaction, 8 µl of tau5 antibody was

added to each reaction, then 75 µl of agarose protein A/G beads. Lysates were rotated on a wheel

at 20 rpm at 4 °C overnight. The following day, lysates were centrifuged at 1,500 rcf for 5 minutes.

Most of the supernatant was carefully removed, followed by washing with 500 µl of RIPA buffer

with inhibitors, and lysates were centrifuged at 1,500 rcf for 5 minutes. This wash was repeated

four times. After the final wash, loading dye with DTT was directly added to the beads and heated

for 10 minutes at 98 °C with intermittent mixing. These were centrifuged for 30 seconds at 1,500

rcf prior to storage at -80 °C.

Purification of recombinant tau

Tau purification methods were adapted from Barghorn et al, 2004[69]. Untagged wild-type and

mutant versions of tau were cloned into pRK172 backbone and transformed into BL21(DE3)-RIL

cells. Cells were grown in LB media at 37 °C until the OD600 reached 0.8. Protein expression was

then induced with IPTG and continued shaking at 37 °C for 3-5 hrs. After the cells were harvested

and resuspended in lysis buffer (20 mM MES pH 6.8, 1 mM EDTA, 5 mM DTT, protease

inhibitors benzamidine and leupeptin, and PMSF), the cells were lysed via sonication. Lysed cells

were heat shocked at 80 °C for 20 minutes, then centrifuged at 17,500 rpm (36,635 xg) for 30

minutes to remove cell debris. Clarified lysate was dialyzed overnight at 4 °C into wash buffer (20

mM MES pH 6.8, 50 mM NaCl, 1 mM EDTA, 1 mM $MgCl_2$, 2 mM DTT). Following overnight

dialysis, lysates were centrifuged at 17,500 rpm (36,635 xg) for 30 minutes to remove precipitated

proteins and then subjected to cation exchange chromatography. The protein was eluted by

increasing salt concentration in buffer to 250 mM. The samples were further purified via size

exclusion chromatography on a Superdex 200 Increase 10/300 GL column in sizing buffer (25 mM

HEPES pH 7.5, 100 mM NaCl, 5 mM $MgCl_2$, 1 mM DTT). Samples were then concentrated as

needed using centrifugal filters and flash frozen for later use.

Heparin-induced tau aggregation assays:

Tau aggregation assays were carried out in a PolarStar Omega microplate reader using Thioflavin

T (ThT) fluorescence to monitor formation of tau aggregates. Aggregation reactions were started

by combining 5 µM tau, 2 µM heparin, and 20 µM ThT in reaction buffer (1x PBS, 2 mM TCEP)

and transferred to a costar 96-well half-area microplate (black polystyrene, flat bottom, non-

treated). The microplate was covered with an ELISA plate sealer and placed in the microplate

reader at 37 °C. Fluorescence measurements were taken every 5 minutes for 12 hrs using a 440-

10 nm excitation and 485-12 nm emission filter.

HSP70 and HSP90 shRNA knockdown

All shRNAs were obtained from UNC Lenti-shRNA core facility and cloned into the pUltra

vectors. Sequences: HSP70 shRNA: GCAGGTGAACTACAAGGGCGA; eGFP control shRNA:

TACAACAGCCACAACGTCTAT (Y145 to Y151 residues of eGFP). HSP70 shRNAs were

added to the neuron culture four days after transduction with tau lentiviruses at a titer of ~$3.0x10^{12}$

viral particles per mL. Media was half exchanged for fresh neuronal media every two days. After

an additional four days in culture, primary neurons lysates were collected for experiments. We

noted some toxicity after more than 4 days in culture with HSP70 shRNAs. HSP90 shRNAs were

applied for 7 days in culture (~3.0×10^{12} viral particles per mL) prior to collection for experiments

and were not overtly toxic.

Multi-electrode array recordings and analysis

Following dissection, primary mouse cortical neurons were directly plated onto poly-D lysine

coated 24-well BioCircuit MEA plates (Axion Biosystems). A 10 µl droplet of neurons at a

concentration of 16,000,0000 neurons/mL was added directly to the center of each well. Neurons

were incubated at 37 °C for 1 hr then 250 µl of plating media (BrainPhys media (Stemcell), 5%

fetal bovine serum, 1X B27 (Gibco), 1X penicillin/streptomycin (Gibco), and 1X Glutamax) was

carefully added to each well in a circular motion. Following another 1-hr incubation, another 250

µl of plating media was added to each well (total volume = 500 µl). The following day, ~90% of

plating media was aspirated and fresh neuronal (BrainPhys media (Stemcell), B27 (Gibco), 1X

penicillin/streptomycin (Gibco), and 1X Glutamax) media was added to each well. Half of the

media was exchanged for fresh neuronal media every two days. At DIV18, neurons underwent an

initial, 10-minute pre-recording using the Axion Maestro Edge (Axion Biosystems) and AxIS

Navigator software (Axion Biosystems). Immediately following this recording, neurons were

transduced with lentiviruses at ~3.0×10^{12} viral particles per mL (four replicate wells per condition).

The following day, neuronal activity was recorded for 10 minutes, then lentivirus-containing

neuronal media was completely replaced with half fresh neuronal media and half conditioned

neuronal media from the same culture. Neurons were recorded nearly every day for 10 minutes

and half of the neuronal media was replaced every 2-3 days after recording until DIV71 (day 53

of recording). Post-hoc analysis was performed using the NeuralMetricTool (Axion Biosystems)

using default analysis options. Data was exported into Microsoft Excel, then graphed via GraphPad

Prism 10 for statistical analyses and graphing. Raster plots were generated from the same data

using the AxIS Metric Plotting Tool. 96 well plates were plated at the same time using the same

primary neurons, underwent identical media changes, and were used for immunocytochemistry at

DIV29 and DIV62 in Figure 5.

Post-synaptic density isolation

Primary mouse cortical neurons from wild-type C57/BL6 mice were plated on poly-D-lysine

coated 6-well plates at 600,000 cells/well. Neurons were treated with lentivirus at DIV4 (1 6-well

plate per condition) and half-fed every two days until DIV14. At DIV14, neurons were rinsed in

cold PBS and all 6-wells of each plate were collected in a total of 900 µl of homogenization buffer

(320 mM sucrose, 5 mM sodium pyrophosphate, 1 mM EDTA, 10 mM HEPES pH 7.4) with

protease/phosphatase inhibitor cocktail 200 nM okadaic acid, 1 mM orthovanadate, 1X protease

inhibitor cocktail (Roche). Neuronal lysates were homogenized by passing the lysate 12 times

using a 26g needle. For each condition, 70 µl of lysate was kept and stored as whole cell lysate.

The remaining lysate was centrifuged at 1,000 g for 10 minutes at 4 °C *to* generate pelleted nuclear

fraction (P1) and supernatant fraction (S1). The S1 fraction was centrifuged at 15,000 g for 20

minutes at 4 °C to generate the membrane/crude synaptosome pelleted fraction (P2) and cytosolic

supernatant fraction (S2). The P2 fraction was resuspended in 500 µl of ultrapure H_2O with

protease/phosphatase cocktail. 20 µl of 100 mM HEPES was added to the 500 µl of suspension to

yield 4 mM HEPES pH 7.4. These samples were then spun on a wheel for 30 minutes at 4°C then

centrifuged for 20 minutes at 4 °C at 25,000 g using a Sorvall mTX 150 ultracentrifuge (Thermo

Scientific) to generate pelleted synaptic fraction (LP1) and supernatant fraction (LS2). The LP1

pellet was resuspended in 250 µl mM HEPES pH7.4 using repeated trituration with a micropipette.

This was then mixed with 250 µl of 1% Triton X-100 with protease/phosphatase cocktail and spun

on a wheel for 10 minutes at 4 °C. This was then centrifuged in Sorvall mTX 150 ultracentrifuge

(Thermo Scientific) at 32,000 g for 20 minutes at 4 °C. This pellet was resuspended in 80 µL of

50 mM HEPES pH7.4 using micropipette trituration to yield the PSD fraction. Only total

homogenate and PSD fraction were processed and analyzed by SDS-PAGE.

Imaging quantification

All immunocytochemistry (aside from organotypic slice cultures) quantification was performed

using Celleste 5.0 and 6.0 software (Invitrogen). The number of inclusions were counted using the

count/size function. Puncta and inclusions were manually segmented based on intensity (Segment:

Manual) and size (Region: Area) thresholds for each tau probe (e.g. AT8, MC1, ThioS). These

threshold settings were then applied to all images used for that specific experiment, automatically

batched processed using a segment-count-export macro generated in Celleste, then exported to

Microsoft Excel for subsequent analysis.

Hippocampal slice cultures

Sprague-Dawley rats (Charles River Laboratories, Wilmington, MA) were housed and maintained

in accordance with the recommendations from the Guide for the Care and Use of Laboratory

Animals from the National Institutes of Health, and in accordance with an approved protocol from

the Institutional Animal Care and Use Committee of the University of North Carolina-Pembroke.

Brain tissue from postnatal 12-day-old rats was rapidly prepared and hippocampi positioned on a

McIlwain tissue chopper for assembling slices for interface cultures as previously described[70,71].

Three-dimensional organotypic cultures of specific brain regions are well known for studying

different types of physiologies and pathogenic processes[72–74]. Transverse hippocampal slices (400-

µm thickness) were immediately placed in ice-cooled-buffer solution (pH 7.2) then transferred in

groups of eight to nine onto the Biopore membrane of culture inserts (Millipore, Billerica, MA).

The media below the culture inserts contained 50% basal medium eagle (Sigma-Aldrich, St. Louis,

MO), 25% Earle's balanced salts (Sigma-Aldrich), 25% horse serum (Gemini Bio-Products,

Sacramento, CA), and defined supplements as previously described[75] . The media were initially

changed 24 hrs after the slicing procedure, followed by subsequent changes every 3-4 days. The

cultivated hippocampal slices were maintained under conditions of a 5% CO_2-enriched

atmosphere at a constant temperature of 37 °C for a duration of nine weeks.

Lentiviral infection of organotypic cultures

Lentiviruses expressing GFP or GFP-tagged tau variants (wild-type, PL-SF, and PL-SF 4Δ tau).

For each condition, 1.5 µl of lentivirus were applied onto the surface of individual organotypic

slices (~7.5×10^{12} viral particles per slice) approximately 3 hrs after the placement of the slices on

Millipore inserts. Following a nine-week incubation period in vitro, both the infected organotypic

slices and their corresponding controls were fixed with 4% PFA for 3-4 hrs. Subsequently, the

slices were rinsed multiple times with PBS and stored in PBS at 4 ºC.

Immunofluorescence staining and imaging of organotypic brain slice cultures

The fixed organotypic hippocampal explants underwent permeabilization in PBS containing 0.5%

Triton-X100 for 30 minutes. Following this, the slices were blocked with a solution of 5% BSA in

PBS containing 0.1% Triton-X100 for 90 minutes at room temperature. Subsequently, the

organotypic slices were incubated for 48 hrs at 4 ºC with primary antibodies targeting AT8 (1/400;

ThermoFisher Scientific- MN1020) or MC1 (1/400; Dr. Peter Davies, diluted in a solution of 3%

BSA containing 0.1% Triton-X100. Next, brain explants were washed 3-4 times with PBS

containing 0.1% Triton-X100 and then exposed to Goat anti-mouse IgG Alexa-FluorTM 568

(ThermoFisher, Waltham, MA) for 120 minutes at room temperature. Following this incubation,

the organotypic slices were rinsed three times with PBS containing 0.1% Triton-X100. During the

second wash, NucBlue stain (Invitrogen, MA, USA) was introduced to the PBS to promote DAPI

nuclear staining. Subsequently, the slices were rinsed in PBS and were mounted onto glass slides,

covered by 1.5 mm coverslips, using ProlongTM diamond antifade mountant (Thermo Fisher

Scientific, NC, USA). Immunofluorescence images were captured with Andor Dragonfly spinning

disk confocal using the Zyla Plus 4.2MP sCMOS (Oxford Instruments, Abingdon, United

Kingdom). Image our blinded processed adopting the equal adjust using the Imaris

Multidimensional visualization and analysis software (Schlieren-Zurich, Switzerland).

Human brain fractionation

Human control and Alzheimer's disease (AD) cortical brain tissues (Braak stages V–VI) were

obtained from the Center for Neurodegenerative Disease Research brain bank at the University of

Pennsylvania. Gray matter from the frontal cortex of each subject was isolated and homogenized

in 3 volumes per gram (vol/g) of cold high-salt RAB buffer (0.75 M NaCl, 100 mM Tris, 1 mM

EGTA, 0.5 mM $MgSO_4$, 0.02 M NaF, 2 mM DTT, pH 7.4), supplemented with deacetylase,

phosphatase, and protease inhibitors as previously described. Homogenates were incubated at 4 °C

for 20 minutes to facilitate microtubule depolymerization, followed by ultracentrifugation at

100,000 × g for 30 minutes at 4 °C. The resulting supernatant was designated the high-salt fraction.

The pellet was then re-homogenized and subjected to a second round of centrifugation under

identical buffer and conditions. After removal of the supernatant, the remaining pellets were

homogenized in 5 vol/g of cold RIPA buffer (50 mM Tris, pH 8.0, 150 mM NaCl, 1% NP-40, 5

mM EDTA, 0.5% sodium deoxycholate, 0.1% SDS), and centrifuged once more at 100,000 × g

for 30 minutes at 4 °C. Subsequent to myelin flotation, the pellets were re-extracted in RIPA buffer

containing 20% sucrose to eliminate residual myelin and labeled as the RIPA fraction. Finally, the

insoluble material was extracted using 1 vol/g of urea extraction buffer (7 M urea, 2 M thiourea,

4% CHAPS, 30 mM Tris, pH 8.5). Both the high-salt and urea-soluble fractions were subjected to

SDS-PAGE followed by immunoblotting with antibodies targeting Hsp70, Hsc70, and GAPDH.

Human brain immunohistochemistry and confocal microscopy

Human hippocampal tissue sections mounted on coverslips were obtained from the UNC

Translational Pathology Laboratory (TPL), which oversees a cataloged and archived repository of

Alzheimer's disease (AD) brain specimens, including neuropathologically confirmed cases from

UNC-affiliated donors. The 10 um sections were de-paraffinized with two 45-minute washes in

xylene followed by one-minute washes in 100% ethanol (x2), 95% ethanol, 80% ethanol, 75%

ethanol, water, and finally allowed to soak O/N in 1x PBS. Antigen retrieval is performed using a

citrate-based antigen unmasking solution (Vector Labs, H-3300-250) in boiling Mill-Q water for

10-minutes. The slides were then washed 3x5-min with TBS, followed by 1x wash in TBS + 0.1%

TritonX-100 for 20 minutes, 3x5-min with TBS, and 1x5 min in 3% Goat Serum in TBS (Southern

Biotech, 0060-01). Sections were then stained 2x O/N in a humidified chamber at room

temperature using anti-AT8 (1:500, Invitrogen, MN1020) and anti-Hsp70 (1:500, Abcam,

ab181606). Slides were then washed 3x5-min with TBS followed by secondary staining with goat

anti-mouse IgG 488 (1:1000, Invitrogen, A-11001) and goat anti-rabbit IgG 594 (1:1000,

Invitrogen, A-11012) for 24 hours at 4°C in a dark, humidified chamber. Slides were then washed

3x5-min with TBS, allowed to dry, and mounted with Fluoromount-G with DAPI (Invitrogen, 00-

4959-52).

Confocal microscopy was performed on an LSM980 (Zeiss) inverted laser scanning confocal

microscope. The LSM980 is equipped with a motorized X, Y, stage and Z-focus with high-speed

Piezo insert along with four diode lasers (405, 488, 561, and 633). Images were acquired with 1x

Nyquist sampling using 63x 1.4 NA Plan-Apochromat oil objective (Zeiss) and 4 channel GaAsP

detectors on Zen Blue 3.6 software. Images were acquired with 1x Nyquist sampling using 20x

0.8 NA Plan-Apochromat air objective (Zeiss) and 4 channel GaAsP detectors on Zen Blue 3.6

software.

Statistical analysis

GraphPad Prism 10 software was used for all statistical analyses. For comparisons between two

groups, two-tailed unpaired Student's t-tests were used. For multiple group comparisons, one-way

ANOVA followed by Dunnet's or Sidak's post hoc tests were applied as appropriate. Results were

polled from a minimum of n=3 independent experiments and presented as mean ± standard error

of the mean (SEM). P-values < 0.05 were considered statistically significant. The number of

replicates (n) and details of the specific tests applied are provided in figure legends.

Funding Declaration

Support for this work was provided by National Institutes of Health (NIH) grants F32AG072826 (MRB) and R01AG068063 (TJC). This study was supported by the National Institute on Aging of the National Institutes of Health under Award Number P30AG072958 (NIAP30AG072958) (MRB, MA, and JHT), Alzheimer's Association grants AARG-22-924321 (TJC) AARF22926617 (JHT), and CurePSP grant 656-2018-06 (TJC). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Acknowledgements

The Andor Dragonfly microscope was funded with support from National Institutes of Health grant S10OD030223. The Zeiss LSM 980 microscope was funded with support from NIH grant S10 OD032388. We would also like to thank Kinsley Adams for her expertise and assistance with the hippocampal organotypic slice cultures. We would also like to thank the University of Pennsylvania (Center for Neurodegenerative Disease Research brain bank) for providing human control and AD cortex lysates. Lastly, we would like to thank the University of North Carolina Translational Pathology Laboratory for providing AD hippocampal brain sections.

Conflict of interest

The authors declare that they have no conflicts of interest with the contents of this article.

Author Contributions

MRB designed and performed most experiments. MA, JS, and BB designed and carried out ex vivo organotypic slices experiments and imaging. KP performed all immunohistochemistry and

confocal imaging of human post-mortem brain. CO assisted with co-immunoprecipitation

experiments and post-synaptic density preparations. CO and JHT performed human brain

fractionation and immunoblots. EP performed all protein purification and carried out the

Thioflavin T aggregation assays. JMM assisted with multi-electrode array recordings and

analyses. XT prepared all plasmid and lentiviral constructs for all experiments. DK and HT

designed and performed the QBI293 experiments comparing various P301 and S320 tau mutants.

JHT, JS, NB, and BB provided additional experimental guidance and interpretation. MRB and TJC

prepared the manuscript and figures. TJC directed and supervised the study. All authors consented

and contributed to the final version of the manuscript.

Figures and Figure Legends

Bryan et al., Figure 1.

Figure 1: Abrogating tau binding to heat shock proteins alleviates pathology in a PL-SF tau aggregation model.

A) Representative immunoblots (RIPA fraction) of QBI-293 cells transfected with 2N4R WT, P301L, S320F, or P301L-S320F tau. **B)** Immunoblot quantification of total tau, p-tau (AT8 and AT100), and cleaved tau (TauC3) in RIPA fraction. N=4 per condition; **C)** Immunocytochemistry quantification of QBI-293 cells transduced with WT, P301L, S320F, or PL-SF tau labeled with AT8, TauC3, or ThioS. N=48 FOVs per condition. *= p-value significant by one-way ANOVA. Representative images found in Supplemental Figure 1. **D)** Representative immunoblots (RIPA fraction) of QBI-293 cells transfected with 2N4R P301L-S320F tau or P301L-S320F 4Δ (deletion of I277/278, I308/V309 residues). *AT100 was run on a separate blot with the exact same lysates to improve detection. **E)** Immunoblot quantification of total tau, p-tau (AT8 and AT100), and cleaved tau (TauC3) in RIPA soluble fractions. N=3 per condition. **F)** Immunocytochemistry quantification of QBI-293 cells transduced with 2N4R PL-SF or PL-SF 4Δ tau labeled with AT8, TauC3, or ThioS. N=16 FOVs per condition. *= p-value significant by unpaired Student's t-test. Representative images found in Supplemental Figure 3. **G)** Co-immunoprecipitation of QBI-293 cells transfected with WT, P301L, PL-SF, or PL-SF 4Δ tau. Tau5 antibody was used to IP tau and immunoblots were probed with total tau (rabbit polyclonal) and Hsp70, Hsc70, and Hsp90 antibodies. **H)** Representative immunoblots of QBI-293 cells transfected with 2N4R tau variants and probed with acetyl-tubulin and total alpha-tubulin antibodies. Note: Acetyl tubulin and alpha tubulin blots were run separately with the same lysates to improve detection. **I)** Quantification of acetyl-tubulin across all 2N4R tau variants. -- line signifies separate sets of transfections in panel I and J. N=4 per condition; *= p-value significant by one-way ANOVA. Control= empty plasmid vector (pcDNA3.1). **J)** Quantification of in vitro Thioflavin T aggregation assay using purified

708   protein from PLSF and PLSF 4Δ. N=4 replicates per condition. *= significant by unpaired student

709   t-test at 12hr timepoint. Coomassie (right) shows equal protein loading into reactions.

Bryan et al., Figure 2.

Figure 2: Development and characterization of a PL-SF tau aggregation model in primary cortical neurons.

A) Representative RIPA-soluble immunoblots of DIV10 primary cortical neurons transduced with

lentiviruses expressing 2N4R WT, P301L, S320F, P301L-S320F, or PL-SF 2Δs (R2 and R3) and

4Δ tau variants. **B)** Quantification of AT8, AT100, AT180, and total tau immunoblots in primary

cortical neurons (RIPA fraction). N=3 per condition; *= p-value significant by one-way ANOVA.

C) Representative SDS-soluble immunoblots of DIV10 primary cortical neurons transduced with

lentiviruses expressing 2N4R WT, P301L, S320F, P301L-S320F, or PL-SF 2Δ and 4Δ tau variants.

N=2 for control, WT, and P301L; N=3 for other variants. **D)** Representative immunocytochemistry

images of 2N4R tau variants labeled with AT8 (p-tau), MC1 (conformational tau), ThioS

(aggregated tau) in GFP. Scale bar= 125 μm. **E)** Quantification of the number of AT8, MC1, and

ThioS inclusions in PL-SF and PL-SF 2Δ (R2 and R3) and 4Δ variants. N=12-24 FOVs per

condition. *= significant by one-way ANOVA. **F, G)** Representative immunocytochemistry

images of PL-SF tau displaying colocalization between F) AT8 (p-tau) or G) MC1 (conformational

tau) with ThioS (aggregated tau). Scale bar= 50um. **H)** Representative immunocytochemistry of

PL-SF GFP colocalized with T49 tau (mouse-specific). Scale bar= 125 μm. Control= empty viral

vector (pUltra)**.**

Figure 3: Hsp70 exacerbates tau hyperphosphorylation and aggregation.

A) Representative immunoblots of 2N4R PL-SF tau co-transduced with HSP70, HSC70, and HSP90 in DIV10 primary cortical neurons (RIPA fraction). **B)** Quantification of AT8 (p-tau), total tau (rabbit polyclonal), or Hsp70, Hsc70, Hsp90 (FLAG-tagged) in the RIPA fraction. N=6 per condition. *= p-value significant by one-way ANOVA. **C)** Representative immunoblots of 2N4R PL-SF tau co-transduced with HSP70, HSC70, and HSP90 in DIV10 primary cortical neurons (SDS fraction). **D)** Quantification of AT8 (p-tau), total tau (rabbit polyclonal) in the SDS fraction. N=6 per condition. ns= p-value is not significant by one-way ANOVA. **E)** Representative immunocytochemistry images of primary cortical neurons co-transduced 2N4R PL-SF tau and HSP70, HSC70, or HSP90 labeled with AT8, MC1, Tau12 (RFP) or ThioS (GFP). Scale bar= 125 μm. **F)** Quantification of MC1, AT8, and ThioS positive inclusions in primary cortical neurons co-transduced 2N4R PL-SF tau and either HSP70, HSC70, or HSP90. n=12 FOVs per condition. *= p-value significant by one-way ANOVA. Control= empty viral vector (pUltra)**.**

Figure 4: Reducing Hsp70 can suppress tau hyper-phosphorylation and aggregation.

A) Representative immunoblots of 2N4R PL-SF tau co-transduced with a shRNA against HSP70 in DIV10 primary cortical neurons (RIPA fraction). **B)** Quantification of AT8 (p-tau), total tau (rabbit polyclonal) and endogenous Hsp70 and Hsc70 in the RIPA fraction. N=3. **C)** Representative immunoblots of 2N4R PL-SF tau co-transduced a shRNA against HSP70 in DIV10 primary cortical neurons (SDS fraction). **D)** Quantification of AT8 (p-tau), total tau (rabbit polyclonal) in the SDS fraction. N=3. *= p-value significant by unpaired Student's t-test. Control= empty viral vector (pUltra**).**

Bryan et al., Figure 5.

A)

Mean Firing Rate
(Activity)

B)

Number of Network Bursts
(Connectivity)

C)

Area Under Normalized Cross-Correlation
(Synchrony)

DIV56

D)

Control   PLSF   PLSF 4Δ

E)

F)

G)

	Homogenate			PSD		
	PL	PLSF	PLSF 4Δ	PL	PLSF	PLSF 4Δ

PSD95 — 100kD
AT8 — 75kD
Total Tau — 75kD
Total Protein

H)

DIV29

Control   PLSF   PLSF 4Δ

DIV62

Control   PLSF   PLSF 4Δ

ThioS

MC1

AT8

Tau12
GFAP

I)

DIV29

MC1

PLSF   PLSF 4Δ

AT8

PLSF   PLSF 4Δ

DAPI

HSP70

MC1

AT8

Merge

Figure 5: Mature tau aggregates cause functional deficits in neurons that are reversed by abrogating tau-Hsp binding.

A-C) Quantification of neuronal activity comparing PL-SF to PL-SF 4Δ across 53 days of multi-electrode array recording starting at DIV18 following initial pre-recording. A) Mean firing rate-activity B) Number of network bursts- connectivity C) Area under normalized cross-correlation-synchrony. N=4 replicate wells per condition. *= p-value significant by one-way ANOVA. **D-F)** Representative raster and network burst plots at DIV54 (day 36 of recording) comparing empty control (pUltra) to PL-SF and PL-SF 4Δ. D) Number of network bursts across electrodes over time (pink lines) E) number of spikes across electrodes over time F) Average number of spikes per network burst over time. **G)** Representative immunoblot of isolated post synaptic density (PSD) in DIV18 primary cortical neurons transduced with lentiviruses expressing P301L, PL-SF, or PL-SF 4Δ. Immunoblot was probed with PSD95, AT8 (p-tau), and rabbit polyclonal tau (total tau). **H)** Representative 20X immunocytochemistry images of primary cortical neurons comparing empty vector control (pUltra) to PL-SF and PL-SF 4Δ tau variants at DIV29 and DIV62. Neurons were labeled with ThioS (aggregated tau), MC1 (conformational tau), AT8 (p-tau), and Tau12 (total tau) in GFP. Tau12 was co-labeled with GFAP (RFP) to label astrocytes. Scale bar= 125 μm. **I)** Representative immunocytochemistry images of primary cortical neurons comparing PL-SF and PL-SF 4Δ at DIV29. Neurons were labeled with endogenous Hsp70 (GFP), MC1 or AT8 (RFP), and DAPI (BFP). Scale bar= 50 μm. Control= empty viral vector (pUltra**).**

Bryan et al., Figure 6.

 Figure 6: Abrogating tau-Hsp binding suppresses tau pathology in three-dimensional

organotypic hippocampal slice cultures.

Rat hippocampal slices were processed for immunocytochemistry after 9 weeks in culture

following lentiviral infection of eGFP empty vector control, wild-type tau, PL-SF tau, and PL-SF

4Δ. **A, B)** Immunocytochemistry of the entire hippocampal slice following labeling of reveal

sufficient expression of PL-SF and PL-SF 4Δ lentiviruses (GFP). Left- AT8 staining; Right- MC1

staining. CA1, CA3, and DG regions are labeled in white font. White boxes denote magnified

images in panel C, D. Scale bar= 500 μm. **C, D)** Representative immunocytochemistry images of

C) AT8 and D) MC1 labeled in RFP for empty vector control, wild-type tau, PL-SF tau, and PL-

SF 4Δ taken from the dentate gyrus (DG) region of the hippocampal slice culture (inset white

boxes from A, B). Scale bar= 100 μm.

Figure 7: HSP70 is upregulated in human AD brain and intercalates within tau aggregates.

A) Hi-Salt soluble immunoblots from control (n=5) and AD (n=6) cortical lysates probed for

Hsp70, Hsc70, and GAPDH. **B)** Quantification of hi-salt immunoblots. *= p-value significant by

unpaired Student's t-test. **C)** Urea soluble immunoblots from control (n=4) and AD (n=5) probed

for Hsp70 and Hsc70. **D)** Quantification of urea immunoblots. *= p-value significant by unpaired

Student's t-test. Overexposed Hsp70 AD sample in urea was excluded from quantification. **E)**

Representative 63X confocal single-plane z-stacks from AD human hippocampus stained with

AT8 (p-tau) and Hsp70. White arrow label neuritic plaques and white arrowheads label NFTs.

A)

Supplemental Figure 1: Accelerated PL-SF aggregation compared to single tau mutants in immortalized cell lines.

A) Representative immunocytochemistry images from QBI-293 cells expressing empty vector (pcDNA) or WT, PL, SF, and PL-SF tau variants. Cells were labeled with AT8 (p-tau) or TauC3 (C-terminal cleaved tau) with rabbit polyclonal tau (total tau) and ThioS with Tau12 (human tau). Quantification of images is found in Figure 1C. Control= empty plasmid vector (pcDNA3.1). Scale bar= 125 µm.

Supplemental Figure 2: Immunoblotting analysis of a panel of P301 and S320 single and

combination mutants.

A-G) Quantification of AT8, MC1, TauC3, and ThioS immunocytochemistry in QBI-293 cells

expressing various P301 and S320 mutants alone or in combination. Panels E-G are subsets of data

from A-C to perform additional comparisons among combinatorial mutants. Representative

images not shown. Error bars= SEM. *= significant difference by one-way ANOVA. **H)**

Representative immunoblots from QBI-293 cells expressing wild-type, P301S/S320F,

P301T/S320F, and P301L/S320F tau variants. RIPA (soluble) and urea (insoluble) fractions were

analyzed by immunoblotting for p-tau epitopes (AT8, AT180, AT270, p-396) and total tau (rabbit

polyclonal). **I, J)** Quantification of RIPA soluble and urea (insoluble) fractions. Error bars= SEM.

*= significant difference by one-way ANOVA.

Bryan et al., Supplemental Figure 3.

Supplemental Figure 3: Abrogating Hsp binding via the PL-SF 4Δ variant suppresses the accumulation of pathological tau.

Representative immunocytochemistry images from QBI-293 cells expressing tau PL-SF and PL-SF 4Δ tau variants. Cells were probed with AT8 (p-tau), TauC3 (C-terminal cleaved tau) with rabbit polyclonal tau (total tau), and ThioS with Tau12 (human tau). Quantification of images is found in Figure 1F. Control= empty plasmid vector (pcDNA3.1). Scale bar= 125 μm.

Bryan et al., Supplemental Figure 4.

Supplemental Figure 4: PLSF containing the β-structure breaking mutations, I277P/I308P, prevents tau aggregation similar to PLSF 4Δ.

A) Representative RIPA immunoblot of primary neurons transduced with empty control, PLSF 4Δ-eGFP, PLSF I277P/I308P (2IP), or PLSF-eGFP and probed for p-tau (AT8) and total tau (rabbit polyclonal). **B)** Quantification of AT8 and total tau in RIPA fraction. **C)** Representative SDS immunoblot of primary neurons transduced with empty control, PLSF 4Δ-eGFP, PLSF I277P/I308P (2IP), or PLSF-eGFP and probed for p-tau (AT8) and total tau (rabbit polyclonal. **D)** Quantification of SDS fraction. *= significant difference by one-way ANOVA.

Bryan et al., Supplemental Figure 5.

Supplemental Figure 5: PL-SF aggregates emerge similarly in either wild-type or tau knockout neurons lacking endogenous mouse tau.

A) RIPA (soluble) immunoblot of wild-type tau and tau knockout neurons probed with p-tau (AT8, AT180, AT100) and total tau (rabbit polyclonal). **B)** SDS (insoluble) immunoblot of wild-

848   type tau and tau knockout neurons probed with p-tau (AT8) and total tau (rabbit polyclonal). **C)**

849   Representative immunoblotting images of wild-type, P301L, S320F, and P301L/S320F tau

850   variants in wild-tau-expressing or tau knockout neurons. Neurons were labeled with AT8 (GFP)

851   and rabbit polyclonal total tau (RFP), MC1 (GFP), or ThioS (GFP) and Tau12 (RFP). Scale bar=

852   125 μm. Control= empty viral vector (pUltra)**.**

Bryan et al., Supplemental Figure 6.

Supplemental Figure 6: Overexpression of HSP70 in P301L-expressing neurons does not promote tau phosphorylation or aggregation.

A) RIPA (soluble) immunoblot of primary neurons following 1-week of lentiviral co-transduction with P301L tau and HSP70. Blot was probed with p-tau (AT8), total tau (rabbit polyclonal), and M2 FLAG (Hsp70-FLAG). B) SDS (insoluble) immunoblot of primary neurons was unable to detect insoluble tau species with total tau antibody (rabbit polyclonal). Control= empty viral vector (pUltra).

Bryan et al., Supplemental Figure 7.

Supplemental Figure 7: HSP90 depletion does not alter phosphorylated or aggregated PL-

SF tau.

A) RIPA (soluble) immunoblot of primary neurons probed with p-tau (AT8) and Hsp90 following

1-week of lentiviral HSP90 knockdown with two shRNAs (called TRC91 and TRC94). **B)** SDS

(insoluble) immunoblot of primary neurons probed with p-tau (AT8) and total tau (rabbit

polyclonal) following 1-week of lentiviral HSP90 knockdown. **C)** Quantification of AT8 and

Hsp90 in the RIPA fraction. Error bars= SEM. **D)** Quantification of AT8 and total tau (rabbit

polyclonal) in SDS fraction. *Note: Only TRC94, and not TRC91, HSP90 shRNA was able to

reduce Hsp90 expression.

Bryan et al., Supplemental Figure 8.

Supplemental Figure 8: Hsp70 foci accumulation within neuritic plaques and neurofibrillary

tangles in human AD brain

A) Representative stitched widefield images of a human AD patient hippocampus labeled with

AT8 (GFP) and Hsp70 (RFP). AT8/Hsp70 positive plaques are labeled with white arrowheads.

AT8/Hsp70 positive tangles are labeled with white arrows. * = a single AT8 positive tangle that is

889   absent for Hsp70 foci.

890   **Table 1: Primers used for vector construction**

Primer	Sequence
d277-278F	AATAAGAAGCTGGATCTTAGCAACGTC
d277-278R	CTGCACCTTCCCGCCTCC
d308-309F	TACAAACCAGTTGACCTGAGCAAG
d308-309R	TTGCACACTGCCGCCTCC
Tau F	CCGACCGGTGCCACCATGGCTGAGCCCCGCCAG
Tau R	CGCGTCGACTCACAAACCCTGCTTGGCCAGG
I308P-F	CAGTGTGCAAccgGTCTACAAACCAGTTGACCTGAGcaagg
I308P-R	CCGCCTCCCAGGACGtgtttg
I277P-F	GAAGGTGCAGccgATTAATAAGAAGCTGGATCTTAGCAACGtccagtcc
HSP70-F	GCGACCGGTGCCACCATGGATTACAAGGATGACGACGATAAGATGGCCAAAGCCGCGGCGATCG
HSP70-R	TCGGTCGACCTAATCTACCTCCTCAATGGTGGGGCC
HSC70-F	GCGACCGGTGCCACCATGGATTACAAGGATGACGACGATAAGATGTCCAAGGGACCTGCAGTTGG
HSC70-R	TCGGTCGACATCAACCTCTTCAATGGTGGGCCC
Hsp90-F	GCGACCGGTGCCACCATGGATTACAAGGATGACGACGATAAG
Hsp90-R	GCGGTCGACGCTAGTCTACTTCTTCCATGCGTGATGTG

References

1. Strang, K. H., Golde, T. E. & Giasson, B. I. MAPT mutations, tauopathy, and mechanisms of
neurodegeneration. *Lab. Investig.* **99**, 912–928 (2019).

2. Trzeciakiewicz, H. *et al.* An HDAC6-dependent surveillance mechanism suppresses tau-
mediated neurodegeneration and cognitive decline. *Nat Commun* **11**, 5522 (2020).

3. Koller, E. J. *et al.* Combining P301L and S320F tau variants produces a novel accelerated
model of tauopathy. *Hum Mol Genet* **28**, 3255–3269 (2019).

4. Strang, K. H. *et al.* Distinct differences in prion-like seeding and aggregation between tau
protein variants provide mechanistic insights into tauopathies. *J Biological Chem* **293**, 2408–
2421 (2017).

5. Croft, C. L. *et al.* rAAV-based brain slice culture models of Alzheimer's and Parkinson's
disease inclusion pathologies. *J Exp Medicine* **216**, 539–555 (2019).

6. Xia, Y. *et al.* Pathogenic tau recruits wild-type tau into brain inclusions and induces gut
degeneration in transgenic SPAM mice. *Commun Biology* **5**, 446 (2022).

7. Beckman, D. *et al.* Temporal progression of tau pathology and neuroinflammation in a rhesus
monkey model of Alzheimer's disease. *Alzheimer's Dement.* **20**, 5198–5219 (2024).

8. Kaniyappan, S. *et al.* FRET-based Tau seeding assay does not represent prion-like templated
assembly of Tau fibers. *Biorxiv* 2020.03.25.998831 (2020) doi:10.1101/2020.03.25.998831.

9. Xu, P. *et al.* Allosteric inhibition of HSP70 in collaboration with STUB1 augments
enzalutamide efficacy in antiandrogen resistant prostate tumor and patient-derived models.
Pharmacol. Res. **189**, 106692 (2023).

10. Li, X. *et al.* Analogues of the Allosteric Heat Shock Protein 70 (Hsp70) Inhibitor, MKT-077,
As Anti-Cancer Agents. *Acs Med Chem Lett* **4**, 1042–1047 (2013).

11. Miyata, Y. *et al.* Synthesis and Initial Evaluation of YM-08, a Blood-Brain Barrier
Permeable Derivative of the Heat Shock Protein 70 (Hsp70) Inhibitor MKT-077, Which Reduces
Tau Levels. *Acs Chem Neurosci* **4**, 930–939 (2013).

12. Moradi-Marjaneh, R., Paseban, M. & Marjaneh, M. M. Hsp70 inhibitors: Implications for the
treatment of colorectal cancer. *IUBMB Life* **71**, 1834–1845 (2019).

13. Abisambra, J. *et al.* Allosteric Heat Shock Protein 70 Inhibitors Rapidly Rescue Synaptic
Plasticity Deficits by Reducing Aberrant Tau. *Biol Psychiat* **74**, 367–374 (2013).

14. Biamonte, M. A. *et al.* Heat Shock Protein 90: Inhibitors in Clinical Trials. *J. Med. Chem.*
53, 3–17 (2010).

15. Jinwal, U. K. *et al.* Hsc70 Rapidly Engages Tau after Microtubule Destabilization*. *J Biol*
Chem **285**, 16798–16805 (2010).

16. Kundel, F. *et al.* Hsp70 Inhibits the Nucleation and Elongation of Tau and Sequesters Tau
Aggregates with High Affinity. *ACS Chem. Biol.* **13**, 636–646 (2018).

17. Patterson, K. R. *et al.* Heat Shock Protein 70 Prevents both Tau Aggregation and the
Inhibitory Effects of Preexisting Tau Aggregates on Fast Axonal Transport. *Biochemistry* **50**,
10300–10310 (2011).

18. Petrucelli, L. *et al.* CHIP and Hsp70 regulate tau ubiquitination, degradation and aggregation.
Hum Mol Genet **13**, 703–714 (2004).

19. Fontaine, S. N. *et al.* DnaJ/Hsc70 chaperone complexes control the extracellular release of
neurodegenerative-associated proteins. *EMBO J.* **35**, 1537–1549 (2016).

20. Abisambra, J. F. *et al.* DnaJA1 Antagonizes Constitutive Hsp70-Mediated Stabilization of
Tau. *J. Mol. Biol.* **421**, 653–661 (2012).

21. Thompson, A. D. *et al.* Analysis of the Tau-Associated Proteome Reveals That Exchange of
Hsp70 for Hsp90 Is Involved in Tau Degradation. *ACS Chem. Biol.* **7**, 1677–1686 (2012).

22. Sarkar, M., Kuret, J. & Lee, G. Two motifs within the tau microtubule-binding domain
mediate its association with the hsc70 molecular chaperone. *J Neurosci Res* **86**, 2763–2773
(2008).

23. Joseph, M., Anglada-Huguet, M., Paesler, K., Mandelkow, E. & Mandelkow, E.-M. Anti-
aggregant tau mutant promotes neurogenesis. *Mol Neurodegener* **12**, 88 (2017).

24. Pir, G. J. *et al.* Suppressing Tau Aggregation and Toxicity by an Anti-Aggregant Tau
Fragment. *Mol Neurobiol* **56**, 3751–3767 (2019).

25. Sydow, A. *et al.* Tau-Induced Defects in Synaptic Plasticity, Learning, and Memory Are
Reversible in Transgenic Mice after Switching Off the Toxic Tau Mutant. *J Neurosci* **31**, 2511–
2525 (2011).

26. Mocanu, M.-M. *et al.* The Potential for -Structure in the Repeat Domain of Tau Protein
Determines Aggregation, Synaptic Decay, Neuronal Loss, and Coassembly with Endogenous
Tau in Inducible Mouse Models of Tauopathy. *J Neurosci* **28**, 737–748 (2008).

27. Bergen, M. von *et al.* Mutations of Tau Protein in Frontotemporal Dementia Promote
Aggregation of Paired Helical Filaments by Enhancing Local β-Structure. *J Biol Chem* **276**,
48165–48174 (2001).

28. Smith, E. D. *et al.* Tau from SPAM Transgenic Mice Exhibit Potent Strain-Specific Prion-
Like Seeding Properties Characteristic of Human Neurodegenerative Diseases. *NeuroMolecular*
Med. **27**, 44 (2025).

29. Morito, T. *et al.* Human MAPT knockin mouse models of frontotemporal dementia for the
neurodegenerative research community. *Cell Rep. Methods* **5**, 101024 (2025).

30. Beckman, D. *et al.* A novel tau-based rhesus monkey model of Alzheimer's pathogenesis.
Alzheimer's Dement. **17**, 933–945 (2021).

31. Scheres, S. H., Zhang, W., Falcon, B. & Goedert, M. Cryo-EM structures of tau filaments.
Curr. Opin. Struct. Biol. **64**, 17–25 (2020).

32. Fitzpatrick, A. W. P. *et al.* Cryo-EM structures of tau filaments from Alzheimer's disease.
Nature **547**, 185–190 (2017).

33. Tarutani, A. *et al.* Cryo-EM structures of tau filaments from SH-SY5Y cells seeded with
brain extracts from cases of Alzheimer's disease and corticobasal degeneration. *FEBS Open Bio*
13, 1394–1404 (2023).

34. Arakhamia, T. *et al.* Posttranslational Modifications Mediate the Structural Diversity of
Tauopathy Strains. *Cell* (2020) doi:10.1016/j.cell.2020.01.027.

35. Schweighauser, M. *et al.* Cryo-EM structures of tau filaments from the brains of mice
transgenic for human mutant P301S Tau. *Acta Neuropathol. Commun.* **11**, 160 (2023).

36. Zhao, W. *et al.* Cryo-EM structures reveal variant Tau amyloid fibrils between the rTg4510
mouse model and sporadic human tauopathies. *Cell Discov.* **10**, 27 (2024).

37. Chen, D. *et al.* FTD-tau S320F mutation stabilizes local structure and allosterically promotes
amyloid motif-dependent aggregation. *Nat. Commun.* **14**, 1625 (2023).

38. Nixon, R. A. Autophagy–lysosomal-associated neuronal death in neurodegenerative disease.
Acta Neuropathol. **148**, 42 (2024).

39. Ihara, Y., Morishima-Kawashima, M. & Nixon, R. The Ubiquitin–Proteasome System and
the Autophagic–Lysosomal System in Alzheimer Disease. *Cold Spring Harb. Perspect. Med.* **2**,
a006361 (2012).

40. Keller, J. N., Hanni, K. B. & Markesbery, W. R. Impaired Proteasome Function in
Alzheimer's Disease. *J. Neurochem.* **75**, 436–439 (2000).

41. Bobkova, N. V. *et al.* Therapeutic Effect of Exogenous Hsp70 in Mouse Models of
Alzheimer's Disease. *J. Alzheimer's Dis.* **38**, 425–435 (2014).

42. Voss, K., Combs, B., Patterson, K. R., Binder, L. I. & Gamblin, T. C. Hsp70 Alters Tau Function and Aggregation in an Isoform Specific Manner. *Biochemistry* **51**, 888–898 (2012).

43. Nachman, E. *et al.* Disassembly of Tau fibrils by the human Hsp70 disaggregation machinery generates small seeding-competent species. *J Biol Chem* **295**, 9676–9690 (2020).

44. Young, Z. T. *et al.* Stabilizing the Hsp70-Tau Complex Promotes Turnover in Models of Tauopathy. *Cell Chem. Biol.* **23**, 992–1001 (2016).

45. Fontaine, S. N. *et al.* The active Hsc70/tau complex can be exploited to enhance tau turnover without damaging microtubule dynamics. *Hum Mol Genet* **24**, 3971–3981 (2015).

46. Fontaine, S. N. *et al.* Isoform-selective Genetic Inhibition of Constitutive Cytosolic Hsp70 Activity Promotes Client Tau Degradation Using an Altered Co-chaperone Complement*. *J Biol Chem* **290**, 13115–13127 (2015).

47. Yang, X. & Tohda, C. Heat Shock Cognate 70 Inhibitor, VER-155008, Reduces Memory Deficits and Axonal Degeneration in a Mouse Model of Alzheimer's Disease. *Front. Pharmacol.* **9**, 48 (2018).

48. Jinwal, U. K. *et al.* Chemical Manipulation of Hsp70 ATPase Activity Regulates Tau Stability. *J Neurosci* **29**, 12079–12088 (2009).

49. Yu, A. *et al.* Tau protein aggregates inhibit the protein-folding and vesicular trafficking arms of the cellular proteostasis network. *J Biol Chem* **294**, 7917–7930 (2019).

50. Dou, F. *et al.* Chaperones increase association of tau protein with microtubules. *Proc National Acad Sci* **100**, 721–726 (2003).

51. Vries, L. E. de *et al.* Gene-expression profiling of individuals resilient to Alzheimer's disease reveals higher expression of genes related to metallothionein and mitochondrial processes and no changes in the unfolded protein response. *Acta Neuropathol. Commun.* **12**, 68 (2024).

52. Hondius, D. C. *et al.* The proteome of granulovacuolar degeneration and neurofibrillary tangles in Alzheimer's disease. *Acta Neuropathol.* **141**, 341–358 (2021).

53. Dickey, C. A. *et al.* The high-affinity HSP90-CHIP complex recognizes and selectively degrades phosphorylated tau client proteins. *J Clin Invest* **117**, 648–658 (2007).

54. Salminen, A., Ojala, J., Kaarniranta, K., Hiltunen, M. & Soininen, H. Hsp90 regulates tau pathology through co-chaperone complexes in Alzheimer's disease. *Prog Neurobiol* **93**, 99–110 (2011).

55. Moll, A. *et al.* Hsp multichaperone complex buffers pathologically modified Tau. *Nat Commun* **13**, 3668 (2022).

56. Karagöz, G. E. *et al.* Hsp90-Tau Complex Reveals Molecular Basis for Specificity in Chaperone Action. *Cell* **156**, 963–974 (2014).

57. Cook, C. *et al.* Loss of HDAC6, a novel CHIP substrate, alleviates abnormal tau accumulation. *Hum Mol Genet* **21**, 2936–45 (2012).

58. Dickey, C. A. *et al.* Akt and CHIP coregulate tau degradation through coordinated interactions. *P Natl Acad Sci Usa* **105**, 3622–7 (2008).

59. Weickert, S., Wawrzyniuk, M., John, L. H., Rüdiger, S. G. D. & Drescher, M. The mechanism of Hsp90-induced oligomerizaton of Tau. *Sci Adv* **6**, eaax6999 (2020).

60. Rutledge, B. S., Choy, W.-Y. & Duennwald, M. L. Folding or holding?—Hsp70 and Hsp90 chaperoning of misfolded proteins in neurodegenerative disease. *J. Biol. Chem.* **298**, 101905 (2022).

61. Jinwal, U. K. *et al.* Imbalance of Hsp70 family variants fosters tau accumulation. *FASEB J.* **27**, 1450–1459 (2013).

62. Miyata, Y. *et al.* Cysteine Reactivity Distinguishes Redox Sensing by the Heat-Inducible and Constitutive Forms of Heat Shock Protein 70. *Chem. Biol.* **19**, 1391–1399 (2012).

63. Shelton, L. B. *et al.* Hsp90 activator Aha1 drives production of pathological tau aggregates. *Proc National Acad Sci* **114**, 9707–9712 (2017).

64. Jinwal, U. K. *et al.* The Hsp90 Cochaperone, FKBP51, Increases Tau Stability and Polymerizes Microtubules. *J. Neurosci.* **30**, 591–599 (2010).

65. Blair, L. J. *et al.* Accelerated neurodegeneration through chaperone-mediated oligomerization of tau. *J. Clin. Investig.* **123**, 4158–4169 (2013).

66. Webster, J. M. *et al.* Hsp22 with an N-Terminal Domain Truncation Mediates a Reduction in Tau Protein Levels. *Int. J. Mol. Sci.* **21**, 5442 (2020).

67. Ospina, S. R. *et al.* Small Heat Shock Protein 22 Improves Cognition and Learning in the Tauopathic Brain. *Int J Mol Sci* **23**, 851 (2022).

68. Abisambra, J. F. *et al.* Phosphorylation Dynamics Regulate Hsp27-Mediated Rescue of Neuronal Plasticity Deficits in Tau Transgenic Mice. *J. Neurosci.* **30**, 15374–15382 (2010).

69. Barghorn, S., Biernat, J. & Mandelkow, E. Amyloid Proteins, Methods and Protocols. *Methods Mol. Biol. (Clifton, NJ)* **299**, 35–51 (2004).

70. Farizatto, K. L. G., Almeida, M. F., Long, R. T. & Bahr, B. A. Early Synaptic Alterations and Selective Adhesion Signaling in Hippocampal Dendritic Zones Following Organophosphate Exposure. *Sci. Rep.* **9**, 6532 (2019).

71. Almeida, M. F. *et al.* Distinct and dementia-related synaptopathy in the hippocampus after
military blast exposures. *Brain Pathol.* **31**, e12936 (2021).

72. McKenna, M. *et al.* Organotypic whole hemisphere brain slice models to study the effects of
donor age and oxygen-glucose-deprivation on the extracellular properties of cortical and striatal
tissue. *J. Biol. Eng.* **16**, 14 (2022).

73. Bahr, B. A. *et al.* Amyloid β protein is internalized selectively by hippocampal field CA1 and
causes neurons to accumulate amyloidogenic carboxyterminal fragments of the amyloid
precursor protein. *J. Comp. Neurol.* **397**, 139–147 (1998).

74. Caba, E. *et al.* Excitotoxic stimulation activates distinct pathogenic and protective expression
signatures in the hippocampus. *J. Cell. Mol. Med.* **25**, 9011–9027 (2021).

75. Farizatto, K. L. G., Ikonne, U. S., Almeida, M. F., Ferrari, M. F. R. & Bahr, B. A. Aβ42-
mediated proteasome inhibition and associated tau pathology in hippocampus are governed by a
lysosomal response involving cathepsin B: Evidence for protective crosstalk between protein
clearance pathways. *PLoS ONE* **12**, e0182895 (2017).

1092