Title

Caecilians maintain a functional long-wavelength-sensitive cone opsin gene despite signatures of

relaxed selection and more than 200 million years of fossoriality

Running title

A long-wavelength cone opsin in caecilians

Authors

María José Navarrete Méndez[1,*], Sina S. Amini[1,*], Juan Carlos Santos[2], Jacob Saal[1], Marvalee H.

Wake[1], Santiago R. Ron[3], and Rebecca D. Tarvin[1]

[1] Department of Integrative Biology and Museum of Vertebrate Zoology, University of

California Berkeley, Berkeley, CA, USA 94720

[2] Department of Biological Sciences, St John's University, NY, USA 11439

[3] Museo de Zoología, Facultad de Ciencias Exactas y Naturales, Pontificia Universidad Católica

del Ecuador, Quito, Ecuador

[*] Co-first authors

Corresponding author contact information

Rebecca Tarvin: rdtarvin@berkeley.edu

Author contributions

RDT, MJNM, SSA, JCS, SRR conceptualized the study; RDT, MJNM, SRR, SSA, JCS, MHW,

JS collected data; MJNM, RDT, SSA, SSR performed analyses; MJNM, SSA, RDT, MHW, JS,

JCS, SSR performed the investigation; RDT, SRR, JCS acquired funding; SRR, MJNM, MHW

contributed samples and helped with permits; MJNM, SSA, JS, MHW, JCS, RDT, SSR prepared

the original draft; RDT coordinated and supervised the project; all authors reviewed and revised

the manuscript.

Conflict of interests

The authors declare no conflict of interests.

Acknowledgments. We thank Rayna Bell and Lauren Sheinberg at the California Academy of

Sciences, Carol Spencer at the Museum of Vertebrate Zoology, and Fernando Ayala and Diego

Paucar at QCAZ for facilitating access to caecilian tissues for sequencing. We thank José

Simbaña and Yanayacu Biological Station for permission and facilitation of specimen collection,

Alexander Stubbs and Diego Paucar for their assistance in caecilian dissections and specimen

preparation, Todd Oakley and Seth Frazer for their support in running the machine learning

models to estimate λ_{max} values for our sampled genes, Alex Somlyay for assistance with

microscopy, Mike Boots for lending us use of his microscope and camera, Valeria Ramírez-

Castañeda and Lydia Smith for conducting PCRs on MVZ tissues, Claudia Terán for laboratory

work at QCAZ in Quito. Yocelyn Gutiérrez-Guerrero provided advice and guidance in the

transcriptome analysis. For specimen collection in Ecuador, we thank William Booker, Verónica

Crespo, Juan Guayasamin, Giovanna Romero, Yerka Sagredo, Scott Tiegs, and David

Velalcazar. We thank Ryan Schott for his guidance on selection analyses, Simon Scarpetta, the

two anonymous reviewers and the editor for their suggestions that improved the manuscript.

Funding. Funding was provided by Secretaría Nacional de Educación Superior, Ciencia,

Tecnología e Innovación del Ecuador SENESCYT (Arca de Noé initiative; SRR and Omar

Torres principal investigators), NIH NIGMS R35GM150574 to RDT, UC Berkeley start-up

funding to RDT, fellowships from the American Association of University Women and

Philomathia to MJN, and long-term support from the USA National Science Foundation to

MHW. The Ecuadorian Ministry of Environment provided permits for specimen collection and

genetic research: 008-09 IC-FAU-DNB/MA, 005-12 IC-FAU-DNB/MA, 003-15 IC-FAU-

DNB/MA, MAE-DNB-CM-2015-0025-M-0001, and MAAE-DBI-CM-2022-0230. Finally, we

acknowledge and express gratitude for the animals involved in this study.

Data accessibility statement

Sequences have been deposited in GenBank under accession numbers PQ541071–PQ541084 and

in the Sequence Read Archive under project number PRJNA1181370. Voucher specimens are

housed in the Museo de Zoología (QCAZ) at the Pontificia Universidad Católica del Ecuador,

the Museum of Vertebrate Zoology (MVZ) at UC Berkeley, and the Herpetology Collection of

the California Academy of Sciences. All other data are provided as supplementary material.

Supplementary datasets have been updated and archived in Dryad (DOI:

10.5061/dryad.h18931zxf)

65
66

67 **TITLE**

68 Caecilians maintain a functional long-wavelength-sensitive cone opsin gene despite signatures of

69 relaxed selection and more than 200 million years of fossoriality

TEASER TEXT

The colonization of low-light environments exerts selective pressure to enhance non-visual senses while relaxing selection on vision, often leading to opsin gene loss. We present compelling evidence for the retention of a cone opsin gene (*LWS*) in caecilians—a fossorial amphibian lineage with a 200-million-year history of burrowing and reduced eyes. Despite signs of relaxed selection, *LWS* appears functional, echoing patterns in other low-light-adapted vertebrates. Additionally, the loss of key cone phototransduction genes and absence of clear cone cells raises the possibility of *LWS* expression in rods or a functional change in its use. Our study provides an evolutionary framework for investigating *LWS* retention in fossorial vertebrates.

ABSTRACT

Vision is tuned to animals' ecologies, evolving in response to specific light environments and visual needs. Transitions to fossorial lifestyles impose strong selective pressures favoring adaptations for underground life, such as increased skull ossification and reduced eye protrusion. Fossoriality may simultaneously relax constraints on vision leading to diminished visual capabilities. Caecilians (Gymnophiona)—specialized, fossorial amphibians—possess reduced eyes covered by skin or bone. For years, these traits, along with the presence of a single photoreceptor expressing one functional opsin gene, have been interpreted as evidence of limited vision, including an inability to focus or perceive color. Our results challenge these assumptions: we identified the long-wavelength-sensitive (*LWS*) opsin gene in 13 species of caecilians spanning 8 of 10 recognized families. Molecular evidence indicates that *LWS* is intact and transcribed in the eye of at least one species (*Caecilia orientalis*). However, the specific

photoreceptor type expressing *LWS* remains uncertain, as our survey of cone phototransduction

genes revealed a mosaic of losses, and anatomical observations from five families did not

conclusively identify cone-like cells, though they revealed highly organized retinae even in

families with vestigial eyes. Altogether, our results suggest that vision in caecilians may be

underestimated and the role of light perception in their ecology is possible.

KEYWORDS: visual opsins, relaxed selection, vision, OPN1LW, *LWS*, phototransduction

cascade

1. INTRODUCTION

Strong selection for adaptation to new habits, such as fossoriality (i.e., organisms that live

underground), results both in gain and loss of traits (Nevo, 1979). For example, some burrowing

frogs evolved spades through the enlargement of metatarsal tubercles (Emerson, 1976), yet their

altered body and limbs trade off with jumping ability (Vidal☐García et al., 2014). The star-nosed

mole (*Condylura cristata*) evolved a novel mechanosensory structure that allows them to "see"

prey (Catania, 1999; Holmberg, 2022) despite having small, underdeveloped eyes (Carmona et

al., 2010). In these and other animals, subterranean life intensifies selection pressures for

adaptations that facilitate life underground and relaxes selection for traits that are no longer as

relevant.

Although relaxed selection can lead to trait degeneration, it is rare that traits are ever

completely lost because they may be developmentally constrained, still possess some functional

value, or cost little to maintain (Sadier et al., 2022). Partial eye degeneration without complete

loss is a common characteristic of fossorial, deep-sea, and cave-dwelling vertebrates. For

example, the European mole (*Talpa europaea*) and Iberian mole (*Talpa occidentalis*) have

organized retinae with rod and cone cells despite being fully subterranean (Carmona et al., 2009;

Glösmann et al., 2008); *T. occidentalis* never even opens its eyelids. Eyeless populations of the

cave-dwelling fish *Astyanax mexicanus* develop a layered retina during their embryonic stage,

retaining the ability to exhibit light-entrained retinomotor rhythms even as their eyes undergo

extensive ontological degeneration (Emam et al., 2020; Espinasa & Jeffery, 2006).

Eye degeneration can be coupled at the molecular level with the loss of visual opsin

genes that encode sensitive photoreceptor proteins (Bowmaker, 2008; Emerling & Springer,

2014; Lin et al., 2024). For example, the Cape golden mole (*Chrysochloris asiatica*), which is

almost exclusively subterranean and has subcutaneous eyes, exhibits one of the highest numbers

of pseudogenes among fossorial mammals, including losses of cone opsins and several other

genes involved in the phototransduction cascade (Emerling & Springer, 2014). However, opsin

genes may be lost even when ocular anatomy remains well-developed, including features such as

an organized and layered retina (Foureaux et al., 2010; Wake, 1985; Warrant, 2000). Across

vertebrates, independent losses of opsins have been proposed as adaptations to specific visual

demands shaped by behavioral and ecological shifts (W. L. Davies, 2011). For instance, at least

10% of mammalian species are thought to lack functional copies of the short-wavelength

sensitive opsin gene (*SWS1*), and all placental mammals have lost the middle-wavelength

sensitive opsin gene (*RH2*)–patterns attributed to nocturnal lifestyles and/or ancestry (W. I. L.

Davies et al., 2012; Hunt et al., 2009; Jacobs, 2013).

Interestingly, although opsins have been lost multiple times in vertebrates, others tend to

persist (W. L. Davies, 2011), even when environmental conditions seem unsuitable for their

spectral sensitivity. Among the five visual opsin genes found in vertebrates (i.e. *RH1*, *RH2*, *LWS*,

SWS1, *SWS2*), the *LWS* opsin exhibits peak sensitivity to the longest wavelengths within the

visible spectrum (560 – ~570 nm, corresponding to red-green hues, though sensitivity can extend

to wavelengths beyond the peak absorbance; (Liebman & Entine, 1968; Margetts et al., 2024)

and is one of the most consistently retained across vertebrates, including species adapted to low-

light environments (W. I. L. Davies et al., 2012; Ford et al., 2024; Musilova et al., 2021).

Ancestors of caecilians, a group of limbless amphibians, began their transition to a

subterranean lifestyle more than 250 million years ago (MYA) through elongation of the trunk,

fusion of cranial bones, and reduction of limbs (Jenkins & Walsh, 1993; Santos et al., 2020). In

extant caecilians (with most recent common ancestor 190 MYA; Kligman et al., 2023) all species

lack limbs, have eye sockets that are covered by skin or by skin and bone, and lack the eye

muscles for accommodation (Wake, 1985). Visual system reduction correlates broadly with

fossorial specialization (Himstedt, 1995; Kamei, San Mauro, et al., 2012): early-diverging, less

fossorial caecilian lineages like ichthyophiids and rhinatrematids retain better-developed eyes,

while taxa with more derived fossorial adaptations–such as *Boulengerula boulengeri*–often

exhibit extreme ocular reduction (Mohun & Wilkinson, 2015; Wake, 1985). However, certain

aspects of visual system anatomy—such as the degree of degeneration in extrinsic eye muscles,

lens structure, and even retinal organizations—display a mosaic pattern of trait loss across

species, independent of phylogeny, ecology, or cranial ossification (Mohun & Wilkinson, 2015;

Wake, 1985).

In spite of these morphological constraints, evidence from anatomy, physiology, and

molecular biology has shown that visual systems in caecilians have remained somewhat

functional (Dünker, 1998; Mohun et al., 2010; Mohun & Wilkinson, 2015; Wake, 1985). For

example, negative phototactic behavior has been reported in *Ichthyophis cf. kohtaoensis*

(Himstedt, 1995; Mohun et al., 2010). Lens morphology appears to be adapted to environment,

as the aquatic/semi-aquatic family Typhlonectidae possesses spherical lenses (similar to most

aquatic vertebrates), in contrast to the ellipsoid lenses typical of most terrestrial caecilians

(Wake, 1985). Furthermore, the rod phototransduction pathway used for scotopic vision remains

functional in Gymnophiona, as evidenced by electroretinography and microspectrophotometry

analyses that detected a visual pigment with peak absorbance near 500 nm (consistent with the

spectral properties of *RH1*) in *Ichthyophis* cf. *kohtaoensis* (Himstedt, 1995; Mohun et al., 2010)*,*

Rhinatrema bivittatum, Geotrypetes seraphini, and *Typhlonectes natans* (Himstedt, 1995; Mohun

et al., 2010). The identity of this pigment was later confirmed in two of these species (i.e. *I.* cf.

kohtaoensis and *T. natans*) as the rod opsin *RH1* (Himstedt, 1995; Mohun et al., 2010).

Moreover, although caecilian retinas are thought to lack cone cells (Mohun & Wilkinson, 2015;

Wake, 1985), a gene encoding the long-wavelength-sensitive cone opsin (*LWS*) was recently

identified in three published caecilian genomes from the families Rhinatrematidae,

Dermophiidae, Siphonopidae (Lin et al., 2024).

Here we investigated the evolution of the two visual opsin genes known to be present

(*LWS* and *RH1*) in caecilians and explore why *LWS* has been maintained without loss or

pseudogenization for hundreds of millions of years despite significant eye degeneration and an

ecological context wherein it would otherwise appear superfluous. Given that intact copies of the

LWS gene have been identified in three distantly related caecilian species exhibiting varying

degrees of skull ossification and eye reduction (Lin et al., 2024), we hypothesize that *LWS* was

present in the common ancestor of caecilians and has been maintained in all extant species,

independent of eye exposure or anatomical complexity. To test this, we reconstructed the

evolution of eye musculature and exposure using ancestral state reconstruction to infer the

ancestral condition, trace the trajectory of eye degeneration, and evaluate the persistence of *LWS*

in this anatomical context.

Additionally, given that caecilian *LWS* genes exhibit intact open reading frames without

premature stop codons, we hypothesize that they remain functional and may be expressed in the

eye, with absorbance properties resembling those of long-wavelength-sensitive opsins in

amphibian cones. To test this, we sequenced the eye transcriptome of one caecilian species and

used a machine learning approach to estimate the absorbance peak of caecilian *LWS* based on its

amino acid sequence.

Finally, to investigate whether cone photoreceptors capable of expressing *LWS* persist in

caecilians, we used two complementary approaches. First, we examined the genomes and

transcriptomes of caecilian species for genes associated with the cone signaling cascade, under

the hypothesis that if *LWS* is expressed in cones, the downstream components of the cone

phototransduction pathway should also be present. Second, we reviewed histological material to

search for evidence of cone photoreceptors. We further hypothesize that selection analyses will

reveal evolutionary constraints on *LWS* that parallel patterns observed for the rod opsin gene

RH1, thereby supporting its continued physiological relevance. As one of the earliest vertebrate

lineages to adopt a subterranean lifestyle, caecilians provide a unique opportunity to explore the

evolution and persistence of light perception in fossorial environments.

2.   METHODS

2.1 ANCESTRAL CHARACTER STATE RECONSTRUCTION

We estimated ancestral character states for eye musculature and exposure using character

data mainly from Wake (1993), but also, for a few species, from Norries & Hughes (1918),

Taylor (1968, 1969), and Maciel and Hoogmoed (2011) (see Supplementary Table 1). The

reconstructions were performed using stochastic character mapping in *phytools* v2.0 (Revell,

2024) on a time-tree obtained from the San Mauro et al. (2014) matrix of mitochondrial

genomes. The time-tree was inferred using BEAST v2.7.8 (Bouckaert et al., 2019) under a strict

clock and the model of DNA evolution HKY. See Supplementary Methods section 2.1 for more

details.

2.2 MORPHOLOGY AND SEARCH FOR CONE PHOTORECEPTORS

2.2.1 Histology. The examined sections are archival material accessioned in the MVZ

that were originally prepared by MHW in the 1980s. Details on sample preparation, staining, and

storage are provided in Supplementary Methods section 2.2.1. Sampling was opportunistic,

representing all species with available histological sections suitable for visualizing eye

morphology. More information about these species and accession information can be found in

Supplementary Table 2 (see also (Wake, 1985).

2.2.2 Photomicrography. Eyes were observed with a Laxco microscope, where we

examined sections with intact photoreceptors and retina. Photographs were taken at standard low,

medium, and high magnifications using a Laxco SeBaCam5C digital microscope camera, and

rendered using SeBaView software v3.7 (Laxco, Inc.). A stage micrometer was used for

measurements and to calibrate scale bars.

2.2.3 Search for cones and description of the retina. We visually surveyed all images

for photoreceptor cells with morphology characteristic of cones (i.e., bulbous ellipsoids, tapered

outer segments that are much thinner than their inner segments) or otherwise distinct from the

rods that predominate the retina. Slides of *Epicrionops* sp., *Typhlonectes compressicauda*,

Ichthyophis kohtaoensis, *Dermophis mexicanus*, and *Scolecomorphus kirkii* specimens were

selected to highlight taxonomic and morphological diversity of retinae across Gymnophiona due

to their taxonomic disparity and well-preserved morphology. Standard descriptions of the retinae

were made following (Mohun & Wilkinson, 2015; Wake, 1985) and (Himstedt, 1995; Mohun et

al., 2010). Details on measurement methodology are provided in Supplementary Methods section

2.2.3.

2.3 FURTHER VALIDATION OF A CAECILIAN LWS AND ASSESSMENT OF ITS

POTENTIAL FUNCTIONALITY

2.3.1 Generating an eye transcriptome. We collected one individual of *Caecilia*

orientalis in 2020 at Yanayacu Biological Station in Napo, Ecuador (S 0.5984097, W

77.8907409, 2143 m.a.s.l.) under permit MAE-DNB-CM-2015-0025-M-0001 issued by the

Ministerio del Ambiente of Ecuador to Universidad Católica del Ecuador.

We euthanized the specimen using commercial Roxicaine anesthetic (lidocaine) and

immediately dissected both eyes which were preserved in RNAlater. The specimen is deposited

in the collection of Museo de Zoología of the Pontificia Universidad Católica del Ecuador

(QCAZ), with specimen number QCAZ:A:77338. Tissue samples were sent to Macrogen Inc.

(Seoul, South Korea) for RNA extraction, library preparation using the Illumina TruSeq Stranded

mRNA LT Sample Preparation Kit, and sequencing on the Illumina NovaSeq6000 to 100 million

paired-end 100-bp reads.

Read quality was assessed using FastQC (Andrews, 2010). We used TrimGalore!

(https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/) to remove adapters, low-

quality bases, and short reads. Transcriptome assembly was then conducted *de novo* using

Trinity v2.13.2 (Grabherr et al., 2011), incorporating all paired-end reads with default settings.

The resulting assembly was filtered using SeqClean (Chen et al., 2007). To annotate protein-

coding genes, TransDecoder v2.026 (Haas, n.d.) was used with default parameters and homology

to UniProt was then used to infer functions of predicted genes with NCBI-BLASTP v2.2.30+ (E-

value = 1e–6) and Interproscan 5.35 (Jones et al., 2014) to detect proteins with known functional

domains. Finally, we used Salmon v1.10.3 (Patro et al., 2017) to quantify transcript abundances

of opsin genes. First, we built a Salmon index for our transcriptome using the index command

with default parameters for k-mer length. Next, we quantified the paired-end reads against the

previously generated index using the quant command. The output gene counts were summarized

as normalized transcript abundances using DESeq2 v3.11 (Love et al., 2014).

2.3.2 PCR of **LWS** *in additional caecilian species.* To validate whether *LWS* existed in a

broader phylogenetic and ecological sampling of caecilians, we obtained tissue samples from

terrestrial and aquatic species exhibiting varying degrees of fossoriality and reports of above-

ground behavior. We included samples of species from early branching lineages that exhibit

lesser degrees of adaptation to fossoriality including *Epicrionops bicolor*, *E. petersi*

(Rhinatrematidae), as well as more phenotypically derived species such as *Caecilia orientalis*

(Caecilidae) and *Microcaecilia albiceps* (Siphonopidae) from specimens at QCAZ. Additional

tissues were obtained from the Museum of Vertebrate Zoology (MVZ) at the University of

California, Berkeley, including terrestrial species that inhabit moist, loose soils with varying

levels of eye coverage such as *Dermophis mexicanus*, *Geotrypetes seraphini*, *Gymnopis*

multiplicata (Dermophiidae), as well as *Ichthyophis kohtaoensis* (Ichthyophiidae) which has

aquatic larvae and terrestrial adults, and the fully aquatic *Typhlonectes natans* (Typhlonectidae).

Tissues from species with markedly reduced eyes, such as *Boulengerula boulengeri* (Herpelidae)

and *Scolecomorphus vittatus* (Scolecomorphidae), were kindly loaned from the California

Academy of Sciences. We attempted to obtain tissues that match our morphological material to

the extent possible (at least to genus). Together, these samples represent the full range of

diversity in caecilian eye morphology. See Supplementary Table 3 for the accession numbers.

DNA extraction, amplification, and sequencing was carried out at PUCE for QCAZ

specimens, and at the Evolutionary Genomics Lab (EGL), UC Berkeley, for MVZ and CAS

specimens. Fragments of *LWS* were amplified using new primer sets designed from an alignment

of amphibian opsins. The new LWS_542R (5'-GCA GAC CAA ACC CAG GAG AA-3') and

LWS_117F (5' TTT TGA GGG MCC CAA CTA CC -3') primers amplify a region that includes

part of exon I, the intron between exon I and II, and part of exon II. Specifics about the

extraction and amplification protocols are provided in the Supplementary Material Methods

section 2.3.2. Additionally, an evaluation of primers used in previous attempts that did not yield

successful amplification of *LWS* in caecilians (Mohun et al., 2010) is provided in Supplementary

Material Methods section 2.4.

PCR products from QCAZ samples were sequenced at Macrogen Inc. (Seoul, South

Korea); MVZ and CAS samples were sequenced at the UC Berkeley DNA Sequencing Facility.

We assembled forward and reverse sequences into consensus sequences using default parameters

in Geneious Prime 2022.1.1 and then aligned them against each other using MUSCLE (Edgar,

2004). To confirm the identity and quality of the new *LWS* sequences, we inferred gene trees for

LWS and *RH1* in IQTREE2. Sequences were partitioned by coding and non-coding regions and

also by codon position. The best-partition model was estimated using the -TESTMERGE option

(Chernomor et al., 2016; Kalyaanamoorthy et al., 2017). Bootstrap values were calculated using

500 replicates (-b 500) and were plotted on the best likelihood tree. See Supplementary Data 1

for input and output files (Supplementary Data 1-7 are archived at Dryad—DOI:

10.5061/dryad.h18931zxf).

***2.3.3 Estimation of the wavelength of maximum absorbance (λ_{max}) of* LWS *and* RH1.**

Given that the *LWS* copies identified in caecilian genomes contain open reading frames without

stop codons (i.e., considereded intact), we hypothesize that the gene is functional and that the

protein encoded by *LWS* will exhibit absorbance spectra within the expected ranges (560–570

nm; (W. I. L. Davies et al., 2012). To test this hypothesis, we used a machine-learning model

(Frazer et al., 2024) to estimate the wavelength of maximum absorbance (λ_{max}) for five full-

length *LWS* sequences from the caecilian species included in our study incorporating data

available on GenBank (*Rhinatrema bivittatum*, XM_029609805.1; *Microcaecilia unicolor,*

XM_030192059.1; *Geotrypetes seraphini*, XM_033922369.1; *Ichthyophis bannanicus*,

GCA_033557465.1 hereafter *I. kohtaoensis* following Nishikawa et al., 2021) and our *C.*

orientalis eye transcriptome. Additionally, we estimated absorbance values for the *RH1*

sequences of the same species listed above to validate the accuracy of the machine-learning

model, since experimental absorbance estimates for this opsin have been reported for *G.*

seraphini and *R. bivittatum* (Mohun et al., 2010). We expected the estimated λ_{max} values for *RH1*

to fall within the typical range for this opsin (~500 nm; W. L. Davies, 2011) and to closely

approximate experimental MSP data. Furthermore, we increased our phylogenetic sampling by

including *LWS* and *RH1* sequences from other relevant amphibian and vertebrate species. For

this purpose, we explored three types of sequence archives: NCBI sequence references,

unannotated transcriptomes, and published genomes. The methods used to retrieve *LWS* from

each of these datasets are described in more detail in the Supplementary Material section 2.3.3

Our final dataset included 19 species of Anura and 8 species of Caudata (see Supplementary

Table 3 for details). The nucleotide matrices were manually inspected to remove any ambiguities

and the sequences were translated to amino acids for downstream analysis (see Supplementary

Data 2).

Point estimations of λ_{max} for *LWS* and *RH1* were predicted using the vertebrate model

trained with the VPOD_vert_het_1.0 dataset available in the Visual Physiology Opsin Database

(https://github.com/VisualPhysiologyDB/visual-physiology-opsin-db). This model demonstrated

the best predictive ability, with the highest 10-fold cross-validation R^2 (0.968) and the lowest

mean absolute error (6.56 nm) among all the models compared in the machine learning analysis

(Frazer et al., 2024). The training dataset included 721 vertebrate opsins (UVS, SWS, MWS,

LWS, RH1, RH2) spanning wild-type, mutant, and chimeric sequences with λ_{max} values ranging

from 350 to 611 nm (Frazer et al., 2024).

Additionally, a literature search was conducted to identify available

microspectrophotometry (MSP) data for the species included in our machine-learning analysis.

In some instances, when values were not available for a particular species, we used data from a

closely related species within the same genus (Supplementary Table 4). We then performed a

direct comparison between the MSP data and the machine-learning estimations using a linear

correlation analysis (Pearson's correlation) using the *lm* and *cor* test functions in Rstudio

v2024.9.1.394 and calculated the mean absolute error to quantify the discrepancy between the

absorbance values obtained from the two methodologies.

2.3.4 Search for presence and expression of rod and cone phototransduction cascade

genes. The presence of *LWS* in caecilians contrasts with the lack of morphological evidence for

cone photoreceptors, which are the typical cellular structure for *LWS* expression. We

hypothesized that if *LWS* is expressed and functional in cones, intact cone phototransduction

genes should also be present and expressed, and thus detectable in both the genomes and

transcriptomes of caecilians. Alternatively, if caecilian retinas contain only rod photoreceptors,

as previously reported, and *LWS* is potentially co-expressed with *RH1* in rods, we would expect

to find only rod-associated phototransduction genes, with cone-specific genes either

pseudogenized, absent, or not expressed in the eye transcriptome. To test this, we searched for

rod and cone phototransduction genes in NCBI reference databases and publicly available

genomes of three anuran species: *Xenopus tropicalis* (GCF_901001135.1), *Nanorana parkeri*

(GCF_000935625.1), and *Pyxicephalus adspersus* (GCA_032062135.1); two salamanders:

Ambystoma mexicanus (GCA_040938575.1) and *Pleurodeles waltl* (GCA_031143425.1); and

three caecilians: *Microcaecilia unicolor* (GCF_901765095.1), *Rhinatrema bivittatum*

(GCF_901001135.1), and *Geotrypetes seraphini* (GCF_902459505.1). Additionally, we searched

for these genes in our newly generated eye transcriptome from *Caecilia orientalis*

(PRJNA1181370) (Supplementary Table 3). The canonical visual phototransduction cascade in

vertebrates involves three main components: (1) the heterotrimeric G-protein transducins (in

rods: *GNAT1*, *GNB1*, *GNGT1*; in cones: *GNAT2*, *GNB3*, *GNGT2*); (2) the phosphodiesterase

enzymes (in rods: *PDE6A*, *PDE6B*, *PDE6G*; in cones: *PDE6C*, *PDE6H*), and (3) the cyclic

nucleotide-gated channels (in rods: *CNGA1*, *CNGB1*; in cones: *CNGA3*, *CNGB3*) (Lamb, 2013).

The identified sequences of our genes of interest were downloaded and used as reference

queries to search our *de novo*–assembled transcriptome data, which had been formatted as a

custom BLAST database using the default parameters for *makeblastdb* (v2.16.0+). All the

obtained sequences were translated into amino acids and aligned using MAFFT v7.490 (Katoh &

Standley, 2013), and the alignments were manually inspected in Geneious Prime 2022.1.1

(https://www.geneious.com) for the presence of premature stop codons. Finally, following the

methodology described in section 2.2.1, we used the previously generated Salmon index of the

eye transcriptome to quantify transcript abundances of visual phototransduction genes and assess

whether their expression levels mirror those of the opsin genes. When multiple isoforms were

reported for a given gene, we summed the expression levels of all isoforms. Although some

isoforms may result from assembly artifacts, we opted to include all of them to avoid making an

arbitrary choice about which isoform to retain.

2.4 SELECTION ANALYSES

The presence of intact *LWS* genes without nonsense and frameshift mutations in five

phylogenetically diverse caecilian species supports the idea that *LWS* is an ancestrally retained

feature despite a long fossorial history. Thus, we hypothesized that *LWS* is under evolutionary

constraint, similar to *RH1*, indicating a physiological role similar to its ancestral function. To test

this, we conducted a series of selection analysis on *LWS* and *RH1* in Gymnophiona and other

amphibian species using the sequence data described in section 2.2.3, including transcriptomic

data from *C. orientalis* and *LWS* fragments obtained via Sanger sequencing. In total, our

sampling included 13 caecilian species spanning eight families (Supplementary Table 3). An

updated and unrooted version of the phylogenetic tree presented by (Jetz & Pyron, 2018) served

as a backbone for conducting selection analysis in the codeml program of PAML v4 (Yang,

2007).

Sequences for *LWS* and *RH1* were aligned using MAFFT (Katoh & Standley, 2013)

followed by manual editing to remove terminal stop codons in Mesquite v3.6 (Maddison &

Maddison, 2015). For *LWS*, we ran analyses on two types of datasets, a complete dataset

including the full coding sequence of the gene with smaller sampling for Gymnophiona (368 aa;

Anura = 19 spp, Caudata = 8 spp, Gymnophiona = 5 spp) and an incomplete dataset with more

species for Gymnophiona but fewer sites (132 aa; Anura = 19 spp; Caudata = 8 spp,

Gymnophiona = 13 spp). For *RH1*, we only ran analyses using the complete sequences (354 aa;

Anura = 18 spp, Caudata = 6 spp, Gymnophiona = 5 spp).

We initially conducted a broad phylogeny-wide analysis to assess the strength of

purifying selection on *LWS* and *RH1*. We used the M0 model, which assumes a single ω (dN/dS)

ratio across all sites and lineages. To represent the null hypothesis of neutral evolution, we fixed

ω at 1 (fix_omega = 1; omega = 1), thereby constraining the model to assume no selection. To

test our hypothesis that *LWS* has been maintained under purifying selection, we ran an alternative

model in which ω was estimated from the data (fix_omega = 0; omega = 0.5). We then compared

the likelihoods of the two models using a likelihood ratio test (LRT).

We also inferred selection patterns and tested for site-specific positive selection for each

gene comparing the support for different models (M0, M1a, M2a, M2a_rel, M3, M7, M8a, and

M8). We used the Bayes Empirical Bayes (BEB) approach to identify individual sites with a

posterior probability >90% of being in the positively selected class of sites. Additionally, we

compared the fit of clade models C and D (CmC vs. CmD) (Bielawski & Yang, 2004) to M3 and

M2a_rel to investigate long-term shifts in selection pressures associated with the evolution of

fossoriality in caecilians (Weadick & Chang, 2012). Both clade models assume three site classes,

with the third class modeling divergent selection among clades; however, unlike CmC, CmD

does not constrain the ω estimates for any site class. The best model was identified using

likelihood ratio tests (LRTs) with a χ^2 distribution.

Additionally, we conducted a series of analyses in HYPHY, implemented on the

Datamonkey web server (Delport et al., 2010). FEL (Fixed Effect Likelihood; Kosakovsky Pond

& Frost, 2005) and FUBAR (Fast, Unconstrained Bayesian AppRoximation; Murrell et al., 2013)

were employed to estimate positive or purifying selection across sites by comparing single

estimates of dN and dS values for each site. FUBAR uses a hierarchical Bayesian method rather

than maximum likelihood, making it more sensitive than FEL to weaker signatures of positive

selection (i.e., when ω is close to 1) (Murrell et al., 2013). We used BUSTED (Branch-site

Unrestricted Statistical Test for Episodic Diversification; Murrell et al., 2013) to identify

episodic positive selection at a fraction of sites. BUSTED allows for branch-to-branch variation

across the entire phylogeny. Finally, we ran RELAX (Wertheim et al., 2015) to detect relaxation

of selective constraint. This methodology requires an *a priori* specification (e.g., partition) of a

subset of branches into foreground and background. In our case, the foreground comprised the

crown Gymnophiona and its stem branch, while anurans and salamanders were designated as the

background. RELAX estimates a selection intensity parameter (k), where $k < 1$ indicates relaxed

selection and $k > 1$ indicates intensified selection, based on shifts in ω values between test and

reference lineages.

We report results from the selection analyses using amino acid site numbers based on

bovine rhodopsin (NP_001014890.1; alignments with bovine rhodopsin are available in the

Supplementary Data 3). Selection results are reported by referring to the location of each amino

acid according to the 3D structure of opsins, which encompasses seven transmembrane domains

(TMD I–VII), three extracellular domains (ECD I–III), and the amino- and carboxyl- termini (N

and C) (Palczewski et al., 2000). All input and output files for PAML and HYPHY selection

analyses are available in Supplementary Data 4.

Finally, we examined *LWS* amino acid alignments at key spectral tuning sites to

characterize the residue identity and variation at positions previously shown to influence the

absorption properties of *LWS* and *RH1* opsins in vertebrates (Asenjo et al., 1994; Yokoyama,

2002a; Yokoyama & Radlwimmer, 2001a). We then compared the amino acid variants at these

sites with those reported for amphibians in the recent comprehensive analysis by (Schott et al.,

2024).

3. RESULTS

3.1 ANCESTRAL CHARACTER STATE RECONSTRUCTION
441

Ancestral character reconstruction of eye musculature shows that the ancestral condition

for caecilians was the presence of all six eye muscles (Figure 1A). The most highly supported

scenario suggests a retention of all muscles across most lineages, except for at least six

independent losses of muscles. Notably, there was a loss from six to zero muscles,

independently, in the MRCA of *Boulengerula* and *Gymnophis* The preferred model was equal

rates (AIC 69.60), followed by ordered (AIC 80.46), and all rates different (AIC 114.98). The

transition rate for the equal rates model was 0.0012.

Reconstruction of eye exposure shows that the ancestral condition for Gymnophiona is

eye covered by skin (Figure 1B). From that ancestral condition, there have been five independent

gains of eye covered by bone in *Boulengerula*, *Caecilia gracilis*, *Gymnopis*, *Oscaecilia* and

Scolecomorphus. Note that the character is labile in the caecilids *C. gracilis* and *O. ochrocephala*

(See Supplementary Material section 4.3). The preferred model was equal rates (AIC 46.09),

followed by all rates different (AIC 46.37). The transition rate for the equal rates model was

0.00171.

FIGURE 1. Posterior probabilities at each ancestral node for eye exposure and eye

musculature condition. Probabilities were obtained from stochastic character mapping using

model weights to sample three (A) or two (B) different Mk trait evolution models. The vertical

bar indicates habitat type: black for terrestrial burrowers, white for aquatic and semi-aquatic

burrowers. See Supplementary Methods section 2.1 for details.

3.2 CAECILIAN RETINAL MORPHOLOGY

282 photomicrographs of retinae representing 8 of the 10 families of Gymnophiona were

examined (see Supplementary Table 2 for specimen information and Supplementary Data 5 for

all photomicrographs). A subset of images with clear, well preserved morphological characters

were selected to highlight retinal diversity across Gymnophiona as well as the finding of

scattered cone-like photoreceptors. (Figures 2 and 3). All specimens examined had organized

retinae containing all the cellular layers typical for visual vertebrates except *Boulengerula*

boulengeri and *Scolecomorphus uluguruensis,* but this may be a result of poor preservation,

particularly in *S. uluguruensis*. Species of the primarily aquatic family Typhlonectidae

(*Chthonerpeton indistinctum, Typhlonectes compressicauda*) possess spherical lenses, while

terrestrial species exhibit ellipsoid lenses (e.g., *Epicrionops* sp*., Dermophis mexicanus)* except in

Caecilia occidentalis and all species which have the orbit covered by bone (*Scolecomorphus*

kirkii, S. uluguruensis, Boulengerula boulengeri, Gymnopis multiplicata), where the lens is

amorphous. Quantitative measures of the retina varied considerably among these taxa (Table 1).

We found cone-like cells in 4 of 14 species (*Epicrionops* sp*., Dermophis mexicanus, Ichthyophis*

kohtaoensis, *Scolecomorphus kirkii)*, but note that the sections for most specimens did not

preserve the photoreceptor layer sufficiently to allow observation of individual photoreceptor

morphology (See Supplementary Data 5). Detailed descriptions of eye morphology for the

evaluated species are presented in Supplementary Material section 3.2.

FIGURE 2. Photomicrographs of retinae from five species of caecilians at low and medium

magnification. (A, B) *Epicrionops sp.* (C, D) *Ichthyophis kohtaoensis* (E, F) *Dermophis*

mexicanus (G, H) *Typhlonectes compressicauda* (I, J) *Scolecomorphus kirkii.* Scale bars are 100

µm (A, C, E, G, I) and 20 µm (B, D, F, H, J). Letter abbreviations in (B): GCL - Ganglion cell

layer; IPL - Inner Plexiform Layer; INL - Inner Nuclear Layer; OPL - Outer Plexiform Layer;

ONL - Outer Nuclear Layer; IS - Inner Rod Segment; OS - Outer Rod Segment; PE - Pigment

Epithelium.

Table 1. Summary of characters reported from sections shown in Figure 1.

Species	Inner Rod Segment Length (µm)	Outer Rod Segment Length (µm)	Outer Plexiform Layer width (µm)	Inner Plexiform Layer width (µm)	Number of nuclei rows in ONL	Number of nuclei rows in INL	Number of nuclei rows in GCL	Lens Shape
Epicrionops sp.	10	35–40	NA	40	2–3	3–4	1–2	ellipsoid
Ichthyophis kohtaoensis	5	8–14	2–4	7–10	1–2	2–3	1–2	ellipsoid
Dermophis mexicanus	10	30–35	NA	46	2–3	3–4	2–3	ellipsoid
Typhlonectes compressicauda	N/A (see text)	N/A (see text)	< 3	5–8	1–2	3–4	1-2	spherical

Scolecomorphus kirkii	8	15–20	2–5	12	2–3	2–3	1, discontiguous	amorphous

FIGURE 3. Photomicrographs demonstrating photoreceptor cell diversity. Black arrows

point to photoreceptors that exhibit characteristics typical of cone photoreceptors (bulbous

ellipsoids, tapered outer segments that are much thinner than their inner segments); blue arrow

indicates a photoreceptor reminiscent of a double rod. These are morphologically distinct from

the rods that predominate the caecilian retina. Beside each panel, we include diagrams and

images of cone photoreceptors and one double rod diagram that resemble the structures we

identified. (A) *Epicrionops sp.* (B) *Epicrionops sp.* (C) *Dermophis mexicanus* (D) *Ichthyophis*

kohtaoensis (E) *Scolecomorphus kirkii* (F) *Dermophis mexicanus;* Dissociation of the

photoreceptors from the pigment epithelium (PE) is a frequent artifact of preservation. Scale bars

are 5 µm (A, B, C, D, E) and 20 µm (F). The retinal cell drawings are from (Walls, 1942). The

blue diagrams were adapted from Donner and Yovanovich (2020). The grey images were

adapted from Isayama et al. (2014). Abbreviations for the diagram associated with (B): o - outer

segment; d - oil droplet; e - ellipsoid; m - myoid; l - external limiting membrane; n - nucleus; f -

foot-piece.

3.3 RETRIEVAL OF LWS FROM CAECILIANS

We retrieved *LWS* sequences from four species of Gymnophiona (*Rhinatrema bivittatum*,

XM_029609805.1; *Microcaecilia unicolor,* XM_030192059.1; *Geotrypetes seraphini,*

XM_033922369.1; *Ichthyophis kohtaoensis*, GCA_033557465.1). The sequences are publicly

available in GenBank from Torres-Sanchez et al. (2019) and early results from the Vertebrate

Genome Project ([https://vertebrategenomesproject.org/]; (Haussler et al., 2009)). Previous

attempts to detect *LWS* were likely unsuccessful (Mohun et al., 2010) due to primer mismatch

(see Supplementary Table 5 in section 3.3 of the Supplementary Material).

FIGURE 4. (A) Maximum wavelength absorbance values (λ_{max}) for *LWS* and *RH1* estimated

using a regression-based machine learning model (ML; solid markers; Frazer et al., 2024) or

microspectrophotometry (MSP; hollow markers; Mohun et al., 2010) for five caecilian species.

Because *LWS* was previously thought to be absent in caecilians, no MSP data is available for this

opsin. The color scale at the top of the figure represents the visible light spectrum for the range

of estimated absorbance values (470–570 nm). The image corresponds to *Microcaecilia albiceps*.

(B) Linear regression between ML and MSP data for *LWS* (Pearson's correlation test: $r = 0.13$,

$P = 0.66$) and *RH1* ($r = 0.38$, $P = 0.09$). The images show three caecilian species included in

our study: *C. orientalis*, *M. albiceps* (© Santiago Ron; BIOWEB, https://bioweb.bio), and *G.*

seraphini (© Brian Freiermuth, with permission).

3.4 FURTHER VALIDATION OF A CAECILIAN* LWS *AND ASSESSMENT OF ITS

POTENTIAL FUNCTIONALITY

***3.4.1 Transcriptomics and presence of* LWS.** In total, we obtained 131,623,768 reads

from *C. orientalis* eye tissue (PRJNA1181370). The reads were assembled into a reference eye

transcriptome containing 175,181 transcripts. For *RH1*, one transcript was assembled, while for

LWS, two were assembled. Compared to LWS_v1, LWS_v2 is missing exon 5 which encodes

transmembrane domains VI and VII as well as the C-terminal region, based on comparison with

bovine rhodopsin. We found that *RH1* was expressed at much higher levels (17.53 TPM)

compared to LWS_v1 (0.22 TPM) or LWS_v2 (0.02 TPM; 80.3X and 818.5X, respectively). We

note that, given our expression data is based on a transcriptome of the whole eye, it does not

necessarily reflect mRNA expression exclusive to the retina, though significant expression of

LWS and *RH1* is not expected from extraretinal tissues of the eye. No sequences for *SWS1* or

SWS2 were assembled.

3.4.2 PCR. We obtained 568 bp of *LWS* sequences from 11 species of caecilians. After

aligning with *C. orientalis*, all sequences had open reading frames that lacked stop codons

(Supplementary Data 6).

***3.4.3 λ_{max} of* LWS *and* RH1.** The estimated λ_{max} for the complete *LWS* gene sequences of

five caecilian species (*G. seraphini*, *I. kohtaoensis*, *M. unicolor*, *R. bivittatum*, and *C. orientalis*)

ranged from 508.57 to 550.58 nm. The shortest predicted absorbance value corresponds to

*LWS*_v2 from *C. orientalis*. The predicted λ_{max}values for *RH1* in caecilians, using the complete

gene sequences, ranged from 492.07 to 513.54 nm (Figure 4; Supplementary Table 4).

We estimated absorbance values for *LWS* and *RH1* in 39 additional vertebrate species,

with *LWS* values ranging from 531.39 nm in the lungfish *Protopterus annectens* to 565.3 nm in

the chicken *Gallus gallus*, and *RH1* values ranging from 493.72 nm in the lizard *Pogona*

vitticeps to 513.54 nm in the frog *Scaphiopus couchii* (Supplementary Table 4).

Of the 44 species we reviewed, microspectrophotometry (MSP) data were available for

14 *LWS* sequences and 21 *RH1* sequences, including for two caecilian species. Our results

suggest a weak relationship between the λ_{max} values estimated using machine learning and

experimental MSP data from the literature, particularly for *LWS* (Pearson's correlation: $\square = 0.13$,

$\square = 0.66$). Although not significant, the correlation is stronger for *RH1* ($\square = 0.38$, $\square = 0.09$;

Figure 4B). Additionally, the machine-learning estimates showed greater discrepancies from

MSP data for *LWS* (mean absolute error: 26.4 nm) compared to *RH1* (2.8 nm). For the two

caecilian species with *RH1* MSP data, the difference between MSP and machine learning data

was ~3 nm (Figure 4A; Supplementary Table 4); MSP data for *LWS* was not available to

compare.

FIGURE 5. Rod and cone phototransduction cascade components and expression levels in

caecilian eye tissue. (A) In blue, schematic of the rod phototransduction cascade, showing the

key proteins involved: the rod opsin *RH1*, the heterotrimeric transducin complex (*GNAT1*,

GNB1, *GNGT1*), phosphodiesterase components (*PDE6A*, *PDE6B*, *PDE6G*), and the cyclic

nucleotide-gated channel subunits (*CNGA1*, *CNGB1*). (B) In red, schematic of the cone

phototransduction cascade, highlighting the cone opsin *LWS* and its associated components:

transducin complex (*GNAT2*, *GNB3*, *GNGT2*), phosphodiesterases (*PDE6C*, *PDE6H*), and cyclic

nucleotide-gated channel subunits (*CNGA3*, *CNGB3*). (C) Bar plots showing transcript

expression levels (transcripts per million, TPM) for the rod- and cone-associated

phototransduction genes that are present in the eye transcriptome of *Caecilia orientalis.* Blue

bars indicate rod-associated genes; red bars indicate cone-associated genes. White shapes in (A)

and (B) indicate genes not detected in the genomes or transcriptome, while light blue and light

red indicate genes present in at least one, but not all, of the assessed species (see Table 2).

578   *3.4.4 Assessing phototransduction cascade genes to infer* LWS *expression*

location. Our results indicate that most components of the rod phototransduction pathway are

intact in caecilian genomes and are expressed in the eye transcriptome of *C. orientalis* (with only

1 of 8 genes absent in all species; Figure 5A, Table 2). In contrast, the cone pathway displays a

more mosaic pattern, with 2 of 7 genes absent from all species and 3 others missing in one or

more species (Figure 5B, Table 2).

Most genes are represented by two or more isoforms, though these may be assembly artifacts.

We provide alignments including all detected transcriptomic isoforms in Supplementary Data 7.

Within the rod phototransduction pathway, *PDE6B*—a subunit of the phosphodiesterase

complex—is the only gene absent from all available caecilian genomes and from the salamander

P. waltl, suggesting a possible gene loss. A truncated *PDE6B*-like sequence (127 amino acids;

typical length ~890) was detected in the *C. orientalis* transcriptome (Figure 5A, Table 2), sharing

26.8% similarity with *C. orientalis PDE6A*. Due to its short length and absence in other

caecilians, this annotation remains uncertain. For comparison, we included the putative *PDE6B*

sequence in both *PDE6A* and *PDE6B* alignments (Supplementary Data 7). Although the

remaining rod genes were present in caecilians, several exhibited sequence irregularities (Table

2), which are described in detail in Supplementary Material Results section 3.4.4.

In contrast, the cone phototransduction pathway shows more instances of gene loss and a

mosaic pattern of gene presence and absence in caecilians (Figure 5B, Table 2). The genes

GNGT2 and *CNGB3*, which encode components of the cone transducin complex and the cyclic

nucleotide-gated (CNG) channels respectively, are entirely absent from all available genomes

and transcriptomes of Gymnophiona. Meanwhile *GNB3* and *CNGA3*, which are also part of the

transducin complex and CNG channels, are the only cone phototransduction genes consistently

present across all caecilian species, though both exhibit irregularities in *M. unicolor* and *C.*

orientalis, respectively (Table 2). Additional details on cone phototransduction gene

presence/absence and sequence irregularities are provided in Supplementary Material Results

section 3.4.4.

In terms of expression levels of rod- versus cone-specific phototransduction genes in *C.*

orientalis, the pattern observed mirrors that of the opsins, in that genes associated with the rod

phototransduction pathway are expressed at higher levels than those associated with the cone

phototransduction pathway (Figure 5C).

Table 2. Presence and characteristics of rod and cone phototransduction cascade

genes in four caecilian species. Symbols and cell colors indicate detection (+; black cell),

absence (-; white cell), or presence with sequence irregularities (e.g., early stop codon, shorter or

longer sequence, ambiguous sequence; grey cell).

Protein complex	Gene Symbol	Cell type	*R. bivittatum*	*M. unicolor*	*G. seraphini*	*C. orientalis* transcript
Transducin	GNAT1	Rod	Early stop codon	+	+	+
	GNB1		+	+	+	+
	GNGT1		+	+	+	+
	GNAT2	Cone	+	-	-	-
	GNB3		+	Variable sequence	+	+
	GNGT2		-	-	-	-
Phosphodiesterase	PDE6A	Rod	+	+	+	Shorter sequence

	Gene						
	PDE6B		-	-	-	Ambiguous sequence
	PDE6G		+	+	+	+
	PDE6C	Cone	+	-	+	Shorter sequence
	PDE6H		+	-	-	Longer sequence
Cyclic Nucleotide Gated Channel	*CNGA1*	Rod	+	+	+	+
	CNGB1		+	+	+	+
	CNGA3	Cone	+	+	+	Shorter sequence
	CNGB3		-	-	-	-

3.5 SELECTION ANALYSES

Results from the complete and incomplete *LWS* analyses were congruent (Supplementary

Material File S1 and S2). Thus, we only present the complete dataset. Results from the

incomplete dataset are provided in Supplementary Material Section 3.5.1. We initially tested

whether purifying selection better described the data than neutral evolution, which would be

consistent with long-term maintenance of *LWS*. In this test, the unconstrained M0 model was

favored over the constrained M0 model (null hypothesis), in which omega was fixed at 1

(simulating neutral selection), for both *LWS* (likelihood ratio (LR) = 2687.642, P << 0.0001) and

RH1 (LR = 2244.112, P <<0.0001), rejecting the null hypothesis. Additionally, we found that the

M3 model was a better fit than M0 for both genes (*RH1*: LR = 615.675, P < 0.0001; *LWS*: LR =

825.605, P < 0.0001; File S2), suggesting significant ω variation across sites. To test for

positively selected sites, we compared the fit of the M8 model, which allows for sites with ω > 1,

to that of M7 and M8a. For *RH1*, the M8 model was a better fit than M7 (LR = 40.181, P <

0.0001; File S2) and M8a (LR = 5.978, P = 0.015; File S2). Two sites were inferred to be under

positive selection with a BEB posterior probability >90% (positions 213 and 217; bovine *RH1*

numbering), while one site was close to but did not meet the significance threshold (position

107). Consistently, BUSTED found nearly significant evidence for gene-wide episodic

diversifying selection of *RH1* in caecilians (P = 0.059) and identified two sites with evidence

ratios (ER) ≥ 100 (positions 107 and 213, File S1), aligning with CODEML.

In contrast, for *LWS*, M8 was a better fit than M7 (LR = 44.241, P < 0.0001; File S2) but

not M8a (LR = 0.0, P = 1.0; File S2), indicating only a weak signal for positive selection. Only

one site exceeded the 90% BEB significance threshold (site 66; 49 in bovine *RH1*). Meanwhile,

BUSTED did not detect gene-wide episodic diversifying selection in caecilian *LWS* sequences (P

= 0.081) and found no sites under positive selection (File S1).

The FEL and FUBAR analyses can detect signals of positively selected sites across the

entire phylogeny. However, FUBAR did not identify any sites for *LWS* or *RH1*. On the other

hand, FEL identified one site under diversifying positive selection in *LWS* when the stem

caecilian clade was used as foreground (site 109; 92 in bovine *RH1;* P = 0.04), but not across the

entire phylogeny. In contrast, for *RH1*, FEL identified one site under positive selection across the

entire phylogeny (119; P = 0.05) and a different site when the stem caecilian clade was analyzed

as foreground (292; P = 0.04). Interestingly, FEL results showed that 272 out of 368 codon sites

(when no foreground was identified) and 175 out of 368 codon sites (when caecilians were

defined as the test group) in *LWS* were under purifying selection at p ≤ 0.05. This pattern of

purifying selection is similar to that observed in the functional *RH1*, where 247 out of 354 codon

sites (no foreground) and 165 out of 354 codon sites (caecilians foreground) were under

purifying selection at the same significance threshold. These findings suggest that selection is

acting to preserve the integrity of visual opsin gene sequences and maintain their function over

evolutionary time.

Our analysis using RELAX revealed an overall relaxation of selective constraint in *LWS*

but not *RH1* when crown Gymnophiona and its stem branch were designated as the foreground

(*LWS:* K = 0.74, LR = 15.47, P < 0.0001; *RH1*: K = 1.37, LR = 1.67, P = 0.196; Supplementary

Figure 1 and File S1). This pattern was further supported by CODEML. Specifically, both clade

model C (CmC vs M2a_rel: LRT = 8.507, P = 0.004; File S2) and clade model D (CmD vs. M3:

LRT = 11.055, P = 0.0009; File S2) were supported over other models, providing significant

evidence for shifts in selection pressures, with some sites in *LWS* showing a transition from

relatively stronger to weaker purifying selection within Gymnophiona + stem. For *RH1*,

however, neither the CmC nor CmD models were favored over the alternative models (CmC vs

M2a_rel: LRT = 0.508, P = 0.476; CmD vs M3: LRT = 3.053, P = 0.08), suggesting that

selective pressure on *RH1* has not undergone significant evolutionary shifts in caecilians (File

S2).

4. DISCUSSION

4.1 Evolution of caecilian eyes. Caecilians (Gymnophiona) are an ancient order of

amphibians whose ancestor began transitioning to a fossorial lifestyle more than 250 million

years ago (Kligman et al., 2023; Santos et al., 2020). Caecilians have undergone a degeneration

of visual organs associated with an increasingly fossorial lifestyle (Himstedt, 1995; Kamei,

Mauro, et al., 2012; Taylor, 1968), including losing two of three cone opsin genes (Lin et al.,

2024; Mohun et al., 2010; Musser & Arendt, 2017). As evidenced by our ancestral state

reconstructions of eye musculature and eye exposure (Figure 1, Supplementary Material

Methods section 2.1), patterns in trait loss are not necessarily correlated with species ecology or

phylogeny. For example, *Dermophis mexicanus* and *Gymnopis multiplicata* are closely related

and ecologically similar members of Dermophiidae from Central America: they are viviparous,

live as litter burrowers in shallow soil, and emerge at dawn and dusk, especially in light rains, to

forage (MHW pers. obs.). *Dermophis mexicanus* have relatively well-developed eyes in open

sockets, covered by skin, with all six extrinsic muscles, a cell-rich, organized retina, and a

crystalline lens that retains some cells peripherally. In contrast, *Gymnopis multiplicata* has eyes

in sockets covered by bone as well as skin, no extrinsic muscles, and an amorphous cellular lens

that is usually flattened against the cornea and retina (Wake, 1985). In another example, the

covering of the eye socket by bone is a homoplastic character that occurs in members of several

families: Caeciliidae (*Oscaecilia* and *Caecilia*), Dermophiidae (*Gymnopis*, *Dermophis*),

Herpelidae (*Boulengerula*), and Scolecomorphidae (*Scolecomorphus*, *Crotaphatrema*).

Scolecomorphus has an exceptional morphology: the eyes in at least three species are not in the

bone-covered sockets but embedded in the tentacle (De Jager, 1940; Taylor, 1968; Wake, 1985)

see also Supplementary Material section 4.3 (De Jager, 1940; Taylor, 1968; Wake, 1985). Given

what appears to be a general association between a bone-covered socket and reduced eye

musculature (Figure 1), it is tempting to consider the covered eye as a factor driving further

degeneration of the visual apparatus in these taxa (Wake, 1985), yet the morphological patterns

are not exactly similar, with different elements being more reduced or lost among the species in

the set, including among *Scolecomorphus* species. These examples illustrate the mosaic loss of

visual features across the history of caecilians, likely because of the relaxed selection that

appears to be a feature of the clade.

4.2 Expression of **LWS***, phototransduction cascade genes, and morphology of*

caecilian retinae. For such a unique and ancient vertebrate order, caecilians are woefully

understudied, and their genomes are no exception (Torres-Sánchez et al., 2019). Prior to work by

Lin et al (2024), caecilian eyes were thought to have rod-based retinae that only contain

rhodopsin (Mohun et al., 2010; Wake, 1985). Here we report full-length transcriptomic *LWS* and

RH1 sequences from one additional caecilian family (Caecilidae), as well as partial *LWS*

sequences from four additional families (Caecilidae, Herpelidae, Typhlonectidae, and

Scolecomorphidae), expanding coverage to eight of the ten families in the Order Gymnophiona.

These findings indicate that *LWS* has likely been maintained across all extant lineages, despite

varying degrees of fossoriality and eye reduction.

Given that all these *LWS* sequences have a translatable open reading frame and a stop

codon position homologous to other amphibian *LWS* sequences (Supplementary Data 3), and that

the gene is expressed in the eye of at least one caecilian species (Figure 5C), we hypothesize that

LWS is functionally expressed. The maintenance of *LWS* is further supported by our selection

analyses, which indicates that although there is a signal of relaxed evolution, a model of

purifying selection provides a significantly better fit than a fully neutral model—unsurprising

given the gene's retention over 200 million years of fossorial evolution. However, results from

two complementary approaches (CmC and CmD, and RELAX) indicate that *LWS* in caecilians is

experiencing a relaxation in the strength of purifying selection relative to other amphibians. This

pattern is not observed in the other retained opsin, *RH1*, and may suggest that the physiological

role of *LWS* is under less stringent selective constraint. We explore this possibility further in the

next section.

Our *LWS* expression data comes from *Caecilia orientalis*, a member of Caeciliidae

known for derived fossorial adaptations and relatively reduced eyes (*Taylor 1968*, n.d.). In this

species, *LWS* expression is significantly lower than *RH1*. There is little eye transcriptome data

available for amphibians but these results align with data from the nocturnal frog *Rana*

sphenocephala, where adult individuals express *RH1* at about 50 times the level of *LWS* (Schott

et al., 2022). Similarly, in some dim-light adapted (e.g., deep-water, nocturnal) fish species, *LWS*

expression relative to *RH1* can be very low (Fogg et al., 2022; Luehrmann et al., 2019).

However, some diurnal species also show relatively low *LWS* expression, such as the spotted

unicorn fish (*Naso brevirostris*) (Tettamanti et al., 2019). The relatively low expression of *LWS*

may be a result of a rod-biased (or rod-only) retina, a pattern observed across tetrapods, but

especially pronounced in nocturnal lineages or those that were adapted to nocturnality at some

point in their evolutionary history (de Busserolles et al., 2021; Kim et al., 2016; even at relatively

young evolutionary time scales, e.g., in colubrid snakes; Walls, 1942). The common ancestor of

Amphibia was likely nocturnal (Anderson & Wiens, 2017), and salamanders and anurans possess

a higher proportion of rods to cones in their retinae (Donner & Yovanovich, 2020; Fain, 1976;

Nilsson, 1964; Walls, 1942).

While the presence of intact and expressed *LWS* and *RH1* genes suggests the possibility

of a duplex retina, opsin expression alone does not confirm the maintenance of two functionally

distinct photoreceptor cell types. To further test this hypothesis, we pursued two complementary

lines of evidence: we examined the presence and expression of key rod and cone

phototransduction cascade genes and evaluated the morphology of caecilian retinas for structures

resembling cones. Our results revealed that most components of the rod phototransduction

pathway are present in caecilian genomes and expressed in the eye of *C. orientalis* (Figure 5A),

supporting the retention of rod function. In contrast, we observed a mosaic of gene loss in the

cone phototransduction pathway, consistent with relaxed selection on color vision in caecilians

(Figure 5B). The only rod-specific gene we were unable to detect in any caecilian genome was

PDE6B. This gene encodes one of the catalytic subunits of the PDE6 holoenzyme, which is

essential for signal transmission in rods as it hydrolyzes cGMP, leading to the closure of cyclic

nucleotide-gated channels and subsequent membrane hyperpolarization (Lagman et al., 2016).

Notably, *PDE6B*, *PDE6A*, and *PDE6G* have also been reported as lost in nocturnal geckos (Pinto

et al., 2019; Schott et al., 2019). Instead, these species are thought to utilize the cone protein

PDE6C to transduce rod signaling, supporting the hypothesis that ancestral cones were

transmuted into rods for scotopic vision (Kojima et al., 2021; Zhang et al., 2006). Similarly,

PDE6A is lost in birds and reptiles (Lagman et al., 2016), where this loss may be compensated

by the formation of a *PDE6B/B* homodimer rather than the *PDE6A*/*B* heterodimer typically used

by other vertebrates (Huang et al., 2004). As such, the absence of *PDE6B* in caecilian genomes

does not necessarily imply that the rod pathway is non-functional; instead, it raises the possibility

that an alternative catalytic subunit—potentially even *PDE6C*—may compensate for cGMP

hydrolysis in caecilian rods. It is important to note, however, that *M. unicolor* appears to lack

both *PDE6B* and *PDE6C*, though this observation could reflect incomplete sequencing or

assembly artifacts. Alternatively, if these absences are real, they raise further questions about

how cGMP hydrolysis is achieved in this species.

In contrast, cone phototransduction genes exhibit a mosaic pattern of presence and

absence across caecilian species. The only cone phototransduction gene consistently present in

all species analyzed is the Guanine nucleotide-binding protein β3 (*GNB3*) (5B, Table 3).

However, in addition to its expression in cone photoreceptors, *GNB3* has also been reported in

subsets of bipolar cells and rod photoreceptors across vertebrates, so it is possible that the

transcripts we detect originate from these other retinal cell types (Dhingra et al., 2012; Ritchey et

al., 2010). Many critical components are missing in all four surveyed species or detected only in

one or, at most, two caecilian species (Figure 5B, Table 3). The potential impact of these gene

absences in caecilians can be inferred from experimental studies in mice. For example, the

absence of *CNGA3* and *CNGB3* leads to cone degeneration, affecting both cell viability and

function (Michalakis et al., 2010). Moreover, although the effects of *GNGT2* loss have not been

directly studied, findings from *GNAT2* suggest that any disruption to the transducin complex

would likely impair cone phototransduction without necessarily compromising cone cell survival

(Chang et al., 2006; Ronning et al., 2018).

A similar patchy pattern of cone phototransduction gene losses has been reported in other

species that are either adapted to low-light conditions or with a history of subterranean lifestyles

during their evolution. For example, one species of armadillo (*Dasypus novemcinctus*) lacks

CNGB3 (Emerling & Springer, 2015) and one sloth species (*Choloepus hoffmanni*) and a group

of subterranean snakes lack *GNGT2* (Emerling & Springer, 2015; Gower et al., 2021). These

losses, along with the inactivation of other cone-specific genes, have been interpreted as

evidence for rod monochromacy in these groups. Given the extent of cone phototransduction

gene loss in Gymnophiona—with all species lacking at least two of the seven genes examined,

and in some cases, such as *M. unicolor*, lacking five genes—it is tempting to speculate that

caecilians may lack functional cones altogether, resulting in a rod-only retina. Taken together,

our findings suggest that while caecilians have retained *LWS*, their cone phototransduction

machinery may be at least partially degraded. This raises the possibility that *LWS* is expressed in

rod photoreceptors, and that some cone phototransduction components may have been co-opted

in rod cells to support *RH1* function and/or some rod phototransduction components may enable

LWS functionality.

Furthermore, existing literature and our morphological review of caecilian retinae do not

provide conclusive support for the presence of cone cells in any caecilian species. Cone cells

could be overlooked if they are inconspicuous and highly outnumbered by large rod cells, which

is consistent with our 80:1 *RH1* to *LWS* mRNA transcript ratio. Thus, observations of

photoreceptors with cone-like characteristics could be dismissed as anomalous variation,

damaged cells, or rod outer segments in a state of regeneration. We carefully inspected over 280

images in order to encounter a handful of putative cone-like cells. There is considerable

morphological variation among these cone-like photoreceptors, and, given the extreme variation

in cone morphology observed among vertebrates (even at relatively young evolutionary time

scales, e.g., in colubrid snakes; Walls, 1942); it is impossible to know what caecilian cones might

look like and how similar or different they might be among families, some of which likely

diverged nearly 200 MYA (Kligman et al., 2023). We acknowledge that given the distortions and

artifacts observed in retinal sections, the cone-like photoreceptors we highlighted may be rod

cells with anomalous or misleading morphologies as described above.

The expression of cone opsins in rod photoreceptors, or the co-expression of multiple

visual opsins within a single photoreceptor cell, is not uncommon. For example, the cone opsin

SWS2 is expressed in 'green rods' of frogs and salamanders (W. I. L. Davies et al., 2012), which,

in conjunction with 'red rods' expressing *RH1,* is thought to aid in nocturnal color vision

(Yovanovich et al., 2017). Additionally, Gower et al. (2021) found that *LWS* was co-expressed

with *RH1* in rod cells of *Anilios bicolor,* a fossorial scolecophidian snake with highly reduced

eyes. Thus, a similar phenomenon could be occurring in caecilians. Immunolabeling studies

employing antibodies against *LWS* or other proteins specific to cones and the cone

phototransduction pathway will be necessary to confirm their presence in the caecilian retina.

As far as we are aware, our study presents the first set of eye transcriptome data for any

caecilian, so we are not certain if the two *LWS* transcripts that we identified are artifacts of the

assembly process or represent alternatively spliced transcripts. The genome assemblies of *R.*

bivittatum, *M. unicolor*, and *G. seraphini* only contain one annotated transcript, but as far as we

are aware, the assembly process did not incorporate eye transcriptome data (Ovchinnikov et al.,

2023). Based on our alignments with bovine rhodopsin, exon 5 encodes portions of

transmembrane domains VI and VII and the C-terminal region of the opsin protein—regions that

play essential roles in chromophore binding (Palczewski et al., 2000) and spectral tunning .

Therefore, the absence of exon 5 in one of the *LWS* transcripts likely renders that isoform non-

functional, at least in its expected purpose.

4.3 Evidence for and implications of a putatively functional **LWS** *in caecilians.* When

analyzing the amino acid composition of known spectral tuning sites in available *LWS* and *RH1*

sequences for caecilians, we found that they are consistent with data reported for other

amphibians (Schott et al., 2024; Wan et al., 2023), containing at least one of the known amino

acid variants at each site (see Supplementary Table 6 in section 4.2). For example, as highlighted

by Lin et al. (2024) the specific combination of S164, H181, Y261, T269, and A292 found in the

LWS sequence of *R. bivitatum*, *G. seraphini*, *M. unicolor,* and *C. orientalis* has been shown to

generate a long-wavelength-sensitive photopigment with λ_{max} at ~560 nm in other vertebrates

(Asenjo et al., 1994; Hauzman et al., 2017; Yokoyama, 2002b; Yokoyama & Radlwimmer,

2001b). While this prediction is relatively close to our machine learning estimates (546.87–

550.58 nm; excluding *C. orientalis LWS*_v2), our estimates fall on the shorter end of the whole

range of absorption observed in the vertebrate *LWS* (~549–626 nm; Schott et al., 2024), similar

to the absorbance properties observed in nocturnal terrestrial species (Margetts et al., 2024;

Veilleux & Cummings, 2012). If accurate, these short absorbance values could enhance low-light

vision by improving the signal-to-noise ratio through reduced thermal noise susceptibility (Luo

et al., 2011).

It is important to note, however, that our machine learning (ML) model showed only a

weak correlation with MSP data for *LWS* ($\square$ = 0.13, $\square$ = 0.66), possibly because of an

underrepresentation of opsins in the ~600-nm range within the ML training dataset (Frazer et al.,

2024). Moreover, the model was trained using absorbance values based on 11-cis retinal (A1) as

the chromophore (Frazer et al., 2024), which may result in estimates that differ from MSP values

obtained using alternative chromophores such as A2. Additionally, variation in MSP data cannot

be ruled out; for instance, red cones from *Ambystoma tigrinum* show a 30–40 nm range in MSP

data (Makino & Dodd, 1996). This variation likely stems from the challenges of estimating

absorbance in single cone photoreceptors, especially for *LWS* in amphibians, as it is more

susceptible to noise and generally has smaller sample sizes compared to rod pigments like *RH1*

or *SWS2* (see Supplementary material in Schott et al., 2024).

LWS is present in a significant proportion of fossorial and low-light adapted taxa, such as

blind mole rats, cryptozooic Australian scincid lizards, and some deep-sea fish species (W. I. L.

Davies et al., 2012; Ford et al., 2024; Musilova et al., 2021). Most reported instances of *LWS*

gene loss or pseudogenization occur in fishes adapted to deep sea environments (Lin et al., 2017;

Musilova et al., 2021), where light in the long wavelength end of the visual spectrum is minimal

or absent. To our knowledge, pseudogenization of *LWS* in tetrapods has only been observed in

the subterranean naked mole-rat (*Heterocephalus glaber*), the Cape golden mole (*Chrysochloris*

asiatica), armadillos, and deep diving whales (Emerling & Springer, 2014; Meredith et al.,

2013). Dysfunction or complete loss of *LWS* is possible in the obligate subterranean Texas blind

salamander (*E. rathbuni*), given that Tovar et al. (2021) were unable to label any of the visual

opsins in the retina. However, failure to immunochemically label LWS in the retina is not

necessarily confirmation of gene loss or dysfunction. For example, in one population of the blind

cave salamander (*Proteus anguinus anguinus*), Kos et al. (2001) found the LWS protein in the

pineal gland but not the eye, while both tissues from other populations were successfully

immunolabeled for LWS (Kos et al., 2001). These data show that the *LWS* gene tends to persist

in ecological and evolutionary contexts under which other cone opsins are readily lost, strongly

suggesting that *LWS* retains some function even in species with highly reduced visual systems.

Additional evidence for the potential functionality of *LWS* in caecilians arises from our

selection analyses. With the shift to a fossorial lifestyle, caecilians underwent a simplification of

their visual systems, likely leading to relaxed selection on the molecular mechanisms underlying

vision, and eventually to the loss of some cone opsins and phototransduction genes. While the

loss of *SWS1* and *SWS2* (Lin et al., 2024; Mohun et al., 2010) might be the result of relaxed

selection, the same selective environment did not result in the loss of *LWS*, although the signal of

relaxation remains detectable in caecilian *LWS* sequences (see Results section 3.3). Importantly,

this relaxation in selective pressure did not result in an overabundance of deleterious mutations,

as the *LWS* sequences retrieved from caecilian genomes and transcriptome have intact open

reading frames. This contrasts with the other cone phototransduction genes that we examined,

where 5 of 7 appear lost in at least one species, several are truncated in the *C. orientalis*

transcriptome, and those that are variably present are always present in the earliest branching

lineage, *R. bivittatum* (Table 3). Given this long-term retention of *LWS* (over 240 million years;

Kligman et al., 2023), we hypothesize that some degree of purifying selection has preserved its

functionality, despite the relaxed constraints associated with reduced visual demands. One

possible function for *LWS* could be tied to behavioral ecology. Light between 520 and 610 nm is

most prominent in the forest understory while in more open woodland environments wavelengths

below 500 nm are more prevalent (Endler, 1993). Thus, slight variations in blue to red

wavelengths may provide caecilians information about canopy cover, habitat type, and exposure

to predators when they are at or near the surface in forested habitats. Furthermore, there are

significant shifts in light intensity within forests at long and short wavelengths in the transition

from daylight, to sunset, to twilight (Endler, 1993). Dichromatic vision is possible with only rods

(*RH1*) and red cones (*LWS*) under mesopic conditions (W. I. L. Davies et al., 2012; Hunt et al.,

2009) and could afford caecilians the spectral resolution necessary to discriminate between these

structural and temporal variables of their environment. Alternatively, If *LWS* is expressed in rods

in caecilians, similar information may be available to them at dawn/dusk/twilight and while

moving in the leaf litter or just under the surface. As caecilians appear to be primarily nocturnal

(Prakash et al., 2024) and most observations of surface activity occur at night, twilight or after

heavy rains, when flooded underground habitats force animals to the surface to breathe,

distinguishing light from different ends of the spectrum may be important for informing activity

patterns, entraining circadian rhythm, and survival. If expressed in cones, *LWS* may also provide

some visual information when caecilians find themselves at the surface under bright daylight

conditions that would render rods useless due to saturation. Light intensity and spectra may be

the only visual information available to caecilians. Therefore, retaining some spectral

discrimination may allow them to maximize the visual information available given the

constraints imposed by adaptation for head-first burrowing.

5. CONCLUSION

Although previous work suggested that the loss of *LWS* might be a synapomorphy of caecilians,

our results show that these amphibians retain the *LWS* gene along with most rod-specific

phototransduction genes and a subset of cone-pathway components. While it is possible that *LWS*

is expressed in rod photoreceptors, its precise function remains uncertain. Selection analyses

indicate that *LWS* is broadly under purifying selection, but with signs of relaxed constraint.

Although purifying selection appears strong enough to preserve gene integrity, this relaxation

may reflect reduced reliance on canonical cone-mediated vision. Additionally, we detected *LWS*

expression in the eye of one caecilian species but cannot confirm whether it is translated into a

functional protein. The maintenance of the open reading frame, however, suggests some function

of the mRNA, and/or of the protein, if it exists. We propose that *LWS* may function to transmit

light signals to other genes that regulate activity patterns (Boyette et al., 2024) or that it is used to

complement wavelength or luminance information obtained with *RH1*. Although traits are almost

never completely lost (Sadier et al., 2022), it is interesting that *LWS*, which is a terminal gene in

the visual cascade, has been maintained for such a long time under fossorial conditions. The

maintenance of *LWS* in caecilians and other dim-light-adapted animals begs the question of what

this opsin does and why it is still useful – i.e., why *LWS* and not the other opsins? These data

pose new questions regarding the selective pressures associated with dim-light or fossorial

lifestyles, the biology of caecilians, including the presence of bright coloration and above-ground

behavior in some species, as well as present an interesting example of the mosaic nature of trait

loss.

References

Anderson, S. R., & Wiens, J. J. (2017). Out of the dark: 350 million years of conservatism and evolution in diel activity patterns in vertebrates. *Evolution; International Journal of Organic Evolution*, *71*(8), 1944–1959.

Andrews, S. (2010). *FastQC: a quality control tool for high throughput sequence data.*

Asenjo, A. B., Rim, J., & Oprian, D. D. (1994). Molecular determinants of human red/green color discrimination. *Neuron*, *12*(5), 1131–1138.

Bielawski, J. P., & Yang, Z. (2004). A maximum likelihood method for detecting functional divergence at individual codon sites, with application to gene family evolution. *Journal of Molecular Evolution*, *59*(1), 121–132.

Bouckaert, R., Vaughan, T. G., Barido-Sottani, J., Duchêne, S., Fourment, M., Gavryushkina, A., Heled, J., Jones, G., Kühnert, D., De Maio, N., Matschiner, M., Mendes, F. K., Müller, N. F., Ogilvie, H. A., du Plessis, L., Popinga, A., Rambaut, A., Rasmussen, D., Siveroni, I., … Drummond, A. J. (2019). BEAST 2.5: An advanced software platform for Bayesian evolutionary analysis. *PLoS Computational Biology*, *15*(4), e1006650.

Bowmaker, J. K. (2008). Evolution of vertebrate visual pigments. *Vision Research*, *48*(20), 2022–2041.

Boyette, J. L., Bell, R. C., Fujita, M. K., Thomas, K. N., Streicher, J. W., Gower, D. J., & Schott, R. K. (2024). Diversity and molecular evolution of nonvisual opsin genes across environmental, developmental, and morphological adaptations in frogs. *Molecular Biology and Evolution*, *41*(6). https://doi.org/10.1093/molbev/msae090

Carmona, F. D., Glösmann, M., Ou, J., Jiménez, R., & Collinson, J. M. (2009). Retinal

development and function in a "blind" mole. *Proceedings of the Royal Society B: Biological Sciences*, *277*(1687), 1513–1522.

Carmona, F. D., Ou, J., Jiménez, R., & Collinson, J. M. (2010). Development of the cornea of true moles (Talpidae): morphogenesis and expression of PAX6 and cytokeratins. *Journal of Anatomy*, *217*(5), 488–500.

Catania, K. C. (1999). A nose that looks like a hand and acts like an eye: the unusual mechanosensory system of the star-nosed mole. *Journal of Comparative Physiology. A, Sensory, Neural, and Behavioral Physiology*, *185*(4), 367–372.

Chang, B., Dacey, M. S., Hawes, N. L., Hitchcock, P. F., Milam, A. H., Atmaca-Sonmez, P., Nusinowitz, S., & Heckenlively, J. R. (2006). Cone photoreceptor function loss-3, a novel mouse model of achromatopsia due to a mutation in Gnat2. *Investigative Ophthalmology & Visual Science*, *47*(11), 5017–5021.

Chen, Y.-A., Lin, C.-C., Wang, C.-D., Wu, H.-B., & Hwang, P.-I. (2007). An optimized procedure greatly improves EST vector contamination removal. *BMC Genomics*, *8*(1), 416.

Chernomor, O., von Haeseler, A., & Minh, B. Q. (2016). Terrace Aware Data Structure for Phylogenomic Inference from Supermatrices. *Systematic Biology*, *65*(6), 997–1008.

Davies, W. I. L., Collin, S. P., & Hunt, D. M. (2012). Molecular ecology and adaptation of visual photopigments in craniates. *Molecular Ecology*, *21*(13), 3121–3158.

Davies, W. L. (2011). Adaptive gene loss in vertebrates: Photosensitivity as a model case. In *Encyclopedia of Life Sciences*. John Wiley & Sons, Ltd. https://doi.org/10.1002/9780470015902.a0022890

de Busserolles, F., Cortesi, F., Fogg, L., Stieb, S. M., Luehrmann, M., & Marshall, N. J. (2021). The visual ecology of Holocentridae, a nocturnal coral reef fish family with a deep-sea-like

multibank retina. *The Journal of Experimental Biology*, *224*(Pt 1), jeb233098.

De Jager, E. F. J. (1940). *Cranial Morphology of the Gymnophiona*. University of Stellenbosch.

Delport, W., Poon, A. F. Y., Frost, S. D. W., & Kosakovsky Pond, S. L. (2010). Datamonkey 2010: a suite of phylogenetic analysis tools for evolutionary biology. *Bioinformatics* , *26*(19), 2455–2457.

Dhingra, A., Ramakrishnan, H., Neinstein, A., Fina, M. E., Xu, Y., Li, J., Chung, D. C., Lyubarsky, A., & Vardi, N. (2012). Gβ3 is required for normal light ON responses and synaptic maintenance. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *32*(33), 11343–11355.

Donner, K., & Yovanovich, C. A. M. (2020). A frog's eye view: Foundational revelations and future promises. *Seminars in Cell & Developmental Biology*, *106*, 72–85.

Dünker, N. (1998). Colocalization of serotonin and GABA in retinal neurons of Ichthyophis kohtaoensis (Amphibia; Gymnophiona). *Anatomy and Embryology*, *197*(1), 69–75.

Edgar, R. C. (2004). MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, *32*(5), 1792–1797.

Emam, A., Yoffe, M., Cardona, H., & Soares, D. (2020). Retinal morphology in Astyanax mexicanus during eye degeneration. *The Journal of Comparative Neurology*, *528*(9), 1523–1534.

Emerling, C. A., & Springer, M. S. (2014). Eyes underground: regression of visual protein networks in subterranean mammals. *Molecular Phylogenetics and Evolution*, *78*, 260–270.

Emerling, C. A., & Springer, M. S. (2015). Genomic evidence for rod monochromacy in sloths and armadillos suggests early subterranean history for Xenarthra. *Proceedings. Biological Sciences*, *282*(1800), 20142192.

Emerson, S. B. (1976). Burrowing in frogs. *Journal of Morphology*, *149*(4), 437–458.

Endler, J. A. (1993). The color of light in forests and its implications. *Ecological Monographs*, *63*(1), 1–27.

Espinasa, L., & Jeffery, W. R. (2006). Conservation of retinal circadian rhythms during cavefish eye degeneration. *Evolution & Development*, *8*(1), 16–22.

Fain, G. L. (1976). Sensitivity of toad rods: Dependence on wave-length and background illumination. *The Journal of Physiology*, *261*(1), 71–101.

Fogg, L. G., Cortesi, F., Lecchini, D., Gache, C., Marshall, N. J., & de Busserolles, F. (2022). Development of dim-light vision in the nocturnal reef fish family Holocentridae. I: Retinal gene expression. *The Journal of Experimental Biology*, *225*(17), jeb244513.

Ford, M. J. R., Ludington, A. J., Bradford, T., Sanders, K. L., Hutchinson, M. N., & Simões, B. F. (2024). Retention of the full visual opsin repertoire in Australia's cryptozoic lizards. In *bioRxiv* (p. 2024.02.19.581082). https://doi.org/10.1101/2024.02.19.581082

Foureaux, G., Egami, M. I., Jared, C., Antoniazzi, M. M., Gutierre, R. C., & Smith, R. L. (2010). Rudimentary eyes of squamate fossorial reptiles (amphisbaenia and serpentes). *Anatomical Record (Hoboken, N.J.: 2007)*, *293*(2), 351–357.

Frazer, S. A., Baghbanzadeh, M., Rahnavard, A., Crandall, K. A., & Oakley, T. H. (2024). Discovering genotype-phenotype relationships with machine learning and the Visual Physiology Opsin Database (VPOD). *GigaScience*, *13*, giae073.

Glösmann, M., Steiner, M., Peichl, L., & Ahnelt, P. K. (2008). Cone photoreceptors and potential UV vision in a subterranean insectivore, the European mole. *Journal of Vision*, *8*(4), 23.1–12.

Gower, D. J., Fleming, J. F., Pisani, D., Vonk, F. J., Kerkkamp, H. M. I., Peichl, L., Meimann,

S., Casewell, N. R., Henkel, C. V., Richardson, M. K., Sanders, K. L., & Simões, B. F. (2021). Eye-transcriptome and genome-wide sequencing for Scolecophidia: Implications for inferring the visual system of the ancestral snake. *Genome Biology and Evolution*, *13*(12). https://doi.org/10.1093/gbe/evab253

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, A., Rhind, N., di Palma, F., Birren, B. W., Nusbaum, C., Lindblad-Toh, K., … Regev, A. (2011). Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology*, *29*(7), 644–652.

Haas, B. J. (n.d.). *TransDecoder*. TransDecoder v2.026. Retrieved 2021, from https://github.com/TransDecoder/TransDecoder

Haussler, D., O'Brien, S. J., Ryder, O. A., Keith Barker, F., Clamp, M., Crawford, A. J., Hanner, R., Hanotte, O., Johnson, W. E., McGuire, J. A., Miller, W., Murphy, R. W., Murphy, W. J., Sheldon, F. H., Sinervo, B., Venkatesh, B., Wiley, E. O., Allendorf, F. W., Amato, G., … Turner, S. (2009). Genome 10K: A Proposal to Obtain Whole-Genome Sequence for 10 000 Vertebrate Species. *The Journal of Heredity*, *100*(6), 659–674.

Hauzman, E., Bonci, D. M. O., Suárez-Villota, E. Y., Neitz, M., & Ventura, D. F. (2017). Daily activity patterns influence retinal morphology, signatures of selection, and spectral tuning of opsin genes in colubrid snakes. *BMC Evolutionary Biology*, *17*(1), 249.

Himstedt, W. (1995). Structure and function of the eyes in the caecilian *Ichthyophis kohtaoensis (Amphibia, Gymnophiona)*. *Zoology-Analysis of Complex Systems*, *99*(1), 81–94.

Holmberg, B. J. (2022). Ophthalmology of Eulipotyphla: Moles, Shrews, Hedgehogs, and Relatives. In F. Montiani-Ferreira, B. A. Moore, & G. Ben-Shlomo (Eds.), *Wild and Exotic*

Animal Ophthalmology: Volume 2: Mammals (pp. 355–366). Springer International Publishing.

Huang, D., Hinds, T. R., Martinez, S. E., Doneanu, C., & Beavo, J. A. (2004). Molecular determinants of cGMP binding to chicken cone photoreceptor phosphodiesterase. *The Journal of Biological Chemistry*, *279*(46), 48143–48151.

Hunt, D., Carvalho, L. S., Cowing, J. A., & Davies, W. L. (2009). Evolution and spectral tuning of visual pigments in birds and mammals. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *364*(1531), 2941–2955.

Isayama, T., Chen, Y., Kono, M., Fabre, E., Slavsky, M., DeGrip, W. J., Ma, J.-X., Crouch, R. K., & Makino, C. L. (2014). Coexpression of three opsins in cone photoreceptors of the salamander *Ambystoma tigrinum*: Pigment mixtures in salamander photoreceptors. *The Journal of Comparative Neurology*, *522*(10), 2249–2265.

Jacobs, G. H. (2013). Losses of functional opsin genes, short-wavelength cone photopigments, and color vision--a significant trend in the evolution of mammalian vision. *Visual Neuroscience*, *30*(1-2), 39–53.

Jenkins, P. a., & Walsh, D. M. (1993). An Early Jurassic caecilian with limbs. *Nature*, *365*, 246–250.

Jetz, W., & Pyron, R. A. (2018). The interplay of past diversification and evolutionary isolation with present imperilment across the amphibian tree of life. *Nature Ecology & Evolution*, *2*(5), 850–858.

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R., & Hunter, S. (2014). InterProScan 5: genome-scale protein function

classification. *Bioinformatics (Oxford, England)*, *30*(9), 1236–1240.

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., & Jermiin, L. S. (2017). ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods*, *14*(6), 587–589.

Kamei, R. G., Mauro, D. S., Gower, D. J., Van Bocxlaer, I., Sherratt, E., Thomas, A., Babu, S., Bossuyt, F., Wilkinson, M., & Biju, S. D. (2012). Discovery of a new family of amphibians from northeast India with ancient links to Africa. *Proceedings of the Royal Society B: Biological Sciences*, *279*, 2396–2401.

Kamei, R. G., San Mauro, D., Gower, D. J., Van Bocxlaer, I., Sherratt, E., Thomas, A., Babu, S., Bossuyt, F., Wilkinson, M., & Biju, S. D. (2012). Discovery of a new family of amphibians from northeast India with ancient links to Africa. *Proceedings. Biological Sciences*, *279*(1737), 2396–2401.

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4), 772–780.

Kim, J.-W., Yang, H.-J., Oel, A. P., Brooks, M. J., Jia, L., Plachetzki, D. C., Li, W., Allison, W. T., & Swaroop, A. (2016). Recruitment of rod photoreceptors from short-wavelength-sensitive cones during the evolution of nocturnal vision in mammals. *Developmental Cell*, *37*(6), 520–532.

Kligman, B. T., Gee, B. M., Marsh, A. D., Nesbitt, S. J., Smith, M. E., Parker, W. G., & Stocker, M. R. (2023). Triassic stem caecilian supports dissorophoid origin of living amphibians. *Nature*, *614*(7946), 102–107.

Kojima, K., Matsutani, Y., Yanagawa, M., Imamoto, Y., Yamano, Y., Wada, A., Shichida, Y., &

Yamashita, T. (2021). Evolutionary adaptation of visual pigments in geckos for their photic environment. *Science Advances*, *7*(40), eabj1316.

Kosakovsky Pond, S. L., & Frost, S. D. W. (2005). Not so different after all: a comparison of methods for detecting amino acid sites under selection. *Molecular Biology and Evolution*, *22*(5), 1208–1222.

Kos, M., Bulog, B., Szél, A., & Röhlich, P. (2001). Immunocytochemical demonstration of visual pigments in the degenerate retinal and pineal photoreceptors of the blind cave salamander (*Proteus anguinus*). *Cell and Tissue Research*, *303*(1), 15–25.

Lagman, D., Franzén, I. E., Eggert, J., Larhammar, D., & Abalo, X. M. (2016). Evolution and expression of the phosphodiesterase 6 genes unveils vertebrate novelty to control photosensitivity. *BMC Evolutionary Biology*, *16*(1), 124.

Lamb, T. D. (2013). Evolution of phototransduction, vertebrate photoreceptors and retina. *Progress in Retinal and Eye Research*, *36*(C), 52–119.

Liebman, P. A., & Entine, G. (1968). Visual pigments of frog and tadpole (Rana pipiens). *Vision Research*, *8*(7), 761–775.

Lin, J.-J., Wang, F.-Y., Chung, W.-Y., & Wang, T.-Y. (2024). The genomic evolution of visual opsin genes in amphibians. *Vision Research*, *222*, 108447.

Lin, J.-J., Wang, F.-Y., Li, W.-H., & Wang, T.-Y. (2017). The rises and falls of opsin genes in 59 ray-finned fish genomes and their implications for environmental adaptation. *Scientific Reports*, *7*(1), 15568.

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.

Luehrmann, M., Carleton, K. L., Cortesi, F., Cheney, K. L., & Marshall, N. J. (2019).

Cardinalfishes (Apogonidae) show visual system adaptations typical of nocturnally and diurnally active fish. *Molecular Ecology*, *28*(12), 3025–3041.

Luo, D.-G., Yue, W. W. S., Ala-Laurila, P., & Yau, K.-W. (2011). Activation of visual pigments by light and heat. *Science (New York, N.Y.)*, *332*(6035), 1307–1312.

Maciel, A. O., & Hoogmoed, M. S. (2011). Taxonomy and distribution of caecilian amphibians (Gymnophiona) of Brazilian Amazonia, with a key to their identification. *Zootaxa*, *2984*(1), 1.

Maddison, W. P., & Maddison, D. R. (2015). *Mesquite: a modular system for evolutionary analysis*. http://mesquiteproject.org

Makino, C. L., & Dodd, R. L. (1996). Multiple visual pigments in a photoreceptor of the salamander retina. *The Journal of General Physiology*. https://rupress.org/jgp/article-abstract/108/1/27/26499

Margetts, B. M., Stuart-Fox, D., & Franklin, A. M. (2024). Red vision in animals is broadly associated with lighting environment but not types of visual task. *Ecology and Evolution*, *14*(2), e10899.

Meredith, R. W., Gatesy, J., Emerling, C. A., York, V. M., & Springer, M. S. (2013). Rod monochromacy and the coevolution of cetacean retinal opsins. *PLoS Genetics*, *9*(4), e1003432.

Michalakis, S., Mühlfriedel, R., Tanimoto, N., Krishnamoorthy, V., Koch, S., Fischer, M. D., Becirovic, E., Bai, L., Huber, G., Beck, S. C., Fahl, E., Büning, H., Paquet-Durand, F., Zong, X., Gollisch, T., Biel, M., & Seeliger, M. W. (2010). Restoration of cone vision in the CNGA3-/- mouse model of congenital complete lack of cone photoreceptor function. *Molecular Therapy: The Journal of the American Society of Gene Therapy*, *18*(12), 2057–

2063.

Mohun, S. M., Davies, W. L., Bowmaker, J. K., Pisani, D., Himstedt, W., Gower, D. J., Hunt, D. M., & Wilkinson, M. (2010). Identification and characterization of visual pigments in caecilians (Amphibia: Gymnophiona), an order of limbless vertebrates with rudimentary eyes. *The Journal of Experimental Biology*, *213*(20), 3586–3592.

Mohun, S. M., & Wilkinson, M. (2015). The eye of the caecilian *Rhinatrema bivittatum* (Amphibia: Gymnophiona: Rhinatrematidae). *Acta Zoologica* , *96*(2), 147–153.

Murrell, B., Moola, S., Mabona, A., Weighill, T., Sheward, D., Kosakovsky Pond, S. L., & Scheffler, K. (2013). FUBAR: a fast, unconstrained bayesian approximation for inferring selection. *Molecular Biology and Evolution*, *30*(5), 1196–1205.

Musilova, Z., Salzburger, W., & Cortesi, F. (2021). The visual opsin gene repertoires of teleost fishes: Evolution, ecology, and function. *Annual Review of Cell and Developmental Biology*, *37*(1), 441–468.

Musser, J. M., & Arendt, D. (2017). Loss and gain of cone types in vertebrate ciliary photoreceptor evolution. *Developmental Biology*, *431*(1), 26–35.

Nevo, E. (1979). Adaptive Convergence and Divergence of Subterranean Mammals. *Annual Review of Ecology and Systematics*, *10*, 269–308.

Nilsson, S. E. (1964). An electron microscopic classification of the retinal receptors of the leopard frog (*Rana pipiens*). *Journal of Ultrastructure Research*, *10*(5-6), 390–416.

Nishikawa, K., Matsui, M., Yoshikawa, N., Khonsue, W., & Sanamxay, D. (2021). *Ichthyophis bannanicus Yang, 1984*, a junior subjective synonym of I. kohtaoensis Taylor, 1960 (Amphibia, Gymnophiona, Ichthyophiidae). *Alytes*, *38*(1-4), 1–17.

Norris, H., & Hughes, S. P. (1918). The cranial and anterior spinal nerves of the caecilian

amphibians. *Journal of Morphology*, *31*. https://doi.org/10.1002/jmor.1050310304

Ovchinnikov, V., Uliano-Silva, M., Wilkinson, M., Wood, J., Smith, M., Oliver, K., Sims, Y., Torrance, J., Suh, A., McCarthy, S. A., Durbin, R., & O'Connell, M. J. (2023). Caecilian genomes reveal the molecular basis of adaptation and convergent evolution of limblessness in snakes and caecilians. *Molecular Biology and Evolution*, *40*(5). https://doi.org/10.1093/molbev/msad102

Palczewski, K., Kumasaka, T., Hori, T., Behnke, C. A., Motoshima, H., Fox, B. A., Le Trong, I., Teller, D. C., Okada, T., Stenkamp, R. E., Yamamoto, M., & Miyano, M. (2000). Crystal structure of rhodopsin: A G protein-coupled receptor. *Science*, *289*(5480), 739–745.

Patro, R., Duggal, G., Love, M. I., Irizarry, R. A., & Kingsford, C. (2017). Salmon provides fast and bias-aware quantification of transcript expression. *Nature Methods*, *14*(4), 417–419.

Pinto, B. J., Nielsen, S. V., & Gamble, T. (2019). Transcriptomic data support a nocturnal bottleneck in the ancestor of gecko lizards. *Molecular Phylogenetics and Evolution*, *141*(106639), 106639.

Prakash, A., Gower, D. J., Vengot, R., & Kotharambath, R. (2024). Circadian rhythm and surface activity in soil-dwelling caecilians (Amphibia: Gymnophiona). *Scientific Reports*, *14*(1), 9950.

Revell, L. J. (2024). phytools 2.0: an updated R ecosystem for phylogenetic comparative methods (and other things). *PeerJ*, *12*, e16505.

Ritchey, E. R., Bongini, R. E., Code, K. A., Zelinka, C., Petersen-Jones, S., & Fischer, A. J. (2010). The pattern of expression of guanine nucleotide-binding protein beta3 in the retina is conserved across vertebrate species. *Neuroscience*, *169*(3), 1376–1391.

Ronning, K. E., Allina, G. P., Miller, E. B., Zawadzki, R. J., Pugh, E. N., Jr, Herrmann, R., &

Burns, M. E. (2018). Loss of cone function without degeneration in a novel Gnat2 knock-out mouse. *Experimental Eye Research*, *171*, 111–118.

Sadier, A., Sears, K. E., & Womack, M. (2022). Unraveling the heritage of lost traits. *Journal of Experimental Zoology. Part B, Molecular and Developmental Evolution*, *338*(1-2), 107–118.

San Mauro, D., Gower, D. J., Müller, H., Loader, S. P., Zardoya, R., Nussbaum, R. A., & Wilkinson, M. (2014). Life-history evolution and mitogenomic phylogeny of caecilian amphibians. *Molecular Phylogenetics and Evolution*, *73*, 177–189.

Santos, R. O., Laurin, M., & Zaher, H. (2020). A review of the fossil record of caecilians (Lissamphibia: Gymnophionomorpha) with comments on its use to calibrate molecular timetrees. *Biological Journal of the Linnean Society. Linnean Society of London*, *131*(4), 737–755.

Schott, R. K., Bell, R. C., Loew, E. R., Thomas, K. N., Gower, D. J., Streicher, J. W., & Fujita, M. K. (2022). Transcriptomic evidence for visual adaptation during the aquatic to terrestrial metamorphosis in leopard frogs. *BMC Biology*, *20*(1), 138.

Schott, R. K., Bhattacharyya, N., & Chang, B. S. W. (2019). Evolutionary signatures of photoreceptor transmutation in geckos reveal potential adaptation and convergence with snakes. *Evolution; International Journal of Organic Evolution*, *73*(9), 1958–1971.

Schott, R. K., Fujita, M. K., Streicher, J. W., Gower, D. J., Thomas, K. N., Loew, E. R., Bamba Kaya, A. G., Bittencourt-Silva, G. B., Guillherme Becker, C., Cisneros-Heredia, D., Clulow, S., Davila, M., Firneno, T. J., Jr, Haddad, C. F. B., Janssenswillen, S., Labisko, J., Maddock, S. T., Mahony, M., Martins, R. A., … Bell, R. C. (2024). Diversity and Evolution of Frog Visual Opsins: Spectral Tuning and Adaptation to Distinct Light Environments.

Molecular Biology and Evolution, *41*(4). https://doi.org/10.1093/molbev/msae049

Taylor, E. H. (1968). *The caecilians of the world*. University of Kansas Press.

Taylor, E. H. (1969). A new Panamanian caecilian. *The University of Kansas Science Bulletin*, *48*(12), 315–323.

Tettamanti, V., de Busserolles, F., Lecchini, D., Marshall, N. J., & Cortesi, F. (2019). Visual system development of the spotted unicornfish, *Naso brevirostris* (Acanthuridae). *The Journal of Experimental Biology*, *222*(Pt 24), jeb209916.

Torres-Sánchez, M., Creevey, C. J., Kornobis, E., Gower, D. J., Wilkinson, M., & San Mauro, D. (2019). Multi-tissue transcriptomes of caecilian amphibians highlight incomplete knowledge of vertebrate gene families. *DNA Research: An International Journal for Rapid Publication of Reports on Genes and Genomes*, *26*(1), 13–20.

Tovar, R. U., Cantu, V., Fremaux, B., Gonzalez, P., Jr, Spikes, A., & García, D. M. (2021). Comparative development and ocular histology between epigean and subterranean salamanders (Eurycea) from central Texas. *PeerJ*, *9*(e11840), e11840.

Veilleux, C. C., & Cummings, M. E. (2012). Nocturnal light environments and species ecology: implications for nocturnal color vision in forests. *The Journal of Experimental Biology*, *215*(Pt 23), 4085–4096.

Vidal□García, M., Byrne, P. G., Roberts, J. D., & Keogh, J. S. (2014). The role of phylogeny and ecology in shaping morphology in 21 genera and 127 species of Australo□Papuan myobatrachid frogs. *Journal of Evolutionary Biology*, *27*(1), 181–192.

Wake, M. H. (1985). The comparative morphology and evolution of the eyes of caecilians (Amphibia, Gymnophiona). *Zoomorphology*, *105*(5), 277–295.

Wake, M. H. (1993). Non-traditional characters in the assessment of caecilian phylogenetic

relationships. *Herpetological Monograph*, *7*, 42.

Walls, G. L. (1942). *The vertebrate eye and its adaptive radiation [by] Gordon Lynn Walls*. Cranbrook Institute of Science,.

Wan, Y. C., Navarrete Méndez, M. J., O'Connell, L. A., Uricchio, L. H., Roland, A.-B., Maan, M. E., Ron, S. R., Betancourth-Cundar, M., Pie, M. R., Howell, K. A., Richards-Zawacki, C. L., Cummings, M. E., Cannatella, D. C., Santos, J. C., & Tarvin, R. D. (2023). Selection on visual opsin genes in diurnal Neotropical frogs and loss of the SWS2 opsin in poison frogs. *Molecular Biology and Evolution*, *40*(10). https://doi.org/10.1093/molbev/msad206

Warrant, E. (2000). The eyes of deep-sea fishes and the changing nature of visual scenes with depth. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *355*(1401), 1155–1159.

Weadick, C. J., & Chang, B. S. W. (2012). An improved likelihood ratio test for detecting site-specific functional divergence among clades of protein-coding genes. *Molecular Biology and Evolution*, *29*(5), 1297–1300.

Wertheim, J. O., Murrell, B., Smith, M. D., Kosakovsky Pond, S. L., & Scheffler, K. (2015). RELAX: detecting relaxed selection in a phylogenetic framework. *Molecular Biology and Evolution*, *32*(3), 820–832.

Yang, Z. (2007). PAML 4: phylogenetic analysis by maximum likelihood. *Molecular Biology and Evolution*, *24*(8), 1586–1591.

Yokoyama, S. (2002a). Molecular evolution of color vision in vertebrates. *Gene*, *300*(1-2), 69–78.

Yokoyama, S. (2002b). Molecular evolution of color vision in vertebrates. *Gene 300*, 69–78.

Yokoyama, S., & Radlwimmer, F. (2001a). The molecular genetics and evolution of red and

green color vision in vertebrates. *Genetics*, *158*(4), 1697–1710.

Yokoyama, S., & Radlwimmer, F. B. (2001b). The molecular genetics and evolution of red and green color vision in vertebrates. *Genetics*, *158*, 1697–1710.

Yovanovich, C. A. M., Koskela, S. M., Nevala, N., Kondrashev, S. L., Kelber, A., & Donner, K. (2017). The dual rod system of amphibians supports colour discrimination at the absolute visual threshold. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, *372*(1717), 20160066.

Zhang, X., Wensel, T. G., & Yuan, C. (2006). Tokay gecko photoreceptors achieve rod-like physiology with cone-like proteins. *Photochemistry and Photobiology*, *82*(6), 1452–1460.

A. EYE MUSCULATURE

All muscles lost
Two muscles
Three muscles
Four muscles
Six muscles, *s.o.* attenuate
Six muscles

B. EYE EXPOSURE

Eye exposed
Eye under skin
Eye under bone

GCL
IPL
INL
OPL
ONL
IS
OS
PE