Use of massive DNA barcoding to monitor biodiversity: a test on forest soil macrofauna

Authors: Franck Jabot[a*], Gwenaëlle Auger[a], Pauline Bonnal[a], Mathilde Pizaine[a], Marilyn Roncoroni[a], Sandrine Revaillot[a], Julien Pottier[a]

Affiliation: [a]INRAE, VetAgro Sup, UMR Ecosystème Prairial, Université Clermont Auvergne, Clermont-Ferrand, France.

Abstract

Biodiversity monitoring primarily focuses on particular taxonomic groups for which identification expertise is widely available, such as vascular plants or birds. As a result, the response of many taxonomic groups to forest management is much less documented. With the advent of low-cost next generation barcoding approaches, it is now possible to envisage monitoring strategies based on molecular approaches, beyond metabarcoding/eDNA strategies that do not give access to species abundances and integrate species presence over largely unknown spatio-temporal scales. In this contribution, we demonstrate the use of massive DNA barcoding – also referred as megabarcoding – as a promising solution to overcome identification difficulties in demanding taxonomic groups. We performed a proof of concept study in a mountainous beech forest in the Massif Central, France. We sampled soil macrofauna at 25 sampling sites and managed to visually identify at the species level 130 out of the 1413 sampled individuals. Using megabarcoding on the 1283 remaining non-identified individuals, we managed to assign 1124 additional individuals to an operational taxonomic species at a competitive cost. We present the summary statistics of barcoding success in the different taxonomic groups encountered and in larvae versus adult individuals. We demonstrate that larvae individuals, which can hardly be visually identified at the species level, make a substantial contribution to overall macrofauna diversity. We finally showcase how this megabarcoding approach provides a concrete avenue for forest biodiversity monitoring, by assessing the cost and labour intensity of this approach.

32 **Keywords**

Biodiversity, monitoring, DNA barcoding, soil fauna, megabarcoding.

1. Introduction

Forest biodiversity monitoring mainly focuses on a limited number of taxonomic groups, among which plants, birds and beetles have attracted much attention (Paillet et al. 2010). Rationale for monitoring such particular taxonomic groups have been abundantly developed: these groups may harbour some ecological and/or conservation importance; they can potentially be used as indicator species; and their monitoring is practical, or at least feasible. Indeed, plants constitute habitats and resources for a wide-range of often specialized phytophagous invertebrates (Jaenike 1990), hence monitoring plants is also a way to indirectly monitor the potential presence of a series of interacting organisms (Basset et al. 2012, Castagneyrol and Jactel 2012). Plants can also be used as indicator species of a wide range of environmental (Tichy et al. 2023) and historic (Hermy et al. 1999) variables. Birds have also been proposed as potential indicator species of forest disturbance (Canterbury et al. 2000), as well as wood-inhabiting beetles (Nilsson et al. 2001). Wood-living beetles in particular are important components of forest biodiversity and play a key role in wood decomposition and nutrient cycling (Stokland et al. 2012). The functional composition of this taxonomic group is further influenced by forest management practices (Gossner et al. 2013) and is thus a group of conservation concern.

Focusing forest biodiversity monitoring on particular taxonomic groups also comes with a number of shortcomings. Indicator species remain surrogates of variables of interest and are also often influenced by the environmental, historical and landscape contexts in which they are applied (Gossner et al. 2014, Lalechère et al. 2017). Besides, several taxonomic groups may respond differently to human management or natural perturbations, rendering the generalization from one group to the others risky (Barlow et al. 2007, Cours et al. 2023). It is thus important to expand forest biodiversity monitoring to a larger number of taxonomic groups than what is currently performed (Nilsson et al. 2001, Burrascano et al. 2023). Among the often-neglected taxa, soil organisms account for ca. 60% of the total number of species on Earth (Anthony et al. 2023) and have an essential role in ecosystem functioning (Bardgett and Van der Putten 2014). Yet, relatively few monitoring programs focus on forest soil biodiversity (Korboulewsky et al. 2016).

A major obstacle to the study of the diversity of soil organisms is the rarity of suitable

68   taxonomic expertise (Eisenhauer et al. 2017), and the difficulty or impossibility to identify

larval stages that can substantially contribute to soil fauna communities (Paquin and Coderre

70   1997).   Molecular approaches have been developed for more than two decades for

taxonomic identification of a wide range of taxa (Hebert et al. 2003), including soil organisms

72   (Rougerie et al. 2009, Decaens et al. 2013). They consist in sequencing a small region of the

genome of the studied organism that is called the DNA barcode. The cost of DNA barcoding

74   was however prohibitive until recently, limiting its widespread use in biodiversity monitoring

programs. But this cost has dramatically decreased recently, thanks to the use of nanopore

76   sequencing technologies and of low-cost protocols for DNA extraction and amplification

(Srivathsan et al. 2021). In the meantime, reference libraries are continuously completed for

78   a wide range of taxa (e.g., Rougerie et al. 2015, Santos-Perdomo et al. 2024). For instance,

the Barcode of Life Database (BOLD) now gathers more than 19 million barcode sequences,

80   and coordinated efforts to fill the remaining gaps are underway (Roslin et al. 2022). The

massive barcoding of individual specimens therefore constitutes a promising avenue to

82   speed-up and largely develop biodiversity monitoring, in particular for less studied taxa.

84   Barcoding a large number of individual specimens simultaneously has been recently coined

as "megabarcoding" (Chua et al. 2023) to distinguish it from the "metabarcoding" approach

86   in which mixtures of organisms or their living media (environmental DNA) are sequenced.

The potential of metabarcoding, and in particular environmental DNA, for biodiversity

88   monitoring has been abundantly developed in the literature (Rodriguez-Ezpeleta et al. 2021).

The great advantage of metabarcoding is its speed of use: an environmental sample is

90   quickly processed and its metabarcoding informs on a list of species that have left DNA

traces in the sample. The main limitation of this approach is that the inference of species

92   abundances from metabarcoding results remains challenging (Elbrecht and Leese 2015, Beng

and Corlett 2020). Another limitation of the environmental DNA approach is that it

94   integrates the DNA signal over spatio-temporal scales that are still difficult to appraise (Bloor

et al. 2021). In contrast, megabarcoding enables to assess species abundances by processing

96   individual specimens and to control the spatiotemporal scale of sampling, as traditional

inventory methods. In particular, it should enable to track spatial and temporal changes in
98   species relative abundances, which usually prelude species turnovers.

100   Given the rapid and promising developments of low-cost megabarcoding, comparisons of its
efficiency among a full range of taxa composing each ecosystem compartments is now
102   needed. Low-cost megabarcoding has been successfully applied to Dipteran specimens
(Srivathsan et al. 2021, Hartop et al. 2024). A recent work using a higher cost protocol
104   extended this application to a wider range of taxa, but mainly epigeal ones (Hebert et al.
2025). Soil macrofauna gathers various taxonomic groups with varied physiological and
106   anatomical attributes that may compromise the ability of low-cost protocols to efficiently
extract and amplify DNA, such as the presence of protective shells (in Styllommatophora), of
108   dense exoskeletal cuticles (in Insecta, Isopoda and Diplopoda) or the presence of mucus (in
Styllommatophora and Haplotaxida). For instance, mollusc slime has been shown to
110   compromise amplification success (Chakraborty et al. 2020). Our aim in this study was thus
to test the applicability of low-cost megabarcoding to monitor soil fauna communities in a
112   temperate forest. We specifically assessed 1) whether the low-cost protocol proposed by
Srivathsan et al. (2021) for the extraction, amplification and purification of DNA can be
114   effectively adapted to soil fauna taxa, 2) the completeness of the BOLD database for the taxa
encountered in our survey, and 3) the relative ability to monitor larval stages with DNA
116   barcoding, compared to the one obtained for adult individuals.

2. Material and methods

120   2.1 Soil fauna collection and barcoding protocol

We performed an inventory of soil macrofauna in a beech forest stand located in Laqueuille
122   in the French Massif Central (exact location: 45.6236 N, 2.7393 E, altitude: 1182m) in June
2022. This stand was rather homogeneous with low micro-habitat diversity, except some
124   patches of *Vaccinum myrtillus* near the stand edge. Twenty-five soil samples were gathered
within a 50 m square. Each sample consisted in hand sorted individuals larger than 3 mm

126   extracted from soil monoliths of 18 cm x 18 cm and 10 cm deep together with the litter

above. These individuals were then immediately stored in 96° ethanol. Individuals were first

128   visually identified using a stereomicroscope (Zeiss Stemi 2000-C). This visual identification

enabled us to sort the 1413 individuals into morpho-species that had variable taxonomic

130   resolutions. Among them, 130 individuals were either identified at the species level, or were

singletons of a morpho-species and were thus treated as a distinct species in the following.

132   The remaining 1283 partially identified individuals were barcoded following the protocol

detailed in Srivathsan et al. (2021).

134   This barcoding protocol consisted in rinsing each individual with purified water (Elga

Labwater Purelab Ultra MK2) and placing it in one well of a 96-well PCR microplate. If the

136   individual was too large to fit into a well, one of its legs (or anus for earthworms) was

detached and put into the well. DNA extraction was performed using a HotShot solution

138   (Truett et al. 2000) with a heating at 65°C for 18 minutes and 98°C for 2 minutes. Extracted

DNA was then amplified within a mixture of 3 µL of extracted DNA, 7 µL of 2x Mastermix

140   (CWBio CW0682L), 1 µL of BSA, 2 µL of tagged primers. Left and right primers were tagged

with 9 bp tags developed by Srivathsan et al. (2024) so that each well had a unique

142   combination of left and right tags. We used the BF3-BR2 primer set (Elbrecht and Leese

2017) to amplify a 418 bp region of the COI barcode for 808 individuals and the LCO1490-

144   HCO2198 primer set (Folmer et al. 1994) to amplify a 658 bp region of the COI barcode for

the remaining 475 individuals. We used these two different sets to compare their efficiency

146   to amplify the COI "Folmer" region of soil fauna. A previous benchmark on 21 primer sets by

Elbrecht et al. (2019) revealed that the BF3-BR2 primer set performed better than the

148   classical LCO1490-HCO2198 set when analysing a Malaise trap sample. We wanted to test

whether this enhanced performance was also recovered on soil taxa. We used the following

150   PCR settings: 15 min at 95°C followed by 25 cycles of 30 s at 94°C, 1 min 30 s at 50°C and 1

min 30 s at 72 °C, followed by 10 min at 72°C. Amplified DNA was then pooled (fixed volume

152   of 5 µL for each well) and a subset of 500 µL of this pooled DNA was purified using Ampure

XP beads at 0.5 X. Library preparation for sequencing on a flongle followed the ONT protocol

154   (with the modifications explained in Srivathsan et al. 2021). Srivathsan et al. (2021)

recommended to pool up to 3 PCR plates on a flongle, while Hebert et al. (2025)

156   demonstrated that up to 20 PCR plates could be pooled on the same device. We thus used

an intermediate number and pooled between 6 and 10 PCR plates on each flongle. Note that
to decrease barcoding costs, PCR success was not assessed with gel electrophoresis.

2.2 Bioinformatic processing

Raw sequences were then processed with the software ONTbarcoder 2.0 (Srivathsan et al. 2024) and only quality-checked consensus barcodes (i.e., no ambiguities nor insertion/deletion) for each well were retained. The individuals for which the first sequencing round failed were re-processed from scratch (extraction, amplification, purification and library preparation) for a second round of sequencing, and those still in failure were processed a third time. The barcodes obtained were compared to the BOLD database (Ratnasingham and Hebert 2007) using Boldigger3 (Buchner and Leese 2020). We used the animal library (public and private), the rapid search option, and similarity thresholds of 99%, 97% and 90% for species, genus and family respectively. This step was performed on the twenty first of May 2025. This choice of using this user-friendly identification engine was motivated to keep the process simple and easy to reproduce, although it comes with a potential cost of neglecting some reference sequences that may be present in other databases, notably GenBank. We then clustered barcodes that could not be matched to a species on the BOLD database into distinct OTUs (operational taxonomic units) using a UPGMA method and a similarity threshold of 99% (Buchner and Leese 2020). We replicated our analyses using alternative thresholds of 98% and 97% to assess the sensitivity of our results to this arbitrary threshold and obtained very similar results (Figure S9).

2.3 Data analysis

We computed the barcoding efficiency as the percentage of consensus barcodes obtained (the number of obtained barcodes divided by the number of sequenced individuals). We repeated this computation for each of the three rounds of sequencing and separately for this list of taxonomic groups: Arachnida, Chilopoda, Diplopoda, Diplura, Haplotaxida, Insecta, Isopoda and Stylommatophora.

We also computed the barcoding efficiency for each set of primers (BF3/BR2 and
186   LCO1490/HCO2198) separately. To test whether the BF3-BR2 primer set lead to a larger
barcoding efficiency than the LCO1490-HCO2198 set, we fitted a Bernouilli GLM explaining
188   barcoding success of an individual as a function of the primer set used and of the taxonomic
group of the individual. This analysis was restricted on data of the first round of barcoding
190   and was replicated with and without interaction terms between primer set and taxonomic
groups.

192   We computed OTUs accumulation curves to assess the completeness of our species
inventory, distinguishing the adult-only and total assemblages of soil fauna individuals. This
194   was performed with the function "specaccum" of the vegan R package (Oksanen et al. 2007).
We finally compared the summary statistics of inventories based either on visual
196   (morphospecies) or on barcoding-based identification by computing species/OTU
abundance distributions, richness, Shannon's index, Pielou's equitability index and average
198   abundance-based Bray-Curtis dissimilarity between samples.

Datasets to replicate the analyses are included in the Supplementary Information (Tables S1
200   and S2) as well as the corresponding R script (Text S3). This material is also available on
public repositories (urls can be found at the end of the manuscript).

3. Results

204   Using a low-cost DNA barcoding protocol (see Tables S4 and S5 for detailed costs), we
obtained an overall barcoding efficiency of 88 %: we recovered 1124 DNA barcodes out of
206   the 1283 individuals processed (Fig. 1A). These barcodes clustered into 122 distinct
Operational Taxonomic Units (OTUs) among which 65 (53 %) had a species name in the BOLD
208   database (Fig. 1B). Interestingly, OTUs without species names in the BOLD database were
not solely the rarest taxa in our samples (Fig. 1C). We obtained a larger barcoding efficiency
210   on larvae (96 % of the 234 larvae processed) than on adult individuals (86 % of the 1049
adults processed). Barcodes of larvae had moreover a larger percentage of species name
212   presence in the BOLD database (77 % versus 37 %).

Figure 1: DNA barcoding efficiency of soil macrofauna. Panel A: number of individuals for

216   which DNA barcodes have been obtained and the corresponding matches at various

taxonomic levels in the BOLD database. Panel B: number of Operational Taxonomic Units

218   (OTUs) and the corresponding matches at various taxonomic levels in the BOLD database.

Panel C: Distribution of OTU abundances among OTU with a species name present or absent

220   in the BOLD database.

222   DNA barcoding efficiency was highly heterogeneous among taxonomic groups (Figure 2),

varying between 22 % (for Stylommatophora) and 100 % (for Arachnida). The percentage of

224   species name presence in the BOLD database was also highly heterogeneous among these

groups, varying between 0.3 % (for Diplura) and 82 % (for Insecta). We observed some

226   variation in these numbers depending on the primer set used; the barcoding success among

taxonomic groups for each primer set separately is depicted in Supplementary Figure S8.

230 **Figure 2**: Heterogeneity of DNA barcoding efficiency among taxonomic groups. Barcoding efficiency is equal to the proportion of individuals in each taxonomic group for which a DNA

232 barcode could be obtained. Species match is the proportion of individuals for which a species match was obtained in the BOLD database. Ar: Arachnida, In: Insecta, Diu: Diplura, Is:

234 Isopoda, Ch: Chilopoda, Ha: Haplotaxida, Dio: Diplopoda, St: Stylommatophora. The numbers above the bars correspond to the number of individuals processed in each taxonomic group.

Our study also revealed that DNA barcoding failure could largely be reduced by re-

238 processing failed individuals (Figure 3). Using this approach, barcoding efficiency sequentially improved from 59 % (using a single barcoding round for all individuals), to 81 % (using a

240 second barcoding round for individuals in failure) up to 88 % (using a third barcoding round for individuals in failure after the first two rounds). The barcoding efficiency of the first two

242 rounds was close (59 % and 56 % for round 1 and 2 respectively), while the one of the third round was lower (36 %). Finally, we observed a sizeable difference in barcoding efficiency

244 among the two pairs of primers used (68 % and 43 % for BF3/BR2 and LCO1490/HC02198 respectively). This difference was significant ($p < 0.001$) and did not interact significantly with

246 taxonomic groups (see Supplementary Text S7 for the full results).

Figure 3: Improvement in DNA barcode efficiency brought by multiple rounds of barcoding.
250   Each bar height stands for the barcoding efficiency of the barcoding round. The broken line
represents the cumulative barcoding efficiency after a given number of barcoding rounds.

Our study further revealed that 25 samples of soil monoliths of 18 cm x 18 cm x 10 cm were
254   not sufficient to reach an asymptote in the OTUs accumulation curve (Figure 4).
Interestingly, OTUs accumulated faster when including larvae individuals, although larvae
256   only accounted for 18 % of the total number of individuals. This was linked to the fact that
39 of the 148 OTUs (26 %) were only found among larvae individuals.

Figure 4: OTU accumulation curves as a function of the number of samples considered. The plain blue curve represents the OTU accumulation curve for the total community (adult and larvae individuals) while the dashed orange curve represents the OTU accumulation curve for the adult individuals only. Light colour areas depict confidence intervals.

Our analysis finally revealed that the barcoding-based species inventory strongly modified the initial morphospecies-based inventory, by more than doubling the number of OTUs detected (from 74 morphospecies to 148 OTUs, Figure 5B) and strongly altering biodiversity indices such as Shannon's index (Fig. 5C), Pielou's equitability index (Fig. 5D) and average Bray-Curtis dissimilarity between samples (Fig. 5E). The barcoding-based OTU abundance distribution was consequently strongly altered by being much more even than the original morphospecies-based abundance distribution (Figure 5A). The taxonomic resolution of the barcoding-based inventory was also sharply improved with the number of individuals with a species-level identification multiplied by more than three, from 130 individuals to 488 individuals.

Figure 5: Comparison of the morphospecies-based and the barcoding-based abundance
distributions (A), richness (B), Shannon's index (C), Pielou's equitability index (D) and average
abundance-based Bray-Curtis dissimilarity index between samples (E).

4. Discussion

282 Our study demonstrates a clear success of megabarcoding to assess the diversity of forest
soil macrofauna. Nearly 90% of the processed individuals were successfully barcoded and we
284 were able to attribute a species name to nearly half of them (Fig. 1). Since taxonomic
expertise of soil organisms is particularly rare, megabarcoding of sampled individuals could
286 therefore constitute an operational way to perform soil biodiversity monitoring. Another
major advantage of megabarcoding of soil macrofauna is the identification of larvae. In our
288 study case, we found that larvae accounted for 18% of sampled individuals, and contained a
large number of OTUs not found among adult individuals (26% of the total OTU number).
290 Therefore, the inclusion of soil larvae in biodiversity assessment, enabled by megabarcoding,
may profoundly alter our understanding of the rich forest soil fauna diversity (Fig. 4 and 5).

292 Our study further reveals that barcoding efficiency is variable among taxonomic groups, as
previously demonstrated by Hebert et al. (2025). Despite this variation, barcoding efficiency

294   was good to very good (above 80 % or 90 %) for most taxonomic groups, except for

Stylommatophora and to a lesser extent Diplopoda (Fig. 2). This may explain why barcoding

296   efficiency was remarkably large (96.1 %) for larvae individuals that mostly belonged to

Coleoptera, Araneae and Diptera and thus not to these more problematic taxonomic groups

298   (Table S1). The lower barcoding efficiency for the Stylommatophora and Diplopoda

individuals echoes previous studies that reported barcoding difficulties in these taxonomic

300   groups (Spelda et al. 2011, Chakraborty et al. 2020). It may be due to a variety of reasons

(notably, failure of or poor DNA extraction, lack of or low DNA amplification) that would

302   need to be further investigated. Note that Stylommatophora individuals were not extracted

from their shells before DNA extraction. This may be part of the explanation for the lower

304   barcoding efficiency of this group, on top of mucus-related amplification difficulties

(Chakraborty et al. 2020). The number of processed individuals for these two groups was

306   nevertheless low (23 and 42 respectively, Fig. 2) so that additional tests will be required to

confirm or not this lower barcoding efficiency. And solutions to this problem may be sought,

308   for instance by lengthening the heat extraction of DNA in the HotShot solution, by testing

alternative primer sets for these groups (e.g., "C_LepFolF- C_LepFolR" as in Spelda et al.

310   2011), or by sequencing individuals of these groups separately from the other taxonomic

groups. This last solution may decrease the among-individual variability in numbers of

312   sequencing reads, thereby increasing the number of reads obtained for the individuals of

these difficult groups.

314   The barcoding efficiency that we reached on the first two rounds was slightly above 50 %

(Fig. 3). This is lower than the ones reported by previous studies that were mostly in the

316   range of 80-90 % (Srivathsan et al. 2024, Hebert et al. 2025). Various factors can be put

forward to explain the lower barcoding efficiency that we obtained in this study. A first

318   explanation lays in the taxonomic groups studied. Previous megabarcoding studies have

notably targeted Dipteran individuals (Srivathsan et al. 2021) that lead to larger barcoding

320   efficiencies than Coleopteran and Arachnid individuals (Hebert et al. 2025). A second

explanation linked to the first one is that we pooled individuals of different taxonomic

322   groups in the same flongle. Sorting individuals of different taxonomic groups and sequencing

them separately may enable some enhancement in barcoding efficiency, since the DNA

324   sequences of more productive taxonomic groups would not overwhelm the ones of less

producing groups. A third explanation lays in the number of microplates that we pooled on each flongle for sequencing. Srivathsan et al. (2024) recommended to pool up to 3 microplates on a flongle, while Hebert et al. (2025) demonstrated that up to 20 microplates could be pooled on a single flongle, but with the use of high-fidelity Taq polymerases for DNA amplification. We chose intermediate numbers and pooled between six and ten microplates on each flongle. Although this strategy led to a smaller barcoding efficiency, we found that it was a good compromise to speed up the treatment of samples, since the library preparation steps are long relative to the extraction and amplification ones (Table S6), and to further decrease barcoding costs (Table S4). Importantly, by re-sequencing individuals in failure of barcode recovery, we demonstrated that final barcoding efficiency was comparable to the one of previous studies (Fig. 3). For this, we recommend using two rounds of barcoding, since the third round provides comparatively fewer barcodes (Fig. 3).

As argued before by Srivathsan et al. (2021), the cost of barcode sequencing has dramatically decreased thanks to the use of nanopore technologies. But the total cost of DNA barcoding also incorporates the costs of reagents involved in the extraction and amplification of DNA. The protocol of Srivathsan et al. (2021) put forward a number of innovations to cut down these costs, including the use of low-cost HotShot solutions to extract DNA instead of commercial kits and the use of low PCR reaction volumes. We provide in Table S4 the details of the reagent costs implied in the full protocol and compute that the production of a DNA barcode in this study costed around one euro. Although this cost depends on the commercial arrangements each lab may get with reagent suppliers, this nevertheless gives the order of magnitude of megabarcoding costs. These costs could be further decreased by using MinION flow cells instead of Flongles for larger sample sizes. We also detail the equipment costs to perform megabarcoding and demonstrate that it amounts to around 20,000 euros for minimal equipment needs (Table S5). Finally, following this protocol, a single lab technician can barcode around 1000 (2000) individuals per week if the lab is equipped with a single (two) thermocycler(s) and thus obtain around 500 (1000) barcodes (Table S6).

A current limitation of megabarcoding is that reference databases are still lacunar, especially for soil fauna (Recuero et al. 2024). In our study, nearly half of barcodes could not be matched to a referenced species in the BOLD database. This limitation is however not

definitive. Indeed, reference databases are updated at a rapid pace: the number of sequences in the BOLD database has increased by 67 % in 2024. Besides, coordinated
regional and national efforts are under way to barcode missing taxa (e.g., Spelda et al. 2011, Hendrich et al. 2015, Roslin et al. 2022). This implies that currently unmatched sequences
could be matched in the future, so that inventories based on megabarcoding can be progressively updated, a feature especially important for longitudinal and/or comparative
studies. To illustrate this point, during the time interval of the review process, the number of individuals that could be matched to a species in the BOLD database increased by 10% (i.e.,
44 additional individuals).

5. **Conclusion**

We demonstrated that the low-cost megabarcoding protocol of Srivathsan et al. (2024)
368 could be fruitfully applied to forest soil macrofauna and that it led to an overall very good barcoding efficiency of nearly 90 % (Fig. 1). Two taxonomic groups, Styllommatophora and
370 Diplopoda, would require further study to reach such high level of barcoding efficiency (Fig. 2). Megabarcoding of soil fauna also enables the identification of larvae individuals. Our
372 study demonstrates that these larvae individuals substantially contribute to macrofauna biodiversity (Fig. 4), with many larvae OTUs not encountered among adult individuals.
374 Finally, megabarcoding enables to keep track of OTUs abundances and therefore provides a strongly renewed assessment of soil macrofauna community structure (Fig. 5). With a cost of
376 ca. one euro per barcode, megabarcoding constitutes a concrete avenue to vastly develop the monitoring of forest soil macrofauna.

CRediT authorship contribution statement

380 **Franck Jabot**: Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation, Methodology, Project administration, Supervision, Writing – original draft.
382 **Gwenaëlle Auger**: Data curation, Investigation, Writing – review and editing. **Pauline Bonnal**: Data curation, Methodology, Investigation, Writing – review and editing. **Mathilde**
384 **Pizaine**: Data curation, Investigation, Writing – review and editing. **Marilyn Roncoroni**:

Investigation, Writing – review and editing. **Julien Pottier**: Conceptualization, Supervision,

386   Writing – review and editing.

388   **Declaration of Competing Interest**

The authors declare that they have no known competing financial interests or personal

390   relationships that could have appeared to influence the work reported in this paper.

392   **Data availability statement**

Datasets to replicate the analyses are included in the Supplementary Information (Tables S1

394   and S2) as well as the corresponding R script (Text S3). This material has also been deposited

on public repositories, the datasets can be found at doi:10.57745/LPYOC1 and the R script at

396   Link2.

398   **Acknowledgements**

This work was supported by the French Agence Nationale de la Recherche [grant number

400   ANR-20-CE32-0004], by the IDEX-ISITE initiative [grant number 16-IDEX-0001(CAP 20-25)]

and by Clermont Auvergne Metropole. We also thank Amrita Srivathsan for her advices to

402   replicate their low-cost barcoding protocol and Isabelle Bosio for her help during field work.

404   **References**

-Anthony, M. A., Bender, S. F., & van der Heijden, M. G. (2023). Enumerating soil biodiversity.

406   *Proceedings of the National Academy of Sciences*, *120*(33), e2304663120.

- Bardgett, R. D., & Van Der Putten, W. H. (2014). Belowground biodiversity and ecosystem

408   functioning. *Nature*, *515*(7528), 505-511.

- Barlow, J., Gardner, T. A., Araújo, I. S., et al. (2007). Quantifying the biodiversity value of tropical

410   primary, secondary, and plantation forests. *Proceedings of the National Academy of Sciences*,
        104(47), 18555-18560.

412   - Basset, Y., Cizek, L., Cuénoud, P., et al. (2012). Arthropod diversity in a tropical forest. *Science*, *338*(6113), 1481-1484.

414   - Beng, K. C., & Corlett, R. T. (2020). Applications of environmental DNA (eDNA) in ecology and conservation: opportunities, challenges and prospects. Biodiversity and conservation, 416   29(7), 2089-2121.

- Bloor, J. M., Si-Moussi, S., Taberlet, P., Carrère, P., & Hedde, M. (2021). Analysis of complex trophic 418   networks reveals the signature of land-use intensification on soil communities in agroecosystems. *Scientific Reports*, *11*(1), 18260.

420   - Buchner, D., & Leese, F. (2020). BOLDigger–a Python package to identify and organise sequences with the Barcode of Life Data systems. *Metabarcoding and Metagenomics*, *4*, e53535.

422   - Burrascano, S., Chianucci, F., Trentanovi, G., et al. (2023). Where are we now with European forest multi-taxon biodiversity and where can we head to?. *Biological conservation*, *284*, 110176.

424   - Canterbury, G. E., Martin, T. E., Petit, D. R., Petit, L. J., & Bradford, D. F. (2000). Bird communities and habitat as ecological indicators of forest condition in regional monitoring. *Conservation Biology*, 426   *14*(2), 544-558.

- Castagneyrol, B., & Jactel, H. (2012). Unraveling plant–animal diversity relationships: a 428   meta-regression analysis. *Ecology*, *93*(9), 2115-2124.

- Chakraborty, S., Saha, A., & Neelavar Ananthram, A. (2020). Comparison of DNA extraction methods 430   for non-marine molluscs: is modified CTAB DNA extraction method more efficient than DNA extraction kits?. *3 Biotech*, *10*(2), 69.

432   - Chua, P. Y., Bourlat, S. J., Ferguson, et al. (2023). Future of DNA-based insect monitoring. *Trends in Genetics*, *39*(7), 531-544.

434   - Cours, J., Bouget, C., Barsoum, N., et al. (2023). Surviving in changing forests: abiotic disturbance legacy effects on arthropod communities of temperate forests. *Current Forestry Reports*, *9*(4), 189- 436   218.

- Decaëns, T., Porco, D., Rougerie, R., Brown, G. G., & James, S. W. (2013). Potential of DNA barcoding 438   for earthworm research in taxonomy and ecology. *Applied Soil Ecology*, *65*, 35-42.

- Eisenhauer, N., Antunes, P. M., Bennett, A. E., et al. (2017). Priorities for research in soil ecology. 440   *Pedobiologia*, *63*, 1-7.

- Elbrecht, V., & Leese, F. (2015). Can DNA-based ecosystem assessments quantify species 442   abundance? Testing primer bias and biomass—sequence relationships with an innovative metabarcoding protocol. *PloS one*, *10*(7), e0130324.

444   - Elbrecht, V., & Leese, F. (2017). Validation and development of COI metabarcoding primers for freshwater macroinvertebrate bioassessment. *Frontiers in Environmental Science*, *5*, 11.

446   - Elbrecht, V., Braukmann, T. W., Ivanova, N. V., et al. (2019). Validation of COI metabarcoding primers for terrestrial arthropods. *PeerJ*, *7*, e7745.

448   - Folmer, O., Hoeh, W. R., Black, M. B., & Vrijenhoek, R. C. (1994). Conserved primers for PCR amplification of mitochondrial DNA from different invertebrate phyla. *Molecular Marine Biology and*
450   *Biotechnology*, *3*(5), 294-299.

   - Gossner, M. M., Lachat, T., Brunet, J., et al. (2013). Current near-to-nature forest management
452   effects on functional trait composition of saproxylic beetles in beech forests. *Conservation Biology*, *27*(3), 605-614.

454   - Gossner, M. M., Fonseca, C. R., Pašalić, E., Türke, M., Lange, M., & Weisser, W. W. (2014). Limitations to the use of arthropods as temperate forests indicators. *Biodiversity and Conservation*,
456   *23*, 945-962.

   - Hartop, E., Lee, L., Srivathsan, A., Jones, M., et al. (2024). Resolving biology's dark matter: species
458   richness, spatiotemporal distribution, and community composition of a dark taxon. *BMC biology*, *22*(1), 215.

460   - Hebert, P. D., Ratnasingham, S., & De Waard, J. R. (2003). Barcoding animal life: cytochrome c oxidase subunit 1 divergences among closely related species. *Proceedings of the Royal Society of*
462   *London. Series B: Biological Sciences*, *270*(suppl_1), S96-S99.

   - Hebert, P. D., Floyd, R., Jafarpour, S., & Prosser, S. W. (2025). Barcode 100k specimens: in a single
464   nanopore run. *Molecular Ecology Resources*, *25*(1), e14028.

   - Hendrich, L., Morinière, J., Haszprunar, G., Hebert, P. D., Hausmann, A., Köhler, F., & Balke, M.
466   (2015). A comprehensive DNA barcode database for Central European beetles with a focus on Germany: adding more than 3500 identified species to BOLD. *Molecular Ecology Resources*, *15*(4),
468   795-818.

   - Hermy, M., Honnay, O., Firbank, L., Grashof-Bokdam, C., & Lawesson, J. E. (1999). An ecological
comparison between ancient and other forest plant species of Europe, and the implications for forest conservation. *Biological conservation*, *91*(1), 9-22.

- Jaenike, J. (1990). Host specialization in phytophagous insects. *Annual Review of Ecology and Systematics*, 243-273.

- Korboulewsky, N., Perez, G., & Chauvat, M. (2016). How tree diversity affects soil fauna diversity: a review. *Soil Biology and Biochemistry*, *94*, 94-106.

- Lalechère, E., Jabot, F., Archaux, F., & Deffuant, G. (2017). Non-equilibrium plant metapopulation dynamics challenge the concept of ancient/recent forest species. *Ecological modelling*, *366*, 48-57.

- Nilsson, S. G., Hedin, J., & Niklasson, M. (2001). Biodiversity and its assessment in boreal and nemoral forests. *Scandinavian Journal of Forest Research*, *16*(S3), 10-26.

- Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M. H. H., Oksanen, M. J., & Suggests, M. A. S. S. (2007). The vegan package. *Community ecology package*, *10*(631-637), 719.

- Paillet, Y., Bergès, L., Hjältén, J., et al. (2010). Biodiversity differences between managed and unmanaged forests: Meta-analysis of species richness in Europe. *Conservation biology*, *24*(1), 101-
112.

   - Paquin, P., & Coderre, D. (1997). Changes in soil macroarthropod communities in relation to forest
maturation through three successional stages in the Canadian boreal forest. *Oecologia*, *112*, 104-111.

- Ratnasingham, S., & Hebert, P. D. (2007). BOLD: The Barcode of Life Data System (http://www. barcodinglife. org). *Molecular ecology notes*, *7*(3), 355-364.

- Recuero, E., Etzler, F. E., & Caterino, M. S. (2024). Most soil and litter arthropods are unidentifiable based on current DNA barcode reference libraries. *Current Zoology*, *70*(5), 637-646.

- Rodríguez-Ezpeleta, N., Zinger, L., Kinziger, A., et al. (2021). Biodiversity monitoring using environmental DNA. *Molecular Ecology Ressources*, 21(5), 1405-1409.

- Roslin, T., Somervuo, P., Pentinsaari, M., et al. (2022). A molecular-based identification resource for the arthropods of Finland. *Molecular Ecology Resources*, *22*(2), 803-822.

- Rougerie, R., Decaëns, T., Deharveng, L., et al. (2009). DNA barcodes for soil animal taxonomy. *Pesquisa Agropecuária Brasileira*, *44*, 789-802.

- Rougerie, R., Lopez-Vaamonde, C., Barnouin, T., et al. (2015). PASSIFOR: A reference library of DNA barcodes for French saproxylic beetles (Insecta, Coleoptera). *Biodiversity data journal*, (3).

- Santos-Perdomo, I., Suárez, D., Moraza, M. L., Arribas, P., & Andújar, C. (2024). Towards a Canary Islands barcode database for soil biodiversity: revealing cryptic and unrecorded mite species diversity within insular soils. *Biodiversity Data Journal*, *12*.

- Spelda, J., Reip, H. S., Oliveira–Biener, U., & Melzer, R. R. (2011). Barcoding Fauna Bavarica: Myriapoda–a contribution to DNA sequence-based identifications of centipedes and millipedes 504 (Chilopoda, Diplopoda). *ZooKeys*, (156), 123.

- Srivathsan, A., Lee, L., Katoh, K., et al. (2021). ONTbarcoder and MinION barcodes aid biodiversity discovery and identification by everyone, for everyone. *BMC biology*, *19*, 1-21.

- Srivathsan, A., Feng, V., Suárez, D., Emerson, B., & Meier, R. (2024). ONTbarcoder 2.0: rapid species discovery and identification with real-time barcoding facilitated by Oxford Nanopore R10. 4. *Cladistics*, *40*(2), 192-203.

- Stokland, N., Siitonen, J. & Jonsson, B.G. (2012). Biodiversity in dead wood. *Cambridge University Press*.

- Tichý, L., Axmanová, I., Dengler, J., et al. (2023). Ellenberg-type indicator values for European vascular plant species. *Journal of Vegetation Science*, *34*(1), e13168.

- Truett, G. E., Heeger, P., Mynatt, R. L., Truett, A. A., Walker, J. A., & Warman, M. L. (2000). Preparation of PCR-quality mouse genomic DNA with hot sodium hydroxide and tris (HotSHOT). 516 *Biotechniques*, *29*(1), 52-54.

Supplementary Information

520 -Table S1: Barcoding-enhanced inventory data. This dataset includes the identification of 1124 forest soil macrofauna individuals by DNA barcoding and the visual identification of 130 additional individuals.

522 -Table S2: Visual inventory data. This dataset includes the visual identification of 1413 forest soil macrofauna individuals.

524   -Text S3: R script to replicate the analyses and figures.

-Table S4: Data on reagent costs.

526   -Table S5: Data on equipment costs.

-Table S6: Duration of each barcoding step.

528   -Supplementary Text S7: Full results of the statistical analysis.

-Supplementary Figure S8: Barcoding success among taxonomic groups for each primer set.

530   -Supplementary Figure S9: Summary statistics of macrofauna biodiversity for different similarity thresholds to delineate OTUs.