Experimentally mimicking long term infections of *Magallana gigas* with the OsHV-1 virus reveals evolution through positive selection

Camille Pelletier[1], Nicole Faury[1], Mickaël Mege[1,#], Lionel Dégremont[1], Maelle Hattinguais[1], Jeremie Vidal-Dupiol[2], Germain Chevignon[1], Maude Jacquot[1,*], Benjamin Morga[1,*]

Affiliations

[1] Ifremer, RBE-ASIM, Unité Adaptations Santé des Invertébrés Marins, Avenue de Mus de Loup, 17390 La Tremblade, France.

[2] IHPE, Univ Montpellier, CNRS, IFREMER, Univ Perpignan Via Domitia, Montpellier, France

Current address: INRAE, UMR BIOEPAR, TiBoDi Team, Oniris – site de la Chantrerie, Route de Gachet - CS40706, 44307 Nantes cedex 3

*Corresponding authors:

- Benjamin Morga; E-mail address: benjamin.morga@ifremer.fr; Tel: +33 5 46 76 26 49

- Maude Jacquot; E-mail address: maude.jacquot@ifremer.fr; Tel: +33 5 46 76 26 82

Abstract

Ostreid herpesvirus 1 (OsHV-1) pose a significant threat to the global oyster farming industry, causing substantial economic losses due to mortality outbreaks. While OsHV-1 primarily affects the Pacific oyster *Magallana gigas*, it has also been associated with mortality events in various other host species. Despite progress in understanding OsHV-1 epidemiology, important knowledge gaps remain regarding its evolutionary mechanisms and adaptation to host genetic backgrounds. This study uses experimental evolution and extensive genomic analysis to investigate the dynamics of OsHV-1 evolution in response to oyster host genetic variation. Our results show that genetic mutations, particularly transitions and transversions, play a key role in shaping viral populations, contributing to a trend toward genetic homogenization. Notably, stronger positive selection signals were observed in oyster populations with higher susceptibility, suggesting adaptation of viral genotypes to specific host genetic backgrounds. These findings shed light on the complex evolutionary dynamics of OsHV-1 and its interactions with oyster hosts. Understanding how this virus adapts to host genetic diversity is crucial for developing strategies to mitigate its impact on the oyster farming industry and provides valuable insights into the broader mechanisms of viral evolution in response to host variation.

1. Introduction

Ostreid herpesvirus 1 (OsHV-1), a double-stranded DNA virus belonging to the *Malacoherpesviridae*, pose a significant threat to the global oyster farming industry by initiating a polymicrobial disease known as the Pacific Oyster Mortality Syndrome (POMS, de Lorgeril et al., 2018). In recent years, mortality outbreaks associated with POMS have caused substantial economic losses. Among several factors (*i.e.* temperature and food) driving windows of oyster permissivity to disease, oyster age is a key determinant, with individuals being susceptible up to ~18 months old (Azéma, Lamy, et al., 2017; Burioli et al., 2017; Nicolas et al., 1992; Renault et al., 1994). Meanwhile, oyster farmers have reported mortality associated with OsHV-1 mainly in spat, as survivors demonstrate genetic and epigenetic resistance upon subsequent exposure to the virus (Evans et al., 2017; Gawra et al., 2023).

The first comprehensive characterization of the OsHV-1 genome was published in 2005 (Davison et al., 2005). The genome of OsHV-1 spans approximately 207 kilobase pairs (kb) in length and exhibits a distinctive genomic organization comprising unique regions (U_L/U_S: unique long or short) flanked by repeated regions (TR_L/TR_S: terminal repeat long or short and IR_L/IR_S: inverted repeat long or short). The architecture of the OsHV-1 genome can be summarized as TR_L-U_L-IR_L-X-IR_S-U_S-TR_S (Davison et al., 2005). Advances in high-throughput sequencing have facilitated the genomic characterization of several OsHV-1 lineages, including OsHV-1 µVar (Burioli et al., 2017), OsHV-1 PT (Abbadi et al., 2018), and OsHV-1 SB (Xia et al., 2015). Currently, 28 complete genomes are available in public databases (Delmotte-Pelletier et al., 2022; Morga-Jacquot et al., 2021; Pelletier et al., 2025). These genomic datasets have revealed a spatiotemporal structuring of viral lineages in France (Delmotte-Pelletier et al., 2022) and globally (Morga-Jacquot et al., 2021), consistent with oyster farming practices. Furthermore, a recent study has highlighted a differentiation between lineages infecting *M. gigas* and *O. edulis*, with a greater viral diversity observed in *M. gigas* (Pelletier et al., 2025). However, despite significant advancements in understanding OsHV-1 epidemiology and evolution, several knowledge gaps persist regarding its evolutionary mechanisms and how it adapts to host genetic backgrounds. Indeed, substantial genetic and epigenetic variation for resistance to OsHV-1 has been reported worldwide,

enabling selective breeding programs to sustain oyster production (Camara et al., 2017; Dégremont et

al., 2015b; Divilov et al., 2019; Gutierrez et al., 2020; Gawra et al., 2023; Valdivieso et al., 2025).

The evolution of viruses is driven by a range of processes, including mutation, genetic recombination,

genetic drift, and natural selection. RNA viruses provide vivid examples of high mutation rates and rapid

evolution. These viruses are known for their ability to accumulate genetic changes swiftly, resulting in

substantial levels of viral polymorphism (Sanjuán, 2012; Sanjuán & Domingo-Calap, 2016, 2016). This

genetic diversity empowers them to adapt to newly infected cellular environments and develop strategies

to evade vaccines and antiviral drugs (Lauring, 2020). Although large dsDNA viruses exhibit lower

mutation rates (*i.e.* between 10^{-03} and 10^{-08} nucleotide substitution/site/year, e.g. 5.9×10^{-08} for the Herpes

simplex virus) compared to RNA viruses (*i.e.* between 10^{-02} and 10^{-04} nucleotide substitution/site/year)

due to their use of high-fidelity proofreading polymerases (Sanjuán et al., 2010; Sanjuán, 2012; Sanjuán

& Domingo-Calap, 2016), their genome stability is increasingly controversial as they sometimes display

noteworthy genetic diversity (Sanjuán et al., 2016).

Viral evolutionary processes can be investigated through various approaches, including sequence

analyses, metagenomics, phylogenetics, phylodynamics, and even Experimental Evolution (EE). This

latter approach involves subjecting viruses to controlled laboratory conditions, enabling the real-time

observation and manipulation of evolutionary processes (Kawecki et al., 2012). The emergence and

dynamics of mutations, immune escape variants, and evolutionary trade-offs can be assessed by

monitoring viral populations under different selective pressures, such as antiviral drug treatments or

interactions with immune systems (Kawecki et al., 2012).

While most studies using EE focus on easily replicable model organisms such as *Escherichia coli*,

Drosophila spp., *Saccharomyces cerevisiae* or various bacteriophages (Kawecki et al., 2012), recent

research has expanded EE to diverse viruses, including Herpesviruses (Fuandila et al., 2022; Kuny et

al., 2020).

For instance, *in vitro* EE conducted with populations of Herpes Simplex Virus type 1 (HSV-1) has

revealed the accumulation of minor genetic variants and changes in population diversity. Although

minor, these changes have been observed to lead to relatively rapid modifications in the viral population at the consensus level (Kuny et al., 2020). Another EE study, conducted on Koi herpesvirus (KHV), which infects the common carp *Cyprinus carpio*, has highlighted the key role of structural variations in driving rapid viral evolution *in vitro* (Fuandila et al., 2022). By driving genetic diversity, adaptation, and pathogenicity changes, these large-scale genomic alterations, including deletions, insertions, duplications, and inversions, can rapidly reshape viral genomes, influencing key functional regions (Fuandila et al., 2022).

In the absence of oyster cell lines for OsHV-1 propagation, we were forced to use *in vivo* techniques, which require high-throughput sequencing to efficiently and comprehensively analyze the evolution of viral populations. This however offered the advantage to study this host pathogen interaction as a whole in an ecologically relevant context.

In the context of OsHV-1, our understanding of the evolutionary processes contributing to the diversity of the viral population remains limited. The primary objective of the present study is to gain insights into the evolutionary mechanisms driving the adaptation and diversification of OsHV-1 lineages based on oyster susceptibility to infection. To achieve this goal, we used an EE approach spanning up to 28 generations under controlled conditions, combined with a comprehensive sequencing analysis of 96 OsHV-1 genomes.

2. Material and methods

2.1. Oysters production

To explore the impact of oyster genetic backgrounds on the evolutionary dynamics of OsHV-1, we generated three oyster populations with different levels of resistance to OsHV-1 infection. In December 2021, wild oysters were collected from two sites in Charente-Maritime (France) based on their proximity to oyster farms and history of viral exposure.

The first site, La Floride (Marennes-Oléron Bay, Charente-Maritime, Lat: 45.803; Long.: -1.153) is

surrounded by oyster farms, densely populated by wild oyster beds and is exposed to OsHV-1 outbreaks

annually (Dégremont et al., 2019). Oysters from this site have shown high resistance to POMS

(Valdivieso et al., 2025).

The second site, Chaucre (Charente-Maritime, Lat: 45.982; Long: -1.396), is located on the west coast

of Oléron Island in a low-density, non-farming area. Wild oysters collected from Chaucre are less

frequently exposed to OsHV-1 µVar than oysters from densely populated areas and are therefore

considered moderately susceptible to OsHV-1 µVar infection.

In addition, a control oyster population used in previous studies was included (Gawra et al., 2023;

Valdivieso et al., 2025). These oysters were initially sampled in 2008 prior to the emergence of OsHV-

1 µVar, and have since been bred for six generations using oysters from second-generation individuals

never exposed to OsHV-1 µVar. This lineage has been maintained in a secure facility with UV-treated

seawater (40 mJ/cm²), and is considered naive to OsHV-1 µVar since 2009. It has been described as

highly susceptible to OsHV-1 µVar, with mortality rates exceeding 80% (Valdivieso et al., 2025).

The three oyster populations (La Floride, Chaucre, and the naïve control) were reared separately at the

Ifremer hatchery in La Tremblade (December 2021) to prevent cross-contamination with pathogens such

as OsHV-1 and *Vibrio aestuarianus* (Dégremont et al., 2005). Seawater temperature was gradually

increased from 11°C to 20°C and oysters were fed with a cultured phytoplankton mix (*Isochrysis*

galbana, Tetraselmis suecica and *Skeletonema costatum*) to induce gametogenesis (Dégremont et al.,

2005).

In March 2022, 30 oysters per population were sexed, and gametes were collected via gonad stripping.

Female gametes were pooled and filtered through 20-µm and 100-µm meshes, then fertilized with sperm

from individual males. After 10 minutes, all fertilized eggs were pooled and transferred to 30 L tanks to

minimize sperm competition and maximize effective population size (Boudry et al., 2002). Larval and

spat culture followed established protocols (Dégremont et al., 2005, 2007). Progenies were reared in

UV-treated seawater until experimental OsHV-1 infections and designated as FA, NFA, and C,

corresponding to farming area (high density with the typical oyster bed of 100 individuals/m² and annual

POMS reported), non-farming area (low-density beds, less than 20 individuals/m² and no POMS

reported), and control populations (naïve laboratory-reared population, highly sensitive to OsHV-1

infection).

Before the experiment, spat were acclimated in 120 L tanks with a continuous flow of UV-treated

seawater heated to 19°C. The seawater was supplemented with a phytoplankton mixture containing *I.*

galbana, T. suecica and *S. costatum*. The acclimation period lasted at least 2 weeks to ensure optimal

growth and feeding conditions, critical for efficient OsHV-1 replication and observable oyster mortality

(Azéma, Maurouard, et al., 2017).

2.2. Viral suspension preparation

To generate a diversified viral inoculum for infecting all oyster populations, nine OsHV-1 suspensions

were produced between 2020 and 2021. These suspensions were obtained from spat oysters naturally

infected during field monitoring of OsHV-1 µVar outbreaks in various areas in France. Briefly,

approximately 1000 specific-pathogen-free (SPF) spat oysters, reared at the Ifremer experimental

facility in Argenton (Brittany, France), following a standardized protocol (Petton et al., 2015), were

deployed in four oyster farming areas during active disease outbreaks (Figure 1). These areas included:

i) Brest harbor (BR) (used to produce two viral suspensions) (Logonna-Daoulas, lat.: 48.335 long.: -

4.318), *ii)* Marennes-Oléron Bay (MO) (used to produce four viral suspensions) (La Floride, lat.: 45.803

and long.: -1.153), *iii)* Arcachon Basin (ARC) (used to produce two viral suspensions) (Grahude, lat.:

44.653 and long.: -1.073), *iv)* Leucate lagoon (LEU) (used to produce one viral suspension) (Leucate,

lat.: 43.379 and long.: 3.571). Seven days after deployment of SPF spat, once mortalities began, oysters

were retrieved and kept at 20°C for 7 days in the lab. During this period, moribund oysters were sampled

daily and stored at -80°C.

In total, nine viral suspensions were prepared following a standardized protocol (Schikorski et al., 2011)

(Figure 1, Table S1).

Figure 1: Overview of the bioinformatic workflow used to monitor OsHV-1 genomic diversity and to detect selection patterns during experimental evolution. The central panel shows the sampling map with the four French oyster-farming areas involved in the study between 2020 and 2021: BR (Bittany), MO (Marennes-Oléron), ARC (Arcachon), and LEU (Leucate). The number of samples processed in each region is indicated for both 2020 and 2021, with color-coded oyster illustrations representing each site. The left section (blue panels) describes the analytical pipeline applied to naturally infected oysters collected between 2020 and 2021 from four farming areas in France. DNA was individually extracted from mantle tissue of moribund oysters and sequenced using Illumina technology. Raw reads were assembled de novo into OsHV-1 NR-genomes using a dedicated bioinformatic pipeline previously developed for OsHV-1 genomic analysis (Dotto-Maurel-Pelletier et al., 2019). The right section (orange panels) illustrates the experimental evolution design and associated bioinformatic analyses. Nine viral suspensions were prepared from moribund oysters collected in the field, then pooled in equimolar proportions to create a standardized inoculum. This inoculum was used to infect oysters in a controlled evolution experiment (see Figure 2 for details). DNA extracted from infected oysters at successive time points was sequenced using Illumina technology. A specific bioinformatic pipeline was developed to detect and characterize signatures of selection.

2.3. Inoculum viral load quantification and viral suspension mix setup.

To prepare an equimolar viral inoculum for injection into oysters, the viral load of each of the nine viral suspensions was quantified. DNA was extracted from 100 µl of each viral suspension using the MagAttract® HMW DNA kit according to the manufacturer's instructions. DNA purity and concentration were measured using a Nano-Drop ND-1000 spectrophotometer (Thermo Scientific) and Qubit® dsDNA BR assay kits (Molecular Probes Life Technologies), respectively. Viral DNA copy number was then quantified by quantitative PCR using a Mx3005 P Thermocycler (Agilent) (Pepin et al. 2008). Each reaction contained 1 µl of Milli-Q water, 2 µl of each primer at 5.5 µM, OsHVDP For (forward) 5′-ATTGATGATGTGGATAATCTGTG-3′ and OsHVDP Rev (reverse) 5′-GGTAAATACCATTGGTCTTGTTCC-3′, 10 µl of Brilliant III Ultra-Fast SYBR® Green PCR Master Mix (Agilent), and finally 25 ng of DNA sample from infected oysters or Milli-Q water (non-template control) in a total volume of 20 µl. The qPCR program consisted of 3 min at 95 °C followed by 40 cycles of amplification at 95 °C for 5s and 60 °C for 20s. Results were expressed as the log-transformed copy number of viral DNA per microliter of seawater (cp/µL).

The nine viral suspensions were pooled to create the initial viral inoculum (Table S1). To ensure equal contribution from each suspension, the pool was adjusted to obtain an equimolar concentration of viral DNA (Table S1).

2.4. Experimental evolution design

The EE was conducted under controlled conditions in a Type 2 laboratory at Ifremer's La Tremblade facility. To initiate the experimental infections, 50 oysters from each population were first myorelaxed using hexahydrate $MgCl_2$ (50 g/L) (Suquet et al., 2009). Each oyster was then individually injected with 100 µL of pooled inoculum into the abductor muscle using a 26-gauge needle attached to a multi-dispensing hand pipette (Figure 2, step 1). At the same time, 10 oysters per population were myorelaxed and injected with artificial seawater to serve as control. The experiment was conducted in duplicate (*i.e.* two tanks per condition), with oysters placed in 5-L plastic tanks filled with seawater at 20°C and

continuously supplied with a phytoplankton mixture (*Isochrysis galbana, Tetraselmis suecica* and *Skeletonema costatum*). Mortality was monitored daily for seven days.

Following this initial infection, nine tanks were prepared: two containing infected oysters and one containing ASW-injected control oysters for each oyster population. These tanks served as the starting point, referred as "Generation 0" (Figure 2, step 2). To initiate the first transmission generation (G1) through cohabitation, 25 healthy oysters from each population were introduced into the corresponding infected tanks 24 hours after donor injection (Figure 2, step 3). For instance, 25 FA oysters were added to each FA-infected tank, and so on. After 24 hours of cohabitation, these healthy oysters were transferred to fresh, pathogen-free seawater to allow viral excretion and initiate the next infection generation (Figure 2, step 4). This procedure was repeated to produce a second generation, following the same protocol. During each cohabitation period, donor oysters were monitored for seven days, and all moribund oysters were collected for DNA extraction (Figure 2, step 5). In total, we completed 13, 18, and 28 generations for the FA, C, and NFA oyster populations, respectively. Mortality gradually declined across all experimental populations, indicating the cessation of active viral transmission and marking the end of the experimental infection.

2.5. DNA extraction and sequencing

DNA extraction was performed on oysters collected during mortality monitoring (Figure 2, step 5) at specific infection generations: G0, G5, G10, and G13 for the FA population, at generations G0, G5, G10, G15, and G18 for the C population, and at generations G0, G5, G10, G15, G20, G24, and G28 for the NFA population. Sampling every five generations was chosen to limit the number of samples processed while ensuring consistent infection dynamics. This approach explains the discrete distribution of viral copy numbers and confirms that viral loads remained relatively stable across generations.

For each oyster population, infection generation, and replicate, seven samples of 35 mg of mantle tissue were collected for DNA extraction, which were performed using the MagAttract® HMW DNA kit (Qiagen) following manufacturer's instructions. DNA purity and concentration were measured using a

Nano-Drop ND-1000 spectrophotometer (Thermo Scientific) and Qubit® dsDNA BR assay kits

(Molecular Probes Life Technologies), respectively. Viral DNA copy number were quantified by

quantitative PCR using a Mx3005 P Thermocycler (Agilent), as previously described. From each oyster

population and replicate, three individuals were selected based on their viral load and DNA

concentration. In total, 96 samples were sequenced by DNA-seq Illumina by IntegraGen SA (Evry,

France). PCR-free libraries were prepared using the Twist library Preparation Enzymatic Fragmentation

(EF) Kit 1.0 (Twist Bioscience) according to the supplier's recommendations. Briefly, after

quantification of double-stranded genomic DNA, 400 ng of DNA were enzymatically fragmented to

generate inserts of approximately 400 bp. Following fragmentation, libraries underwent end-repair, A-

tailing, ligation with unique dual-index (UDI) Illumina adapters. Libraries were then purified and size-

selected using SPRI beads, quantified by qPCR and sequenced on the Illumina NovaSeq platform to

generate paired-end 150 bp reads.

2.6. Statistical analysis of mortality and OsHV-1 viral load across generations and oyster

populations

Mortality and viral load data were analyzed using R v4.2.2. Raw data were graphically represented using

ggplot2 v3.5.2. Differences in mortality and viral load among generations and populations were assessed

using an ANOVA analysis computed using anova_test(factor1 ~ factor2) function from the rstatix

v0.7.2. package.

Figure 2: Experimental evolution design for one oyster population. The experimental evolution was conducted independently on three oyster populations using a serial cohabitation infection design. Healthy oysters are shown in dark blue, infected individuals in dark blue with red viral particles, and moribund oysters in light red with red viral particles. Experiments were performed in duplicate for each oyster population. At generation G0, 25 healthy oysters were injected intramuscularly with a standardized viral inoculum to initiate infection (step 1). After 24 hours, the infected oysters were transferred to tanks containing clean seawater (step 2). A new batch of 25 healthy oysters was then introduced to allow cohabitation and passive exposure to viral particles released into the water (step 3). After 24 hours, exposed oysters were moved to fresh tanks with pathogen-free seawater (step 4) to allow the of infection to progress under controlled conditions. Moribund individuals were monitored over a 7-day period (step 5), and 25 of them were selected to initiate the next generation (G1). This serial transfer process was repeated for each generation until the designated endpoint, with each round involving exposure, infection, and mortality monitoring.

2.7. Assembly of a global consensus for each viral suspension

To reconstruct the overall genetic composition of the viral inocula, we assembled 126 de novo OsHV-1 µVar NR-genomes from infected oyster tissues corresponding to the batches used for viral suspension preparation (Figure 1, part blue). DNA was extracted, quantified, and sequenced as previously described (see "DNA extraction and sequencing" section). DNA libraries were prepared using the Shotgun PCR-free library preparation kit (Lucigen) and sequenced on an Illumina NovaSeq™ 6000 system (paired-ends, 150 bp reads) by Genome Quebec Innovation Center (McGill University, Montreal, Canada).

Raw sequencing data are publicly available in the NCBI SRA database under Bioproject PRJNA1216400 with accession numbers SAMN46433918 to SAMN46434013 to ensure data accessibility and reproductibility (Table S2).

De novo OsHV-1 NR-genome assemblies were generated from sequencing reads as previously described using a custom bioinformatic pipeline (see Dotto-Maurel et al., 2022 for details on the bioinformatic pipeline). Briefly, for each library, sequencing reads were quality-filtered (PHRED score > 31) and trimmed using Fastp v0.20.1 (with parameter -q 31; Chen et al., 2018). Filtered reads were mapped to the *M. gigas* reference genome (accession number: GCF_902806645.1; Peñaloza et al., 2021) using Bowtie2 v2.4.1 (parameter --local; Langmead & Salzberg, 2012) to remove host sequences. The remaining reads were assembled *de novo* using metaSPAdes v3.15.4 (parameter --meta; Bankevich et al., 2012). OsHV-1 contigs were identified with BLASTn v2.12.0 (options -max_target_seqs 1, -evalue "1e-10"; Altschul et al., 1990), manually re-ordered in Geneious v2022.2.2 based on read overlaps, and reconstructed as NR-genomes to remove redundant inverted repeats. Assembly accuracy and coverage uniformity were validated by dotplot similarity and read realignment with Bowtie2.

To summarize the genetic diversity of the viral suspensions, the 126 assembled NR-genomes were aligned using MAFFT v1.4.0 (Katoh et al., 2002, Figure 1). In this manuscript, the resulting consensus is referred as "artificial OsHV-1 major NR-consensus".

2.8. Variant calling

To assess the genomic variability among individuals, intra-individual Single Nucleotide Variants (iSNVs) - defined as sites displaying at least two alleles with frequencies above 10% in at least one sample – were identified for each sequencing library based on the "artificial OsHV-1 major NR-consensus" previously generated. Variant calling was conducting using a modified version of a previously described pipeline for detecting polymorphisms (Figure 1, see Delmotte-Pelletier et al., 2022 for details of the bioinformatic workflow). Briefly, reads were filtered using Fastp v0.20.1 (Chen et al., 2018), mapped to the "artificial OsHV-1 major NR-consensus" using Bowtie2 v2.4.1 (Langmead & Salzberg, 2012) and variants were called using Freebayes v1.3.5 (Garrison & Marth, 2012), with the following settings: *-use-mapping-quality, -min-repeat-entropy 1, -haplotype-length 0, -min-alternate-count 5, -pooled-continuous, -hwe-priors-off, -allele-balance-priors-off*. The resulting variant calling outputs were normalized, decomposed, sort and deduplicated using vt v2005.11.10 (Tan et al., 2015). Only libraries with a minimum mean genome-wide coverage of 110× were retained for downstream analyses (Delmotte et al., 2022; Pelletier et al., 2025). iSNVs were included in the analysis only if their frequency was ≥10% in the viral population. Detected iSNVs were then mapped to their respective consensus genomes and aligned using MAFFT v1.4.0 (Katoh et al., 2002) implemented in Geneious Prime to re-coordinate all iSNVs position across samples. Each re-coordinated iSNV was assigned a unique identifier based on *(i)* its position within the multiple alignment, *(ii)* the reference allele (REF) and *(iii)* the alternative allele (ALT) (*i.e.* position_REF >ALT) following the approach described by Delmotte-Pelletier et al. (2022).

2.9. Analysis of substitutions types and clustering of individuals based on iSNVs

To obtain detailed information on the number of transitions (*i.e.* substitution of a purine by another purine or a pyrimidine by another pyrimidine) and transversions (*i.e.* substitution of a purine by a pyrimidine or a pyrimidine by a purine) within OsHV-1 genome across generations and among host

oyster populations, iSNVs detected in all libraries were analyzed using BCFtools stats v1.17 (Figure 1, Danecek et al., 2021). In the absence of a true ancestral genome, substitution directionality cannot be inferred with certainty, consequently, complementary substitutions (*e.g.* A>G and G>A) were collapsed into base pair substitution classes (*e.g.* A|G). In addition, iSNVs were analyzed separately depending on whether they were identified as ancestral or *de novo*, allowing distinction between newly emerging iSNVs and those inherited from previous viral generations. Substitution counts were averaged per generation and per oyster population and visualized using R v4.2.2 and the ggplot2 package v3.4.0 (Wickham, 2009). The ts:tv ratios were calculated to compare substitution patterns among samples, and a baseline ts:tv ratio of 0.5 was used to represent the expected value under random sequencing errors (Zook et al., 2014).

2.10. *Exploring sample clustering and iSNV data quality assessment through Principal Component Analysis*

In the context of EE studies, Principal Component Analysis (PCA) was employed to assess sample clustering and evaluate the quality of intra-individual Single Nucleotide Variants (iSNVs) data, allowing detection of underlying population structure and potential data anomalies (Patterson et al., 2006).

Based on iSNVs frequencies, individuals were projected into a two-dimensional space to assess sample clustering. PCA was performed using the pcrcomp function from the factoextra v1.0.7 package (Kassambara & Mundt, 2020) in R v4.2.2 (Figure 1). Cluster groupings were visualized using ellipses drawn with stat_ellipse() in ggplot2. This analysis incorporated information on oyster populations, infection generations and replicate identity.

2.11. *Selection detection*

Selection patterns were detected using a SNV-based approach. Since the experiment was conducted in duplicate for each oyster population, selected iSNVs could appear in both replicates or only in specific

ones. Selection of particular OsHV-1 lineages occurred independently within each oyster population, and consequently, all analyses were performed separately for each population. To detect iSNVs showing significant shifts in allele frequency (AF), we compared the initial generation (G0) with the evolved generations (up to G13, G18, and G28 for FA, C and NFA oyster populations, respectively). For this purpose, an adapted Cochran-Mantel-Haenszel (CMH) test was applied (Figure 1) using the ACER v1.0 package (Barghi et al., 2020; Spitzer et al., 2020) in R, which computed p-values for each iSNV, reflecting frequency changes over time while accounting for genetic drift and sampling variance (Spitzer et al., 2020).

P-values were then adjusted for multiple testing using the false discovery rate (FDR) method, with the p.adjust function from the stats v4.2.2 package (R Development Core Team, 2005). Adjusted p-values were visualized on the OsHV-1 NR-genome using Manhattan plots generated with the ggplot2 v3.4.0 package (Wickham, 2009). iSNVs were considered under selection if adjusted p-values exceeded the CMH threshold corresponding to a 5% FDR. Finally, allele frequency trajectories across generations were plotted for each replicate using the ggplot2 v3.4.0 package (Wickham, 2009), based on adjusted p-values and frequency data.

2.12.   Genotypes proportions

A Muller plot was used to visualize the emergence or disappearance of lineages specific to the origin of the initial viral suspension used to infect oysters. As previously described, the G0 viral inoculum was produced by pooling nine viral suspensions. To identify iSNVs specific to each origin, libraries from individuals sampled during the same mortality events within each basin were analyzed. Raw reads were filtered and trimmed as previously described, then aligned to the "artificial OsHV-1 major NR-consensus". Variants were called using Freebayes and assigned unique identifiers as described above (Figure 1).

Because each sample originated from a specific location, each detected iSNV could be linked to its geographical origin. iSNVs identified throughout the EE of OsHV-1 µVar were grouped by origin and

342 used to visualize their relative abundance using the MullerPlot package v0.13 (Farahpour et al., 2022)

343 in R (Figure 1).

3. Results

To investigate how OsHV-1 evolves during successive infection cycles, we conducted a

multigenerational experimental evolution using *M. gigas* oysters. Natural pools of viral populations

were used to infect specific-pathogen-free (SPF) oysters, which were subsequently collected from either

oyster farming (FA) or non-farming areas (NFA) along the French Atlantic coast. This protocol was

repeated over successive generations to simulate natural viral transmission under controlled

environmental conditions.

In each generation, moribund oysters were sampled, and intra-host viral diversity was assessed through

high-throughput sequencing. iSNVs were called to monitor changes in viral diversity and composition

over time and across different host population. Specifically, we examined patterns of iSNV

accumulation, substitution biases, and lineage dynamics between oyster generations.

3.1. Evolution of survival rates and viral load in Pacific Oyster spat infected with OsHV-

1 across generations

To assess the impact of OsHV-1 µVar infection on Pacific oyster spat, survival was monitored over 7

days post-infection (dpi) in both infected replicates and control tanks across three oyster populations.

No mortality was observed in control tanks (*i.e.* ASW-injected oysters) throughout the experiment.

Cohabitation-based infection depends on effective within-host viral propagation. Challenges in

maintaining reinfection across generations led to the termination of experimental evolution at the 13[th]

generation for the farming area (FA) population, the 18[th] for the control (C) population, and the 28[th] for

the non-farming area (NFA) population.

Mortality rates varied across generations but showed an overall decreasing trend (Figure 3A). At G0,

mortality reached 88% in FA, 96% in C, and 84% in NFA oyster populations. In the C population,

mortality decreased steadily to 42% at G15 before rising to 56% at G18, after which the experiment was

stopped due to the absence of mortality at G19.

In contrast, FA and NFA populations exhibited more variable dynamics. In FA oysters, mortality

declined to 58% at G5, then rose to 87% at G10, before falling sharply to 2%, which led to experimental

termination. NFA oysters followed a similar pattern, with mortality initially decreasing to 50% at G5,

peaking at 99% at G10, and subsequently fluctuating between 28% and 89%.

Viral DNA copy number also varied among individuals and across oyster populations (Figure 3B). In

FA oysters, the mean viral DNA copy number increased slightly over generations, rising from 8.38×10^5

$\pm 5.49 \times 10^5$ viral DNA copies/ng of DNA at G0 to $1.13 \times 10^6 \pm 4.81 \times 10^5$ viral DNA copies/ng of DNA at

G13 (Figure 3B). Conversely, C and NFA populations showed slight decreases in viral DNA copy

number across generations. In C oysters, viral DNA copy number decreased from $6.62 \times 10^5 \pm 4.49 \times 10^5$

viral DNA copies/ng of DNA at G0 to $3.34 \times 10^5 \pm 2.54 \times 10^5$ viral DNA copies/ng of DNA at G18. For

NFA oysters, viral DNA copy number decreased from $2.09 \times 10^6 \pm 1.20 \times 10^6$ viral DNA copies/ng of

DNA at G0 to $9.35 \times 10^5 \pm 1.11 \times 10^6$ viral DNA copies/ng of DNA at G28 (Figure 3B). Only selected

generations are displayed to provide a clear overview of temporal trends, and when experimental

evolution ended before G30 for a population, only the final available generation is shown.

ANOVA results revealed that infection generation had a significant effect on mortality ($F_{(29,36)} =$

2.798, $p = 0.002$, ges = 0.693), while oyster population had no effect ($F_{(2,63)} = 0.222$, $p = 0.802$, ges =

0.007) (Table 1). In contrast, viral DNA copy number was significantly influenced by oyster population

($F_{(2,216)} = 13.216$, $p < 0.001$, ges = 0.109), whereas the effect of infection generation approached but

did not reach significance ($F_{(8,210)} = 1.979$, $p = 0.051$, ges = 0.07). Together, these results indicate a

significant decline in mortality across infection generations and a significant difference in viral DNA

copy number among oyster populations.

Table 1. Summary of ANOVA results testing the effects of oyster population and infection
generation on mortality and viral load. Mortality and viral DNA copy number were analyzed as a
function of oyster population and infection generation. Significant effects were considered if $p < 0.05$.
Generalized eta-squared (ges) values are reported as effect size estimates. DFn: degrees of freedom
(numerator); DFd = degrees of freedom (denominator); F = F statistic; ges = generalized eta-squared
(effect size).

	DFn	DFd	F	p-value	Significance	ges
Mortalities ~ oyster populations	2	63	0.222	0.802		0.007
Mortalities ~ infection generations	29	36	2.798	0.002	*	0.693
Viral DNA copy number ~ oyster populations	2	216	13.216	3.85e-06	*	0.109
Viral DNA copy number ~ infection generations	8	210	1.979	0.051		0.07

Figure 3: Oyster mortalities and viral load in the three oyster populations at generations 0, 5, 10, 13, 15, 18, 20, 24, and 28. A) Percentage of mortality observed at each generation in oysters from the Farming Area (FA, blue), Control (C, orange), and Non-Farming Area (NFA, green) populations. Mortality was monitored daily, and the proportion of dead oysters per generation is represented by solid lines. Dashed lines indicate the linear regression trend for each population across generations. Asterisks denote the final generation sequenced for each population. B) Individual viral DNA loads measured in infected oysters, expressed as $\log_{10}$-transformed viral DNA copy number. Viral loads were quantified from seven oysters per replicate per generation. Dots represent individual oysters, while box plots show the median, interquartile range, and variability within each group. Dotted lines connect mean viral loads across generations for each population

3.2. Dynamics of substitutions and transition:transversion ratio

Studying substitution patterns across viral generations in EE is crucial to understand evolutionary dynamics, genetic diversity, and adaptation (Bromham, 2020). Based on the iSNVs identified from all sequencing libraries, we examined substitution dynamics across viral generations and oyster populations by analyzing ancestral iSNVs - defined as iSNVs already present in the initial inoculum - and *de novo* iSNVs - defined as iSNVs arising during the experimental evolution process - counts and transition:transversion (ts:tv) ratios. A clear trend emerges across all three oyster populations, showing a rapid decline in the number of ancestral iSNVs in the OsHV-1 µVar genome over successive infection generations (Figure 4A). This decline is most pronounced in the FA population, with iSNVs counts dropping from 346 ± 26 iSNVs at G0 to 139 ± 59 iSNVs at G13 (Figure 4A). The number of ancestral iSNVs remains comparatively high in the NFA population (*e.g.* 329 ± 53 at G0 and 236 ± 12 at G28), with a slight increase observed at G10 (Figure 4A). *De novo* substitutions were systematically fewer than ancestral ones (*i.e.* typically fewer than 40 per generation across populations) and their numbers increased over successive infection generations (Figure 4C). For all three populations, most *de novo* iSNVs arise between G0 and G5, and remain relatively constant in subsequent generations.

A statistical analysis of the identified iSNVs was performed to assess the ts:tv ratio (Figure 4B and 4D). For ancestral iSNVs, the ts:tv ratio remained consistently between 1 and 0.4 across generations and populations, indicating a slight excess of transitions. The only exception was the FA population, where the ratio reached 1.26 at G0, reflecting a stronger transition bias, and then declined to below 0.4 from G5 onward, indicating a shift toward an excess of transversions (Figure 4B). In addition, the ts:tv ratio tended to decrease over generations in both the FA and C populations. For the NFA population, although the overall trend also points to a decline in the ts:tv ratio, a temporary increase was observed at G10. The ts:tv ratio for *de novo* iSNVs shows an almost opposite trend compared with ancestral iSNVs. In the FA population, values range from 0 to 0.4, whereas in the C population they span from 0.4 to 2.5, and from 0.4 to 1 in the NFA population (Figure 4D). Overall, the ratio tends to increase between G0 and G5 or G10, followed by a subsequent decline.

Figure 4 : Substitution types and transition:transversion (ts:tv) ratio observed within OsHV-1 populations.
A) Mean number of ancestral substitution types per generation and oyster population. Transitions (pyrimidine ↔ pyrimidine or purine ↔ purine) are shown in red to orange, and transversions (purine ↔ pyrimidine) in blue to green. Numbers above stacked bars indicate counts of transitions and transversions while numbers in brackets represent the mean per substitution type across samples. B) Variation of the ts:tv ratio across generations for each oyster population. Line colors correspond to the different oyster populations. C) Mean number of de novo substitution types per generation and oyster population. Colors and annotations follow the same scheme as in panel A. D) Variation of the ts:tv ratio for de novo substitutions across generations and populations.

3.3. Unveiling OsHV-1 population structure and anomalies in experimental evolution through iSNVs data analysis and principal component projection

To investigate viral population structure across generations and oyster populations, we performed a Principal Component Analysis (PCA) on iSNVs frequency data, incorporating metadata on host population, infection generation, and replicates (Figure 5). To address potential concerns about restricting the analysis to only two components, we further extended the PCA to the first 25 components (Figure S1). This extended analysis confirms that the major clustering patterns observed in Figure 5 are consistent with the broader genomic variation, while also showing that subsequent components contribute relatively little to the explained variance. We therefore retained the first two components in the main figure for clarity, as they best capture the main biological signals of infection progression and host population differences. The first principal component accounted for 12.61% of the variance, while the second component explains 9.89% of the variance (Figure 5). The first component reflects infection progression over generations, whereas the second is associated with oyster population or susceptibility traits. The plot reveals four distinct clusters corresponding to the interplay between oyster populations and infection generations.

The first cluster includes samples from all oyster populations at G0, along with those from replicate 1 of the C population at G0, G5, G10, G15, and G18 (Figure 5, group 1) suggesting that the initial viral diversity was maintained across generations in replicate 1 of the C population.

The second cluster (Figure 5, group 2) includes samples mainly from the FA population at G5, G10, and G13, as well as from replicate 2 of the C population at G5, G10, G15, and G18.

Finally, the third cluster comprises samples from the NFA population, which includes samples from G15 and G28 (Figure 5).

Figure 5 : Principal Component Analysis (PCA) of OsHV-1 genetic variation among viral populations from different oyster populations and infection generations. PCA was performed on iSNVs detected within OsHV-1 genomes to investigate patterns of genetic differentiation among viral populations and infection generations. Each point represents an individual viral genome, projected onto the first two principal components (PC1 and PC2). Colors indicate the host oyster population: blue for FA (Farming Area), orange for C (Control), and green for NFA (Non-Farming Area). Replicates are distinguished by point shapes (squares for replicate 1 and triangles for replicate 2), and infection generations are represented by a color gradient. The ellipses delineate three major clusters of genetically related viral populations, labeled Clusters 1 to 3.

3.4. Detection of regions under selection pressure and allelic frequency trajectories

To detect genomic regions under selection, we applied the Cochran-Mantel-Haenszel (CMH) test independently to iSNVs identified in each oyster population, focusing on allele frequency changes between the initial and evolved generations. The CMH test is widely used to detect selection pressures by analyzing allele frequency shifts in experimental evolution and time-series data (Spitzer et al., 2020).

The CMH test, applied to iSNVs identified in each viral population from the three oyster populations independently (Figure 6), showed that no genomic regions were significantly affected by selection in the FA (Figure 6A) and C (Figure 6B) populations. In contrast, the NFA population exhibited 117 candidate iSNVs under selection (Figure 6C). These iSNVs were distributed across the OsHV-1

genome, with a concentration in repeated regions and in the U_L region between 50 kb and 80 kb. More

precisely, 28 potentially selected positions were located in intergenic regions, including one in the stem

loop, and 89 iSNVs were located within ORFs (Figure 6C, Table S3).

Among the 89 iSNVs located within ORFs, 58 were found in ORFs of unknown function, and 12 in

ORFs encoding transmembrane proteins (Table S3). In addition, 8 iSNVs were located in ORFs

encoding secreted proteins, four in ORF 100 coding for a DNA polymerase, three within ORFs encoding

Zinc-finger, Ring-type proteins, two within ORFs encoding BIR repeat motifs, one within ORF115

associated with an Origin-binding protein, and the final iSNV within ORF20 encoding a ribonucleotide

reductase (Table S3).

Figure 6: Genomic distribution of candidate iSNVs under selection in oyster populations. The scatter plots show the negative log10-transformed p-values of iSNVs along the OsHV-1 genome for A) Farming Area, B) Control and C) Non-Farming Area experimental oyster populations. Each point represents an iSNV identified by comparing the viral populations from the founder and evolved generations using the Cochran–Mantel–Haenszel (CMH) test. The red dashed line indicates the significance threshold corresponding to a 5% false discovery rate (CMH cutoff). iSNVs above this threshold are considered candidates for selection and are color-coded based on their potential functional impact. The OsHV-1 genome structure is shown below the plots, including unique regions (U_L and U_S) and inverted repeat regions (IR_L and IR_S). Colored blocks correspond to Open Reading Frames (ORFs) containing candidate iSNVs, with colors reflecting the predicted function or protein domain of each ORF.

An allele frequency trajectory (AFT) plot was used to track changes in the frequencies of candidate

iSNVs identified in the NFA oyster population, providing insights into evolutionary dynamics and

signals of positive selection (Barghi et al., 2020). The analysis was conducted separately for both

replicates (Figure 7).

The AFT plot shows a general trend of increasing allele frequencies, often leading to fixation of selected

alleles in both replicates. In replicate 1, allele frequencies of candidate iSNVs rise from G0 to G5, drop

between G5 and G10, and finally increase again between G10 and G15 until fixation (Figure 7A). A

similar trend is observed in replicate 2, but only for 77 iSNVs (Figure 7B). Moreover, 66 iSNVs in

replicate 1 and 26 iSNVs in replicate 2, shown at the bottom of the plot, also exhibit increasing

frequencies, though to a lesser extent (Figure 7A and B).

Figure 7: Temporal trajectories of allelic frequencies for top 100 candidate OsHV-1 iSNVs during serial infections in the NFA oyster population. Each blue line represents the allelic frequency trajectory of a specific SNP across generations. Red line represents the median allelic frequency trajectory.

3.5. Relative abundance of iSNVs from specific origin

483   To monitor the persistence and dynamics of viral lineages from specific origins across generations of

484   infection, a Muller plot was generated (Figure 8). The results show that multiple lineages from different

origins coexist from the start to the end of the experiment within each oyster population. At generation

0, genotypes specific to MO and LEU had a higher relative abundance (RA) than those from ARC and

BR. This general trend was maintained across generations of infections in all oyster populations, though

RA varied depending on the population. In the FA population, OsHV-1 iSNVs specific to MO are

present at a RA of 0.2 at G0, which increased to 0.4 by G10 then slightly declined. A similar trend was

observed for iSNVs from ARC and BR, but with much lower RA values (between 0.02 and 2.0). In

contrast, the RA of LEU-specific iSNVs declined from 0.4 at G0 to 0.1 at G5.

New iSNVs, absent from all initial viral suspensions, began to emerge during the experiment. At G0,

their RA was low in both farming and non-farming populations (RA = 0.1), but increased over

generations. In contrast, in the control population, the RA of these "other" iSNVs started at 0.3, dropped

to 0.2 by G5, rose to 0.5 at G10, then declined again to 0.2 by G15.

Figure 8 : Muller plot representing the relative abundance of viral iSNVs collected from specific origins for each oyster population during the experimental evolution. The relative abundance of viral SNPs is color-coded according to their geographical origin of the viral suspension: the orange area corresponds to genotypes collected specifically in MO, the red area corresponds to genotypes collected specifically in BR, the blue area corresponds to genotypes collected specifically in ARC, the green area corresponds to genotypes collected specifically in LEU, and the grey area corresponds to genotypes that appeared during the experimental evolution.

4. Discussion

To mitigate the impact of infectious diseases, it is essential to understand the mechanisms shaping the

evolution and spread of pathogenic microorganisms (Gu et al., 2021; NIH, 2007). Variation in host

resistance plays a major role in driving these processes (Gandon & Michalakis, 2000; Kubinak & Potts,

2013). This study provides valuable insights into the evolutionary dynamics of OsHV-1 and its

interactions with Pacific oysters, focusing on how the virus adapts and diversifies in response to host

genetic backgrounds. These findings are crucial to address the devastating impact of OsHV-1 on oyster

farming.

During this experimental evolution, we observed a decrease in both oyster mortalities and viral load

across successive generations of OsHV-1 infection. Interestingly, despite the distinct infection histories

of the three oyster populations, mortality patterns were unexpectedly similar during the early generations

of infection, contrary to the anticipated gradient. This convergence in mortality rates suggests that the

initial infection dynamics may have been strongly influenced by experimental conditions rather than by

prior host exposure history. For instance, standardized infection doses and cohabitation settings could

have minimized differences in host susceptibility or viral virulence between populations. The decline

in mortalities and viral load results from several, non-exclusive factors. First, it can be explained by the

selection of less virulent viral lineages. Host-pathogen interactions are shaped by a balance between

virulence and transmission (Anderson & May, 1982). In conditions with limited host numbers, such as

in our setup with 25 oysters per tank, rapid host death could threaten viral persistence. A plausible

strategy for the virus may be to reduce virulence, and maintaining a stable host population while

ensuring replication. This trade-off between virulence and transmission is well-documented, with higher

transmission usually associated with lower virulence, and conversely, excessive virulence reduces

transmission (Anderson & May, 1982; Gandon & Michalakis, 2000; Kun et al., 2023). Comparable

results were reported in EE study involving tobacco etch potyvirus (TEV) in plants, where viral load

and virulence decreased across generations, strongly influenced by host lineage (Cuevas et al., 2015).

However, virulence does not necessarily decrease during host–virus co-evolution. In some systems,

virulence has increased over time, as observed in Marek's disease virus (Trimpert et al., 2017, 2019),

illustrating that the direction of virulence evolution is context-dependent.

Secondly, the reduction in mortality rates can be attributed to a lack of complementation between

lineages (Froissart et al., 2004; Montville et al., 2005). Host infection rarely occurs with a single lineage,

but often involves co-infection, with multiple viral lineages contributing to the infection and associated

symptoms. Certain lineages carrying disadvantageous mutations can be complemented by co-infecting high-fitness lineages, allowing low-fitness lineages to gain a phenotypic advantage (Froissart et al., 2004; Montville et al., 2005). In our study, where genetic diversity has likely been drastically reduced, possibly due to a bottleneck effect, some lineages may have been purged, limiting complementation and reducing virulence over successive generations. Bottlenecks can drastically reduce genetic diversity by eliminating rare variants and fixing certain alleles through genetic drift (Zwart & Elena, 2015). In viral populations, this can compromise cooperative interactions between lineages, diminishing overall viral fitness and reduction in mortality rates (Zwart & Elena, 2015).

A third hypothesis to explain declines in mortality rates and viral loads is a temporal shift in the peak of viral particle release into seawater across infection generations. OsHV-1 excretion peaks between 24 and 48 hours post-infection (hpi) (Delmotte et al., 2020), but timing can vary with oyster susceptibility, initial viral load, donor numbers and age, food availability, water temperature, and salinity (Dégremont, 2011; Delmotte et al., 2020; Evans et al., 2017; Pernet et al., 2018, 2018; Petton et al., 2015; Schikorski et al., 2011). Consequently, the viral load would be high when cohabitation begins and progressively decreasing over time. In our experiment, successive infections involved oysters from the same cohort, whose age increased across generations, potentially altering susceptibility and timing of viral excretion peak. Cohabitation infections can also lead to fewer fatalities due to progressive decline in viral load across cycles (Cain et al., 2021). While it may not be the optimal method for EE, successive infection by cohabitation was chosen to mimic the natural infection of oysters by OsHV-1 in the field.

A fourth complementary hypothesis relates to latent or persistent OsHV-1 infections. OsHV-1 may establish a latent or persistent state in oysters, allowing persistence without acute disease (Divilov et al., 2024), as seen in other herpesviruses (Dotto-Maurel et al., 2025; White et al., 2012). FA oysters indeed exhibited high viral copy numbers even during low mortality periods, suggesting a shift toward non-lytic or less virulent states, potentially mediated by host immune regulation or viral adaptation favoring long-term coexistence (Divilov et al., 2024). In contrast, the C population show lower viral loads than FA population indicating that viral replication efficiency, rather than immediate host mortality, can vary among populations. These patterns may reflect differences in host immune responses or in the balance

between lytic and persistent infection. Although our experimental design did not include latency-specific

assays, these observations highlight the need to further investigate the latent potential of OsHV-1 under

recurrent exposures and genetically selected oyster populations.

Genetic mutations, including transitions and transversions, shape viral evolution by contributing to

genetic diversity and adaptation and molecular evolutionary hypothesis suggests that natural selection

promotes amino acid substitutions mostly through transitions (Stoltzfus & Norris, 2016). The decline in

intra-individual single nucleotide variants (iSNVs) and in the ancestral transition:transversion (ts:tv)

ratio across generations suggest progressive genetic homogenization. Nevertheless, *de novo* iSNVs

increased sharply at G5 before stabilizing, indicating ongoing mutation emergence despite reduced

overall diversity. Interpretation must consider potential technical artefacts, including sequencing errors

or library preparation bias (e.g. oxidative DNA damage generate G|T and C|A transversions), low

frequency false-positive calls particularly for viral quasispecies, or misalignments in repetitive, low

complexity of poor mapability genomic regions can produce spurious iSNVs, often enriched in

transversions (Jensen et al., 2021; Lenz et al., 2014; López-Muñoz et al., 2021; Phan et al., 2015;

Poetsch, 2020; Van Poelvoorde et al., 2022). Biologically, the fidelity of the OsHV-1 DNA polymerase,

as well as host-mediated processes such as oxidative stress or deamination, could also favor

transversions (Mourier et al., 2021). Natural selection may further modulate which lineages persist.

Analyses based on a global consensus genome (i.e. 126 viral NR-genomes) may additionally introduce

reference biases affecting iSNVs detection. Indeed, the ts:tv ratios observed here are lower than those

reported for humans (~2) or for Varicella-Zoster virus (~2.3), consistent with technical influences, but

they may also reflect virus- or host-specific mutation biases (Jeon et al., 2016; Wang et al., 2015). These

observations align with patterns in other viral systems, where bottlenecks and genetic drift or selective

sweeps reduce diversity across generations (Kutnjak et al., 2017; Li & Roossinck, 2004). Given the

context of OsHV-1 infecting oyster of distinct genetic backgrounds, a combination of technical artefacts,

host-mediated processes, and genuine evolutionary bottlenecks likely contributed to reduced viral

lineages.

PCA analysis of iSNV frequencies revealed clusters corresponding to oyster populations and infection

generations, indicating genetic differentiation and adaptability. Notably, clustering of NFA and FA

individuals was clearly delineated, while control oysters exhibited intermediate clustering, suggesting

susceptibility-dependent evolution (Kubinak & Potts, 2013). Within the NFA oyster population, distinct

clusters emerged, indicating evolutionary changes across infection generations in response to selective

pressures.

The dynamics of iSNVs revealed clear differences among viral lineages of distinct geographic origins,

highlighting how host environment and exposure history shape OsHV-1 diversity and persistence.

Throughout the experiment, iSNVs from MO remained dominant in all oyster populations, particularly

in those from FA and in control oysters, whereas iSNVs originating from ARC progressively increased

in relative abundance within NFA oysters. This contrasting pattern likely reflects both the

epidemiological history and ecological context of the host populations. All viral lineages used in the

experiment originated from farming areas and were initially sampled during mortality events, suggesting

that NFA oysters represent the most geographically and epidemiologically distant hosts from these viral

sources. The expansion of ARC-like iSNVs in NFA oysters could indicate that these hosts, having

experienced limited or no prior exposure to OsHV-1 lineages from ARC, and impose weaker selective

constraints on viral replication. Such relaxed selection may allow for the persistence or even

amplification of lineages that are not favored in aquaculture environments, where repeated exposure and

strong host-virus coevolution tend to filter viral diversity (Kennedy et al., 2016). Conversely, the

predominance of MO-like iSNVs in FA oysters likely results from repeated selective pressures imposed

by ongoing viral circulation in dense farming contexts, favoring lineages already adapted to these hosts

(Kennedy et al., 2016).

Interestingly, new iSNVs that were absent from all initial inocula also emerged during the course of the

experiment, suggesting ongoing within-host diversification and potential *de novo* mutation events. The

accumulation of these new variants, particularly in NFA oysters, may reflect broader environmental

tolerance or increased mutational persistence under relaxed selective pressure. In contrast, the lower

diversity observed in FA oysters supports the idea that repeated exposure to similar viral lineages and

stronger immune selection in farmed populations drive the convergence toward dominant MO-like
lineages.

The identification of 117 candidate selection signatures in the OsHV-1 genomes and allelic frequency
trajectories (AFT) plots in the NFA population, provides strong evidence of selective pressures during
experimental evolution. Allele frequency increases and fixation indicate positive selection (Barghi et
al., 2020). Selection was detected exclusively in NFA oysters, suggesting enhanced viral replication that
may increase mutation rates and facilitate the fixation of advantageous alleles in these more susceptible
hosts (Dégremont, 2011; Elena & Sanjuán, 2005; Peck & Lauring, 2018). Conversely, no selection
signals were detected in FA oysters, despite their reduced viral diversity, supporting the influence of
bottleneck and drift effects. Given that viral evolution are typically shaped by environmental changes
and host immune pressures acting as selective filters to constrain viral diversity (Kubinak & Potts, 2013),
this absence of detectable selection in FA oysters is unexpected, suggesting that stochastic processes
such as bottlenecks and genetic drift dominated viral evolution in these populations. This may result
from the repeated transmission and amplification of a limited number of viral lineages within farming
environments, reducing opportunities for adaptive lineages to emerge and spread.

Selection signatures were distributed across the viral genome, particularly within repetitive regions and
in the U_L region. Such a widespread pattern likely reflects a combination of immune-driven and relaxed
selection processes acting in under contrasting host environments. According to the "relaxed selection
hypothesis" (Mikonranta et al., 2012), a neutral virulence-related trait in environments where host
density and immune pressure are low, lead to alteration or loss through the random accumulation of
mutations rather than active negative selection. In the context of OsHV-1, NFA oysters, characterized
by reduced host density and weaker immune activation, may thus favor the persistence of neutral or
mildly deleterious mutations, potentially giving rise to attenuated viral lineages. This relaxation of
selective constraints could therefore contribute to the progressive decline in mortality rates observed
during the experiment, acting in parallel with adaptive processes occurring in farming environments.

These selection signals were further enriched in ORFs encoding proteins critical for viral attachment,
replication, packaging, and membrane functions. Viral entry proteins, even with single amino acid

changes, can alter infection success across hosts with different susceptibilities (van Sluijs et al., 2017). Acquired, duplicated or lost domains in Herpesviridae evolution, including envelope (such as membrane glycoproteins and transmembrane receptors), auxiliary domains (like Zinc-finger, RING type, dinucleoside kinase), and modulation domains (such as Interleukin, Interferon-regulatory factor, or Zinc-finger), enhance specific binding and immune evasion (Brito & Pinney, 2020). In OsHV-1, most selection signals were within these domains, indicating adaptation of essential proteins to host genetic backgrounds. Two previous studies support immune-driven adaptation in OsHV-1 across host species and environments (Delmotte et al., 2020; Pelletier et al., 2025). Based on the results obtained in this study, it is evident that adaptation can also occur at a finer scale within the host's genetic background, which is influenced in this case by the geographic origin of the parental individuals (*i.e.* farmed or not farmed area).

5. Conclusion

This study provides valuable insights into the evolutionary driving mechanisms of OsHV-1 and its interactions with Pacific oysters, with a particular emphasis on viral adaptation and diversification in response to host infection susceptibility. Employing an experimental evolution approach, it sheds light on the dynamic nature of host-virus interactions, the potential for viral adaptation, and the possible role of genetic diversity in shaping the outcome of these interactions. The nucleotide evolution of OsHV-1, particularly a decrease in transition and transversion numbers, appears to be mainly driven by both genetic drift and positive selection linked to the oyster genetic background, resulting in a reduction in viral diversity and the fixation of specific alleles within ORFs dedicated to host-virus interactions and virus functional maintenance. Further research into the functional consequences of these changes is necessary. These findings contribute to our understanding of the mechanisms governing viral evolution and hold significant implications for mitigating the devastating impacts of OsHV-1 on oyster farming industries. Future studies should continue to explore the intricate dynamics of host-virus interactions and their implications for viral evolution and disease management.

Data availability

Raw data have been deposited on the SRA database under Bioproject PRJNA1216400 accession numbers SAMN46433918 to SAMN46434013 for future reference and accessibility (Table S2). All the scripts are available at https://gitlab.ifremer.fr/asim/experimental_evolution_public.

Acknowledgment

We thank the staff of the Ifremer station at La Tremblade (ASIM), Frédéric Girardin and his team (Plateforme des Mollusques Marins de La Tremblade PMMLT), and Virginie François and his team (Plateforme des Mollusques Marins de Bouin PMMB). We also thank the SEBIMER team for maintaining bioinformatics tools.

C.P. was financially supported by a grant from the Ifremer Scientific Board and the Nouvelle-Aquitaine region. The present study was supported by scientific direction of Ifremer in context of the GT Huître project and by the FEAMP Gestinnov (FFEA470020FA1000007) GESTINNOV project and by DGAL (French General Directorate for Food) through the National Reference Laboratory for Mollusc Diseases and by the European-Union Reference Laboratory for Mollusc Diseases, Ifremer, La Tremblade. The authors acknowledge the Pôle de Calcul et de Données Marines (PCDM; https://wwz.ifremer.fr/en/Research-Technology/Research-Infrastructures/Digital-infrastructures/Computation-Centre) for providing DATARMOR computing and storage resources. The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

C.P. and co-authors thank J.M.E. for the discussions and exchanges. We would like to thank Bruno Petton and his team PHYTNESS for sampling the virus in the Brittany area, as well as Johan Vieira and the CAPENA team for sampling OsHV-1 in Arcachon Bay. B.M., N.F., and C.P. designed the study; L.D., B.M., and J.V.D. provided the biological materials; N.F., M.H., M.M., and C.P. collected and processed the samples. C.P. performed the bioinformatics analysis of the Illumina data.

687   B.M., G.C., M.J., J.V.D., L.D., and C.P. drafted the manuscript. All authors read and approved the final

688   version of the manuscript.

689   We declare that we have no competing interests.

References

Abbadi, M., Zamperin, G., Gastaldelli, M., Pascoli, F., Rosani, U., Milani, A., Schivo, A., Rossetti, E.,

Turolla, E., Gennari, L., Toffan, A., Arcangeli, G., & Venier, P. (2018). Identification of a newly

described OsHV-1 µvar from the North Adriatic Sea (Italy). *Journal of General Virology*, *99*(5),

Article 5. https://doi.org/10.1099/jgv.0.001042

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment

search tool. *Journal of Molecular Biology*, *215*(3), Article 3. https://doi.org/10.1016/S0022-

2836(05)80360-2

Anderson, R. M., & May, R. M. (1982). Coevolution of hosts and parasites. *Parasitology*, *85*(2), 411–

426. https://doi.org/10.1017/S0031182000055360

Azéma, P., Lamy, J.-B., Boudry, P., Renault, T., Travers, M.-A., & Dégremont, L. (2017). Genetic

parameters of resistance to Vibrio aestuarianus, and OsHV-1 infections in the Pacific oyster,

Crassostrea gigas, at three different life stages. *Genetics, Selection, Evolution : GSE*, *49*, 23.

https://doi.org/10.1186/s12711-017-0297-2

Azéma, P., Maurouard, E., Lamy, J.-B., & Dégremont, L. (2017). The use of size and growing height to

improve Crassostrea gigas farming and breeding techniques against OsHV-1. *Aquaculture*, *471*,

121–129. https://doi.org/10.1016/j.aquaculture.2017.01.011

Bankevich, A., Nurk, S., Antipov, D., Gurevich, A. A., Dvorkin, M., Kulikov, A. S., Lesin, V. M.,

Nikolenko, S. I., Pham, S., Prjibelski, A. D., Pyshkin, A. V., Sirotkin, A. V., Vyahhi, N., Tesler,

G., Alekseyev, M. A., & Pevzner, P. A. (2012). SPAdes: A New Genome Assembly Algorithm

and Its Applications to Single-Cell Sequencing. *Journal of Computational Biology*, *19*(5),

Article 5. https://doi.org/10.1089/cmb.2012.0021

Barghi, N., Hermisson, J., & Schlötterer, C. (2020). Polygenic adaptation: A unifying framework to

understand positive selection. *Nature Reviews Genetics*, *21*(12), 769–781.

https://doi.org/10.1038/s41576-020-0250-z

Boudry, P., Collet, B., Cornette, F., Hervouet, V., & Bonhomme, F. (2002). *High variance in*

reproductive success of the Pacific oyster žCrassostrea gigas, Thunberg/ revealed by

microsatellite-based parentage analysis of multifactorial crosses.

Bromham, L. (2020). *Substitution Rate Analysis and Molecular Evolution.*

Burioli, E. A. V., Prearo, M., & Houssin, M. (2017). Complete genome sequence of Ostreid herpesvirus

type 1 μVar isolated during mortality events in the Pacific oyster Crassostrea gigas in France

and Ireland. *Virology*, *509*, 239–251. https://doi.org/10.1016/j.virol.2017.06.027

Cain, G., Liu, O., Whittington, R. J., & Hick, P. M. (2021). Reduction in Virulence over Time in Ostreid

herpesvirus 1 (OsHV-1) Microvariants between 2011 and 2015 in Australia. *Viruses*, *13*(5), 946.

https://doi.org/10.3390/v13050946

Chen, S., Zhou, Y., Chen, Y., & Gu, J. (2018). fastp: An ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics, *34*(17), Article 17. https://doi.org/10.1093/bioinformatics/bty560

Cuevas, J. M., Willemsen, A., Hillung, J., Zwart, M. P., & Elena, S. F. (2015). Temporal Dynamics of

Intrahost Molecular Evolution for a Plant RNA Virus. *Molecular Biology and Evolution*, *32*(5),

1132–1147. https://doi.org/10.1093/molbev/msv028

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane,

T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools and BCFtools.

GigaScience, *10*(2), giab008. https://doi.org/10.1093/gigascience/giab008

Davison, A. J., Trus, B. L., Cheng, N., Steven, A. C., Watson, M. S., Cunningham, C., Deuff, R.-M. L.,

& Renault, T. (2005). A novel class of herpesvirus with bivalve hosts. *Journal of General*

Virology, *86*(1), Article 1. https://doi.org/10.1099/vir.0.80382-0

de Lorgeril, J., Lucasson, A., Petton, B., Toulza, E., Montagnani, C., Clerissi, C., Vidal-Dupiol, J.,

Chaparro, C., Galinier, R., Escoubas, J.-M., Haffner, P., Dégremont, L., Charrière, G. M.,

Lafont, M., Delort, A., Vergnes, A., Chiarello, M., Faury, N., Rubio, T., … Mitta, G. (2018).

Immune-suppression by OsHV-1 viral infection causes fatal bacteraemia in Pacific oysters.

Nature Communications, *9*(1), Article 1. https://doi.org/10.1038/s41467-018-06659-3

Dégremont, L. (2011). Evidence of herpesvirus (OsHV-1) resistance in juvenile Crassostrea gigas

selected for high resistance to the summer mortality phenomenon. *Aquaculture*, *317*(1–4), 94–

98. https://doi.org/10.1016/j.aquaculture.2011.04.029

Dégremont, L., Bédier, E., Soletchnik, P., Ropert, M., Huvet, A., Moal, J., Samain, J.-F., & Boudry, P.

(2005). Relative importance of family, site, and field placement timing on survival, growth, and

yield of hatchery-produced Pacific oyster spat (Crassostrea gigas). *Aquaculture*, *249*(1), 213–

229. https://doi.org/10.1016/j.aquaculture.2005.03.046

Dégremont, L., Ernande, B., Bédier, E., & Boudry, P. (2007). Summer mortality of hatchery-produced

Pacific oyster spat (Crassostrea gigas). I. Estimation of genetic parameters for survival and

growth. *Aquaculture*, *262*(1), 41–53. https://doi.org/10.1016/j.aquaculture.2006.10.025

Dégremont, L., Maurouard, E., Ledu, C., & Benabdelmouna, A. (2019). Synthesis of the "PLAN DE

SAUVEGARDE" using selected all-triploid oysters to reduce the shortage of spat in France due

to OsHV-1–associated mortality in Crassostrea gigas. *Aquaculture*, *505*, 462–472.

https://doi.org/10.1016/j.aquaculture.2019.03.014

Delmotte, J., Chaparro, C., Galinier, R., de Lorgeril, J., Petton, B., Stenger, P.-L., Vidal-Dupiol, J.,

Destoumieux-Garzon, D., Gueguen, Y., Montagnani, C., Escoubas, J.-M., & Mitta, G. (2020).

Contribution of Viral Genomic Diversity to Oyster Susceptibility in the Pacific Oyster Mortality

Syndrome. *Frontiers in Microbiology*, *11*. https://doi.org/10.3389/fmicb.2020.01579

Delmotte, J., Pelletier, C., Morga, B., Galinier, R., Petton, B., Lamy, J.-B., Kaltz, O., Avarre, J.-C.,

Jacquot, M., Montagnani, C., & Escoubas, J.-M. (2022). Genetic diversity and connectivity of

the Ostreid herpesvirus 1 populations in France: A first attempt to phylogeographic inference

for a marine mollusc disease. *Virus Evolution*, *8*(1), veac039.

https://doi.org/10.1093/ve/veac039

Divilov, K., Wang, X., Swisher, A. E., Yeoman, P. C., Rintoul, M., Fleener, G. B., Schoolfield, B.,

Langdon, C., & Jin, L. (2024). Ostreid herpesvirus 1 latent infection and reactivation in adult

Pacific oysters, *Crassostrea gigas*. *Virus Research*, *339*, 199245.

https://doi.org/10.1016/j.virusres.2023.199245

Dotto-Maurel, A., Arzul, I., Morga, B., & Chevignon, G. (2025). Herpesviruses: Overview of

systematics, genomic complexity and life cycle. *Virology Journal*, *22*(1), 155.

https://doi.org/10.1186/s12985-025-02779-7

Dotto-Maurel, A., Pelletier, C., Morga, B., Jacquot, M., Faury, N., Dégremont, L., Bereszczynki, M.,

Delmotte, J., Escoubas, J.-M., & Chevignon, G. (2022). Evaluation of tangential flow filtration

coupled to long-read sequencing for ostreid herpesvirus type 1 genome assembly. *Microbial*

Genomics, *8*(11). https://doi.org/10.1099/mgen.0.000895

Elena, S. F., & Sanjuán, R. (2005). Adaptive Value of High Mutation Rates of RNA Viruses: Separating

Causes   from   Consequences.   *Journal   of   Virology*,   *79*(18),   11555–11558.

https://doi.org/10.1128/JVI.79.18.11555-11558.2005

Evans, O., Hick, P., & Whittington, R. J. (2017). Detection of Ostreid herpesvirus -1 microvariants in

healthy Crassostrea gigas following disease events and their possible role as reservoirs of

infection.   *Journal   of   Invertebrate   Pathology*,   *148*,   20–33.

https://doi.org/10.1016/j.jip.2017.05.004

Farahpour, F., Saeedghalati, M., & Hoffmann, D. (2022). *MullerPlot: Generates Muller Plot from*

Population/Abundance/Frequency Dynamics Data. R package version 0.1.3. (Version 0.1.3)

[Computer software].

Froissart, R., Wilke, C. O., Montville, R., Remold, S. K., Chao, L., & Turner, P. E. (2004). Co-infection

Weakens Selection Against Epistatic Mutations in RNA Viruses. *Genetics*, *168*(1), 9–19.

https://doi.org/10.1534/genetics.104.030205

Fuandila, N. N., Gosselin-Grenet, A.-S., Tilak, M.-K., Bergmann, S. M., Escoubas, J.-M., Klafack, S.,

Lusiastuti, A. M., Yuhana, M., Fiston-Lavier, A.-S., Avarre, J.-C., & Cherif, E. (2022).

Structural variation turnovers and defective genomes: Key drivers for the in vitro evolution of

the large double-stranded DNA koi herpesvirus (KHV). *Peer Community Journal*, *2*, e44.

https://doi.org/10.24072/pcjournal.154

Gandon, S., & Michalakis, Y. (2000). Evolution of parasite virulence against qualitative or quantitative

host   resistance.   *Proceedings.   Biological   Sciences*,   *267*(1447),   985–990.

https://doi.org/10.1098/rspb.2000.1100

Garrison, E., & Marth, G. (2012). Haplotype-based variant detection from short-read sequencing.
arXiv:1207.3907 [q-Bio]. http://arxiv.org/abs/1207.3907

Gawra, J., Valdivieso, A., Roux, F., Laporte, M., de Lorgeril, J., Gueguen, Y., Saccas, M., Escoubas, J.-
M., Montagnani, C., Destoumieux-Garzón, D., Lagarde, F., Leroy, M. A., Haffner, P., Petton,
B., Cosseau, C., Morga, B., Dégremont, L., Mitta, G., Grunau, C., & Vidal-Dupiol, J. (2023).
Epigenetic variations are more substantial than genetic variations in rapid adaptation of oyster
to Pacific oyster mortality syndrome. *Science Advances*, *9*(36), eadh8990.
https://doi.org/10.1126/sciadv.adh8990

Gu, J., Zhou, Z., & Wang, Y. (2021). Editorial: Evolutionary Mechanisms of Infectious Diseases.
Frontiers in Microbiology, *12*. https://doi.org/10.3389/fmicb.2021.667561

Jensen, A., Lillie, M., Bergström, K., Larsson, P., & Höglund, J. (2021). Whole genome sequencing
reveals high differentiation, low levels of genetic diversity and short runs of homozygosity
among Swedish wels catfish. *Heredity*, *127*(1), 79–91. https://doi.org/10.1038/s41437-021-
00438-5

Jeon, J. S., Won, Y. H., Kim, I. K., Ahn, J. H., Shin, O. S., Kim, J. H., & Lee, C. H. (2016). Analysis of
single nucleotide polymorphism among Varicella-Zoster Virus and identification of vaccine-
specific sites. *Virology*, *496*, 277–286. https://doi.org/10.1016/j.virol.2016.06.017

Kassambara, A., & Mundt, F. (2020). *Extract and Visualize the Results of Multivariate Data Analyses*
[R package factoextra v1.0.7]. https://rpkgs.datanovia.com/factoextra/index.html

Katoh, K., Misawa, K., Kuma, K., & Miyata, T. (2002). MAFFT: A novel method for rapid multiple
sequence alignment based on fast Fourier transform. *Nucleic Acids Research*, *30*(14), Article
14. https://doi.org/10.1093/nar/gkf436

Kawecki, T. J., Lenski, R. E., Ebert, D., Hollis, B., Olivieri, I., & Whitlock, M. C. (2012). Experimental
evolution. *Trends in Ecology & Evolution*, *27*(10), 547–560.
https://doi.org/10.1016/j.tree.2012.06.001

Kennedy, D. A., Kurath, G., Brito, I. L., Purcell, M. K., Read, A. F., Winton, J. R., & Wargo, A. R.
(2016). Potential drivers of virulence evolution in aquaculture. *Evolutionary Applications*, *9*(2),
344–354. https://doi.org/10.1111/eva.12342

Kubinak, J. L., & Potts, W. K. (2013). Host resistance influences patterns of experimental viral

adaptation and virulence evolution. *Virulence*, *4*(5), 410–418.

https://doi.org/10.4161/viru.24724

Kun, Á., Hubai, A. G., Král, A., Mokos, J., Mikulecz, B. Á., & Radványi, Á. (2023). Do pathogens

always evolve to be less virulent? The virulence–transmission trade-off in light of the COVID-

19 pandemic. *Biologia Futura*, 1–12. https://doi.org/10.1007/s42977-023-00159-2

Kuny, C. V., Bowen, C. D., Renner, D. W., Johnston, C. M., & Szpara, M. L. (2020). In vitro evolution

of herpes simplex virus 1 (HSV-1) reveals selection for syncytia and other minor variants in cell

culture. *Virus Evolution*, *6*(1), veaa013. https://doi.org/10.1093/ve/veaa013

Kutnjak, D., Elena, S. F., & Ravnikar, M. (2017). Time-Sampled Population Sequencing Reveals the

Interplay of Selection and Genetic Drift in Experimental Evolution of *Potato Virus Y*. *Journal*

of Virology, *91*(16), e00690-17. https://doi.org/10.1128/JVI.00690-17

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nature Methods*,

9(4), Article 4. https://doi.org/10.1038/nmeth.1923

Lauring, A. S. (2020). Within-host Viral Diversity, a Window into Viral Evolution. *Annual Review of*

Virology, *7*(1), 63–81. https://doi.org/10.1146/annurev-virology-010320-061642

Lenz, C., Haerty, W., & Golding, G. B. (2014). Increased Substitution Rates Surrounding Low-

Complexity Regions within Primate Proteins. *Genome Biology and Evolution*, *6*(3), 655–665.

https://doi.org/10.1093/gbe/evu042

Li, H., & Roossinck, M. J. (2004). Genetic Bottlenecks Reduce Population Variation in an Experimental

RNA Virus Population. *Journal of Virology*, *78*(19), 10582–10587.

https://doi.org/10.1128/JVI.78.19.10582-10587.2004

López-Muñoz, A. D., Rastrojo, A., Kropp, K. A., Viejo-Borbolla, A., & Alcamí, A. (2021). Combination

of long- and short-read sequencing fully resolves complex repeats of herpes simplex virus 2

strain MS complete genome. *Microbial Genomics*, *7*(6), 000586.

https://doi.org/10.1099/mgen.0.000586

Mikonranta, L., Friman, V.-P., & Laakso, J. (2012). Life History Trade-Offs and Relaxed Selection Can Decrease Bacterial Virulence in Environmental Reservoirs. *PLOS ONE*, *7*(8), e43801. https://doi.org/10.1371/journal.pone.0043801

Montville, R., Froissart, R., Remold, S. K., Tenaillon, O., & Turner, P. E. (2005). Evolution of Mutational Robustness in an RNA Virus. *PLOS Biology*, *3*(11), e381. https://doi.org/10.1371/journal.pbio.0030381

Morga, B., Jacquot, M., Pelletier, C., Chevignon, G., Dégremont, L., Biétry, A., Pepin, J.-F., Heurtebise, S., Escoubas, J.-M., Bean, T. P., Rosani, U., Bai, C.-M., Renault, T., & Lamy, J.-B. (2021). Genomic Diversity of the Ostreid Herpesvirus Type 1 Across Time and Location and Among Host Species. *Frontiers in Microbiology*, *12*, 711377. https://doi.org/10.3389/fmicb.2021.711377

Mourier, T., Sadykov, M., Carr, M. J., Gonzalez, G., Hall, W. W., & Pain, A. (2021). Host-directed editing of the SARS-CoV-2 genome. *Biochemical and Biophysical Research Communications*, *538*, 35–39. https://doi.org/10.1016/j.bbrc.2020.10.092

Nicolas, J.-L., Comps, M., & Cochennec, N. (1992). *Herpes-like virus infecting Pacific oyster larvae.* (No. 1). *12*(1), Article 1.

NIH. (2007). Understanding Emerging and Re-emerging Infectious Diseases. In *NIH Curriculum Supplement Series [Internet]*. National Institutes of Health (US). https://www.ncbi.nlm.nih.gov/books/NBK20370/

Patterson, N., Price, A. L., & Reich, D. (2006). Population Structure and Eigenanalysis. *PLOS Genetics*, *2*(12), e190. https://doi.org/10.1371/journal.pgen.0020190

Peck, K. M., & Lauring, A. S. (2018). Complexities of Viral Mutation Rates. *Journal of Virology*, *92*(14), 10.1128/jvi.01031-17. https://doi.org/10.1128/jvi.01031-17

Pelletier, C., Chevignon, G., Faury, N., Arzul, I., Garcia, C., Chollet, B., Renault, T., Morga, B., & Jacquot, M. (2025). Phylogenomic evidence for host specialization and genetic divergence in OsHV-1 infecting *Magallana gigas* and *Ostrea edulis*. *Infection, Genetics and Evolution*, 105803. https://doi.org/10.1016/j.meegid.2025.105803

879   Peñaloza, C., Gutierrez, A. P., Eöry, L., Wang, S., Guo, X., Archibald, A. L., Bean, T. P., & Houston,

880   R. D. (2021). A chromosome-level genome assembly for the Pacific oyster Crassostrea gigas.

881   *GigaScience*, *10*(3), giab020. https://doi.org/10.1093/gigascience/giab020

882   Pepin, J. F., Riou, A., & Renault, T. (2008). Rapid and sensitive detection of ostreid herpesvirus 1 in

883   oyster samples by real-time PCR. *Journal of Virological Methods*, *149*(2), Article 2.

884   https://doi.org/10.1016/j.jviromet.2008.01.022

885   Pernet, F., Fuhrmann, M., Petton, B., Mazurié, J., Bouget, J.-F., Fleury, E., Daigle, G., & Gernez, P.

886   (2018). Determination of risk factors for herpesvirus outbreak in oysters using a broad-scale

887   spatial epidemiology framework. *Scientific Reports*, *8*(1), Article 1.

888   https://doi.org/10.1038/s41598-018-29238-4

889   Petton, B., Boudry, P., Alunno-Bruscia, M., & Pernet, F. (2015). Factors influencing disease-induced

890   mortality of Pacific oysters Crassostrea gigas. *Aquaculture Environment Interactions*, *6*(3),

891   Article 3. https://doi.org/10.3354/aei00125

892   Phan, V., Gao, S., Tran, Q., & Vo, N. S. (2015). How genome complexity can explain the difficulty of

893   aligning reads to genomes. *BMC Bioinformatics*, *16*(Suppl 17), S3.

894   https://doi.org/10.1186/1471-2105-16-S17-S3

895   Poetsch, A. R. (2020). The genomics of oxidative DNA damage, repair, and resulting mutagenesis.

896   *Computational and Structural Biotechnology Journal*, *18*, 207–219.

897   https://doi.org/10.1016/j.csbj.2019.12.013

898   R Development Core Team. (2005). *R: A language and environment for statistical computing.*

899   [Computer software]. http://www.R-project.org

900   Renault, T., Cochennec, R., Le Deuff, R. M., & Chollet, B. (1994). *HERPES-LIKE VIRUS INFECTING*

901   *JAPANESE OYSTER (CRASSOSTREA GIGAS) SPAT*. 3.

902   Sanjuán, R. (2012). From Molecular Genetics to Phylodynamics: Evolutionary Relevance of Mutation

903   Rates Across Viruses. *PLOS Pathogens*, *8*(5), e1002685.

904   https://doi.org/10.1371/journal.ppat.1002685

905   Sanjuán, R., & Domingo-Calap, P. (2016). Mechanisms of viral mutation. *Cellular and Molecular Life*

906   *Sciences*, *73*(23), 4433–4448. https://doi.org/10.1007/s00018-016-2299-6

Sanjuán, R., Nebot, M. R., Chirico, N., Mansky, L. M., & Belshaw, R. (2010). Viral Mutation Rates.

Journal of Virology, *84*(19), 9733–9748. https://doi.org/10.1128/JVI.00694-10

Sanjuán, R., Pereira-Gómez, M., & Risso, J. (2016). Chapter 3—Genome Instability in DNA Viruses.

In I. Kovalchuk & O. Kovalchuk (Eds.), *Genome Stability* (pp. 37–47). Academic Press.

https://doi.org/10.1016/B978-0-12-803309-8.00003-3

Schikorski, D., Renault, T., Saulnier, D., Faury, N., Moreau, P., & Pépin, J.-F. (2011). Experimental

infection of Pacific oyster Crassostrea gigas spat by ostreid herpesvirus 1: Demonstration of

oyster spat susceptibility. *Veterinary Research*, *42*(1), Article 1. https://doi.org/10.1186/1297-

9716-42-27

Spitzer, K., Pelizzola, M., & Futschik, A. (2020). Modifying the Chi-square and the CMH test for

population genetic inference: Adapting to overdispersion. *The Annals of Applied Statistics*,

14(1). https://doi.org/10.1214/19-AOAS1301

Stoltzfus, A., & Norris, R. W. (2016). On the Causes of Evolutionary Transition:Transversion Bias.

Molecular Biology and Evolution, *33*(3), 595–602. https://doi.org/10.1093/molbev/msv274

Suquet, M., Kermoysan, G. de, Araya, R. G., Queau, I., Lebrun, L., Souchu, P. L., & Mingant, C. (2009).

Anesthesia in Pacific oyster, Crassostrea gigas. *Aquatic Living Resources*, *22*(1), Article 1.

https://doi.org/10.1051/alr/2009006

Tan, A., Abecasis, G. R., & Kang, H. M. (2015). Unified representation of genetic variants.

Bioinformatics, *31*(13), Article 13. https://doi.org/10.1093/bioinformatics/btv112

Trimpert, J., Groenke, N., Jenckel, M., He, S., Kunec, D., Szpara, M. L., Spatz, S. J., Osterrieder, N., &

McMahon, D. P. (2017). A phylogenomic analysis of Marek's disease virus reveals independent

paths to virulence in Eurasia and North America. *Evolutionary Applications*, *10*(10), 1091–

1101. https://doi.org/10.1111/eva.12515

Trimpert, J., Groenke, N., Kunec, D., Eschke, K., He, S., McMahon, D. P., & Osterrieder, N. (2019). A

proofreading-impaired herpesvirus generates populations with quasispecies-like structure.

Nature Microbiology, *4*(12), 2175–2183. https://doi.org/10.1038/s41564-019-0547-x

Valdivieso, A., Morga, B., Degremont, L., Mege, M., Courtay, G., Dorant, Y., Escoubas, J.-M., Gawra,

J., de Lorgeril, J., Mitta, G., Cosseau, C., & Vidal-Dupiol, J. (2025). DNA methylation

landscapes before and after Pacific Oyster Mortality Syndrome are different within and between resistant and susceptible *Magallana gigas*. *Science of The Total Environment*, *962*, 178385. https://doi.org/10.1016/j.scitotenv.2025.178385

Van Poelvoorde, L. A. E., Delcourt, T., Vuylsteke, M., De Keersmaecker, S. C. J., Thomas, I., Van Gucht, S., Saelens, X., Roosens, N., & Vanneste, K. (2022). A general approach to identify low-frequency variants within influenza samples collected during routine surveillance. *Microbial Genomics*, *8*(9), mgen000867. https://doi.org/10.1099/mgen.0.000867

Wang, J., Raskin, L., Samuels, D. C., Shyr, Y., & Guo, Y. (2015). Genome measures used for quality control are dependent on gene function and ancestry. *Bioinformatics*, *31*(3), 318–323. https://doi.org/10.1093/bioinformatics/btu668

White, D. W., Beard, R. S., & Barton, E. S. (2012). Immune Modulation During Latent Herpesvirus Infection. *Immunological Reviews*, *245*(1), 189–208. https://doi.org/10.1111/j.1600-065X.2011.01074.x

Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis*. Springer New York. https://doi.org/10.1007/978-0-387-98141-3

Xia, J., Bai, C., Wang, C., Song, X., & Huang, J. (2015). Complete genome sequence of Ostreid herpesvirus-1 associated with mortalities of Scapharca broughtonii broodstocks. *Virology Journal*, *12*(1), Article 1. https://doi.org/10.1186/s12985-015-0334-0

Zook, J. M., Chapman, B., Wang, J., Mittelman, D., Hofmann, O., Hide, W., & Salit, M. (2014). Integrating human sequence data sets provides a resource of benchmark SNP and indel genotype calls. *Nature Biotechnology*, *32*(3), 246–251. https://doi.org/10.1038/nbt.2835

Zwart, M. P., & Elena, S. F. (2015). Matters of Size: Genetic Bottlenecks in Virus Infection and Their Potential Impact on Evolution. *Annual Review of Virology*, *2*(Volume 2, 2015), 161–179. https://doi.org/10.1146/annurev-virology-100114-055135