Versatile Modular Antibodies for Sensitive and Specific Detection of Poly-ADP-Ribose

Helen Dauben[1], Ivan Matić[1,2]*

[1] Research Group of Proteomics and ADP-Ribosylation Signaling, Max Planck Institute for Biology of Ageing, 50931 Cologne, Germany
[2] Cologne Excellence Cluster for Stress Responses in Ageing-Associated Diseases (CECAD), University of Cologne, 50931 Cologne, Germany

* Corresponding author (IMatic@age.mpg.de)

Abstract

ADP-ribosylation is a chemically versatile modification of proteins, RNA and DNA that regulates various important signaling pathways, many of which are implicated in human diseases. Despite being discovered 60 years ago in the form of poly-ADP-ribosylation - the most studied form of this modification - investigating ADP-ribosylation at the molecular level has historically been challenging. By applying serine ADP-ribosylation-based antibody engineering technology, we have developed the first site-specific, as well as sensitive, mono-ADP-ribosylation modular antibodies. Here, we extend the scope of this technology to poly-ADP-ribosylation. By combining serine poly-ADP-ribosylated peptides as antigens, PARP1 serine mono- and poly-ADP-ribosylation for validation, phage display and the SpyTag protein ligation systems, we developed modular antibodies that are highly specific for poly-ADP-ribosylation. SpyTag-based coupling of horseradish peroxidase at specific positions, distant from the antigen-binding region of the antibody, yields a format that simplifies immunoblotting while dramatically enhancing poly- ADP-ribosylation detection sensitivity. Additionally, the creation of synthetic immunoglobulin formats – mouse, rabbit and human – enables straightforward co-detection of mono- and poly-ADP-ribosylation in cells by immunofluorescence. Our new tools improve and simplify the detection of poly-ADP-ribosylation, particularly in immunoblotting, enabling specific investigations of this key cellular signal in the context of other distinct forms of ADP-ribosylation.

Introduction

ADP-ribosylation is a versatile post-translational modification (PTM) that regulates numerous cellular processes, including DNA repair, transcription, signal transduction, and cell death[1-4]. It involves the enzymatic transfer of ADP-ribose units from nicotinamide adenine dinucleotide (NAD+) to target molecules, mediated primarily by members of the ADP-ribosyltransferase (ART) enzyme family[5]. This dynamic modification can occur as mono-ADP-ribosylation (mono-ADPr), where a single ADP-ribose unit is present, or poly-ADP-ribosylation (poly-ADPr), involving the formation of linear or branched chains of ADP-ribose. The reversibility of this process, mediated by ADP-ribosyl hydrolases, underscores its regulatory significance in maintaining cellular homeostasis[6,7]. First identified in the 1960s, ADP-ribosylation has since emerged as a critical player in cellular stress responses, particularly in the context of DNA damage repair[8,9]. Poly(ADP-ribose) polymerase 1 (PARP1), the most studied member of the PARP family, is activated in response to DNA strand breaks, orchestrating the recruitment of DNA repair machinery and chromatin relaxation. Beyond DNA repair, ADP-ribosylation influences chromatin remodeling, RNA processing, and immune signaling. Notably, dysregulation of ADP-ribosylation is implicated in various diseases, including cancer, neurodegenerative disorders, and inflammatory conditions, making it an attractive target for therapeutic intervention[10-14].

The biological complexity of ADP-ribosylation stems from the diversity of its enzymes, substrates, and cellular contexts. The PARP family alone consists of 17 members in humans, of which four are known to synthesize poly-ADPr (PARP1, PARP2, PARP5a/TNKS1, and PARP5b/TNKS2), each exhibiting distinct substrate specificities and cellular functions[15]. Moreover, emerging evidence highlights non-canonical roles of ADP-ribosylation, such as its involvement in liquid-liquid phase separation and the regulation of stress granules[16-18].

ADPr is one of the most chemically complex PTMs, with conjugation chemistries ranging from O-glycosidic (serine and tyrosine), S-glycosidic (cysteine), N-glycosidic (arginine) to ester (aspartate and glutamate) linkages. Strikingly, some of the main conjugation chemistries have remained elusive for decades. A prime example of this is serine ADP-ribosylation, which, despite being the most abundant - though not the exclusive target residue in PARP1/2 signaling - was only discovered less than ten years ago[19-22].

The structural complexity of ADP-ribosylation arises from the unique structure of ADP-ribose and its ability to form both mono-ADPr and intricate polymer chains. The polymeric forms, particularly poly-ADPr, can adopt linear or branched conformations, creating a diverse array of structural motifs[23,24]. These variations influence the binding affinity and functionality of

effector proteins, known as "readers," which recognize specific poly-ADPr structures through
macrodomains, PAR-binding zinc fingers, and other motifs[25-28]. Additionally, the rapid
synthesis and degradation of these chains add a temporal layer of complexity, making ADP-
ribosylation a highly dynamic and versatile regulatory mechanism.

The inherent biological and chemical/structural complexity of ADPr has historically hindered
its study and the development of appropriate analytical tools. This lack of tools has historically
remained a significant limitation in many studies. Nevertheless, the critical role of poly-ADPr
in modulating essential biological processes, particularly PARP1 signaling in the DNA damage
response, has driven the development of various approaches and tools to study this
modification since its discovery in the 1960s[29].

ADPr research methods and tools can be broadly categorized into indirect and direct
approaches. Indirect methods include the use of small molecule inhibitors to block poly-ADPr
writers[30], or genetic and RNA interference-based techniques to modulate the occurrence and
persistence of this modification by targeting readers or erasers. Direct approaches aim to
study poly-ADPr either *in-vitro* or *in-vivo*, including artificial construction or modification of poly-
ADPr[31,32] and its targets to unravel its structural complexity and biological functions. Early
techniques utilized NAD^+ analog probes[33], which remain in use today with further
optimizations, such as Et-DTB-NAD^{+}[34-36]. In addition to chemical probes, enrichment
strategies originally designed for other post-translational modifications (e.g.,
phosphoenrichment) have been adapted for poly-ADPr[37]. Elegant methods such as Enzymatic
Labeling of Terminal ADP-Ribose (ELTA) offer another avenue for studying this modification
by leveraging enzymes like OAS1 and modified dATP analogs to chemically tag and track
poly-ADPr chains[38].

Among direct methods to study endogenous poly-ADPr levels, antibodies and reagents
specific to poly-ADPr have emerged as promising tools to advance the field. Notably, many of
the most frequently cited poly-ADPr antibodies have been polyclonal, which are challenging
to produce consistently, leading to their discontinuation. This left the field reliant on monoclonal
antibodies, such as 10H and the much less commonly used 16B. The 10H antibody, which
was the most widely used tool in the early decades of ADPr research, primarily recognizes
long poly-ADPr chains starting at 10–20 units and has low sensitivity compared to reagents
developed in the last decade.

Recently, the Kraus laboratory pioneered the conversion of established ADP-ribose reader
domains into Fc fusion, antibody-like recombinant reagents. One widely used tool is the
AF1521 domain, which has been applied in various studies. Its mutated from, with increased

affinity is used particularly for enrichment followed by mass spectrometry analysis[39,40]. While

this approach has identified numerous ADPr sites, it cannot distinguish between mono- and

poly-ADPr because the domain binds to the terminal unit, enriching both forms

indiscriminately. Under certain conditions, both the original and affinity-matured AF1521

domain exhibit hydrolytic activity towards aspartate and glutamate ADPr[39]. Another approach

employs the WWE domain of RNF146, which specifically binds poly-ADPr. More precisely, it

targets iso-ADPr, a structure unique to poly-ADPr chains, thereby avoiding misinterpretation

by not interacting with mono-ADPr[41]. This innovation added a more selective tool for studying

poly-ADPr[42,43]. However, the diversity of poly-ADPr-specific domains, each potentially evolved

to recognize distinct poly-ADPr structures, suggests that individual domains might exhibit

specificity for certain chain types or protein targets. Even among WWE domains, differences

in binding specificity and affinity have been observed, as exemplified by the two WWE

domains in PARP13[44]. As a result, the application of these protein domain-based tools may

be beneficial to study specific research questions rather than offering the broad applicability

of antibodies. In fact, a so-called 'Poly/Mono-ADP Ribose' rabbit monoclonal antibody is

frequently used to detect poly-ADPr. While it is sensitive and specific for ADP-ribose, this

antibody is not specific for poly-ADP-ribose, as it also recognizes mono-ADP-ribose[45].

By converting our biological discovery of serine ADPr by HPF1/PARP1[19,20] into a technology

for generating ADPr peptides we have developed several anti-ADPr modular antibodies[21,46-

49]. This includes the first - and so far only - site-specific ADPr antibodies[46], as well as a

sensitive and specific mono-ADPr antibody that has proven effective in detecting not only

serine mono-ADPr[47] but also beyond PARP1 signaling, including interferon signaling[50-52].

Importantly, serine ADPr reactions not only yield antigens for the initial selection of antibody

candidates via phage display but also play a crucial role in subsequent steps, enabling in-

depth validation and characterization of the best antibody candidates. While phage display

used in our approach offers several advantages compared to traditional animal-based

approaches for generating antibodies, including the rapid initial identification of multiple

binders to the antigen of choice, it has the drawback of yielding recombinant antibodies in non-

standard formats. While conversion to standard immunoglobulins and other formats is

feasible, cloning-based conversion is impractical and time-consuming, counterbalancing some

of the advantages of recombinant antibodies. This has changed dramatically with the

application of SpyTag/SpyCatcher protein ligation technology to recombinant antibodies. This

biochemical format conversion enables the almost instantaneous generation of a versatile

family of formats, including rabbit, mouse and human synthetic immunoglobulins, as well as

site-specific labeling with horseradish peroxidase (HRP), biotin, and other tags[53]. In particular,

a bivalent modular antibody carrying three HRP molecules has proven to be a much more

sensitive, simpler and faster immunoblotting approach compared to standard protocols based

on secondary antibodies[53], significantly improving detection of ADPr[21,47].

Here, we extend the scope of our serine ADPr technology to poly-ADPr, combining it with a

proven phage display recombinant antibody platform and the SpyTag protein ligation system

to develop sensitive and specific poly-ADPr antibodies. This advance enables versatile,

precise and consistent detection across a variety of experimental applications, such as

immunofluorescence, immunoblotting and immunoprecipitation. In this study, we present the

characterization and validation of new poly-ADPr recombinant antibodies, reasoning that the

modular nature of this antibody, as well as our previously introduced complementary

antibodies, will enable more versatile and sensitive ADPr detection. Despite recent advances,

many aspects of ADP-ribosylation remain poorly understood, with outstanding questions

regarding its substrate specificity, spatial and temporal regulation. Furthermore, recent

discoveries of novel ADP-ribosylation readers, writers, and erasers promise to reshape our

understanding of this modification and its broader impact on cellular physiology.

Material and Methods

<u>Resource availability</u>

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Dr. Ivan Matić (imatic@age.mpg.de)

Material availability

Antibodies, Reagents and BiSpyCatcher

The α-Poly-ADPr antibodies generated in this study (AbD64138 and AbD64235) are not yet commercially available but requests can be forwarded to the lead contact. Previously published antibodies converted to the SpyTag formats (AbD43647) and SpyCatchers are available through BioRad Laboratories (unconjugated, requires conjugation to a SpyCatcher: TZA020; HRP-conjugated, ready-to-use: TZA020P)

Uncoupled SpyTagged antibodies were conjugated as described below to following SpyCatchers: bivalent-HRP Format (BiSpyCatcher2 #TZC002P, Bio-Rad); bivalent synthetic IgG Format mouse (Mouse IgG2a-FcSpyCatcher3 # TZC012, Bio-Rad); bivalent synthetic IgG Format rabbit (Rabbit IgG-FcSpyCatcher3 # TZC013, Bio-Rad)

Commercial antibodies and reagents used in this study: α-Poly/Mono-ADP Ribose (D9P7Z) (Cell Signaling Technology, #89190), α-Poly (ADPr-ribose) monoclonal antibody (10H) (Enzo Life Sciences, ALX-804-220-R100), α-Poly-ADP-ribose binding reagent (Merck, MABE1031)

<u>Experimental Model</u>

Cell Culture and drug treatments

U2OS cell lines were obtained, authenticated by STR profiling and confirmed mycoplasma free by ATCC cell line authentication service. Cells were cultured in Glutamax-DMEM supplemented with 10% bovine serum and 100U/ml penicillin/Streptomycin at 37°C and 5% CO_2.

To induce DNA damage, the cell medium was aspirated and replaced with 37°C complete DMEM containing 2mM H_2O_2 for 10 min. For PARP inhibition (Olaparib, Cayman Chemical) cells were treated with 1 µM Olaparib in complete DMEM for 30 min prior to DNA damage treatment, followed by 10 min 2mM H_2O_2 & 1µM Olaparib treatment.

<u>Method details</u>

Antibody SpyCatcher Coupling

All Bio-Rad antibodies have been obtained unconjugated in a monovalent format and were conjugated to BiSpyCatchers inhouse leading to bivalent HRP conjugated or bivalent

synthetic IgG antibody formats. In brief antibody stocks were diluted to 1 mg/ml in PBS and

incubated with 1/10th the volume of unconjugated antibody with BiSpyCatcher for 1 hr at RT,

i.e. 10 µl of BiSpyCatcher was added to 100 µl of 1mg/ml unconjugated antibody. Afterwards

they were aliquoted and stored at -20°C.

Expression and purification of recombinant proteins

Proteins were expressed in *Escherichia coli* (*E.coli*) and purified essentially as previously

described[54], but with addition of a size-exclusion chromatography purification using HiLoad

16/60 Superdex 75 column. PARP1 wild-type was expressed and purified as previously

reported[54,55]. PARG was purified as described[56]. HPF1 was expressed and purified as

reported[57].

In vitro Modification of Proteins

Recombinant PARP1 protein (0.5 µM) was incubated with 2 mM NAD^+ in the presence of

HPF1 (5 µM). The reaction buffer contained 50 mM Tris-HCl pH 8.0, 100 mM NaCl, 2 mM

$MgCl_2$ and 1 ng/µL sonicated DNA. Reactions were allowed to proceed for at 30 min at RT,

before being stopped by the addition of 1 µM Olaparib. To create mono- and poly-ADP-

ribosylated protein samples were split and one half was further incubated with 3 µM PARG for

1 hr at RT and stopped by the addition of Laemmli buffer. Samples were boiled and run on an

8% Bis-Tris gel (Invitrogen).

For Dot-blotting the same reaction as above was performed without splitting the reaction after

stopping with Olaparib. No loading buffer was added. A dilution series of the sample was

performed using reaction buffer and the same volumes (1 µl) of the probe was pipetted onto

a dry nitrocellulose membrane. The membrane was left to dry and was proceeded with 5%

non-fat dried milk blocking and regular immunoblotting steps as described below.

SDS Sample Preparation and Immunoblotting

U2OS cells were treated as indicated, lysed in SDS buffer (4% SDS; 50 mM HEPES pH 7.2,

150 mM NaCl, 5 mM $MgCl_2$) and incubated for 5 min at RT with recombinant benzonase

(smDNase, 750U per sample). Samples were spun at max. speed for 10 min and supernatant

was transferred into fresh tube. Sample concentration was measured using a Pierce Dilution-

Free BCA Protein Assay Kit (#A55864, ThermoScientific). Samples were prepared with

NuPAGE LDS sample buffer with final concentration of 5 mM DTT (Sigma) and loaded on 4-

12% Bis-Tris gel (Invitrogen). After SDS-Page the gels are transferred onto PVDF membranes

(Amersham) using wet transfers at 90 mA overnight on ice. The precooled transfer buffer is 1x

NuPAGE transfer buffer (Invitrogen), 20% ethanol in water.

For immunoblotting membranes were blocked in 5% non-fat dried milk before primary

antibodies were added. Primary Antibodies (commercial antibodies and HRP-formats of Bio-

Rad conjugated antibodies) were prepared in milk using concentrations found in figures. The
membranes were incubated overnight at 4°C before being washed in TBS-T (25 mM Tris-HCl
pH 7.5, 150 mM NaCl, 0.05% Tween-20) 3x10 min. Commercial antibodies (Enzo, Merck, Cell
Signaling Technologies) were incubated with a secondary antibody (Anti-mouse IgG HRP-
conjugated secondary Amersham Cat# NA931V, Anti-rabbit IgG HRP-conjugated secondary
Merck Cat# GENA934-1ML) for 1 hr at RT, followed by 3x10 min TBS-T wash. All membranes
were developed using ECL Select Western Blotting Detection Reagent (Amersham).

Immunofluorescence
U2OS cells were grown on sterile glass coverslips in 24-well plates. treated as indicated and
fixed for 20 min with Methanol at -20°C. The fixed cells were washed twice with PBS, and
blocked for 1hr, RT with 5% Normal-Goat-Serum in PBS supplemented with 0.3% Triton.
Primary antibodies (commercial antibodies and synthetic IgG formats of Bio-Rad conjugated
antibodies) were diluted as indicated and incubated overnight at 4°C in 1x PBS, 1% BSA and
0.3% Triton. Coverslips were washed in PBS 3x5 min before adding the secondary antibody
and DAPI, again in 1x PBS, 1% BSA and 0.3% Triton for 1 hr, RT protected from light.
Coverslide were washed again in PBS 3x5 min before mounted on microscopy slides using
Prolong   Diamond   Antifade   Mountant   (ThermoFisher).   Cells   were   scanned   for
immunofluorescence using a Leica SP8-X inverted laser-scanning confocal microscope.

Results

Generation of a serine poly-ADPr peptide

Previously, we established the PARP1/HPF1 complex[20,57,58] as an efficient and specific biocatalytic tool for the rapid and scalable generation of mono-ADPr peptides, enabling the development, validation and characterization of site-specific, mono-ADPr and pan-ADPr antibodies[46,47]. Here, we aimed to adapt our strategy to generate modular antibodies specific for poly-ADPr. Inspired by the possibility of extending the initial ADP-ribose attached to serine of the peptide[40], we employed a simple protocol that does not require separation of different ADP-ribosylated peptide species, reasoning that this approach would be sufficient for the development of poly-ADPr antibodies. Specifically, we generated and purified the mono-ADPr peptide from proteins using C18 StageTips[46] and subsequently added PARP1 without HPF1 to extend the mono-ADPr modification to di- and tri-ADPr. This process proved to be as scalable as the previously described serine mono-ADPr peptide generation method[46]. The resulting peptides were then employed in a phage display screen, leading to the identification of several positive clones. From these, we selected the two most promising candidates for further validation and comparison, as detailed in the following sections (AbD64138, AbD64235).

Modular Antibodies for versatile, specific and sensitive detection of Poly-ADPr

Given the demonstrated value of the SpyTag system in the generation of modular antibodies[46,47], we also chose to implement it for our new set of poly-ADPr antibodies. As illustrated in the scheme (Fig. 1), the monovalent poly-ADPr antibody can be easily converted into a bivalent format in a fast highly reliable reaction. This conversion adds an avidity effect in addition to the high affinity of the antibody, further enhancing signal intensity and specificity. Depending on the application, a variety of catchers can be chosen for coupling. For example, an HRP-catcher can be used, resulting in an HRP-coupled bivalent poly-ADPr antibody for immunoblotting. This eliminates the need for a secondary antibody, reducing costs, antibody incubation time and increasing signal intensity, as demonstrated previously[47,53]. Alternatively, for applications not reliant on HRP, such as immunofluorescence, a catcher fused with the mouse or rabbit Fc region can be selected, creating a synthetic IgG antibody that, like a regular IgG, is detectable by fluorophore-coupled secondary antibodies.

After coupling, we first assessed the specificity of the poly-ADPr antibodies (AbD64138, AbD64235) to ensure there was no off-target binding to other forms of ADPr. To test this, we took advantage of a key feature of serine ADPr: the inability of PARG to cleave the bond between serine and the initial ADP-ribose, despite its ability to hydrolyze tyrosine and

aspartate/glutamate ADPr[21,59]. Specifically, we automodified PARP1 in the presence of HPF1,
resulting in nearly complete serine Poly-ADPr of PARP1. Addition of PARG to a portion of the
reaction completely reduced Poly-ADPr-PARP1 to mono-ADPr-PARP1[46], resulting in pure
mono- and poly-ADPr standards for antibody validation. As expected, our mono-ADPr
antibody[47] recognized exclusively mono-ADPr (Fig. 1A). Our newly developed poly-ADPr
antibodies did not detect any mono-ADPr, but produced an intense signal for poly-ADPr
PARP1. In contrast, pan-ADPr antibodies, such as D9P7Z from Cell Signaling Technology
(CST), detected both mono- and poly-ADPr (Fig. 1,A). We also compared our antibodies with
other commercially available poly-ADPr reagents, including 10H (Enzo) antibody and the
WWE-based poly-ADPr reagent from Merck (MABE1031)[42]. All poly-ADPr probes were shown
to be specific for poly-ADPr, although 10H only detected long ADPr polymers as previously
reported[60] (Fig. 1, A). Like MABE1031, our new antibodies were able to detect the full range
of poly-ADPr chains in immunoblotting. To ensure a fair comparison of signal intensities, we
tested the reagents at two different dilutions, starting with those recommended by the
manufacturers/publications for optimal performance. However, we found that using our
antibodies at the same concentration (1 µg/ml) led to signal bleaching before detection.
Consequently, we lowered the concentration by a factor of ten, which allowed for signal
detection without compromising the poly-ADPr detection by commercially available reagents.
Despite this adjustment, the sensitivity of our antibodies is so high that oversaturation occurred
after just one second of exposure.

To allow for a more accurate comparison without signal saturation, we performed a dot blot
using a 1:2 dilution series of poly-ADPr PARP1 (Fig. 1,B). Again, using antibody dilutions as
recommended by the manufacturer and a common dilution of 0.1 µg/ml, we compared the
signal intensities of our new tools to those of other reagents. The results showed that our new
antibodies provided significantly higher signal intensity compared to the MABE1031 reagent
and D9P7Z antibody. The 10H antibody did not produce any detectable signal at this dilution.

These results demonstrate a high specificity for poly-ADPr of the new antibodies, with no
detection of mono-ADPr, confirming that our antibodies selectively recognize poly-ADPr
chains.

<u>Sensitive detection of Poly-ADPr targets in cells</u>

After establishing their poly-ADPr specificity and sensitivity in *in vitro* biochemical reactions,
we applied the new antibodies to cell lysates to evaluate their performance on cellular poly-
ADPr targets and assess potential off-target binding. To induce high levels of poly-ADPr, cells
were treated H_2O_2 for ten minutes and additionally treated with the PARP1 inhibitor Olaparib.

We used optimal antibody dilutions for all probes to ensure peak performance. First, we

demonstrated that there was no nonspecific binding to proteins, as shown in lanes 2 and 4 of

each blot, where no signal was detected in olaparib-treated samples (Fig. 2,A). Second, even

when using ten times higher amounts of commercially available reagents, our new poly-ADPr

antibodies still exhibited superior signal intensity. Third, our antibodies also improved the

detection of poly-ADPr in unperturbed cells, where its generation has been shown to be

induced by unligated Okazaki fragments during DNA replication[61], even with low exposures,

as seen in lane 1.

We then tested the specificity of our antibodies in cells using immunofluorescence and the

synthetic IgG format. Again, we applied optimal antibody dilutions and tailored microscope

settings for each reagent to achieve high performance. While intensity levels are not directly

comparable, the focus was on the specificity of the signal and the signal-to-noise ratios. Our

poly-ADPr antibodies AbD64138 and AbD64235 provided a superior signal-to-noise ratio in

untreated versus H_2O_2-treated cells (Fig. 2,B), outperforming other poly-ADPr reagents. The

signal specificity was comparable to the poly-/mono-ADPr antibody (CST) and our mono-ADPr

reagent (AbD43647), with antibody AbD64138 performing slightly better over AbD64235.

In conclusion, these experiments validate the high signal-to-noise ratio of our new tools for

both immunoblotting and immunofluorescence, demonstrating their versatility and potential for

multiple applications.

<u>SpyTag System Enables Flexible Co-Detection of Mono- and Poly-ADPr In Cells</u>

Finally, to assess the potential for simultaneous use of our poly-ADPr antibodies alongside

our previously published mono-ADPr antibodies, we performed co-staining of both poly-ADPr

(AbD64138) and mono-ADPr (AbD43647) in cells (Fig. 3,B). For this, we coupled the poly-

ADPr antibody with a mouse Fc catcher and the mono-ADPr antibody with a rabbit Fc catcher.

To prevent channel bleed-through, we used two fluorophores with distinct emission peaks and

employed line sequencing during imaging. The results demonstrate that, following a ten-

minute H_2O_2 treatment, both mono- and poly-ADPr can be detected simultaneously in the

same cells. Notably, the staining patterns revealed distinct distributions: the mono-ADPr signal

appeared more homogenously distributed throughout the nucleus, while the poly-ADPr signal

exhibited a more punctate pattern, forming small foci.

The ability to co-stain both forms of ADPr enhances the depth of analysis in

immunofluorescence, allowing for the use of smaller sample sizes and reduced experimental

times. Additionally, the flexible coupling of different IgG species to either poly- or mono-ADPr

antibodies provides (Fig.3, B) versatility for co-staining with other targets, such as full-length
PARP1, facilitating broader applications in cellular analysis.

To summarize the results of our antibody validation, we demonstrate superior performance in
both signal intensity and specificity for immunoblotting and immunofluorescence applications.
While both of our antibodies are highly specific for poly-ADPr and exceptionally sensitive,
especially in immunoblotting, AbD64138 shows slightly better performance compared to
AbD64235.

Discussion

Despite decades of research on ADPr and recent technological advances[29], the availability of specific and sensitive tools for studying all forms of this challenging PTM remains limited. This study aimed to extend the reach of our serine ADPr technology, which enabled the development of the first site-specific ADPr antibodies as well as specific and sensitive mono-ADPr antibodies[21,46-49], to poly-ADPr, resulting in a highly sensitive, specific and versatile modular antibody. As a technology, Serine ADPr offers several advantages inherent to the nature of the physiological reaction. First, PARP1 is the most catalytically active PARP enzyme, strongly activated by DNA breaks both in cells and in biochemical reactions. Therefore, the modification is highly scalable, allowing reactions to be upscaled to milligram quantities suitable for antibody panning. Second, HPF1 enables not only precise targeting of PARP1 activity toward serine residues, but also modification of peptides rather than just PARP1 automodification[20]. Third, the biological interplay between serine ADPr and canonical histone marks[62] can be harnessed to program the HPF1/PARP1 complex, directing it to target a specific serine using phosphorylation as an unconventional protecting group[46]. Fourth, a key advantage of serine-ADPr is its chemical stability: the O-glycosidic linkage remains intact across a broad range of pH conditions and withstands heat exposure up to 95°C. This stability facilitates efficient handling and minimizes sample loss during experimental procedures, a problem observed with other forms of ADPr, particularly aspartate/glutamate ADPr[21,63]. Fifth, while the poly-ADPr eraser PARG can cleave the bonds between ADP-ribose and tyrosine, aspartate and glutamate[21,59], it does not remove serine ADPr. Here, beyond peptide generation for phage display and antibody screening, we harness the resistance of serine ADPr to PARG to achieve nearly complete modification of PARP1 at the protein level, either by poly-ADPr or, upon PARG treatment, of mono-ADPr. This enables comprehensive antibody specificity testing, ensuring the selection of clones with high specificity for poly-ADPr and minimal off-target binding.

The newly developed poly-ADPr antibodies presented in this study offer several key advantages. They enable the detection of a broad range of poly-ADPr, from short to long polymeric chains, as demonstrated by full signal coverage in immunoblotting analysis. The modular nature of these antibodies, derived from the SpyTag technology results in strong signal intensity, especially in immunoblotting, and offers versatility across multiple applications, including immunofluorescence and immunoprecipitation, allowing for the detection of poly-ADPr even under untreated conditions. The modular SpyTag system, as illustrated previously for our mono-ADPr and site-specific antibodies[47], provides significant benefits. The reaction used to couple various catchers to the monovalent antibodies is fast,

efficient, and highly reliable. For immunoblotting, the HRP Catcher eliminates the need for a secondary antibody, significantly increasing signal intensity, reducing costs by lowering the required antibody amount per experiment and shortening the overall immunoblotting procedure. Additionally, the SpyTag system enables the creation of our poly-ADPr antibodies as synthetic immunoglobulins, which can be used in immunofluorescence assays. This flexibility allows for the selection of species-specific antibodies (e.g., mouse, rabbit, or human) tailored to experimental needs, allowing co-staining with other ADPr- or protein-specific antibodies.

Beyond their high affinity and modularity, the antibodies described in this study also offer high reliability. As monoclonal recombinant antibodies, they ensure consistent quality and reproducibility, unlike previously available polyclonal poly-ADPr antibodies, which were subject to batch variability and eventual discontinuation. This homogeneity and consistency further reinforce their utility as a long-term tool for ADPr research.

The most critical factor when selecting a tool to study ADPr is that its effectiveness is inherently dependent on the research question being asked. Choosing the appropriate tool is essential to obtaining meaningful and accurate insights. Given the high variability of ADPr arising from modifications at different sites, in diverse forms, by various proteins across multiple cellular signaling pathways, multiple complementary tools are required to study this PTM in an in-depth manner. Broad yet ADPr-specific tools, such as the highly sensitive poly-/mono-ADPr antibody from Cell Signaling Technology, allow for the general detection of ADPr. However, their inability to distinguish between different ADPr forms can complicate result interpretation, especially as it is emerging that mono- and poly-ADPr act as distinct signals, even within pathways where they co-occur[21,47]. In this study, we aimed to develop antibodies that provide both broad detection and high specificity, fully complementing our previously developed mono-ADPr antibodies[46,47] to enable a clear differentiation between mono- and poly-ADPr. The tools presented here contribute to expanding the available toolkit for ADPr research at this level. To further investigate mono- and poly-ADPr in greater depth, additional specialized tools may be necessary. For instance, site-specific mono-ADPr and poly-ADPr antibodies may enable a more detailed examination of ADPr modifications, while engineered protein domains such as WWE fused to an Fc chain[39,43,64], offer an alternative approach to studying certain forms of poly-ADPr.

In conclusion, we present the first application of our technology for generating poly-ADPr-specific antibodies. As with the development of site-specific and mono-ADPr antibodies, we incorporated serine ADPr at nearly every phase of the antibody generation process, from

precisely modified peptide antigens to multiple validation steps and in-depth characterization.
Obtaining purely mono- and poly-ADP-ribosylated PARP1 was particularly useful for
unambiguously establishing the specificity of the new antibodies. In this process, phage
display and the SpyTag protein ligation system were essential for generating numerous initial
candidate clones and for conferring modularity to the final, validated antibodies, increasing
their versatility and sensitivity. We anticipate that these new tools, along with previously
developed reagents, will play key roles in deciphering the distinct roles of the two primary
forms of ADPr - its monomeric and polymeric signals - in various signaling pathways.

Acknowledgements

This work was funded by the Max Planck Society, the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany´s Excellence Strategy (CECAD, EXC 2030-390661388) and by the European Research Council (ERC-CoG-864117) to I.M.

Author Contributions

I.M. conceived the project. H.D. and I.M. designed research and wrote the manuscript. H.D. performed all the experiments.

Declaration of Interest

The authors I.M. and H.D. declare the following competing financial interest: Max-Planck-Innovation, which is responsible for technology transfer from Max Planck Institutes, has licensed the antibody AbD43647 to Bio-Rad Laboratories, which markets it for research purposes. Additionally, a similar agreement is in preparation for the commercialization of AbD64138.

References

1   Messner, S. *et al.* PARP1 ADP-ribosylates lysine residues of the core histone tails.
Nucleic Acids Research **38**, 6350-6362 (2010). https://doi.org/10.1093/nar/gkq463
2   Chou, H. Y., Chou, H. T. & Lee, S. C. CDK-dependent activation of poly(ADP-ribose)
polymerase member 10 (PARP10). *J Biol Chem* **281**, 15201-15207 (2006).
https://doi.org/10.1074/jbc.M506745200
3   Gupte, R., Liu, Z. & Kraus, W. L. PARPs and ADP-ribosylation: recent advances
linking molecular functions to biological outcomes. *Genes & development* **31**, 101-
126 (2017). https://doi.org/10.1101/gad.291518.116
4   Lüscher, B. *et al.* ADP-Ribosylation, a Multifaceted Posttranslational Modification
Involved in the Control of Cell Physiology in Health and Disease. *Chem Rev* **118**,
1092-1136 (2018). https://doi.org/10.1021/acs.chemrev.7b00122
5   Lüscher, B. *et al.* ADP-ribosyltransferases, an update on function and nomenclature.
The FEBS Journal **289**, 7399-7410 (2022).
https://doi.org/https://doi.org/10.1111/febs.16142
6   O'Sullivan, J. *et al.* Emerging roles of eraser enzymes in the dynamic control of
protein ADP-ribosylation. *Nat Commun* **10**, 1182 (2019).
https://doi.org/10.1038/s41467-019-08859-x
7   Prokhorova, E. *et al.* Unrestrained poly-ADP-ribosylation provides insights into
chromatin regulation and human disease. *Mol Cell* **81**, 2640-2655.e2648 (2021).
https://doi.org/10.1016/j.molcel.2021.04.028
8   Chambon, P., Weill, J. D. & Mandel, P. Nicotinamide mononucleotide activation of
new DNA-dependent polyadenylic acid synthesizing nuclear enzyme. *Biochem
Biophys Res Commun* **11**, 39-43 (1963).
9   Collier, R. J. & Pappenheimer, A. M., Jr. STUDIES ON THE MODE OF ACTION OF
DIPHTHERIA TOXIN. II. EFFECT OF TOXIN ON AMINO ACID INCORPORATION IN
CELL-FREE SYSTEMS. *J Exp Med* **120**, 1019-1039 (1964).
https://doi.org/10.1084/jem.120.6.1019
10   Challa, S. *et al.* Ribosome ADP-ribosylation inhibits translation and maintains
proteostasis in cancers. *Cell* **184**, 4531-4546.e4526 (2021).
https://doi.org/10.1016/j.cell.2021.07.005
11   Vermehren-Schmaedick, A. *et al.* Characterization of PARP6 Function in Knockout
Mice and Patients with Developmental Delay. *Cells* **10** (2021).
https://doi.org/10.3390/cells10061289
12   Bütepage, M., Eckei, L., Verheugd, P. & Lüscher, B. Intracellular Mono-ADP-
Ribosylation in Signaling and Disease. *Cells* **4**, 569-595 (2015).
https://doi.org/10.3390/cells4040569
13   Palazzo, L., Mikolčević, P., Mikoč, A. & Ahel, I. ADP-ribosylation signalling and
human disease. *Open Biology* **9**, 190041 (2019). https://doi.org/10.1098/rsob.190041
14   Rosado, M. M. & Pioli, C. ADP-ribosylation in evasion, promotion and exacerbation of
immune responses. *Immunology* **164**, 15-30 (2021).
https://doi.org/10.1111/imm.13332
15   Vyas, S. *et al.* Family-wide analysis of poly(ADP-ribose) polymerase activity. *Nature
communications* **5**, 4426-4426 (2014). https://doi.org/10.1038/ncomms5426
16   Alemasova, E. E. & Lavrik, O. I. Poly(ADP-ribose) in Condensates: The PARtnership
of Phase Separation and Site-Specific Interactions. *International Journal of Molecular
Sciences* **23** (2022).
17   Catara, G. *et al.* PARP1-produced poly-ADP-ribose causes the PARP12 translocation
to stress granules and impairment of Golgi complex functions. *Sci Rep* **7**, 14035
(2017). https://doi.org/10.1038/s41598-017-14156-8
18   Leung, Anthony K. L. *et al.* Poly(ADP-Ribose) Regulates Stress Responses and
MicroRNA Activity in the Cytoplasm. *Molecular Cell* **42**, 489-499 (2011).
https://doi.org/https://doi.org/10.1016/j.molcel.2011.04.015

19 Leidecker, O. *et al.* Serine is a new target residue for endogenous ADP-ribosylation
on histones. *Nature Chemical Biology* **12**, 998 (2016).
https://doi.org/10.1038/nchembio.2180
20 Bonfiglio, J. J. *et al.* Serine ADP-Ribosylation Depends on HPF1. *Molecular Cell* **65**,
932-940.e936 (2017). https://doi.org/https://doi.org/10.1016/j.molcel.2017.01.003
21 Longarini, E. J. & Matić, I. Preserving ester-linked modifications reveals glutamate
and aspartate mono-ADP-ribosylation by PARP1 and its reversal by PARG. *Nature*
Communications **15**, 4239 (2024). https://doi.org/10.1038/s41467-024-48314-0
22 Longarini, E. J. & Matic, I. The fast-growing business of Serine ADP-ribosylation.
DNA Repair (Amst) **118**, 103382 (2022).
https://doi.org/10.1016/j.dnarep.2022.103382
23 Kiehlbauch, C. C., Aboul-Ela, N., Jacobson, E. L., Ringer, D. P. & Jacobson, M. K.
High resolution fractionation and characterization of ADP-ribose polymers. *Anal*
Biochem **208**, 26-34 (1993). https://doi.org/10.1006/abio.1993.1004
24 Krietsch, J. *et al.* Reprogramming cellular events by poly(ADP-ribose)-binding
proteins. *Mol Aspects Med* **34**, 1066-1087 (2013).
https://doi.org/10.1016/j.mam.2012.12.005
25 Rack, J. G. M. *et al.* (ADP-ribosyl)hydrolases: Structural Basis for Differential
Substrate Recognition and Inhibition. *Cell Chemical Biology* **25**, 1533-1546.e1512
(2018). https://doi.org/https://doi.org/10.1016/j.chembiol.2018.11.001
26 Gagné, J.-P. *et al.* Proteome-wide identification of poly(ADP-ribose) binding proteins
and poly(ADP-ribose)-associated protein complexes. *Nucleic acids research* **36**,
6959-6976 (2008). https://doi.org/10.1093/nar/gkn771
27 Ahel, I. *et al.* Poly(ADP-ribose)-binding zinc finger motifs in DNA repair/checkpoint
proteins. *Nature* **451**, 81-85 (2008). https://doi.org/10.1038/nature06420
28 Aravind, L. The WWE domain: a common interaction module in protein ubiquitination
and ADP ribosylation. *Trends in Biochemical Sciences* **26**, 273-275 (2001).
https://doi.org/https://doi.org/10.1016/S0968-0004(01)01787-X
29 Dasovich, M. & Leung, A. K. L. PARPs and ADP-ribosylation: Deciphering the
complexity with molecular tools. *Molecular Cell* **83**, 1552-1572 (2023).
https://doi.org/https://doi.org/10.1016/j.molcel.2023.04.009
30 Kirby, I. T. & Cohen, M. S. Small-Molecule Inhibitors of PARPs: From Tools for
Investigating ADP-Ribosylation to Therapeutics. *Curr Top Microbiol Immunol* **420**,
211-231 (2019). https://doi.org/10.1007/82_2018_137
31 Liu, Q., van der Marel, G. A. & Filippov, D. V. Chemical ADP-ribosylation: mono-
ADPr-peptides and oligo-ADP-ribose. *Organic & Biomolecular Chemistry* **17**, 5460-
5474 (2019). https://doi.org/https://doi.org/10.1039/c9ob00501c
32 Hagino, R. *et al.* Straightforward Synthesis of the Poly(ADP-ribose) Branched Core
Structure. *ACS Omega* **7**, 32795-32804 (2022).
https://doi.org/10.1021/acsomega.2c04732
33 Oei, S. L. *et al.* NAD+ analogs substituted in the purine base as substrates for
poly(ADP-ribosyl) transferase. *FEBS Letters* **397**, 17-21 (1996).
https://doi.org/https://doi.org/10.1016/S0014-5793(96)01137-4
34 Rieth, S. *et al.* A desthiobiotin labelled NAD+ analogue to uncover Poly(ADP-ribose)
polymerase 1 protein targets. *ChemBioChem* **25**, e202300797 (2024).
https://doi.org/https://doi.org/10.1002/cbic.202300797
35 Depaix, A. & Kowalska, J. NAD Analogs in Aid of Chemical Biology and Medicinal
Chemistry. *Molecules* **24** (2019).
36 Rodriguez, K. M. & Cohen, M. S. Chemical genetic methodologies for identifying
protein substrates of PARPs. *Trends in Biochemical Sciences* **47**, 390-402 (2022).
https://doi.org/https://doi.org/10.1016/j.tibs.2021.07.002
37 Daniels, C. M., Ong, S.-E. & Leung, A. K. L. in *Poly(ADP-Ribose) Polymerase:*
Methods and Protocols (ed Alexei V. Tulin) 79-93 (Springer New York, 2017).
38 Ando, Y. *et al.* ELTA: Enzymatic Labeling of Terminal ADP-Ribose. *Molecular Cell* **73**,
845-856.e845 (2019). https://doi.org/https://doi.org/10.1016/j.molcel.2018.12.022

39 Nowak, K. *et al.* Engineering Af1521 improves ADP-ribose binding and identification
of ADP-ribosylated proteins. *Nature Communications* **11**, 5199 (2020).
https://doi.org/10.1038/s41467-020-18981-w
40 Mohapatra, J. *et al.* Serine ADP-ribosylation marks nucleosomes for ALC1-
dependent chromatin remodeling. *eLife* **10**, e71502 (2021).
https://doi.org/10.7554/eLife.71502
41 Wang, Z. *et al.* Recognition of the iso-ADP-ribose moiety in poly(ADP-ribose) by
WWE domains suggests a general mechanism for poly(ADP-ribosyl)ation-dependent
ubiquitination. *Genes Dev* **26**, 235-240 (2012).
https://doi.org/10.1101/gad.182618.111
42 Gibson, B. A., Conrad, L. B., Huang, D. & Kraus, W. L. Generation and
Characterization of Recombinant Antibody-like ADP-Ribose Binding Proteins.
Biochemistry **56**, 6305-6316 (2017). https://doi.org/10.1021/acs.biochem.7b00670
43 Chiu, S. P., Camacho, C. V. & Kraus, W. L. Development and characterization of
recombinant ADP-ribose binding reagents that allow simultaneous detection of mono
and poly ADP-ribose. *J Biol Chem* **300**, 107609 (2024).
https://doi.org/10.1016/j.jbc.2024.107609
44 Kuttiyatveetil, J. R. A. *et al.* Crystal structures and functional analysis of the ZnF5-
WWE1-WWE2 region of PARP13/ZAP define a distinctive mode of engaging
poly(ADP-ribose). *Cell Reports* **41**, 111529 (2022).
https://doi.org/https://doi.org/10.1016/j.celrep.2022.111529
45 Lu, A. Z. *et al.* Enabling drug discovery for the PARP protein family through the
detection of mono-ADP-ribosylation. *Biochemical Pharmacology* **167**, 97-106 (2019).
https://doi.org/https://doi.org/10.1016/j.bcp.2019.05.007
46 Bonfiglio, J. J. *et al.* An HPF1/PARP1-Based Chemical Biology Strategy for Exploring
ADP-Ribosylation. *Cell* **183**, 1086-1102.e1023 (2020).
https://doi.org/10.1016/j.cell.2020.09.055
47 Longarini, E. J. *et al.* Modular antibodies reveal DNA damage-induced mono-ADP-
ribosylation as a second wave of PARP1 signaling. *Mol Cell* **83**, 1743-1760.e1711
(2023). https://doi.org/10.1016/j.molcel.2023.03.027
48 Longarini, E. J. & Dauben, H. Deciphering ADP-ribosylation signalling. *Nature*
Reviews Molecular Cell Biology **25**, 3-3 (2024). https://doi.org/10.1038/s41580-023-
00630-x
49 Dauben, H., Longarini, E. J. & Matic, I. A chemical biology/modular antibody platform
for ADP-ribosylation signaling. *Trends Biochem Sci* **48**, 910-911 (2023).
https://doi.org/10.1016/j.tibs.2023.06.005
50 Kar, P. *et al.* PARP14 and PARP9/DTX3L regulate interferon-induced ADP-
ribosylation. *The EMBO Journal* **43**, 2929-2953-2953 (2024).
https://doi.org/https://doi.org/10.1038/s44318-024-00126-0
51 Đukić, N. *et al.* PARP14 is a PARP with both ADP-ribosyl transferase and hydrolase
activities. *Science Advances* **9**, eadi2687 (2023).
https://doi.org/10.1126/sciadv.adi2687
52 Ribeiro, V. C., Russo, L. C. & Hoch, N. C. PARP14 is regulated by the PARP9/DTX3L
complex and promotes interferon γ-induced ADP-ribosylation. *The EMBO Journal* **43**,
2908-2928-2928 (2024). https://doi.org/https://doi.org/10.1038/s44318-024-00125-1
53 Hentrich, C. *et al.* Periplasmic expression of SpyTagged antibody fragments enables
rapid modular antibody assembly. *Cell Chem Biol* **28**, 813-824.e816 (2021).
https://doi.org/10.1016/j.chembiol.2021.01.011
54 Dunstan, M. S. *et al.* Structure and mechanism of a canonical poly(ADP-ribose)
glycohydrolase. *Nature Communications* **3**, 878 (2012).
https://doi.org/10.1038/ncomms1889
55 Langelier, M.-F., Planck, J. L., Servent, K. M. & Pascal, J. M. in *Poly(ADP-ribose)*
Polymerase: Methods and Protocols   (ed Alexei V. Tulin) 209-226 (Humana Press,
2011).

56 Lambrecht, M. J. *et al.* Synthesis of Dimeric ADP-Ribose and Its Structure with
Human Poly(ADP-ribose) Glycohydrolase. *Journal of the American Chemical Society*
137, 3558-3564 (2015). https://doi.org/10.1021/ja512528p
57 Gibbs-Seymour, I., Fontana, P., Rack, J. G. M. & Ahel, I. HPF1/C4orf27 Is a PARP-1-
Interacting Protein that Regulates PARP-1 ADP-Ribosylation Activity. *Molecular cell*
62, 432-442 (2016). https://doi.org/10.1016/j.molcel.2016.03.008
58 Suskiewicz, M. J. *et al.* HPF1 completes the PARP active site for DNA damage-
induced ADP-ribosylation. *Nature* **579**, 598-602 (2020).
https://doi.org/10.1038/s41586-020-2013-6
59 Rack, J. G. M. *et al.* Reversal of tyrosine-linked ADP-ribosylation by ARH3 and
PARG. *J Biol Chem* **300**, 107838 (2024). https://doi.org/10.1016/j.jbc.2024.107838
60 Kawamitsu, H. *et al.* Monoclonal antibodies to poly(adenosine diphosphate ribose)
recognize different structures. *Biochemistry* **23**, 3771-3777 (1984).
https://doi.org/10.1021/bi00311a032
61 Hanzlikova, H. *et al.* The Importance of Poly(ADP-Ribose) Polymerase as a Sensor
of Unligated Okazaki Fragments during DNA Replication. *Mol Cell* **71**, 319-331.e313
(2018). https://doi.org/10.1016/j.molcel.2018.06.004
62 Bartlett, E. *et al.* Interplay of Histone Marks with Serine ADP-Ribosylation. *Cell*
Reports **24**, 3488-3502.e3485 (2018).
https://doi.org/https://doi.org/10.1016/j.celrep.2018.08.092
63 Weixler, L. *et al.* Protein and RNA ADP-ribosylation detection is influenced by sample
preparation and reagents used. *Life Science Alliance* **6**, e202201455 (2023).
https://doi.org/10.26508/lsa.202201455
64 Challa, S. *et al.* Development and characterization of new tools for detecting
poly(ADP-ribose) in vitro and in vivo. *eLife* **11**, e72464 (2022).
https://doi.org/10.7554/eLife.72464
623

Figures

Figure 1: Modular Antibodies for versatile, specific and sensitive detection of Poly-ADPr

Scheme describing the coupling process of monovalent SpyTagged antibody, using various BiSpyCatchers to create a variety of formats for multiple approaches.

A: *In Vitro* modified recombinant PARP1, treated w/o PARG to create Poly-/ and Mono-ADPr PARP1 immunoblotted with new anti-Poly-ADPr antibodies in comparison to other available antibodies and reagents. Red represents oversaturated signal. **B:** *In Vitro* modified recombinant PARP1 dot-blotted in dilution series and incubated with new anti-Poly-ADPr antibodies in comparison to competitors. Red represents oversaturated signal.

Figure 2: New highly sensitive Poly-ADPr-specific Antibodies

A: Immunoblotting of U2OS cells treated or not with H_2O_2, w/o Olaparib, lysed in SDS buffer and proceeded with immunoblotting. Antibodies and Reagents used in indicated concentration recommended by manufacturer or publication. HRP-format was used for new anti-poly-ADPr antibodies. Red represents oversaturated signal **B:** Immunofluorescence images of U2OS cells treated or not with H_2O_2, w/o Olaparib followed by methanol fixation. Antibodies and Reagents used in indicated concentration recommended by manufacturer or publication. Synthetic IgG format was used for new anti-poly-ADPr antibodies. Error bar represents SEM. Scale bar represents 10 µm.

Figure 3: Versatile SpyTag System Enables Flexible Co-Detection of Mono- and Poly-ADPr In Cells

A: Coomassie Staining of the same AbD64138 antibody in monovalent unconjugated format (first lane) and bivalent conjugates formats (lane 2-4). **B:** Immunofluorescence co-staining of mono- and poly-ADPr in the same cell using synthetic mouse/rabbit IgG antibody formats.