Cholesterol-mediated Lysosomal Dysfunction in *APOE4* Astrocytes Promotes α-Synuclein Pathology in Human Brain Tissue

Louise A. Mesentier-Louro[1-6,13], Camille Goldman[1-6,13], Alain Ndayisaba[6-9], Alice Buonfiglioli[1-5], Rikki B. Rooklin[1-5], Braxton R. Schuldt[1-5], Abigail Uchitelev[1-5,12], Vikram Khurana[6-11], Joel W. Blanchard[1-6,14, *]

1. Icahn School of Medicine, Mount Sinai, New York, NY, USA
2. Nash Family Department of Neuroscience, Mount Sinai, New York, NY, USA
3. Friedman Brain Institute, Mount Sinai, New York, NY, USA
4. Ronald M. Loeb Center for Alzheimer's Disease, Mount Sinai, New York, NY USA
5. Black Family Stem Cell Institute, Mount Sinai, New York, NY, USA
6. Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD 20815, USA
7. Ann Romney Center for Neurologic Diseases, Brigham and Women's Hospital, Boston, MA, USA
8. Division of Movement Disorders, American Parkinson Disease Association (APDA) Center for Advanced Research and MSA Center of Excellence, Department of Neurology, Brigham and Women's Hospital, Boston, MA, USA
9. Harvard Medical School, Boston, MA, USA
10. The Broad Institute of MIT and Harvard, Cambridge, MA, USA
11. Harvard Stem Cell Institute, Cambridge, MA, USA
12. Macaulay Honors College at Hunter College, New York, NY, USA
13. These authors contributed equally
14. Lead contact

*Correspondence: joel.blanchard@mssm.edu

Summary

The pathological hallmark of neurodegenerative disease is the aberrant post-translational modification and
aggregation of proteins leading to the formation of insoluble protein inclusions. Genetic factors like *APOE4*
are known to increase the prevalence and severity of tau, amyloid, and α-Synuclein inclusions. However, the
human brain is largely inaccessible during this process, limiting our mechanistic understanding. Here, we
developed an iPSC-based 3D model that integrates neurons, glia, myelin, and cerebrovascular cells into a
functional human brain tissue (miBrain). Like the human brain, we found pathogenic phosphorylation and
aggregation of α-Synuclein is increased in the *APOE4* miBrain. Combinatorial experiments revealed that
lipid-droplet formation in APOE4 astrocytes impairs the degradation of α-synuclein and leads to a pathogenic
transformation that seeds neuronal inclusions of α-Synuclein. Collectively, this study establishes a robust
model for investigating protein inclusions in human brain tissue and highlights the role of astrocytes and
cholesterol in *APOE4*-mediated pathologies, opening therapeutic opportunities.

Keywords:

α-Synuclein; iPSC-derived brain model; Lewy Body Dementia; Alzheimer's Disease; Astrocytes; *APOE4*;
Lysosomal dysfunction; Cholesterol metabolism; Neurodegeneration; miBrain.

Introduction.

Alzheimer's disease (AD) is canonically associated with amyloid-β and tau pathology[1]. However, neuronal intracellular inclusions of aggregated α-synuclein (α-Syn) are present in 50-90% of AD cases[2-5]. Phosphorylated α-Syn often aggregates forming Lewy Bodies and Lewy neurites, which are frequently found in the brains of individuals with AD[2-4]. The occurrence of α-Syn inclusions with amyloid-β and tau exacerbates neurodegeneration particularly in brain regions associated with memory and executive functions[5-10]. Indeed, clinical studies show that AD patients with α-Syn pathology exhibit faster cognitive decline compared to those with only amyloid-β and tau pathology[11,12]. The strongest genetic risk factor for late-onset AD, *APOE4*, significantly increases both the prevalence and severity of α-Syn pathology in AD[12,13] and is one of the most well replicated genetic risk factors for Lewy Body Dementia (LBD)[14,15]. However, the mechanisms by which genetic factors like *APOE4* influence the presence and severity of α-Syn pathology are largely unclear. Insight into the mechanisms underlying non-amyloid-β co-pathologies could lead to much needed therapeutic and diagnostic opportunities in AD and LBD.

The development of model systems that faithfully recapitulate α-Syn pathology and the genetic and environmental context of the human brain is essential to uncovering how ancillary genetic factors, such as APOE4, modify α-Syn-driven neurodegeneration. Addition of pre-formed fibrils (PFFs) of α-Syn are commonly employed in mice[16,17] and *in vitro* models[18-20] to induce α-Syn pathology. PFFs are reported to spread across the brain in a prion-like manner corrupting endogenous α-Syn and promoting the propagation of pathology[18,21,22]. However, animal models and *in vitro* systems with α-Syn PFFs are plagued with high variability and often fail to induce phosphorylated α-Syn-rich inclusions while having significant biohazards concerns from the use of prion-like particles[23]. This highlights the need for more physiological models that do not rely on PFFs and can explore multicellular mechanisms of disease. For instance, the induction of neurodegenerative phenotypes in cells cultured in traditional two-dimensional (2D) conditions may be limited by a chemical and mechanical microenvironment that is vastly different from *in vivo* conditions[24]. In contrast, three-dimensional (3D) *in vitro* systems have proven more efficient than 2D systems at replicating key pathological features such as amyloid-β plaques and tau tangles[25,26]. Combining genetic approaches with 3D tissue engineering is a promising alternative for developing more physiological models of neurodegenerative disease.

We recently developed the human **m**ulti-cellular **i**ntegrated **Brain** (**miBrain**), a fully induced pluripotent stem cell (iPSC)-derived 3D human brain tissue that incorporates neurons, glia, and microvasculature into an anatomically precise tissue capable of modeling neurodegeneration in a dish[27]. The miBrain is an *in vitro* human brain tissue containing the major cells and tissues found in human brain tissue including a blood-brain barrier, active neurons with oligodendrocytes, and the brain immune cells- such as microglia and astrocytes. The miBrain can be generated from patient-derived iPSCs. Because it is an engineered tissue, this permits the generation of genetically mixed tissue. For example, coupled with CRISPR-edited iPSC lines, we can generate genetically identical miBrains with AD risk genes such as *APOE4* in specific cell types. We have developed methods to model neuropathological phenotypes associated with amyloid-β and tau in the miBrains[27]. Here, we expanded the miBrain technology, by first developing methods to cryopreserve the miBrain, enabling the production of large batches of full or partial miBrains and dramatically reducing the batch-to-batch variability and improving scalability of the model system. Leveraging this advanced approach, we developed the first highly reproducible model of α-Syn neuropathological phenotypes in an iPSC-derived human brain tissue. We then apply the miBrain to investigate the mechanisms by which *APOE4* promotes increased α-Syn phosphorylation and aggregation in human brain tissue. Our results revealed that *APOE4* increases α-Syn phosphorylation and neuronal inclusions via non-cell-autonomous mechanisms driven by lipid accumulation in APOE4 astrocytes. Our results establish robust methods for modeling α-synuclein

pathological phenotypes in human brain tissue and highlight a causal role of astrocytes and lipids in *APOE4*-
mediated α-Syn pathologies, opening new therapeutic opportunities AD, LBD, and other neurodegenerative
diseases.

Results

Induction of α-Syn intracellular inclusions in a multi-cellular integrated Brain (miBrain) tissue.

Intracellular neuronal inclusions of phosphorylated and aggregated α-Syn have been challenging to
reproduce in human brain cells or tissue, typically requiring very long maturation protocols[28] and the
concomitant use of PFFs[29]. Therefore, we sought to develop a robust, high-fidelity model of α-Syn
intracellular inclusions leveraging the multi-cellular integrated Brain (miBrain). The miBrain is a human brain
tissue generated by encapsulation of iPSC-derived neurons[29,30], astrocytes[31,32], endothelial cells[33-36], mural
cells[33,37], oligodendrocyte precursor cells (OPCs)[38], and microglia[39]. miBrains displayed homogenous TUJ1⁺
neuronal networks and PECAM1⁺ vascular structures (**Fig. 1a**). Staining for multiple markers revealed
interactions and co-localization between S100β⁺ and AQP4⁺ astroglia and PECAM1⁺ vascular structures,
vascular coverage with NG2⁺ mural cells, myelin basic protein (MBP)⁺/neurofilament⁺ neuro-myelin
structures, and the presence of IBA1⁺ microglia in a grid-like pattern (**Fig. 1b**), consistent with the specific
cell types encapsulated in the miBrain. In other platforms such as brain organoids, significant variability in
cell composition have been reported, leading to a high-degree of experimental variability and consistency[40].
Therefore, to minimize miBrain batch-to-batch variability and increase the scalability of the miBrain, we
developed methods to cryopreserve large batches of miBrain tissue containing multiple cell types at a defined
ratio. Off-the-shelf cryopreserved miBrains retain more than 90% cell viability upon thaw and expressed
specific cell markers two weeks after thawing into the 3D culture system (**Extended Data Fig. 1a, b**). The
ratio of neurons to nuclei did not significantly differ between three different batches of thawed miBrain tissue
(p = 0.29, **Extended Data Fig. 1b**).

Phosphorylation of α-Syn on S129 is the predominant pathological modification associated with α-Syn
aggregation and neuronal inclusions, and, therefore, is an established method for detecting pathogenic
transformation of α-Syn[41,42]. Consequently, we first investigated whether the conventionally employed α-
synuclein PFFs[16-20] can increase phosphorylated α-Syn (p-Syn) in the miBrains. Inoculating the miBrain
culture media with α-Syn PFFs significantly increased p-Syn levels when compared to control miBrains (p =
0.0175, **Extended Data Fig. 1c**), indicating the suitability of the tissue to develop α-Syn pathological
phenotypes. However, using this approach, p-Syn was mostly seen as dispersed puncta (**Extended Data
Fig. 1c**) rather than within the typical neuronal inclusions that are hallmarks of synucleinopathies[43]. In
addition, there was a high degree of variability in the presence and abundance of p-Syn (**Extended Data
Fig. 1c**) consistent with other reports comparing the pattern of α-Syn aggregate spreading in PFF models
across different research groups[17]. Therefore, we sought to identify more robust methods for inducing
pathogenic α-Syn phenotypes in the miBrain.

A recent study found that iPSC-derived neurons express low levels of the gene encoding α-Syn (*SNCA)*
compared to adult human brain tissue and, therefore, used *SNCA* overexpression to increase α-Syn to
physiological levels and induce its phosphorylation and neuronal inclusions[29]. Thus, to achieve brain-like
levels of *SNCA* and induce pathological phenotypes, we generated neurons from iPSCs with inducible
expression of *SNCA* bearing the A53>T mutation (A53T), known to increase α-Syn's aggregation
propensity[29,44,45]. SNCA-A53T was fused to a small folding green fluorescent protein (sfGFP, diagram in **Fig.
1c**), enabling live imaging and direct visualization of α-Syn accumulation[46]. We confirmed that α-Syn
phosphorylation is dependent on the α-Syn protein non-amyloid component (NAC) domain and not

influenced by sfGFP (**Extended Data Fig. 1d**). In 2D monocultures, wild-type (WT) and A53T neurons have similar levels of p-Syn, which significantly increased in A53T neurons with the addition of PFFs (**Extended Data Fig. 1d**). In contrast, in the absence of exogenously added PFFs, miBrains with A53T neurons already showed a significant increase in neuronal p-Syn compared to isogenic miBrains harboring WT neurons (p = 0.0007), suggesting that the more physiological 3D environment of the miBrain is more permissive to α-Syn pathology. We found the addition of PFFs in A53T miBrains further increased p-Syn levels compared to WT miBrains with PFFs (p < 0.0001) and A53T miBrains without PFFs (p = 0.038, **Fig. 1c**). The SNCA-A53T-sfGFP signal co-localized with p-Syn (**Fig. 1d** and **Supplementary Video 1**; Control: 13.54% ± 2.37, PFFs: 17.79% ± 2.61, mean of the percentage of colocalization ± SEM, n = 4), indicating that SNCA-A53T undergoes phosphorylation at a similar extent with or without PFFs (p = 0.27, unpaired t-test). Although most of the p-Syn co-localized with sfGFP (Control: 86.7% ± 1.9, PFFs: 88.0% ± 1.4, n = 4), we also found p-Syn that did not co-localize with sfGFP (Control: 13.3% ± 1.9, PFFs: 12.0% ± 1.4, n = 4), indicating that endogenous, non-A53T α-Syn is also phosphorylated. This data shows that A53T miBrains developed p-Syn-rich inclusions via corruption of both induced (A53T) and endogenous (WT) α-Syn without requiring the use of PFFs.

In the human brain, lipid droplets[47] and mitochondria[48] are frequently observed within α-Syn$^+$ inclusions. In A53T miBrains, the neutral lipid marker Lipid Spot overlapped with SNCA-A53T-sfGFP and with p-Syn (**Fig. 1e**). The co-localized volume between Lipid Spot and p-Syn was significantly increased (p = 0.02) in A53T miBrains compared with WT (**Fig. 1e**), consistent with previous reports on p-Syn$^+$/Lipid Spot$^+$ neurotoxic inclusions in iPSC-derived neurons[29]. Furthermore, the volume of aggregated α-Syn overlapping with mitochondrial marker Tom20 was significantly increased in A53T miBrains compared to WT miBrains (p = 0.04, **Extended Data Fig. 1e**), consistent with reports of aggregated α-Syn bound to mitochondria in α-Syn$^+$ inclusions in the human brain[49].

Neuronal inclusions of α-Syn present as spherical dense Lewy Bodies, less dense "pale bodies", and Lewy neurites that can be thread-like or attain a varicose appearance[43]. Through serial live imaging, we observed somatic and neuritic presentations of the SNCA-A53T-sfGFP signal in miBrains, resembling various morphological presentations found in the human brain (**Fig. 1f, i-iii**). During the first 12 weeks, there was a significant increase in the number of α-Syn somatic inclusions (2 weeks: $4.7 \times 10^{-5} \pm 1.9 \times 10^{-6}$, 12 weeks: $8.5 \times 10^{-5} \pm 4.0 \times 10^{-6}$, number of sfGFP$^+$ inclusions normalized by total sfGFP volume, p < 0.0001, unpaired t-test), and no significant changes in the number of α-Syn neuritic inclusions (2 weeks: $1.5 \times 10^{-7} \pm 4.4 \times 10^{-8}$, 12 weeks: $3.4 \times 10^{-7} \pm 1.6 \times 10^{-7}$, p= 0.3; **Fig. 1f**). *In vivo* aggregation of α-Syn within neuronal inclusions promotes neuronal death[50,51]. Soluble biomarkers of cell death (lactate dehydrogenase (LDH)) were significantly higher in media of A53T miBrains compared to WT miBrains (p = 0.0018) and the sfGFP volume was significantly reduced between 2 and 22 weeks of miBrain assembly (p < 0.0001, **Fig. 1f**). Consistent with α-Syn-induced neuronal death we observed a significant reduction in the volume of TUJ1 staining at 2 weeks (p = 0.002) and 24 weeks (p = 0.011) in A53T miBrains when compared to WT (**Fig. 1g**). By 24 weeks, A53T miBrains had significantly increased percentages of the sfGFP volume overlapping with pS129 Syn (p = 0.0095), and aggregated α-Syn (p = 0.0381) in comparison with 2 weeks miBrains (**Fig. 1h**), consistent with aggregation of α-Syn over time. Collectively, these results show that the miBrain can effectively model critical aspects of α-Syn pathology.

APOE4 **increases the phosphorylation and aggregation of** α-**Syn in the miBrain**

APOE4 is a genetic risk factor for α-Syn in LBD[14] and AD[52,53], and it is associated with increased disease severity in human studies, animal models[54,55], and *in vitro* systems[56]. To model and investigate the mechanisms by which *APOE4* promotes α-Syn pathology, we utilized isogenic iPSC lines obtained from an *APOE3/3* individual CRISPR-edited to *APOE4/4*[57,58]. Monocultures of *APOE4/4* A53T neurons showed similar levels of p-Syn as isogenic *APOE3/3* SNCA-A53T neurons with or without PFF exposure (**Extended Data Fig. 2a**). Therefore, we differentiated the isogenic iPSCs into astrocytes, OPCs, endothelial cells, and mural cells observing similar cell type-specific marker expression between the *APOE3/3* and *APOE4/4* genotypes for each cell type (**Extended Data Fig. 2b-c**). We next generated isogenic *APOE3/3* and *APOE4/4* miBrains containing SNCA-A53T neurons. Consistent with clinical studies, *APOE4/4* miBrains with SNCA-A53T neurons showed significantly increased neuronal p-Syn compared to isogenic control *APOE3/3* miBrains (p = 0.0074, **Fig. 2a**). The volume of p-Syn outside the sfGFP mask was also significantly increased in *APOE4/4* miBrains compared to *APOE3/3* (p = 0.0156, **Extended Data Fig. 2d**), indicating that endogenous non-A53T α-Syn is also phosphorylated in the *APOE4/4* miBrains. To assess whether increased phosphorylation of α-Syn in the *APOE4/4* miBrain is a direct cell-autonomous effect of *APOE4/4* neurons, we generated miBrains that were all *APOE4/4* except for *APOE3/3* SNCA-A53T neurons. Strikingly, we observed a significant increase in p-Syn even when *APOE3/3* SNCA-A53T neurons were placed in an otherwise *APOE4/4* miBrain (p = 0.0001, **Extended Data Fig. 2e**).

***APOE4/4 astrocytes are responsible for increased α-Syn pathological phenotypes in human brain tissue**.*

Our results suggest that non-neuronal cell types are responsible for increased p-Syn in the *APOE4/4* miBrains. To identify which *APOE4* cell types promote α-Syn pathology, we generated permutations of the *APOE3/3* miBrains where each cell type was replaced with isogenic *APOE4/4* cells (**Fig 2b**). Consistent with previous results, all *APOE4/4* miBrain had significantly increased p-Syn staining compared to isogenic *APOE3/3* miBrains. Replacing *APOE3/3* neurons, endothelial cells, mural cells, or OPCs with their *APOE4/4* isogenic counterpart did not significantly increase p-Syn immunoreactivity in the miBrain (p = 0.99, **Fig. 2b**). However, selectively replacing *APOE3/3* astrocytes with isogenic *APOE4/4* astrocytes led to a significant increase in p-Syn immunoreactivity (p = 0.014) reaching levels similar to the all *APOE4/4* miBrains (**Fig. 2b**), suggesting that *APOE4/4* astrocytes are responsible for increasing α-Syn pathology in *APOE4* miBrain. Since microglia are implicated in the clearance of α-Syn[59,60], we investigated the effect of the presence or absence of microglia in the isogenic miBrains. The presence of microglia in the *APOE3/3* and *APOE4/4* miBrains did not significantly alter α-Syn phosphorylation (**Extended Data Fig. 2f**). Astrocytes are responsible for uptake and degradation of α-Syn released by neurons[61-63]. We found that GFAP+ *APOE4/4* astrocytes in the *APOE4/4* miBrain have a significantly reduced overlap with SNCA-A53T-sfGFP (p = 0.0348) and adopt an amoeboid-like morphology with increased circularity (p = 0.0398) and GFAP expression (p = 0.0100) compared with *APOE3/3* miBrain astrocytes (**Fig. 2c and Extended Data Videos 2 and 3**), suggesting changes in astrocytic uptake of α-Syn associated with increased reactivity. Collectively, these results highlight a critical role for *APOE4/4* astrocytes in the phosphorylation and aggregation of neuronal α-Syn.

APOE4 astrocytes have impaired processing of exogenous α-Syn.

To investigate the mechanisms by which *APOE4* astrocytes increase the phosphorylation and aggregation
of neuronal α-Syn, we first examined whether altered expression of *SNCA* in astrocytes contributes to
increased abundance and phosphorylation of α-Syn in the *APOE4/4* miBrain and human brain tissue.
Analysis of single-nucleus post-mortem transcriptomics data[64] revealed that *SNCA* mRNA expression is
significantly (p = 0.019) down-regulated in astrocytes from *APOE4* carriers (n = 10) compared to age-
matched *APOE3/3* individuals (n = 8) (**Fig. 3a**). We observed a similar decrease in *SNCA* mRNA expression
in *APOE3/3* and *APOE4/4* iPSC-derived astrocytes[57] (p = 0.017) (**Extended Data Fig. 3a**) suggesting that
transcriptional upregulation of *SNCA* in astrocytes is likely not responsible for increased phosphorylation and
aggregation of α-Syn in the *APOE4* human brain tissue. However, immunoblotting astrocyte monocultures
for α-Syn revealed that *APOE4/4* astrocytes have more total and phosphorylated α-Syn protein compared to
isogenic *APOE3/3* astrocytes from two different individuals (**Fig. 3b; Extended Data Fig. 3b**) suggesting
that post-translational mechanisms underlie increased α-Syn abundance in *APOE4/4* astrocytes.

Astrocytes have a well-described protective and homeostatic function of taking up and degrading neuronal
α-Syn[61-63]. Given this role and our findings that *APOE4/4* astrocytes increase neuronal α-Syn phosphorylation
and co-localize less with SNCA-A53T-sfGFP in the miBrain (**Fig. 2c**), we hypothesized that *APOE4/4*
astrocytes have impaired processing of exogenous α-Syn. To investigate this, we incubated isogenic
APOE3/3 and *APOE4/4* astrocytes with fluorescently labeled α-Syn monomers (αSyn-HiLyte) for 24 hours.
APOE3/3 astrocytes showed a robust ability to take up αSyn within 4.5 hours (p = 0.001) and 6 hours (p <
0.0001) (**Figure 3c**). In contrast, α-Syn monomer uptake by *APOE4/4* astrocytes was significantly impaired.
At 4.5 hours *APOE4/4* astrocytes had no significant (p > 0.99) intracellular α-Syn-HiLyte signal, but it began
to increase by 6 hours (p = 0.005) (**Fig. 3c**). Despite delayed kinetics, at 24 hours *APOE4/4* astrocytes
reached similar uptake levels as *APOE3/3* astrocytes. We then removed the media containing α-Syn and
assessed the degradation of intracellular α-Syn-HiLyte over the following 24h. *APOE3/3* astrocytes had a
significant decrease (p < 0.0001) in the α-Syn-HiLyte signal. In contrast, the signal in *APOE4/4* did not
significantly decrease (p = 0.98) (**Fig. 3c**) suggesting that *APOE4/4* astrocytes have impaired degradation of
α-Syn.

APOE4 astrocytes have impaired lysosomal function and release pathogenic α-Syn.

Since the two main systems for the degradation of intracellular proteins are the lysosomal and proteasomal
pathways[65], we asked whether α-Syn was cleared via lysosomal or proteasomal degradation in astrocytes.
We treated astrocytes with bafilomycinA1, a lysosomal inhibitor, or with MG-132, a proteasomal inhibitor.
Lysosomal inhibition blocked the degradation of α-Syn-HiLyte (p = 0.02), unlike proteasomal inhibition, which
had no significant effect (p = 0.07) on the degradation of α-Syn-HiLyte signal (**Extended Data Fig. 3c**). This
confirms previous reports that α-Syn is primarily degraded through the endolysosomal pathway in astrocytes
and demonstrates that *APOE4*-driven lysosomal dysfunction leads to α-Syn accumulation.

We hypothesized that the endolysosomal pathway of *APOE4/4* astrocytes is impaired in comparison to
APOE3/3 astrocytes. To assess lysosomal function, we measured the lysosome-specific proteolytic cleavage
of DQ-BSA and found *APOE4/4* astrocytes have significantly decreased lysosomal proteolytic activity
compared to isogenic control *APOE3/3* astrocytes in two isogenic donor lines (p= 0.003 and p < 0.0001, **Fig.**
3d). Consistent with this, immunoreactivity against the lysosomal-associated protein LAMP1 was significantly
reduced in *APOE4/4* astrocytes compared to isogenic *APOE3/3* astrocytes (p = 0.0002 and p = 0.0001,
Extended Data Fig. 3d). Likewise, staining with LysoSensor Green, a lysosomal dye that fluoresces most
brightly at an acidic lysosomal pH of ~5.2, revealed that *APOE4/4* astrocytes have significantly decreased

LysoSensor Green signal, indicating decreased lysosomal acidity compared to *APOE3/3* across isogenic *APOE3/3* and *APOE4/4 astrocytes* generated from 3 different individuals (p < 0.0001, p = 0.002, p = 0.0002, **Extended Data Fig. 3e**). These results suggest *APOE4/4* astrocytes have decreased LAMP1 positive lysosomes with decreased lysosomal acidity, contributing to impaired lysosomal proteolytic function.

Since phosphorylated α-Syn is used as a measurement of α-Syn pathology, we next determined whether astrocytes can phosphorylate neuronal α-Syn. Therefore, we exposed *APOE3/3* and *APOE4/4* astrocytes to fresh media or conditioned media from *SNCA*-A53T neurons. At baseline, *APOE4/4* astrocytes have more endogenous p-Syn compared to *APOE3/3* astrocytes (**Fig. 3e;** p < 0.0001). *APOE4/4* astrocytes incubated with media previously exposed to *SNCA*-A53T neurons for 3 days, showed a further increase in p-Syn (p = 0.007) while *APOE3/3* astrocytes were unaffected, (p = 0.232) (**Fig. 3e**) indicating that *APOE4/4* astrocytes can increase α-Syn phosphorylation. However, in the miBrain and post-mortem human brain, inclusions of α-Syn are primarily found inside neurons. We found that cell culture media collected from *SNCA*-A53T neurons and then inoculated onto fresh *SNCA*-A53T neuronal cultures itself does not increase phosphorylation of neuronal α-Syn (**Fig. 3f**), suggesting that non-cell-autonomous mechanisms likely modify α-synuclein's pathogenicity. Therefore, we hypothesized that *APOE4/4* astrocytes fail to degrade α-Syn due to impaired endolysosomal function. Instead, *APOE4/4* astrocytes phosphorylate α-Syn and release the more pathogenic forms of a-Syn into the extracellular space allowing them to be taken up by neurons and promote the formation of neuronal α-Syn inclusions. To test this hypothesis, we first exposed *SNCA*-A53T neurons to conditioned media from only *APOE3/3* or *APOE4/4* astrocytes. Astrocyte-conditioned media itself did not induce significant p-Syn in *SNCA-A53T* neurons, and no significant difference was observed between *APOE3/3* and *APOE4/4* conditioned media (**Extended Data Fig. 3f**). We reasoned the low levels of α-Syn produced by astrocytes alone is not sufficient to induce phosphorylation and aggregation of neuronal α-Syn. Therefore, we performed a double conditioned media experiment where conditioned media was first collected from *SNCA*-A53T neurons and subsequently cultured with either *APOE3/3* or *APOE4/4* astrocytes. This double-conditioned media was then collected and inoculated onto fresh *SNCA*-A53T- neuron monocultures (**Fig 3f**). Neurons grown in *APOE3/3* double-conditioned media did not show a significant increase in α-Syn phosphorylation (p = 0.601). In contrast, *APOE4/4* double-conditioned media led to a 2.371-fold increase (± 0.315; SEM) in α-Syn phosphorylation (p = 0.0016). To assess whether the induction of p-Syn by *APOE4/4* double-conditioned media was a failure of astrocytes to degrade α-Syn or a modification of α-Syn, we analyzed the α-Syn species in the double-conditioned media via dot blot. Although the total levels of α-Syn in the media were not different between *APOE3/3* and *APOE4/4* astrocytes (p = 0.894), p-Syn and aggregated α-Syn were both significantly increased (p < 0.0001 and p = 0.0051) in media conditioned by *APOE4/4* astrocytes compared to *APOE3/3* astrocytes (**Extended Data Fig. 3g**). *APOE3/3* astrocyte double-conditioned media appeared not to change having similar p-Syn and aggregated α-Syn levels to the original *SNCA*-A53T media that was not conditioned by astrocytes (p = 0.951 and p = 0.894; **Extended Data Fig. 3g**). Collectively, these results suggest that *APOE4/4* astrocytes fail to degrade neuronal a-Syn, instead increase its phosphorylation and aggregation, and secrete these more pathological forms of α-Syn which are taken up by neurons and promote the formation of a-Syn inclusions.

Cholesterol accumulation in APOE4 astrocytes leads to dysfunctional α-Syn processing and neuronal pathology.

We next investigated how APOE4 in astrocytes leads to lysosomal dysfunction and the spread of αSyn pathogenic forms. APOE is primarily expressed in astrocytes in the brain,[66] shuttling cholesterol to neurons to maintain membranes and synapses[67,68]. APOE4 displays lower lipid transport affinity and binding

capacity[69], associated with cholesterol accumulation in astrocytic lysosomes and impaired lysosomal function[58,70]. Consistent with previous reports, transcriptomic analysis showed that genes in pathways associated with lipid storage and transport are differentially expressed compared to isogenic APOE3/3 astrocytes (**Extended Data Fig. 4a**)[58], as well as response to lipopolysaccharide and oxidative stress, which have been associated with a neurotoxic reactive phenotype[71,72]. Consistent with a dysfunctional lipid metabolism, we found that *APOE4/4* astrocytes have a significant increase in the accumulation of BODIPY-cholesterol compared to isogenic *APOE3/3* astrocytes generated from two different individuals (p = 0.001 and p = 0.002, **Fig. 4a**).

Given our findings that *APOE4/4* astrocytes have increased cholesterol and impaired endolysosomal function, resulting in increased pathogenic α-Syn secretion, we hypothesized that intracellular accumulation of cholesterol leads to impaired degradation of α-Syn by astrocytes. To test this hypothesis, we treated *APOE4/4* astrocytes with various compounds to reduce cholesterol bioavailability, either by sequestering cholesterol (2-hydroxypropyl-β-cyclodextrin (2HβCD) and methyl-β-cyclodextrin (MβCD), inhibiting cholesterol biosynthesis (atorvastatin), or promoting cholesterol efflux (efavirenz, LXR-623, and T0901317). BODIPY staining confirmed a decrease in lipid droplets, which include cholesterol, in *APOE4/4* astrocytes following treatment with 2HβCD (p = 0.0013) and MβCD (p = 0.0063), to levels comparable to *APOE3/3* astrocytes (**Fig. 4b**). 2HβCD and MβCD significantly increased *APOE4/4* lysosomal proteolytic activity, as measured by the BSA-DQ assay, compared to vehicle treated *APOE4/4* astrocytes (2HβCD: p = 0.0005; MβCD: p = 0.046) (**Fig. 4c**). However, treatment with the other compounds had no effect, or a negative effect, on *APOE4/4* lysosomal proteolytic activity. We confirmed that MβCD increases lysosomal proteolytic activity in *APOE4/4* astrocytes generated from a second individual (p = 0.0093), although the effect of 2HβCD was not significant (p = 0.357; **Extended Data Fig 4b**). These findings suggest that improvement of proteolytic function may be selective to cyclodextrin treatments.

We further evaluated the effect of 2HβCD and MβCD on lysosomes by staining treated astrocytes with the live-cell endolysosomal dye, LysoTracker. *APOE4/4* astrocytes treated with MβCD had an increase in lysosomal staining intensity compared to control *APOE4/4* astrocytes (p = 0.044) (**Fig. 4d**). However, 2HβCD treatment did not have a significant effect on endolysosome intensity (p = 0.51). In astrocytes differentiated from a second individual, both cyclodextrin treatments significantly increased endolysosomal staining intensity in *APOE4/4* astrocytes (2HβCD: p < 0.0001; MβCD: p < 0.0001) to levels comparable to *APOE3/3* astrocytes (**Extended Data Fig. 4c**). These findings suggest that lowering cholesterol burden in *APOE4/4* astrocytes with 2HβCD or MβCD treatment is sufficient to increase lysosomal and proteolytic function in *APOE4/4* astrocytes.

To evaluate the effect of 2HβCD and MβCD treatment on α-Syn uptake and degradation, we measured α-Syn-HiLyte uptake after 24 hours in astrocytes. The intensity of intracellular α-Syn-HiLyte after 24 hours significantly increased in 2HβCD and MβCD treated *APOE4/4* astrocytes (2HβCD: p = 0.002; MβCD: p = 0.0003) compared to vehicle-treated *APOE4/4* astrocytes (**Fig 4e**). This effect was replicated in isogenic astrocytes from a second individual (2HβCD: p = 0.0004; MβCD: p = 0.0003) (**Extended Data Fig 4d**). To validate that the α-Syn was being endocytosed into the endolysosomal pathway, we incubated astrocytes with α-Syn-HiLyte for 4 hours and co-stained with LysoTracker. In agreement with our other findings that *APOE4/4* astrocytes have impaired α-Syn uptake, the percentage of α-Syn-HiLyte that co-localized with LysoTracker decreased in *APOE4/4* astrocytes compared to *APOE3/3* astrocytes (p = 0.0395; **Extended Data Fig 4e**). However, *APOE4/4* astrocytes treated with MβCD showed increased colocalization of α-Syn-HiLyte with LysoTracker compared to vehicle-treated *APOE4/4* astrocytes (p = 0.0343). Strikingly APOE4/4 astrocytes treated with MβCD had similar α-Syn lysosomal localization as vehicle-treated *APOE3/3* astrocytes (p = 0.946; **Extended Data Fig 4e**). Taken together, these results suggest that pharmacologically

reducing intracellular cholesterol in *APOE4/4* astrocytes increase α-Syn uptake and degradation potentially reducing the accumulation of neurotoxic forms of α-Syn.

Our findings implicate astrocytes as a critical cell type in α-Syn processing in the genetic context of *APOE4.* However, α-Syn pathology in Lewy body diseases is characterized by inclusions found within neuronal projections and soma. To investigate whether improved lysosomal activity in cyclodextrin-treated astrocytes affected neuronal α-Syn phosphorylation, we returned to the miBrain model. After 7 days of treatment with MβCD, the levels of neuronal phosphorylated α-Syn were significantly reduced in MβCD treated *APOE4/4* miBrains (p = 0.031) when compared to untreated *APOE4/4* miBrains and restored phosphorylation to levels that were not statistically different from *APOE3/3* miBrains (p = 0.099) (**Fig. 4f**). These results show that MβCD treatment reduces α-Syn phosphorylation in neurons likely via restoration of lysosomal activity and α-Syn processing in astrocytes, alleviating cells of the cytotoxic burden of α-Syn aggregation. This may open a therapeutic avenue for cholesterol-lowering pharmacological interventions in the treatment of synucleinopathies and AD with Lewy Bodies, particularly in *APOE4* carriers.

Discussion:

By combining stem cell and genetic engineering, we extended our recently developed miBrain system[27] to model α-Synuclein pathological phenotypes and dissect disease-relevant cellular and molecular mechanisms in human brain tissue in a dish. miBrains containing neurons, glia, and vascular cells displayed trackable α-Syn pathological forms in neurons and neuronal death as early as 2 weeks and as late as 6 months in culture. Cryopreservation of pre-assembled tissue allowed better reproducibility and decreased variability of our methods, enabling complex multivariate experiments. Using a genetic mix-and-match permutation approach that is unique to miBrains, we found that *APOE4* promotes neuronal α-Syn accumulation specifically via astrocytes. *APOE4*-induced cholesterol accumulation causes endolysosomal dysfunction and impaired α-Syn processing in astrocytes. This leads to the astrocytic secretion of pathogenic α-Syn that seeds α-Syn aggregation and phosphorylation on neurons. We found that pharmacologically increasing cholesterol efflux restored astrocytic endolysosomal function and prevented the accumulation of neuronal p-Syn in human brain tissue. These results establish a causal link between cholesterol dysregulation and α-Syn pathology in *APOE4* carriers, which may influence the progression of classical synucleinopathies and AD with Lewy Bodies.

Cholesterol dysregulation in *APOE4* astrocytes has been related to a reduced expression of lipid transport genes[57], reduced cholesterol efflux[58], and increased accumulation of cholesterol in lysosomes[70]. This was associated with increased secretion of inflammatory cytokines and reduced amyloid-β processing[73]. Genetically induced cholesterol accumulation has been shown to promote an increased secretion of exosomes[74]. Cholesterol accumulation might convert astrocytes into a secretory phenotype, including releasing pathogenic α-Syn and other cargo. While it is known that neurons can secrete pathogenic α-Syn[75] that is taken and processed by astrocytes[61-63], to our knowledge, the release of pathogenic α-Syn by dysfunctional astrocytes has not been previously described.

Our results suggest that *APOE4* leads to a disease-relevant astrocytic phenotype characterized by endolysosomal dysfunction driven by cholesterol accumulation, leading to suboptimal processing and secretion of pathogenic α-Syn that is further taken by neurons, exacerbating Lewy-like pathology. We found pharmacological cholesterol reduction using methyl-β-cyclodextrin prevented α-Syn accumulation in *APOE4* human brain tissue. Clinical trials on cyclodextrins have primarily focused on NPC1 disease, with a recent

Phase I/II trial describing improved clinical symptoms and mild-to-moderate adverse effects after intravenous administration of hydroxypropyl-β-cyclodextrin[76]. Other cholesterol-lowering drugs such as statins remain the primary approach to reduce cholesterol in patients without NPC1. In a mouse model of α-Syn pathology, feeding a high-fat diet increased α-Syn aggregation and that was reduced by treatment with brain-penetrating statins, suggesting that statins can lower α-Syn aggregation by reducing brain cholesterol[77]. A meta-analysis study reported that statins can reduce the risk of Parkinsonism in older adults and that this effect is mediated by reduced atherosclerosis[78], but a randomized clinical trial evaluating the benefit of the most used type of statin, simvastatin, found no evidence to support its use as a disease-modifying therapy in Parkinson's disease[79]. We discovered that *APOE4* increases α-Syn pathology via cholesterol dysregulation, suggesting that the *APOE* genotype may impact whether cholesterol-lowering medications can benefit classic synucleinopathies and co-pathological presentations of pathogenic α-Syn such as AD with Lewy Bodies.

We present a novel, scalable stem cell-based tissue platform capable of modeling human neurodegenerative phenotypes in a dish. Collectively, our findings establish a mechanistic link between cholesterol accumulation and α-Syn pathology mediated by *APOE4* astrocytes and offer therapeutic opportunities for *APOE4* carriers with α-Syn pathology in classical synucleinopathies like LBD, as well as in AD with Lewy Bodies and other neurodegenerative conditions. Future optimization of the miBrain system to enable microvascular perfusion and support extended experimentation and drug testing will further enhance its utility, paving the way for deeper exploration of neurodegenerative mechanisms and therapeutic testing.

Acknowledgments

We would like to thank Tatyana Kareva for technical support and all members of the Blanchard Lab for scientific discussions. We would like to thank Elena Coccia for revising the manuscript. We acknowledge the use of BioRender for generation of illustrative cartoons.

This work has been funded in part with federal funds from NASA under contract #80ARC022CA004 titled "Identification of Biomarkers and Pathological Mechanisms via Longitudinal Analysis of Neurological and Cerebrovascular Responses to Neurotoxic Stress Using a Multi-cellular Integrated Model of the Human Brain." This research was funded in whole or in part by Aligning Science Across Parkinson's (ASAP-00458) through the Michael J. Fox Foundation for Parkinson's Research (MJFF). For open access, the authors have applied a CC-BY public copyright license to all Author Accepted Manuscripts arising from this submission. This work was also supported by funding from the NIH/NINDS (R01NS114239 and UH3NS115064), and the CureAlz Fund. C.G. was supported by funding from the NIH/NIA (T32AG04968) and NIH/NINDS (F31NS13090).

Author Contributions

Conceptualization: L.A.M.L, C.G., and J.W.B.; Methodology: L.A.M.L., C.G., J.W.B., A.N., and V.K.; Investigation: L.A.M.L., C.G., A.B., R.B.R., B.R.S., and A.U.; Writing – Original Draft: L.A.M.L and C.G.; Writing – Review & Editing: L.A.M.L., C.G., J.W.B., A.N., and V.K; Supervision: J.W.B.; Funding Acquisition: J.W.B.

Declaration of Interests

V.K. is a cofounder of and senior advisor to DaCapo Brainscience and Yumanity Therapeutics, companies focused on CNS diseases. L.A.M.L, J.W.B., C.G., and R.B.R. are inventors on patent applications filed by Mount Sinai Innovation Partners on the methods described in this study. A.N. and V.K. are inventors on a patent application filed by Brigham and Women's Hospital related to the induced inclusion iPSC models.

402 **Methods:**

403 Key Resources Table

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Chemicals, peptides, and recombinant proteins		
2-hydroxypropyl-β-cyclodextrin	Sigma	Cat# C0926
Accutase	Stemcell	Cat# 07920
Activin A	Peprotech	Cat# 120-14P
Astrocyte Growth Supplement	ScienCell	Cat#1852
Astrocyte Medium (AM)	ScienCell	Cat# 1801
a-Synuclein (1-140) HiLyte Fluor 488	Anaspec	Cat# AS-55457
Atorvastatin	Sigma	Cat# SML3030
B27	Gibco	Cat# 17504044
B27 without Vitamin A	Gibco	Cat# 12587010
Bafilomycin A1	Millipore	Cat# B1793
BDNF	Peprotech	Cat# 450-02
Biotin	Sigma	Cat#B4639
Blasticidin	Gibco	Cat # A1113903
BMP4	Peprotech	Cat# 120-05ET
BODIPY-Cholesterol	Cayman	Cat# 24618
CD Lipids	Gibco	Cat# 11905031
CHIR99021	Tocris	Cat# 4423
CNTF	Peprotech	Cat# 450-13
DAPT	Cayman	Cat# 13197
Dibutyryl cAMP	Biogems	Cat# 1698950
DMEM	Gibco	Cat# 11965092
DMEM/F12 with GlutaMAX	Gibco	Cat# 10565018
Doxycyline	Millipore	Cat# D3072
DQ-BSA Red	Thermo Scientific	Cat # D12051
DRAQ5	Thermo Scientific	Cat# 52251
Efavirenz	MedChem Express	Cat# HY10572
FGF-basic	Peprotech	Cat# 100-18B
Forkskolin	R&D	Cat# 1099/10
GDNF	Peprotech	Cat# 450-10
Gelxtrex	Gibco	Cat# A1413201
GlutaMAX	Gibco	Cat# 35050061
HGF	Peprotech	Cat# 100-39H
Hoechst33342	Thermo Scientific	Cat# 62249
Human Endothelial SFM	Gibco	Cat# 11111044
IGF-1	Peprotech	Cat# 100-11
Insulin	Sigma	Cat# I9278
Laminin	Gibco	Cat# 23017015

L-Ascorbic Acid	Fisher Scientific	Cat# BP351
LipidSpot	Biotium	Cat# 70069-T
LDN193189	Tocris	Cat# 6053
Lipofectamine Stem Transfection Reagent	Invitrogen	Cat# STEM00001
LXR-623	MedChem Express	Cat# HY-10629
LysoSensor Green DND-189	Thermo Scientific	Cat# L7535
LysoTracker DeepRed	Thermo Scientific	Cat# L12492
MEM-Non-Essential Amino Acids	Gibco	Cat#111400050
Methyl-β-cyclodextrin	Cayman	Cat #21633
N2	Gibco	Cat# 17502048
Neurobasal	Gibco	Cat# 21103049
NT3	Peprotech	Cat# 450-03
PDGF-AA	Peprotech	Cat# 100-13A
PDGF-BB	Peprotech	Cat# AF-100-14B
Penicillin-Streptomycin	Gibco	Cat# 15140122
Puromycin	Gibco	Cat# A1113803
Retinoic Acid	MIllipore	Cat# R2625
SAG	Cayman	Cat# 11914
SB431542	Stemgent	Cat# 04-0010
StemFlex	Gibco	Cat# A3349401
T0901317	MedChem Express	Cat# HY-10626
TrypLE Select	Gibco	Cat# 12563011
VEGF-A	Peprotech	Cat#100-20
Y-27632	Tocris	Cat# 1254
Antibody		
a-Synuclein (MJFR1)	Abcam	Cat# ab138501
a-Synuclein (phospoho S129) (MJF-R13)	Abcam	Cat# ab168381
a-Synuclein pS129 (D1R1R)	CellSignaling	Cat# 23706
a-Synuclein pS129 (EP1536Y)	Abcam	Cat# ab51253
aSynuclein aggregate (MJFR-14-6-4-2)	Abcam	Cat# ab209538
AQP4	Invitrogen	Cat# PA5-53234
CD31/PECAM-1	R&D Systems	Cat# AF806
CD49f	BioLegend	Cat# 313602
GFAP	Invitrogen	Cat# 14-9892-82
IBA1	NovusBio	Cat# NB100-1028
LAMP1	NovusBio	Cat# NBP2-25183
MAP2	Biolegend	Cat# 822501
MBP	EMD Millipore	Cat# AB9348
MRF	EMD Millipore	Cat# ABN45
Neurofilament-H	BioLegend	Cat# 822601
NG2	EMD Millipore	Cat# Ab5320
S100β	Sigma-Aldrich	Cat# 2532

Tom20 (F-10)	Santa Cruz Biotechnologies	Cat# sc-17764
Tubulin β 3 (TUBB3) (TUJ1)	BioLegend	Cat# 801202
VECAD	R&D Systems	Cat# AF938
β-Actin	Invitrogen	Cat# am4302
Critical commercial assays		
Criterion 4-15% TGX precast gel	BioRad	Cat# 5671084
LDH-Glo Cytotoxicity Assay	Promega	Cat# J2381
Pierce BCA Protein Assay	Thermo Scientific	Cat# 23223
Experimental models: Cell lines		
A53T-1A CORR28	Harvard Medical School	RRID: CVCL_E3CE
AG09173 iPSCs	Massachusetts Institute of Technology	RRID: CVCL_4L66
ADRC5 iPSCs	UCI	
KOLF2.1J iPSCs	Jackson Laboratories	RRID:CVCL_B5P3
Recombinant DNA		
piggyBac-rtTA (4th_Gen)-NGN2-2A-PURO-IRES-SNAP	Addgene	Cat# 209077
piggyBac-rtTA (4th_Gen)-SNCA(A53T)-sfGFP-IRES-NGN2-puro	Addgene	Cat# 209080
piggyBac-rtTA (4th_Gen)-SNCA(A53T-DNAC)-sfGFP-IRES-NGN2-puro	Addgene	Cat# 209081
PB_iETV2_P2A_GFP_Puro	Addgene	Cat# 168805
Software and algorithms		
Fiji	Schindelin, J. et al., 2012	imagej.net/ software/fiji/
Nikon NIS Elements software	Nikon	microscope.healthcare.nikon.com
GraphPad Prism	GraphPad Software	graphpad.com
Other		
48-well plate	P48G-1.5-6-F	MatTek
96-well plate with cover glass thickness polystyrene bottom	Greiner Bio-One	655096
Ti2E Ax R Confocal Microscope	Nikon	microscope.healthcare.nikon.com

Resource availability

Lead Contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the
lead contact, Dr. Joel W. Blanchard (joel.blanchard@mssm.edu).

Material availability

This study did not generate new unique reagents

Data and code availability

- All raw data will be made available on Mendeley upon paper acceptance and will be made publicly available.
- Original code relating to scRNAseq pseudobulk analysis will be made available on Mendeley upon paper acceptance and will be publicly available.
- Any additional information required to reanalyze the data reported in this paper is available from the lead contact upon request.

Human iPSC cultures

All human iPSCs were maintained in feeder-free conditions in StemFlex™ medium (Gibco) on Geltrex™ Matrix (Thermo Fisher Scientific) pre-coated plates. All iPSC lines used in this study are listed in the Key Resource Table. CRISPR/Cas9 genome editing was performed as previously described[57]. Human iPSCs were grown as colonies in StemFlex™ medium until they reached 60-70% confluency. At this point, iPSCs were either passaged for maintenance using 0.5 mM EDTA to gently lift colonies or harvested using Accutase™ cell detachment solution for 5-10 minutes at 37C to start a differentiation protocol as singularized cells.

Differentiation of human iPSCs into neurons

Neuron differentiation was adapted from Zhang *et al*.[30] and Lam *et al*.[29]. Briefly, iPSCs were transfected with PiggyBac plasmids to confer doxycycline-inducible expression of the Neurogenin- 2 gene (*NGN2*, Addgene Plasmid #209077) alone or combination with SNCA-A53T-sfGFP (Addgene Plasmid: 209080) or A53T-DNAC-SNCA-sfGFP (Addgene Plasmid: 209081), using Lipofectamine™ Stem Transfection Reagent. Briefly, dissociated iPSCs were plated at ~104,000 cells/cm^2 onto Geltrex™-coated plates, in StemFlex™ supplemented with 10 μM Y27632 and 5 μg/mL doxycycline (day 0). At day 1, medium was replaced with Neurobasal N2B27 medium (Neurobasal, 1x B-27, 1x N-2, 1x MEM-NEAA, 1x GlutaMAX, 1% penicillin-streptomycin) supplemented with 10uM SB431542, 100 nM LDN, 5 μg/mL doxycycline, 5 μg/mL Blasticidin. On day 2, the medium was replaced with Neurobasal N2B27 media supplemented with 10uM SB431542, 100 nM LDN, 5 μg/mL doxycycline, 1 μg/mL puromycin. On days 3-6, the medium was replaced daily with Neurobasal N2B27 media supplemented with 5 μg/mL doxycycline, and 1 μg/mL puromycin. At day 7, cells were dissociated with Accutase and either seeded into miBrains (see bellow) or seeded into 2D monocultures on Poly-L-Ornithine and Laminin pre-coated plates at 156,250 cells/cm^2, in Neurobasal N2B27 with 5 μg/mL doxycycline and 10 μM Y27632. For 2D cultures, on day 8, wells were gently topped with Neurobasal N2B27 supplemented 20ng/mL BDNF, 20ng/mL GDNF, 1mM dcAMP, 2ug/mL Laminin, and 1 uM AraC, using the same volume of medium in the wells. At day 11, media is replaced with Neurobasal N2B27 supplemented with 10ng/mL BDNF, 10ng/mL GDNF, 0.5 mM dcAMP, 1ug/mL Laminin. This medium was used for half-media changes every 3-4 days.

Differentiation of human iPSCs into astrocytes

Astrocytes were generated using previously published protocols for iPSC-derived NPC (Chambers *et al*.)[31] and astrocyte (TCW *et al*.)[32] differentiation. Briefly, dissociated iPSCs were plated at 100,000 cells/cm^2 onto Geltrex™-coated plates, in pre-warmed StemFlex™ supplemented with 10 μM Y27632. Cells were fed every other day with StemFlex™ until they reached >95% confluence. Once cells reached confluence, the medium was replaced with NPC medium (1:1 DMEM/F12: Neurobasal Medium, 1x N-2 Supplement, 1x B-27 Serum-Free supplement, 1x GlutaMAX Supplement, 1x MEM-NEAA, 1% penicillin-streptomycin) supplemented with 10 μM SB43152 and 100 nM LDN193189 (day 0). From days 1 to 9, cells are fed daily with NPC medium

plus 10 µM SB43152 and 100 nM LDN193189. At day 10, cells were split with Accutase and replated onto
fresh Geltrex™-coated plates, in NPC media supplemented with 20 ng/mL bFGF and 10 µM Y27632. From
days 11 to 13, cells were fed with NPC media plus 20 ng/mL bFGF. At day 14, cells were split with Accutase
and re-seed onto fresh Geltrex™-coated plates, in NPC media plus 20 ng/mL bFGF and 10 µM Y27632.
Starting from day 15, cells were fed every 2-3 days with Astrocyte Medium (AM, ScienCell) and passaged
using Accutase once they reached 90% confluence. From this point, NPCs were fully differentiated into
astrocytes in 30 days. NPCs and fully differentiated astrocytes were cryopreserved in the freezing medium
consisting of 90% knockout serum replacement (KSR) and 10% dimethyl sulfoxide (DMSO).

Differentiation of human iPSC into brain microvascular endothelial cells

Brain endothelial cell differentiation was adapted from the protocols from Blanchard *et al.*, Qian *et al.,* and
Wang *et al*. [33,36,80]. Briefly, iPSCs were transfected with a PiggyBac plasmid to confer doxycycline-inducible
expression of the ETS variant transcription factor 2 (*ETV2*, Addgene Plasmid #168805), using
Lipofectamine™ Stem Transfection Reagent. Inducible ETV2-iPSCs were grown until 60-70% confluency,
dissociated with Accutase, and plated at 20,800 cells/cm^2 onto Geltrex™-coated plates in StemFlex™
supplemented with 10 µM Y27632 (day 0). On day 1, the medium was replaced with DeSR1 medium
(DMEM/F12 with GlutaMAX, 1× MEM-NEAA, 1× penicillin-streptomycin) supplemented with 10 ng/mL BMP4,
6 µM CHIR99021, and 5 µg/mL doxycycline. On day 3, the medium was replaced with DeSR2 medium
(DeSR1 media, 1x N-2, 1× B-27) supplemented with 5 µg/mL doxycycline. At days 5 and 7, the medium was
replaced with hECSR medium (Human Endothelial Serum-free Medium, Gibco, 1× MEM-NEAA, 1× B-27,
1% penicillin-streptomycin) supplemented with 50 ng/mL VEGF-A, 2 µM Forskolin, and 5 µg/mL doxycycline.
At day 8, cells were dissociated using Accutase and re-seed onto fresh Geltrex™-coated plates in hECSR
supplemented with 50 ng/mL VEGF-A and 5 µg/mL doxycycline. This medium was used for every 2-3 days
medium change to maintain cells for up to 1 week until ready for tissue assembly (miBrain or JAMs).

Differentiation of human iPSCs into mural cells

Mural cells were differentiated using previously published protocol from Patsch *et al.*[37]. Dissociated iPSCs
were plated at 37,000 to 40,000 cells/cm^2 onto Geltrex™-coated plates, in StemFlex™ supplemented with
10 µM Y27632 (day 0). On day 1, the medium was replaced with N2B27 medium (1:1 DMEM/F12:
Neurobasal media, 1x B-27, 1x N-2, 1x MEM-NEAA, 1x GlutaMAX, 1% penicillin-streptomycin)
supplemented with 25 ng/mL BMP4 and 8 µM CHIR99021. At days 3 and 4, the medium was replaced with
N2B27 media supplemented with 10 ng/mL Activin A and 10 ng/mL PDGF-BB. At day 5, mural cells were
dissociated with Accutase and re-seeded onto fresh 0.1% gelatin-coated plates at 35,000 cells/cm^2, in N2B27
supplemented with 10 ng/mL PDGF-BB. This medium was used every 2-3 days medium was change for
another 5–7 days. Cells were then banked in freezing medium (90% KSR/ 10% DMSO) and expanded in
N2B27 until ready for tissue assembly (miBrain or JAMs).

Differentiation of human iPSCs into oligodendrocyte progenitor cells (OPCs)

OPC differentiation was adapted from Douvaras et al, 2014[81]. Briefly, iPSCs were dissociated into single
cells using Accutase and seeded at near-confluent density. Differentiation began the next day (designated
as day 0) by culturing the cells in DMEM/F12 (1:1) medium supplemented with N2, 10 µM SB431542, 100
nM LDN 193189, and 100 nM all-trans retinoic acid (RA), with daily medium changes. On day 8, 1 µM SAG
was added to the differentiation medium, maintaining the presence of 10 µM SB431542 and 100 nM LDN
193189. By day 12, adherent cells were detached and transferred to low-attachment plates to form cell
spheres. These spheres were cultured in DMEM/F12 (1:1) medium containing N2, RA, and SAG. On day 30,

spheres were plated onto poly-L-ornithine/laminin-coated plates to allow cells to migrate outward. At this stage, the medium was replaced with DMEM/F12 (1:1) supplemented with N2, B27, 10 ng/ml PDGF-AA, 10 ng/ml IGF, 5 ng/ml HGF, 10 ng/ml NT3, 25 µg/ml insulin, 100 ng/ml biotin, 1 µM cAMP, and 60 ng/ml T3. By day 75, cells were harvested, dissociated, and purified using NG2-specific magnetic-activated cell sorting (MACS). The enriched cells were expanded in DMEM/F12 (1:1) medium supplemented with N2, B27 without Vitamin A, 10 ng/ml PDGF-AA, 10 ng/ml β-FGF, and 10 ng/ml NT3 until ready for tissue assembly (miBrain or JAMs).

Differentiation of human iPSCs into microglia

iPSC-derived microglia were differentiated as previously shown[39,82] via an intermediate differentiation step into hematopoietic progenitor cells (HPCs). For the generation of HPCs, the STEMdiff Hematopoietic Kit (cat#: 05310; StemCell Technologies) was used, according to the manufacturer's manual. Briefly, when 70% confluent (day 0), iPSCs were harvested and passaged at a density of 20-40 colonies per well in a 6-well coated with 0.1mg/mL Matrigel (cat#: 354234; Corning). On day 1, Medium A was added to the culture, and on day 4 it was switched to Medium B until complete HPC maturation on days 11-13. Fully differentiated HPCs, floating in the medium and detached from the colonies, were collected for microglial differentiation or frozen in Stem-CellBanker (cat#: 11924; AMSBIO) supplemented with microglial cytokines. For the generation of mature microglia, differentiated HPCs were collected and transferred into Matrigel coated 6-well plate at a confluency of 350k HPCs per well. The differentiation takes 25-28 days, during which HPCs are cultured in microglia media consisting of DMEM/F12 (cat#: 11320-033; Gibco), with 2X B27 (cat#: 17504044; Thermo Fisher Scientific), 0.5X N2 (cat#: 17502048; Thermo Fisher Scientific), 1X Glutamax (cat#: 35050061; Gibco), 1X non-essential amino acids (cat#: 11-140-050; Gibco, 400 mM Monothioglycerol (cat#: M6145; Millipore Sigma), and 5 mg/mL human insulin (cat#: I9278; Millipore Sigma), freshly supplemented with 100 ng/mL IL-34 (cat#: 200-34; PeproTech), 50 ng/mL TGFβ1 (cat#: 100-21; PeproTech), and 25 ng/mL M-CSF (cat#: 300-25; PeproTech). Microglia were added to miBrains within the pool of Geltrex encapsulated cells at the time of miBrain assembly. MiBrains were maintained in miBrain media supplemented with 100 ng/mL IL-34 and 25 ng/mL M-CSF for 1 week and then switched to miBrain media supplemented with 25 ng/mL M-CSF until downstream experiments.

3D Tissue Assembly for miBrains, co-cultures or Cryopreservation

Neurons, astrocytes, endothelial cells, mural cells, and OPCs were dissociated using Accutase or TryplE Select (astrocytes). Cells were resuspended in corresponding media, counted, and resuspended at 1×10^6 cells/ mL. For miBrains, a tube was prepared to contain 5×10^6 neurons, 5×10^6 endothelial cells, 1×10^6 astrocytes, 1×10^6 OPCs, and 1×10^6 mural cells per 1 mL. Microglia was added for a subset of miBrains at the ratio of 1.67×10^6 per 1 mL. Pooled cells were spun down at 200 x g for 5 min at RT. Media was aspirated carefully, leaving the cell pellet undisturbed. The cell pellet was placed on ice and resuspended in 1 mL Geltrex™ supplemented with 10 µM Y27632 and 5 µg/mL doxycycline, avoiding air bubbles and keeping it on ice to prevent premature Geltrex polymerization and inability to seed miBrains properly. To generate miBrain tissue that adopted a free-floating, organoid-like morphology over time, 25-50 µL of encapsulated cell suspension were seeded per inner glass-bottom well of a 48-well MatTek plate (MatTek). To generate miBrain tissue that remained attached to the plate (more suitable for automated imaging) while conserving 3D morphology, 10 µL of encapsulated cell suspension were seeded per well of a 96-well µClear plastic-bottom plate (Greiner). For JAMs assembly and cryopreservation, a tube containing 5×10^6 endothelial cells, 1×10^6 astrocytes, 1×10^6 OPCs, and 1×10^6 mural cells per 1 mL was prepared. Pooled cells were spun down at 200 x g for 5 min at RT and cryopreserved in miBrain freezing media (60% KSR, 30% hECSR medium, 10% DMSO, 10 µM Y27632, 50 ng/mL VEGF-A). Upon thaw, cell viability was assessed, and the

appropriate volume of neurons needed to conserve the original miBrain cell-to-cell ratio was added to the
pooled cell suspension, which was spun again, encapsulated, and seeded as described above. After
miBrains were seeded, the plates were transferred into a 37 ˚C 95%/5% Air/CO2 incubator for 30 minutes to
allow the Geltrex™ to polymerize. After polymerization of the gel, miBrain week-1 medium (Human
Endothelial Serum-free Medium, 1x Pen/Strep, 1X MEM-NEAA, 1X CD Lipids, 1x Astrocyte Growth
Supplement (ScienCell), 1x B27 Supplement, 10ug/mL Insulin, 1μM cAMP-dibutyl, 50μg/mL Ascorbic acid,
10ng/mL NT3, 10ng/mL IGF, 100ng/mL Biotin, 60 ng/mL T3, 50 ng/mL VEGF, 1μM SAG, 5 μg/mL
doxycycline) was added to each well, ensuring complete submersion of the culture in media (250-500uL per
well of a 48-well matTek plate, 100 to 200 uL per well of a 96-well plate). Half media change was performed
every 2-3 days. On day 8 after miBrain seeding, the media was changed to miBrain week-2 medium (Human
Endothelial Serum-free Medium, 1x Pen/Strep, 1X MEM-NEAA, 1X CD Lipids, 1x Astrocyte Growth
Supplement (ScienCell), 1x B27 Supplement, 10ug/mL Insulin, 1μM cAMP-dibutyl, 50μg/mL Ascorbic acid,
10ng/mL NT3, 10ng/mL IGF, 100ng/mL Biotin, 60 ng/mL T3, 5 μg/mL doxycycline). Cultures were used for
downstream assays after 2 weeks.

Exposure of tissue to exogenous α-Synuclein

miBrains were exposed to 4 mg/ml of Human Recombinant Alpha Synuclein Protein Aggregates (Pre Formed
Fibrils, PFFs, from StressMarq, #SPR-322, or Abcam, #ab218819). PFFs were sonicated in a water bath
(VWR Ultrasonic Cleaner) immediately before adding to the cultures, using 10 cycles of 30 sec on, and 30
sec off. Free-floating miBrains incubated with Alpha-Synuclein had a media change after 48h and were fixed
in 4% paraformaldehyde after 2 weeks.

Induction of α-Syn pathology via expression of SNCA-A53T

Neurons with inducible expression of *SNCA* with the A53>T mutation (A53T), which increases α-Syn's
propensity to aggregate[44,45], were generated from iPSC as described above. Cells were harvested on day 7
of the differentiation and cultured in 2D on Poly-L-Ornithine and Laminin pre-coated plates at 156,250
cells/cm^2, or encapsulated for 3D cultures in Geltrex at 50,000 cells for 10 uL of Geltrex, or added to JAMs
for miBrain assembly as described above. The cultures were maintained for 18 days with or without PFFs.
For the cultures that received PFFs, 4 mg/ml PFFs were added to the media on day 4 after assembly. The
media was half-changed every 2-3 days. As an additional control, we generated neurons where SNCA-A53T
was overexpressed without the non-amyloid component (NAC) domain, which is required for α-Syn
aggregation.[83]

Immunofluorescence

2D cultures were fixed in 4% paraformaldehyde for 15 min at room temperature and rinsed with PBS. 2D
cultures were blocked in 0.3% Triton-X100, 5% normal donkey serum in PBS for 30 min., and then with
primary antibodies diluted in blocking buffer overnight at 4°C. Cultures were rinsed 3 times with 0.3% Triton-
X100 in PBS for 15 min each, and incubated with secondary antibodies (and Hoechst 33342) diluted at
1:1000 in blocking buffer for 2h at room temperature. Cultures were rinsed 3 times with PBS for 15 minutes
and left in PBS for image acquisition. 2D neuronal cultures were blocked in 0.1% Triton-X100, and 10%
normal donkey serum in PBS, antibodies were diluted in 0.02% Triton-X100 and 2% normal donkey serum
in PBS, and washes were done with PBS. All other conditions were kept the same as the other cell types.

3D cultures and miBrains were fixed in 4% paraformaldehyde overnight at 4°C and rinsed with PBS. 3D
cultures and miBrains were incubated in blocking solution (0.3% Triton-X100, 5% normal donkey serum, in
PBS) overnight at 4°C and then with primary antibodies diluted at 1:500 in blocking solution for 2-3 nights at

4°C. Cultures were rinsed 5 times with 0.3% Triton-X100 PBS for 30 min each and incubated with secondary
antibodies and Hoechst 33342 (Thermo 62249) diluted at 1:1000 in blocking solution for 2-3 nights at 4°C.
Cultures were rinsed 5 times with 0. 3% Triton-X100 PBS for 30 min each then rinsed and left in PBS for
image acquisition.

Image acquisition and quantification

Images were acquired using a confocal microscope (Leica Stellaris or Nikon AX R). For quantification, 10-
20 μm Z-stack images were acquired at 10x or 20x, 4 fields per well or free-floating miBrains. Volume
measurements on Z-stacks were performed using Nikon AX R built-in quantification software. Statistical
analyses were performed using GraphPad Prism Software. Normality and Lognormality tests D'Agostino &
Pearson, Anderson-Darling, Shapiro-Wilk and Kolmogorov-Smirnov were performed to determine the choice
for parametric or non-parametric testing. Post hoc tests on ANOVAs were conducted based on GraphPad
reccomendations.

Human iPSC-derived astrocyte bulk RNAseq analysis

Fragments Per Kilobase of transcript per Million mapped reads (FPKM) values were obtained from published
bulk RNA-seq data of isogenic human iPSC-derived astrocytes expressing either *APOE3/3* or *APOE4/4*[57].
FPKM values were log2-transformed with an offset of 0.1 and genes with zero variance across all the samples
were excluded. To focus on genes with meaningful variability, an additional filtering step was applied to retain
genes above the 10th percentile (variance > 0.033) after the removal of the zero-variance genes. Differential
expression analysis was performed using the limma package in R. A design matrix was constructed to model
the experimental conditions (APOE3 vs. APOE4) and sample-specific array weights were estimated using
the array Weights function with a prior.n value of 100 to help stabilize weight estimation. A linear model was
then fit to the log-transformed, filtered expression data using lmFit, and empirical Bayes moderation was
applied via the eBayes function, with trend and robust both set to TRUE. Differentially expressed genes
(DEGs) were defined as having an absolute log2 fold change > 0.5 and adjusted p-value (false discovery
weight) < 0.05. A volcano plot was then generated using the ggplot2 package and genes meeting the DEG
thresholds were highlighted. Enrichment analysis was performed on the upregulated and downregulated
DEGs separately using the clusterProfiler package. Pathways were identified through Gene Ontology (GO)
database. The top 10 pathways for each direction (upregulated and downregulated) were selected based on
adjusted p-values.

Human astrocyte scRNAseq pseudobulk analysis

The processed dataset was downloaded from Haney et al[64]. Unless otherwise stated, all following analyses
were performed in R using the package Seurat[84]. Aligning with the quality control protocol from the source
publication, cells with nFeature < 500, nCount < 1000, and mitochondrial and ribosomal reads > 10% were
discarded. Doublets were removed using the DoubletFinder package[85]. Following the standard Seurat
pipeline with default parameters, raw gene counts were normalized, the top 2,500 highly variable genes were
identified, and the data was scaled. Harmony was used to integrate the patient samples, and the top 20
principal components were used in Seurat's FindNeighbors, FindClusters (0.2 resolution), and RunUMAP
functions. Cell-type annotation was manually performed using the marker genes described in Haney et al.
and clusters not enriched for these marker genes were removed from further analysis. Pseudobulked
samples were generated by summing the raw counts for each patient sample and normalized using Seurat's
AggregateExpression() function.

Isogenic iPSC-derived astrocyte RNAseq analysis

623 The raw counts from iPSC-derived astrocyte bulk RNAseq were downloaded from Lin et al.[57] Raw counts
624 were processed using the DeSeq2 package[86] and normalized using the counts() function with normalized set
625 to TRUE.

DQ-BSA proteolytic activity assay

Astrocytes were seeded at 10,000 cells/well of a 96 well µClear plastic bottom plate (Greiner 655090) in AM.
The following day, media was changed to FBS-free maturation media (50% DMEM/F12, 50% Neurobasal,
1X B27 without Vitamin A, 1X N2, 1X NEAA, 1X GlutaMAX, 1% penicillin-streptomycin). Half of the wells
were treated with 100 nM BafilomycinA1 to inhibit lysosomal function as a negative control. The next day,
cells were pulsed with 1mM DQ-BSA (Thermo D12051) in maturation media for 30 minutes. Media was
replaced with fresh media (and fresh bafilomycin in the appropriate wells), and the red fluorescence and
bright field were imaged at 20x with an Incucyte (Sartorius) every hour for 24 hours.

Alternatively, astrocytes were seeded at 50K cells/well of a 12-well plate. Media changes were followed as
above. Cells were lifted to single-cell suspension with TrypLE Select 24 hours after the DQ-BSA pulse and
filtered into flow cytometry tubes with DAPI. Red fluorescence in live cells was analyzed with a BD Celesta
flow cytometer.

LysoSensor by flow cytometry

Astrocytes were seeded at 50,000 cells/well of a 12-well plate in AM. The following day, media was changed
to FBS-free maturation media (50% DMEM/F12, 50% Neurobasal, 1X B27 without Vitamin A, 1X N2, 1X
NEAA, 1X GlutaMAX, 1% penicillin-streptomycin), and cultured for 3 more days. Cells were incubated in 100
nM LysoSensor Green DND-189 (Thermo L7535) for 1 minute. Cells were washed and lifted to single cells
suspension with media changes followed as above. Cells were lifted to single-cell suspension with TrypLE
Select and filtered into flow cytometry tubes with DAPI. Green fluorescence in live cells was analyzed with a
BD Celesta flow cytometer.

Live Imaging of α-Syn-HiLyte Uptake and Degradation

Astrocytes were seeded 10,000 cells/well of a 96 well µClear plastic bottom plate (Greiner) in AM. The
following day, the media was changed to FBS-free maturation media (50% DMEM/F12, 50% Neurobasal, 1X
B27 without Vitamin A, 1X N2, 1X NEAA, 1X GlutaMAX, 1% penicillin-streptomycin), and cultured for 2 more
days. α-Syn HiLyte (AnaSpec AS-55457) was added to the media at a final concentration of 2 µg/mL and
nuclei were stained with 10 µg/mL Hoechst 33342. Images were acquired every 30-90 minutes for the first 6
hours on a Nikon AX R. At 24 h, the media was changed to fresh FBS-free maturation media to remove any
excess α-Syn. Images were acquired at 24 h and 48 h using the same parameters and laser settings.

Live Imaging of BODIPY-Cholesterol

Astrocytes were seeded 10,000 cells/well of a 96 well µClear plastic bottom plate (Greiner) in AM. The
following day, the media was changed to FBS-free maturation media (50% DMEM/F12, 50% Neurobasal, 1X
B27 without Vitamin A, 1X N2, 1X NEAA, 1X GlutaMAX, 1% penicillin-streptomycin), and cultured for 3 more
days. Astrocytes were incubated with 2 µM BODIPY-Cholesterol (Cayman 24618) for 2 hours. Cells were
then incubated with 10 µg/mL Hoechst 33342 for 10 minutes to stain nuclei and then washed. Media was
replaced with fresh media and cells were imaged using a CX7 High Content Screening Platform with a 20x
objective lens (Thermo; LUCPLFLN20x).

Western blotting and dot blotting

For Western blots, the protein concentration of each sample was measured using Pierce BCA Protein Assay (Thermo Fisher). Volumes corresponding to 20 μg of protein for each sample were loaded into Criterion Precast gels (BioRad), and a current of 120V was applied for approximately 60 min. The gel proteins were transferred to a PVDF membrane using BioRad TransBlot Turbo Transfer System, fixed in 4% PFA for 40 min and blocked in 5% w/v non-fat milk in 0.1% Tween in TBS (TBST) for 1 hour. For antibodies against phosphorylated proteins, blots were blocked in 5% w/v bovine serum albumin (BSA) in 0.1% TBST. Blots were then incubated with primary antibodies diluted at 1:1000 in blocking buffer overnight at 4°C. Membranes were washed in 0.1% TBST 3 times for 5 min each, incubated with secondary antibodies conjugates with horseradish peroxidase for 2h at room temperature in blocking buffer, and exposed to chemiluminescence activator before imaging using LI-COR Odyssey XF system. For dot blots, 3 mg of cell protein lysate or 3 ml of media were added onto nitrocellulose membranes and allowed to dry. Blocking and antibody incubation was performed as described for Western blots.

Collection and treatment with conditioned media

To generate neuron-conditioned media neurons were plated at ~150,000/cm2 on poly-L-ornithine and laminin coated plates on day 7 of differentiation. Media was changed following the differentiation protocol described above. Starting day 14 of differentiation, the media was collected, centrifuged at 2000xg for 5 minutes to pellet debris, and stored at -20°C. Media was collected with each half media change, every 3-4 days.

To generate astrocyte conditioned media, astrocytes were plated at ~30,000/cm2 on 0.1% gelatin-coated plates in AM. The following day, media was changed to either fresh neuron media (Neurobasal, 1x B-27, 1x N-2, 1x MEM-NEAA, 1x GlutaMAX, 1% penicillin-streptomycin, 10ng/mL BDNF, 10ng/mL GDNF, 0.5 mM dcAMP, 1ug/mL Laminin) supplemented with 10ng/mL CNTF or neuron conditioned media supplemented with 10ng/mL CNTF. Media was collected, centrifuged at 2000xg for 5 minutes to pellet debris, and stored at -20°C after 3 days.

To treat neurons with conditioned media, SNCA-A53T overexpressing neurons were plated at ~150,000/cm2 on poly-L-ornithine and laminin-coated 96-well μClear plates (Greiner) on day 7 of differentiation, following the normal neuron differentiation protocol described above. On day 11 of differentiation, the media was changed to conditioned media supplemented with 5 μg/mL doxycycline and 1 μg/mL laminin. Half media changes with conditioned media supplemented with doxycycline and laminin were continued every 3-4 days until day 25 of differentiation when cultures were fixed.

Treatment with cholesterol-lowering drugs

Astrocytes were seeded into 0.1% gelatin-coated 6 well plates in AM (ScienCell). After 2-3 days, the media was changed to AM without FBS supplemented with cholesterol-lowering drugs or DMSO as a vehicle. After 4 days, cells were seeded for final assays in 0.1% gelatin-coated 96-well μClear plates (Greiner) in AM with treatment. The following day, media was changed to FBS-free maturation media (50% DMEM/F12, 50% Neurobasal, 1X B27 without Vitamin A, 1X N2, 1X NEAA, 1X GlutaMAX, 1% penicillin-streptomycin) with treatment. Assays were started 2-3 days later, as described above. miBrains were treated with methyl-β-cyclodextrin (concentration below) or DMSO from day 0 of assembly to day 18, with half-media changes performed 3 times a week.

The following drug concentrations were used:
        2-hydroxypropyl-β-cyclodextrin: 100 μM (Sigma C0926)

703   methyl-β-cyclodextrin: 100 µM (Cayman 21633)
704   atorvastatin: 50 nM (Sigma SML3030)
705   efavirenz: 10 µM (MedChemExpress HY10572)
706   LXR-623: 10 µM (MedChemExpress HY-10629)
707   T0901317: 10 µM (MedChemExpress HY-10626)

Supplemental information

709 Document S1. Figures S1-S4; Legends for Videos S1, S2, and S3
710 Video S1. Neurons with α-Synuclein pathological phenotypes within the miBrain; related to Figure 1
711 Video S2. *APOE3/3* astrocytes co-localize with neuronal-derived α-Synuclein; related to Figure 2
712 Video S3. *APOE4/4* reactive astrocytes poorly co-localize with neuronal-derived α-Synuclein; related to
713 Figure 2

Figure Legends

Figure 1. *Induction of α-Syn intracellular inclusions in a multi-cellular integrated Brain (miBrain)*
tissue. **a**. Cartoon and representative images of our **m**ulticellular **i**ntegrated human **brain** (miBrain) tissue
generated from human iPSCs differentiated into six brain cell types, including neurons, glia, and vascular
cells. Four-week-old miBrains showing neurons (TUJ1, cyan) and vascular networks (PECAM-1, red). Scale
bar: 500 µm. Nuclei: blue. **b**. miBrains stained for specific markers of neurons (cyan; Neurofilament-H, TUJ1,
NGN2-sfGFP), astrocytes (green; S100b, AQP4), endothelial cells (red; PECAM-1), mural cells (green;
NG2), myelin (green; MBP), and microglia (green; IBA1). Scale bar: 50 µm, nuclei: blue. **c.** miBrains
containing neurons generated through direct iPSC reprograming via *NGN2* expression, with or without the
overexpression of *SNCA-A53T*. The tissue was cultured for a total of 18 days, with or without α-synuclein
PFFs added on day 4. Immunofluorescence for neuronal marker TUJ1 and for α-Syn phosphorylated at
Serine 129 (pS129) shows a robust expression of pS129-Syn in A53T neurons within the miBrains, but not
in WT tissue. Scale bar: 50 µm, nuclei: blue. Neuronal pS129 expression was significantly increased in
miBrains with A53T neurons. Exposure to PFFs exacerbated this effect. Bars represent means of pS129-
Syn[+] volume within TUJ1[+] neurons normalized to control WT. Error bars represent standard error (n = 4
biological replicates). P-values were calculated using a 2-way ANOVA followed by a Fisher's LSD test. **d**.
SNCA-A53T was fused to small folding green fluorescent protein (sfGFP). pS129-Syn[+] inclusions co-localize
with sfGFP expression in TUJ1[+] cells. Scale bar: 50 µm, nuclei: blue. **e**. Representative images depicting the
co-localization of SNCA-A53T-sfGFP (green) and neutral lipid marker Lipid Spot (magenta) in A53T miBrains,
or p-Syn (red) and Lipid Spot (cyan) in wild type and A53T miBrains cultured for two weeks. Arrowhead points
to a lipid droplet within a p-Syn[+] inclusion. The violin plot shows the percentage of sfGFP-SNCA volume
occupied by the overlapping signal of Lipid Spot. The sfGFP-SNCA volume overlapped at 5.2 % ± 0.7 with
lipid droplet marker Lipid Spot (mean ± standard error, n = 6 biological replicates). Dot plots represent the
median overlapping volume between p-Syn and Lipid Spot normalized by nuclei (n = 5-6 biological
replicates). P-values were calculated by Mann-Whitney test. Scale bar: 50 µm. **f.** Live imaging of SNCA-
A53T-sfGFP in cell bodies (i), neurites (ii), and varicose-like inclusions (iii). Scale bars: 25 µm (i, ii) and 15µm
(iii). Line graphs show the number of somatic and neuritic inclusions normalized by the sfGFP volume over
time (n = 4 biological replicates). Compared to WT, miBrains with A53T neurons had a significant increase
in the levels of lactate dehydrogenase (LDH) in the media, indicating cell death. Bars represent mean LDH
luminescence normalized to WT control, and error bars represent standard error (n = 4 biological replicates).
P-values were calculated using an unpaired t-test. Representative images of live tracking of SNCA-A53T-
sfGFP imaging and quantification at 2 and 22 weeks after miBrain assembly revealed a reduction in sfGFP
area between 2 and 22 weeks (p < 0.0001). Bars represent mean sfGFP volume. Error bars represent

standard error (n = 4 biological replicates). Scale bar: 100 μm. **g**. Representative images of miBrains with
A53T or WT neurons. The volume occupied by neurons (cyan; TUJ1) was significantly reduced in miBrains
with A53T neurons when compared with WT at 2 and 24 weeks. Bars represent TUJ1[+] volume and error
bars represent standard error (n = 3 biological replicates). P-values were calculated by 2-way ANOVA
followed by a Tukey test. Scale bar: 50 μm. **h.** Representative images of A53T miBrains stained for pS129
α-synuclein (red) and aggregated α-synuclein (cyan) 24 weeks after assembly. Bars represent the percent
volume of sfGFP-SNCA occupied by either pS129 α-Syn or aggregated α-synuclein staining. P-values were
calculated by Mann-Whitney test. Scale bar: 25 μm. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001

Figure 2. *APOE4* increases the phosphorylation and aggregation of α-Syn via astrocytes. a. Using
CRISPR-Cas9 on an *APOE3/3* iPSC line, we generated isogenic iPSCs harboring *APOE4/4* and
differentiated them into neurons with SNCA-A53T overexpression. We generated miBrains with *APOE3/3* or
APOE4/4 isogenic cells. Representative images of phosphorylated α-Syn immunoreactivity in *APOE3/3* and
APOE4/4 isogenic miBrains. Higher magnification panel depicts co-localized immunoreactivity between
pS129 (red), SNCA-A53T-sfGFP (green) and TUJ1 (magenta). Scale bars: 50 μm. *APOE4/4* miBrains had
significantly higher levels of pS129 (red) compared with *APOE3/3*. Bars represent mean values of percent
of sfGFP volume immunoreactive for phosphorylated α-Syn, normalized by *APOE3/3*. Error bars represent
standard error (n = 8 biological replicates). P-values were calculated using unpaired t-test. **b.** We generated
miBrains with *APOE3/3* or *APOE4/4* isogenic cells, including miBrains with one cell type at a time harboring
APOE4/4 (yellow boxes) and the remaining cells *APOE3/3* (blue boxes). Representative images of
phosphorylated α-Syn in a combinatorial screen of *APOE3/3* and *APOE4/4* isogenic cell types. *APOE4/4*
and *APOE3/3* miBrains with *APOE4/4* astrocytes had equivalent levels of pS129 α-Syn (red). These levels
were significantly higher than in miBrains where all cells harbored *APOE3/3* or where any cell type except
astrocytes harbored *APOE4/4*. Bars represent mean values of percent of sfGFP volume immunoreactive for
phosphorylated α-Syn, and error bars represent standard error (n = 8 biological replicates per combination).
P-values were calculated using one-way ANOVA followed by a Dunnett's Multiple Comparisons Test. Scale
bar: 50 μm. **c.** Representative images of GFAP immunoreactivity in *APOE3/3* and *APOE4/4* miBrains. Scale
bar: 50 μm. *APOE4/4* miBrain astrocytes (GFAP[+]) had reduced overlap with sfGFP-SNCA, increased
circularity and significantly higher levels of GFAP (magenta) compared with *APOE3/3*. Bars represent mean
values of GFAP volume and circularity and error bars represent standard error (n = 4 biological replicates).
P-values were calculated using unpaired t-test. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001

Figure 3. Impaired lysosomal function of *APOE4/4* astrocytes seeds α-Syn phosphorylation in
neurons. a. Pseudobulk analysis of human astrocytes (Haney et al. 2024) for *SNCA* expression. Bars
represent mean normalized psuedobulk gene count and error bars represent standard error (n = 8 APOE3/3
and n = 10 APOE4/4). P-value was calculated using unpaired t-test. **b** Western blots of *APOE3/3* and
APOE4/4 astrocytes for total α-Syn protein and phosphorylated α-Syn protein. Total and phosphorylated
αSyn is increased in *APOE4/4* astrocytes. The expected band size of α-Syn monomers is approximately
18kDa. β-Actin was used as a loading control. (n = 3 replicates). **c.** Representative images of uptake and
degradation of exogenous, fluorescently labeled α-Syn (green). α-Syn was removed from the culture media
after 24 hours. *APOE3/3* astrocytes uptake and degrade αSyn more readily than APOE4/4 astrocytes. Data
points represent mean values of α-Syn-HiLyte mean intensity normalized to nuclei area and *APOE3/3* at the
first time point. Error bars represent standard error (n = 4 replicates). P-values were calculated using 2-way
ANOVA followed by a Tukey test. **d.** Left: DQ-BSA Red integrated intensity per cell confluency measured
over 24 hours on an Incucyte (Sartorius). BafilomycinA1 at 100nM was used as a control for non-lysosomal
proteolysis of DQ-BSA. Data points represent mean values and error bars represent standard error (n = 4
replicates). Right: DQ-BSA Green mean fluorescence intensity measured in DAPI negative cell population
by flow cytometry after 24 hours in two different isogenic iPSC lines. Bars represent mean values normalized

to APOE3/3 and error bars represent standard error (n = 3 replicates). P-values were calculated by unpaired
t-tests. **e.** Representative images of pS129 α-Syn (red) in *APOE3/3* and *APOE4/4* astrocytes after exposure
to fresh or conditioned neuron media. *APOE4/4* astrocytes have more pS129 a-Syn, which is further
increased upon incubation with neuronal media. Data points represent mean values of pS129 a-Syn
normalized to cell area (CD44; cyan), and error bars represent standard error (n = 6 replicates). P-values
were calculated using 2-way ANOVA followed by a Fisher's LSD test. **f.** Left: schematic of the experimental
paradigm generating "double conditioned media" from *APOE/3* or *APOE4/4* astrocytes previously exposed
to neuronal media. Center, right: Representative images of *SNCA*-A53T neurons treated with neuron
conditioned media or neuron and astrocyte conditioned media. Neurons treated with *APOE4/4* double
conditioned media showed significant increase in pS129 α-Syn compared to all other conditions. Bars
represent mean pS129 volume normalized by sfGFP volume. Error bars represent standard error (n = 5
replicates). P-values were calculated using 1-way ANOVA followed by a Tukey test. All scale bars = 50 µm.
*P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Figure 4. Reducing cholesterol improves *APOE4/4* astrocyte lysosomal and α-Syn homeostasis. a.
Representative images of live astrocytes with BODIPY-cholesterol in two isogenic lines. *APOE4/4* astrocytes
have significantly more BODIPY-cholesterol staining than *APOE3/3* astrocytes. Bars represent BODIPY-
cholesterol mean intensity (green) normalized by Hoechst area (blue). Error bars represent standard error (n
= 3 replicates). P-values were calculated by unpaired t-tests. **b.** Representative images of BODIPY staining
in astrocytes treated with 2HβCD or MβCD. *APOE4/4* astrocytes have more BODIPY staining than *APOE3/3*
astrocytes, which is reduced after treatment. Bars represent BODIPY punctae (green) per cell number
(Hoechst; blue). Error bars represent standard error (n = 6 replicates). P-values were calculated by 2-way
ANOVA followed by a Tukey test. **c.** Top: DQ-BSA Red integrated intensity, in astrocytes with 2HβCD,
MβCD, atorvastatin, efavirenz, LXR-623, or T0901317 treatment, measured over 24 hours on an Incucyte
(Sartorius). Data points represent mean values and error bars represent standard error (n = 4 replicates).
Bottom: area under the curve calculation for DQ-BSA. Treatment of *APOE4/4* astrocytes with cyclodextrins,
but not other drugs, improved lysosomal proteolytic activity. Bars represent mean value and error bars
represent standard error (n = 4 replicates). P-values were calculated by 2-way ANOVA followed by a Tukey
test. **d.** Representative images of LysoTracker in astrocytes treated with 2HβCD or MβCD. Treatment with
cyclodextrins increased endolysosomal intensity. Bars represent LysoTracker mean intensity (magenta)
normalized by Hoechst area (blue). Error bars represent standard error (n = 5-6 replicates) P-values were
calculated by 2-way ANOVA followed by a Sidak test. **e.** Representative images of astrocytes treated with
cyclodextrins after a 24h incubation with fluorescently labeled α-Syn, in two isogenic lines. Cyclodextrin
treated astrocytes have increased α-Syn uptake compared to untreated *APOE4/4* astrocytes. Bars represent
a-Syn mean intensity (green) normalized by Hoechst area (blue). Error bars represent standard error (n = 6
replicates) P-values were calculated by 2-way ANOVA followed by a Tukey test. **f.** Representative images
of phosphorylated α-Syn in isogenic *APOE3/3* and *APOE4/4* miBrains. *APOE4/4* miBrains had significantly
higher levels of pS129 (red) than APOE3/3. These levels were significantly reduced in *APOE4/4* miBrains
treated with MβCD. Bars represent mean values of percent of sfGFP volume immunoreactive for
phosphorylated α-Syn and error bars represent standard error (n = 8 replicates). P-values were calculated
using two-way ANOVA followed by a Tukey test. All scale bars = 50µm. *P < 0.05, **P < 0.01, ***P < 0.001,
****P < 0.0001.

Extended Data Figure 1. **miBrain cryopreservation and development of α-synuclein intracellular**
inclusions. **a.** Cartoon of the cryopreservation approach to preserve miBrains or smaller tissue units
(miVasC: microvascular combo; BBB: blood-brain barrier; JAM: just add missing cell type). **b.** Top: cell

viability upon thawing the cryopreserved tissue compared with fresh cells harvested and counted. Bars
represent cell viability (%), and error bars represent standard deviation (n = 3 biological replicates). Bottom:
quantification of the ratio between neurons and nuclei volume in 18 days-old thawed miBrain tissue from
three different batches. Bars represent ratio between TUJ1 and Hoechst volumes, and error bars represent
standard error (n = 3 batches with 4 to 8 biological replicates each). Representative images of two-week-old,
thawed tissue stained with markers of neurons (cyan; neurofilament), astrocytes (green; S100b, GFAP), and
endothelial cells (red; PECAM-1). Scale bars: 50 μm. **c.** Representative images of miBrains exposed to α-
synuclein PFFs for 48h and stained at two weeks. Scale bars: 500 µm and 50 μm on lower and higher
magnification, respectively. The levels of α-Syn phosphorylated at Serine 129 (pS129, red) were significantly
increased in miBrains exposed to PFFs. Bars represent the intensity of pS129-Syn immunostaining
normalized by nuclei and by control. Left graph shows the quantification of the experiment represented by
the images. Error bars represent standard error (n = 4 biological replicates). P-values were calculated using
an unpaired t-test. Right graph shows the quantification for the PFF treated group in three different
experiments (n = 4 biological replicates each). Error bars represent standard error. Only experiments 1 and
2 had mean levels higher than control. Point dispersion in the bar graphs display high variability. **d**.
Representative images of pS129 α-Syn in neurons generated from SNCA-A53T iPSCs, wild type (WT)
iPSCs, or SNCA-A53T iPSCs without the non-amyloid component (NAC) domain. SNCA-A53T neurons had
increased pS129 only in the presence of PFFs. Bars represent neuron volume, and error bars represent
standard error (n = 4 biological replicates). P-values were calculated by 2-way ANOVA followed by a Tukey
test. Scale bars: 50 μm. **e.** In A53T miBrains, the overlap between aggregated α-synuclein and Tom20 was
significantly increased. Dot plot represents the median volume of aggregated α-Syn overlapping with Tom20
(n = 4 biological replicates). P-value was calculated using a Mann-Whitney test. Scale bar: 25 μm. *P < 0.05,
**P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Figure 2. Non-neuronal cells promote α-synuclein pathological phenotypes in APOE4
tissue. a. Representative images of 2D *APOE3/3* and *APOE4/4* isogenic neuronal cultures overexpressing
SNCA-A53T-sfGFP and exposed to PFFs for 2 weeks. Scale bar: 50 μm. pS129 was increased in both
APOE3/3 and *APOE4/4* SNCA-A53T neurons after exposure to PFFs when compared to WT controls. Bars
represent mean volume of phosphorylated α-Syn per β-III tubulin volume and normalized to control *APOE3/3*
neurons; error bars represent standard error (n = 4 biological replicates). **b.** Cartoon depicting the
experimental paradigm for generating isogenic *APOE3/3* and *APOE4/4* cells for miBrains. **c.** Cells were
harvested on the day of miBrain assembly, and a subset of the cells was plated in monocultures and fixed
after 24h. Representative images of *APOE3/3* and *APOE4/4* iPSC-derived cells stained for specific markers
(green) of each cell type. Nuclei: blue. Scale bar: 20 μm. Bars represent mean values of the area
immunoreactive for each cell marker normalized by nuclei and error bars represent standard error (n = 3
biological replicates). P-values were calculated using unpaired t-test. **d.** *APOE4/4* miBrains had significantly
higher levels of phosphorylated α-Syn outside the sfGFP volume compared with *APOE3/3*. Bars represent
mean values of percent of sfGFP volume outside the sfGFP volume, normalized by *APOE3/3*. Error bars
represent standard error (n = 8 biological replicates). P-values were calculated using unpaired t-test. **e.**
Representative images of *APOE3/3* and *APOE4/4* miBrains with WT or SNCA-A53T-sfGFP neurons. WT
miBrains were generated with isogenic *APOE3/3* or *APOE4/4* cells. A53T miBrains were generated with
A53T neurons that were differentiated from an iPSC line from an *APOE3/3* donor with familial Parkinson's
disease. All other cells were isogenic between each other except for the APOE locus. Scale bar: 50 μm. The
presence of the SNCA-A53T neurons significantly increased the levels of pS129-Syn, but this effect was
exacerbated when the non-neuronal cells harbored *APOE4/4*. Bars represent mean volume immunoreactive
for phosphorylated α-Syn per nuclei volume and normalized to *APOE3/3* miBrains with WT neurons. Error
bars represent standard error (n = 4 biological replicates). P- values were calculated using 2-way ANOVA

followed by a Fisher's LSD test. **f.** Representative images of *APOE3/3* and *APOE4/4* miBrains with isogenic
microglia and SNCA-A53T-sfGFP neurons. Scale bar: 50 μm. The presence of microglia in either APOE3/3
or APOE4/4 tissue did not significantly change the levels of pS129-Syn. Bars represent mean values of
percent of sfGFP volume immunoreactive for phosphorylated α-Syn, and error bars represent standard error
(n = 6 biological replicates per combination). P-values were calculated using 2-way ANOVA followed by a
Fisher's LSD test. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Figure 3. Characterization of lysosomal function and α-Synuclein uptake and
accumulation in astrocytes. a. RNAseq analysis of isogenic, iPSC derived astrocytes for SNCA expression.
Bars represent mean normalized counts and error bars represent standard error (n = 3 replicates). P-value
was calculated by unpaired t-test. **b.** Western blots of *APOE3/3* and *APOE4/4* astrocytes for total α-Syn
protein and phosphorylated α-Syn protein. Total and phosphorylated αSyn is increased in *APOE4/4*
astrocytes. The expected band size of α-Syn monomers is approximately 18kDa. β-Actin was used as a
loading control. (n = 3 replicates). **c.** Representative images of astrocytes with α-Syn-HiLyte (green) after 24
hours of uptake and 24 hours of degradation after treatment with BafilomycinA1 and MG-132. αSyn
degradation is only affected by lysosomal disruption in BafilomycinA1. Bars represent the mean α-Syn-HiLyte
intensity per nuclei area, normalized to control astrocytes. Error bars represent standard error (n = 5
replicates). P-values were calculated by one-way ANOVA followed by Tukey test. **d.** Representative images
of LAMP1 immunostaining (magenta) in two different isogenic iPSC lines. APOE4/4 astrocytes have
significantly less LAMP1⁺ lysosomes. Bars represent mean intensity per nuclei area normalized to *APOE3/3*.
Error bars represent standard error (n = 4 replicates). P-values were calculated using unpaired t-tests. **e.**
LysoSensor mean fluorescence intensity after 1 minute incubation measured by flow cytometry in DAPI
negative cells in three different isogenic iPSC lines. *APOE4/4* astrocytes have significantly reduced
LysoSensor signal compared to *APOE3/3* astrocytes. Bars represent mean values normalized to APOE3/3
and error bars represent standard error (n = 3 replicates). P-values were calculated using unpaired t-tests.
f. Top: Schematic of the experimental paradigm generating astrocyte conditioned media from *APOE3/3* and
APOE4/4 astrocytes. Bottom: Representative images of *SNCA*-A53T neurons treated with conditioned media
from naïve *APOE3/3* or *APOE4/4* astrocytes, or with fresh neuron media. There was no differential effect
between conditions. Bars represent mean pS129 volume normalized by sfGFP volume. Error bars represent
standard error (n = 6 replicates). P-values were calculated using 1-way ANOVA followed by a Tukey test. **g.**
Dot blots of *SNCA*-A53T conditioned media or double conditioned media from SNCA-A53T neurons and
isogenic *APOE3/3* or *APOE4/4* astrocytes for total αSyn, phosphorylated αSyn, and aggregated αSyn. All
dots were on one blot; the image was cropped to fit for the figure. Bars represent mean intensity and error
bars represent standard error (n = 4 replicates). P-values were calculated by unpaired, t-tests. All scale bars
= 50 μm. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Figure 4. MβCD treatment in *APOE4/4* astrocytes improves lysosomal uptake of α-
Synuclein. a. Volcano plot of *APOE4/4* vs. *APOE3/3* astrocytes showing DEGs (p val adj < 0.5, abs(logFC
> 0.5). Top 10 Gene Ontology (GO) upregulated (red) and downregulated (blue) pathways ranked by
adjusted p-value. Genes highlighted in the volcano plot correspond to GO terms related to lipid metabolism,
including "fatty acid oxidation", "cellular response to fatty acid", "regulation of lipid storage", "monocarboxylic
acid transport", "fatty acid transport", "cholesterol transfer activity", and "cholesterol metabolic process". **b.**
Left: DQ-BSA Red integrated intensity in astrocytes with 2HβCD or MβCD treatment, measured over 24
hours on an Incucyte (Sartorius) in a second isogenic donor line. Data points represent mean values and
error bars represent standard error (n = 4 replicates). Right: Area under the curve calculation for DQ-BSA.
Treatment of *APOE4/4* astrocytes with MβCD improved lysosomal proteolytic activity in a second donor line.
Bars represent mean value and error bars represent standard error (n = 4 replicates). P-values were
calculated by 2-way ANOVA followed by a Tukey test. **c.** Representative images of LysoTracker in astrocytes

treated with cyclodextrins in a second isogenic donor line. Treatment with cyclodextrins increased endolysosomal area. Bars represent LysoTracker mean intensity (magenta) normalized by Hoechst area (blue). Error bars represent standard error (n = 5-6 replicates) P-values were calculated by 2-way ANOVA followed by a Sidak test. **d.** Representative images of astrocytes treated with cyclodextrins after a 24h incubation with fluorescently labeled α-Syn in a second isogenic donor line. Cyclodextrin treated astrocytes have increased α-Syn uptake compared to untreated *APOE4/4* astrocytes. Bars represent α-Syn mean intensity (green) normalized by Hoechst area (blue). Error bars represent standard error (n = 6 replicates) P-values were calculated by 2-way ANOVA followed by a Tukey test. **e.** Representative images of fluorescently labeled α-Syn co-localized with endolysosomes in astrocytes treated with cyclodextrins. Cyclodextrin treated astrocytes had increased co-localization between α-Syn and lysosomes compared to untreated *APOE4/4,* indicative of improved α-Syn uptake. Bars represent mean Mander's coefficient M2 (amount of a-Syn signal in overlapping LysoTracker). Error bars represent standard error (n = 8 replicates). P-values were calculated by 2-way ANOVA followed by a Fisher's LSD test. All scale bars = 50μm. *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Extended Data Video 1. Neurons with α-synuclein pathological phenotypes within the miBrain. 3D rendering with animated orthogonal projection of SNCA-A53T neurons within the miBrain. Green: SNCA-A53T-sfGFP; red: pS129 α-Syn; magenta: TUJ1, blue: Hoechst.

Extended Data Video 2. APOE3/3 astrocytes co-localize with neuronal-derived α-synuclein. 3D rendering with animated orthogonal projection of astrocytes within an *APOE3/3* miBrain. Astrocytes display elongated morphology along neuronal networks. Green: SNCA-A53T-sfGFP; magenta: GFAP, blue: Hoechst.

Extended Data Video 3. *APOE4/4* **reactive astrocytes poorly co-localize with neuronal-derived α-synuclein.** 3D rendering with animated orthogonal projection of astrocytes within an *APOE4/4* miBrain. Astrocytes display amoeboid morphology not aligned with neuronal networks. Green: SNCA-A53T-sfGFP; magenta: GFAP, blue: Hoechst.

References

1. Hardy JA, Higgins GA. Alzheimer's disease: the amyloid cascade hypothesis. *Science*. Apr 10
1992;256(5054):184-5. doi:10.1126/science.1566067

2. Kotzbauer PT, Trojanowsk JQ, Lee VM. Lewy body pathology in Alzheimer's disease. *J Mol Neurosci*.
Oct 2001;17(2):225-32. doi:10.1385/jmn:17:2:225

3. Uchikado H, Lin WL, DeLucia MW, Dickson DW. Alzheimer disease with amygdala Lewy bodies: a
distinct form of alpha-synucleinopathy. *J Neuropathol Exp Neurol*. Jul 2006;65(7):685-97.
doi:10.1097/01.jnen.0000225908.90052.07

4. Popescu A, Lippa CF, Lee VM, Trojanowski JQ. Lewy bodies in the amygdala: increase of alpha-
synuclein aggregates in neurodegenerative diseases with tau-based inclusions. *Arch Neurol*. Dec
2004;61(12):1915-9. doi:10.1001/archneur.61.12.1915

5. Lippa CF, Fujiwara H, Mann DM, et al. Lewy bodies contain altered alpha-synuclein in brains of many
familial Alzheimer's disease patients with mutations in presenilin and amyloid precursor protein genes.
Am J Pathol. Nov 1998;153(5):1365-70. doi:10.1016/s0002-9440(10)65722-7

6. Hansen LA, Masliah E, Galasko D, Terry RD. Plaque-only Alzheimer disease is usually the lewy body
variant, and vice versa. *J Neuropathol Exp Neurol*. Nov 1993;52(6):648-54. doi:10.1097/00005072-
199311000-00012

7. Iwai A, Masliah E, Sundsmo MP, et al. The synaptic protein NACP is abnormally expressed during the
progression of Alzheimer's disease. *Brain Res*. May 13 1996;720(1-2):230-4. doi:10.1016/0006-
8993(96)00014-5

8. Hamilton RL. Lewy bodies in Alzheimer's disease: a neuropathological review of 145 cases using alpha-
synuclein immunohistochemistry. *Brain Pathol*. Jul 2000;10(3):378-84. doi:10.1111/j.1750-
3639.2000.tb00269.x

9. Arai Y, Yamazaki M, Mori O, Muramatsu H, Asano G, Katayama Y. Alpha-synuclein-positive structures
in cases with sporadic Alzheimer's disease: morphology and its relationship to tau aggregation. *Brain
Res*. Jan 12 2001;888(2):287-296. doi:10.1016/s0006-8993(00)03082-1

10. Marui W, Iseki E, Ueda K, Kosaka K. Occurrence of human alpha-synuclein immunoreactive neurons
with neurofibrillary tangle formation in the limbic areas of patients with Alzheimer's disease. *J Neurol
Sci*. Mar 15 2000;174(2):81-4. doi:10.1016/s0022-510x(99)00327-5

11. Olichney JM, Galasko D, Salmon DP, et al. Cognitive decline is faster in Lewy body variant than in
Alzheimer's disease. *Neurology*. Aug 1998;51(2):351-7. doi:10.1212/wnl.51.2.351

12. Chung EJ, Babulal GM, Monsell SE, Cairns NJ, Roe CM, Morris JC. Clinical Features of Alzheimer
Disease With and Without Lewy Bodies. *JAMA Neurol*. Jul 2015;72(7):789-96.
doi:10.1001/jamaneurol.2015.0606

13. Capouch SD, Farlow MR, Brosch JR. A Review of Dementia with Lewy Bodies' Impact, Diagnostic
Criteria and Treatment. *Neurol Ther*. Dec 2018;7(2):249-263. doi:10.1007/s40120-018-0104-1

14. Guerreiro R, Ross OA, Kun-Rodrigues C, et al. Investigating the genetic architecture of dementia with
Lewy bodies: a two-stage genome-wide association study. *Lancet Neurol*. Jan 2018;17(1):64-74.
doi:10.1016/S1474-4422(17)30400-3

15. Tsuang D, Leverenz JB, Lopez OL, et al. APOE epsilon4 increases risk for dementia in pure
synucleinopathies. *JAMA Neurol*. Feb 2013;70(2):223-8. doi:10.1001/jamaneurol.2013.600

16. Luk KC, Kehm V, Carroll J, et al. Pathological alpha-synuclein transmission initiates Parkinson-like
neurodegeneration in nontransgenic mice. *Science*. Nov 16 2012;338(6109):949-53.
doi:10.1126/science.1227157

17. Chung HK, Ho HA, Perez-Acuna D, Lee SJ. Modeling alpha-Synuclein Propagation with Preformed
Fibril Injections. *J Mov Disord*. Sep 2019;12(3):139-151. doi:10.14802/jmd.19046

18. Volpicelli-Daley LA, Luk KC, Lee VM. Addition of exogenous alpha-synuclein preformed fibrils to
primary neuronal cultures to seed recruitment of endogenous alpha-synuclein to Lewy body and Lewy
neurite-like aggregates. *Nat Protoc*. Sep 2014;9(9):2135-46. doi:10.1038/nprot.2014.143

19. Volpicelli-Daley LA, Luk KC, Patel TP, et al. Exogenous alpha-synuclein fibrils induce Lewy body
pathology leading to synaptic dysfunction and neuron death. *Neuron*. Oct 6 2011;72(1):57-71.
doi:10.1016/j.neuron.2011.08.033

20. Pediaditakis I, Kodella KR, Manatakis DV, et al. Modeling alpha-synuclein pathology in a human brain-
chip to assess blood-brain barrier disruption. *Nat Commun*. Oct 8 2021;12(1):5907.
doi:10.1038/s41467-021-26066-5

21. Jia L, Liu Y, Wang W, et al. Molecular Mediation of Prion-like alpha-Synuclein Fibrillation from Toxic
PFFs to Nontoxic Species. *ACS Appl Bio Mater*. Sep 21 2020;3(9):6096-6102.
doi:10.1021/acsabm.0c00684

22. Jan A, Goncalves NP, Vaegter CB, Jensen PH, Ferreira N. The Prion-Like Spreading of Alpha-
Synuclein in Parkinson's Disease: Update on Models and Hypotheses. *Int J Mol Sci*. Aug 3
2021;22(15)doi:10.3390/ijms22158338

23. Bousset L, Brundin P, Bockmann A, Meier B, Melki R. An Efficient Procedure for Removal and
Inactivation of Alpha-Synuclein Assemblies from Laboratory Materials. *J Parkinsons Dis*.
2016;6(1):143-51. doi:10.3233/JPD-150691

24. Duval K, Grover H, Han LH, et al. Modeling Physiological Events in 2D vs. 3D Cell Culture. *Physiology
(Bethesda)*. Jul 2017;32(4):266-277. doi:10.1152/physiol.00036.2016

25. Sierksma A, Escott-Price V, De Strooper B. Translating genetic risk of Alzheimer's disease into
mechanistic insight and drug targets. *Science*. Oct 2 2020;370(6512):61-66.
doi:10.1126/science.abb8575

26. Lancaster MA, Renner M, Martin CA, et al. Cerebral organoids model human brain development and
microcephaly. *Nature*. Sep 19 2013;501(7467):373-9. doi:10.1038/nature12517

27. Stanton AE, Bubnys A, Agbas E, et al. Engineered 3D Immuno-Glial-Neurovascular Human Brain
Model. *bioRxiv*. Aug 17 2023;doi:10.1101/2023.08.15.553453

28. Mazzulli JR, Zunke F, Isacson O, Studer L, Krainc D. alpha-Synuclein-induced lysosomal dysfunction
occurs through disruptions in protein trafficking in human midbrain synucleinopathy models. *Proc Natl
Acad Sci U S A*. Feb 16 2016;113(7):1931-6. doi:10.1073/pnas.1520335113

29. Lam I, Ndayisaba A, Lewis AJ, et al. Rapid iPSC inclusionopathy models shed light on formation,
consequence, and molecular subtype of alpha-synuclein inclusions. *Neuron*. Jul 23
2024;doi:10.1016/j.neuron.2024.06.002

30. Zhang Y, Pak C, Han Y, et al. Rapid single-step induction of functional neurons from human pluripotent
stem cells. *Neuron*. Jun 5 2013;78(5):785-98. doi:10.1016/j.neuron.2013.05.029

31. Chambers SM, Fasano CA, Papapetrou EP, Tomishima M, Sadelain M, Studer L. Highly efficient neural
conversion of human ES and iPS cells by dual inhibition of SMAD signaling. *Nat Biotechnol*. Mar
2009;27(3):275-80. doi:10.1038/nbt.1529

32. Tcw J, Wang M, Pimenova AA, et al. An Efficient Platform for Astrocyte Differentiation from Human
Induced Pluripotent Stem Cells. *Stem Cell Reports*. Aug 8 2017;9(2):600-614.
doi:10.1016/j.stemcr.2017.06.018

33.   Blanchard JW, Bula M, Davila-Velderrain J, et al. Reconstruction of the human blood-brain barrier in
1043   vitro reveals a pathogenic mechanism of APOE4 in pericytes. *Nat Med*. Jun 2020;26(6):952-963.
1044   doi:10.1038/s41591-020-0886-4

1045   34.   Mesentier-Louro LA, Suhy N, Broekaart D, Bula M, Pereira AC, Blanchard JW. Modeling the Blood-
1046   Brain Barrier Using Human-Induced Pluripotent Stem Cells. *Methods Mol Biol*. 2023;2683:135-151.
1047   doi:10.1007/978-1-0716-3287-1_11

1048   35.   Goldman C, Suhy N, Schwarz JE, et al. Reconstruction of the Blood-Brain Barrier In Vitro to Model and
1049   Therapeutically Target Neurological Disease. *J Vis Exp*. Oct 20 2023;(200)doi:10.3791/65921

1050   36.   Wang K, Lin RZ, Hong X, et al. Robust differentiation of human pluripotent stem cells into endothelial
1051   cells via temporal modulation of ETV2 with modified mRNA. *Sci Adv*. Jul 2020;6(30):eaba7606.
1052   doi:10.1126/sciadv.aba7606

1053   37.   Patsch C, Challet-Meylan L, Thoma EC, et al. Generation of vascular endothelial and smooth muscle
1054   cells from human pluripotent stem cells. *Nat Cell Biol*. Aug 2015;17(8):994-1003. doi:10.1038/ncb3205

1055   38.   Douvaras P, Fossati V. Generation and isolation of oligodendrocyte progenitor cells from human
1056   pluripotent stem cells. *Nat Protoc*. Aug 2015;10(8):1143-54. doi:10.1038/nprot.2015.075

1057   39.   McQuade A, Coburn M, Tu CH, Hasselmann J, Davtyan H, Blurton-Jones M. Development and
1058   validation of a simplified method to generate human microglia from pluripotent stem cells. *Mol*
1059   *Neurodegener*. Dec 22 2018;13(1):67. doi:10.1186/s13024-018-0297-x

1060   40.   Quadrato G, Nguyen T, Macosko EZ, et al. Cell diversity and network dynamics in photosensitive
1061   human brain organoids. *Nature*. May 4 2017;545(7652):48-53. doi:10.1038/nature22047

1062   41.   Anderson JP, Walker DE, Goldstein JM, et al. Phosphorylation of Ser-129 is the dominant pathological
1063   modification of alpha-synuclein in familial and sporadic Lewy body disease. *J Biol Chem*. Oct 6
1064   2006;281(40):29739-52. doi:10.1074/jbc.M600933200

1065   42.   Fujiwara H, Hasegawa M, Dohmae N, et al. alpha-Synuclein is phosphorylated in synucleinopathy
1066   lesions. *Nat Cell Biol*. Feb 2002;4(2):160-4. doi:10.1038/ncb748

1067   43.   Braak H, Del Tredici K, Rub U, de Vos RA, Jansen Steur EN, Braak E. Staging of brain pathology
1068   related to sporadic Parkinson's disease. *Neurobiol Aging*. Mar-Apr 2003;24(2):197-211.
1069   doi:10.1016/s0197-4580(02)00065-9

1070   44.   Narhi L, Wood SJ, Steavenson S, et al. Both familial Parkinson's disease mutations accelerate alpha-
1071   synuclein aggregation. *J Biol Chem*. Apr 2 1999;274(14):9843-6. doi:10.1074/jbc.274.14.9843

1072   45.   Giasson BI, Duda JE, Quinn SM, Zhang B, Trojanowski JQ, Lee VM. Neuronal alpha-synucleinopathy
1073   with severe movement disorder in mice expressing A53T human alpha-synuclein. *Neuron*. May 16
1074   2002;34(4):521-33. doi:10.1016/s0896-6273(02)00682-7

1075   46.   Altay MF, Kumar ST, Burtscher J, et al. Development and validation of an expanded antibody toolset
1076   that captures alpha-synuclein pathological diversity in Lewy body diseases. *NPJ Parkinsons Dis*. Dec
1077   7 2023;9(1):161. doi:10.1038/s41531-023-00604-y

1078   47.   Gai WP, Yuan HX, Li XQ, Power JT, Blumbergs PC, Jensen PH. In situ and in vitro study of
1079   colocalization and segregation of alpha-synuclein, ubiquitin, and lipids in Lewy bodies. *Exp Neurol*. Dec
1080   2000;166(2):324-33. doi:10.1006/exnr.2000.7527

1081   48.   Reeve AK, Park TK, Jaros E, et al. Relationship between mitochondria and alpha-synuclein: a study of
1082   single substantia nigra neurons. *Arch Neurol*. Mar 2012;69(3):385-93.
1083   doi:10.1001/archneurol.2011.2675

1084   49.   Wang X, Becker K, Levine N, et al. Pathogenic alpha-synuclein aggregates preferentially bind to
1085   mitochondria and affect cellular respiration. *Acta Neuropathol Commun*. Mar 14 2019;7(1):41.
1086   doi:10.1186/s40478-019-0696-4

1087   50.   Cookson MR. alpha-Synuclein and neuronal cell death. *Mol Neurodegener*. Feb 4 2009;4:9.
1088   doi:10.1186/1750-1326-4-9

1089   51.   Calabresi P, Mechelli A, Natale G, Volpicelli-Daley L, Di Lazzaro G, Ghiglieri V. Alpha-synuclein in
1090   Parkinson's disease and other synucleinopathies: from overt neurodegeneration back to early synaptic
1091   dysfunction. *Cell Death Dis*. Mar 1 2023;14(3):176. doi:10.1038/s41419-023-05672-9

1092   52.   Zhao N, Liu CC, Qiao W, Bu G. Apolipoprotein E, Receptors, and Modulation of Alzheimer's Disease.
1093   *Biol Psychiatry*. Feb 15 2018;83(4):347-357. doi:10.1016/j.biopsych.2017.03.003

1094   53.   Yamazaki Y, Zhao N, Caulfield TR, Liu CC, Bu G. Apolipoprotein E and Alzheimer disease:
1095   pathobiology and targeting strategies. *Nat Rev Neurol*. Sep 2019;15(9):501-518. doi:10.1038/s41582-
1096   019-0228-7

1097   54.   Davis AA, Inman CE, Wargel ZM, et al. APOE genotype regulates pathology and disease progression
1098   in synucleinopathy. *Sci Transl Med*. Feb 5 2020;12(529)doi:10.1126/scitranslmed.aay3069

1099   55.   Zhao N, Attrebi ON, Ren Y, et al. APOE4 exacerbates alpha-synuclein pathology and related toxicity
1100   independent of amyloid. *Sci Transl Med*. Feb 5 2020;12(529)doi:10.1126/scitranslmed.aay1809

1101   56.   Zhao J, Lu W, Ren Y, et al. Apolipoprotein E regulates lipid metabolism and alpha-synuclein pathology
1102   in human iPSC-derived cerebral organoids. *Acta Neuropathol*. Nov 2021;142(5):807-825.
1103   doi:10.1007/s00401-021-02361-9

1104   57.   Lin YT, Seo J, Gao F, et al. APOE4 Causes Widespread Molecular and Cellular Alterations Associated
1105   with Alzheimer's Disease Phenotypes in Human iPSC-Derived Brain Cell Types. *Neuron*. Jun 27
1106   2018;98(6):1141-1154 e7. doi:10.1016/j.neuron.2018.05.008

1107   58.   Tcw J, Qian L, Pipalia NH, et al. Cholesterol and matrisome pathways dysregulated in astrocytes and
1108   microglia. *Cell*. Jun 23 2022;185(13):2213-2233 e25. doi:10.1016/j.cell.2022.05.017

1109   59.   Choi I, Zhang Y, Seegobin SP, et al. Microglia clear neuron-released alpha-synuclein via selective
1110   autophagy and prevent neurodegeneration. *Nat Commun*. Mar 13 2020;11(1):1386.
1111   doi:10.1038/s41467-020-15119-w

1112   60.   Scheiblich H, Dansokho C, Mercan D, et al. Microglia jointly degrade fibrillar alpha-synuclein cargo by
1113   distribution through tunneling nanotubes. *Cell*. Sep 30 2021;184(20):5089-5106 e21.
1114   doi:10.1016/j.cell.2021.09.007

1115   61.   Lee HJ, Suk JE, Patrick C, et al. Direct transfer of alpha-synuclein from neuron to astroglia causes
1116   inflammatory responses in synucleinopathies. *J Biol Chem*. Mar 19 2010;285(12):9262-72.
1117   doi:10.1074/jbc.M109.081125

1118   62.   Loria F, Vargas JY, Bousset L, et al. alpha-Synuclein transfer between neurons and astrocytes
1119   indicates that astrocytes play a role in degradation rather than in spreading. *Acta Neuropathol*. Nov
1120   2017;134(5):789-808. doi:10.1007/s00401-017-1746-2

1121   63.   Tsunemi T, Perez-Rosello T, Ishiguro Y, et al. Increased Lysosomal Exocytosis Induced by Lysosomal
1122   Ca(2+) Channel Agonists Protects Human Dopaminergic Neurons from alpha-Synuclein Toxicity. *J
1123   Neurosci*. Jul 17 2019;39(29):5760-5772. doi:10.1523/JNEUROSCI.3085-18.2019

1124   64.   Haney MS, Palovics R, Munson CN, et al. APOE4/4 is linked to damaging lipid droplets in Alzheimer's
1125   disease microglia. *Nature*. Apr 2024;628(8006):154-161. doi:10.1038/s41586-024-07185-7

1126   65.   Stefanis L, Emmanouilidou E, Pantazopoulou M, Kirik D, Vekrellis K, Tofaris GK. How is alpha-
1127   synuclein cleared from the cell? *J Neurochem*. Sep 2019;150(5):577-590. doi:10.1111/jnc.14704

66. Konings SC, Torres-Garcia L, Martinsson I, Gouras GK. Astrocytic and Neuronal Apolipoprotein E
Isoforms Differentially Affect Neuronal Excitability. *Front Neurosci*. 2021;15:734001.
doi:10.3389/fnins.2021.734001

67. Pfrieger FW. Role of cholesterol in synapse formation and function. *Biochim Biophys Acta*. Mar 10
2003;1610(2):271-80. doi:10.1016/s0005-2736(03)00024-5

68. Lane-Donovan C, Philips GT, Herz J. More than cholesterol transporters: lipoprotein receptors in CNS
function and neurodegeneration. *Neuron*. Aug 20 2014;83(4):771-87.
doi:10.1016/j.neuron.2014.08.005

69. Jeong W, Lee H, Cho S, Seo J. ApoE4-Induced Cholesterol Dysregulation and Its Brain Cell Type-
Specific Implications in the Pathogenesis of Alzheimer's Disease. *Mol Cells*. Nov 30 2019;42(11):739-
746. doi:10.14348/molcells.2019.0200

70. Lee H, Cho S, Kim MJ, et al. ApoE4-dependent lysosomal cholesterol accumulation impairs
mitochondrial homeostasis and oxidative phosphorylation in human astrocytes. *Cell Rep*. Oct 31
2023;42(10):113183. doi:10.1016/j.celrep.2023.113183

71. Liddelow SA, Guttenplan KA, Clarke LE, et al. Neurotoxic reactive astrocytes are induced by activated
microglia. *Nature*. Jan 26 2017;541(7638):481-487. doi:10.1038/nature21029

72. Chen Y, Qin C, Huang J, et al. The role of astrocytes in oxidative stress of central nervous system: A
mixed blessing. *Cell Prolif*. Mar 2020;53(3):e12781. doi:10.1111/cpr.12781

73. de Leeuw SM, Kirschner AWT, Lindner K, et al. APOE2, E3, and E4 differentially modulate cellular
homeostasis, cholesterol metabolism, and inflammatory response in isogenic iPSC-derived astrocytes.
Stem Cell Reports. Jan 11 2022;17(1):110-126. doi:10.1016/j.stemcr.2021.11.007

74. Strauss K, Goebel C, Runz H, et al. Exosome secretion ameliorates lysosomal storage of cholesterol
in Niemann-Pick type C disease. *J Biol Chem*. Aug 20 2010;285(34):26279-88.
doi:10.1074/jbc.M110.134775

75. Stykel MG, Humphries KM, Kamski-Hennekam E, et al. alpha-Synuclein mutation impairs processing
of endomembrane compartments and promotes exocytosis and seeding of alpha-synuclein pathology.
Cell Rep. May 11 2021;35(6):109099. doi:10.1016/j.celrep.2021.109099

76. Sharma R, Hastings C, Staretz-Chacham O, et al. Long-term administration of intravenous Trappsol(R)
Cyclo (HP-beta-CD) results in clinical benefits and stabilization or slowing of disease progression in
patients with Niemann-Pick disease type C1: Results of an international 48-week Phase I/II trial. *Mol
Genet Metab Rep*. Sep 2023;36:100988. doi:10.1016/j.ymgmr.2023.100988

77. Min JO, Ho HA, Lee W, et al. Statins suppress cell-to-cell propagation of alpha-synuclein by lowering
cholesterol. *Cell Death Dis*. Jul 27 2023;14(7):474. doi:10.1038/s41419-023-05977-9

78. Oveisgharan S, Yu L, Barnes LL, et al. Association of Statins With Cerebral Atherosclerosis and
Incident Parkinsonism in Older Adults. *Neurology*. May 10 2022;98(19):e1976-e1984.
doi:10.1212/WNL.0000000000200182

79. Stevens KN, Creanor S, Jeffery A, et al. Evaluation of Simvastatin as a Disease-Modifying Treatment
for Patients With Parkinson Disease: A Randomized Clinical Trial. *JAMA Neurol*. Dec 1
2022;79(12):1232-1241. doi:10.1001/jamaneurol.2022.3718

80. Qian T, Maguire SE, Canfield SG, et al. Directed differentiation of human pluripotent stem cells to blood-
brain barrier endothelial cells. *Sci Adv*. Nov 2017;3(11):e1701679. doi:10.1126/sciadv.1701679

81. Douvaras P, Wang J, Zimmer M, et al. Efficient generation of myelinating oligodendrocytes from primary
progressive multiple sclerosis patients by induced pluripotent stem cells. *Stem Cell Reports*. Aug 12
2014;3(2):250-9. doi:10.1016/j.stemcr.2014.06.012

82.   McQuade A, Blurton-Jones M. Human Induced Pluripotent Stem Cell-Derived Microglia (hiPSC-
Microglia). In: Nagy A, Turksen K, eds. *Induced Pluripotent Stem (iPS) Cells: Methods and Protocols*.
Springer US; 2022:473-482.

83.   Giasson BI, Murray IV, Trojanowski JQ, Lee VM. A hydrophobic stretch of 12 amino acid residues in
the middle of alpha-synuclein is essential for filament assembly. *J Biol Chem*. Jan 26 2001;276(4):2380-
6. doi:10.1074/jbc.M008919200

84.   Hao Y, Hao S, Andersen-Nissen E, et al. Integrated analysis of multimodal single-cell data. *Cell*. Jun
24 2021;184(13):3573-3587 e29. doi:10.1016/j.cell.2021.04.048

85.   McGinnis CS, Murrow LM, Gartner ZJ. DoubletFinder: Doublet Detection in Single-Cell RNA
Sequencing Data Using Artificial Nearest Neighbors. *Cell Syst*. Apr 24 2019;8(4):329-337 e4.
doi:10.1016/j.cels.2019.03.003

86.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data
with DESeq2. *Genome Biol*. 2014;15(12):550. doi:10.1186/s13059-014-0550-8

1185

Figure 1

miBrain System

Human iPSC

Neurons   OPCs   Astrocytes   Endothelial Cells   Mural Cells   Microglia

miBrain

TUJ1 PECAM-1 Hoechst

b

Neurovascular unit

Neurofilament-H
PECAM-1

TUJ1 PECAM-1 S100β Hoechst

Blood-brain barrier

PECAM-1 AQP4

PECAM-1 NG2 Hoechst

Neuro-myelin

Neurofilament-H MBP Hoechst

Microglia

NGN2-sfGFP IBA1 Hoechst

Synuclein Pathological Phenotypes in the miBrain

c

miBrain → iPSC → TetON NGN2 → ... → iPSC → TetON A53T SCNA-sfGFP IRES NGN2 → ...

SNCA WT   SNCA A53T

Control

PFFs

pS129 Syn TUJ1 Hoechst

Neuronal p-Syn Normalized

100 80 60 40 20 0

*** **** *

WT   SNCA-A53T

Control   PFFs

d

pS129 Syn SNCA-A53T sfGFP TUJ1 Hoechst

e

LipidSpot SNCA-A53T sfGFP Hoechst

%sfGFP with LipidSpot

15 10 5 0

WT   A53T

pS129 Syn LipidSpot Hoechst

pSyn w/ Lipid Spot Normalized

20 15 10 5 0

*

WT   A53T

f Live imaging SNCA-A53T sfGFP

i   ii   iii

Somatic inclusions

2.0 1.5 1.0 0.5 0.0
0 4 8 12 Weeks

Neuritic inclusions

4 3 2 1 0 -1
0 4 8 12 Weeks

Relative LDH Luminescence

2.0 1.5 1.0 0.5 0.0
**
WT   A53T

2 weeks   22 weeks

sfGFP volume (μm³)

9×10⁶ 6×10⁶ 3×10⁶
****
2   22 Weeks

g

WT   SNCA-A53T

2 weeks

22 weeks

TUJ1

TUJ1 Volume Normalized

1.5 1.0 0.5 0.0
** *
2   24 Weeks

WT   SNCA-A53T

h

SNCA-A53T sfGFP pS129 Syn Aggregated Syn Hoechst

%sfGFP with p-Syn

60 40 20 0 -20
**
2   24 Weeks

%sfGFP with Agg. Syn

30 20 10 0 -10
*
2   24 Weeks

Figure 2

Figure 3

a *SNCA* Normalized Counts — *APOE3/3* vs *APOE4/4*, *

b pS129 α-Syn (~35kDa, ~18kDa), α-Syn (~18kDa), Actin (~42kDa); *APOE3/3* vs *APOE4/4*

c 6 hours, 24 hours, 48 hours; *APOE3/3*, *APOE4/4*
αSyn-HiLyte Hoechst
α-Synuclein HiLyte Intensity vs Elapsed Time (hrs)
Remove α-syn-488

d DQ-BSA Integrated Intensity vs Time (hrs); *APOE3/3*, *APOE4/4*, + BafA1
DQ-BSA Normalized MFI; *APOE3/3* vs *APOE4/4* Donor #1, Donor #3; **

e Control Media → *APOE3/3* Astrocytes, *APOE4/4* Astrocytes; SNCA-A53T Cond. Media → *APOE3/3* Astrocytes, *APOE4/4* Astrocytes
Control, Neuron Conditioned Media; *APOE3/3*, *APOE4/4*
pS129 αSyn CD44 Hoechst
phospho-α-Syn (S129)/Cell Area; *APOE3/3* Astrocytes, *APOE4/4* Astrocytes

f *SNCA*-A53T Neurons; *APOE3/3* Astrocytes; *APOE4/4* Astrocytes; Conditioned Media
Neuron Cond. Media: No AC; Double Cond. Media: *APOE3/3* AC, *APOE4/4* AC
pS129 αSyn sfGFP-SNCA Hoechst
%sfGFP with p-Syn normalized; No Astrocytes, *APOE3/3* Astrocytes, *APOE4/4* Astrocytes

Figure 4

Extended Data Fig. 1

Extended Data Fig. 2

Extended Data Fig. 3

Extended Data Fig. 4