Functional Specialization for Language Processing in Inferior Frontal Regions During Early Childhood: Evidence from functional near-infrared spectroscopy individual functional channels of interest approach

Haolun Luo[1], Tao Yu[2*], Qun Li[3*], Li Sheng[1*]

[1] Department of Chinese and Bilingual Studies, Faculty of Humanities, The Hong Kong Polytechnic University, Hong Kong SAR, China.

[2] Bio-X Institutes, Shanghai Jiao Tong University, Shanghai, China

[3] Department of Otorhinolaryngology-Head and Neck surgery, West China Hospital of Sichuan University, Chengdu, China.

Abstract

Significance: Early language acquisition represents a fundamental achievement in cognitive development, yet the neural mechanisms underlying this process remain debated, particularly whether specialized language circuits exist from early life or emerge gradually through development.

Aim: To investigate functional specialization for language processing in bilateral inferior frontal regions during early childhood, examining whether language-selective regions are distinct from domain-general cognitive networks in toddlers aged 2-4 years.

Approach: Using functional near-infrared spectroscopy (fNIRS) with an innovative functional Channel of Interest (fCOI) approach, we conducted two experiments involving adults (N=20) and toddlers (N=22, ages 2-4 years) who completed language processing and cognitive control tasks.

Results: We demonstrated early functional specialization in the language-selective region of left inferior frontal gyrus, which showed selective responses to linguistic content while remaining insensitive to cognitive demand manipulations in both age groups. However, language selectivity in the homologous right hemisphere region was present only in adults. The multiple demand regions showed complementary patterns, with right-hemispheric selectivity for cognitive control emerging early.

Conclusions: These findings provide evidence for early neural specialization of language processing in the left hemisphere, while revealing ongoing development in right hemispheric organization. Our results support models of early language-specific neural circuits rather than gradual differentiation from domain-general mechanisms, while highlighting the protracted development of language organization.

KEYWORDS : cognitive load, development, fNIRS, language, multiple demand

[*]Correspondence:
YU Tao, Bio-X Institutes, Shanghai Jiao Tong University, Shanghai, China
Email: yutao@sjtu.edu.cn

LI Qun, Department of Otorhinolaryngology-Head and Neck surgery, West China Hospital of Sichuan University, Chengdu, China
Email: liqun2021@wchscu.cn

SHENG Li, Department of Chinese and Bilingual Studies, Faculty of Humanities, The Hong Kong Polytechnic University, Hong Kong SAR, China
Email: dr-li.sheng@polyu.edu.hk

Introduction

Language acquisition in early childhood represents a remarkable achievement of human cognitive development, occurring naturally through exposure without formal training[1]. A fundamental question in developmental cognitive neuroscience is how the brain supports early language acquisition. One of the most debated aspects of language development is whether the brain contains specialized circuits for language processing from early in life, or whether these circuits emerge gradually over development. Specifically, there are two competing hypotheses about the neural mechanisms underlying language processing in young children. One possibility is that language processing relies on domain-specific neural mechanisms from early on, suggesting an early neural specialization for language that is distinct from other cognitive functions[2,3]. Alternatively, language processing might initially emerge through domain-general cognitive processes, such as working memory and executive function, with specialized language circuits developing gradually through experience[4,5].

Studies in adults have identified two distinct neural networks potentially relevant for understanding language development: the 'core language network' and the 'multiple demand' (MD) network. The language network, which includes regions primarily in left frontal and temporal areas, shows consistent engagement during language processing with limited activation during non-linguistic tasks[6–8]. In contrast, the MD network, comprising bilateral frontal, parietal, cingulate, and insular regions, responds to increased cognitive demands across various cognitive tasks[9,10]. The relationship between these networks and their respective roles in language processing have been debated. Some researchers argue that language processing relies substantially on domain-general cognitive mechanisms implemented in the MD network[11,12] , while others propose more distinct functional roles[13]. Recent adult neuroimaging evidence provides insights into how these networks contribute to language processing. Despite apparent similarities between mental operations involved in language comprehension (such as

memory retrieval, prediction, and structure building) and those required by other cognitive

domains, the MD network's role in basic language comprehension appears more subtle than

initially theorized[14–16]. By contrast, the MD network displays more robust engagement when

language tasks introduce extra cognitive demands, such as retaining information in working

memory or answering comprehension questions[17–19]. This can be understood as the language

network implementing core computations for linguistic structure and meaning, while MD areas

are recruited when task requirements demand increased cognitive control[15,20]. This dissociation

is supported by evidence from multiple methodologies, including functional connectivity

analyses[14,21], studies of patients with focal brain lesions[22], and investigations using naturalistic

language stimuli[23,24]. These findings suggest that while the language network may implement

certain computations on linguistic representations somewhat independently, it likely interacts

with domain-general circuits in the MD network when tasks require additional cognitive

control[20,25] .

While studies with adults have suggested relatively clear dissociation between

language-specific and domain-general networks, documenting this dissociation in adults alone

is insufficient for understanding the origin of uniquely human cognitive abilities. Several

competing developmental trajectories could lead to the same adult organization: cortical

regions might start as domain-general and gradually specialize through competitive

interactions[4], with language regions initially participating in broader cognitive functions like

working memory before gaining linguistic selectivity. Alternatively, domain-general processes

might bootstrap the development of language abilities[26–28]. A third pattern could be that there

is an innate proto-organization of domain-specificity. Understanding how and when these

networks dissociate during development is therefore critical for distinguishing between these

possibilities and illuminating the origins of human cognitive architecture. Recent fMRI studies

have demonstrated that this selective and specific language network is already established in

children as young as 4 years of age, showing adult-like functional dissociation from domain-

general cognitive processes[29,30]. However, due to the methodological constraints of fMRI,

particularly the requirement for participants to remain still, the emergence and organization of

these networks in even younger children remains largely unexplored. To the best of our

knowledge, there is no existing exploration of the double dissociation of language and domain-

general cortex in children younger than 4.

To address this gap, we employ functional near-infrared spectroscopy (fNIRS), which

allows measurements while participants move relatively freely. fNIRS uses absorption of near-

infrared light between emitters and detectors placed against the scalp to measure changes in

oxygenated and deoxygenated hemoglobin concentrations that reflect neural activity[31]. This

method has been successfully used to study early language processing in infants. Several

studies have demonstrated that newborns already show specialized neural responses to speech

in temporal and frontal regions[32–34], while later work has suggested that language-specific

responses continue to develop throughout infancy[35]. While these studies provide evidence for

differential neural responses to speech versus reversed or degraded speech stimuli, they have

not fully characterized the functional architecture of language processing in the developing

brain. Specifically, showing greater activation to speech than non-speech does not establish

whether these regions are specialized for language processing or serve more domain-general

functions that happen to be engaged during speech processing. As Kanwisher[36] emphasizes, in

studies of functional specificity, demonstrating true specialization requires not only evidence

of preferential responses to one category but should also test whether regions are exclusively

engaged in processing that category. For example, the right temporoparietal junction (rTPJ)

activates selectively when thinking about others' thoughts, but not during closely related tasks

like processing physical appearances or even other mental states such as hunger or pleasure.

In additional to these conceptual challenges, mapping fNIRS measurement channels to specific

cortical regions presents several methodological challenges, particularly in developmental

populations. These challenges arise from anatomical variability in the relationship between

external landmarks and underlying cortical structures, developmental variations in head size

affecting measurement array coverage, and practical constraints in precise optode placement[37].

Moreover, even when measurement channels can be accurately mapped to specific anatomical

regions, this anatomical localization does not guarantee functional correspondence. A given

anatomical region often contains multiple functionally distinct sub-regions that vary in size and

precise location across individuals[38,39].

To overcome these methodological challenges, we employ a functional Channel of

Interest (fCOI) approach for analyzing fNIRS data[37]. Unlike traditional approaches, which

assume that measurement channels correspond to the same cortical regions across participants,

the fCOI method identifies channels of interest by examining each individual's functional

activation patterns. This approach recognizes that the same channel position relative to scalp

landmarks may measure different cortical regions across individuals due to variations in array

placement, head size, and the relationship between surface landmarks and underlying cortical

anatomy - a particularly important consideration in developmental research. The fCOI

approach offers several key advantages for developmental studies[40]: it accounts for individual

variability in brain anatomy and functional organization[41], reduces noise from neighboring

regions with different functional profiles, and provides a principled way to limit multiple

comparisons while maintaining statistical power. This approach has been validated across

multiple domains of cognitive development, including face processing and theory of

mind[37,42,43].

In the present study, we investigate the functional organization of language and domain-

general networks in toddlers aged 2-4 years using fNIRS with the fCOI approach. Following

Kanwisher's[36] framework for establishing functional specificity, our experimental design

carefully contrasts responses across cognitive domains rather than simply identifying regions

that respond preferentially to language. We focus on bilateral inferior frontal gyrus (IFG)

regions given their well-documented heterogeneity in adult neuroimaging studies[13].

Specifically, prior research has established that Broca's area within the left IFG contains two

functionally distinct regions: a language-selective region responding specifically to linguistic

input, and a domain-general region responding to diverse cognitive demands[13]. While

anatomically adjacent, these regions belong to different networks - the left-lateralized

frontotemporal language network and the bilateral frontoparietal multiple-demand (MD)

network, respectively. Moreover, IFG's anatomical characteristics[44] (e.g. minimal hair

interference, relatively thin skull in this region, proximity to cortical surface) maximize fNIRS

signal quality. Therefore, IFG provides an optimal starting point for investigating whether

specialized circuits emerge in core regions before extending to broader networks. Our

experimental design includes a child-friendly language localizer adapted from Scott et al.[45] that

contrasts responses to auditorily presented engaging narrative clips versus acoustically

degraded versions of the same stimuli. This localizer has been demonstrated to reliably identify

language-selective regions across 45 languages from 12 language families, showing consistent

activation patterns in the fronto-temporal language network regardless of the specific language

being processed[46]. The passive listening nature of this task makes it particularly suitable for

young children who may have difficulty with reading or cognitively demanding tasks with the

confounding effects of task difficulty being controlled. The narrative stimuli can be easily

customized with age-appropriate content to maintain children's attention. For the cognitive

control task, we used a spatial working memory task to engage the MD network for adults, as

this paradigm has been consistently shown to be the most reliable and widely used task for

localizing the MD network across numerous studies[10,22]. Using the same paradigm allows us

to directly compare our fNIRS findings with the extensive fMRI literature. For children, we

employed a Go/No-go task that probes inhibitory control - another key component of executive

function that has been shown to engage the MD network[47]. While these tasks tap different

aspects of executive function (working memory versus inhibitory control), both have been

demonstrated to reliably activate the MD network and show similar patterns of increased

activation under higher cognitive demands[9,10]. We implemented two conditions in the Go/No-

go task: an easier condition with only "go" trials and a more demanding condition mixing "go"

and "no-go" trials, paralleling the easy versus difficult manipulation in the adult working

memory task. This modification was necessary as young children are unable to perform

complex spatial working memory tasks[48] but can engage with the more developmentally

appropriate Go/No-go paradigm[49] while still allowing us to probe domain-general cognitive

control functions supported by the MD network. Through this approach, we aim to investigate

whether these regions show distinct functional profiles for language processing versus domain-

general cognitive demands in young children.

To validate our methodology, we first applied the individual fCOI approach to fNIRS

data analysis in a sample of adults in experiment 1, ensuring that our source-detector array was

capable of detecting activation from known functional regions in adult cortex. Then, in

experiment 2, we examined whether the same language-preferring regions with functional

profiles matching those established in the adult literature (i.e., selective responses to linguistic

content and insensitivity to cognitive demands) could be observed in toddlers between 2-4

years of age using the same approach of identifying fCOIs in individual participants.

Materials and Methods

Experiment1

Participants. Twenty adults (between 18 and 26 years, 10 female) were recruited from the student population at Sichuan University. All participants had normal or corrected to normal vision, no history of neurological or psychiatric disorders, head trauma, seizures, or current use of psychoactive medications. None reported physical impairments that could affect task performance. All were right-handed and gave written informed consent. This study was approved by the Biomedical Ethics Review Committee of West China Hospital, Sichuan University (Protocol #2023-2376) and was conducted in accordance with the Declaration of Helsinki and CIOMS International Ethical Guidelines.

Procedure. Participants were seated in front of a 68 cm monitor at a distance of approximately 50 cm. Each participants completed (1) an auditory language localizer task[45]; (2) one spatial working memory task[6].

In the language localizer task, participants listened to blocks of intact meaningful sentences and acoustically degraded speech following the procedure introduced in Scott et al.[45]. All the materials for this localizer were adopted to Mandarin from Gweon et al.[50] and read by a native female speaker. We used female speakers because children tend to pay attention to female voices[51]. Thirty-two audio clips were created, consisting of 24 intact clips and 8 degraded clips (see the full list of stimuli in supplementary material). Each clip was 19-20 seconds in duration. Degraded speech clips were created from the intact versions using the procedure described below, resulting in muffled speech where the linguistic content was no longer intelligible. The degraded versions were created by first applying a low-pass filter (350 Hz pass-band) to the intact audio clips, then creating a noise track by randomizing the time-points of the original audio and multiplying it by the amplitude envelope of the intact clip. The

noise track was low-pass filtered (8,000 Hz pass-band, 10,000 Hz stop frequency) to soften the

highest frequencies and added to the filtered speech at a level that rendered it unintelligible.

Each block lasted around 20 seconds. Each run included three blocks corresponding to three

language conditions plus one block of degraded speech condition, with 10-second fixation

blocks between blocks and additional fixation blocks at the beginning and end of the run. There

are a total of eight runs. The task was originally designed to serve a dual purpose: localizing

language processing and theory of mind networks. However, pilot testing revealed that children

could not sustain this lengthy protocol, leading us to simplify the design to a single language

condition for child participants in Experiment 2. For adult participants, we maintained all three

language conditions but combined them into a single intact speech condition for analysis,

contrasting it with the degraded speech condition. This resulted in an uneven distribution of

stimuli between conditions (24 intact vs. 8 degraded clips). Participants were told that they

would listen to some fun audio clips and some clips that were distorted in a way that makes it

impossible to understand what the speaker is saying. They were instructed to listen attentively.

Prior to the experiment, it was ensured that the volume level was sufficiently loud yet

comfortable.

In the MD localizer task, participants performed a spatial working memory (WM) task

involving a 3x4 grid. During each trial, squares appeared in different locations on the grid,

either one at a time (easy condition) or two at a time (hard condition). Participants were

required to remember these sequential locations. At the end of each trial, they were shown two

sets of locations and had to select the set that matched the sequence they had just seen. They

received feedback indicating whether their response was correct or incorrect[6]. Each trial

followed a precise timing structure: beginning with a 500 ms fixation cross, followed by

sequential grid presentations where each location was shown for 1000 ms. Participants then

had up to 3750 ms to make their choice between the two location sets. After responding, they

received feedback for 250 ms, followed by a fixation cross that filled the remaining time to

ensure a consistent trial length (calculated as 4000 ms minus the response time minus 250 ms).

Each run consisted of two blocks (one easy, one hard) with four trials per block, plus three 10-

second fixation blocks interspersed at the beginning, middle, and end. Participants completed

four runs in total. To ensure task comprehension, all participants completed a practice run

before beginning the main experiment.

After each run the display paused until the participant initiated the next run by pressing

a key on a keyboard located to his/her right. Participants were instructed to remain still and to

focus on the screen throughout each run but were not asked to fixate on any point on the screen

and could adjust their position between runs.

Data acquisition. fNIRS measurements were collected with a continuous wave system

NirSmart (Danyang Huichuang Medical Equipment Co., Ltd., China) using wavelengths of

760 nm and 850 nm with a sampling rate of 11 Hz. 29 sources and 30 probes constituted 79

measurement channels of 3 cm (figure 1). The emitter and detector were placed according to

the 10–20 system. The optodes were stabilized using a plastic holder and then affixed to

participants' heads over the frontal, temporal, occipital and parietal lobes in the left and right

hemisphere using custom headgear. Prior to data collection, head circumference measurements

were taken to ensure all participants fell within the suitable range for our standard cap size (54-

58 cm). When wearing the optode cap, it was ensured that the Cz point of the electrode cap

coincided with the Cz point of the scalp surface measurement. The spatial coordinates of

sources, detectors, and anchor points (Nz, Cz, Al, Ar, lz, and other points from the international

10-20 system) were digitized using an electromagnetic 3D digitizer (Patriot, Polhemus, USA).

These coordinates were then transformed into Montreal Neurological Institute (MNI)

coordinates and mapped onto the MNI standard brain template using the spatial registration

method in NirSpace (Danyang Huichuang Medical Equipment Co., Ltd, China).

FIGURE 1 : (a) Schematic representation of the fNIRS optode placement. Red circles

indicate sources, blue circles indicate detectors, and yellow lines represent measurement

channels. (b) Spatial registration result of the probe array. Red dots with number indicate

channel positions on the MNI standard brain template .

Data preprocessing. Initial measurements of incident light reaching the detector for

each channel were trimmed to remove excess time points from the beginning and end of data

collection and were then preprocessed using the HomER NIRS processing package[52]. First,

each run of the experiment was isolated from the continuous data and processed as a separate

file to ensure that normalization of a given run was not skewed by more or less noisy portions

of the data occurring at other times in the experiment[42]. Then, the negative intensity values

resulting from noisy data were corrected using the hmrR_PreprocessIntensity_Negative

function. Subsequently, channels were visually inspected and those lacking clear cardiac signal

were manually excluded from individual participants' data sets. Raw intensity measurements

were then converted to optical density changes (ΔOD). Then a linear fit function was used to

further remove the baseline drift. Next, a Temporal Derivative Distribution Repair (TDDR)

was used to filter out motion artifacts, which can effectively remove baseline shift and spike artifacts without the need for any user-supplied parameters[53] (Fishburn et al., 2019). The resulting time courses were then band-pass filtered with cut-off frequencies of 0.01–0.09 Hz to further remove slow drifts and high frequency and physiological noise[54] (Pinti et al., 2019, butter filter 3 order for lowpass, 5 order for high pass). After that the ΔOD values were transformed into changes in oxyhemoglobin (HbO2) and deoxyhemoglobin (HHb) concentration according to the modified Beer-Lambert law (hmrOD2Conc, ppf = [1.0 1.0]). Then, the global components were removed using a principal component analysis spatial filter[55,56](σ=46 deg; Zhang et al., 2016; Zhang et al., 2017), which is comparable to using short-source channels when the coverage of the measurement is much larger than the regions of interest[57]. And the concentration of each block is baseline corrected by 2s prior to the onset of stimuli.

Data analysis. We focused our analysis on changes in oxygenated hemoglobin concentration, as it has been demonstrated to be a more sensitive and reliable measure compared to deoxygenated or total hemoglobin concentration changes, which do not provide additional information beyond HbO2[58]. For each participant and channel, we computed the mean change in HbO2 concentration during each block, beginning 2 seconds post-stimulus onset to account for hemodynamic response lag. The analysis window extended to 25 seconds post-onset for the language localizer task and 38 seconds post-onset for the spatial working memory task, since different tasks' blocks have different durations[58,59]. The differences in these responses across conditions were then analyzed using the fCOI approach.

First, we selected a set of channels ("search space")[37] that plausibly covered regions of the scalp plausibly overlaying bilateral inferior frontal lobe across participants (see table 1 for probabilistic mapping of selected fNIRS channels to underlying cortical regions).

261   TABLE 1 Probabilistic mapping of fNIRS channels to underlying cortical regions

262 showing Brodmann areas (based on MRIcro atlas) and corresponding MNI coordinates for each

263 source-detector pair. Percentages indicate the probability of channel placement over the

264 specified anatomical regions.

Channel	MNI coordinates	Anatomic label	Percentage	Search Space
5	64.9643, 5.7288, 27.8566	4 - Primary Motor Cortex, 6 - Pre-Motor and Supplementary Motor Cortex, 43 - Subcentral area, 44 - pars opercularis	6.35%, 59.53%, 27.42%, 6.69%	RIFG
13	61.2752, 22.4516, 12.4687	6 - Pre-Motor and Supplementary Motor Cortex, 44 - pars opercularis, 45 - pars triangularis, 48 - Retrosubicular area	5.75%, 31.95%, 51.12%, 11.18%	RIFG
14	57.049, 37.9202, -2.4401	38 - Temporopolar area, 45 - pars triangularis, 46 - Dorsolateral prefrontal cortex, 47 - Inferior prefrontal gyrus	3.34%, 66.22%, 21.40%, 9.03%	RIFG
17	53.2021, 33.3479, 29.3424	44 - pars opercularis, 45 - pars triangularis, 46 - Dorsolateral prefrontal cortex	6.69%, 92.57%, 0.74%	RIFG
19	49.1243, 47.253, 15.4105	45 - pars triangularis, 46 - Dorsolateral prefrontal cortex	48.70%, 51.30%	RIFG

20	36.6861, 51.5968, 29.9235	9 - Dorsolateral prefrontal cortex, 45 - pars triangularis, 46 - Dorsolateral prefrontal cortex	4.17%, 3.70%, 92.13%	RIFG
68	-38.2423, 52.8662, 27.4047	45 - pars triangularis, 46 - Dorsolateral prefrontal cortex	12.45%, 87.55%	LIFG
69	-43.5347, 37.7257, 38.1227	9 - Dorsolateral prefrontal cortex, 44 - pars opercularis, 45 - pars triangularis, 46 - Dorsolateral prefrontal cortex	25.45%, 4.55%, 39.09%, 30.91%	LIFG
70	-50.5051, 46.6936, 10.4438	45 - pars triangularis, 46 - Dorsolateral prefrontal cortex	53.45%, 46.55%	LIFG
71	-54.2983, 33.6329, 22.7694	45 - pars triangularis	100%	LIFG
72	-56.3003, 37.7076, -7.1488	38 - Temporopolar area, 45 - pars triangularis, 46 - Dorsolateral prefrontal cortex, 47 - Inferior prefrontal gyrus	8.14%, 48.06%, 19.77%, 24.03%	LIFG
73	-59.7137, 21.7772, 5.0031	6 - Pre-Motor and Supplementary Motor Cortex, 38 - Temporopolar area, 44 - pars opercularis, 45 - pars triangularis, 48 - Retrosubicular area	4.03%, 15.44%, 14.09%, 36.24%, 30.20%	LIFG

| **75** | -66.014, 6.3358, 19.328 | 6 - Pre-Motor and Supplementary Motor Cortex, 43 - Subcentral area, 44 - pars opercularis, 48 - Retrosubicular area | 59.79%, 31.12%, 4.90%, 4.20% | LIFG |

The leave-one-run-out cross-validation procedure[42] was implemented within each

search space, which has demonstrated better sensitivity than other selection methods[40]. For

each participant, one run was held out while the remaining runs were averaged. Within the

predefined search space, we identified the eligible channel showing maximum contrast

between intact speech and degraded speech for Language fCOI, and maximum contrast

between hard and easy conditions for MD fCOI. This process was repeated iteratively for each

run and participant. The identified channel was used as the fCOI definition to extract response

values for each condition from each held-out run.

Our fCOI selection method does not guarantee the identification of reliably preferential

channels in either search space. If none of the available channels recorded from cortex

exhibited distinct responses between conditions, the selection process would just operate on

random noise, resulting in no consistent response patterns in the independent data[37]. In other

words, in cases where no robust preferential activation emerges during fCOI selection, any

subsequent analyse may be driven by random variance rather than reflecting a true functional

specialization, warranting cautious interpretation.

To compare the functional profile across the two networks, we also estimated the

responses of the fCOIs to conditions that were not used to define them (i.e., hard and easy WM

for the language fCOI, intact speech and degraded speech for the MD fCOI). For this analysis,

we used all runs from the localizer task to define the fCOIs and all data from the other task to

estimate their responses.

fCOIs were excluded from analyses if more than one-third of the channels in their

search space were excluded[43]. Participants were excluded if they had fewer than 2 runs.

Our main analysis addressed three primary questions. First, regarding functional

selectivity, we examined whether identified language fCOIs show selectivity to the properties

of high-level language, specifically stronger activation in response to intact versus degraded

speech[45]. Second, for functional specificity, we investigated whether language fCOIs exhibit

specificity to language. A lack of difference in the language fCOI between the Easy and Hard

conditions of the spatial working memory task , or a reverse pattern showing stronger responses

to the Easy than the Hard condition, would suggest that the language network is not selectively

engaged in this working memory task. Third, concerning domain-general processing, we

needed to verify that the language fCOIs' lack of response to cognitive load was not simply due

to a failure of our tasks to engage domain-general function[60]. Therefore, we tested whether the

MD fCOIs in our sample exhibited the expected selectivity and specificity for cognitive

demands.

We employed a generalized linear mixed-effects model (GLMM) on individual trials to

account for the unbalanced nature of the dataset (different numbers of runs across participants

due to exclusion criteria) and individual differences in global signal strength. The analysis was

implemented using MATLAB's "fitglme" function with Maximum Penalized Likelihood (MPL)

method. Our model structure was: HbO data ~ condition + (1|subject) where condition

represented the experimental conditions (intact vs degraded speech for language task; hard vs

easy for cognitive/MD task). While inclusion of random slopes for condition by subject would

be theoretically appropriate[61], attempts to include these led to convergence failures. Separate

models were run for two fCOI types in each search space (LIFG language, RIFG language,

LIFG MD, RIFG MD) of two experiment contrasts (Intact > Degraded; Hard > Easy). To

account for multiple comparisons, we applied Bonferroni correction separately for each task

(language vs. cognitive control). In each task, the left and right IFG fCOIs formed a single

family of two comparisons, yielding a family-wise threshold of $\alpha = 0.025$ (0.05/2).

Experiment 2

Participants. Thirty-five children between 2 and 4 years of age were recruited from

preschools in Shanghai. Thirteen children were excluded because of failing to follow the

experimental instructions. Twenty-two children (mean: 3 years 6 months; range: 2 years 5

months–4 years 11 months; 14 male, 8 female) were finally included in the study. Two children

were aged 2 years 5 months to 2 years 11 months, seventeen were aged 3 years 0 months to 3

years 11 months, and three were aged 4 years 0 months to 4 years 11 months. According to

parent reports, all participants had no developmental disorders, had normal or corrected-to-

normal vision and hearing, and could understand basic instructions in Mandarin. Additionally,

all children were full-term births (≥ 37 weeks gestation) with no significant medical

complications at birth. Written informed consent was obtained from parents/legal guardians of

all child participants prior to their inclusion in the study, in accordance with the requirements

of the Biomedical Ethics Review Committee of West China Hospital, Sichuan University

(Protocol #2023-2376). The study was conducted following the principles of the Declaration

of Helsinki and CIOMS International Ethical Guidelines

Procedure. The procedure was similar to that of experiment 1 except that children

under 3 years sat on their parents' laps while viewing the display, and the experiment was

discontinued, either during or between runs, if the participant became fussy, inattentive or if

the parent indicated their wish to end the experiment. Each participants completed an auditory

language localizer task and a Go/no-go task.

In the language localizer task, participants listened to blocks of intact meaningful

sentences and acoustically degraded speech following the same procedure described in

experiment 1. The materials were adapted from experiment 1 by selecting a subset of the adult

stimuli based on a pre-experiment assessment of comprehension in a separate group of toddlers

(n=3). While the adult version contained three language conditions per run, we selected only

the material that showed the highest comprehension rates in our pre-experiment (accurate

responses to simple comprehension questions. E.g. "Was the girl helping her father?" or "Was

the rabbit in the garden?"). The selected narratives were shortened from the original 20-second

duration to 18-second segments to accommodate shorter attention spans, while maintaining the

core narrative structure. Each block lasted 18 seconds, with 6-second fixation blocks between

blocks and additional fixation blocks at the beginning and end of the run. Each run included

one language condition block plus one degraded speech condition block. There are four total

runs. As in experiment 1, participants were told that they would listen to some fun audio clips

and some clips that were distorted in a way that makes it impossible to understand what the

speaker is saying. They were instructed to listen attentively. Prior to the experiment, it was

ensured that the volume level was sufficiently loud yet comfortable.

In the MD localizer task, participants completed a Go/no-go task. A training session

before the experiment was conducted to ensure task comprehension. The training included

practice trials for both "go" (simple response) and "go/no-go" (response inhibition) conditions

with performance feedback, continuing until participants demonstrated reliable response

patterns. The formal experiment consisted of four runs, each containing alternating "go" and

"go/no-go" blocks separated by 10-second fixation periods. Each block began with a 3-second

instruction screen followed by 12 trials lasting 12 seconds total. In "go" blocks, participants

pressed a key with their right index finger in response to both cat and dog images. In "go/no-

go" blocks, participants pressed the key for chicken images (go) but withheld responses for

duck images (no-go). A study personnel monitored participant performance during scanning.

Data acquisition. fNIRS measurements were collected with a continuous wave system

NirSmart (Danyang Huichuang Medical Equipment Co., Ltd., China) using wavelengths of

730nm, 808 nm and 850 nm with a sampling rate of 11 Hz. 26 sources and 27 probes constituted

81 measurement channels of 3 cm (see figure 2). The optodes were stabilized using a plastic

holder and then affixed to participants' heads over the frontal, temporal, occipital and parietal

lobes in the left and right hemisphere using custom headgear. When wearing the optode cap, it

was ensured that the Cz point of the electrode cap coincided with the Cz point of the scalp

surface measurement.

FIGURE 2 : (a) Schematic representation of the fNIRS optode placement. Red circles

indicate sources, blue circles indicate detectors, and yellow lines represent measurement

channels. (b) Spatial registration result of the probe array. Red dots with number indicate

channel positions on the MNI standard brain template .

Data preprocessing. Data were preprocessed in the same manner as Experiment 1,

with the exception of the partial pathlength factor (ppf) in the modified Beer-Lambert law

calculation. Given the different optical properties of toddler brain tissue compared to adults,

we used a ppf value of [2.0 2.0 2.0], which better accounts for the light scattering and

absorption characteristics in infant participants[62].

Data analysis. For each participant and channel, mean HbO2 concentration changes

were calculated per trial. The analysis window started 4 seconds after stimulus onset and lasted

until 17 seconds post-onset for the language localizer task or 12 seconds post-onset for the

Go/no-go task. This timing accounts for the HRF lag time commonly assumed in children's

NIRS research[63,64]. Language and MD selective regions were identified and mean HbO

responses were extracted using individual fCOI approaches within the same search space (see

table 2 for probabilistic mapping of selected fNIRS channels to underlying cortical regions),

following Experiment 1's methodology. Inclusion criteria required children to complete at least

two runs per experiment, ensuring sufficient trials for the leave-one-run-out procedure.

Additional exploratory analyses incorporating age as a fixed effect in the generalized linear

mixed models are reported in the supplementary material.

TABLE 2 Probabilistic mapping of fNIRS channels to underlying cortical regions

showing Brodmann areas (based on MRIcro atlas) and corresponding MNI coordinates for each

source-detector pair. Percentages indicate the probability of channel placement over the

specified anatomical regions.

Channel	MNI coordinates	Anatomic label	Percentage	Search Space
17	77.67, 19.00, 17.33	6 – Pre-Motor and Supplementary Motor Cortex, 43 – Subcentral	51.44%, 0.32%,	RIFG

		area, 44 – pars opercularis, 45 – pars triangularis Broca's area, 48 – Retrosubicular area	42.49%, 4.79%, 0.96%	
18	69.67, 20.33, 44.67	6 – Pre-Motor and Supplementary Motor Cortex, 9 – Dorsolateral prefrontal cortex, 44 – pars opercularis	36.02%, 9.20%, 54.79%	RIFG
25	69.33, 42.67, 13.33	45 – pars triangularis Broca's area, 46 – Dorsolateral prefrontal cortex	99.32%, 0.68%	RIFG
26	60.33, 42.67, 40.33	9 – Dorsolateral prefrontal cortex, 44 – pars opercularis, 45 – pars triangularis Broca's area, 46 – Dorsolateral prefrontal cortex	0.81%, 6.07%, 75.30%, 17.81%	RIFG
27	59.67, 60.33, 5.67	10 – Frontopolar area, 45 – pars triangularis Broca's area, 46 – Dorsolateral prefrontal cortex	4.14%, 7.52%, 88.35%	RIFG
56	46.00, 42.33, 59.67	10 – Frontopolar area, 45 – pars triangularis Broca's area, 46 – Dorsolateral prefrontal cortex	0.85%, 4.68%, 94.47%	LIFG
57	33.00, 36.00, 73.00	44 – pars opercularis, 45 – pars triangularis Broca's area, 46 – Dorsolateral prefrontal cortex	1.23%, 90.95%, 7.82%	LIFG
59	33.67, 62.67, 50.67	45 – pars triangularis Broca's area, 46 – Dorsolateral prefrontal cortex	99.65%, 0.35%	LIFG

64	-57.67, -46.33, 79.33	6 – Pre-Motor and Supplementary Motor Cortex, 44 – pars opercularis	33.45%, 66.55%	LIFG
66	-70.33, 18.33, 39.00	6 – Pre-Motor and Supplementary Motor Cortex, 44 – pars opercularis, 45 – pars triangularis Broca's area, 48 – Retrosubicular area	49.49%, 31.99%, 0.67%, 17.85%	LIFG

<div align="center">**Results**</div>

Experiment 1

No participants were excluded based on our criteria for channel rejection and fCOI

rejection. For the subsequent analyses, we included all data after removing the excluded noisy

channel and any runs where noisy channels comprised more than one-third of the search space.

Table 3 provides a summary of the available runs across participants for different fCOI types

(See the distribution of identified channel for each type of fCOI in supplementary material in

figure S4-S8).

TABLE 3 Summary of the available runs across participants for different fCOI types

fCOI Type	Total LAN Subjects Included	Mean LAN Runs (SDs)	Total MD Subjects Included	Mean MD Runs (SDs)
LAN LIFG	20	6.95(0.683)	20	3.85(0.489)

LAN RIFG	20	7.15(0.3663)	20	3.85(0.489)
MD LIFG	20	7.10(0.4472)	20	3.90(0.4472)
MD RIFG	20	7.15(0.3663)	20	3.90(0.4472)

The language fCOIs showed clear selectivity for language processing. In both

hemispheres, these channels demonstrated significantly stronger responses to intact versus

degraded speech. The LIFG language fCOI showed a significant effect of condition (β = -8.92e-

06, SE = 3.04e-06, t = -2.9365, p = 0.0036), with higher activation for intact speech. Similarly,

the right inferior frontal gyrus (RIFG) language fCOI exhibited a significant preference for

intact speech (β = -1.18e-05, SE = 3.36e-06, t = -3.5248, p = 0.00049). Importantly, when tested

on the spatial working memory task, neither the left nor right language fCOIs showed

significant modulation by cognitive demand (LIFG: β = -4.66e-06, SE = 5.31e-06, t = -0.87838,

p = 0.381; RIFG: β = -2.1894e-06, SE = 4.9866e-06, t = -0.43906, p = 0.661). This pattern

suggests that the language fCOIs are specifically selective for language processing and do not

respond systematically to increased cognitive load in non-linguistic tasks (figure 3a & 3b).

The MD fCOIs demonstrated a complementary pattern of activation. Both left and right

MD fCOIs showed significant sensitivity to cognitive demand in the spatial working memory

task. The LIFG MD fCOI exhibited stronger activation for harder versus easier conditions (β

= 1.34e-05, SE = 5.09e-06, t = 2.6279, p = 0.0095), as did the RIFG MD fCOI (β = 1.75e-05,

SE = 5.26e-06, t = 3.3278, p = 0.0011). In contrast, when tested on the language task, neither

MD fCOI showed significant differentiation between intact and degraded speech (LIFG: β =

5.04e-08, SE = 3.25e-06, t = 0.015523, p = 0.988; RIFG: β = 1.01e-06, SE = 3.25e-06, t =

0.31183, p = 0.756). This finding indicates that the MD network responds selectively to

cognitive demand but not to linguistic properties per se (figure 3c & 3d).

FIGURE 3 Results from Experiment 1: Mean HbO2 responses in the selected fCOIs

(a) Left hemisphere language fCOI responses showing significant selectivity for intact versus

degraded speech (p = 0.0036) and no significant modulation by cognitive demand (p = 0.381).

(b) Right hemisphere language fCOI responses showing significant selectivity for intact versus

degraded speech (p = 0.00049) and no significant modulation by cognitive demand (p = 0.661).

(c) Left hemisphere MD fCOI responses showing significant sensitivity to cognitive demand

(p = 0.0095) and no significant difference between speech conditions (p = 0.988). (d) Right

hemisphere MD fCOI responses showing significant sensitivity to cognitive demand (p =

0.0011) and no significant difference between speech conditions (p = 0.756). Error bars

represent standard error of the mean.

These results demonstrate a double dissociation between the language and MD networks, with language fCOIs showing selectivity for linguistic processing but not cognitive demand, and MD fCOIs showing the opposite pattern.

Experiment 2

Table 4 provides a summary of the included participants and available runs across participants for different fCOI types based on the criteria for fCOIs rejection we described in

440   experiment 1 (See the distribution of identified channel for each type of fCOI in supplementary

441   material, Figure S1-S4).

442   TABLE 4 Summary of the available runs across participants for different fCOI types

fCOI Type	Total LAN Subjects Included	Mean LAN Runs (SDs)	Total MD Subjects Included	Mean MD Runs (SDs)
LAN LIFG	22	3.54(0.510)	21	2.24(0.436)
LAN RIFG	22	3.50(0.597)	17	2.23(0.437)
MD RIFG	22	3.50(0.597)	22	2.27(0.457)
MD LIFG	21	3.38(0.74)	21	2.24(0.539)

Analysis of the language fCOIs in children showed a partially overlapping pattern with

adults. The LIFG language fCOI maintained significant selectivity for language processing,

showing stronger responses to intact versus degraded speech (β = -5.21e-07, SE = 1.69e-07, t

= -3.0781, p = 0.0025). However, unlike adults, the RIFG language fCOI did not show

significant differentiation between intact and degraded speech (β = -7.49e-08, SE = 1.90e-07,

t = 0.1211, p = 0.693). Consistent with the adult findings, neither language fCOI showed

significant modulation by cognitive demand in the Go/no-go task (LIFG: β = -3.22e-08, SE =

1.46e-07, t = -0.21987, p = 0.826; RIFG: β = -8.69e-08, SE = 2.01e-07, t = -0.72473, p = 0.667).

This suggests that even in children, the language fCOI we identified maintains specificity for

language processing over general cognitive demands (figure 4a & 4b).

The MD fCOI we identified in children showed a more complex pattern than expected.

Behavioral performance confirmed the task difficulty manipulation, with participants showing

higher accuracy on go trials (M = 81%, SD = 12%) compared to no-go trials (M = 73%, SD =

15%; t(21) = 2.25, p = 0.035). Only the right MD fCOI displayed higher sensitivity to cognitive

demand in the Go/no-go task (significantly stronger responses to the harder versus easier

condition; RIFG: β = 4.49e-07, SE = 1.48e-07, t = 3.0185, p = 0.00323). By contrast, the left

MD fCOI did not show a significant difference between these conditions (LIFG: β = -8.63e-

08, SE = 1.25e-07, t = -0.69016, p = 0.4917). When we tested these fCOI in the language task,

an unexpected pattern emerged: although the left MD fCOI was not strongly modulated by

cognitive control demands, it nonetheless responded more robustly to degraded than to intact

speech (β = 4.2037e-07, SE = 1.7142e-07, t = 2.4523, p = 0.015313). The right MD fCOI

showed a similar trend though not reaching statistical significance after correction for multiple

comparisons (β = 3.6637e-07, SE = 1.8245e-07, t = 2.0081, p = 0.046). These findings suggest

that only the right MD fCOI displays the characteristic MD network profile of responding to

cognitive control demands but not to linguistic properties. The left MD fCOI's lack of

modulation by task difficulty suggests that an adult-like MD network component may not yet

be established in the left IFG in our sample, warranting caution in interpreting its preferential

response to degraded speech as we have discussed in the method part (figure 4c & 4d).

FIGURE 4 Results from Experiment 2: Mean HbO2 responses in the selected fCOIs (a)

Left hemisphere language fCOI responses showing significant selectivity for intact versus

degraded speech (p = 0.0025) and no significant modulation by cognitive demand (p =

0.826). (b) Right hemisphere language fCOI responses showing no significant selectivity for

speech conditions (p = 0.693) and no significant modulation by cognitive demand (p =

0.667). (c) Left hemisphere MD fCOI responses showing no significant sensitivity to

cognitive demand (p = 0.4917) but significant preference for degraded speech (p = 0.015). (d)

Right hemisphere MD fCOI responses showing significant sensitivity to cognitive demand (p

= 0.00323) and marginal preference for degraded speech (p = 0.046). Error bars represent

standard error of the mean.

<center>**Discussion**</center>

Our study investigates how the brain supports language processing in early childhood,

focusing on whether neural circuits are specialized for language or share with other cognitive

domains. By employing fNIRS with an individual functional channel of interest approach, we

assessed both adults and toddlers on language and cognitive control tasks to seek evidence for

distinct functional profiles in the left and right inferior frontal regions. Our adult findings

validated the fCOI method by replicating well-established dissociations between language-

specific and domain-general processor, providing a foundation for investigating the

developmental period. Our toddler results suggest that left-hemisphere language selectivity

emerges early and remains robust, while right-hemisphere organization shows signs of ongoing

maturation.

In line with adult findings and recent work in older children[29,65], toddlers demonstrated

robust language selectivity in the left IFG. The language fCOI within this area showed specific

responses to intact versus degraded speech, mirroring the adult pattern, while remaining

insensitive to the manipulation of cognitive demands in the Go/No-Go task. This supports the

hypothesis of early domain specificity in language processing, as the language selective region

within left IFG region exhibited both selectivity and specificity for language[36,60]. Our toddler

data align with evidence that even very young children recruit the left frontal cortex for high-

level linguistic processing[33,35,66]. Moreover, our findings revealed a developmental asymmetry

in language processing between hemispheres. While our adult participants showed bilateral

IFG fCOIs with selectivity for meaningful speech, toddlers did not exhibit statistically reliable

language responses in the RIFG language fCOI. This asymmetry aligns with previous

developmental research and recent findings in older children. Hiersche et al.[29], using a similar

speech localizer task comparing sentences to acoustically degraded speech, also found no

significant language selectivity in the right IFG. Early in development, infant studies have

documented limited right-hemisphere sensitivity to language in speech versus control

conditions[33,66]. A developmental shift occurs later: children ages 5-11 show right hemisphere

engagement, particularly during complex linguistic tasks requiring greater syntactic processing

or elevated linguistic demands[67–69]. Our observation of strong LIFG but nonsignificant RIFG

selectivity in toddlers (ages 2-4) suggests that RIFG language specialization follows a

prolonged developmental trajectory. While LIFG supports core linguistic computations from

early development, RIFG may serve complementary functions that become increasingly

important as language abilities mature. Studies suggest RIFG involvement in prosodic

processing, pragmatic aspects of language, and discourse-level comprehension[70,71]. The

absence of RIFG language selectivity in toddlers suggest that these higher-level linguistic

functions develop later, potentially aligning with children's growing ability to process complex

narrative[72] and social aspects of language[73]. These findings support a developmental model

where left-hemisphere language selectivity emerges early and remains stable, while right-

hemisphere language involvement develops gradually throughout childhood[67,74].

With regard to domain-general processing, our findings revealed a distinct pattern in

toddlers from adults: only the right MD fCOI showed selective sensitivity to increased

cognitive demands during the Go/No-Go task. These MD fCOIs demonstrated functional

profiles that contrasted with language fCOIs - while language regions responded selectively to

intact versus degraded speech but showed no sensitivity to cognitive demands, MD fCOIs

exhibited the opposite pattern. Importantly, only the right MD fCOI displayed the characteristic

MD network profile of responding to cognitive control demands but not to linguistic properties.

This toddler-specific pattern diverges from established findings in both adults and older

children (approximately 7-12 years). Adults show bilateral MD network responses to escalating

task difficulty, and older children demonstrate bilateral prefrontal recruitment during inhibitory

control tasks[75–78]. The developmental differences likely reflect the ongoing structural and

functional specialization of the MD network in early childhood. Our findings of right MD

responsiveness in toddlers, coupled with previous fNIRS and fMRI studies, suggest that very

young children's prefrontal cortex involvement in executive tasks may initially manifests in

right hemisphere circuits compared to older groups[79,80]. This aligns with evidence of protracted

maturation for left frontal functions[75,77,81]. The left MD fCOI's lack of modulation by task

difficulty suggests that an adult-like MD network component may not yet be established in the

left IFG in our sample. This developmental pattern is further corroborated by recent work from

Schettini et al.[30] who similarly found that the left IFG in children did not show significant

selectivity for cognitive demand. While the left MD fCOI exhibited stronger activation to

degraded compared to intact speech, this pattern should be interpreted cautiously, as it lacks

the hallmark feature of the MD network—sensitivity to cognitive demand—and may reflect

immature functional engagement in this region. An alternative possibility is that degraded

speech, which is more challenging to understand than intact speech, elicits more effortful

processing in children, thereby driving up activity in what would eventually become a domain-

general region. In other words, although the left MD fCOI does not show the classic adult-like

sensitivity to cognitive demands, it may nonetheless be recruited when children strain to extract

meaning from ambiguous input. Together, these findings support previous research indicating

ongoing development of prefrontal involvement in domain-general control[30,74,76], with bilateral

integration emerging later in development as children increasingly recruit both hemispheres

for executive tasks.

Our use of the fCOI approach addressed several methodological challenges in fNIRS

studies. First, it circumvents the limitations of the 'reverse inference' problem which plagues

many fNIRS studies that rely on inferring cognitive functions solely from anatomical

correspondence with prior findings. This practice is problematic as the same anatomical

location may support different functions across individuals[82], particularly in developmental

research where variations in head size, brain organization, and probe placement complicate

anatomical localization[41]. Our fCOI approach addresses these limitations by identifying

regions based on functional properties rather than assumed anatomical locations, effectively

accommodating individual variability in neural functional anatomy across development.

Second, it provided enhanced statistical power while controlling for multiple comparisons,

which is crucial when working with limited developmental data[40]. Third, the implementation

of the same localizer task used by fMRI studies[7,29,65] allowed for direct comparison of

properties of specific functional regions over developmental stages and imaging modalities.

Our successful identification of functionally distinct networks in both adults and toddlers

demonstrates the robustness of this approach across developmental stages.

Our experimental design also addressed several methodological challenges. The

implementation of a leave-one-run-out cross-validation procedure for fCOI selection avoided

the "double dipping" issue in neuroimaging studies[83], enhancing the reliability of our findings.

By employing passive language tasks contrasting meaningful versus degraded speech, we

controlled for confounding effects of task difficulty while maintaining prosodic and rhythmic

features[45]. This paradigm, extensively validated in adult studies across different languages,

allowed us to specifically probe the processing of high-level linguistic features such as

semantics and syntax[46]. The careful selection of age-appropriate tasks for probing the MD

network—spatial working memory for adults and Go/No-go for children—enabled us to

examine domain-general processing while accommodating developmental constraints. By

comparing responses to both linguistic and non-linguistic tasks within the same participants,

we directly assessed the specificity of these networks to language processing.

Despite these methodological strengths, several limitations warrant discussion. First,

like most fNIRS experiments, our study faced challenges of limited and non-continuous

cortical coverage. While our probe arrangement provided good coverage of bilateral inferior

frontal regions, some cortical areas may have fallen into measurement "blind spots"[84] . Second,

we did not have access to short-separation channels in our study. Short-separation channels are

efficient in accounting for and removing superficial hemodynamic signals originating from the

scalp and skull, thereby isolating the cortical activity of interest[85,86]. The absence of short-

separation channels may have led to contamination of our signals by systemic physiological

artifacts. Although we have applied global regression method to filter out systemic

physiological artifacts[55], systemic physiological artifacts may still potentially affect the

specificity and accuracy of our findings. Third, different cognitive tasks were used for adults

and children. While both tasks engage the MD network, the cognitive demands they place on

participants differ. However, this adaptation was necessary given the age-appropriate task

requirements, and both tasks share the feature of manipulating cognitive control demands.

Fourth, while our peak-channel selection approach within predefined anatomical regions of

interest was optimized for our specific hypotheses, alternative approaches might be valuable

for different research questions. Additionally, we did not examine behavioral metrics of

language development, which limits our ability to differentiate between changes in language

selectivity due to maturation versus language skill development[87]. Finally, our focus was

primarily on the inferior frontal cortex, which constrained our ability to fully characterize both

the MD network and the language-selective network. While the IFG is a critical hub for

domain-general cognitive control[9] and language processing[10], the broader language network

encompasses distributed temporal, parietal, and superior frontal regions. By restricting our

measurements to the IFG, we may have overlooked crucial activity in other regions of the

language network (e.g., the posterior temporal lobe) that could further elucidate the transition

from domain-general to domain-specific processes in language development. Consequently,

caution is warranted when generalizing our findings to the broader MD and language networks.

Future research should extend these findings in two main ways. First, longitudinal

studies beginning at earlier ages[5,66] are needed to clarify whether language networks transition

from domain-general to domain-specific functioning over time. Second, studies should

examine neural specialization at key language acquisition milestones by collecting brain

measures when children reach specific linguistic achievements (e.g., first words, word

combinations, complex sentences). This milestone-based approach[7,87] would illuminate how

maturing brain circuits support children's emerging linguistic abilities.

To conclude, using the methodologically improved fCOI approach with fNIRS, we

replicated well-established patterns of bilateral selectivity and specificity in adult participants

during linguistic and cognitive control tasks. In toddlers, we found early specialization of the

language-selective component within LIFG, while the homologous right hemisphere region

had not yet developed language selectivity. On cognitive control tasks, toddlers showed adult-

like patterns in the within RIFG only. Notably, the selected MD fCOI in toddler's LIFG only

showed engagement with degraded speech rather than the broader domain-general profile seen

in adults, suggesting ongoing maturation of domain-general circuits. These findings advance

our understanding of how specialized brain circuits for language emerge during early childhood

and establish the utility of the fCOI approach for studying functional organization in the

developing brain.

Disclosures

The authors declare no competing financial interests.

Code and Data Availability

The data that support the findings of this study are available from the corresponding author

upon reasonable request. The data are not publicly available due to privacy or ethical

restrictions involving data from minor participants.

Acknowledgments

We express our sincere gratitude to all participating families and children for their invaluable

contributions to this study. We thank WEN Jian for her dedicated efforts in data collection. We

are deeply grateful to Evelina Fedorenko, Arturo E. Hernandez, CHEN Xiaocong, and QI

Zhenghan for their insightful discussions and guidance. Our thanks extend to CHEN Jiahan,

YANG Jingdan, and GENG Luyuan for their assistance in adapting the materials into Chinese.

We also acknowledge the crucial support of the preschools affiliated with the Shanghai

Maternal and Child Health Center in participant recruitment.

This research was supported by The Hong Kong Polytechnic University Presidential PhD

Fellowship Scheme (PPPFS) and Ordos Clinical Medicine Translational Innovation Center

Program Science and Technology Cooperation Program of Shanghai Jiaotong University in

Inner Mongolia Autonomous Region—Action Plan of Shanghai Jiaotong University for

"Revitalizing Inner Mongolia through Science and Technology" (KJXM2024-11-01).

References

1.   Sharp HM, Hillenbrand K. Speech and Language Development and Disorders in
Children. *Pediatric Clinics of North America*. 2008;55(5):1159-1173.
doi:10.1016/j.pcl.2008.07.007

2.   Chomsky N. Three Factors in Language Design. *Linguistic Inquiry*. 2005;36(1):1-22.
doi:10.1162/0024389052993655

3.   Spelke ES. Core Knowledge, Language, and Number. *Language Learning and
Development*. 2017;13(2):147-170. doi:10.1080/15475441.2016.1263572

4.   Johnson MH. Interactive Specialization: A domain-general framework for human functional brain development? *Developmental Cognitive Neuroscience*. 2011;1(1):7-21. doi:10.1016/j.dcn.2010.07.003

5.   Karmiloff-Smith A, Birkbeck Centre for Brain & Cognitive Development, University of London, London WC1N 7HX, UK. An Alternative to Domain-general or Domain-specific Frameworks for Theorizing about Human Evolution and Ontogenesis. *AIMS Neuroscience*. 2015;2(2):91-104. doi:10.3934/Neuroscience.2015.2.91

6.   Fedorenko E, Behr MK, Kanwisher N. Functional specificity for high-level linguistic processing in the human brain. *Proc Natl Acad Sci USA*. 2011;108(39):16428-16433. doi:10.1073/pnas.1112937108

7.   Fedorenko E, Ivanova AA, Regev TI. The language network as a natural kind within the broader landscape of the human brain. *Nat Rev Neurosci*. 2024;25(5):289-312. doi:10.1038/s41583-024-00802-4

8.   Vigneau M, Beaucousin V, Hervé PY, et al. Meta-analyzing left hemisphere language areas: Phonology, semantics, and sentence processing. *NeuroImage*. 2006;30(4):1414-1432. doi:10.1016/j.neuroimage.2005.11.002

9.   Duncan J. The multiple-demand (MD) system of the primate brain: mental programs for intelligent behaviour. *Trends in Cognitive Sciences*. 2010;14(4):172-179. doi:10.1016/j.tics.2010.01.004

10.   Fedorenko E, Duncan J, Kanwisher N. Broad domain generality in focal regions of frontal and parietal cortex. *Proc Natl Acad Sci USA*. 2013;110(41):16616-16621. doi:10.1073/pnas.1315235110

11.   Hagoort P. Nodes and networks in the neural architecture for language: Broca's region and beyond. *Current Opinion in Neurobiology*. 2014;28:136-141. doi:10.1016/j.conb.2014.07.013

12.   Novick JM, Trueswell JC, Thompson-Schill SL. Cognitive control and parsing: Reexamining the role of Broca's area in sentence comprehension. *Cognitive, Affective, & Behavioral Neuroscience*. 2005;5(3):263-281. doi:10.3758/CABN.5.3.263

13.   Fedorenko E, Blank IA. Broca's Area Is Not a Natural Kind. *Trends in Cognitive Sciences*. 2020;24(4):270-284. doi:10.1016/j.tics.2020.01.001

14.   Blank I, Kanwisher N, Fedorenko E. A functional dissociation between language and multiple-demand systems revealed in patterns of BOLD signal fluctuations. *Journal of Neurophysiology*. 2014;112(5):1105-1118. doi:10.1152/jn.00884.2013

15.   Fedorenko E, Shain C. Similarity of Computations Across Domains Does Not Imply Shared Implementation: The Case of Language Comprehension. *Curr Dir Psychol Sci*. 2021;30(6):526-534. doi:10.1177/09637214211046955

16.   Mineroff Z, Blank IA, Mahowald K, Fedorenko E. A robust dissociation among the
language, multiple demand, and default mode networks: Evidence from inter-region
correlations   in   effect   size.   *Neuropsychologia*.   2018;119:501-511.
doi:10.1016/j.neuropsychologia.2018.09.011

17.   Diachek E, Blank I, Siegelman M, Affourtit J, Fedorenko E. The Domain-General
Multiple Demand (MD) Network Does Not Support Core Aspects of Language Comprehension:
A   Large-Scale   fMRI   Investigation.   *J   Neurosci*.   2020;40(23):4536-4550.
doi:10.1523/JNEUROSCI.2036-19.2020

18.   Quillen IA, Yen M, Wilson SM. Distinct Neural Correlates of Linguistic and Non-
Linguistic Demand. *Neurobiology of Language*. 2021;2(2):202-225. doi:10.1162/nol_a_00031

19.   Wehbe L, Blank IA, Shain C, et al. Incremental Language Comprehension Difficulty
Predicts Activity in the Language Network but Not the Multiple Demand Network. *Cerebral*
Cortex. 2021;31(9):4006-4023. doi:10.1093/cercor/bhab065

20.   Fedorenko E, Varley R. Language and thought are not the same thing: evidence from
neuroimaging and neurological patients: Language versus thought. *Ann NY Acad Sci*.
2016;1369(1):132-153. doi:10.1111/nyas.13046

21.   Paunov AM, Blank IA, Fedorenko E. Functionally distinct language and Theory of
Mind networks are synchronized at rest and during language comprehension. *Journal of*
Neurophysiology. 2019;121(4):1244-1265. doi:10.1152/jn.00619.2018

22.   Woolgar A, Duncan J, Manes F, Fedorenko E. Fluid intelligence is supported by the
multiple-demand system not the language system. *Nat Hum Behav*. 2018;2(3):200-204.
doi:10.1038/s41562-017-0282-3

23.   MacGregor LJ, Gilbert RA, Balewski Z, et al. Causal Contributions of the Domain-
General (Multiple Demand) and the Language-Selective Brain Networks to Perceptual and
Semantic Challenges in Speech Comprehension. *Neurobiology of Language*. 2022;3(4):665-
698. doi:10.1162/nol_a_00081

24.   Shain C, Blank IA, Van Schijndel M, Schuler W, Fedorenko E. fMRI reveals language-
specific predictive coding during naturalistic sentence comprehension. *Neuropsychologia*.
2020;138:107307. doi:10.1016/j.neuropsychologia.2019.107307

25.   Campbell KL, Tyler LK. Language-related domain-specific and domain-general
systems in the human brain. *Current Opinion in Behavioral Sciences*. 2018;21:132-137.
doi:10.1016/j.cobeha.2018.04.008

26.   Mcmullen E, Saffran JR. Music and Language: A Developmental Comparison. *Music*
Perception. 2004;21(3):289-311. doi:10.1525/mp.2004.21.3.289

27.   Rakison DH, Yermolayeva Y. How to Identify a Domain-General Learning Mechanism
When You See One. *Journal of Cognition and Development*. 2011;12(2):134-153.
doi:10.1080/15248372.2010.535228

28.   Thiessen ED, Saffran JR. Learning to Learn: Infants' Acquisition of Stress-Based Strategies for Word Segmentation. *Language Learning and Development*. 2007;3(1):73-100. doi:10.1207/s15473341lld0301_3

29.   Hiersche KJ, Schettini E, Li J, Saygin ZM. Functional dissociation of the language network and other cognition in early childhood. *Human Brain Mapping*. 2024;45(9):e26757. doi:10.1002/hbm.26757

30.   Schettini E, Hiersche KJ, Saygin ZM. Individual Variability in Performance Reflects Selectivity of the Multiple Demand Network among Children and Adults. *J Neurosci*. 2023;43(11):1940-1951. doi:10.1523/JNEUROSCI.1460-22.2023

31.   Gervain J, Mehler J, Werker JF, et al. Near-infrared spectroscopy: A report from the McDonnell infant methodology consortium. *Developmental Cognitive Neuroscience*. 2011;1(1):22-46. doi:10.1016/j.dcn.2010.07.004

32.   May L, Gervain J, Carreiras M, Werker JF. The specificity of the neural response to speech at birth. *Developmental Science*. 2018;21(3):e12564. doi:10.1111/desc.12564

33.   Peña M, Maki A, Kovačić D, et al. Sounds and silence: An optical topography study of language recognition at birth. *Proc Natl Acad Sci USA*. 2003;100(20):11702-11705. doi:10.1073/pnas.1934290100

34.   Sato H, Hirabayashi Y, Tsubokura H, et al. Cerebral hemodynamics in newborn infants exposed to speech sounds: A whole-head optical topography study. *Human Brain Mapping*. 2012;33(9):2092-2103. doi:10.1002/hbm.21350

35.   Minagawa-Kawai Y, Van Der Lely H, Ramus F, Sato Y, Mazuka R, Dupoux E. Optical Brain Imaging Reveals General Auditory and Language-Specific Processing in Early Infant Development. *Cerebral Cortex*. 2011;21(2):254-261. doi:10.1093/cercor/bhq082

36.   Kanwisher N. Functional specificity in the human brain: A window into the functional architecture of the mind. *Proc Natl Acad Sci USA*. 2010;107(25):11163-11170. doi:10.1073/pnas.1005062107

37.   Powell LJ, Deen B, Saxe R. Using individual functional channels of interest to study cortical development with FNIRS. *Developmental Science*. 2018;21(4):e12595. doi:10.1111/desc.12595

38.   Amunts K, Schleicher A, Bürgel U, Mohlberg H, Uylings HBM, Zilles K. Broca's region revisited: Cytoarchitecture and intersubject variability. *J Comp Neurol*. 1999;412(2):319-341.

39.   Wang L, Mruczek REB, Arcaro MJ, Kastner S. Probabilistic Maps of Visual Topography in Human Cortex. *Cereb Cortex*. 2015;25(10):3911-3931. doi:10.1093/cercor/bhu277

40. Liu Y, Sánchez Hernández F, Ting F, Hyde DC. Comparing fixed-array and functionally-defined channel of interest approaches to infant functional near-infrared spectroscopy data. *NeuroImage*. 2022;261:119520. doi:10.1016/j.neuroimage.2022.119520

41. Kabdebon C, Leroy F, Simmonet H, Perrot M, Dubois J, Dehaene-Lambertz G. Anatomical correlations of the international 10–20 sensor placement system in infants. *NeuroImage*. 2014;99:342-356. doi:10.1016/j.neuroimage.2014.05.046

42. Hyde DC, Simon CE, Ting F, Nikolaeva JI. Functional Organization of the Temporal–Parietal Junction for Theory of Mind in Preverbal Infants: A Near-Infrared Spectroscopy Study. *J Neurosci*. 2018;38(18):4264-4274. doi:10.1523/JNEUROSCI.0264-17.2018

43. Richardson H, Taylor J, Kane-Grade F, Powell L, Bosquet Enlow M, Nelson CA. Preferential responses to faces in superior temporal and medial prefrontal cortex in three-year-old children. *Developmental Cognitive Neuroscience*. 2021;50:100984. doi:10.1016/j.dcn.2021.100984

44. Hoshi Y. Functional near-infrared optical imaging: Utility and limitations in human brain mapping. *Psychophysiology*. 2003;40(4):511-520. doi:10.1111/1469-8986.00053

45. Scott TL, Gallée J, Fedorenko E. A new fun and robust version of an fMRI localizer for the frontotemporal language system. *Cognitive Neuroscience*. 2017;8(3):167-176. doi:10.1080/17588928.2016.1201466

46. Malik-Moraleda S, Ayyash D, Gallée J, et al. An investigation across 45 languages and 12 language families reveals a universal language network. *Nat Neurosci*. 2022;25(8):1014-1019. doi:10.1038/s41593-022-01114-5

47. Duncan J, Owen AM. Common regions of the human frontal lobe recruited by diverse cognitive demands. *Trends in Neurosciences*. 2000;23(10):475-483. doi:10.1016/S0166-2236(00)01633-7

48. Pailian H, Libertus ME, Feigenson L, Halberda J. Visual working memory capacity increases between ages 3 and 8 years, controlling for gains in attention, perception, and executive control. *Atten Percept Psychophys*. 2016;78(6):1556-1573. doi:10.3758/s13414-016-1140-5

49. Simpson A, Riggs KJ. Conditions under which children experience inhibitory difficulty with a "button-press" go/no-go task. *Journal of Experimental Child Psychology*. 2006;94(1):18-26. doi:10.1016/j.jecp.2005.10.003

50. Gweon H, Dodell-Feder D, Bedny M, Saxe R. Theory of Mind Performance in Children Correlates With Functional Specialization of a Brain Region for Thinking About Thoughts. *Child Development*. 2012;83(6):1853-1868. doi:10.1111/j.1467-8624.2012.01829.x

51. Wolff P. Observations on the early development of smiling. *Determinants of infant behavior*. 1963;2:113-138.

52. Huppert TJ, Diamond SG, Franceschini MA, Boas DA. HomER: a review of time-series analysis methods for near-infrared spectroscopy of the brain. *Appl Opt*. 2009;48(10):D280. doi:10.1364/AO.48.00D280

53. Fishburn FA, Ludlum RS, Vaidya CJ, Medvedev AV. Temporal Derivative Distribution Repair (TDDR): A motion correction method for fNIRS. *NeuroImage*. 2019;184:171-179. doi:10.1016/j.neuroimage.2018.09.025

54. Pinti P, Scholkmann F, Hamilton A, Burgess P, Tachtsidis I. Current Status and Issues Regarding Pre-processing of fNIRS Neuroimaging Data: An Investigation of Diverse Signal Filtering Methods Within a General Linear Model Framework. *Front Hum Neurosci*. 2019;12:505. doi:10.3389/fnhum.2018.00505

55. Zhang X, Noah JA, Hirsch J. Separation of the global and local components in functional near-infrared spectroscopy signals using principal component spatial filtering. *Neurophoton*. 2016;3(1):015004. doi:10.1117/1.NPh.3.1.015004

56. Zhang X, Noah JA, Dravida S, Hirsch J. Signal processing of functional NIRS data acquired during overt speaking. *Neurophoton*. 2017;4(04):1. doi:10.1117/1.NPh.4.4.041409

57. Noah JA, Zhang X, Dravida S, et al. Comparison of short-channel separation and spatial domain filtering for removal of non-neural components in functional near-infrared spectroscopy signals. *Neurophoton*. 2021;8(01). doi:10.1117/1.NPh.8.1.015004

58. Luke R, Larson E, Shader MJ, et al. Analysis methods for measuring passive auditory fNIRS responses generated by a block-design paradigm. *Neurophoton*. 2021;8(02). doi:10.1117/1.NPh.8.2.025008

59. Yücel MA, Lühmann AV, Scholkmann F, et al. Best practices for fNIRS publications. *Neurophoton*. 2021;8(01). doi:10.1117/1.NPh.8.1.012101

60. Aslin RN. Questioning the questions that have been asked about the infant brain using near-infrared spectroscopy. *Cognitive Neuropsychology*. 2012;29(1-2):7-33. doi:10.1080/02643294.2012.654773

61. Barr DJ, Levy R, Scheepers C, Tily HJ. Random effects structure for confirmatory hypothesis testing: Keep it maximal. *Journal of Memory and Language*. 2013;68(3):255-278. doi:10.1016/j.jml.2012.11.001

62. Whiteman AC, Santosa H, Chen DF, Perlman S, Huppert T. Investigation of the sensitivity of functional near-infrared spectroscopy brain imaging to anatomical variations in 5- to 11-year-old children. *Neurophoton*. 2017;5(01):1. doi:10.1117/1.NPh.5.1.011009

63. Emberson LL, Richards JE, Aslin RN. Top-down modulation in the infant brain: Learning-induced expectations rapidly affect the sensory cortex at 6 months. *Proc Natl Acad Sci USA*. 2015;112(31):9585-9590. doi:10.1073/pnas.1510343112

64. Lloyd-Fox S, Blasi A, Volein A, Everdell N, Elwell CE, Johnson MH. Social Perception in Infancy: A Near Infrared Spectroscopy Study. *Child Development*. 2009;80(4):986-999. doi:10.1111/j.1467-8624.2009.01312.x

65. Ozernov-Palchik O, O'Brien AM, Lee EJ, et al. Precision fMRI reveals that the language network exhibits adult-like left-hemispheric lateralization by 4 years of age. Published online May 15, 2024. doi:10.1101/2024.05.15.594172

66. Dehaene-Lambertz G, Hertz-Pannier L, Dubois J, et al. Functional organization of perisylvian activation during presentation of sentences in preverbal infants. *Proc Natl Acad Sci USA*. 2006;103(38):14240-14245. doi:10.1073/pnas.0606302103

67. Szaflarski JP, Schmithorst VJ, Altaye M, et al. A longitudinal functional magnetic resonance imaging study of language development in children 5 to 11 years old. *Annals of Neurology*. 2006;59(5):796-807. doi:10.1002/ana.20817

68. Holland SK, Vannest J, Mecoli M, et al. Functional MRI of language lateralization during development in children. *International Journal of Audiology*. 2007;46(9):533-551. doi:10.1080/14992020701448994

69. Weiss-Croft LJ, Baldeweg T. Maturation of language networks in children: A systematic review of 22years of functional MRI. *NeuroImage*. 2015;123:269-281. doi:10.1016/j.neuroimage.2015.07.046

70. Friederici AD. The Brain Basis of Language Processing: From Structure to Function. *Physiological Reviews*. 2011;91(4):1357-1392. doi:10.1152/physrev.00006.2011

71. Vigneau M, Beaucousin V, Hervé PY, et al. What is right-hemisphere contribution to phonological, lexico-semantic, and sentence processing? *NeuroImage*. 2011;54(1):577-593. doi:10.1016/j.neuroimage.2010.07.036

72. Lynch JS, Van Den Broek P, Kremer KE, Kendeou P, White MJ, Lorch EP. The Development of Narrative Comprehension and Its Relation to Other Early Reading Skills. *Reading Psychology*. 2008;29(4):327-365. doi:10.1080/02702710802165416

73. Airenti G. Pragmatic Development. In: Cummings L, ed. *Research in Clinical Pragmatics*. Vol 11. Perspectives in Pragmatics, Philosophy & Psychology. Springer International Publishing; 2017:3-28. doi:10.1007/978-3-319-47489-2_1

74. Bunge SA, Crone EA. Neural correlates of the development of cognitive control. In: Rumsey JM, Ernst M, eds. *Neuroimaging in Developmental Clinical Neuroscience*. 1st ed. Cambridge University Press; 2009:22-37. doi:10.1017/CBO9780511757402.005

75. Bunge SA, Dudukovic NM, Thomason ME, Vaidya CJ, Gabrieli JDE. Immature Frontal Lobe Contributions to Cognitive Control in Children. *Neuron*. 2002;33(2):301-311. doi:10.1016/S0896-6273(01)00583-9

76.   Casey BJ, Trainor RJ, Orendi JL, et al. A Developmental Functional MRI Study of
Prefrontal Activation during Performance of a Go-No-Go Task. *Journal of Cognitive*
Neuroscience. 1997;9(6):835-847. doi:10.1162/jocn.1997.9.6.835

77.   Crone EA, Steinbeis N. Neural Perspectives on Cognitive Control Development during
Childhood and Adolescence. *Trends in Cognitive Sciences*. 2017;21(3):205-215.
doi:10.1016/j.tics.2017.01.003

78.   Rubia K, Smith AB, Taylor E, Brammer M. Linear age-correlated functional
development of right inferior fronto-striato-cerebellar networks during response inhibition and
anterior cingulate during error-related processes. *Human Brain Mapping*. 2007;28(11):1163-
1177. doi:10.1002/hbm.20347

79.   Mehnert J, Akhrif A, Telkemeyer S, et al. Developmental changes in brain activation
and functional connectivity during response inhibition in the early childhood brain. *Brain and*
Development. 2013;35(10):894-904. doi:10.1016/j.braindev.2012.11.006

80.   Moriguchi Y, Hiraki K. Longitudinal development of prefrontal function during early
childhood.   *Developmental   Cognitive   Neuroscience*.   2011;1(2):153-162.
doi:10.1016/j.dcn.2010.12.004

81.   Steinbeis N, Bernhardt BC, Singer T. Impulse Control and Underlying Functions of the
Left DLPFC Mediate Age-Related and Age-Independent Individual Differences in Strategic
Social Behavior. *Neuron*. 2012;73(5):1040-1051. doi:10.1016/j.neuron.2011.12.027

82.   Poldrack R. Can cognitive processes be inferred from neuroimaging data? *Trends in*
Cognitive Sciences. 2006;10(2):59-63. doi:10.1016/j.tics.2005.12.004

83.   Kriegeskorte N, Simmons WK, Bellgowan PSF, Baker CI. Circular analysis in systems
neuroscience: the dangers of double dipping. *Nat Neurosci*. 2009;12(5):535-540.
doi:10.1038/nn.2303

84.   Lloyd-Fox S, Blasi A, Elwell CE. Illuminating the developing brain: The past, present
and future of functional near infrared spectroscopy. *Neuroscience & Biobehavioral Reviews*.
2010;34(3):269-284. doi:10.1016/j.neubiorev.2009.07.008

85.   Brigadoi S, Cooper RJ. How short is short? Optimum source–detector distance for
short-separation channels in functional near-infrared spectroscopy. *Neurophoton*.
2015;2(2):025005. doi:10.1117/1.NPh.2.2.025005

86.   Gregg. Brain specificity of diffuse optical imaging: improvements from superficial
signal regression and tomography. *Front Neuroenerg*. Published online 2010.
doi:10.3389/fnene.2010.00014

87.   Skeide MA, Friederici AD. The ontogeny of the cortical language network. *Nat Rev*
Neurosci. 2016;17(5):323-332. doi:10.1038/nrn.2016.23

Biography

Haolun Luo is a PhD student at the Hong Kong Polytechnic University. He received his BS
degree in Hearing and Speech Rehabilitation from Sichuan University in 2022. His research
interests focus on using fNIRS to assess children's cognitive and language abilities, with the
goal of developing objective biomarkers for child development.

Tao Yu is an Assistant Professor at the Bio-X Institutes, Shanghai Jiao Tong University, and
a PI at Shanghai Mental Health Center. He received his PhD from Sichuan University and
completed postdoctoral research at Shanghai Jiao Tong University and King's College
London. His research includes work on precision medicine, chronic non-communicable
disease prevention, and cognitive neuroscience.

Qun Li received her PhD from The Chinese University of Hong Kong. She is recognized as a
high-level overseas talent in Sichuan Province. Her research focuses on language assessment,
child language acquisition, language and cognition in children with language disorders, and
speech rehabilitation for hearing-impaired children. She currently leads a Sichuan Provincial
International Science and Technology Innovation Cooperation project and collaborates with
Hong Kong's Child Assessment Service.

Li Sheng is a faculty member in the Department of Chinese and Bilingual Studies at the
Hong Kong Polytechnic University. Previously, she held faculty positions at the University
of Texas Austin and the University of Delaware. She directs the Language Learning and
Bilingualism Lab, conducting research on language development mechanisms and improving
quality of life for individuals with language disorders through developing diagnostic tools
and intervention programs.