Brain bases for navigating acoustic features

Alexander J. Billig[a], William Sedley[b], Phillip E. Gander[c,d], Sukhbinder Kumar[d], Meher Lad[b], Maria Chait[a], Yousef Mohammadi[e], Joel I. Berger[d], & Timothy D. Griffiths[d,e,f]

[a] UCL Ear Institute, University College London
332 Grays Inn Road, London, WC1X 8EE, United Kingdom

[b] Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University.
Framlington Place, Newcastle upon Tyne, NE2 4HH, United Kingdom

[c] Department of Radiology, The University of Iowa
Iowa City, IA, 52242, United States of America

[d] Department of Neurosurgery, The University of Iowa
Iowa City, IA, 52242, United States of America

[e] Department of Imaging Neuroscience, University College London
12 Queen Square, London, WC1N 3AR, United Kingdom

[f] Biosciences Institute, Faculty of Medical Sciences, Newcastle University.
Framlington Place, Newcastle upon Tyne, NE2 4HH, United Kingdom

[*] Correspondence concerning this article should be addressed to Alexander Billig, UCL Ear Institute, 332 Grays Inn Road, London WC1X 8EE, United Kingdom. E-mail: a.billig@ucl.ac.uk

Short running title: Navigating acoustic features
Number of **pages**: 34
Number of **figures**: 8
Number of **tables**: 7

Data availability statement: The data that support the findings of this study are openly available in BIDS format at OpenNeuro at https://openneuro.org/datasets/ds006211.

Conflict of interest disclosure: The authors declare no competing financial interests.

Acknowledgments: This work was supported by the Wellcome Trust, United Kingdom (WT106964MA), and Medical Research Council, United Kingdom (MR/T032553/1). The Wellcome Centre for Human Neuroimaging was supported by core funding from the Wellcome Trust (203147/Z/16/Z). We thank Oliver Josephs, Nadège Corbin, and Martina Callaghan for assistance with MRI sequence design, preprocessing, and denoising; Eleanor Maguire and Karl Friston for advice on experimental design; and the radiographers at the The Wellcome Centre for Human Neuroimaging (now UCL Department of Imaging Neuroscience).

Abstract

Whether physical navigation shares neural substrates with mental travel in other behaviourally relevant

domains is debated. With respect to sound, pure tone working memory in humans elicits hippocampal

as well as auditory cortical and inferior frontal activity, and rodent work suggests that hippocampal cells

that usually track an animal's physical location can also map to tone frequency when task-relevant. We

generated a sound dimension based on the density of random-frequency tones in a stack, resulting in a

percept ranging from low- ("beepy") to high-density ("noisy"). We established that unlike tone

frequency, which listeners automatically associate with vertical position, this density dimension elicited

no consistent spatial mapping. During functional magnetic resonance imaging, human participants held

in mind the density of a series of tone stacks and, after a short maintenance period, adjusted further stacks

to match the target ("navigation"). Density was represented most strongly in bilateral non-primary

auditory cortex, specifically bilateral planum polare. Encoding and maintenance activity in

hippocampus, inferior frontal gyrus, planum polare and posterior cingulate was positively associated

with subsequent navigation success. Bilateral inferior frontal gyrus and hippocampus were among

regions with elevated activity during adjustment, compared to a parity-judgment condition with closely

matched acoustics and motor demands. Bilateral orbitofrontal cortex was more active when navigation

was toward a target density than when participants adjusted density in a control condition with no

particular target. We find that self-initiated travel along a non-spatial auditory dimension engages a brain

system overlapping with that supporting physical navigation.

Keywords: Auditory memory, working memory, navigation, hippocampus, sound, auditory cognition

66   **Key Points**

•   Work in rodents suggests that navigation in physical space and the active analysis of sounds share
a neural substrate in the hippocampus, supporting the use of common computational
mechanisms.

•   We examined the human brain system for navigation through an acoustic environment to a
remembered target.

•   In addition to high-level auditory cortex we demonstrate involvement of the hippocampus along
with sites in frontal and cingulate cortex that also support physical navigation.

Introduction

Finding our way in the physical world is supported by computations involving hippocampus as well as entorhinal, parietal, cingulate, and prefrontal cortices (O'Keefe and Dostrovsky, 1971; Ekstrom et al., 2003; Hafting et al., 2005; Spiers, 2008; Doeller et al., 2010; Bicanski and Burgess, 2020; Basu et al., 2021). Recent theoretical and empirical work suggests that this neural machinery may operate over more general sensory or knowledge spaces (Tavares et al., 2015; Constantinescu et al., 2016; Aronov et al., 2017; Bao et al., 2019; Whittington et al., 2020). In the auditory world a principal organising dimension is frequency. Linguistic and musical communication often depend on generating and making sense of trajectories in frequency, which is mapped tonotopically at multiple levels of the auditory system from the cochlea to sensory cortex. When tone frequency is controlled by a rat to match a target and obtain a reward, neurons tuned to frequency are also found in hippocampus and entorhinal cortex (Aronov et al., 2017). Many of these neurons also represent individual or geometrically related spatial locations (place and grid cells, respectively) during physical navigation, suggestive of a flexible code for traversing different dimensions of experience. Such a correspondence may arise simply because frequency has a natural mapping to spatial location: higher frequencies are implicitly associated with higher points in physical space (Rusconi et al., 2006). It may also be that ensembles of sequentially firing hippocampal cells map progress toward a goal or reward, rather than encoding the particular dimension along which such progress is made (see Zutshi et al., 2025 for an example using tones).

Navigation involves not only adjusting one's position but also maintaining a representation of the destination. In the spatial domain, hippocampal activity and place coding are more pronounced for navigation toward a target than during aimless movement or foraging (Cornwell et al., 2008; Gulli et al., 2020). Maintaining spatial targets engages frontal sites including ventromedial prefrontal and orbitofrontal cortex (Spiers, 2008; Basu et al., 2021). Holding a tone frequency in mind over seconds engages auditory cortex, frontoparietal regions (particularly inferior frontal gyrus) and hippocampus (Zatorre et al., 1994; Gaab et al., 2003; Kumar et al., 2016; Uluç et al., 2018; Kumar et al., 2021; Deutsch et al., 2023). A similar network supports maintenance of more complex auditory objects, such as spectral ripples or tones defined by a combination of frequency and either location or presentation rate (Czoschke et al., 2021; Ahveninen et al., 2023; Pomper et al., 2023; Uluç et al., 2023). We designed a stimulus and task that allowed us to probe the neural substrates of what we define as non-spatial "navigation" toward a spectrotemporally complex target sound in humans.

We used successions of tone stacks that varied in the number of concurrently presented tones of

random frequency (Figure 1A). This gave rise to a percept varying from sparse ("beepy") to dense

("noisy"). Using objective and subjective behavioural measures we established that this acoustic

dimension does not consistently elicit a spatial association. We then had participants undergo functional

magnetic resonance imaging (fMRI) while performing tasks that separately tapped processes of working

memory and navigation in this auditory dimension. The main condition of interest involved holding in

mind a series of tone stacks of fixed density, then adjusting a new set of stacks to match the target. In

order to establish whether any coding for density was distinct from proximity to the target we used

different starting and target densities in different trials. In control conditions we removed one or both of

(a) the working memory component by having participants randomly adjust tone stacks with no

particular target, and (b) the navigation component by having participants respond as to whether each in

a series of spoken numbers was odd or even while listening passively to the tone stacks. As far as possible

we matched the acoustics and number of button presses across conditions.

Regardless of task, we expected the spectral density of sounds that were matched for overall

spectrum to be represented in higher-order auditory cortex on the superior temporal plane. Based on

previous imaging work with a range of acoustic stimuli we hypothesized that auditory cortex, inferior

frontal gyrus and hippocampus would be active when such sounds were maintained as targets. If

mechanisms of spatial navigation and auditory analysis overlap in humans as they do in rodents we

anticipated hippocampal activation and representation of the auditory space during adjustment, but not

for the control task. We also expected any such hippocampal engagement to be more pronounced when

auditory adjustment was toward a stored target than when it was aimless.

Materials and Methods

Stimuli

The primary experimental stimuli were 190-ms tone stacks (with 2-ms cosine on- and offset ramps)

interspersed with 10 ms of silence. Varying the number of concurrent pure tones in a stack gave rise to

a "density" percept described to participants as ranging from "beepy" to "noisy" (see Figure 1A). In the

main experiment each stack contained between 2 and 196 tones, this value taking one of 26

logarithmically spaced levels. In the pilot study, an upper density limit of 400 tones was used, and

density levels were determined by the staircase described in "Behavioural pilot experimental procedure".

Figure 1: Stimuli and experimental design. A) Four example tone stacks ranging in spectral density from sparse ("beepy", left) to dense ("noisy", right) based on the number of pure tone components. In the pilot behavioural experiment participants heard two such stacks and judged whether the second was beepier or noisier than the first. **B)** Example of an MA (memory adjust) trial in the fMRI experiment, showing target sound (starting at 0s), maintenance period (starting at 2s), and adjustment period (starting at 4s). Tone components are shown as short horizontal grey lines. Button presses (reflecting attempts to match the target density) are shown as 'd' (make denser) and 's' (make sparser) with black horizontal lines representing their durations. **C)** Structure of trials in each of the four conditions. Conditions with labels containing 'M' involved maintaining a target density in <u>m</u>emory (purple lines) and reporting its similarity to that at the end of a subsequent series of stacks; conditions with labels containing 'N' did <u>n</u>ot involve target maintenance - instead participants reported how beepy or noisy the sound at

the end of the series of stacks was. Conditions with labels containing 'A' (green lines) involved adjusting the
ongoing series of stacks (either to match the target density in the MA condition, or at will in the NA condition);
conditions with labels containing 'P' (blue lines) instead required participants to report the parity of each a series
of spoken digits (i.e. whether it was odd or even). Tone stacks and digits were present in all conditions, including
when task-irrelevant. **D**) Factorization of the design into four conditions based on the presence or absence of
memory and adjustment processes. Instx. = Instructions, MA = Memory Adjust, MP = Memory Parity, NA = No-
memory Adjust, NP = No-memory Parity.

In both the pilot and the main experiment, tone frequencies were randomly drawn from a logarithmic

distribution between 250 and 4000 Hz. Tone stacks were equated for root-mean-square intensity, which

gave a reliable loudness match during piloting. We also created recordings of spoken digits from 1 to 10

using a text-to-speech tool (notevibes.com; voices: Daniel, Gabriela; speed: 1.6). These were trimmed

to have duration below 400 ms and normalised for root mean square intensity.

Participants

Ten individuals (aged 21-40 years, median 26.5 years, 4 males, 6 females) with self-reported normal

hearing took part in the behavioural pilot experiment. A different group of 30 right-handed individuals

with no reported neurological or hearing impairment took part in the fMRI experiment. Data from 25 of

these (aged 19-40 years, median 24 years, 12 males, 13 females) were included in the analysis. Of the

remaining five participants, one did not complete the experiment, two performed more than two standard

deviations below the group mean in the adjustment task (see fMRI experiment procedure), and three

(including one of the poor performers) moved more than 3 mm (one voxel) in the scanner. Participants

were not selected on the basis of musical training in either experiment. The study was approved by the

University College London Research Ethics Committee. Informed consent was obtained from all

participants, who were compensated for their time and travel.

Behavioural pilot experimental procedure

The purposes of the behavioural pilot experiment were to establish (1) how well participants could

discriminate different levels of tone stack density and (2) whether reaction times were affected by the

layout of response keys, potentially reflecting an implicit mapping between density and physical space.

After familiarization with the density feature and instruction that avoided language relating to space,

participants heard pairs of 200-ms tone stacks separated by 1s of silence. Participants had to respond

whether the second stack was "beepier" or "noisier" than the first, as quickly and accurately as possible.

Different pairs of keys were mapped to "beepier" and "noisier" responses in eight blocks. In two blocks

the "down" arrow was used for a "beepier" response and the "up" arrow for a "noisier" response; in

another two blocks this was reversed; in two further blocks the "left" arrow key was assigned to "beepier"
and the "right" arrow to "noisier"; in a final two blocks this was reversed. One of the two stacks in each
trial (selected at random) took one of four logarithmically spaced reference density levels (3.5, 9.1, 25.2,
66.5 tone components/octave). The density difference between the two stacks in a trial was adaptively
tracked using a two-down, one-up staircase procedure converging on the participant's 70.7% correct
threshold (averaging over the final two of six reversals, step size 1/3 until second reversal then 1/6). Two
adaptive tracks, each featuring a particular reference density, were run in parallel in each of the eight
blocks, resulting in sixteen tracks. The assignment of reference density levels to button mappings was
counterbalanced over participants, as was the block order. Accuracy feedback was provided after each
response.

Online training for fMRI participants
To ensure that participants were proficient at the main experimental tasks before scanning, we provided
online training (presented in Gorilla with custom JavaScript code) during the preceding week.
Participants were familiarized with the density manipulation and then practised each element of each
task (see next section) with feedback, before performing several trials of the full experiment. Further
training trials were administered in person immediately prior to scanning.

fMRI experimental procedure
Participants took part in four experimental runs in the scanner, each lasting 8.5 minutes and comprising
28 trials. Each run contained seven trials of each of four conditions, which separately manipulated the
presence of memory and adjustment task components while balancing acoustics and button presses.
Figure 1C illustrates the trial structure for each condition, and videos of example trials are available at
in Supporting Information. In both the "Memory Adjust" (MA) and "Memory Parity" (MP) conditions,
participants held in mind the density of a target sound over a maintenance period. In the MA condition
participants then adjusted the density of a series of stacks (with starting density chosen randomly) to
match the target density (Figure 1B). At the end of the trial, they rated how close in density the final
sound was to the target. In the MP condition after hearing the target they performed an "odd/even" task
on a series of spoken digits. They then turned their attention to the end of an ongoing series of tone stacks
and rated its similarity to the initial target. The "No-Memory Adjust" (NA) and "No-Memory Parity"
(NP) conditions did not include a target. In the NA condition, after hearing a series of stacks of mixed
density, participants adjusted a new series of stacks at will. They then rated how "beepy" or "noisy" the

final adjusted stack was. The NP condition also began with a series of stacks of mixed density, was

followed by the "odd/even" task on spoken digits and ended with a rating of how "beepy" or "noisy" the

final stack of an ongoing series was. Trials began with two seconds of silence while instructions were

displayed on screen. These varied with condition, included a reminder of the available response options,

were shown in black text on a light grey background, and covered all three phases of the trial (see Table

1). Once the sounds started, the instructions for the relevant phase were highlighted in white.

Table 1. Participant instructions for each condition

Condition code	Condition description	Phase 1 instructions	Phase 2 instructions	Phase 3 instructions
MA	Memory, Adjust	Store	Adjust	Compare
MP	Memory, Parity	Store	Odd/Even	Compare
NA	Non-memory, Adjust	Ignore	Adjust	Beepy/Noisy
NP	Non-memory, Parity	Ignore	Odd/Even	Beepy/Noisy

The initial sound in all conditions was a series of ten 190-ms stacks, each followed by 10 ms of silence.

In the MA and MP conditions the density of these stacks was set to a fixed target value, the specific

frequencies varying from stack to stack over the resulting 2-s target. In the NA and NP conditions these

initial stacks instead alternated between the lowest- and highest-density values. This was to emphasise

that they formed a "dummy" sound that was to be ignored, while matching the average spectrum and

density across conditions. In all conditions, the 2-s sound was followed by 1-3 s of silence, sampled from

a rectangular distribution – in the MA and MP conditions this silence acted as a working memory

maintenance phase. Over the next 8 s two simultaneous sound streams were heard: (1) a series of 40

stacks and (2) between one and six spoken digits. The value of each digit was randomly selected between

1 and 10 and spoken by either a male or female voice, also randomly selected. Digits could appear in

slots beginning at one-second intervals from second two of the eight-second phase, leaving the first and

final second free of digits such that only the tone stacks were audible. In the MA and NA (adjustment)

conditions participants ignored the digits and directed attention to the tone stacks throughout. They held

down one right-hand button to increase the density of the subsequent tone stacks, and another right-hand

button to reduce it - a new random set of frequency components was selected for each tone stack

regardless of whether the density changed. In the MA condition they were instructed to try to match the

target as closely as possible, using as much of the eight seconds as needed. In the NA condition they

adjusted the sounds at will and were asked to use a similar number of presses as in the MA trials.

In the MP and NP (parity) conditions, participants attended to spoken digits and responded via

one of two right-hand buttons as to whether each digit was odd or even (a similar task has served as an

appropriate baseline for hippocampal studies in the visual domain; Stark and Squire, 2001). To try to

match the number of button presses across conditions we set the number of digits presented in each trial

to that during the previous MA or NA trial plus the mean number of digits for which no response was

made across all previous MP and NP trials (or 0.5 when no such trials were available at the start of a

block), rounded up or down (at random) to the nearest integer. For the last second of this 8-s section of

MP and NP trials, participants redirected their attention to the final few tone stacks. To match acoustics

across conditions, in these MP and NP trials the densities of the ongoing tone stacks matched those

generated in the most recent MA or NA trial respectively. Identity, speaker, and timing of (ignored)

digits in the MA and NA trials were identical to those in the most recent MP or NP trials respectively

(or, when no previous trials were available at the start of a block, to those in a randomly generated

dummy trial). In the MA and MP conditions, the target density and starting density of the 8-s tone series

of stacks for each trial were randomly drawn from a predetermined set of 28 pairs that would generate a

range of expected trajectories spanning the density space with equal sampling of high- and low-density

sounds. In the NA and NP conditions the starting density of the 8-s series of tone stacks for each trial

was drawn from the same set of 28 pairs (with the target density disregarded).

The final phase of each trial required a judgment of either how different the density of the final

tone stack was to that of the target (MA and MP trials; response options: "Very similar", "Somewhat

similar", "Somewhat different", "Very different") or how beepy/noisy the final tone stack was (NA and

NP trials; response options: "Very beepy", "Somewhat beepy", "Somewhat noisy", "Very noisy").

Participants had three seconds to respond using one of four left-hand buttons. After a pause of between

0 and 2 s (sampled from a rectangular distribution) the next trial began. No feedback was provided. The

28 trials in a block were grouped into seven sets of four (not made explicit to participants), each set

containing one trial of each condition. Ordering within each set was pseudo-randomized, with the

constraint that the MA trial appeared earlier than the NA trial, and the MP trial appeared earlier than the

NP trial. This ensured that the 8-s series of tone stacks in the NA and NP trials could be matched to those

in the previous MA or NA trials, while preventing two trials of the same condition occurring in

succession.

The first two experimental blocks were separated by a break of up to two minutes. These were

followed by field mapping and anatomical scans (apart from for one participant whose anatomical scan

was obtained at the start of the session), lasting a total of around ten minutes. The final two experimental

blocks followed and were also separated by a break of up to two minutes. Sounds were presented through

an Edirol UA-4FX (Roland Corporation, Shuzuoka) USB soundcard and Ear-Tone Etymotics tube
earphones (Etymotic Research, Inc.) with foam tips. Levels were tested and adjusted at the start of the
session to ensure that both the tone stacks and spoken digits were audible over the scanner noise.
Instructions were displayed via an Epson EB-L1100U projector and viewed with a head-coil-mounted
mirror. The experiment was controlled using Psychtoolbox and custom MATLAB code.

MRI data acquisition

Data were collected using a 3-Tesla Siemens MAGNETOM Prisma MR scanner (Siemens Healthcare)
with a 32-channel receive head coil at the Wellcome Centre for Human Neuroimaging (London, UK).
Whole-brain T2*-weighted functional images were acquired using 3D echo-planar imaging with field of
view 192 x 192 x 144 mm, voxel size 3 mm isotropic, 48 transverse slices, 3D acceleration factor 2,
repetition time (TR) 44 ms, volume acquisition time 1056 ms, anterior-to-posterior phase encoding, and
bandwidth 2298 Hz/Px. The accelerated functional images were reconstructed using a motion-robust
higher-order sensitivity encoding approach (https://github.com/fil-physics/gadgetron-
matlab/tree/master/FIL-recon). Respiratory signals were obtained with a breathing belt. Field maps
(double-echo gradient echo sequence with field of view 192 x 192 x 144 mm, voxel size 3 x 3 x 2 mm,
64 transverse slices, TR 1020 ms, short echo time (TE) 10.00 ms, long TE 12.46 ms, flip angle 90°) and
a whole-brain T1-weighted anatomical image (MPRAGE, field of view 256 x 256 x 176 mm, voxel size
1 mm isotropic, 176 transverse slices, GRAPPA acceleration factor 2, TR = 2530 ms, TE = 3.34 ms)
were acquired during the same scanning session.

MRI preprocessing

MRI data were processed in SPM12 (Wellcome Centre for Human Neuroimaging). Each participant's
functional images were unwarped using their field map and realigned to the first volume. Functional and
anatomical images were coregistered to the mean functional image. For the univariate analyses, the
coregistered images were normalized to the SPM12 template (avg305T1) in MNI space and the
functional images were spatially smoothed using a Gaussian kernel with 6 mm full-width at half-
maximum.

Behavioural data analysis

For the behavioural experiment, participants' just-noticeable-differences (JNDs) were obtained as a
function of density. Reaction times were log-transformed and analysed as a function of button mapping

using a repeated-measures 4-way ANOVA. For the fMRI experiment, navigation success was scored

using the absolute number of steps in density space between the final adjusted sound and the target.

Parity judgments were scored by dividing the number of matches by the larger of two values: the number

of digits presented or the number of responses made. When determining matches, the series of responses

in a trial was allowed to be dislocated by an extra or missing response. Accuracy of responses to the final

question in each trial was assessed by calculating the Spearman's rank correlation of the responses (1-4)

with the true values (closeness in density for MA and MP conditions; actual density for NA and NP

conditions), after replacing missing responses with a value of 2.5 (the midpoint of both rating scales).

Proportions were compared following arcsine square root transformation, and correlation coefficients

following the Fisher *z*-transformation.

Univariate fMRI analyses

In a first level analysis a General Linear Model (GLM) was created for each participant's fMRI time

series data. Regressors of no interest were included for motion and respiration. Task phases of interest

were modelled as boxcar functions and convolved with the canonical haemodynamic response function.

A high pass filter of 1/230 Hz (corresponding to twice the longest interval between trials of the same

condition) was applied to remove low frequency noise, and SPM12's "FAST" option (appropriate for

rapid imaging sequences; Corbin et al., 2018) was used to model temporal correlations. To distinguish

activity relating to the different task components we first modelled the whole trial period and tested for

main effects of memory [contrast: (MA+MP)-(NA+NP)] and adjustment [contrast: (MA+NA)-

(MP+NP)], and for their interaction [contrast: (MA-MP)-(NA-NP)]. In a control analysis we modelled

these effects only during the portion of the MA trials during which adjustment responses were made (and

matching portions of trials in other conditions). Effects of density (all trials), distance from target (MA

trials), time in trial (MA trials), and performance in adjustment (MA trials) and parity judgment (MP

trials) tasks were examined by entering centered and orthogonalized linear parametric modulators into

the GLM over the relevant time intervals.

For each comparison of interest, participants' contrast images were entered into a second level

analysis, with one-sample *t*-tests conducted at the group level. Statistical parametric maps were obtained

and thresholded using a family-wise error- (FWE-) corrected *p*-value of .05, and a minimum cluster size

matching the "expected voxels per cluster" value in SPM for the corresponding height threshold.

Additional tests with small volume corrections were conducted in three a priori regions of interest,

namely bilateral auditory cortex (voxels assigned at least 20% probability of lying in Heschl's gyrus,

planum polare, planum temporale, or superior temporal gyrus in the Harvard-Oxford Cortical Structural

Atlas), bilateral inferior frontal gyrus pars opercularis and pars triangularis (voxels assigned at least 20%

probability of lying in one of these two regions in the Harvard-Oxford Cortical Structural Atlas), and

bilateral hippocampus (based on segmentation of the average T1w image across participants using the

Automatic Segmentation of Hippocampal Subfields (ASHS) software package, Yushkevich et al., 2014).

Multivariate fMRI analyses

The unsmoothed, unnormalized functional images were entered into two further GLMs. One contained

regressors for lower- versus higher-density sounds (splitting the 26 density levels into two groups) and

used trials of all conditions. The other contained regressors for navigation in the direction of decreasing

versus increasing density and used MA and NA trials. In both models these binary labels were applied

to each stack during the adjustment period and could therefore change within a trial. In both GLMs the

same regressors of no interest were entered as described in Univariate fMRI analyses above. Multivariate

analysis was performed using The Decoding Toolbox (Hebart et al., 2015) to obtain cross-validated

whitened Euclidean ("crossnobis") distances between neural representations corresponding to the two

category labels in a given model. Further analyses used the same approach to find distances between

representations of lower- versus higher-density targets, either during the encoding and maintenance

period only, or with inclusion of the adjustment period (in MA and MP conditions). Neural

representations were beta weights scaled for each run separately using residuals from the GLM with the

"Ledoit-Wolf retain variances" option selected. A spherical volumetric searchlight with 4 mm radius was

used over the whole brain. Leave-one-run-out cross-validation was applied and the lower diagonal values

of the dissimilarity matrices averaged over cross-validation iterations. Each participant's resulting

distance image was normalised to the SPM12 template (avg305T1) in MNI space and spatially smoothed

using a Gaussian kernel with 6 mm full-width at half-maximum. These were entered into a second level

analysis, which proceeded as described in Univariate fMRI analyses.

Results

Tone stack density: discrimination and associations with physical space

In the initial behavioural experiment, just-noticeable differences for density increased with density [$t(9)$

$= 13.717$, $p < .001$, $d = 4.338$; Figure 2A]. For this reason, density values in the fMRI experiment were

logarithmically spaced to ensure steps along the density scale were similarly discriminable. Reaction

times when comparing the density of two tone stacks was not affected by the spatial mapping of response

keys [$F(3, 27) = 0.549$, $p = .653$, $\eta^2_p = .057$; Figure 2B]. When prompted, twelve of the participants in

the fMRI experiment reported forming a spatial association with the stimuli (nine associated beepier

sounds with "left", one with "right", and two with "up"). To confirm that the group neural results were

not driven by these spatial associations we also report the relevant analyses by excluding the

corresponding participants, obtaining qualitatively similar results.

Behavioural performance in the scanner

In the MA (memory adjustment) condition of the main experiment, participants finished closer to the

target than they started in 94% of trials on average (range: 82-100%; Figure 2C), with a mean final

distance of 16% of the density scale (range: 9-33%). Participants tended to undershoot the target, that is

to remain on the same side of the target on the density scale as they began; on average this occurred in

70% of trials (range: 32-86%; Supplemental Figure S1). Performance did not differ between participants

who reported forming spatial associations and those who did not [$t(23) = 0.503$, $p = .620$, $d = .100$].

Parity judgment performance did not differ between the memory and non-memory conditions [MP and

NP; $t(24) = 1.132$, $p = .269$, $d = 0.227$; Figure 2D], with an average score of 89% (range: 59-98%).

Accuracy in comparing the target density to the final density did not differ significantly between the MA

and MP conditions [$t(24) = 1.475$, $p = .139$, $d = 0.298$; Figure 2E], while accuracy in reporting the final

density was significantly greater in the NA than the NP condition [$t(24) = 2.441$, $p = .022$, $d = 0.584$].

Matching of motor activity across conditions

There was a significant effect of condition on the number of presses despite our attempt to eliminate this

potential confound [$F(3, 72) = 10.256$, $p < .001$, $\eta^2_p = .299$]. Although the majority of participants were

well matched (on average a difference of less than one press per trial between pairs of conditions of

interest, namely MA versus NA, and MP versus NP), eight participants had more than one extra press

on average in adjustment trials (MA, NA) compared to parity trials (MP, NP), and six participants had

more than one extra press on average in non-memory trials (NA, NP) compared to memory trials (MA,

MP). Eleven participants met both criteria. To confirm that the group neural results were not confounded

by activity relating to motor preparation or execution we also report the relevant analyses excluding the

corresponding participants, obtaining qualitatively similar results.

Figure 2: Behavioural results. A) Just-noticeable differences in density increase as a function of reference density; values are the number of tone components per octave. Grey lines: individual participants in the behavioural pilot experiment. Black line: group mean. **B)** Raincloud and box-and-whisker plots of reaction times for correct trials in the behavioural pilot experiment. Each plot corresponds to a different button mapping (the arrow key that was mapped to "beepier" responses is shown; the opposite arrow key was mapped to "noisier" responses). Distributions did not differ by condition. **C)** Percentage of MA trials in the main experiment in which participants (grey dots) adjusted the ongoing density to be closer to the target than before adjustment. Black line represents mean performance and dotted line represents 50% chance performance. **D)** Percentage accuracy of parity judgments, which did not differ significantly between MP (left) and NP (right) conditions. Joined grey dots represent individual participants. Joined black squares represent group mean performance and dotted line represents 50% chance performance. **E)** Correlation between actual and reported target-versus-final density

similarity, which did not differ significantly between MA (left) and MP (right) conditions. Joined grey dots
represent individual participants. Joined black squares represent group mean performance and dotted line
represents chance performance.

Neural responses to sound and representation of tone stack density

Sites that were more active during sound compared to silence were bilateral auditory cortices in Heschl's

gyrus, planum temporale, and posterior superior temporal gyrus, along with parietal, somatomotor,

frontal, parietal and cerebellar areas (see Table 2). This contrast did not reflect purely sensory processes

since cognitive and motor demands also differed when sound was presented compared to when it was

not. During the adjustment phase (or equivalent period in the parity judgment task), activity in a more

restricted set of brain regions, shown in Figure 3A and Supplemental Table S1, was parametrically

modulated by the density of the tone stacks. Activity was greater for the sparser (less noise-like) tone

stacks - particularly in bilateral planum polare but also in Heschl's gyrus and along the temporal

convexity. During the same phase, multivoxel patterns for sparser versus denser sounds were also most

distinct at these sites (Figure 3B, Table 3). After small volume correction, a significant representation of

density was also found in right inferior frontal gyrus (pars opercularis and triangularis; Supplemental

Table S2), but not hippocampus.

413
414
415 **Figure 3: Representation of tone stack density. A)** Sagittal slice showing negative parametric effect of density
416 in auditory cortex, i.e. a lower univariate response to progressively denser tone stacks. **B)** Sagittal slice showing
417 accuracy (transformed to *t*-score) of multivariate decoding of density in auditory cortex. All clusters significant at
418 *p* < .05 (familywise error-corrected, whole brain).
419
420

422 **Table 2. Clusters more active during sound presentation than silence**

Regions	Voxels	Peak MNI coordinates			Peak t-value
Right planum temporale, Heschl's gyrus, superior temporal gyrus (posterior)	407	56	-14	6	10.29
Left planum temporale, Heschl's gyrus, superior temporal gyrus (posterior)	333	-54	-28	10	9.65
Left supplementary motor area, superior frontal gyrus	131	-6	8	54	9.45
Left precentral gyrus, postcentral gyrus	174	-40	18	60	8.89
Right superior frontal gyrus, supplementary motor area	108	6	10	58	8.11
Left supramarginal gyrus (anterior)	72	-52	-32	50	7.79
Left precentral gyrus	16	-56	6	18	7.55
Right inferior frontal gyrus (pars opercularis)	14	52	10	10	6.85
Left precentral gyrus	7	-28	-6	62	6.49
Right cerebellum (lobe V)	5	20	-52	-24	6.19

Clusters significant at $p < .05$ (FWE-corrected), minimum cluster size: 5 voxels

424 **Table 3. Clusters with multivoxel representation of density**

Regions	Voxels	Peak MNI coordinates			Peak t-value
Right planum polare	63	54	6	-4	8.37
Right planum temporale	125	64	-20	10	7.80
Left planum polare	23	-50	-8	-2	7.71
Left Heschl's gyrus, planum temporale	100	-48	-20	10	7.04
Left central opercular cortex	11	-58	-18	16	6.74
Left superior temporal gyrus (posterior)	6	-52	-16	-4	6.61

Clusters significant at $p < .05$ (FWE-corrected), minimum cluster size: 5 voxels

Memory-related neural activity

Next we contrasted activity in the two conditions in which participants held a target density in mind (MA
and MP) versus those with no such target (NA and NP). We considered activity throughout the trial,
from the start of the instructions to the end of the final report (a period lasting an average of 16 s and
corresponding to all but the final column in Fig. 1C). This revealed memory-related activity in bilateral
anterior insula, as well as frontal pole, middle frontal gyrus, and cerebellum (Figure 4 middle, right;
Table 4). After small volume correction, memory-related activity was also found in bilateral inferior
frontal gyrus (pars opercularis and pars triangularis, Figure 4 (left), Supplemental Table S3), but not
hippocampus. Activity levels for each condition averaged across voxels in each significant cluster are
shown in Supplemental Figure 2. No regions exhibited a multivoxel representation of the target density
in the MA and MP trials, whether considering just the encoding and maintenance periods or the whole
trial. The ability to decode the density of adjustment sounds (previous section) but not target sounds may
be due to targets only being physically present for a quarter of the time (2 s versus 8 s per trial).

Figure 4: Activity associated with target memory. Axial slices showing clusters with greater activity for MA and MP conditions than NA and NP conditions. **Left**: Inferior frontal gyrus clusters significant at $p < .05$ (familywise error-corrected, small volume for bilateral inferior frontal gyrus). **Middle**: Anterior insular clusters significant at $p < .05$ (familywise error-corrected, whole brain). **Right**: Cerebellar clusters significant at $p < .05$ (familywise error-corrected, whole brain). Red-yellow gradient represents t-score. Blue indicates regions of interest for small volume corrections. IFG = Inferior frontal gyrus.

Table 4. Clusters more active during memory than non-memory conditions

Regions	Voxels	Peak MNI coordinates			Peak t-value
Right insular cortex	58	30	24	0	8.92
Right frontal pole	30	46	40	28	8.23
Left insular cortex	43	-32	20	-4	7.50
Left middle frontal gyrus	6	-44	2	54	6.90
Left cerebellum (lobe VI)	13	-30	-64	-28	6.76
Right cerebellum (crus I)	11	34	-62	-28	6.46

Clusters significant at $p < .05$ (FWE-corrected), minimum cluster size: 5 voxels

Adjustment-related neural activity

The contrast of adjustment conditions (MA + NA) minus non-adjustment conditions (MP + NP) revealed activity in our regions of interest of hippocampus and inferior frontal gyrus after small volume correction (Figure 5; Supplemental Table S4). Activity levels for each condition averaged across voxels in each significant cluster are shown in Supplemental Figure 2. The corresponding clusters remained significant when excluding participants who reported noticing a spatial mapping of density (Supplemental Table

S5) or for whom button presses were not well matched across conditions (Supplemental Table S6). We

also considered the fact that even when button presses were comparable across conditions, subjects could

choose to stop adjusting before the end of an adjustment trial whereas they needed to remain alert

throughout parity judgment trials, even if they made no further presses. It is therefore possible that some

differences were driven by task-negative (e.g. mind-wandering) activity toward the end of adjustment

trials. In an analysis including only the interval in a given MA or NA trial during which adjustments

actually occurred and using the same intervals in matching MP and NP trials, the more anterior of the

hippocampal clusters in Figure 5 was no longer significant (Supplemental Table S7). At the whole brain

level, additional frontal, parietal, insular, cingulate and cerebellar sites also showed adjustment-related

activity (Table 5, Figure 6).

Figure 5: Activity associated with density adjustment in regions of interest. Clusters showing greater activity
for MA and NA conditions than MP and NP conditions. **Left**: Sagittal slice with hippocampal clusters significant
at $p < .05$ (familywise error-corrected, small volume for bilateral hippocampus). **Middle**: Axial slice with
hippocampal clusters significant at $p < .05$ (familywise error-corrected, small volume for bilateral hippocampus).
Right: Axial slice with inferior frontal gyrus clusters significant at $p < .05$ (familywise error-corrected, small
volume for bilateral inferior frontal gyrus). Red-yellow gradient represents t-score. Blue indicates regions of
interest for small volume corrections. IFG = Inferior frontal gyrus.

483
484
485
486

Table 5. Clusters more active during adjustment than parity conditions

Regions	Voxels	Peak MNI coordinates			Peak t-value
Right frontal pole	471	44	42	6	13.32
Left cerebellum (crus I, crus II, lobe VI)	945	-28	-72	-28	12.39
Right supramarginal gyrus (posterior, anterior)	618	56	-40	48	10.45
Right precentral gyrus, inferior frontal gyrus (pars opercularis/triangularis)	201	54	10	14	9.02
Left supramarginal gyrus (anterior), angular gyrus	315	-58	-34	44	8.66
Left insular cortex	67	-40	-4	6	8.58
Right frontal pole	59	40	48	22	8.37
Right insular cortex	65	40	0	4	8.34
Left frontal pole	264	-42	52	4	8.27
Right cerebellum (crus II)	280	38	-70	-42	8.07
Right inferior temporal gyrus (posterior)	11	56	-24	-22	7.85
Left frontal pole	10	-26	44	-14	7.32
Left lateral occipital cortex (superior)	54	-44	-64	42	7.25
Right supplementary motor area	16	4	26	42	7.03
Right cingulate gyrus (posterior)	45	6	-34	44	7.02
Right frontal pole	11	16	60	-6	6.82
Right central opercular cortex	7	46	-2	8	6.69
Right superior frontal gyrus	9	22	12	64	6.61
Left middle frontal gyrus	17	-36	30	44	6.48
Left insular cortex	6	-38	-16	-2	6.45
Right middle frontal gyrus	6	44	26	44	6.43
Left frontal pole	7	-42	42	26	6.38

Clusters significant at $p < .05$ (FWE-corrected), minimum cluster size: 6 voxels

488
489
Figure 6: Activity associated with density adjustment in other regions. Clusters showing greater activity for
MA and NA conditions than MP and NP conditions significant at $p < .05$ (familywise error-corrected, whole brain).
Left: Sagittal slice with clusters in posterior cingulate cortex and supplementary motor area. **Right**: Sagittal slice
with clusters in frontal pole, insula, and cerebellum. Red-yellow gradient represents t-score.

Navigation toward a target

We next looked for sites that showed an interaction between memory and adjustment, i.e. for which

adjustment-related activity was greatest when there was a target to be navigated toward [(MA-

MP)>(NA-NP)]. No regions survived a family-wise error correction at $p < .05$, however at a more liberal

threshold ($p < .001$ uncorrected), bilateral orbitofrontal cortex was more active for navigation toward a

target than for aimless navigation (Figure 7A; Supplemental Table S8). Activity levels for each condition

averaged across voxels in each significant cluster are shown in Supplemental Figure 2. This result was

qualitatively unaffected by the exclusion of subjects reporting spatial associations (Supplemental Table

S9). There was no interaction effect in any of our regions of interest.

505
Figure 7: Adjustment activity relating to target presence and direction. A) Axial slice showing interaction
between memory and adjustment in orbitofrontal cortex. Clusters significant at $p < .001$ (uncorrected). **B)** Coronal
slice showing accuracy (transformed to t-score) of multivariate decoding of navigation direction in left postcentral
gyrus. Cluster significant at $p < .05$ (familywise error-corrected, whole brain).

Direction and distance from target

The only areas encoding adjustment direction in MA and NA trials were left postcentral gyrus (Figure

7B) and right intracalcarine sulcus (Supplemental Table S10). The former may relate to the specific

fingers used to adjust sounds to become sparser or denser. To investigate correlates of distance from

target, we had to account for its negative correlation with time in trial. We assessed the unique

contribution of distance from target in an analysis in which elapsed time during the adjustment period
was entered as an initial regressor, and distance from target as a second orthogonal regressor. This
analysis included only the interval in each MA or NA trial during which adjustments actually occurred,
along with matching intervals in the corresponding MP and NP trials. No clusters showed effects of
distance at the corrected whole brain level, nor did our regions of interest after small volume correction.
There were significant clusters at the $p < .001$ uncorrected level in frontal and parietal cortex (Table 6),
with the largest cluster in left postcentral gyrus likely reflecting a greater likelihood of button pressing
when participants were more distant from the target; this peak was close to that shown in Figure 7B at
which adjustment direction was encoded. Effects of elapsed time can be found in Supplemental Tables
S11 and S12.

Table 6. Clusters with adjustment activity modulated by distance from target (uncorrected)

Modulation direction	Regions	Voxels	Peak MNI coordinates			Peak *t*-value
Positive	Left postcentral gyrus	251	-50	-22	50	4.64
Negative	Right supramarginal gyrus	69	60	-32	46	4.38
	Right frontal pole	45	46	48	8	4.29
	Right postcentral gyrus, precentral gyrus	48	40	-26	60	4.18

Clusters significant at $p < .001$ (uncorrected), minimum cluster size: 31 voxels

Navigation performance as a function of encoding and maintenance activity

Navigation to an auditory target requires that the target is first adequately encoded. In a final
analysis we looked for brain areas in which encoding and maintenance activity was parametrically
associated with performance in the MA trials (i.e. negatively related to the final distance from the target;
see Table 7, Figure 8). Two peaks in right auditory cortex were well aligned with those from which
density could be decoded (Figure 8 bottom left, compare Figure 3B). Other sites with activity associated
with subsequent performance were posterior cingulate gyrus (Figure 8 bottom right) and, after small-
volume corrections, hippocampus and inferior frontal gyrus (Figure 8 top, Supplemental Table S13).
These effects were specific to the adjustment task: none of these clusters' encoding/maintenance activity
was modulated by performance in the parity task in the MP trials.

545 **Table 7. Clusters with encoding/maintenance activity positively modulated by adjustment**
546 **performance**

Regions	Voxels	Peak MNI coordinates			Peak t-value
Right middle frontal gyrus	17	-32	12	60	9.26
Right Heschl's gyrus	43	54	-14	4	7.32
Right planum temporale	12	62	-26	12	7.12
Left cingulate gyrus (posterior)	16	-2	-32	30	7.03
Right planum polare	5	52	-2	-6	7.02
Right cingulate gyrus (posterior)	19	2	-34	30	6.96
Left occipital pole	7	-18	-100	10	6.55

Clusters significant at *p* < .05 (FWE-corrected), minimum cluster size: 5 voxels

549
Figure 8: Encoding and maintenance activity positively associated with navigation performance. Top left:
Sagittal slice with hippocampal cluster significant at *p* < .05 (familywise error-corrected, small volume for
bilateral hippocampus). **Top right:** Sagittal slice with inferior frontal gyrus clusters significant at *p* < .05
(familywise error-corrected, small volume for bilateral inferior frontal gyrus). **Bottom left:** Sagittal slice with
clusters in auditory cortex (familywise error-corrected, whole brain). **Bottom right:** Sagittal slice with cluster in
posterior cingulate cortex (familywise error-corrected, whole brain). Red-yellow gradient represents *t*-score. Blue
indicates regions of interest for small volume corrections. IFG = Inferior frontal gyrus.

Discussion

We studied the neural basis for perceiving, storing, and "navigating" toward an auditory target, defined
by the density of its acoustic spectrum, a parameter that participants did not consistently map to any
physical spatial dimension. It was represented most strongly in non-primary auditory cortex, and
adjusting it engaged a broad network including inferior frontal gyrus and hippocampus. Orbitofrontal
cortex was differentially activated when adjustments were made to match a target sound, compared to
aimless navigation. The level of activity in auditory cortex, inferior frontal gyrus, hippocampus, and
posterior cingulate during target encoding and maintenance was positively associated with navigation
performance.

We refer to self-initiated adjustment in this auditory density space as "navigation" but
acknowledge this stretches the standard definition of the term. Our study was partly motivated by
suggestions that hippocampus and neighbouring entorhinal cortex, known to be involved in mapping
physical space, can also represent task-relevant sensory spaces (Killian et al., 2012; Aronov et al., 2017;
Bao et al., 2019). The adjustment-related activity we observed in hippocampus and elsewhere was not
driven by "low-level" stimulus features (which were matched across conditions), nor by motor demands
or direct spatial associations (which control analyses accounted for). An anterior hippocampal cluster
dominated by activity after adjustments finished may reflect greater mind-wandering toward the end of
these trials (Zeidman et al., 2016). Posterior hippocampus showed more robust activation; it is involved
in retrieving sounds from working memory (Kumar et al., 2016) and in spatial navigation (Kühn and
Gallinat, 2014) - for example this region shows experience-dependent structural changes in taxi drivers
(Maguire et al., 2000). Previous work also demonstrated structural changes in hippocampus (although
more anteriorly) associated with a form of navigation through an auditory space in highly trained piano
tuners (Teki et al., 2012). Our study supports hippocampal activation during acoustic navigation in
novice participants.

Although the above suggests parallels in the neural basis for spatial and auditory navigation,
hippocampal activity patterns did not distinguish sparser from denser sounds. Aronov et al. (2017) found
individual hippocampal neurons that tiled frequency space when relevant to a rat engaged in a related
adjustment task. We may not have had a strong enough hippocampal signal for any spectral density
coding to be identified with the low temporal resolution of fMRI; other techniques such as single neuron
recordings from patients may prove fruitful. An alternative explanation is that hippocampus only

represents dimensions readily mapped to physical space, such as frequency. Rusconi et al. (2006) showed
that mapping sounds that were higher or lower in frequency or pitch to keys positioned higher or lower
on a computer keyboard, respectively, led to quicker responses than under the reverse mapping - even
when the task was unrelated to frequency. Under this explanation, spatially tuned machinery in rat
hippocampus would have been readily co-opted by space-associated sounds in Aronov et al. (2017). We
found no such consistent "dimensional overlap" for our stimulus, although a subset of participants
reported spatial associations. Another possibility is that neurons in Aronov et al. (2017) were tracking
progress toward a goal. Frequencies were always presented in the same order, although their rate of
change varied. The observed tuning could in fact reflect firing of hippocampal "time cells" that adapt
their temporal tuning scale in accordance with task demands (Macdonald et al., 2011, Shimbo et al.,
2021). This argument is reinforced by recent findings that apparent frequency-specific firing in
hippocampus or entorhinal cortex is absent in trials in which animals fail to stop at the target sound
(Nguyen et al., 2024; Zutshi et al., 2025). We deliberately chose trial-specific target and start densities,
such that participants' trajectories varied over trials, with a view to identifying tuning specific to the
acoustic feature. Density tuning was most robust in planum polare and also present in planum temporale
and anterior parts of Heschl's gyrus. It was driven by stronger responses to sparser sounds (chords with
individually identifiable frequency components) than denser (more noise-like) ones. This difference
might be due in part to greater adaptation within frequency channels for denser tone stacks. Planum
polare is also known to reflect higher-order acoustic features, such as pitch change (Warren and Griffiths,
2003) and spectrotemporal correlation (Overath et al., 2008), and to respond to music over scrambled
musical sounds (Angulo-Perkins and Concha, 2019).

Whereas previous work found changes in activity relating to working memory for sound in
auditory cortex (e.g. Gaab et al., 2003; Linke et al., 2015; Kumar et al., 2016; Czoschke et al., 2021),
our contrast comparing memory to non-memory conditions did not. Our silent maintenance period was
much shorter (1-3 s) than in other fMRI studies and not designed to be distinguished from the remainder
of the trial. Instead, our analysis encompassed the whole trial, which was dominated by periods of
auditory stimulation. Any memory-related activity in auditory cortex may have been swamped by
responses to the input, present in both memory and no-memory conditions. We did however find
memory-related activity in anterior insula, cerebellum and frontal sites including inferior frontal gyrus,
consistent with previous work (Zatorre et al., 1994; Kurth et al., 2010; Kumar et al., 2016; King et al.,
2019; Brissenden et al., 2021; Llorens et al., 2023).

The role of hippocampus during working memory maintenance has been hard to pin down
(Axmacher et al., 2007; Jenseon and Squire, 2011; Stretton et al., 2012). Recent studies have described
both elevated firing during working memory maintenance in hippocampal cells selective for particular
concepts (Daume et al., 2024) and reduced broadband high frequency power at hippocampal sites at
which spectrotemporal ripples held in working memory can nonetheless be decoded (Uluç et al., 2023).
It has been argued that not only persistent firing (Fuster and Alexander, 1971) but also "activity-silent"
mechanisms involving changes in synaptic weights (Mongillo et al., 2008; Miller et al., 2018) can
support working memory maintenance. Holding a target sound in mind may therefore not require
increases in the ensemble activity observed indirectly in the BOLD signal. We did however find a link
between the level of posterior hippocampal activity and subsequent success in the auditory navigation
task. This contrasts with the finding of Kumar et al. (2016) where participants with better tone working
memory showed a reduced hippocampal signal during maintenance (although activity was elevated
above baseline at the group level). It is possible that our behavioural effect was dominated by activity
during target presentation, rather than in the subsequent short silent maintenance period.

This link between auditory navigation success and encoding/maintenance activity, together with
the navigation-related activity, add to growing evidence for a role of the hippocampus in auditory
cognition (Billig et al., 2022), disruption of which may mediate links between hearing loss and dementia
(Griffiths et al., 2020). A highly connected structure, in our task the hippocampus likely formed part of
a functional network with areas including non-primary auditory cortex where spectral density was
encoded. The posterior cingulate was also more active during encoding and maintenance for trials in
which participants were subsequently more successful in reaching the target. Changes in posterior
cingulate structure and function have been associated with impaired working memory performance in
healthy ageing, schizophrenia, and attention deficit hyperactivity disorder (Leech and Sharp, 2014). In
our task this region was also more active during adjustment than parity-judgment trials. Reduced
connectivity between posterior cingulate and entorhinal cortex is associated with impaired navigation,
and six-fold rotational symmetry in the blood-oxygen-level-dependent (BOLD) response has been
detected there during navigation in virtual and abstract spaces, suggestive of the presence of grid cell
populations (Doeller et al., 2010; Constantinescu et al., 2016). Regions of medial frontal cortex appear
to be essential when rats (Basu et al., 2021) or humans (Ciaramelli, 2008; Spiers, 2008) maintain a goal
in physical space. We found clusters in bilateral orbitofrontal cortex that may serve the same purpose
during auditory navigation, showing a greater memory effect during the adjustment than the non-

adjustment task. Effective connectivity has been demonstrated in humans between auditory cortex and

hippocampus on the one hand and orbitofrontal cortex on the other (Rolls et al., 2023). It is another site

at which grid-like navigation signals have been detected in the BOLD signal (Constantinescu et al.,

2016).

Contrary to one of our hypotheses, the presence of an auditory target did not modulate the amount

of adjustment-related activity in hippocampus. This differs from findings from work comparing

navigation toward a spatial goal with aimless movement or foraging (Cornwell et al., 2008; Gulli et al.,

2019). It is possible that participants in our study set themselves an internal goal in trials in which they

were asked to make random adjustments. A fuller understanding of the network supporting auditory

navigation - both holding targets in mind and making adjustments in auditory space to reach them - will

require experiments teasing apart the contributions of hippocampus, orbitofrontal cortex and other

regions in which we observed related activity, such as cerebellum, insula, and multiple frontoparietal

sites (Tables 4 and 5), some of which may be domain general (Nozari and Martin, 2024). Directly

comparing spatial and auditory navigation correlates in the same participants would also be illuminating.

Distinguishing activity relating to adjustment and working memory processes required a complex

design. Rather than comparing conditions involving these processes to a silent baseline, we used a parity-

judgment task as a control that minimized hippocampal activity while matching motor activity and

sensory input. It required participants to redirect attention within a trial between tone stacks and digits.

This had no significant impact on working memory performance (participants stored the target density

similarly well in MA and MP trials) but the "adjustment" effects reported should be interpreted with the

nature of the baseline task in mind. We also note that some regions, including parts of hippocampus,

showed reduced activity in all conditions compared to the inter-trial rest periods. Despite this, the

factorial design allowed us to identify condition-specific correlates of auditory adjustment and working

memory that existed alongside any general reduction of task-unrelated activity.

673   **Conclusion**

We developed an acoustic stimulus defined by its spectral density, a feature represented in non-

primary auditory cortex and not consistently associated with any spatial dimension. Navigation through

this density space engaged brain regions including inferior frontal gyrus and hippocampus, with

orbitofrontal cortex activated more when navigation was toward a target. Encoding and maintenance in

areas including hippocampus, inferior frontal gyrus, auditory cortex and cingulate cortex was positively

associated with subsequent successful navigation. Overall, we find that self-initiated travel in humans

along a pure auditory dimension engages a brain system overlapping that previously found to support

spatial navigation.

References

Ahveninen J, Uluç I, Raij T, Nummenmaa A, Mamashli F (2023) Spectrotemporal content of human auditory working memory represented in functional connectivity patterns. Commun Biol 6:294. https://doi.org/10.1038/s42003-023-04675-8

Angulo-Perkins A, Concha L (2019) Discerning the functional networks behind processing of music and speech through human vocalizations. PLoS One 14:e0222796. https://doi.org/10.1371/journal.pone.0222796

Aronov D, Nevers R, Tank, DW (2017) Mapping of a non-spatial dimension by the hippocampal–entorhinal circuit. Nat 543:719–722. https://doi.org/10.1038/nature21692

Axmacher N, Mormann F, Fernandez G, Cohen MX, Elger CE, Fell J (2007) Sustained neural activity patterns during working memory in the human medial temporal lobe. J Neurosci 27:7807–7816. https://doi.org/10.1523/JNEUROSCI.0962-07.2007

Bao X, Gjorgieva E, Shanahan LK, Howard, JD, Kahnt T, Gottfried JA (2019) Grid-like neural representations support olfactory navigation of a two-dimensional odor space. Neuron 102:1066-1075. https://doi.org/10.1016/j.neuron.2019.03.034

Basu R, Gebauer R, Herfurth T, Kolb S, Golipour Z, Tchumatchenko T, Ito HT (2021) The orbitofrontal cortex maps future navigational goals. Nat 599:449–452. https://doi.org/10.1038/s41586-021-04042-9

Bicanski A, Burgess N (2020) Neuronal vector coding in spatial cognition. Nat Rev Neurosci 21:453–470. https://doi.org/10.1038/s41583-020-0336-9

Billig AJ, Lad M, Sedley W, Griffiths, TD (2022) The hearing hippocampus. Prog Neurobiol 218:102326. https://doi.org/10.1016/j.pneurobio.2022.102326

Brissenden JA, Tobyne SM, Halko MA, Somers DC (2021) Stimulus-specific visual working memory representations in human cerebellar lobule VIIb/VIIIa. J Neurosci 41: 1033–1045. https://doi.org/10.1523/JNEUROSCI.1253-20.2020

Ciaramelli, E (2008) The role of ventromedial prefrontal cortex in navigation: A case of impaired wayfinding and rehabilitation. Neuropsychologia 46:2099–2105. https://doi.org/10.1016/j.neuropsychologia.2007.11.029

Constantinescu, AO, O'Reilly JX, Behrens TEJ (2016) Organizing conceptual knowledge in humans with a gridlike code. Science 352:1464–1468. https://doi.org/10.1126/science.aaf0941

Corbin N, Todd N, Friston KJ, Callaghan MF (2018) Accurate modeling of temporal correlations in
rapidly sampled fMRI time series. Hum Brain Mapp 39:3884-3997.
https://doi.org/10.1002/hbm.24218
Cornwell BR, Johnson LL, Holroyd T, Carver FW, Grillon C (2008) Human hippocampal and
parahippocampal theta during goal-directed spatial navigation predicts performance on a virtual
Morris water maze. J Neurosci 28:5983-5990. https://doi.org/10.1523/JNEUROSCI.5001-07.2008
Czoschke S, Fischer C, Bahador T, Bledowski C, Kaiser J (2021) Decoding concurrent representations
of pitch and location in auditory working memory. J Neurosci 41:4658-4666.
https://doi.org/10.1523/JNEUROSCI.2999-20.2021
Daume J, Kamiński J, Salimpour Y, Gómez Palacio Schjetnan A, Anderson WS, Valiante TA,
Mamelak AN, Rutishauser U (2024) Persistent activity during working memory maintenance
predicts long-term memory formation in the human hippocampus. Neuron 112:1-12.
https://doi.org/10.1016/j.neuron.2024.09.013
Deutsch P, Czoschke S, Fischer C, Kaiser J, Bledowski C (2023) Decoding of working memory
contents in auditory cortex is not distractor-resistant. J Neurosci 43:3284-3293.
https://doi.org/10.1523/JNEUROSCI.1890-22.2023
Doeller CF, Barry C, Burgess N (2010) Evidence for grid cells in a human memory network. Nat 463:
657–661. https://doi.org/10.1038/nature08704
Ekstrom AD, Kahana MJ, Caplan JB, Fields TA, Isham EA, Newman EL, Fried I (2003) Cellular
networks underlying human spatial navigation. Nat 425:184–188.
https://doi.org/10.1038/nature01964
Fuster JM, Alexander GE (1971) Neuron activity relating to short-term memory. Science 173:652-654.
https://doi.org/10.1126/science.173.3997.652
Gaab N, Gaser C, Zaehle T, Jancke L, Schlaug G (2003) Functional anatomy of pitch memory - an
fMRI study with sparse temporal sampling. Neuroimage 19:1417-1426.
https://doi.org/10.1016/S1053-8119(03)00224-6
Griffiths TD, Lad M, Kumar S, Holmes E, McMurray B, Maguire EA, Billig AJ, Sedley W (2020)
How can hearing loss cause dementia? Neuron, 108:401–412.
https://doi.org/10.1016/j.neuron.2020.08.003

Gulli RA, Duong LR, Corrigan BW, Doucet G, Williams S, Fusi S, Martinez-Trujillo JC (2020) Context-dependent representations of objects and space in the primate hippocampus during virtual navigation. Nat Neurosci 23:103-112. https://doi.org/10.1038/s41593-019-0548-3

Hafting T, Fyhn M, Molden S, Moser M-B, Moser EI (2005) Microstructure of a spatial map in the entorhinal cortex. Nat 436:801–806. https://doi.org/10.1038/nature03721

Hebart MN, Görgen K, Haynes J-D (2015) The Decoding Toolbox (TDT): a versatile software package for multivariate analyses of functional imaging data. Front Neuroinform 8:88. https://doi.org/10.3389/fninf.2014.00088

Jeneson A, Squire, LR (2011) Working memory, long-term memory, and medial temporal lobe function. Learn Mem 19:15-25. https://doi.org/10.1101/lm.024018.111

Killian NJ, Jutras MJ, Buffalo EA (2012) A map of visual space in the primate entorhinal cortex. Nat 491:761-764. https://doi.org/10.1038/nature11587

King M, Hernandez-Castillo CR, Poldrack RA, Ivry RB, Diedrichsen J (2019) Functional boundaries in the human cerebellum revealed by a multi-domain task battery. Nat Neurosci 22:1371-1378. https://doi.org/10.1038/s41593-019-0436-x

Kumar S[*], Gander PE[*], Berger JI[*], Billig AJ, Nourski KV, Oya H, Kawasaki H, Howard MA, Griffiths, TD (2021) Oscillatory correlates of auditory working memory examined with human electrocorticography. Neuropsychologia 150:107691. https://doi.org/10.1016/j.neuropsychologia.2020.107691

Kumar S, Joseph S, Gander PE, Barascud N, Halpern AR, Griffiths TD (2016) A brain system for auditory working memory. J Neurosci 36:4492-4505. https://doi.org/10.1016/j.neuropsychologia.2020.107691

Kurth F, Zilles K, Fox PT, Laird AR, Eickhoff SB (2010) A link between the systems: Functional differentiation and integration within the human insula revealed by meta-analysis. Brain Struct Funct 214:519–534. https://doi.org/10.1007/s00429-010-0255-z

Kühn S, Gallinat J (2014) Segregating cognitive functions within hippocampal formation: A quantitative meta-analysis on spatial navigation and episodic memory. Hum Brain Mapp 35:1129-1142. https://doi.org/10.1002/hbm.22239

Leech R, Sharp DJ (2014) The role of the posterior cingulate cortex in cognition and disease. Brain 137: 12–32. https://doi.org/10.1093/brain/awt162

Linke AC, Cusack R (2015) Flexible information coding in human auditory cortex during perception,
imagery, and STM of complex sounds. J Cog Neurosci 27:1322-1333.
https://doi.org/10.1162/jocn_a_00780
Llorens A, Bellier L, Blenkmann, AO, Ivanovic J, Larsson PG, Lin JJ, Endestad T, Solbakk A-K,
Knight RT (2023) Decision and response monitoring during working memory are sequentially
represented in the human insula. iScience 26:107653. https://doi.org/10.1016/j.isci.2023.107653
MacDonald CJ, Lepage KQ, Eden UT, Eichenbaum, H (2011) Hippocampal "time cells" bridge the
gap in memory for discontiguous events. Neuron 71: 737–749.
https://doi.org/10.1016/j.neuron.2011.07.012
Maguire EA, Gadian DG, Johnsrude IS, Good CD, Ashburner J, Frackowiak RSJ, Frith CD (2000)
Navigation-related structural change in the hippocampi of taxi drivers. Proc Natl Acad Sci U S A
97: 4398–4403. https://doi.org/10.1073/pnas.070039597
Miller EK, Lundqvist M, Bastos AM (2018). Working memory 2.0. Neuron 100: 463–475.
https://doi.org/10.1016/j.neuron.2018.09.023
Mongillo G, Barak O, Tsodyks M (2008) Synaptic theory of working memory. Science 319: 1543–
1546. https://doi.org/10.1126/science.1150769
Nguyen D, Wang G, Wafa T, Fitzgerald T, Gu Y (2024) The medial entorhinal cortex encodes
multisensory spatial information. Cell Rep 43: 114813.
https://doi.org/10.1016/j.celrep.2024.114813
Nozari N, Martin, RC (2024) Is working memory domain-general or domain-specific? Trends Cogn
Sci, 28:1023–1036. https://doi.org/10.1016/j.tics.2024.06.006
O'Keefe J, Dostrovsky J (1971) The hippocampus as a spatial map. Preliminary evidence from unit
activity in the freely-moving rat. Brain Res 34:171–175.
https://doi.org/10.1016/0006-8993(71)90358-1
Overath T, Kumar S, Von Kriegstein K, Griffiths TD (2008) Encoding of spectral correlation over time
in auditory cortex. J Neurosci 28:13268–13273.
https://doi.org/10.1523/JNEUROSCI.4596-08.2008
Pomper U, Curetti LZ, Chait M (2023) Neural dynamics underlying successful auditory short-term
memory performance. Eur J Neurosci 58:3859-3878. https://doi.org/10.1111/ejn/16140

Rolls, ET, Deco G, Huang, C-C, Feng, J (2023) The human orbitofrontal cortex, vmPFC, and anterior cingulate cortex effective connectome: Emotion, memory, and action. Cereb Cortex 33:330–356. https://doi.org/10.1093/cercor/bhac070

Rusconi E, Kwan B, Giordano B, Umilta C, Butterworth B (2006) Spatial representation of pitch height: The SMARC effect. Cognition 99:113-129. https://doi.org/10.1016/j.cognition.2005.01.004

Shimbo A, Izawa E-I, Fujisawa S (2021) Scalable representation of time in the hippocampus. Sci Adv 7:eabd7013. https://doi.org/10.1126/sciadv.abd7013

Spiers HJ (2008) Keeping the goal in mind: Prefrontal contributions to spatial navigation. Neuropsychologia 46:2106–2108. https://doi.org/10.1016/j.neuropsychologia.2008.01.028

Stark CEL, Squire LR (2001) When zero is not zero: The problem of ambiguous baseline conditions in fMRI. Proc Natl Acad Sci U S A 98:12760-12766. https://doi.org/10.1073/pnas.221462998

Stretton J, Winston G, Sidhu M, Centeno M, Vollmar C, Bonelli S, Symms M, Koepp M, Duncan JS, Thompson PJ (2012) Neural correlates of working memory in Temporal Lobe Epilepsy—An fMRI study. Neuroimage 60:1696–1703. https://doi.org/10.1016/j.neuroimage.2012.01.126

Tavares RM, Mendelsohn A, Grossman Y, Williams CH, Shapiro M, Trope Y, Schiller D (2015) A map for social navigation in the human brain. Neuron 87: 231–243. https://doi.org/10.1016/j.neuron.2015.06.011

Teki S[*], Kumar S[*], von Kriegstein K, Stewart L, Lyness CR, Moore BCJ, Capleton B, Griffiths TD (2012) Navigating the auditory scene: An expert role for the hippocampus. J Neurosci 32: 12251–12257. https://doi.org/10.1523/JNEUROSCI.0082-12.2012

Uluç I, Peled N, Paulk AC, Bush A, Gumenyuk V, Kotlarz P, Lankinen K, Mamashli F, Matsuda N, Richardson MR, Stufflebeam SM, Cash SS, Ahveninen J (2023) Decoding auditory working memory content from intracranial high frequency activity in humans. bioRxiv 552073 https://doi.org/10.1101/2023.08.04.552073

Uluç I, Schmidt TT, Wu Y-H, Blankenburg F (2018) Content-specific codes of parametric auditory working memory in humans. Neuroimage 183:254-262. https://doi.org/10.1016/j.neuroimage.2018.08.024

Warren JD, Griffiths TD (2003) Distinct mechanisms for processing spatial sequences and pitch sequences in the human auditory brain. J Neurosci 23:5799–5804. https://doi.org/10.1523/JNEUROSCI.23-13-05799.2003

Whittington JCR, Muller TH, Mark S, Chen G, Barry C, Burgess N, Behrens TEJ (2020) The Tolman-
Eichenbaum Machine: unifying space and relational memory through generalization in the
hippocampal formation. Cell 183:1249-1263. https://doi.org/10.1016/j.cell.2020.10.024
Yushkevich PA, Pluta JB, Wang H, Xie L, Ding SL, Gertje EC, Mancuso L, Kliot D, Das SR, Wolk
DA (2014) Automated volumetry and regional thickness analysis of hippocampal subfields and
medial temporal cortical structures in mild cognitive impairment. Hum Brain Mapp 36:258-287.
https://doi.org/10.1002/hbm.22627
Zatorre R, Evans A, Meyer E (1994) Neural mechanisms underlying melodic perception and memory
for pitch. J Neurosci 14: 1908–1919. https://doi.org/10.1523/JNEUROSCI.14-04-01908.1994
Zeidman P, Maguire E (2016) Anterior hippocampus: the anatomy of perception, imagination and
episodic memory. Nat Rev Neurosci 17:173-182. https://doi.org/10.1038/nrn.2015.24
Zutshi I, Apostolelli A, Yang W, Zheng Z, Dohi T, Balzani E, Williams AH, Savin C, Buzsáki G
(2025) Hippocampal neuronal activity is aligned with action plans. Nature 649:153-161
https://doi.org/10.1038/s41586-024-08397-7