Long-range genomic loci stochastically assemble into combinatorial forms of chromosome skeleton

Jingyu Zhang[1,*], Siyuan Wang[2], Simon C Watkins[3], Jianhua Xing[1,4,5,*]

[1] Department of Computational and Systems Biology, University of Pittsburgh; Pittsburgh, PA 15232, USA.

[2] Department of Cell Biology, Yale School of Medicine; New Haven, CT 06510, USA.

[3] Department of Cell Biology, University of Pittsburgh; Pittsburgh, PA 15232, USA.

[4] UPMC-Hillman Cancer Center, University of Pittsburgh; Pittsburgh, PA, USA.

[5] Department of Physics and Astronomy, University of Pittsburgh; Pittsburgh, PA 15232, USA.

[*] To whom correspondence should be addressed. Email: xing1@pitt.edu, zhangjy4316@gmail.com

Abstract

One fundamental yet open question is how eukaryotic chromosomes fold into segregated territories, a process essential for gene transcription and cell fate. Through analyzing multiple sequencing- and imaging-based datasets, we identify long-range chromosomal skeleton loop structures that span over 100 Mb, extending beyond the reach of several existing DNA loop models. Further epigenetic and spatial density analyses point to assembly formation independent of major nuclear structures. Some long-range loops share a subset of genomic loci, which serve as nucleation centers and drive loop clustering. These complexes are highly stable, as shown by live-cell imaging with sequence-specific fluorescent labeling, and biophysical model analyses reveal a multivalent binding mechanism. Our findings suggest a redundant, distributed cluster mechanism that ensures robustness across cell types and against mutations, guiding both chromosome compaction and the formation of smaller-scale chromosomal structures.

Main Text

A central problem in structural biology and biological physics is how macromolecules fold into three-dimensional structures. An emerging "local-to-global" mechanism highlights the role of local cooperativity among peptide monomers in guiding the formation of a native protein structure from an expansive conformational space[1]. In contrast to typical polypeptides with $300 - 400$ amino acids, the normal human genome is composed of 46 long-chain polymers, ranging from ~ 40 to 250 Mb, with a total linear length extending up to two meters. Given that the 3D structures play a central role in regulating cell identity and gene expression[2], an intriguing question is how these linear DNA molecules compact into segregated chromatin territories within a cell nucleus, typically only about 10 μm in diameter.

Chromatin, unlike a polypeptide, is often considered to lack a stable, ordered three-dimensional structure due to its significantly greater length. When modeled as a linear polymer chain without

structural supports such as protein bridges or long non-coding RNA networks, theoretical analyses
predict that the average contact frequency (f) between two chromosomal fragments depends on
their genomic separation (l) and follows a power-law decay, $f \propto l^{\alpha}$. Previous Hi-C measurements
have confirmed this relationship, with an exponent $\alpha \sim -1$ within the $1 - 5$ Mb range (Fig. 1a,
Extended Data Fig. 1a-c). This result aligns with polymer models describing chromatin as a
crumpled polymer[3-5] or emphasizing the excluded volume effect[6].

However, deviations from the power-law behavior have been observed at both short and long
genomic distances. At genomic distances ~100 – 200 kb, the deviations (Fig. 1a, Extended Data
Fig. 1a-c) suggest the presence of stable chromatin loops, potentially formed through a loop-
extrusion mechanism[7]. Beyond five Mb, Hi-C data reveal further deviations, where the f-l curves
level off (Fig. 1a blue line, Extended Data Fig 1a&b). This phenomenon has been attributed to
polymer models incorporating random-sized loop formation between chromatin segments[8].
Additional factors, such as A/B compartment segregation and nuclear structure tethering, also
contribute to deviations from the crumpled polymer model at large genomic distances[4, 9].

In sum, previous studies suggest that chromosomes form functionally important local structures
such as topologically associated domains (TADs) and enhancer-promoter interactions, typically
spanning within ~ 1 Mb. Reports also exist on the existence of stable, long-range genomic
interactions (>10 Mb) in large genomes of species including human[4, 9-11], and mechanisms such as
liquid-liquid phase separation mediated by Polycomb and HP1 are known to facilitate some such
long-range contacts[12, 13]. However, there is a lack of systematic investigations on the existence of
sequence-specific, conserved, and stable locus-locus interactions >50 Mb and the physical
principles governing them. Here, we address this gap by characterizing spatial colocalization of
pairs of distant loci on chromosomes of several mammalian and non-mammalian species. We
demonstrate the conservation of their spatial proximity across diverse cell lines and elucidate the
physical mechanisms that likely enforce this stable association.

Hi-C data reveals the prevalence of long-range genomic locus colocalization beyond 50 Mb

We first analyzed Hi-C data for chromosome 2 (chr2) and chromosome 14 (chr14) in human
MCF10A cells[14]. While most genomic pairs followed the expected f-l curve, certain pairs (e.g., red
box in Fig. 1a & Extended Data Fig. 1a) showed significant deviations, with contact frequencies
comparable to the median value of 1-Mb pairs despite their much larger genomic separation.
Similar distal locus pairs with high contact frequencies were also observed in chr2 and chr14 of
human IMR90 cells[15] (Extended Data Fig. 1b). Extending this analysis to 11 additional human cell
lines revealed that such deviations are prevalent among genomic locus pairs separated by even
more than 50 Mb (Extended Data Fig. 2), suggesting their stable three-dimensional proximity in
at least one subpopulation of chromatin. Notably, these long-range interactions extend beyond
typical intrachromosomal contacts such as enhancer-promoter interactions, super-enhancers, and
TADs[12, 13]. Given the entropic cost of maintaining substantial colocalization probability over such
long genomic distances, we sought to explore their underlying mechanisms and potential
functional significance.

While the above statistical analyses identified long-range locus pairs with spatial proximity,
detecting them quantitatively from noisy Hi-C data remains challenging, as strong signals from
short-range locus pairs often overshadow weaker signals from long-range ones. To address this
challenge, we developed a computational pipeline, Chromosome Long-range colocalization
identifier through Outlier Detection (CLOD) (Fig. 1b, details in Methods). The pipeline operates
on the premise that colocalized locus pairs increase the contact frequency of their neighboring loci
along the genome. To enhance the signal-to-noise ratio, CLOD employs a sliding-window
averaging approach, allowing for the distinction of colocalization signals from background noise
(Extended Data Fig. 1d&e, details in Methods). It then compares locus pairs with similar genomic
distances and identifies outliers exhibiting abnormally high contact frequencies, defined as locus
pairs whose signals exceed two times the interquartile range above the third quartile of all pairs
with similar genomic distances ($\frac{Signal - Q3}{Q3 - Q1} > 2$).

To assess CLOD's performance in detecting experimentally verified colocalized locus pairs, we
first analyzed an MCF7 Hi-C dataset[14]. CLOD successfully identified a 150 kb loop (Extended
Data Fig. 1f), previously validated as a promoter-enhancer interaction of *PRIP1* using a 3C assay[16].
We then applied CLOD to a Hi-C dataset from the HCT116 cell line[17] and detected a colocalized
genomic pair separated by 27 Mb (Fig. 1c). This locus pair was independently confirmed by
fluorescence in situ hybridization (FISH)[10]. Collectively, these results demonstrate that the CLOD
pipeline can effectively detect colocalized locus pairs across both short and long genomic distances.

We then applied CLOD to Hi-C datasets from a panel of four normal differentiated human cell
lines, five cancer cell lines, and four embryonic cell lines[18-20] (Extended Data Table 1). Across
these datasets, we identified multiple colocalized locus pairs separated by 50 Mb or more in chr2
and chr14, separately (Fig. 1d&e and Extended Data Fig. 3). Notably, some of these long-range
pairs were consistently detected across multiple cell lines of different types (Fig 1f&g). To assess
whether the colocalization is restricted to human chromosomes, we examined colocalizations in
long (>150 Mb) chromosomes of other species, including six mammals (*Mus musculus* (mouse)[21,
22], *Rattus norvegicus* (rat), *Macaca mulatta* (rhesus monkey), *Bos taurus* (cattle), *Ovis aries*
(sheep), *Oryctolagus cuniculus* (rabbit)) and *Gallus gallus* (chicken)[22]. Consistently, we identified
colocalization of a substantial number of genomic locus pairs spanning genomic distances
exceeding 50 Mb (Extended Data Fig. 4, Extended Data Table 1), suggesting this spatial
organization is broadly conserved.

To determine whether the CLOD-identified long-range colocalization was transient or persistent,
we focused on a colocalized genomic pair on human chr14 separated by approximately 80 Mb
(Fig. 1h)[23]. We labeled the loci with two-color CRISPR-dCas9-guided fluorescent proteins in
HEK293T cells (Fig. 1i, Methods, Extended Data Table 2). Live-cell imaging confirmed the
sustained proximity and coordinated movement of green and red puncta, with their co-fluctuations
persisting throughout the observed 10 min in some cases (Fig. 1j &k, and Extended Data Movie
S1). Extended imaging further showed that the neighboring puncta remained in proximity during
an imaging period of about one hour (Fig. 1l and Extended Data Fig. 1g).

MERFISH chromatin tracing data reveal prevalence and conservation of colocalized loci > 100 Mb in human chr2.

While bulk Hi-C provides an ensemble-averaged view of chromosome conformations, it lacks the resolution to capture individual chromosome structures. It can neither distinguish whether colocalized pairs originate from the same or different homologous chromosomes, nor determine whether the observed contact frequencies reflect widespread interactions among cells or rare subpopulations. To cross-validate the prevalence of long-range pair colocalization and assess their distribution at the single-chromatin level, we analyzed previously published IMR90 chromatin tracing data obtained using DNA MERFISH[24].

As a technical control, we first re-analyzed the 3D distances (d_1) of MERFISH-labeled neighboring genomic loci across chr21 with a genomic separation $l_1 = 50\ kb$. The median distance for region chr21:32.45-33.35 Mb (Extended Data Fig. 5a) was found to be 256 nm, consistent with the previous report[24]. The overall distance distribution of neighboring genomic loci across chr21 followed a log-normal distribution, with a median distance of 325 nm (Extended Data Fig. 5b, left).

Next, we calculated the spatial distances of all neighboring genomic locus pairs across 2,991 copies of chr2 that were fluorescently traced with $l_1 = 250\ kb$ (Fig 2a, green line). The distribution also exhibited a log-normal pattern, with a median distance of 462 nm and a first quartile distance $d_r = 299$ nm (Fig. 2b, Extended Data Fig. 5b, right). For subsequent analyses, we adopted d_r as an intrinsic distance reference, classifying a genomic locus pair as in spatial proximity if their measured spatial distance $d \leq d_r$. This criterion aligns with previous studies using MERFISH data at an approximately 30 kb resolution, which employed a cutoff distance of ~ 200 nm to define chromatin interactions[25], and 500 nm for the chr21 data with 50 kb resolution to identify genomic pairs in proximity[24].

Most previous studies have focused on short-range chromosome interactions, such as enhancer-promoter interactions, which occur within genomic distances of ≤ 200 kb, typically in the $20 - 50$ kb range[26, 27]. Few studies have specifically focused on long-range chromatin interactions. Here, we calculated the spatial distances between pairs of loci with genomic separations of $l_2 \in (50, 100]$ Mb and $l_3 > 100$ Mb, respectively (Fig. 2a, purple and orange lines). Compared to d_1, the distributions of corresponding d_2 and d_3 shifted significantly toward larger values, with median distances of 1.86 μm and 2.08 μm, respectively (Fig. 2b). Interestingly, even at such large genomic separations, certain locus pairs exhibited spatial distances $< d_r$ (Fig. 2b, arrow). In the following analyses, we focused on the genomic locus pairs with $l_3 > 100$ Mb and $d_3 < d_r$, denoting them as long-range colocalizations (LRCs).

As expected, individual chr2 copies show diverse conformations (see examples shown in Fig. 2c, Extended Data Fig. 5c&d, Extended Data Movies 2-4), consistent with previous studies[25]. Moreover, a negative correlation exists between the total number of LRCs of an individual chromosome copy and its conformation measured by convex hull volume (Fig. 2d), suggesting a

potential relationship between LRCs and chromosome conformation. Notably, in these
conformations, LRCs either form scattered small clusters or join into larger ones, along with
varying compactness of the chromatin structures. In addition, LRC counts per chr2 copy range
from fewer than 10 to over 10,000, with a peak at ~ 600 per copy (Extended Data Fig. 5e). These
LRCs can be formed due to transient contacts through thermal fluctuations, or via stable structures.
While transient LRCs should appear infrequently among the entire cell population, stable ones are
expected to recur in subpopulations of chr2 copies. To test this hypothesis, we analyzed the
occurrence frequency of 155,009 unique LRCs across 2,991 chr2 copies (Fig. 2e). We identified
7,366 LRCs with occurrence frequencies beyond the 95% population interval, and defined them
as stable LRCs (sLRCs) (Fig. 2e & f). Each sLRC appeared in at least 23 chr2 copies (~0.8% of
the dataset), with some found in over 10% of samples. That is, sLRCs exhibited a significantly
higher probability of occurring within 299 nm compared to other LRCs (Fig. 2g).

Some sLRC detected via MERFISH were also detected with CLOD-identified colocalized pairs
(> 100 Mb separation) from Hi-C (Fig. 2h). Note that Hi-C and MERFISH detect proximity
differently: Hi-C captures direct contact, whereas MERFISH assesses spatial distances, using a
299 nm threshold here. Technical limitations also contribute to discrepancies — Hi-C may miss
repetitive or low-accessibility regions, whereas its much larger sample size (~1 million cells) could
detect rare configurations potentially overlooked by MERFISH (with ~3,000 chr2 copies). Despite
these differences, the strong agreement between Hi-C and MERFISH data reveals that stable LRCs
are prevalent in chromatin structures.

To further validate these findings, we analyzed the locus-pair contact frequencies of IMR90 chr2
from a micro-C dataset[28] and compared them with MERFISH results. Compared to standard Hi-C,
the superior coverage per genomic bin of micro-C[29] provides sufficient signal-to-noise ratios for
direct analysis of raw read counts without using additional algorithms such as CLOD. When
considering micro-C data with contact counts >10 as colocalization pairs, 556 sLRCs that were
identified from MERFISH data are also present in micro-C data (Extended Data Fig. 5e, left). Even
when applying stringent filtering criteria (contact counts >20), we still identified a subset of LRCs
present in both micro-C and MERFISH (Extended Data Fig. 5e, right). Therefore, both
sequencing-based and imaging-based approaches consistently identified sLRCs in IMR90 chr2. In
addition, LRCs and sLRCs were also present in other long chromosomes of IMR90 cells (Extended
Data Fig. 5f).

To assess whether LRCs exist at the single-cell level in other species, we examined a DNA
seqFISH+ dataset of chr1 of mouse NMuMg cells (Fig. 2i) [30]. Applying the same analytical
framework used for MERFISH, we measured spatial distance distributions for genomic loci in the
seqFISH+ data separated by 25 kb, 250 kb, and over 100 Mb, respectively. Those with 25 kb have
the smallest median 3D distance at 253 nm. Notably, the first quartile of spatial distances for loci
250 kb apart was 281 nm in seqFISH+, closely matching the 299 nm threshold from MERFISH
data (Extended Data Fig. 5g). We therefore adopted a universal LRC definition across platforms:
loci with genomic separation >100 Mb and spatial proximity <299 nm. The seqFISH+ dataset

provided substantially lower genomic coverage per chromosome than MERFISH, with most long-
range loci pairs detected in only ~ 200 – 300 chr1 copies out of 617 or more copies (Extended Data
Fig. 5h). To identify robust colocalization, we defined stable LRCs (sLRCs) as those present in at
least 10, or ~5% of, individual chromosomes (Extended Data Fig. 5h&i). All these results further
demonstrate that colocalization of genomically distant loci is a universal phenomenon in long
chromosomes, irrespective of species origin or chromosome identity.

A subset of genomic sites serves as nucleation centers (NCs) for clustering sLRCs.

Our structural analyses revealed that sLRC-associated genomic loci form intertwined networks
with sLRCs acting as links (Fig. 2c&i, Extended Data Fig. 5c&d, Extended Data Movies S5-S8).
Linkage maps from Hi-C, micro-C, and MERFISH data showed that certain genomic loci
colocalize with multiple distant loci (Fig. 2f&h Extended Data Fig. 5e). These loci likely function
as NCs, linking multiple sLRCs and forming the network (Fig. 3a). Some specific locations across
chr2, such as regions at ~28 – 40 Mb and the ends of IMR90 chr2, spatially colocalized with
significantly more well-separated cis-loci compared to other loci on the same chromosome and
thus can serve as NCs (Fig. 3b, left & Fig. 3c). Notably, locus 242 Mb (NC 242 Mb) were
associated with the most sLRCs and was analyzed separately for clear visualization.

Further statistical analyses showed that 26% of sLRC-associated loci formed only one pair of
sLRC, while 50% established fewer than five. However, a subset of loci participated in over 150
sLRCs (Fig. 3b, right and Fig. 3d). Even within individual chr2 copies, some loci can be paired
with up to ~ 90 other distant loci to form sLRCs (Fig. 3d). A striking example was NC 242 Mb,
which paired with loci across the chromosome. Two examples shown in Fig. 3e illustrate the
extensiveness of sLRC partners at the individual chr2 copy level. At the population level, 89.8%
of the 2,991 observed chromosomes had at least one sLRC partner at NC 242 Mb. Other NCs also
showed frequent and diverse long-range interactions. For example, NC 34.25 Mb, which was
found in 38% of chr2 copies, colocalized with mainly three distant regions and formed various
sLRCs in different chr2 copies (Fig. 3f). Extended Data Table 3 lists all loci associated with at
least two sLRCs in chr2 copies.

To reveal the basic patterns of chromosome conformation composed of NCs, we represented
sLRCs as networks. As expected, the results showed that distant regions on chr2 clustered via
sLRCs, where NCs served as hubs in the network (with two representative chr2 copies shown in
Fig. 3g), and most individual chr2 copies had fewer than 30 isolated clusters (Fig. 3h). Further
statistical analyses revealed that most NC-coordinated sLRCs clusters (59.2%) contained only 2 –
5 loci, while 5.4% had over 100 loci as nodes (Fig. 3i). Detailed analyses revealed distinct
structural motifs (Fig. 3j): some NCs formed isolated single-NC clusters, while others coalesced
into multi-NC hubs via direct colocalization or shared pairing loci, forming clusters with 100 –
780 nodes.

To confirm that NCs engage in multiple LRCs across diverse experimental platforms and cellular
contexts, we further analyzed the 3D genome conformation of other cell lines from human and

other species captured by other techniques, including K562 and GM12878 by Linking mRNA to
Chromatin Architecture (LiMCA, Extended Data Fig. 6a) [31], HEK293T by single-cell split-pool
recognition of interactions by tag extension (scSPRITE, Extended Data Fig. 6b) [32], human sperm
Genome Architecture Mapping (GAM, Extended Data Fig. 6c) [33], IMR90 by micro-C (Extended
Data Fig. 6d), and NMuMg by seqFISH+ (Fig. 2i). Hub-like loci appeared in all these cell types
and across different 3D architecture detection techniques. Moreover, some NCs are conserved in
different cell types. For example, loci ~32-35 Mb associated with multiple sLRCs were identified
both in IMR90 by MERFISH and micro-C, and in K562 and GM12878 by LiMCA. Loci around
the 3'-end of chr2 form another NC in IMR90 by MERFISH, human sperm by GAM, and
HEK293T by scSPRITE.

These findings suggest that NCs link multiple genomic regions, assembling hub-like domains
that may be associated with chromosome conformation and territory segregation.

Neighboring LRCs cooperate to form stable complexes through multivalent binding

DNA MERFISH data revealed the widespread presence of sLRCs ($p \geq 0.8\%$) in IMR90 chr2
copies, with some occurring in $5 - 10\%$ of all copies (Fig. 2e). Notably, the genomic distributions
of the two ends of sLRCs are not uniform but clustered along the genome (Fig. 2f). Statistical
analysis confirmed that most sLRC-associated loci have their nearest neighboring sLRC genomic
loci within 500 kb (Extended Data Fig. 7a). Here we defined the genomic proximity distance
between two sLRC pairs a (loci a_1, a_2) and b (loci at b_1, b_2) as $l_{ab} = max\left((|a_1 - b_1|, |a_1 - b_2|), (|a_2 - b_1|, |a_2 - b_2|)\right)$ (Fig. 4b, Extended Data Fig. 7b). Therefore, we grouped sLRCs into
$b_2|), (|a_2 - b_1|, |a_2 - b_2|)$) (Fig. 4b, Extended Data Fig. 7b). Therefore, we grouped sLRCs into
metaLRCs, where each sLRC had at least one other sLRC within 250 kb at both ends (Fig 4c).
This process grouped 72% of sLRCs into 736 metaLRCs, each containing 2 to over 40 sLRCs
(Extended Data Fig. 7c).

We hypothesized that neighboring sLRCs stabilize each other, as revealed from the live-cell
imaging studies (Fig. 1i–l). To examine this hypothesis, we analyzed two classes of minimal
biophysical models with and without cooperativity between sLRCs within a metaLRC (Methods).
In both models, colocalization of an sLRC pair is assumed to be held by some interactions, e.g.,
mediated by proteins or noncoding RNAs. Consequently, a pair exists in either a bound or unbound
state, distinguished by whether the two loci colocalize. In the absence of other sLRC pairs, an
isolated sLRC pair stochastically transits between the two states with association (α) and
dissociation (γ) rate constants (Fig. 4a, left). For simplicity, assuming that the N sLRC pairs within
a metaLRC have the same dynamics and transit independently, one can estimate from the data
$\frac{\alpha}{\gamma} = \frac{p_1}{Np_0}$, where p_0 and p_1 are the frequencies of observing zero or one bound sLRC, respectively.
The probability of observing $i > 1$ bound sLRC within the metaLRC can then be predicted without
any adjustable parameter to be $p_i = \frac{(N-i+1)a}{i\gamma} p_{i-1}$.

The cooperative model assumes that the transition dynamics of neighboring sLRCs are not
independent due to a multivalent binding mechanism. In the presence of other bound sLRC pairs

within the metaLRC, the two loci of an unbound sLRC pair remain spatially close, leading to an increased effective association rate constant. For simplicity, assuming that the effective enhancement δ is the same in the presence of one or more neighboring bounded sLRC pairs, then its value can be determined by fitting the measured distribution of bound sLRCs, p_i vs. i for i > 1. Consequently the metaLRC as a whole would appear as an effective two-conformation system with no or at least one bound sLRC, respectively (Fig. 4a, middle & right).

We tested the predictions of the two models against the MERFISH data. Figure 4d depicted one representative metaLRC. Correlation analyses over MERFISH data confirmed the cooperativity between sLRCs within the metaLRC (Fig. 4e). Note that some sLRCs have higher correlations than others within the same metaLRC, indicating that the assumption of a single value of δ is only crudely valid. The model without cooperativity significantly underestimated the probabilities of p_i for i >1, while a single parameter $\delta > 0$ significantly improves the agreement between the experimental data and theoretical predictions. Two additional examples shown in the Extended Data Fig. 5d&e also support the cooperative model. The bimodal distribution in Extended Data Fig. 5d reveals an even stronger cooperativity than the simple model we considered, suggesting that the enhancement δ increases with the number of bounded sLRCs.

Analysis of individual chr2 copies further supported the cooperative model. The spatial distances between two tagged genomically distant regions in bound form were much shorter compared to those in free form chr2 copies (Fig. 4g, Extended Data Movies 9–12). When one or more sLRCs were bound, the two genomic regions within the metaLRC were held closer, while additional bound sLRCs (> 1) did not significantly alter spatial distance (Fig. 4h). The observed non-monotonic p_i vs. i curves for certain representative metaLRCs (e.g., 33.75–35.5 Mb & 146–146.25 Mb; 119.25–123.75 Mb & 242 Mb) suggested even stronger cooperativity than the minimal model predicted (see Methods) (Extended Data Fig. 7d&e).

Next, we examined all metaLRCs of chr2 with $N > 2$. The α/γ values directly obtained from the FISH data were < 0.007 (Fig. 4i), indicating that associating two distant loci is entropically unfavorable. However, the corresponding $(\alpha + \delta)/\gamma$ values were up to 10 − 100× higher, making the probability of observing a bound metaLRC over 1 – 10% (Extended Data Fig. 7f). That is, pairs of distant loci can remain stably proximate within a chromosome via neighbor-assisted multivalent binding, significantly enhancing structural stability and long-range genomic interactions.

Structural analyses reveal multiple candidate mechanisms for sLRC formation.

Colocalization of distant cis-loci has been intermittently observed. Three general mechanisms have been proposed to explain LRC formation (Fig. 5a, I–III): (I) A/B compartment, or loose/compact region segregation, may bring distant B compartment loci into transient proximity[34] or relatively stable colocalization via polycomb structures[11]; (II) Transcriptionally active loci, around which chromatin is typically loose and accessible, may colocalize via attaching to shared nuclear structures like speckles or nucleoli[35]; (III) The nuclear membrane may recruit multiple loci into

spatial proximity[36]. Alternatively, one may conceive mechanism (IV) as stable sLRC formation
independent of major nuclear structures or compartment segregation. While the first three have
been proposed previously[37], sLRC assemblies formed through Mechanism IV were considered
rare.

sLRC-forming loci are not significantly enriched in protein-coding gene-rich or gene-rare
regions (Extended Data Fig. 8a, top) nor in lncRNA-coding genes (Extended Data Fig. 8a, bottom).
Examination of A/B compartment identities among sLRC-forming loci revealed no clear pattern
or correlation between the two loci of each sLRC pair (Extended Data Fig. 8b–d), while the
majority of sLRCs have two ends at the chomatin loose region (Extended Data Fig. 8c, bottom),
which is consistent with the observation from a previous study[24]. Note that the A/B compartment
was calculated from bulk-level data. That is, an ensemble of chromosomes may be composed of
subpopulations with and without the presence of an sLRC under study; thus, results using such
bulk-averaged data are not conclusive. Bulk averaging also prevented us from concluding whether
sLRCs or the loci linked to sLRCs have specific epigenetic patterns (Extended Data Fig. 8e&f).

To connect the sLRC and associated loci to epigenetic marks and nuclear structures at the single-
chromosome level, we investigated the seqFISH+ dataset[30]. This dataset with single chromosome
resolution combines spatial coordinate information of each location on the mouse genome with its
epigenetic modification marks, such as H3K4me2, H3K27ac, H3K9me3, H3K27me3, etc., as well
as marks for nuclear structures, such as Polycomb (EZH2), speckles (SF3A66), and lamina-
associated domains (Lamin B1). To be consistent with the resolution of MERFISH, we grouped
the DNA seqFISH+ data to a 250 kb resolution, calculated the median mark level, and used the 3rd
quantile as the gate to identify whether a region has a high level of the targeted marker (Extended
Data Fig. 9a).

The epigenetic profiling reveals subsets of sLRCs exhibiting epigenomic features consistent
with the three identified formation mechanisms (Mechanisms I–III, Fig. 5a). For instance, among
the 323 detected copies of the 3.25 and 125.15 Mb locus pair on NMuMg chr1, 15 colocalized
copies exhibited elevated H3K9me3 and uniformly depleted euchromatin marks (H3K4me2 and
H3K27ac) relative to non-colocalized copies (Extended Data Fig. 9b-i). Another pair (15.25 and
151.25 Mb) displayed high H3K27me3 enrichment along with the Polycomb marker EZH2
(Extended Data Fig. 9b–ii). Both examples are consistent with compressed encounters driven by
A/B compartment segregation (Mechanism I).

A colocalized pair in Extended Data Fig. 9c instead showed high levels of active chromatin
markers (H3K4me2 and H3K27ac), enrichment of the nuclear speckle marker SF3A66, and low
H3K9me3 and H3K27me3, consistent with nuclear speckle–associated assembly (Mechanism II).
Extended Data Fig. 9d illustrates an example potentially following Mechanism III, with high
Lamin B1 enrichment indicating association with lamina-associated domains.

Notably, some sLRCs on mouse chr1 could not be assigned to these categories: the regional
epigenetic mark levels of the two ends of an sLRC show no strong correlation (Extended Data Fig.
9e–i); or all epigenetic mark levels of both ends are relatively low or show no clear pattern

(Extended Data Fig. 9e–ii). These observations suggest additional formation mechanisms,
collectively classified as Mechanism IV.

The MERFISH data contain no epigenetic information at single-chromosome level. Instead, we
examined the local density of a locus, noting that the two ends of an sLRC formed through
Mechanism I-III are expected to share similar local packing structures. Specifically, we quantified
local geometric structures within 299 nm of a locus by defining regional density of short-range (<
10 Mb) neighbors (RD-SN) and regional density heterogeneity (RDH), which measures the
uniformity of local spatial distributions (Fig. 5b). Comparing RD-SN to A/B compartments
identified from Hi-C data, an A compartment has a higher probability of having a loose
neighborhood than a B compartment (Extended Data Fig. 8g). Then one expects that Mechanism
I and III lead to high RD-SN for both loci associated with an sLRC, Mechanism II leads to low-
low RD-SN, while the pattern for Mechanism IV is undetermined. Furthermore, a locus next to a
nuclear structure (Mechanism II and III) is expected to exhibit high RDH due to the excluded
volume of the latter. Therefore, a combined RD-SN and RDH analysis can identify sLRCs not
convincingly explainable by Mechanisms I-III, e.g., those with both RD-SN and RDH low on both
loci. Indeed, sLRCs show combinatorial patterns of these two quantities, implying existence of
diverse mechanisms to sLRC formation (Fig. 5c&d), as also reflected from examining
representative structures (Fig. 5e).

Furthermore, we observed spatial colocalization of *cis*-loci that are well separated on the genome
in chromosomes shorter than 150 Mb. Specifically, we analyzed the colocalization state of locus
pairs in chromosome 21 of IMR90 cells at 20 Mb, which is about half of the region (~36.3 Mb)
labeled with MERFISH. The distribution of spatial distances between most locus pairs longer than
20 Mb in chr21 ranged between 0 and 5 μm, and colocalized pairs of loci denote those whose
spatial distances are closer than 299 nm in individual chromosome structures as aforementioned
(Extended Data Fig. 8h). As expected, both linkage maps and 3D structure plots show the existence
and variety of LRCs in individual chr21 (Fig. 5f). While most of 32,341 loci pairs longer than 20
Mb were only observed in fewer than 200 out of 7,591 chr21 copies, 1,718 or 5% of them appeared
in more than 325 copies, which were identified as stable colocalization locus pairs here (Fig. 5g).
Moreover, similar to chr2, a linkage map showing the stable colocalization loci pairs reveals that
clustering of sLRCs and loci serving as nucleation centers also exists in chr21 (Fig. 5h).

Together, all analyses indicate the prevalence of long-range colocalization in chromosomes, and
the epigenetic, positional, and spatial distribution results suggest an additional, previously
underappreciated mechanism contributing to stable sLRC formation.

DISCUSSION

Our analyses of Hi-C and DNA FISH chromosome tracing data reveal widespread colocalization
of genomic loci separated by distances far exceeding a typical TAD. It is important to note that
our study employed stringent criteria to identify representative examples rather than exhaustively

cataloging all stable long-range colocalizations. While we focused on loci separated by >100 Mb,
similar structures exist at shorter genomic distances, particularly in smaller chromosomes such as
chr21 (~48 Mb).

Statistically, the probability of multiple genomic loci with $l \geq 100$ Mb being in physical
proximity without stable structural mechanisms would be negligible. Some sLRCs may form
through known mechanisms such as compartment segregation or anchoring to nuclear structures
such as nucleoli, speckles, or the nuclear envelope. However, many sLRCs and NCs are buried in
the nuclear interior, show no signature of compartment segregation, and are away from large
nuclear structures, suggesting additional nuclear structure-independent mechanisms. generically
assigned to Mechanism IV.

The molecular nature of the interactions holding the assemblies needs further investigation. An
assembly can be liquid-, gel-, or solid-like, similar to what is observed in other contexts. The
existence of NCs suggests that they may play roles in LRC formation different from their
partnering loci. These NC sites can either be nucleation centers for liquid droplet formation or act
as scaffolds for solid assembly mediated by elements such as non-coding RNAs or bridging
proteins. For instance, the human chr2 242 Mb locus is enriched in repetitive sequences, which
may serve as selective binding sites for partnering loci, either directly or through other bridging
factors. Another 32–35 Mb locus on human chr2 is also rich in repetitive sequences and non-
coding RNAs such as LIN1[38]. Multi-way chromosome assemblies have been identified in different
contexts, such as enhancer-promoter hubs[39], superenhancers, and the multi-chromosome enhancer
clusters controlling monoallelic olfactory receptor expression in olfactory sensory neurons[40-42].
The assemblies of sLRCs and metaLRCs may form under similar molecular mechanisms. One
known example is that Polycomb repressive complex 2 uses RNA binding to coordinate long-
range chromosome loop formation and H3K27me3 spreading[42]. Falk et al. argue that attractions
between heterochromatic regions and lamina-chromatin interactions, i.e., Mechanisms I and III,
are critical to recapitulate compartmented chromatin structures[43]. Analyses in this work reveal
existence of long-range colocalization structures showing no clear features of lamina-chromatin
interactions and heterochromatin, and their molecular mechanism(s) of formation need further
investigation.

A key unsolved question is how sLRC loci locate and associate with each other. One plausible
mechanism is that, during post-mitotic DNA decondensation, some genomically distant loci may
be in spatial proximity and associate stochastically, leading to subsequent formation of additional
sLRCs and stabilization of the assembled structures. Liquid-phase droplet fusion can also drive
two regions into proximity to form an sLRC[44].

Besides their varied formation mechanisms, these sLRC structures likely play diverse functional
roles, including transcriptional regulation and broader chromosome 3D organization. Proper DNA
folding into spatially segregated territories with minimal entanglement and knot formation is
essential for eukaryotic genome organization and gene regulation. This process must be robust
across all cell types, including those with abnormal karyotypes, and must function despite cell

type-specific euchromatin/heterochromatin partitioning, folding stochasticity, and mutations.
From an engineering perspective, achieving this through a single centralized mechanism requiring
precise tuning would be challenging. Instead, our findings suggest a redundant, distributed
component mechanism that facilitates chromosome folding into compact structures (Fig. 5i). The
genome harbors an extensive repertoire of NCs and partnering loci separated by long genomic
distances, from which only a subset is available for a given cell type. At the individual chromosome
level, subsets of these loci stochastically form sLRCs and larger structures, contributing to the
observed heterogeneity of chromosome structures[45]. These structures provide a "divide-and-
conquer" mechanism to facilitate forming chromosome structures at smaller length scales, e.g., by
forming segregated globally compact chromosome configurations, and closed-end boundary
conditions for simultaneous crumpled folding at multiple regions (Extended Data Fig. 4d)[5], and
enhanced contact frequencies for forming loops at smaller scales.

In summary, our analyses of chromosome tracing and Hi-C data uncover prevalent, conserved
large-sized loops and their clustering into stable structures. These findings raise important
questions regarding the molecular mechanisms, temporal dynamics, and functional significance of
these structures on chromosome configurations, gene transcriptional activity, and cell type
regulation.

References

1.   Dill, K.A. & MacCallum, J.L. The protein-folding problem, 50 years on. *science* **338**, 1042-1046 (2012).

2.   Bolzer, A. *et al.* Three-dimensional maps of all chromosomes in human male fibroblast nuclei and prometaphase rosettes. *PLoS biology* **3**, e157 (2005).

3.   Lieberman-Aiden, E. *et al.* Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *science* **326**, 289-293 (2009).

4.   Wang, S. *et al.* Spatial organization of chromatin domains and compartments in single chromosomes. *Biophysical Journal* **112**, 217a (2017).

5.   Grosberg, A., Rabin, Y., Havlin, S. & Neer, A. Crumpled globule model of the three-dimensional structure of DNA. *Europhysics letters* **23**, 373 (1993).

6.   Gürsoy, G., Xu, Y., Kenter, A.L. & Liang, J. Spatial confinement is a major determinant of the folding landscape of human chromosomes. *Nucleic acids research* **42**, 8223-8230 (2014).

7.   Polovnikov, K.E. *et al.* Crumpled polymer with loops recapitulates key features of chromosome organization. *Physical Review X* **13**, 041029 (2023).

8.   Mateos-Langerak, J. *et al.* Spatially confined folding of chromatin in the interphase nucleus. *Proceedings of the National Academy of Sciences* **106**, 3812-3817 (2009).

9.   Quinodoz, S.A. *et al.* Higher-order inter-chromosomal hubs shape 3D genome organization in the nucleus. *Cell* **174**, 744-757. e724 (2018).

10.   Jäger, R. *et al.* Capture Hi-C identifies the chromatin interactome of colorectal cancer risk loci. *Nature communications* **6**, 6178 (2015).

11.   Kraft, K. *et al.* Polycomb-mediated genome architecture enables long-range spreading of H3K27 methylation. *Proceedings of the National Academy of Sciences* **119**, e2201883119 (2022).

12.   Pombo, A. & Dillon, N. Three-dimensional genome architecture: players and mechanisms. *Nature reviews Molecular cell biology* **16**, 245-257 (2015).

13.   Whyte, W.A. *et al.* Master transcription factors and mediator establish super-enhancers at key cell identity genes. *Cell* **153**, 307-319 (2013).

14.   Barutcu, A.R. *et al.* Chromatin interaction analysis reveals changes in small chromosome and telomere clustering between epithelial and breast cancer cells. *Genome biology* **16**, 214 (2015).

15.   Rao, S.S. *et al.* A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. *Cell* **159**, 1665-1680 (2014).

16.   Jin, F. *et al.* A high-resolution map of the three-dimensional chromatin interactome in human cells. *Nature* **503**, 290-294 (2013).

17.   Rao, S.S. *et al.* Cohesin loss eliminates all loop domains. *Cell* **171**, 305-320. e324 (2017).

18.   Dixon, J.R. *et al.* Chromatin architecture reorganization during stem cell differentiation. *Nature* **518**, 331-336 (2015).

19.   Rhie, S.K. *et al.* A high-resolution 3D epigenomic map reveals insights into the creation of the prostate cancer transcriptome. *Nature communications* **10**, 4154 (2019).

20.   Ren, B. *et al.* High-resolution Hi-C maps highlight multiscale 3D epigenome reprogramming during pancreatic cancer metastasis. *Journal of Hematology & Oncology* **14**, 120 (2021).

21.   Liu, M. *et al.* Tracing the evolution of single-cell cancer 3D genomes: an atlas for cancer gene discovery. *bioRxiv*, 2023.2007. 2023.550157 (2024).

22. Li, D. *et al.* Comparative 3D genome architecture in vertebrates. *BMC biology* **20**, 99
(2022).
23. Zuin, J. *et al.* Cohesin and CTCF differentially affect chromatin architecture and gene
expression in human cells. *Proceedings of the National Academy of Sciences* **111**, 996-
1001 (2014).
24. Su, J.-H., Zheng, P., Kinrot, S.S., Bintu, B. & Zhuang, X. Genome-scale imaging of the
3D organization and transcriptional activity of chromatin. *Cell* **182**, 1641-1659. e1626
(2020).
25. Bintu, B. *et al.* Super-resolution chromatin tracing reveals domains and cooperative
interactions in single cells. *Science* **362**, eaau1783 (2018).
26. Chepelev, I., Wei, G., Wangsa, D., Tang, Q. & Zhao, K. Characterization of genome-
wide enhancer-promoter interactions reveals co-expression of interacting genes and
modes of higher order chromatin organization. *Cell research* **22**, 490-503 (2012).
27. Furlong, E.E. & Levine, M. Developmental enhancers and chromosome topology.
Science **361**, 1341-1345 (2018).
28. Palikyras, S. *et al.* Rapid and synchronous chemical induction of replicative‐like
senescence via a small molecule inhibitor. *Aging Cell* **23**, e14083 (2024).
29. Krietenstein, N. *et al.* Ultrastructural details of mammalian chromosome architecture.
Molecular cell **78**, 554-565. e557 (2020).
30. Takei, Y. *et al.* Spatial multi-omics reveals cell-type-specific nuclear compartments.
Nature **641**, 1037-1047 (2025).
31. Wu, H. *et al.* Simultaneous single-cell three-dimensional genome and gene expression
profiling uncovers dynamic enhancer connectivity underlying olfactory receptor choice.
Nature Methods **21**, 974-982 (2024).
32. Arrastia, M.V. *et al.* A single-cell method to map higher-order 3D genome organization
in thousands of individual cells reveals structural heterogeneity in mouse ES cells.
bioRxiv, 2020.2008.2011.242081 (2020).
33. Irastorza-Azcarate, I. *et al.* Extensive folding variability between homologous
chromosomes in mammalian cells. *Molecular Systems Biology* **21**, 735-775 (2025).
34. Gibson, B.A. *et al.* Organization of chromatin by intrinsic and regulated phase separation.
Cell **179**, 470-484. e421 (2019).
35. Kim, J., Venkata, N.C., Hernandez Gonzalez, G.A., Khanna, N. & Belmont, A.S. Gene
expression amplification by nuclear speckle association. *Journal of Cell Biology* **219**,
e201904046 (2019).
36. Van Steensel, B. & Belmont, A.S. Lamina-associated domains: links with chromosome
architecture, heterochromatin, and gene repression. *Cell* **169**, 780-791 (2017).
37. Dekker, J. & Mirny, L.A. The chromosome folding problem and how cells solve it. *Cell*
187, 6424-6450 (2024).
38. Percharde, M. *et al.* A LINE1-Nucleolin Partnership Regulates Early Development and
ESC Identity. *Cell* **174**, 391-405.e319 (2018).
39. Lando, D. *et al.* Enhancer-promoter interactions are reconfigured through the formation
of long-range multiway hubs as mouse ES cells exit pluripotency. *Molecular Cell* **84**,
1406-1421.e1408 (2024).
40. Tian, X.-J., Zhang, H., Sannerud, J. & Xing, J. Achieving diverse and monoallelic
olfactory receptor selection through dual-objective optimization design. *Proceedings of*
the National Academy of Sciences **113**, E2889-E2898 (2016).

41.   Pourmorady, A. & Lomvardas, S. Olfactory receptor choice: a case study for gene
533   regulation in a multi-enhancer system. *Current Opinion in Genetics & Development* **72**,
534   101-109 (2022).
535   42.   Kraft, K. *et al.* Polycomb-mediated genome architecture enables long-range spreading of
536   H3K27 methylation. *Proceedings of the National Academy of Sciences* **119**,
537   e2201883119 (2022).
538   43.   Falk, M. *et al.* Heterochromatin drives compartmentalization of inverted and
539   conventional nuclei. *Nature* **570**, 395-399 (2019).
540   44.   Strom, A.R. *et al.* Phase separation drives heterochromatin domain formation. *Nature*
541   **547**, 241-245 (2017).
542   45.   Finn, E.H. *et al.* Extensive heterogeneity and intrinsic variation in spatial genome
543   organization. *Cell* **176**, 1502-1515. e1510 (2019).
544
545

Abbreviations and glossary:

LRC: Long-range colocalization for genomic locus pairs > 100 Mb and within 299 nm in spatial distance.

sLRC: Stable long-range colocalization for LRC pairs with occurrence frequency > 4% in the IMR90 chr2 MERFISH dataset, or the top 5% in IMR90 chr1, chr3 and chr4 MERFISH dataset, or ≥15 chromosomes (~7% in detected chromosomes) in NmuMg chr1 DNA seqFISH+.

metaLRC: a group of sLRCs, in which for each sLRC there was at least one other sLRC with 250 kb for both ends.

NC: Nucleation center referring to locus having multiple sLRC partner loci.

RD-SN: Regional density of short-ranged neighbors (< 10 Mb) within a 299 nm range of a tagged locus.

RDH: Regional density heterogeneity.

Acknowledgments: We thank Tom Mistelli, Harinder Singh, Jing Chen, Jian Liu for helpful discussions.

Funding:

National Institute of General Medical Sciences R01GM148525 (JX)

The Charles E. Kaufman Fund of the Pittsburgh Foundation KA2018-98550 (JX)

Author contributions:

Conceptualization: JZ, JX

Methodology: JZ, SW, SCW, JX

Investigation: JZ, JX

Visualization: JZ, JX

Funding acquisition: JX

Project administration: JX

Supervision: JX

Writing – original draft: JX

Writing – review & editing: JZ, SW, SCW, JX

Competing interests: Authors declare that they have no competing interests.

 Figure captions

Fig. 1 Hi-C data reveals prevalent existence of colocalized pairs of long-range genomic loci.
a, Scatter plot (left) and boxplot (right) showing the decay trend (blue line) of balanced contact
intensity with increasing genomic distance. Red boxed points represent locus pairs > 50 Mb apart
with contact intensity exceeding the third quartile of those observed for ~1 Mb pairs. **b**, Schematic
of the CLOD pipeline. **c**, Experimental validation of colocalized pairs predicted by CLOD using
Hi-C data from HCT116 cells at 50 kb resolution. Left: Raw contact map between the two
segments. Right: $\frac{Signal - Q3}{Q3 - Q1}$ contour map of the same region as the left one. **d-e**, Representative
linkage maps of colocalized locus pairs > 50 Mb on chr2 and chr14 in two different cell lines. **f-**
g, Linkage maps of colocalized pairs > 50 Mb apart, shared across at least two of the following

cell types: four embryonic cell lines (green), four normal differentiated cell lines (pink), and four
cancer cell lines (blue). **h**, CLOD-based identification of colocalized pairs on chr14 from Hi-C
data of IMR90 and HEK293T cells at 50 kb resolution. **i**, CRISPR-dCas9 two-color labeling sites
on chr14. **j**, Live-cell imaging snapshots of a HEK293T cell labeled with RFP and GFP. Scale bar:
2 μm. **k**, 10-min time-resolved trajectories of blue and red puncta in the labeled cell from panel j.
l, Distance fluctuations between blue and red puncta pairs in four cells that were tracked for 60
min. The left and right bars indicate the maximum and minimum distances between a pair of puncta.

Fig. 2 MERFISH chromatin tracing reveals widespread long-range colocalization (LRC). a,
Schematic illustration of genomic pairs and their genomic versus spatial distances. **b**, Density
curves of spatial distance distributions for loci pairs with genomic separations of 250 kb, 50-100
Mb, and >100 Mb. The dashed line represents the first quantile of neighboring locus pairs. **c**, LRC
linkage maps and 3D structures of 3 representative chromosomes. **d**, Scatter plot showing the
relationship between chromosome compaction, quantified by the convex hull volume, and the
number of LRCs. **e**, Histogram (top) and boxplot (bottom) showing the distribution of chr2 copy
counts (n = 2991) containing LRCs. LRCs present in >23 copies (red dashed line) are classified as
stable LRCs (sLRCs). The pie diagram shows the percentage of LRCs classified as sLRCs. **f**,
Identification of chr2 sLRCs from DNA FISH data. **g**, Spatial distance distributions of locus pairs
with genomic separation >100 Mb. Purple lines (n = 6443): sLRCs; blue lines (n = 6443): randomly
selected non-sLRCs. The red dashed line represents the first quantile of spatial distances between
neighbor loci on chr2 (299 nm). **h**, Comparison of colocalized chr2 locus pairs (>100 Mb apart)
identified in IMR90 cells using DNA MERFISH (blue lines) and Hi-C (green lines) data.
Overlapping links were highlighted in brown.

Fig. 3 Nucleation centers (NC) coordinate clustering of sLRCs. a, Left: Schematic illustration of a nucleation center (NC) (orange) forming spatial proximity with multiple partner LRC loci (purple). Right: Ring plot depicting the same interaction network. **b**, Left: Line plot showing the number of sLRCs originating from a given locus, based on 2,991 chr2 copies. Right: Histogram illustrating the distribution of loci (x-axis) connecting to a given number of sLRC (y-axis). **c**, sLRC linkage map with corresponding numbers of potential sLRCs (green bars outside the ring) originating from specific loci in the chr2. **d**, Scatter plot comparing the total number of potential sLRCs from targeted loci across the analyzed chr2 population to the maximum number of sLRCs from a single targeted locus in an individual chr2 copy. Top and right panels: Corresponding distributions of numbers shown in the scatter plot. **e-f**, Linkage maps and 3D structures highlighting two representative NCs (NC 242 Mb and NC 34.25 Mb) on chr2, along with their

630   associated sLRCs. **g**, 3D structural representations and network maps showing connections among
631   NC loci in two representative chr2 copies. **h**, Histogram displaying the number of locus clusters –
632   defined as sets of loci connected to at least one other locus – per individual chr2 copy. **i**,
633   Histogram depicting the number of loci per locus-cluster, as identified in panel h. **j**, Diagrams
634   illustrating single NC elements and various types of connections between NCs.

Fig. 4 DNA FISH data analysis reveals that multivalent binding stabilizes sLRC clusters. a,
Left: Schematic illustration of monovalent binding vs. multivalent binding models. α: association
rate constant of a single pair in the absence of other bound sLRCs; γ: dissociation rate constant of
a single pair; δ: association rate enhancement due to neighboring bound sLRCs. Right: A multi-
valent site is coarse-grained into a two-state system – either free or bound (with one or more
bound sLRCs) two-state system. **b**, Frequency distribution of the genomic separation between two
LRCs. **c**, sLRC and metaLRC linkage map. The outer circle represents chr2, and each line denotes
a sLRC. Gray lines: isolated sLRC; colored lines (except gray): sLRCs belonging to a metaLRC.
d, A representative metaLRC. **e**, Heatmap showing correlation between sLRCs within the
metaLRC from panel d. **f**, Comparison of the probabilities of observing various numbers of bound
sLRCs within the metaLRC (purple bar) vs. predictions based on an independent sLRC binding

model (orange bar), and a multivalent binding model (blue dashed line). p_0 is not shown. **g**,
Representative 3D structures of chr2, highlighting the metaLRC from panel d. Top: Structures
with at least one bound sLRC within the metaLRC. Bottom: Structures with no bound sLRC within
the metaLRC. **h**, Violin plot showing the relationship between 3D distances (between centers of
minimal-sized bounding spheres of the two genomic regions forming the metaLRC and the number
of bound sLRCs per individual chr2 copy. **i**, Multivalent binding model parameters for metaLRCs,
obtained through fitting DNA MERFISH data.

Fig. 5 Possible mechanisms of sLRC formation and functional roles on chromosome folding.
a, Schematic of four possible mechanisms for sLRC formation. I & II: A/B compartment
segregation leads to compressed encounters of genomically distant loci in the B and A
compartments, respectively. III: Colocalization of transcriptionally inactive, genomically distant
loci interacting with common nuclear membrane structures. IV: Specific interactions between
genomically distant loci form an assembly without the involvement of large nuclear structures. **b**,
Definition of regional density of short-range neighbors (RD-SN) and regional density
heterogeneity (RDH) to characterize structural properties of sLRCs formed via mechanisms in
panel a. **c-d**, Scatter plots of RD-SN and RDH around both ends of individual sLRCs in each chr2
copy. Each dot represents one sLRC in a single chr2 copy. **e**, 3D structures of six representative
chr2 copies highlighted in panel c&d. **f**, Representative MERFISH structures of chr21. Black lines
represent LRCs. **g**, Histogram (top) and boxplot (bottom) showing the distribution of chr21 copy

676 counts (n = 7,591) containing LRCs. LRCs present in >325 copies (red dash line) are classified as
677 stable LRCs (sLRCs). **h**, Identification of chr2 sLRCs from DNA FISH data. **i**, Schematic
678 illustration of the proposed excessively redundant distributed component mechanism for
679 chromosome folding.

Methods

Hi-C data analysis

Raw Hi-C data of human were downloaded from GEO, as shown in Extended Data Table 1, and were aligned to the human genome *hg38* using the Burrows-Wheeler Alignment tool[46]. Hi-C contacts were then identified and filtered with Pairtools[47]. The contact matrices of balanced contact intensity used for the CLOD algorithm were created with Cooler[48]. Hi-C counts of other species were downloaded from GEO directly (Extended Data Table 1).

Chromosome Long-range colocalization identifier through Outlier Detection (CLOD) pipeline

To establish a minimalistic and robust pipeline to detect long-range colocalization between cis-loci based on Hi-C results, we smoothed the noisy Hi-C data and then highlighted the robust off-diagonal signals representing chromatin contacts between distal genomic regions. In detail, first we used the balanced Hi-C results from Cooler and converted the original reads to the average of the window to smooth the random noisy signals. Then to give prominence to the signal at off-diagonal regions of contact matrix, we grouped all pairs of loci with the same genomic distances (l) and those with $l \pm resolution$ and calculated the quantiles. The final output of the CLOD is the fold change of the interquartile range above the third quantile of all pairs with similar genomic distances ($\frac{Signal - Q3}{Q3 - Q1}$).

To test the impact of Hi-C resolution and window size on CLOD outlier detection and seek an appropriate combination, we analyzed three Hi-C resolutions (50 kb, 100 kb, and 250 kb) and five sliding window sizes (3×3, 5×5, 7×7, 9×9, and 11×11). The results showed that with different resolutions, the number of CLOD identified outliers varied from a few thousands to close to a hundred of thousands, while the ratios of CLOD identified outliers to total loci pairs ranged from 0.004 to 0.009 (Extended Data Fig. 1d). Since the amount of CLOD outliers obtained from 250 kb resolution was much less than those from resolution 50 kb and 100 kb, we compared the sliding window size using the latter two resolutions. With 50 kb resolution, the identified CLOD outliers were sensitive to window size, with 551 locus pairs consistently present in all sliding window sizes. CLOD outliers identified from 100 kb Hi-C resolution were more robust with diverse window sizes (Extended Data Fig. 1e). Balancing between processing time and identification efficiency, Hi-C long-range colocalization was identified using the 100 kb resolution with a 7×7 window, unless otherwise stated.

MERFISH data

The Euclidean coordinates of chromosome fragments were directly downloaded from Zenodo repository (records:3928890).

DNA seqFISH+ data

The Euclidean coordinates of chromosome fragments and the level of epigenetic marks were directly downloaded from Zenodo repository (records:7693825).

 CRISPR-dCas9 two-color labeling

The plasmid cloning method used for CRISPR-dcas9 fluorescent labeling was adapted from a
previously established protocol[49]. The sgRNAs used in this study are listed in Extended Data Table
2.

Cell culture

HEK293T cells (#ATCC-CRL-3216) were cultured in DMEM (Invitrogen #11995073)
supplemented with 10% FBS (Gbico #10437028), 100 units/ml penicillin, and 100 µg/ml
streptomycin (Invitrogen). Cells were maintained at 37°C in 5% CO2, and used within 15–30
passages for experiments.

For transfection, HEK293T cells were seeded onto 35 mm glass-bottom dishes one day before the
experiments. In the next day, 750 ng of donor template and 750 ng of sgRNA-Cas9 plasmid were
co-transfected using FuGENE HD Transfection Reagent (Promega #E2311). Cells were incubated
for 48 hours before imaging[49].

Time-lapse imaging

For imaging presentation (Fig. 1j), cells were visualized using a Nikon Ti microscope equipped
with a 100x oil objective. Excitation wavelengths were 470 nm for GFP and 555 nm for RFP. Cells
were imaged every 10 seconds for 10 minutes with an exposure time of 50 ms.

For trajectory tracking (Fig. 1l), cells were maintained in a Tokai Hit Microscope Stage Top
Incubator and imaged using a Nikon Ti microscope with FITC and TRITC channels (excitation
wavelength is 470 nm and 555 nm for GFP and RFP, respectively; 60x oil objective). Cells were
imaged every minute for 60 minutes with an exposure time of 100 ms[50].

Imaging processing
Due to the low number of cells, cell segmentation and puncta recognition were performed by
CellProfiler[51] and corrected manually.

Derivation of the minimal multivalent binding model

Consider a metaLRC that contains N pairs of sLRCs. Given the observed abnormally high contact
frequency of a sLRC pair as compared to non-sLRC pairs separated with similar genomic distance,
it is likely that the two loci are held together through some direct interactions or mediated by other
molecular components. Thus, we can assume that an sLRC pair i exists in one of the two possible
configurations, $\sigma_i = 1$ if the pair is bound (with spatial distance $d_r <\, = 299$ nm), and 0 otherwise.
We assume this metaLRC is genomically distant from other metaLRCs and sLRCs, so for a good
approximation there is no need to explicitly consider the influence of the latter on the association-
dissociation dynamics of sLRCs in this tagged metaLRC.

Suppose that without influence from other sLRCs within the tagged metaLRC, an sLRC pair has
a bare association rate α and a dissociation rate γ. Then at kinetic equilibrium, the probability in
the bound state is $\frac{\alpha}{\alpha+\gamma}$, and from the FISH data we expect that $\alpha \ll \gamma$. Here for simplicity we

assume that each sLRC pair has the same association-dissociation kinetic parameters. Let's focus
on one pair i. When there is one or more other bound sLRC pairs within the metaLRC, after
dissociation the two loci of pair i are constrained by the nearby bound sLRCs and cannot diffuse
far away from each other before rebind. Note that the genomic distance between these two loci of
a sLRC $\geq$ 100 Mb, and by definition the genomic distance between two sLRCs i and j within a
metaLRC l_{ij} << 100 Mb. Consequently, at the presence of other bound sLRC pair(s), the two loci
of a dissociated sLRC have an increased probability to confront each other and rebind, so the
effective association constant changes to $\alpha + \delta$, and we expect that $\delta \gg \alpha$. For simplicity, given
l_{ij} << 100 Mb let us assume that each sLRC pair has the same effect on any other one, and the
effect is not additive but δ remains the same value at the presence of one or more bound sLRC
pairs. Given the genomic distance between two loci belonging to two different sLRCs, we further
assume that there is no direct interaction between two sLRC pairs to simplify the analyses. That
is, we assume that the dissociation constant of a sLRC pair, γ, is not affected by the presence of
other bound sLRC pairs.

With the above model, one can write down a set of master equation for finding that the metaLRC
exists with 0, 1,…, N pairs of bound sLRCs,

$\frac{dp_0}{dt} = -Nap_0 + \gamma p_1,$

$\frac{dp_1}{dt} = -\big((N-1)(a+\delta) + \gamma\big)p_1 + 2\gamma p_2 + Nap_0,$

$\frac{dp_2}{dt} = -\big((N-2)(a+\delta) + 2\gamma\big)p_2 + 3\gamma p_3 + (N-1)(a+\delta)p_1,$

...

$\frac{dp_{N-1}}{dt} = -\big((a+\delta) + (N-1)\gamma\big)p_{N-1} + N\gamma p_N + 2(a+\delta)p_{N-2},$

$\frac{dp_N}{dt} = -N\gamma p_N + (a+\delta)p_{N-1}.$

Then at steady state,

$p_1 = \frac{Na}{\gamma}p_0,$

$p_2 = \frac{(N-1)(a+\delta)}{2\gamma}p_1,$

$p_3 = \frac{(N-2)(a+\delta)}{3\gamma}p_2,$

...

$p_N = \frac{a+\delta}{N\gamma}p_{N-1}.$

Define the probability of observing presence of one or more pairs as

$p = 1 - p_0 = \sum_{i=1,\dots,N} p_i.$

One has,

$$p = \left(\frac{N!\,(a+\delta)^{N-1}}{N!\,\gamma^{N-1}} + \cdots + \frac{(N-1)N(a+\delta)}{2\gamma} + N \right) \frac{a}{\gamma} p_0$$

$$= \left(\frac{N!\,(a+\delta)^{N}}{N!\,\gamma^{N}} + \cdots + \frac{(N-1)N(a+\delta)^2}{2\gamma^2} + N\frac{(a+\delta)}{\gamma} \right) \frac{a}{a+\delta} p_0$$

$$= \left(\left(1 + \frac{a+\delta}{\gamma}\right)^N - 1 \right) \frac{a}{a+\delta} p_0 = \frac{\left(\left(1 + \frac{a+\delta}{\gamma}\right)^N - 1 \right) \frac{a}{a+\delta}}{\left(\left(1 + \frac{a+\delta}{\gamma}\right)^N - 1 \right) \frac{a}{a+\delta} + 1}.$$

In the limit $\alpha \ll \delta \ll \gamma$, the above expression can be further simplified as,

$$p \approx \frac{N\left(1 + \frac{(N-1)a+\delta}{2}\frac{a}{\gamma}\right)\frac{a}{\gamma}}{N\left(1 + \frac{(N-1)a+\delta}{2}\frac{a}{\gamma}\right)\frac{a}{\gamma} + 1} \approx \frac{N\left(1 + \frac{(N-1)\delta}{2}\frac{a}{\gamma}\right)\frac{a}{\gamma}}{N\left(1 + \frac{(N-1)\delta}{2}\frac{a}{\gamma}\right)\frac{a}{\gamma} + 1}.$$

Several mechanisms can further increase the cooperativity beyond what described by a constant δ
value in the above model. Presence of additional bound sLRC pair may further constrain the
diffusion of the two loci of a dissociated sLRC, leading to further increase of the value of δ. That
is, the value of δ may increase with the number of bound sLRC pairs. The bound sLRC pairs may
physically contact and stabilize each other with an increased γ value. Indeed, we observed non-
monotonic histograms of p_i for some metaLRCs (Extended Data Fig. 7d&e). With a
thermodynamic equilibrium model, one can introduce some binding free energies to describe the
cooperativity as in allosteric effect and a previous cooperative enhancer binding model[40]. For
simplicity in this work we restricted the analyses to the minimal model discussed above.

Note that all the loci belonging to a metaLRC can be clustered as two clumped genomic regions
that are genomically distant (i.e., ~ 100 Mb or further). Subsequent analyses further coarse-grained
the binding state of a metaLRC as $s = 1$ if $(\sum_{i=1}^{N} \sigma_i) \geq 1$ and 0 otherwise.

Reference
46.   Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows–Wheeler
transform. *bioinformatics* **25**, 1754-1760 (2009).
47.   Open2C *et al*. Pairtools: from sequencing data to chromosome contacts. *PLOS*
Computational Biology **20**, e1012164 (2024).
48.   Abdennur, N. & Mirny, L.A. Cooler: scalable storage for Hi-C data and other
genomically labeled arrays. *Bioinformatics* **36**, 311-316 (2020).
49.   Chen, Y.-J. *et al*. Rapid, modular, and cost-effective generation of donor DNA constructs
for CRISPR-based gene knock-in. *Biology Methods and Protocols* **5**, bpaa006 (2020).

50.   Wang, W., Poe, D., Yang, Y., Hyatt, T. & Xing, J. Epithelial-to-mesenchymal transition
proceeds through directional destabilization of multidimensional attractor. *Elife* **11**,
e74866 (2022).
51.   Stirling, D.R. *et al.* CellProfiler 4: improvements in speed, utility and usability. *BMC*
bioinformatics **22**, 433 (2021).
820
821