Transcriptomic and spatial GABAergic neuron subtypes in zona incerta mediate distinct innate behaviors

Mengyue Zhu[1,2 #], Jieqiao Peng[1,2 #], Mi Wang[1,2 #], Shan Lin[1,2 #], Huiying Zhang[3#], Yu Zhou[1,2], Xinyue Dai[1,2], Huiying Zhao[1,2], Yan-qin Yu[1,2], Li Shen[3], Xiao-Ming Li[1,2,4] *, Jiadong Chen[1,2,5] *

[1] Department of Neurobiology, Departments of Neurosurgery and Neurology of the Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou 310058, China

[2] Nanhu Brain-computer Interface Institute, NHC and CAMS Key Laboratory of Medical Neurobiology, MOE Frontier Science Center for Brain Research and Brain-machine Integration, School of Brain Science and Brain Medicine, Zhejiang University, Hangzhou 310058, China

[3] The MOE Key Laboratory of Biosystems Homeostasis & Protection and Zhejiang Provincial Key Laboratory for Cancer Molecular Cell Biology, Life Sciences Institute, Zhejiang University, Hangzhou 310058, China

[4] Center for Brain Science and Brain-Inspired Intelligence, Research Units for Emotion and Emotion Disorders, Chinese Academy of Medical Sciences/ Guangdong–Hong Kong–Macao Greater Bay Area Center for Brain Science and Brain-Inspired Intelligence, Guangzhou 311305, China

[5] Lead Contact.

[#] These authors contributed equally to this work.

*Corresponding Authors: Xiao-Ming Li, Jiadong Chen

23   Email: lixm@zju.edu.cn; jardongchen@zju.edu.cn

Abstract

Understanding the anatomical connection and behaviors of transcriptomic neuron

subtypes is critical to delineating cell type-specific functions in the brain. Here we

integrated single-nucleus transcriptomic sequencing, *in vivo* circuit mapping,

optogenetic and chemogenetic approaches to dissect the molecular identity and

function of heterogeneous GABAergic neuron populations in the zona incerta (ZI)

in mice, a region involved in modulating various behaviors. By microdissecting ZI

for transcriptomic and spatial gene expression analyses, our results revealed two

non-overlapping *Ecel1-* and *Pde11a*-expressing GABAergic neurons with

dominant expression in the rostral and medial zona incerta (ZIr^{Ecel1} and ZIm^{Pde11a}),

respectively. The GABAergic projection from ZIr^{Ecel1} to periaqueductal gray

mediates self-grooming, while the GABAergic projection from ZIm^{Pde11a} to the oral

part of pontine reticular formation promotes transition from sleep to wakefulness.

Together, our results revealed the molecular markers, spatial organization and

specific neuronal circuits of two discrete GABAergic projection neuron populations

in segregated subregions of the ZI that mediate distinct innate behaviors,

advancing our understanding of the functional organization of the brain.

Introduction

The molecular, morphological and anatomical features of neurons imply distinct

neuronal subtypes, axonal projection pattern, and behavioral functions in the

central nervous system[1, 2]. Recent advances in single-cell transcriptomics reveal

cell type specific genes expression and molecular markers of transcriptomic cell

subtypes[3-5]. Extensive efforts have been applied to classify neuron subtypes by

morphological and electrophysiological characteristics[6-8], in conjunction with

single-cell transcriptomic analysis[9-13]. Spatial transcriptomic analysis discovered

diverse cell subtypes and their anatomical organization across the whole brain[14-

16]. In addition, the integrated retrograde or anterograde transsynaptic tracing with

spatial transcriptomic analyses revealed the long-range synaptic connectivity of

neuronal subtypes, connecting transcriptomic clusters with axonal projections at

scale[4, 17-20]. However, the behavioral functions of different transcriptomic neuronal

subtypes are yet to be determined in behaving animals. Therefore, investigating

the anatomical connection and behavior of transcriptomic and spatial neuron

subtypes is important to reveal the functional organization of the brain.

The GABAergic projection neurons establish long-range efferent projections

and may promote efficient information transfer between distant brain networks

compared with the well-studied cortical interneurons [21, 22]. Recent studies revealed

the presence of long-range GABAergic projection neurons in various brain regions

including the prefrontal cortex[23, 24], entorhinal cortex[25], hippocampus[25, 26], and also

top-down afferents from the zona incerta (ZI)[27], basal ganglia[28], and septum[29], but

their sparse distribution makes it difficult to identify the molecular markers for this

heterogeneous population by conventional single-cell RNA sequencing

approach[21, 22]. The zona incerta (ZI), located in the ventral thalamus,

predominantly comprises GABAergic neurons that project to distant brain regions

and mediate various behavioral functions[30, 31]. The heterogeneous neuron

populations in the ZI can be divided into cytoarchitectonic subregions or sectors

along the anteroposterior axis, which form extensive connections with multiple

brain regions and may function as one of the integrative nodes in the brain[30-33].

Recent studies in rodents and non-human primates revealed various behavioral

functions upon stimulating subregions of the ZI [34-42]. However, the lack of

molecular markers makes it difficult to selectively target or to investigate the

function of GABAergic projection neurons in different subregions of the ZI[21, 22].

In the present study, by microdissecting the ZI for single-nucleus

transcriptomic sequencing, we investigated the molecular subtypes and spatial

localization of GABAergic neuron populations in different subregions of the ZI.

Through constructing transgenic mice to label region-specific GABAergic neuron
subtypes in ZI, integrated with *in vivo* circuit mapping, and optogenetic and
chemogenetic manipulation, we revealed two discrete subpopulations of
GABAergic projection neurons in segregated subregions of the ZI that exert distinct
behavioral functions.

Results

Single-nuclei transcriptomic profiling revealed distinct cell clusters and region specific gene expression in the zona incerta

To analyze the molecular diversity of cell types in ZI, we micro-dissected the brain
region according to the distribution of fluorescent reporter from postnatal day 40
(P40) Gad67-GFP transgenic mice in ZI and the anatomy from the reference
mouse brain atlas of Allen Mouse Brain Common Coordinate Framework[33]. We
dissociated the micro-dissected brain tissue into single nuclei suspension and
analyzed the single-nuclei transcriptome of ZI in the adult mouse brain by 10x
genomics. We harvested 15,375 single nuclei from two independent cohorts of 10x
genomics single-nucleus RNA sequencing experiments and obtained similar cell
clusters by reciprocal principle component analysis (RPCA) (Fig. 1a and
Supplementary Fig. 1a-c). The cell clusters were displayed by uniform manifold
approximation and projection (UMAP) and annotated with canonical cell type
specific marker genes. We identified cell clusters including major cell types such
as inhibitory neurons (*Rbfox3*, *Gad2*), excitatory neurons (*Rbfox3*, *Slc17a6*)[43-45],
dopaminergic neurons (*Slc6a3*)[46, 47], microglia (*Cx3cr1*)[44, 48], astrocytes
(*Aldh1a1*)[49], pericytes (*Pdgfrb*)[44], endothelial cells (*Cldn5*)[46, 47], oligodendrocyte
precursor cells (OPCs, *Pdgfra*), differentiation-committed oligodendrocyte
precursors (COPs, *Tns3*), oligodendrocytes (*Mog*) and mature oligodendrocytes
(MOLs, *Apod*) [50](Fig. 1a and Supplementary Fig. 1d).

We further reclustered the neuronal populations and revealed 10 clusters of
glutamatergic neurons (Glu_N1 – Glu_N10) and 12 clusters of GABAergic neurons

(GABA_N1 – GABA_N12) that showed distinct differentially expressed genes

(DEGs) among neuron subtypes in ZI and neighboring brain regions (Fig. 1b, c).

Based on *in situ* hybridization of gene expression in the adult mouse brain from

Allen brain atlas (https://mouse.brain-map.org/)[51], we annotated the anatomical

locations of neuronal clusters and identified marker genes for ZI and neighboring

brain regions in the subcortical forebrain: ZI (*Pax6, Meis2*), thalamus (*Tcf7l2, Lef1*)

located dorsal to ZI, subthalamus nucleus (*Ttc6, Pitx2*) located ventral to ZI,

paraventricular nucleus (*Sim1*, *Ndst4*) located ventral-medial to ZI, and reticular

nucleus (*Six3*, *Pvalb*) located dorsal-lateral to ZI (Supplementary Fig. 2a).

GABAergic neurons in ZI projected to multiple brain regions and were

implicated in various behaviors, however, the molecular profiles of different

GABAergic neuron subtypes remain incompletely understood. Our data revealed

six GABAergic neuron clusters in ZI (GABA_N4 to GABA_N9, Fig. 1d, e;

Supplementary Fig. 2b, c and Supplementary table 1). Differential gene expression

analysis revealed marker genes for different GABAergic neuron clusters that

exhibited different anatomical distribution pattern in ZI, as demonstrated by *in situ*

hybridization of gene expression in the Allen Mouse Brain Atlas

(https://mouse.brain-map.org/)[51]. The GABA_N9 neuron cluster highly expressed

Ecel1 (endothelin converting enzyme like 1). The GABA_N6, GABA_N7 and

GABA_N8 neuron clusters shared differentially expressed genes and highly

expressed *Cdh23* (cadherin related 23), *Pax6* (paired box 6) and *Pde11a*

(phosphodiesterase 11A). The GABA_N4 neuron cluster highly expressed *Ptprk*

(protein tyrosine phosphatase receptor type K) and *Nos1* (nitric oxide synthase 1)

(Fig. 2a and Supplementary Fig. 2c). These results revealed distinct transcriptomic

profiles and candidate marker genes for GABAergic neurons in the ZI by single-

nucleus RNA sequencing analysis.

Candidate genes for GABAergic neuron subtypes in subregions of zona

incerta

We validated the expression and spatial distribution of the above six

GABAergic marker genes in different anatomical subregions of ZI along the

anteroposterior axis by fluorescence *in situ* hybridization (FISH). We found that

Ecel1 exhibited highest distribution in the rostral ZI (ZIr: 59.56 ± 0.93%; ZIm: 14.41

± 1.33%; ZIc: 26.03 ± 1.76%) compared with other marker genes expression in the

ZIr (*Cdh23*: 22.37 ± 1.40%; *Pax6*: 29.20 ± 1.83%; *Nos1*: 22.98 ± 1.68%; *Ptprk*:

25.59 ± 1.67%). The marker gene *Pde11a* exhibited highest and most selective

distribution in the medial ZI (ZIm: 80.49 ± 1.57%; ZIr: 7.873 ± 0.55%; ZIc: 11.63 ±

2.04%) compared with other marker genes expression in the ZIm (*Cdh23*: 54.60 ±

0.89%; *Pax6*: 43.97 ± 1.79%; *Nos1*: 49.20 ± 5.38%). We found that 33.82 ± 6.63%

of the total *Ptprk* positive cells in the ZI were distributed in the caudal ZI, but

showed no significant difference between different subregions of ZI (ZIr: 25.59 ±

1.67%; ZIm: 40.59 ± 6.35%. Fig. 2b and Supplementary Fig. 2d).

We further demonstrated that the density of $Ecel1^+$ cells is relatively higher

than $Pde11a^+$ or $Ptprk^+$ cells in the ZIr ($Ecel1^+$: 1848 ± 82.52 per mm^2; $Pde11a^+$:

134.8 ± 10.60 per mm^2; $Ptprk^+$: 444.9 ± 38.15 per mm^2)*,* and showed few overlap

with *Pde11a* and *Ptprk* ($Ecel1^+$: 410.0 ± 9.539 cells; $Ecel1^+Pde11a^+$: 9.33 ± 1.20

cells; $Ecel1^+Ptprk^+$: 43.33 ± 5.49 cells; $Pde11a^+Ptprk^+$: 4.33 ± 0.88 cells;

$Ecel1^+Pde11a^+Ptprk^+$: 1.67 ± 0.33 cells)*.* The $Pde11a^+$ cells show higher density

in the ZIm ($Ecel1^+$: 85.77 ± 4.066 per mm^2; $Pde11a^+$: 268.2 ± 24.12 per mm^2; $Ptprk^+$:

136.2 ± 21.35 per mm^2) and also show very few overlap with *Ecel1* and *Ptprk*

($Pde11a^+$: 361.0 ± 23.81 cells; $Ecel1^+Pde11a^+$: 1.67 ± 0.67 cells; $Ecel1^+Ptprk^+$:

4.67 ± 0.88 cells; $Pde11a^+Ptprk^+$: 2.67 ± 0.33 cells; $Ecel1^+Pde11a^+Ptprk^+$: 0 ± 0

cells). In contrast, the density of $Ptprk^+$ cells showed little differences with $Ecel1^+$

or $Pde11a^+$ cells in the ZIc ($Ecel1^+$: 317.8 ± 34.96 per mm^2; $Pde11a^+$: 78.27 ± 14.38

per mm^2; $Ptprk^+$: 228.9 ± 42.81 per mm^2. Fig. 2c - e) and $Ptprk^+$ cells rarely overlap

with $Ecel1^+$ or $Pde11a^+$ cells ($Ptprk^+$: 132.3 ± 16.13 cells; $Ecel1^+Pde11a^+$: 3.33 ±

0.33 cells; $Ecel1^+Ptprk^+$: 7.33 ± 0.88 cells; $Pde11a^+Ptprk^+$: 2.00 ± 0.58 cells;

$Ecel1^+Pde11a^+Ptprk^+$: 2.33 ± 0.88 cells). These results demonstrated that the

expression of *Ecel1* and *Pde11a* are non-overlapping across different subregions

of the ZI along the anteroposterior axis (Fig. 2d, e). Together, these results

revealed that *Ecel1-* and *Pde11a-*expressing GABAergic neurons are non-

overlapping and exhibit dominant expression in the rostral and medial ZI,
respectively.

Ecel1-positive neurons in the rostral ZI and Pde11a-positive neurons in the
medial ZI displayed distinct efferent projections

Previous studies showed that GABAergic neurons in the ZI projected to different
brain regions and exhibited various behavioral functions[30, 31]. We hypothesized
that GABAergic projection neurons in different subregions of the ZI may drive
distinct functions. Firstly, we further confirmed that *Ecel1* and *Pde11a* are co-
expressed with GABAergic neuronal marker *Slc32a1* (Vgat, vesicular GABA
transporter) in the ZIr (*Ecel1$^+$Slc32a1$^+$ /Ecel1$^+$:* 96.39 ± 1.19%;
Ecel1$^+$Slc32a1$^+$/Slc32a1$^+$: 59.69 ± 6.28%) and the ZIm
(*Pde11a$^+$Slc32a1$^+$/Pde11a$^+$*: 100 ± 0.00%*; Pde11a$^+$Slc32a1$^+$/Slc32a1$^+$*: 33.92 ±
9.67%, n = 9 brain slices from 3 mice per group. Fig. 3a-c). And for selectively
target of GABAergic projection neuron subtypes in the rostral or medial ZI, we
constructed two transgenic mice to selectively label *Ecel1*-expressing neurons
(Ecel1-Cre) and *Pde11a*-expressing neurons (Pde11a-Cre), respectively (Fig. 3d).
We verified the specificity of transgenic mice by examining the co-expression of
Cre recombinase with *Ecel1* or *Pde11a*. The Cre recombinase expression in
Ecel1-Cre is predominantly distributed in the ZIr (71.95 ± 1.12%, n = 9 brain slices
from 3 mice) and highly co-expressed with *Ecel1* in ZIr (93.87 ± 1.45% *Ecel1$^+$Cre$^+$*
double positive neurons out of Cre$^+$ neurons. 96.41 ± 0.92% *Ecel1$^+$Cre$^+$* double
positive neurons out of *Ecel1$^+$* neurons). In contrast, the expression of Cre
recombinase in Pde11a-Cre mice is predominantly distributed in the ZIm (90.33 ±
2.91%, n = 9 brain slices from 3 mice) and highly co-expressed with *Pde11a* (99.70
± 0.15% *Pde11a$^+$Cre$^+$* double positive neurons out of Cre$^+$ neurons, 98.91 ± 0.22%
Pde11a$^+$Cre$^+$ double positive neurons out of *Pde11a$^+$* neurons. Fig. 3e-i). These
results demonstrated high fidelity of selective labeling of *Ecel1* and *Pde11a-*
expressing neurons by the Ecel1-Cre and Pde11a-Cre transgenic mice.

Next, we examined the axonal projections of ZIr^{Ecel1} and ZIm^{Pde11a} GABAergic
neurons by stereotaxic injection of AAV virus carrying Cre-dependent expression

of membrane-bound GFP (mGFP) and synaptophysin-mRuby (SYP-mRuby) (Fig.

4a, b and Supplementary Fig. 3a, b). We found the axon terminals of ZIr^{Ecel1}

GABAergic projection neurons distributed in several brain regions including

periaqueductal gray (PAG), lateral septal nucleus (LS), preoptic area (POA),

paraventricular nucleus of the thalamus (PVT), reuniens thalamic nucleus (Re),

lateral hypothalamic area (LH), mediodorsal thalamic nucleus (MD) (Fig. 4c-e and

Supplementary Fig. 3c). In contrast, the axon terminals of ZIm^{Pde11a} GABAergic

neurons distributed in brain regions including pontine reticular formation (PRF),

posterior complex of the thalamus (PO), auditory cortex (AuV), anterior pretectal

nucleus, dorsal part (APTD), nucleus of Darkschewitsch (Dk), red nucleus (RN)

and superior colliculus (SC) (Fig. 4c-e and Supplementary Fig. 3c). These results

demonstrated divergent efferent projections of ZIr^{Ecel1} and ZIm^{Pde11a} GABAergic

neurons to distinct downstream brain regions. For example, the periaqueductal

gray (PAG) received extensive axonal projections from ZIr^{Ecel1} neurons (ZIr^{Ecel1}-

PAG pathway), but very few projections from ZIm^{Pde11a} neurons. In contrast,

pontine reticular formation (PRF) received axonal projections predominantly from

ZIm^{Pde11a} (ZIm^{Pde11a}-PRF pathway), but not ZIr^{Ecel1} GABAergic neurons (Fig. 4c, d).

Therefore, the efferent projection of ZIr^{Ecel1} and ZIm^{Pde11a} GABAergic neurons to

different downstream target brain regions may imply their distinct behavioral

functions.

Activation of ZIr^{Ecel1} neurons and the GABAergic projection from ZIr^{Ecel1} to

PAG pathway mediates self-grooming

To investigate the behavioral function of ZIr^{Ecel1} GABAergic projection neurons, we

bilaterally injected AAV virus expressing Cre-dependent channelrhodopsin2-EYFP

(AAV-DIO-ChR2-EYFP) or EGFP as control (AAV-DIO-EGFP) into the ZIr of

Ecel1-Cre mice (Fig. 5a and Supplementary Fig. 4a). We found optogenetic

activation of ChR2-expressing ZIr^{Ecel1} GABAergic neurons at different stimulation

frequencies (5Hz, 10Hz, 20Hz) elicited robust self-grooming behavior compared

with EGFP-expressing control mice (Fig. 5b-d, Supplementary Fig. 4b,c and

Supplementary Video 1 and 2). High frequency optogenetic stimulation (20 Hz)

also increased food intake or gnawing behavior (Supplementary Fig. 5a-d and
Supplementary Video 3-6), consistent with previous findings on optogenetic
activation of GABAergic neurons in the rostral ZI[34]. Self-grooming is reported to be
associated with stress[52, 53], increased anxiety[54-56] or reward behaviors[57, 58]. We
further measured the Ecel1-Cre mice in real time place preference test (RTPP)
and found that optogenetic activation of ZIr[Ecel1] neurons resulted in aversion to the
optogenetic stimulation paired side (Supplementary Fig. 6e-g). These results
indicate that ZIr[Ecel1] neurons mediate self-grooming behaviors could be associated
with negative emotion.

Next, we examined the downstream brain region of ZIr[Ecel1] neurons that
mediated self-grooming behavior. Our results showed the axonal projection
terminals in periaqueductal gray (PAG) from ZIr[Ecel1] GABAergic neurons (Fig. 4c).
The PAG neurons are reported to be an important sensory processing area which
also involved in the regulation of self-grooming[54, 58], different behaviors could be
mediated by distinct neuron subtypes or ensembles of neurons driven by different
descending afferents. To investigate whether the ZIr[Ecel1] to PAG pathway mediate
self-grooming, we bilaterally injected AAV-DIO-ChR2-EYFP or AAV-DIO-EGFP
into the ZIr of Ecel1-Cre mice (Fig. 5h and Supplementary Fig. 7). We first
examined the functional synaptic connection from the ZIr[Ecel1] to PAG pathway by
whole-cell patch-clamp recording of PAG neurons (Fig. 5e). We found that
optogenetic stimulation of axon terminals from ZIr[Ecel1] GABAergic neurons induced
evoked inhibitory postsynaptic currents (IPSCs) in the PAG neurons in acute brain
slices, that could be completely blocked by the $GABA_A$ receptor antagonist
bicuculline (20 μM). Administration of the Na^+ channel inhibitor tetrodotoxin (TTX,
1 μM) blocked the optogenetically evoked IPSCs, while subsequent application of
the K^+ channel blocker 4-aminopyridine (4-AP, 100 μM) restored the evoked IPSCs
(Fig. 5f, g). These results demonstrated functional monosynaptic GABAergic
synaptic inputs from ZIr[Ecel1] GABAergic neurons to the PAG.

In addition, we investigated the behavioral function of the ZIr[Ecel1] to PAG
pathway. We found optogenetic activation of the ChR2-expressing axon terminals

from ZIr^{Ecel1} neurons to the PAG significantly induced increase in self-grooming

time and grooming bouts (Fig. 5i, j). We also examined whether optogenetic

stimulation of axonal projections from ZIr^{Ecel1} neurons to other target brain regions

could induce self-grooming behavior. We found that optogenetic activation of the

ChR2-expressing axon terminals from ZIr^{Ecel1} neurons to other targets including

LS, POA, LH or SC had no discernible impact on either self-grooming time or self-

grooming bouts in ChR2-group compared with EGFP-group (Supplementary Fig.

8). Together, these results demonstrate that activation of the ZIr^{Ecel1} neurons and

the GABAergic projection from ZIr^{Ecel1} to PAG pathway mediates self-grooming

behavior.

ZIr^{Ecel1} neurons bidirectionally regulate self-grooming behavior

To monitor the activity of ZIr^{Ecel1} GABAergic neurons during self-grooming *in vivo,*

we unilaterally injected AAV virus expressing Cre-dependent GCaMP6s and

mCherry (as control) in the ZIr of Ecel1-Cre mice and recorded the calcium activity

by fiber photometry (Fig. 6a, b). The self-grooming behavior can be efficiently

induced by water spray toward the face[52, 58, 59], we found the grooming time

increased significantly compared to that before the water spray treatment (Fig. 6c,

d). The calcium fluorescent signal of ZIr^{Ecel1} neurons increased before the start of

grooming but decreased throughout the entire self-grooming period, and then

returned to baseline after termination of self-grooming during both water spray

induced and spontaneous self-grooming (Fig. 6e-g and Supplementary Fig. 9a-f).

These results suggest that the increased activity of ZIr^{Ecel1} GABAergic neurons

preceded the self-grooming behavior[53, 60].

We further chemogenetically activated or inhibited the ZIr^{Ecel1} neurons by

bilaterally injecting AAV-DIO-hM3Dq-mCherry or AAV-DIO-hM4Di-mCherry (or

AAV-DIO-mCherry as control) into the ZIr of Ecel1-Cre mice (Fig. 7a, e). We found

that chemogenetic activation of ZIr^{Ecel1} neurons significantly increased the self-

grooming time and self-grooming bouts in hM3Dq-expressing mice compared with

the mCherry-expressing control mice (Fig. 7b-d and Supplementary Fig. 9g), but

chemogenetic inhibition of ZIr^{Ecel1} neurons showed no discernable effect on the

spontaneous self-grooming behavior in hM4Di-expressing mice compared with the

mCherry-expressing control mice (Supplementary Fig. 9h and i). To test whether

ZIr^{Ecel1} neurons are required for water spray induced self-grooming behavior, the

mice were i.p. administered with CNO (3 mg/kg) 30 min before water spray toward

the mice face following by 20min self-grooming test (Fig. 7f). We demonstrated

that chemogenetic inhibition of Ecel1 positive neurons in the ZIr significantly

decreased the total self-grooming time and self-grooming bout duration in hM4Di-

expressing mice compared with the mCherry-expressing control mice in water

spray induced self-grooming (Fig. 7g-i and Supplementary Fig. 9k). Moreover, we

further ablated the ZIr^{Ecel1} neurons by bilaterally injected AAV virus carrying Cre-

dependent DTA (AAV-DIO-DTA) or mCherry (AAV-DIO-mCherry) as control into

the ZIr of Ecel1-Cre mice (Fig. 7j and Supplementary Fig. 9l-n). While ablation of

ZIr^{Ecel1} neurons showed no discernable effect on the spontaneous self-grooming

behavior (Supplementary Fig. 9j), we found that ablation of ZIr^{Ecel1} neurons also

significantly reduced the self-grooming time and grooming bouts induced by water

spray in DTA-expressing mice compared with the mCherry-expressing control

mice (Fig. 7k-m). These results indicates that the ZIr^{Ecel1} neurons can

bidirectionally regulate the self-grooming behavior.

Activation of ZIm^{Pde11a} neurons and the GABAergic projection from ZIm^{Pde11a}

to PRF pathway promotes transition from sleep to wakefulness

We next investigated the behavioral function of ZIm^{Pde11a} GABAergic neurons by

injecting AAV-DIO-ChR2-EYFP or control AAV-DIO-EGFP virus into the ZIm of

Pde11a-Cre mice (Supplementary Fig. 10a). Optogenetic activation of ZIm^{Pde11a}

neurons during the awake state of mice did not have significant impact on the

center time or locomotor activity in open field test, or the time spent in open arm in

EPM test, or preference to light stimulation in RTPP test (Supplementary Fig. 10f-

l). Our result showed extensive axonal projections from the ZIm^{Pde11a} neurons to

the oral part of pontine reticular formation (PRF) (Fig. 4c), which is a component

of the ascending reticular activating system that modulates rapid eye movement

sleep (REM) in sleep and wakefulness cycle[61, 62]. To investigate whether ZIm[Pde11a]

GABAergic projection neurons participated in regulating sleep-wakefulness cycle,

we monitored the activity of ZIm[Pde11a] neurons by AAV mediated Cre-dependent

expression of GCaMP6s or mCherry (as control) using fiber photometry. The

phase of sleep-wakefulness cycle was determined by electroencephalogram

(EEG) and electromyogram (EMG) recordings (Fig. 8a, b). We found the ZIm[Pde11a]

neurons are active during REM sleep and wakefulness, the calcium signal

increases during the transition from NREM to REM sleep and decreases during

the transition from REM sleep to wakefulness and wakefulness to NREM sleep

(Fig. 8c-e).

Optogenetic activation of ChR2-expressing ZIm[Pde11a] GABAergic projection

neurons promotes the transition from NREM or REM sleep to wakefulness in

comparison with the EGFP control group (20 Hz, 120 seconds per trial, applied

every 10–15 minutes. Fig. 9a-d and Supplementary Video 7 and 8). The transition

probability from REM sleep to wakefulness and NREM sleep to wakefulness was

significantly increased upon optogenetic activation of the ZIm[Pde11a] GABAergic

projection neurons, along with a complementary decrease of transition probability

from wakefulness to NREM or NREM to REM sleep (Fig. 9e and Supplementary

Fig. 10b-e). We verified the functional synaptic connectivity from ZIm[Pde11a] to PRF

pathway by whole-cell patch-clamp recording of neurons in the PRF (Fig. 9f).

Optogenetic stimulation of the ChR2-expressing axonal terminals from ZIm[Pde11a]

neurons evoked IPSCs in PRF neurons that can be blocked by bicuculline (20 μM)

or TTX (1 μM). The blockade of IPSCs by TTX was restored following application

of 4-AP (100 μM), suggesting monosynaptic GABAergic inputs from ZIm[Pde11a] to

PRF pathway (Fig. 9g, h). We investigated the behavioral function of the ZIm[Pde11a]

to PRF pathway and found optogenetic activation of the GABAergic projections

from ZIm[Pde11a] to the PRF increased the transition probability from NREM to

wakefulness and decreased the transition probability from wakefulness to NREM

or from NREM to REM sleep (Fig. 9i-l and Supplementary Fig. 11). Thus, these

results demonstrate that activation of the ZIm[Pde11a] GABAergic projection neurons

and the ZIm[Pde11a] to PRF pathway promotes transition from sleep to wakefulness.

ZIm^Pde11a neurons bidirectionally regulate sleep-wakefulness cycle

We further chemogenetically activated ZIm^Pde11a neurons by bilaterally injecting
AAV-DIO-hM3Dq-mCherry or AAV-DIO-mCherry as control into the ZIm of
Pde11a-Cre mice (Fig. 10a). We found that chemogenetic activation of ZIm^Pde11a
neurons significantly increased wakefulness and decreased NREM and REM
sleep in hM3Dq-expressing mice following CNO injection compared with saline
injection group during the light phase (Fig. 10b-h). In addition, the number of
episodes during wakefulness and NREM both increases, but the duration of
average NREM episode significantly decreases, suggesting chemogenetic
activation of ZIm^Pde11a neurons increased wakefulness at the expense of sleep time
and sleep quality in the light phase (Fig. 10i, j). Chemogenetic inhibition did not
alter the sleep-wakefulness state in hM4Di-expressing mice or in mCherry-
expressing control mice (Supplementary Fig. 12a-h). To determine whether
ZIm^Pde11a neurons are required for sleep-wakefulness transition, we selectively
ablated the ZIm^Pde11a neurons by bilaterally injected AAV-DIO-DTA or AAV-DIO-
mCherry as control into the ZIm of Pde11a-Cre mice (Fig. 10k) and verified the
efficiency of DTA ablation (Supplementary Fig. 12i-k). We found that ablation of
ZIm^Pde11a neurons significantly decreased wakefulness and increased the NREM
and REM sleep during dark phase (Fig. 10l-q). In addition, the number of episodes
during wakefulness, NREM and REM all increases, but the duration of average
wakefulness episode significantly decreases, suggesting ZIm^Pde11a neurons may
be required for wakefulness in the dark phase (Fig. 10r, s). These results
demonstrate that the ZIm^Pde11a neurons can bidirectionally regulate the sleep and
wakefulness cycle.

We further explored whether ZIr^Ecel1 and ZIm^Pde11a neurons play distinct roles
in regulating self-grooming or sleep-wakefulness cycle behaviors. We found that
ablation of the ZIm^Pde11a neurons by selective expression of DTA had no effects on
the spontaneous or water spray induced self-grooming behaviors compared with
mCherry control group (Supplementary Fig. 13j-m). Additionally, ablation of ZIr^Ecel1
neurons by DTA had no significant impact on the time or episodes of wakefulness,

NREM sleep or REM sleep compared with mCherry-group mice (Supplementary
Fig. 13a-i). These results suggest that these ZIr^{Ecel1} and ZIm^{Pde11a} neurons
differentially regulate self-grooming or sleep-wakefulness cycle behaviors,
respectively.

Taken together, our study revealed the molecular markers and spatial
distribution of GABAergic neuron subtypes in subregions of zona incerta. By *in*
vivo circuit mapping and manipulating the neuronal activity, we further delineated
distinct functions for two discrete GABAergic projection neuron subtypes: the
ZIr^{Ecel1} GABAergic neurons regulate self-grooming through projection to PAG
pathway and ZIm^{Pde11a} GABAergic neurons promote transition from sleep to
wakefulness through projection to PRF pathway. These results demonstrated the
molecular markers, spatial distribution and circuit mechanisms of GABAergic
projection neuron subtypes in the ZI that mediate distinct innate behaviors.

Discussion

Molecular markers and spatial localization of GABAergic projection neuron
subtypes in zona incerta

Despite the unique long-range axonal projection pattern and distinct functions of
GABAergic projection neurons, the lack of specific molecular markers makes it
difficult to selectively target or to investigate the function of GABAergic projection
neurons *in vivo*. By single-nucleus transcriptomic sequencing of the ZI and
quantifying the spatial expression of marker genes of GABAergic neurons in
different subregions along the anteroposterior axis of the ZI, we found two discrete
Ecel1- and *Pde11a*-expressing GABAergic neurons that are non-overlapping and
exhibit dominant expression in the rostral and medial ZI, respectively (Fig. 2d, e).

By *in vivo* circuit mapping, we further demonstrated that ZIr^{Ecel1} and ZIm^{Pde11a}
neurons sent long-range GABAergic efferent projections to segregated distant
brain regions in the forebrain and the midbrain (Fig. 4 and Supplementary Fig. 3c),

implying distinct behavioral functions. We also found the axonal projections from
the ZIm[Pde11a] GABAergic projection neurons to the layer 1 of auditory cortex
(Supplementary Fig. 3c), consistent with recent finding reporting the top-down
GABAergic projection from ZI in regulating neocortical associative fear learning[27].
The establishment and maintenance of specific projections are determined by both
genetic and environmental factors[63-65]. Recent studies showed heterogeneity of
the projection area and potential functions in transcriptomic neuron subtypes[66].
Therefore, the same transcriptomic neuronal subtypes could establish different
efferent projections during development[66]. The diverse behavioral functions of the
ZI are routed in the heterogenous neuron populations, that are possibly generated
from different progenitor cells during development[67, 68]. Future studies are
warranted to elucidate the developmental origin of GABAergic neuron subtypes in
the ZI.

422 Activation of ZIr[Ecel1] GABAergic projection neurons induced self-grooming
423 behaviors may be associated with stress

Previous reports suggested that ZI neurons could receive information from
different sensory modalities including auditory, whisker and visual stimulation[31, 37,]
[69]. By monitoring the calcium dynamics of ZIr[Ecel1] neurons *in vivo*, we found
increased calcium activity of ZIr[Ecel1] neurons before water spray induced self-
grooming, suggesting these neurons could be activated by water spray induced
unpleasant sensory stress[52, 59] (Fig. 6). Chemogenetic inhibition or ablation of
ZIr[Ecel1] neurons impaired the self-grooming behavior induced by water spray but
had little effect on the spontaneous self-grooming behavior (Fig. 7e-m and
Supplementary Fig. 9h-j). Thus, these results suggest that activation of ZIr[Ecel1]
neurons induced self-grooming behaviors could be associated with sensory stress.

Self-grooming is an innate behavior that could be elicited by comfort or
stressful conditions[70]. Our results demonstrated that optogenetic activation of
ZIr[Ecel1] neurons increased the bouts of self-grooming and induced aversion to laser

stimulation paired side (Fig. 5a-d and Supplementary Fig. 6e-g), resembling with

stress-associated self-grooming behavior that often accompanied by negative

emotional state[52, 59]. Previous study showed that activation of somatostatin (SST)-

expressing neurons or calretinin (CR)-expressing neurons in the ZI differentially

modulate anxiety[40]. We found that only subsets of *Ecel1*-expressing neurons co-

expressed *SOM* (*Ecel1$^+$SST$^+$/Ecel1$^+$*, 23.12 ± 1.12%; *Ecel1$^+$SST$^+$/SST$^+$*, 33.93 ±

0.96%), while *Ecel1*-expressing neurons rarely co-expressed *CR* in the rostral ZI

(*Ecel1$^+$CR$^+$/Ecel1$^+$*, 4.10 ± 0.15%; *Ecel1$^+$CR$^+$/CR$^+$*, 13.80 ± 2.38%. Supplementary

Fig. 14). In addition, optogenetic stimulation of ZIr^{Ecel1} neurons did not affect the

center time in open field behavioral test, or time spent in the open arm in elevated

plus maze (EPM) behavioral test (Supplementary Fig. 6a-d), or the time of

investigating the novel object compared with familiar object in novel object

recognition test (Supplementary Fig. 6h, i). Together, these results suggest that

ZIr^{Ecel1} neurons could be a distinct subpopulation of GABAergic projection neurons

that mediated stress-related self-grooming behavior.

Stress may lead to decreased feeding and increased grooming[71, 72]. Previous

studies demonstrated that hypothalamic neuronal circuit can exert scalable control

over feeding and negative emotional states[52]. Our results showed that low

frequency optogenetic stimulation (5Hz) of ZIr^{Ecel1} neurons only promote self-

grooming, while high frequency optogenetic stimulation (20Hz) of ZIr^{Ecel1} neurons

also increased feeding, suggesting ZIr^{Ecel1} neurons may orchestrate behaviors

related to stress response and feeding (Fig. 5b-d, Supplementary Fig. 4b,c and

Supplementary Fig. 5a-d). In addition, we found ablation of ZIr^{Ecel1} neurons

significantly reduced water spray induced self-grooming (Fig. 7j-m), but had no

discernible impact on food intake or the body weight (Supplementary Fig. 5e-h).

These results demonstrate that ZIr^{Ecel1} neurons are required for sensory stress

related self-grooming.

ZIm^{Pde11a} GABAergic projection neurons participate in regulating sleep-

wakefulness cycle

Extensive GABAergic projections from the diencephalon including ZI, reticular
thalamus and hypothalamus to pontine reticular formation (PRF) have been
identified by retrograde tracing approaches[73]. Endogenous GABA levels were
increased during wakefulness and application of GABA$_A$ receptor agonists in PRF
promotes wakefulness[74-76]. We demonstrated that activation of ZIm^{Pde11a} neurons
and the GABAergic projection from ZIm^{Pde11a} to PRF promote the transition from
sleep to wakefulness (Fig. 9). We found increased Ca^{2+} activity of ZIm^{Pde11a}
neurons during REM sleep and in wakefulness using fiber photometry recordings
(Fig. 8). Recent studies using *in vivo* micro-endoscopic imaging showed that
GABAergic neurons in the ZI demonstrated heterogenous activity in sleep-
wakefulness cycle, including neurons showing maximal Ca^{2+} activity during REM
sleep (REM-active), neurons showing maximal Ca^{2+} activity during wakefulness
(Wake-active), or neurons showing Ca^{2+} activity in both REM sleep and
wakefulness (Wake/REM-active)[77].

Activation of REM-active neurons promote REM sleep, often at the expense
of NREM sleep[78, 79], while activation of Wake/REM-active neurons (arousal
neurons) promote wakefulness[80-82]. The ZIm^{Pde11a} neurons may include both REM-
active and Wake/REM-active neurons as demonstrated by fiber photometry
recording (Fig. 8). Therefore, optogenetic stimulation of ZIm^{Pde11a} neurons
promotes the transition from sleep to wakefulness could be mediated by the
activation of the Wake/REM-active ZIm^{Pde11a} neurons. Additionally, optogenetic
activation of ZIm^{Pde11a} neurons decreased the investigating time of both familiar
and novel object compared with control group in the novel object recognition test,
(Supplementary Fig. 10m,n), suggesting a potential shift to alert state during
wakefulness[83, 84].

The zona incerta has been a target for chronic deep brain stimulation in
treating Parkinson's disease (PD)[85, 86]. Previous studies reported activation of
Lhx6-expressing GABAergic neurons subtypes induced transition from
wakefulness to sleep[35]. Our results showed that *Pde11a*-positive neurons rarely
co-expressed with *Lhx6* (*Pde11a*+*Lhx6*+/*Pde11a*+, 2.23 ± 1.18%;

Pde11a⁺Lhx6⁺/Lhx6⁺, 1.38 ± 0.69%, n = 6 brain slices from 3 mice. Supplementary

Fig. 15), suggesting ZIm^{Pde11a} is a distinct GABAergic projection neuron

subpopulation that promote the transition from sleep to wakefulness. These

findings demonstrate that *Pde11a⁺* and *Lhx6⁺* neuronal subpopulations in the ZIm

may differentially regulate sleep-wakefulness cycle. Intriguingly, sleep

abnormalities are reported in Parkinson's patients and in rodent models of PD[87, 88],

it is plausible that chronic deep brain stimulation of ZI may help improve sleep-

related symptoms in Parkinson's disease. Future studies are warranted to further

elucidate the subtype and the exact role of GABAergic neurons in regulating sleep

disturbances.

In summary, our results revealed the spatial distribution of two discrete

populations of GABAergic projection neurons in segregated subregions of the ZI.

By constructing transgenic mice to selectively target the ZIr^{Ecel1} and ZIm^{Pde11a}

GABAergic projection neurons, we further demonstrated their efferent axonal

projection to non-overlapping distant brain regions and mediate distinct innate

behavioral functions. These findings uncovered the molecular markers, spatial

organization and functions for GABAergic projection neuron subtypes in the zona

incerta that play distinct roles in regulating self-grooming and sleep-wakefulness

cycle.

Methods

Methods

Animals

All experiments were conducted in accordance with the guidelines of Zhejiang

University (ZJU) and were approved by the Animal Advisory Committee at ZJU.

Wild-type C57BL/6J (JAX# Stock No. 000664) and transgenic GAD67-GFP mice

(JAX# Stock No. 007677) were obtained from Jackson Laboratories. Transgenic

mice Ecel1-Cre and Pde11a-Cre (C57/BL6 background) were generated by

CRISPR/Cas9 technology at the Shanghai Model Organisms Center (China) or

Biocytogen (Beijing) company. All mice were bred and housed at the temperature

(23 ± 1 °C) and humidity (40%) controlled animal facility at ZJU under a 12 h
light/dark cycle with light on at zeitgeber time 0 (ZT 0; 7:00), food and water were
available ad libitum. Only adult (>8 weeks old) mice of normal appearance and
weight were used for behavioral tests. Littermate mice were split into random
groups before AAV viral injection.

Viruses

AAV2/9-DIO-mGFP-Synaptophysin-mRuby (titer: 8.9×10^{12} genomic copies per ml),
AAV2/9-EF1α-DIO-hChR2-EYFP (titer: 2×10^{12} genomic copies per ml) were
purchased from Taitool Bioscience, China. AAV2/9-hSyn-DIO-EGFP (titer:
1.44×10^{12} genomic copies per ml) were purchased from WZ Bioscience, China.
AAV2/9-EF1α-DIO-GCamP6s (titer: 8.35×10^{12} genomic copies per ml). AAV2/9-
CAG-DIO-DTA (titer: 6.13×10^{12} genomic copies per ml). AAV2/9-DIO-mCherry
(titer: 5.48×10^{12} genomic copies per ml) were purchased from BrainVTA
Bioscience, China. AAV2/9-hSyn-DIO-hM3Dq-mCherry (titer: 3.27×10^{13} genomic
copies per ml). AAV2/9-hSyn-DIO-hM4Di-mCherry (titer: 8.53×10^{13} genomic
copies per ml) were purchased from Sunbio Medical Biotechnology Company,
China.

Stereotaxic injection

Adult mice were anesthetized with pentobarbital sodium solution (50 mg/kg) via an
intraperitoneal (i.p.) injection. Mice were placed on a stereotaxic frame (RWD Life
Science, 68046) and anesthesia were maintained with 1%-1.5% isoflurane during
surgery. Injection of AAV virus was performed with a manual viral syringe pump
connected to glass micropipette with a 10-50 μm diameter tip (Drummond
Scientific Company, Wiretrol II). After each injection, the syringe was left *in situ* for
an additional 10 min and then withdrawn slowly. Mice were randomly assigned to
experimental and control groups before virus injection.

To map the axonal projection pattern of GABAergic neurons in the rostral or
medial zona incerta, anterograde tracing virus AAV2/9-DIO-mGFP-
Synaptophysin-mRuby (80 nl. nl: nano-liter) was unilaterally injected into the ZIr
(anteroposterior (AP): −1.00 mm, mediolateral (ML): −0.60 mm, dorsoventral (DV):
−4.45 mm) of Ecel1-Cre mice or the ZIm (AP: −2.30 mm, ML: −1.50 mm, DV: −4.20
mm) of Pde11a-Cre mice. Three weeks after injection, mice were sacrificed and
brain sections were collected for fluorescent imaging. The projection pattern was
shown as the percentage of total relative fluorescent density occupied by different
downstream brain areas.

The neuronal activity of GABAergic neurons was monitored by in *vivo* fiber
photometry recording. AAV2/9-EF1α-DIO-GCaMP6s and AAV2/9-DIO-mCherry
were injected as a mixture at 2:1 ratio (80 nl) into the ZIr of Ecel1-Cre mice or ZIm
of Pde11a-Cre mice. Three weeks after injection, optical fibers (200 μm O.D., 0.37
NA, Inper, Hangzhou, China) were implanted above the ZIr in Ecel1-Cre mice or
ZIm in Pde11a-Cre mice, respectively.

To activate GABAergic neuron subtypes by in *vivo* optogenetic stimulation,
AAV2/9-EF1α-DIO-hChR2-EYFP or AAV2/9-hSyn-DIO-EGFP (80 nl) were
bilaterally injected into the ZIr of Ecel1-Cre mice or unilaterally injected into the
ZIm of Pde11a-Cre mice. Three weeks after injection, optical fibers were implanted
at a 10° angle relative to the vertical plane above ZIr bilaterally (AP: −1.00 mm;
ML: ±1.35 mm; DV: −4.13 mm) in Ecel1-Cre mice. The optical fibers were
unilaterally implanted 100 μm above viral injection coordinates of the ZIm in
Pde11a-Cre mice. The optical fibers were implanted at a 10° angle relative to the
vertical plane above PAG bilaterally in Ecel1-Cre mice (AP: −4.40 mm; ML: ±1.00
mm; DV: −2.15 mm) to optogenetically stimulate axon terminals from ZIr to PAG.
The optical fibers were unilaterally implanted into the PRF in Pde11a-Cre mice
(AP: −4.4 mm; ML: ±0.85 mm; DV: −4.6 mm) to optogenetically stimulate axon
terminals from ZIm to PRF.

To chemogenetic manipulation of GABAergic neuron subtypes, AAV2/9-hSyn-
DIO-hM3Dq-mCherry or AAV2/9-hSyn-DIO-hM4Di-mCherry (80 nl) were

bilaterally injected into the ZIr of Ecel1-Cre mice or ZIm of Pde11a-Cre mice. For

the DTA mediated ablation experiments, AAV2/9-CAG-DIO-DTA (80nl) were

bilaterally injected into the ZIr of Ecel1-Cre mice or ZIm of Pde11a-Cre mice.

Dual color fiber photometry

Calcium signals were recorded using a dual-color fiber photometry system

(Thinkertech, Nanjing Bioscience Inc.). GCamP6s was excitated by 488 channels

to record changes of calcium activity, and mCherry fluorescent signal was excited

by 570 nm channel as control to exclude disturbances of fluorescent signals

caused by artificial factors. The laser power at the tip of the optical fiber was

adjusted to be 30 µW for 488 nm excitation channel and 10 µW for the control

channel (570 nm) to decrease laser bleaching of fluorescent signals. Averaged

traces of Ca^{2+} fluorescent signal changes and heatmap were plotted using the

MATLAB program provided by ThinkerTech Nanjing Bioscience. Adult male mice

were used for dual color fiber photometry.

***In vivo* optogenetic manipulation**

Mice were recovered for at least 1 week after optical fiber implantation surgery.

For optogenetic manipulation during the behavioral assessments, pulses of 465

nm laser stimulation were set by laser stimulator (INPER-B1-465, Inper, Hangzhou,

China). Before behavioral tests, mice were habituated to the environments for at

least 30 min after connection to a laser stimulator. The laser power at the tip of the

optical fiber was adjusted to 5-10 mW for the optogenetic stimulation (10 ms pulses

at 20 Hz or 5Hz). Adult male mice were used for optogenetic manipulation.

Electrophysiology

Mice were anesthetized with pentobarbital sodium solution (50 mg/kg) via
intraperitoneal injection (i.p.) and then the brains were removed and placed into
ice-cold oxygenated artificial cerebrospinal fluid (ACSF) containing (in mM) 110
Choline chloride, 2.5 KCl, 0.5 $CaCl_2$, 7 $MgCl_2$, 1.3 NaH_2PO_4, 25 $NaHCO_3$, 20
Glucose, 1.3 Na-Ascorbate, 0.6 Na-Pyruvate. Coronal brain slice sections (300 μm
thickness) including the periaqueductal gray (PAG) or sagittal sections (250 μm
thickness) including the pontine reticular formation (PRF) region were made on a
vibratome (VT1200S, Leica) in ice-cold ACSF. The brain slices were incubated at
30 ± 2 ℃ for 0.5 hour and recovered for 1 hour at room temperature in a recording
solution containing (in mM) 125 NaCl, 2.5 KCl, 2 $CaCl_2$, 7 $MgCl_2$, 1.3 NaH_2PO_4, 25
$NaHCO_3$, 10 Glucose, 1.3 Na-Ascorbate, 0.6 Na-Pyruvate. After recovering,
individual brain slices were transferred to the recording chamber and continuously
perfused (2 – 4 ml/min) with ACSF containing (in mM) 125 NaCl, 2.5 KCl, 2 $CaCl_2$,
7 $MgCl_2$, 1.3 NaH_2PO_4, 25 $NaHCO_3$, 10 Glucose. The incubating and recording
ACSF solutions were continuously bubbled with 95% O_2/5% CO_2.

To test the monosynaptic and inhibitory connection of the ZIr to PAG or the ZIm
to PRF, light-evoked inhibitory postsynaptic currents (eIPSCs) were recorded
using pipettes (3 - 5 MΩ) filled with inner solution containing 133 $CsMeSO_4$, 7 CsCl,
10 HEPES, 4 $MgCl_2$, 0.5 EGTA, 3 CsOH, 4 Na_2ATP, 0.4 Na_4GTP, 5 QX-314 (PH
7.35; Osmolarity 310 ± 10 mOsm). And TTX (1 μM, Absin, China) was used to
block the sodium channel and followed by the 4-AP (100 μM) to confirm the
monosynaptic postsynaptic currents following light activation. Bicuculine (20 μM,
Abcam, UK) was bath applied to validate whether the recorded IPSCs were
mediated by GABA receptors. Whole-cell patch-clamp recordings were conducted
with Multiclamp 700B amplifier (Axon) controlled by DigiData 1550B digitizer and
analyzed with pClamp (Clampex, Clampfit, v10.7). Both male and female mice
were used for electrophysiology.

Nuclei isolation and single-nuclei RNA sequencing analysis

Sagittal brain slice sections (600 µm) from Gad67-GFP knock-in mice at postnatal

day 40 (P40) were made with a vibratome as described above. The zona incerta

region indicated by expression of fluorescent reporters were microdissected under

a fluorescence stereomicroscope (Leica VT1200S, Germany) and collected in ice-

cold artificial cerebrospinal fluid (ACSF). Two independent cohort of micro-

dissected zona incerta brain slice sections (each cohort were collected from 3 male

mice and 3 female mice) were harvested for single-nuclei RNA sequencing.

Nuclei isolation from micro-dissected tissue was performed using the following

procedure: the collected tissue was homogenized in ice-cold RNAase-free

homogenization buffer[89] containing: 0.32 M sucrose, 5 mM $CaCl_2$, 3 mM $MgAC_2$,

0.1 mM EDTA (Invitrogen, AM9260G), 10 mM Tris-HCl pH 7.6 (Invitrogen, 15568-

025), 0.4 U/ul Recombinant RNA inhibitor (Takara, 2312A), 0.1 mM PMSF (Roche,

10837091001), 0.1 mM β-mercaptoethanol, 1% BSA, 0.01% 10% NP40 (Thermo,

85124) in UltraPure distilled-water. Subsequently 1.5 mL 0.32 M Sucrose solution

containing 0.32 M sucrose, 5 mM $CaCl_2$, 3 mM $MgAC_2$, 0.1 mM EDTA, 10 mM

Tris-HCl pH 7.6, 0.4 U/ul Recombinant RNA inhibitor, 0.1 mM β-mercaptoethanol,

1% BSA in UltraPure distilled-water were added to the homogenates, mixed well,

and centrifuged by 900 g for 10 min. After removing the supernatant, another 1.5

mL 0.32 M sucrose solution was added to the tube, resuspended, filtered by 35

µm cell-strainer (FALCON, 352235), and then centrifuged at 900 g for 8 min. The

precipitate was collected and resuspended in 1.5 mL 0.32 M sucrose solution and

was centrifuged at 600 g for two times. The precipitate was resuspended in 0.32

M sucrose solution and diluted with an equal volume of 50% OptiPrep density

gradient medium (60% OptiPrep density gradient medium, 0.32 M sucrose

solution) to give a final concentration of 25% medium solution. Slowly adding the

29% OptiPrep density gradient medium (60% OptiPrep density gradient medium,

0.32 M sucrose solution) under the 25% medium solution without disrupting the

layers in a 1.5 mL centrifuge tube and centrifuged by 3000 g for 20 min[90]. After the

removal of the supernatant, the nuclei pellet was dissolved in 100 µl PBS (Gibco,

10010-031). Nuclei density was analyzed using a cell counting plate (WATSON,

177-112C). Single nuclei capture was performed on 10X Genomics Single-Cell 3'

system with target capture of 8000 nuclei per sample. Single-nuclei libraries from

individual samples were pulled and sequenced on the Illumina HiSeq X Ten

machine. The 10x nuclei capture and library preparation protocol was performed

according to the manufacturer's recommended protocol (10X Genomics).

Sequencing reads were aligned to mouse reference genome mm10 with

CellRanger (version 6.0.2). Low abundant genes (expressed in less than 3 cells)

and cells of potentially low quality (nFeature (number of detected genes) < 1000

or nFeature > 6,000, or counts > 25,000, or percentage of mitochondrial genes >

5%) were removed from downstream analysis. The generated gene expression

matrix files arising from two batches of experiments were further analyzed using

Seurat v4 package. Briefly, SCTransform normalization was performed and RPCA

was used to integrate data. 3,000 variable genes were identified by

FindVariableGenes and selected to function RunPCA for the principal component

analysis (PCA). Next, Uniform Manifold Approximation and Projection (UMAP) was

performed to reduce variation to two dimension and FindClusters was used for

unsupervised clustering of cells by setting resolution to 0.6. Subsequently,

canonical cell type-specific markers were used to annotate major cell types:

neurons, oligodendrocytes, endothelial cells, astrocytes, oligodendrocyte

progenitor cells (OPCs), pericytes, microglia, and differentiation-committed

oligodendrocyte precursors (COPs). Furthermore, we reclustered excitatory

neurons and inhibitory neurons into ten and twelve subclusters, respectively. The

genes differentially expressed in cells of subclusters were analyzed using

FindMarkers function and used the two-sided Wilcoxon Rank-Sum test (default).

Those which with p value < 0.05 were selected as differentially expressed genes

(DEGs) candidates. Heatmap showing differentially expressed genes was

generated by ggplot2 package. The regional information of neuronal subclusters

were determined by the expression of DEGs in brain slices from Allen Mouse Brain

Atlas (https://mouse.brain-map.org).

Histology and immunofluorescent staining

Mice were deeply anesthetized with an intraperitoneal injection of pentobarbital

sodium solution (50 mg/kg) and perfused transcardially with 0.1M phosphate-

buffered saline (PBS, pH 7.4) followed by 4% paraformaldehyde (PFA). Brain

tissues were collected and postfixed in 4% PFA for 6-8 hours at 4 °C. Fixed brains

were then rehydrated with 28-30% sucrose in PBS at 4 °C for 48 hours.

Subsequently, the brains were embedded with Optimal Cutting Temperature

compound and were cut into 40 μm or 16 μm coronal sections using cryomicrotome

(Thermo Scientific, CryoStar NX50). The brain slice sections were stored at -80 °C

or in 0.01% sodium azide in PBS at 4 °C.

For immunofluorescent staining, brain sections were washed with 0.1M PBS for

3 times, 10 min each. Then sections were incubated with blocking buffer containing

10% donkey serum albumin (Jackson ImmunoResearch, USA) dissolved in 0.1%

PBST (0.1% Triton X-100 in PBS) at room temperature for 1 hour. Sections were

then incubated with primary antibody at 4 °C overnight. After washing in PBS,

sections were incubated with secondary antibodies (1:1000, Invitrogen, USA) at

room temperature for 1 hour. Finally, sections were mounted after counterstained

with nuclear dye 4,6-diamidino-2-phenylindole (DAPI, Beyotime, China). All

antibodies were titrated for working solution with blocking buffer. Primary antibody

included anti-Cre recombinase (1:500, Synaptic Systems 257004, Germany) and

anti-Calretinin (1:100, Swant CR6B3, Switzerland). Fluorescent images were

taken with a confocal microscope (IX83-FV3000, Olympus, Japan) and a Virtual

Slide Microscope VS120 (Olympus, Japan).

Fluorescence *in situ* hybridization (FISH) was performed according to the

manufacture's protocol using RNAscope multiplex fluorescent reagent kit and

following designed probes were used: RNAscope® Probe-Mm-Cdh23-C2,

RNAscope® Probe-Mm-Ecel1-C3, RNAscope® Probe-Mm-Ptprk-C1, RNAscope®

Probe-Mm-Pde11a-C1, RNAscope® Probe-Mm-Pax6-C2, RNAscope® Probe-

Mm-Nos1-C3, RNAscope® Probe-Mm-Slc32a1-C4, RNAscope® Probe-Mm-

Lhx6-C2, RNAscope® Probe-Mm-SST-C4 (Advanced Cell Diagnostics, USA). The

density of marker genes was calculated as the cell numbers per μm2. The

distribution of marker genes was shown as the percentage of positive neuron
numbers in different brain areas. Both male and female mice were used for
immunofluorescent staining.

Behavioral analysis

Self-grooming measurement. Mouse self-grooming has a complex sequenced
structure that consists of several syntactic chains which beginning from elliptical
bilateral paw stroke (paw and nose grooming) to unilateral strokes from the
mystacial vibrissae to face following by bilateral head grooming, and finally to body
licking. A 350 mm × 350 mm square box was used for the self-grooming test. Mice
were placed in the chamber to acclimate for at least 10 min before the behavioral
tests. For the optogenetic manipulation, mice were placed in the center of the
apparatus and the behavioral assay was performed with a 3-min Laser OFF, 3-min
Laser ON and 3-min Laser OFF administration pattern. In the self-grooming test
when optogenetic activation of Ecel1 positive neurons in the ZIr, different
frequencies (1Hz, 5Hz, 10Hz and 20 Hz) were applied to ZIr. Self-grooming
behavior was mort robustly induced when optogenetic stimulating Ecel1$^+$ neurons
in the ZIr at 5Hz, we therefore conducted the following experiments at 5Hz. For
chemogenetic manipulation, mice was intraperitoneal administrated with
clozapine-N-oxide (CNO, Sigma-Aldrich; 1 mg/kg in saline for hM3Dq-mediated
activation and 3 mg/kg in saline for hM4Di-mediated inhibition) 30min before self-
grooming test. For water spray in chemogenetic manipulation and DTA ablation
experiments, water spray was started by spraying with water toward the face of
mice 4 times with a sprayer and following with a 20min recording. The total time
spent in self-grooming, self-grooming bouts and duration of self-grooming per bout
were measured during the following 20min.

Open field test. A 450 mm × 450 mm square box was used for the open-field test.
Mice were placed in the experimental room to acclimate for at least 30 min before
the behavioral tests. For the optogenetic manipulation, mice were placed in the

center of the apparatus and a 12-min behavioral assay was performed with a 3-
min Laser OFF and 3-min Laser ON administration pattern. The open field
chamber was cleaned with 75% ethanol between each trial and the moving traces
were tracked by ANY-Maze software. Total distance traveled and time spent in the
center zone were analyzed.

Elevated plus maze (EPM). The EPM apparatus consisted of a central square (50
mm × 50 mm), two open arms without walls (300mm length) and two closed arms
with walls (300mm length, 200mm height). For the optogenetic manipulation, mice
were placed in the center of the EPM facing closed arm and a 12-min behavioral
assay was performed with a 3-min Laser OFF and 3-min Laser ON administration
pattern. Moving traces were tracked by ANY-Maze software. Time spent in the
open arm were analyzed.

Real time place preference or aversion (RTPP). A two-chamber apparatus (500
mm × 250 mm) was used for the test. The mice were allowed to move freely in the
apparatus for 10 minutes on the first day. The mice with over 80% initial place
preference were excluded from the experiment. On the following two days, a 465
nm laser stimulation was delivered when mice entered the laser-coupled side for
a 10min real-time laser coupling test. Moving traces were tracked by ANY-Maze
software. The percentage of time in the laser-coupling side and the preference
index were analyzed.

Food intake analysis. For the food intake test, mice had ad libitum access to chow
in the home cage both prior to and following food intake trials. Each mouse was
housed individually and its intake of normal and high-fat food was measured
separately over a 24-hour period. For the optogenetic manipulation, mice were
connected to a 473 nm blue laser and transferred to a transparent cage for 1-hour
to habituate to the testing environment. Initially, a 10-min food intake test was
conducted without any laser stimulation, followed by a 10 min trial with laser
stimulation. For the 10-min trial, a 473-nm blue laser light with a pulse duration of
10-ms and varying frequencies (1Hz, 5Hz, 10Hz and 20 Hz) were applied to zona
incerta, after which a 30-minute period without stimulation ensued. For the 4 ×10-

min trial, a 10-min session of light stimulation (10 ms, 20 Hz) was followed by 30
min interval without stimulation, repeated four times[34].

Novel object recognition test. The task was conducted under identical
environmental conditions to those of the open-field test, consist of a training phase
(also known as the familiarization phase) followed by a testing phase, with a 1-
hour interval. During the training phase, mice were exposed to two objects that
were identical in terms of shape, color, and odor. These objects were positioned
close to the corners of one wall within the testing arena. Each mouse was placed
in the center of the arena and given 10 minutes to explore the objects. In the
subsequent test phase, one object was a duplicate of the one used in the training
phase, while the other was a new object. Each mouse was again placed in the
arena and optical stimulation was applied during the test phase, which lasted 5
minutes. The arrangement of objects during the test phase was randomly
alternated between the animals within each group. The durations of time spent by
the mice exploring both the familiar and the novel objects were measured.

Adult male mice were used for all the behavioral analysis above.

EEG-EMG recordings and analysis

Electroencephalogram (EEG) recordings were made from three screws on top of
the left and right cortex, at AP –3.5 mm, ML 2.5 mm and AP +1.0 mm, ML ±1.5
mm, respectively. For electromyogram (EMG) recording, two EMG electrodes
(P01965, CALMONT, USA) were inserted into the neck musculature and a
reference screw was implanted into the skull above the right cortex. Insulated leads
from the EEG and EMG electrodes were soldered to a 2 × 3 pin header, which was
secured to the skull by dental cement. For EEG and EMG recordings with fiber
photometry recording or optogenetic manipulation, optical fiber was placed in the
same surgery for EEG and EMG implants. After surgery, mice were allowed to
recover for at least 1week before experiments. Experimental and control animals
were subjected to exactly the same surgical and behavioral manipulations. All the
implanted mice were housed in a 12-h dark - 12-h light cycle (Lights on between

07:00 (ZT0) and 19:00 (ZT12)). For optogenetic activation of Pde11a positive
neurons in ZIm, each trial consisted of a 20 Hz pulse train (10 ms per pulse) lasting
for 120 s and inter-trial interval was chosen randomly between 10 or 15 min. For
chemogenetic activation or inhibition of Pde11a positive neurons in ZIm, we used
the virus-transfected mice with saline as control compared to mice with CNO (1
mg/kg in saline) intraperitoneal injection. The following injection protocol was used
for analysis of all mice injected with AAV virus carrying cre-dependent expression
of hM3Dq or hM4Di: on day 1, we performed two consecutive saline injections at
12-h intervals at ZT5 (12:00) and ZT17 (24:00) and CNO at the same times on day
2. For DTA-mediated ablation of Pde11a positive neurons in ZIm, we recorded the
virus-transfected mice for 24hr. The EEG and EMG signals were recorded and
amplified using PowerLab system (AD Instruments, Australia). NREM, REM and
wake states were analyzed for each 5-s epoch by LabChart8 (wake:
desynchronized EEG and high EMG activity; NREM sleep: synchronized EEG with
high power at 0.5–4 Hz and low EMG activity; REM sleep: desynchronized EEG
with high power at theta frequencies (6–9 Hz) and low EMG activity. Experimenters
were blinded to all training and behavioral assessments. Adult male mice were
used for the EEG-EMG recording experiments.

Quantification and statistical analysis

Statistical analysis was performed using Graph-Pad Prism 6, Matlab 2019a. The
following statistical tests were used: two-tailed t-test and one-way analysis of
variance (ANOVA). Data are presented as mean ± SEM. The number of cells,
animals or experimental replicates are indicated in the figure legends. The animals
were randomly assigned into experimental and control groups. All behavioral
experiments and data analyses were conducted blindly.

Data availability

The raw sequence data reported in this paper have been deposited in the Genome

Sequence Archive in National Genomics Data Center, China National Center for

Bioinformation, Beijing Institute of Genomics, Chinese Academy of Sciences

(GSA: CRA017157) that are publicly accessible at https://ngdc.cncb.ac.cn/gsa.

Source data are provided with this paper.

References

1.   Zeng, H. & Sanes, J.R. Neuronal cell-type classification: challenges,
opportunities and the path forward. *Nat Rev Neurosci* **18**, 530-546 (2017).
2.   Zeng, H. What is a cell type and how to define it? *Cell* **185**, 2739-2755 (2022).
3.   Yao, Z.*, et al.* A taxonomy of transcriptomic cell types across the isocortex and
hippocampal formation. *Cell* **184**, 3222-3241 e3226 (2021).
4.   Tasic, B.*, et al.* Shared and distinct transcriptomic cell types across neocortical
areas. *Nature* **563**, 72-78 (2018).
5.   Yao, Z.*, et al.* A high-resolution transcriptomic and spatial atlas of cell types in
the whole mouse brain. *Nature* **624**, 317-332 (2023).
6.   DeFelipe, J.*, et al.* New insights into the classification and nomenclature of
cortical GABAergic interneurons. *Nat Rev Neurosci* **14**, 202-216 (2013).
7.   Gouwens, N.W.*, et al.* Classification of electrophysiological and morphological
neuron types in the mouse visual cortex. *Nature neuroscience* **22**, 1182-1195 (2019).
8.   Gao, L.*, et al.* Single-neuron analysis of dendrites and axons reveals the
network organization in mouse prefrontal cortex. *Nat Neurosci* **26**, 1111-1126 (2023).
9.   Cadwell, C.R.*, et al.* Electrophysiological, transcriptomic and morphologic
profiling of single neurons using Patch-seq. *Nat Biotechnol* **34**, 199-203 (2016).
10.   Fuzik, J.*, et al.* Integration of electrophysiological recordings with single-cell
RNA-seq data identifies neuronal subtypes. *Nat Biotechnol* **34**, 175-183 (2016).
11.   Gouwens, N.W.*, et al.* Integrated Morphoelectric and Transcriptomic
Classification of Cortical GABAergic Cells. *Cell* **183**, 935-953 e919 (2020).
12.   Scala, F.*, et al.* Phenotypic variation of transcriptomic cell types in mouse
motor cortex. *Nature* **598**, 144-150 (2021).
13.   Kalmbach, B.E.*, et al.* Signature morpho-electric, transcriptomic, and dendritic
properties of human layer 5 neocortical pyramidal neurons. *Neuron* **109**, 2914-2927
e2915 (2021).
14.   Zhang, M.*, et al.* Spatially resolved cell atlas of the mouse primary motor cortex
by MERFISH. *Nature* **598**, 137-143 (2021).
15.   Chen, A.*, et al.* Spatiotemporal transcriptomic atlas of mouse organogenesis
using DNA nanoball-patterned arrays. *Cell* **185**, 1777-1792 e1721 (2022).
16.   Zhang, M.*, et al.* Molecularly defined and spatially resolved cell atlas of the
whole mouse brain. *Nature* **624**, 343-354 (2023).

17.   Yao, S.*, et al.* A whole-brain monosynaptic input connectome to neuron classes
in mouse visual cortex. *Nat Neurosci* **26**, 350-364 (2023).
18.   Zhang, Z.*, et al.* Epigenomic diversity of cortical projection neurons in the
mouse brain. *Nature* **598**, 167-173 (2021).
19.   Huang, L.*, et al.* BRICseq Bridges Brain-wide Interregional Connectivity to
Neural Activity and Gene Expression in Single Animals. *Cell* **183**, 2040 (2020).
20.   Sun, Y.C.,   *et al.* Integrating barcoded neuroanatomy with spatial
transcriptional profiling enables identification of gene correlates of projections. *Nat
Neurosci* **24**, 873-885 (2021).
21.   Caputi, A., Melzer, S., Michael, M. & Monyer, H. The long and short of GABAergic
neurons. *Curr Opin Neurobiol* **23**, 179-186 (2013).
22.   Melzer, S. & Monyer, H. Diversity and function of corticopetal and corticofugal
GABAergic projection neurons. *Nat Rev Neurosci* **21**, 499-515 (2020).
23.   Malik, R., Li, Y., Schamiloglu, S. & Sohal, V.S. Top-down control of hippocampal
signal-to-noise by prefrontal long-range inhibition. *Cell* **185**, 1602-1617 e1617
(2022).
24.   Lee, A.T., Vogt, D., Rubenstein, J.L. & Sohal, V.S. A class of GABAergic neurons
in the prefrontal cortex sends long-range projections to the nucleus accumbens and
elicits acute avoidance behavior. *J Neurosci* **34**, 11519-11525 (2014).
25.   Melzer, S.*, et al.* Long-range-projecting GABAergic neurons modulate
inhibition in hippocampus and entorhinal cortex. *Science* **335**, 1506-1510 (2012).
26.   Szabo, G.G.*, et al.* Ripple-selective GABAergic projection cells in the
hippocampus. *Neuron* **110**, 1959-1977 e1959 (2022).
27.   Schroeder, A.*, et al.* Inhibitory top-down projections from zona incerta mediate
neocortical memory. *Neuron* **111**, 727-738 e728 (2023).
28.   Saunders, A.*, et al.* A direct GABAergic output from the basal ganglia to frontal
cortex. *Nature* **521**, 85-89 (2015).
29.   Schlesiger, M.I.*, et al.* Two septal-entorhinal GABAergic projections
differentially control coding properties of spatially tuned neurons in the medial
entorhinal cortex. *Cell Rep* **34**, 108801 (2021).
30.   Mitrofanis, J. Some certainty for the "zone of uncertainty"? Exploring the
function of the zona incerta. *Neuroscience* **130**, 1-15 (2005).
31.   Wang, X., Chou, X.L., Zhang, L.I. & Tao, H.W. Zona Incerta: An Integrative Node
for Global Behavioral Modulation. *Trends Neurosci* **43**, 82-87 (2020).
32.   Yang, Y.*, et al.* Whole-Brain Connectome of GABAergic Neurons in the Mouse
Zona Incerta. *Neurosci Bull* **38**, 1315-1329 (2022).
33.   Wang, Q.*, et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D
Reference Atlas. *Cell* **181**, 936-953 e920 (2020).
34.   Zhang, X. & van den Pol, A.N. Rapid binge-like eating and body weight gain
driven by zona incerta GABA neuron activation. *Science* **356**, 853-859 (2017).
35.   Liu, K.*, et al.* Lhx6-positive GABA-releasing neurons of the zona incerta
promote sleep. *Nature* **548**, 582-587 (2017).
36.   Shang, C.*, et al.* A subcortical excitatory circuit for sensory-triggered predatory
hunting in mice. *Nat Neurosci* **22**, 909-920 (2019).

37.   Zhao, Z.D.*, et al.* Zona incerta GABAergic neurons integrate prey-related
sensory signals and induce an appetitive drive to promote hunting. *Nat Neurosci* **22**,
921-932 (2019).
38.   Ogasawara, T.*, et al.* A primate temporal cortex-zona incerta pathway for
novelty seeking. *Nat Neurosci* **25**, 50-60 (2022).
39.   Venkataraman, A.*, et al.* Modulation of fear generalization by the zona incerta.
Proc Natl Acad Sci U S A **116**, 9072-9077 (2019).
40.   Li, Z., Rizzi, G. & Tan, K.R. Zona incerta subpopulations differentially encode
and modulate anxiety. *Sci Adv* **7**, eabf6709 (2021).
41.   Ahmadlou, M.*, et al.* A cell type-specific cortico-subcortical brain circuit for
investigatory and novelty-seeking behavior. *Science* **372** (2021).
42.   Lin, S.*, et al.* Somatostatin-Positive Neurons in the Rostral Zona Incerta
Modulate Innate Fear-Induced Defensive Response in Mice. *Neurosci Bull* (2022).
43.   Velmeshev, D.*, et al.* Single-cell genomics identifies cell type-specific molecular
changes in autism. *Science* **364**, 685-689 (2019).
44.   Huang, K.W.*, et al.* Molecular and anatomical organization of the dorsal raphe
nucleus. *Elife* **8** (2019).
45.   Zeisel, A.*, et al.* Molecular Architecture of the Mouse Nervous System. *Cell* **174**,
999-1014 e1022 (2018).
46.   La Manno, G.*, et al.* Molecular Diversity of Midbrain Development in Mouse,
Human, and Stem Cells. *Cell* **167**, 566-580 e519 (2016).
47.   Kee, N.*, et al.* Single-Cell Analysis Reveals a Close Relationship between
Differentiating Dopamine and Subthalamic Nucleus Neuronal Lineages. *Cell Stem Cell*
20, 29-40 (2017).
48.   Chen, R., Wu, X., Jiang, L. & Zhang, Y. Single-Cell RNA-Seq Reveals Hypothalamic
Cell Diversity. *Cell Rep* **18**, 3227-3241 (2017).
49.   Zhang, Y.H.*, et al.* Cascade diversification directs generation of neuronal
diversity in the hypothalamus. *Cell Stem Cell* **28**, 1483-1499 e1488 (2021).
50.   Marques, S.*, et al.* Oligodendrocyte heterogeneity in the mouse juvenile and
adult central nervous system. *Science* **352**, 1326-1329 (2016).
51.   Lein, E.S.*, et al.* Genome-wide atlas of gene expression in the adult mouse brain.
Nature **445**, 168-176 (2007).
52.   Xu, Y.*, et al.* Identification of a neurocircuit underlying regulation of feeding by
stress-related emotional responses. *Nat Commun* **10**, 3446 (2019).
53.   Mu, M.D.*, et al.* A limbic circuitry involved in emotional stress-induced
grooming. *Nat Commun* **11**, 2261 (2020).
54.   Gao, Z.R.*, et al.* Tac1-Expressing Neurons in the Periaqueductal Gray Facilitate
the Itch-Scratching Cycle via Descending Regulation. *Neuron* **101**, 45-59 e49 (2019).
55.   Xie, Z.*, et al.* A brain-to-spinal sensorimotor loop for repetitive self-grooming.
Neuron **110**, 874-890 e877 (2022).
56.   Zhang, Y.F.*, et al.* Ventral striatal islands of Calleja neurons control grooming
in mice. *Nat Neurosci* **24**, 1699-1710 (2021).
57.   Kalueff, A.V.*, et al.* Neurobiology of rodent self-grooming and its value for
translational neuroscience. *Nat Rev Neurosci* **17**, 45-59 (2016).

58.   Sun, J., *et al.* Excitatory SST neurons in the medial paralemniscal nucleus
control repetitive self-grooming and encode reward. *Neuron* **110**, 3356-3373 e3358
(2022).
59.   Mangieri, L.R., *et al.* A neural basis for antagonistic control of feeding and
compulsive behaviors. *Nat Commun* **9**, 52 (2018).
60.   Liu, Q., *et al.* An iterative neural processing sequence orchestrates feeding.
Neuron **111**, 1651-1665 e1655 (2023).
61.   Ahnaou, A., *et al.* Long-term enhancement of REM sleep by the pituitary
adenylyl cyclase-activating polypeptide (PACAP) in the pontine reticular formation of
the rat. *Eur J Neurosci* **11**, 4051-4058 (1999).
62.   Maloney, K.J., Mainville, L. & Jones, B.E. c-Fos expression in GABAergic,
serotonergic, and other neurons of the pontomedullary reticular formation and raphe
after paradoxical sleep deprivation and recovery. *J Neurosci* **20**, 4669-4679 (2000).
63.   Yamahachi, H., Marik, S.A., McManus, J.N., Denk, W. & Gilbert, C.D. Rapid axonal
sprouting and pruning accompany functional reorganization in primary visual cortex.
Neuron **64**, 719-729 (2009).
64.   Bagri, A., Cheng, H.J., Yaron, A., Pleasure, S.J. & Tessier-Lavigne, M. Stereotyped
pruning of long hippocampal axon branches triggered by retraction inducers of the
semaphorin family. *Cell* **113**, 285-299 (2003).
65.   Luo, L. & O'Leary, D.D. Axon retraction and degeneration in development and
disease. *Annu Rev Neurosci* **28**, 127-156 (2005).
66.   Wen, S., *et al.* Spatiotemporal single-cell analysis of gene expression in the
mouse suprachiasmatic nucleus. *Nat Neurosci* **23**, 456-467 (2020).
67.   Kim, D.W., *et al.* Gene regulatory networks controlling differentiation, survival,
and diversification of hypothalamic Lhx6-expressing GABAergic neurons. *Commun
Biol* **4**, 95 (2021).
68.   Puelles, L., *et al.* LacZ-reporter mapping of Dlx5/6 expression and
genoarchitectural analysis of the postnatal mouse prethalamus. *J Comp Neurol* **529**,
367-420 (2021).
69.   Wang, X.Y., *et al.* A cross-modality enhancement of defensive flight via
parvalbumin neurons in zonal incerta. *Elife* **8** (2019).
70.   Kalueff, A.V. & Tuohimaa, P. Grooming analysis algorithm for
neurobehavioural stress research. *Brain Res Brain Res Protoc* **13**, 151-158 (2004).
71.   Kulkosky, P.J., Gibbs, J. & Smith, G.P. Feeding suppression and grooming
repeatedly elicited by intraventricular bombesin. *Brain Res* **242**, 194-196 (1982).
72.   Krahn, D.D., Gosnell, B.A., Grace, M. & Levine, A.S. CRF antagonist partially
reverses CRF- and stress-induced effects on feeding. *Brain Res Bull* **17**, 285-289
(1986).
73.   Rodrigo-Angulo, M.L., Heredero, S., Rodriguez-Veiga, E. & Reinoso-Suarez, F.
GABAergic and non-GABAergic thalamic, hypothalamic and basal forebrain
projections to the ventral oral pontine reticular nucleus: their implication in REM
sleep modulation. *Brain Res* **1210**, 116-125 (2008).
74.   Vanini, G., Wathen, B.L., Lydic, R. & Baghdoyan, H.A. Endogenous GABA levels
in the pontine reticular formation are greater during wakefulness than during rapid
eye movement sleep. *J Neurosci* **31**, 2649-2656 (2011).

75.   Xi, M.C., Morales, F.R. & Chase, M.H. Evidence that wakefulness and REM sleep
are controlled by a GABAergic pontine mechanism. *J Neurophysiol* **82**, 2015-2019
(1999).
76.   Camacho-Arroyo, I., Alvarado, R., Manjarrez, J. & Tapia, R. Microinjections of
muscimol and bicuculline into the pontine reticular formation modify the sleep-
waking cycle in the rat. *Neurosci Lett* **129**, 95-97 (1991).
77.   Blanco-Centurion, C., Vidal-Ortiz, A., Sato, T. & Shiromani, P.J. Activity of GABA
neurons in the zona incerta and ventral lateral periaqueductal grey is biased towards
sleep. *Sleep* **46** (2023).
78.   Chen, K.S.*, et al.* A Hypothalamic Switch for REM and Non-REM Sleep. *Neuron*
97, 1168-1176 e1164 (2018).
79.   Weber, F.*, et al.* Control of REM sleep by ventral medulla GABAergic neurons.
Nature **526**, 435-438 (2015).
80.   Xu, M.*, et al.* Basal forebrain circuit for sleep-wake control. *Nat Neurosci* **18**,
1641-1647 (2015).
81.   Pinto, L.*, et al.* Fast modulation of visual perception by basal forebrain
cholinergic neurons. *Nat Neurosci* **16**, 1857-1863 (2013).
82.   Liu, D. & Dan, Y. A Motor Theory of Sleep-Wake Control: Arousal-Action Circuit.
Annu Rev Neurosci **42**, 27-46 (2019).
83.   McGinley, M.J.*, et al.* Waking State: Rapid Variations Modulate Neural and
Behavioral Responses. *Neuron* **87**, 1143-1161 (2015).
84.   John, J., Wu, M.F., Boehmer, L.N. & Siegel, J.M. Cataplexy-active neurons in the
hypothalamus: implications for the role of histamine in sleep and waking behavior.
Neuron **42**, 619-634 (2004).
85.   Plaha, P., Ben-Shlomo, Y., Patel, N.K. & Gill, S.S. Stimulation of the caudal zona
incerta is superior to stimulation of the subthalamic nucleus in improving
contralateral parkinsonism. *Brain* **129**, 1732-1747 (2006).
86.   Mostofi, A.*, et al.* Outcomes from deep brain stimulation targeting subthalamic
nucleus and caudal zona incerta for Parkinson's disease. *NPJ Parkinsons Dis* **5**, 17
(2019).
87.   Summa, K.C.*, et al.* Disrupted sleep-wake regulation in the MCI-Park mouse
model of Parkinson's disease. *NPJ Parkinsons Dis* **10**, 54 (2024).
88.   Tandberg, E., Larsen, J.P. & Karlsen, K. A community-based study of sleep
disorders in patients with Parkinson's disease. *Mov Disord* **13**, 895-899 (1998).
89.   Corces, M.R.*, et al.* An improved ATAC-seq protocol reduces background and
enables interrogation of frozen tissues. *Nat Methods* **14**, 959-962 (2017).
90.   Tuesta, L.M.*, et al.* In vivo nuclear capture and molecular profiling identifies
Gmeb1 as a transcriptional regulator essential for dopamine neuron function. *Nat*
Commun **10**, 2508 (2019).

1056

Acknowledgement

We thank Research Assistant Zhaoxiaonan Lin from the Core Facilities of Zhejiang University School of Medicine and Dr. Sanhua Fang from the Core Facilities of Zhejiang University Institute of Neuroscience for technical support. We thank Drs. Shumin Duan, Jianhong Luo, Ying Shen (Zhejiang University) for suggestions on the project. This work was supported by grants from the Ministry of Science and Technology (2019YFA0110103 to J.C.), National Natural Science Foundation of China (81870898 to J.C.; 82090030 and 82288101 to X.-M.L.), Zhejiang Provincial Natural Science Foundation (LR18H090002), STI2030-Major Projects (2021ZD0202700 to X.-M.L.; 2021ZD0203400 to Y.-Q.Y.), Key-Area Research and Development Program of Guangdong Province (2019B030335001 to X.-M.L.), Key R&D Program of Zhejiang Province (2020C03009 and 2021C03001 to X.-M.L.), Fundamental Research Funds for the Central Universities (2021FZZX001-37 to X.-M.L.), CAMS Innovation Fund for Medical Sciences (2019-I2M-5-057 to X.-M.L.), Innovative and Entrepreneur Team of Zhejiang for 2020 Biomarker-Driven Basic and Translational Research on Major Brain Diseases (2020R01001 to X.-M.L.), National Natural Science Foundation of China Major Project (T2293733 to Y.-Q.Y.), Postdoctoral Fellowship Program of CPSF (GZC20232285 to M.Z.), and the Non-profit Central Research Institute Fund of Chinese Academy of Medical Sciences (2023-PT310-01).

Author Contributions

J.C. conceived and designed the project, M.Z. M.W. S.L. performed in vivo circuit tracing, optogenetics, behavioral experiments and data analysis. M.Z. M.W. performed immunohistochemistry staining, fluorescence in situ hybridization and confocal imaging, J.P. H.Z. Y.Z. L.S. performed single-nuclei RNA sequencing and data analysis. X.D. performed electrophysiology experiments and data analysis. M.Z. M.W. H.Z. Y.Y. performed EEG/EMG recordings and data analysis. J.C. X.L.

1085 analyzed data and supervised the project. J.C. prepared the manuscript with inputs
1086 from all authors.

1087

Competing Interest

1089 All authors claim that there are no competing interests.

1090

Figure legends

Fig. 1: Single-nucleus transcriptomic sequencing reveals major cell type composition in zona incerta.

a, Schematic diagram showing the workflow for microdissection of zona incerta from brain slices in green fluorescent protein knock-in transporter line Gad67-GFP mice, single nucleus isolation and RNA sequencing by 10x Genomics. UMAP plot of the processed dataset containing 15,375 single nuclei collected from two independent cohorts of 10x Genomics experiments (n = 6 mice for each cohort). The cell clusters were annotated according to canonical cell type-specific marker genes.

b, UMAP plot showing re-clustering the excitatory and inhibitory neurons into ten glutamatergic neuron clusters (Glu_N1 to Glu_N10) and twelve GABAergic neuron clusters (GABA_N1 to GABA_N12).

c, Heatmap showing the differentially expressed genes (DEGs) among neuron clusters from snRNA-seq dataset. The regional identity (top) of the neuron clusters were annotated by the expression of region specific DEGs (right). TH, Thalamus; STN, Subthalamus nucleus; PVN, Paraventricular nucleus; Hypo, Hypothalamus; ZI, Zona incerta; PIL, Posterior intralaminar thalamus nucleus; TRN, Thalamic reticular nucleus; UnA, Un-Annotated region.

d, UMAP plot showing the GABAergic neuron clusters (GABA_N4 to GABA_N9) associated with zona incerta.

1112 **e,** Dotplot showed the differential expressed genes (DEGs) of GABAergic neuron

1113 clusters associated with zona incerta. Source data are provided as a Source Data

1114 file.

1115

Fig. 2: Region-specific genes expression of GABAergic neuron subtypes in

subregions of zona incerta.

a, Violin Plots showed the expression level of selected DEG genes of GABAergic

neuron clusters in zona incerta.

b, Representative images (**b**) and statistical results (**c**) showing the density of

Ecel1 (ZIr *vs* ZIm, ****P < 0.0001; ZIr *vs* ZIc, ****P < 0.0001; ZIm *vs* ZIc, *P =

0.0443), *Cdh23* (ZIr *vs* ZIm, **P =0.0075; ZIr *vs* ZIc, **P =0.0039)*, Pax6* (ZIr *vs*

ZIm, ***P =0.0004; ZIr *vs* ZIc, ***P =0.0007), *Pde11a* (ZIr *vs* ZIm, **P =0.0038;

ZIm *vs* ZIc, ***P =0.0006), *Ptprk* (ZIr *vs* ZIm, **P =0.0020; ZIr *vs* ZIc, *P =0.0119),

Nos1 (ZIr *vs* ZIm, **P =0.0062; ZIr *vs* ZIc, **P =0.0092) in the ZIr, ZIm and ZIc, n

= 9 brain slices from 3 mice per group. Scale bar, 200 μm. One-way ANOVA and

tukey's multiple comparisons test.

d, Representative images (**d**) and statistical results (**e**) showing the overlap

between *Ecel1*, *Pde11a* and *Ptprk* in different brain sections of ZIr, ZIm and ZIc.

The cell numbers in ZIr (*Ecel1*$^+$: 410.0 ± 9.539 cells; *Ecel1*$^+$*Pde11a*$^+$: 9.33 ± 1.20

cells; *Ecel1*$^+$*Ptprk*$^+$: 43.33 ± 5.49 cells; *Pde11a*$^+$*Ptprk*$^+$: 4.33 ± 0.88 cells;

Ecel1$^+$*Pde11a*$^+$*Ptprk*$^+$: 1.67 ± 0.33 cells), ZIm (*Pde11a*$^+$: 361.0 ± 23.81 cells;

Ecel1$^+$*Pde11a*$^+$: 1.67 ± 0.67 cells; *Ecel1*$^+$*Ptprk*$^+$: 4.67 ± 0.88 cells; *Pde11a*$^+$*Ptprk*$^+$:

2.67 ± 0.33 cells; *Ecel1*$^+$*Pde11a*$^+$*Ptprk*$^+$: 0 ± 0 cells) and ZIc (*Ptprk*$^+$: 132.3 ± 16.13

cells; *Ecel1*$^+$*Pde11a*$^+$: 3.33 ± 0.33 cells; *Ecel1*$^+$*Ptprk*$^+$: 7.33 ± 0.88 cells;

Pde11a$^+$*Ptprk*$^+$: 2.00 ± 0.58 cells; *Ecel1*$^+$*Pde11a*$^+$*Ptprk*$^+$: 2.33 ± 0.88 cells) were

shown respectively. N = 9 brain slices from 3 mice per group. Scale bars, 100 μm

and 20 μm (zoom-in image). Data are presented as mean ± SEM. Source data are

provided as a Source Data file.

Fig. 3: Generation and validation of transgenic mice Ecel1-Cre and Pde11a-Cre.

a, Representative images showing the expression of *Ecel1* (**a**) and *Pde11a* (**b**) in *Slc32a1* (Vgat) positive neurons in ZIr or ZIm. Scale bar, 50µm.

c, The percentage of *Ecel1*+*Slc32a1*+ cells relative to all *Ecel1*+ or *Slc32a1*+ cells in ZIr (*Ecel1*+*Slc32a1*+/*Ecel1*+, 96.39±1.19%; *Ecel1*+*Slc32a1*+/*Slc32a1*+, 59.69±6.28%) and the percentage of *Pde11a*+*Slc32a1*+ cells relative to all *Pde11a*+ or *Slc32a1*+ cells in ZIm (*Pde11a*+*Slc32a1*+/*Pde11a*+, 97.7±2.3%; *Pde11a*+*Slc32a1*+/*Slc32a1*+, 33.92±9.67%). n=9 brain slices from 3 mice per group.

d, Schematic diagram showing the generation of Ecel1-Cre or Pde11a-Cre knock-in mice.

e, Representative images showing the Cre and *Ecel1* or the Cre and *Pde11a* expression in ZIr, ZIm and ZIc. Scale bars, 100µm and 20µm (zoom-in image).

f, The percentage of *Ecel1*+Cre+ relative to all Cre+ or *Ecel1*+ neurons in ZIr (*Ecel1*+Cre+/Cre+,93.87±1.45%; *Ecel1*+Cre+/*Ecel1*+, 96.41±0.92%), ZIm (*Ecel1*+Cre+/Cre+, 91.11±2.81%; *Ecel1*+Cre+/*Ecel1*+, 86.46±4.39%) and ZIc (*Ecel1*+Cre+/Cre+, 78.41±4.75%; *Ecel1*+Cre+/*Ecel1*+, 78.00±5.95%) of Ecel1-Cre mice. n = 9 brain slices from 3 mice per group.

g, Statistical results showing the expression and distribution of Cre in the ZIr (71.95±1.12%), ZIm (12.76±0.67%) and ZIc (15.28±0.49%) of Ecel1-Cre mice. ZIr *vs* ZIm, ****$P<0.0001$; ZIr *vs* ZIc, ****$P<0.0001$. N = 9 brain slices from 3 mice per group.

h, The percentage of *Pde11a*+Cre+ relative to all Cre+ or *Pde11a*+ neurons in ZIr (*Pde11a*+Cre+/Cre+, 97.70±2.30%; *Pde11a*+Cre+/*Pde11a*+, 99.48±0.52%), ZIm (*Pde11a*+Cre+/Cre+, 99.70±0.15%; *Pde11a*+Cre+/*Pde11a*+, 98.91±0.22%) and ZIc (*Pde11a*+Cre+/Cre+, 93.00±4.72%; *Pde11a*+Cre+/*Pde11a*+, 89.73±3.06%) of Pde11a-Cre mice. n=9 brain slices from 3 mice per group.

i, Statistical results showing the expression and distribution of Cre in the ZIr (4.43±2.54%), ZIm (90.33±2.91%) and ZIc (5.24±1.27%) of Pde11a-Cre mice. ZIr *vs* ZIm, ****P<0.0001; ZIm *vs* ZIc, ****P< 0.0001. N=9 brain slices from 3 mice per group.

One-way ANOVA and tukey's multiple comparisons test for (**g**) and (**i**). Data are presented as mean±SEM. Source data are provided as a Source Data file.

Fig. 4: Distinct efferent projections of Ecel1 positive neurons in the rostral ZI and Pde11a positive neurons in the medial ZI.

a, Schematic diagram and representative images showing the stereotaxic unilateral injection of Cre-dependent AAV mediated mGFP and synaptophysin-mRuby to specifically label Ecel1 positive neurons in the rostral ZI (ZIr^{Ecel1}) of Ecel1-Cre mice (**a**) or Pde11a positive neurons in the medial ZI (ZIm^{Pde11a}) of Pde11a-Cre mice (**b**). Scale bar, 250 μm. The experiment was independently repeated 3 times with similar results for each mouse strain.

c, Representative images showing dense distribution of mGFP and mRuby positive axons from the ZIr^{Ecel1} neurons to the PAG and from ZIm^{Pde11a} neuron to the PRF. Scale bar, 100 μm.

d, Quantification of integrated fluorescence intensity of mRuby and mGFP-labeled axon terminals in PAG or PRF of ZIr^{Ecel1} and ZIm^{Pde11a} neurons. *P = 0.015, **P = 0.0032, n = 3 mice.

e, Quantification of relative fluorescent density of mRuby and mGFP-labeled axon terminals of ZIr^{Ecel1} and ZIm^{Pde11a} neurons in the target brain regions. N = 9 slices from 3 mice for each brain region. Abbreviations: LS, lateral septal nucleus; POA, preoptic area; PVT, paraventricular thalamic nucleus; Re, reuniens thalamic nucleus; LH, lateral hypothalamic area; MD, mediodorsal thalamic nucleus; Po, posterior thalamic nucleus; Auv, secondary auditory cortex; APTD, anterior pretectal nucleus; Dk, nucleus of Darkschewitsch; RN, red nucleus; PAG,

1195 periaqueductal gray; PRF, pontine reticular nucleus, oral part; SC, superior
1196 colliculus.

1197 Two-tailed unpaired t test for (**d**). Data are presented as mean ± SEM. Source data
1198 are provided as a Source Data file.

1199

Fig. 5: Optogenetic activation of ZIr^{Ecel1} neurons and the ZIr^{Ecel1} to PAG
pathway both induce self-grooming.

a, Schematic diagram showing optogenetic stimulation of the ChR2-EYFP or
EGFP expressing ZIr^{Ecel1} neurons in Ecel1-Cre mice.

b, Schematic diagram showing the behavioral paradigm for measurement of self-
grooming behavior upon optogenetic stimulation.

c, Statistic analysis of time spent for self-grooming (left) and grooming bouts (right)
in 3 min before, during and after optogenetic stimulation of ZIr^{Ecel1} neurons in
ChR2-group (**c**) and EGFP-group (**d**), respectively. Data are presented as mean ±
SEM. **$P = 0.0014$, ****$P < 0.0001$. Blue column indicates laser stimulation period
(5 Hz, 3 min).

e, Schematic diagram showing the recording of inhibitory postsynaptic currents
(IPSCs) from PAG neurons evoked by optogenetic stimulation of ChR2-EYFP
expressing axon terminals from the ZIr^{Ecel1} to the PAG in acute brain slices.

f, Representative traces and statistical analysis of light-evoked IPSCs recorded in
PAG neuron in ACSF and following sequential application of TTX (**f,** 1 µM. **$P =$
0.0082), TTX + 4-AP (**f,** 100 µM. ****$P < 0.0001$), or bicuculline (**g,** 20 µM. **$P =$
0.0039). N = 9 neurons from 5 mice, two-tailed paired t test. Blue bar indicates
optogenetic stimulation (10 ms).

h, Schematic diagram and representative images showing optogenetic stimulation
of the ChR2-EYFP or EGFP expressing terminals from ZIr^{Ecel1} to PAG pathway in
Ecel1-Cre mice. Scale bar, 100 µm.

i, Statistic analysis showing time spent for self-grooming (left) and grooming bouts (right) in 3 min before, during or after optogenetic stimulation of ZIr^{Ecel1} to PAG pathway in ChR2-group (**i**) and EGFP-group (**j**), respectively. **P = 0.0013, ****P < 0.0001. Blue column indicates laser stimulation period (20 Hz, 3min).

One-way ANOVA for (**c**), (**d**), (**i**) and (**j**). Two-tailed paired *t* test for (**f**) and (**g**). Data are presented as mean ± SEM. Source data are provided as a Source Data file.

Fig. 6: Calcium response patterns of Ecel1 positive neurons in the rostral ZI in self-grooming behavior.

a, Schematic (left) and representative images showing GCaMP6s and mCherry expression in ZIr of Ecel1-Cre mice (right). Scale bars, 500 μm and 250 μm (zoom-in image). The experiment was independently repeated 5 times with similar results.

b, Schematic diagram showing the behavioral paradigm for fiber photometry recording in freely moving mice.

c, Schematic diagram (**c**) and statistic analysis (**d**) showing water spray toward the face effectively induced self-grooming behaviors. **P = 0.0013, n = 5 mice.

e, Heatmap (**e**) and average calcium transients (**f**) of ZIr^{Ecel1} neurons before, during and after water-spray induced self-grooming. Red trace: recording of GCaMP6s fluorescent signal; Gray trace: recording of control mCherry fluorescent signal. The dashed lines indicate the start or the end of self-grooming. Shaded areas around means indicate standard error of mean (SEM).

g, Statistic analysis showing area under curve (AUC) of average calcium transients of GCaMP6s channel compared with mCherry control channel before grooming starts (*P = 0.0156) and after grooming starts (*P = 0.0152), before grooming ends (*P = 0.0385) and after grooming ends (*P = 0.0114) in water-spray induced self-grooming. N = 5 mice per group.

1248 Two-tailed paired t-test for (**d**) and (**g**). Data are presented as mean ± SEM. Source
1249 data are provided as a Source Data file.

1250

Fig. 7: Ecel1 positive neurons in the rostral ZI are required for water spray
induced self-grooming.

a, Schematic diagram and representative images showing the stereotaxic bilateral
injection of AAV-DIO-hM3Dq-mCherry or AAV-DIO-mCherry in the ZIr of Ecel1-
Cre mice. Scale bar, 100 µm. The experiment was independently repeated 5 times
with similar results.

b-d, Statistic analysis of time spent for self-grooming (**b**), grooming bouts (**c**) and
grooming duration per bout (**d**) in 20min following injection of CNO in hM3Dq-group
mice compared with mCherry-group mice. **$P = 0.0097$, *$P = 0.0336$.

e, Schematic diagram and representative images showing the injection of AAV-
DIO-hM4Di-mCherry to chemogenetically inhibit ZIr^{Ecel1} neurons of Ecel1-Cre mice.
Scale bar, 100 µm. The experiment was independently repeated 7 times with
similar results.

f, The behavioral paradigm for measurement of water-spray induced self-grooming
behavior following injection of CNO in hM4Di-group or mCherry-group mice.

g-i, Statistic analysis of time spent for water-spray induced self-grooming (**g**),
grooming bouts (**h**) and grooming duration per bout (**i**) in 20 min following injection
of CNO in hM4Di mice. *$P = 0.0117$, **$P = 0.0018$.

j, Schematic diagram showing the stereotaxic bilateral injection of AAV-DIO-DTA
to ablate the ZIr^{Ecel1} neurons of Ecel1-Cre mice.

k-m, Statistic analysis of water-spray induced self-grooming time (**k**), grooming
bouts (**l**) and grooming duration per bout (**m**) in 20 min following water spray in
DTA-group mice. ****$P < 0.0001$, **$P = 0.0036$.

Two-tailed unpaired *t*-test for (**b-d**), (**g-i**) and (**k-m**). Data are presented as mean

± SEM. Source data are provided as a Source Data file.

1276

Fig. 8: Calcium response patterns of Pde11a positive neurons in the medial

ZI during sleep-wakefulness cycle.

a, Schematic and representative images (right) showing *in vivo* recording of

calcium response pattern of GCaMP6s- or mCherry-expressing Pde11a positive

neurons in ZIm. Gray column indicates the position of optical fiber. Scale bars, 500

μm and 250 μm (zoom-in image). The experiment was independently repeated 5

times with similar results.

b, Schematic diagram showing the behavioral paradigm for fiber photometry,

electroencephalogram (EEG) and electromyographic (EMG) recordings in freely

moving mice.

c, Sample trace showing the changes of fluorescent calcium signal of ZIm^{Pde11a}

neurons aligned to sleep- wakefulness state transitions. NREM (yellow bar), non-

rapid eye movement; REM (green bar), rapid eye movement; Wake (blue bar).

d, Average trace and heatmap showing changes of the fluorescent calcium signal

of ZIm^{Pde11a} neurons during sleep-wakefulness state transitions. The red trace

showed average fluorescent GCaMP6s or control mCherry (gray) signals, while

shaded areas indicate standard error of mean (SEM).

e, Statistical analysis of the area under curve (AUC) of average fluorescent calcium

signal of ZIm^{Pde11a} neurons during sleep-wakefulness state transitions. N = 5 mice

per group. *P = 0.0207 (NREM to REM), *P = 0.0332 (REM to Wake), *P = 0.0171

(Wake to NREM), two-tailed paired *t*-test. Data are presented as mean ± SEM.

Source data are provided as a Source Data file.

1299

Fig. 9: Optogenetic activation of ZIm^Pde11a neurons promotes wakefulness through the ZIm^Pde11a to PRF pathway.

a, Optogenetic stimulation of ChR2-EYFP or EGFP expressing ZIm^Pde11a neurons and the behavioral paradigm for EEG-EMG recordings of freely moving Pde11a-Cre mice (**b**).

c, Sample EEG power spectrum, EMG traces and brain states (color coded) during optogenetic activation of ZIm^Pde11a neurons. Dark blue: wake; Yellow: NREM; Green: REM. Freq: frequency.

d, The percentage of wake, NREM, or REM states before, during, and after optogenetic stimulation of ZIm^Pde11a neurons in ChR2-group (left, n = 5 mice) and EGFP-group (right, n = 6 mice). Shaded areas around means indicate standard error of mean (SEM).

e, Probability change of each state 120s before and during optogenetic stimulation of ZIm^Pde11a neurons in ChR2-group compared with EGFP-group. ****$P < 0.0001$, **$P = 0.0021$.

f-h, Schematic diagram (**f**) and representative traces (**g** and **h**, left) showing the recording of evoked IPSCs in PRF neurons in acute brain slices. TTX, *$P = 0.0146$; TTX+4-AP, *$P = 0.0129$ (n = 6 neurons from 4 mice); Bicuculline, *$P = 0.0215$ (n = 7 neurons from 4 mice). Blue bar indicates laser stimulation period (10ms).

i, Optogenetic stimulation of ChR2-EYFP or EGFP expressing terminals of ZIm^Pde11a to PRF pathway together with EEG-EMG recordings in Pde11a-Cre mice. Scale bar, 100µm. The experiment was independently repeated 5 times with similar results.

j, Sample EEG power spectrum, EMG traces and brain states during optogenetic activation of ZIm^Pde11a to PRF pathway.

k, The percentage of wake, NREM, or REM states before, during, and after optogenetic stimulation of ZIm^Pde11a to PRF pathway in ChR2-group (left, n = 5

mice) and EGFP-group (right, n = 5 mice). Shaded areas around means indicate

standard error of mean (SEM).

l, Probability change of each state 120s before and during optogenetic stimulation

of ZImPde11a to PRF pathway in ChR2-group compared with EGFP-group. ****$P <$

0.0001.

Two-tailed unpaired t test for (**e**) and (**l**). Two-tailed paired t test for (**g**) and (**h**).

Data are presented as mean±SEM. Blue bar indicates laser stimulation period

(20Hz, 120s). Source data are provided as a Source Data file.

1335

Fig. 10: Selective chemogenetic activation or ablation of ZImPde11a neurons

bidirectionally regulate sleep-wakefulness cycle.

a, Schematic diagram and representative images showing the injection of AAV-

DIO-hM3Dq-mCherry or AAV-DIO-mCherry into the ZIm of Pde11a-Cre mice.

Scale bar, 250µm. The experiment was independently repeated 10 times with

similar results.

b, Schematic showing sleep recording conditions with saline- or CNO-injection in

hM3Dq-group mice.

c-e, The percentage of time spent in wake (**c**, ZT7, ***P=0.0008), NREM sleep (**d**,

ZT7, ***P=0.0007) and REM sleep (**e**, ZT7, *P=0.0117) in 2-h bins following

injection of saline or CNO in hM3Dq-group mice. Wake, **P=0.0056 (**c**); NREM,

P=0.0080 (d**); REM, **P=0.009 (**e**).

f-h, Time spent in wake (**f**), NREM sleep (**g**) and REM sleep (**h**) across the 24-

hour sleep-wake cycle in hM3Dq-group mice. Wake, **P=0.0083 (**f**); NREM,

*P=0.0180 (**g**); REM, *P=0.0304 (**h**).

i, Number and duration (**j**) of wake, NREM sleep and REM sleep episodes during

the light phase. Number: Wake, **P=0.0027; NREM, **P=0.0026. Duration: NREM,

****P<0.0001.

k, Schematic diagram showing the injection of AAV-DIO-DTA to ablate the

ZImPde11a neurons.

l-n, The percentage of time spent in wake (**l**, ZT21, *$*P$=0.0165), NREM sleep (**m**, ZT21, *$*P$=0.0244) and REM sleep (**n**, ZT21, *$*P$=0.0427) in 2-h bins in DTA-group mice compared with mCherry-group mice. REM, **$**P$=0.0040 (**n**).

o-q, Time spent in wake (**o**), NREM sleep (**p**) and REM sleep (**q**) across the 24-hour sleep-wake cycle in DTA-group mice compared with mCherry-group mice. Wake, **$**P$=0.0045 (**o**); NREM, **$**P$=0.0082 (**p**); REM, **$**P$=0.0013 (**q**).

r, Number and duration (**s**) of wake, NREM sleep and REM sleep episodes during the dark phase. Number: Wake, ***$***P$=0.0004; NREM, ***$***P$=0.0005; REM, **$**P$=0.0016. Duration: Wake, ***$***P$=0.0004.

Two-way ANOVA and Sidak's multiple comparisons test for (**c-e**) and (**l-n**). Two-tailed paired t test for (**c-j**). Two-tailed unpaired t test for (**o-s**). Data are presented as mean±SEM. Grey bar indicates the dark phase (ZT12-ZT0). Source data are provided as a Source Data file.

a AAV-FLEX-mGFP-2A-
Synaptophysin-mRuby

mGFP/mRuby/DAPI

ZIr

Ecel1-Cre mice

b AAV-FLEX-mGFP-2A-
Synaptophysin-mRuby

mGFP/mRuby/DAPI

ZIm

Pde11a-Cre mice

c

Ecel1-Cre mice | Pde11a-Cre mice

mGFP/mRuby/DAPI

PAG | PAG

PRF | PRF

d

Fluorescence intensity in PAG

Ecel1   Pde11a

Fluorescence intensity in PRF

Ecel1   Pde11a

e

% of Total outputs

LS POA PVT Re LH MD PO AuV APTD Dk RN PAG PRF SC

● Ecel1
● Pde11a

a AAV-DIO-GCaMP6s and AAV-DIO-mCherry

Ecel1-Cre mice

GCaMP6s/ mCherry/ DAPI

Merge ZIr | GCaMP6s

mCherry | DAPI

b Amplifier and DAQ

PMT

Filter

Laser

Fiber launch

c Water spray

d

Grooming time (s)

base | water spray

**

e Water spray-induced self-grooming

Grooming starts | Grooming ends

Animals

Time (s) | Time (s)

f

Grooming starts | Grooming ends

ΔF/F (%)

Time (s) | Time (s)

GCaMP6s
mCherry

g

Before grooming | Grooming starts | Grooming ends | After grooming

Area under curve

GCamP | mCherry

*