Cethromycin Pharmacokinetics and Pharmacodynamics for Single Dose Cure of

Plasmodium berghei Liver Stages

Grace Kennedy[1], Rachel M. West[1], Kristin Poti[1], Bryce Bobb[1], Matthew M. Ippolito[1,2],

Mark A. Marzinke[2], Nikola Kaludov[3] and David J. Sullivan Jr.[1,3]#

[1]W. Harry Feinstone Department of Molecular Microbiology and Immunology, Johns

Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

[2]Clinical Pharmacology, Johns Hopkins University School of Medicine, Baltimore

Maryland USA

[3]AliQuantumRx Inc, Baltimore, Maryland, USA

Running Head: Cethromycin malaria action for liver and blood stages

#Corresponding author: David J. Sullivan, Johns Hopkins Bloomberg School of Public

Health. 615 North Wolfe Street, Baltimore, Maryland 21205, USA. Email:

dsulliv7@jhmi.edu

Abstract

Cethromycin combines a quinoline nucleus and a macrolide for broad spectrum antibacterial and antiprotozoan activity. Here we characterized the murine pharmacokinetics and *Plasmodium berghei* lifecycle stage pharmacodynamics for the cethromycin base. Liver pharmacokinetic studies in mice show peak mM drug levels in the liver with 20 hour sustained levels above 10 μM. Peak concentrations in the liver were double the lung and about 440 times that of plasma. Immunofluorescence imaging of *in vitro* cethromycin-treated infected hepatocytes shows complete ablation of the apicoplast. We observed complete cure of *P. berghei* liver stage infection by single oral dose of 60 mg/kg in mice which is equivalent to the 5 mg/kg human dose of 300 mg a day used in bacterial pneumonia studies. Cethromycin at 60 mg/kg daily for 7 days was curative in the high parasitemic *P. berghei* mouse model. Both mosquito membrane feeding of *P. falciparum* gametocytes incubated with 20 μM cethromycin and oral dosing in mice demonstrated no decrease in oocyst numbers. Cethromycin has been evaluated for efficacy against bacterial pneumonia in more than 5,000 patients with good safety profiles. Cethromycin has potential for rapid clinical development for casual malaria prophylaxis and possibly radical cure of dormant liver *P. vivax*.

INTRODUCTION

Cethromycin is a ketolide class macrolide (erythromycin base with a single sugar and quinoline unsubstituted nucleus replacing the cladinose saccharide) with broad spectrum antibacterial, mycobacterial and antiprotozoal activity developed for community acquired pneumonia[1]. The drug was safe, but not superior to clarithromycin in a large phase 3 outpatient study of more than 1000 participants[2] for which the new drug application was later denied for the nonsuperior efficacy results by the FDA[1, 3]. Cethromycin binds to bacterial 23S ribosomal RNA of the 50S ribosome subunit which stops polypeptide amino-acid addition, terminating protein synthesis and growth[4]. Cethromycin specifically binds domains II and V in the wall of the polypeptide exit tunnel[1]. While ketolides like cethromycin have greater activity than azithromycin against some bacterial species[5],both are unable to overcome high level *erm* resistance[6-8].

The macrolide class has demonstrated delayed death activity *in vitro* and *in vivo* against protozoan apicomplexans like *Toxoplasma*[9] and *Plasmodium[10-12)*. The plant-like apicoplast organelle is essential in production of isoprenoids for blood stage parasites. Previous work has shown that isoprenoid supplementation with azithromycin treatment results in loss of the extrachromosomal apicoplast genome and of the organelle in erythrocyte stages[13].

During the liver stages of malaria infection, the macrolide molecule has been shown to also delete the apicoplast and the extrachromosomal apicoplast genome[14]. Thus, while the liver stages of malaria infection progress, loss of the apicoplast prevents infection of erythrocytes, thus transition to the blood stage [15]. Azithromycin and

doxycycline can cure mouse liver stages(16), but in controlled human malaria challenge

studies, only 4 of 10 patients treated with azithromycin(17) and 8 of 12 patients treated

with doxycycline (18)were cured after 7 days of liver stage treatment initiated on

sporozoite challenge.

In a previous report, we described liver-stage *Plasmodium* inhibition in the

quantitative PCR mouse liver-stage sporozoites-challenge model with single oral low

doses of cethromycin at 15, 25 and 50 mg/kg, because of access to only a few

milligrams of newly synthesized cethromycin(19). Here we report more extensive dose-

ranging and schedules for cethromycin pharmacokinetics and pharmacodynamics in a

murine malaria liver, blood and gametocyte stage models.

Results

Pharmacokinetics

We compared cethromycin concentrations in liver, lung and plasma with single

oral doses in mice at 30, 60 or 120 mg/kg, which is allometrically equivalent to 150, 300

and 600 mg total oral dose in humans. The maximum concentration (C_{max}) in the liver

was double that of the lung and approximately 440 times more than plasma. The 765

gm/mole cethromycin molecular weight indicates 1 μM levels at 0.76 mcg/mL or gram in

the liver. The time above 1 μM in the liver was 24 hours for the 60 mg/kg dose. Plasma

concentrations never exceeded 10 μM and stayed above 1 μM for 12 hrs for the 60

mg/kg dose which is allometrically equal to 300 mg single oral dose in humans (Figure

1). The tissue disposition of cethromycin measured as C_{max}, area under the curve

(AUC_{0-72}), and elimination half-life ($t_{1/2\alpha}$) are shown in Table 1.

Liver stage apicoplast disruption

We investigated maturation of the liver stage *P. berghei* in vitro in 48-hour

hepatocyte cultures treated with primaquine, azithromycin and cethromycin exposure at

20 mM resulted in parasites with predominately no observed apicoplasts. Detailed

counting of infected hepatocytes noted that those exposed to primaquine had parasites

with no apicoplast observed in all hepatocytes, but single dysmorphic apicoplasts were

present in parasites in cethromycin and azithromycin treated hepatocytes. In the

controls, all apicoplasts were highly branched indicating normal hepatocyte

development of the parasite. The heat shock protein (hsp) 70 reporter also indicated

almost normal parasite development with azithromycin and cethromycin exposure

nearly indistinguishable from control while primaquine-exposed parasites contained

small pyknotic *P. berghei* hsp 70 signal (Fig 2).

Liver stage pharmacodynamics

In the 3000 sporozoite tail vein injection murine liver-stage model repeated on

three separate occasions, all 9 controls had patent bloodstream parasites, while none of

the mice treated with cethromycin at 60 mg/kg single oral dose 24 hours after

sporozoite injection were patent. All 6 mice dosed with single oral 15 mg/kg cethromycin

were patent and 1 of 2 at the 30 mg/kg dose. Utilizing a more natural lower dose of

sporozoites with 5-8 infected mosquito bites, all controls were positive in the 8 separate

biologic dosing experiments shown in Table 2 and Fig 3. Cethromycin dosed at 30 or 60

mg/kg 24 hours prior to infection cured zero mice. Both azithromycin and cethromycin

dosed 2 hours after mosquito infection were curative at 60 mg/kg oral dose. Doses 24

hours after infection demonstrated 47% patency with azithromycin when combining

results of the 30 mg/kg (3/6 patent), 45 mg/kg (1/3 patent) and 60 mg/kg (3/6 patent)

doses. For cethromycin 30 mg/kg (1/9 patent), 45 mg/kg (0/3 patent) and 60 mg/kg (1/9

patent) more mice were cured. Cethromycin 20 mg/kg dosed at day -1, 0 and 1 was

curative in all mice (n=3). Quinoline nucleus and erythromycin ethyl succinate, the two

components of cethromycin, given alone and in combination were not curative when

dosed on day 0 and 1 at 120 mg/kg. C57Bl/6 mice are more permissive/susceptible to

murine malaria liver stages than Balb/c with higher liver loads given the same inoculum.

In C57Bl/6 mice azithromycin 45 mg/kg had one patent mouse while at same dose of

cethromycin 0 of three were patent. At the 60 mg/kg dose all nine mice were cured.

Blood-stage

We measured the efficacy of cethromycin and clindamycin in *P. falciparum*

isolates 3D7. The 72 hour IC_{50}s were in the low mM range while the 144 hour assays

indicated 100-fold more potency to near 60 nM IC_{50} (Fig 4A). In the murine *in vivo* 4 day

suppression test, drug doses started after a low (one million) parasite intraperitoneal

inoculum. Both azithromycin and cethromycin at 60 mg/kg showed a similar >99%

inhibition (more than 3 log difference in parasite number) at day 4 but both subsequently

rebounded to one million parasites per μL on day 11 post treatment (Fig 4B). The high

parasitemia model with 1 million parasites per μL azithromycin showed greater parasite

reduction than cethromycin at 2 and 4 days of oral drugs (Fig 4C). In the same high

parasitemia model 6 days of cethromycin was partially curative (2/3 mice), while 7 days

cured all mice (Fig 4D).

Gametocyte stages

Investigation of cethromycin and azithromycin activity against late stage IV and V

gametocytes where 15 day old gametocyte cultures were incubated for 48 hours with 10

μM drugs followed by transmembrane feeding noted similar oocyst numbers and

infection rates (Fig 5).

DISCUSSION

In this work, we show that cethromycin demonstrates potent causal prophylactic

activity against *P. berghei* malaria liver stages with superior cure rates compared to

azithromycin at equivalent doses, while exhibiting typical macrolide delayed-death

activity in blood stages and no significant effect on late-stage gametocytes. This activity

profile aligns with the known mechanism of macrolide antimalarials, where the inability

of the apicoplast to successfully replicate results in parasite death in the second

generation of parasites and leads to the phenomenon of delayed generational

death[20]. Macrolides target both the *Toxoplasma* and *Plasmodium* apicoplast where

the tissue stages appear to progress normally, but the progeny all are nonviable and

lack functional apicoplasts for isoprenoid production[13]. Cethromycin cures liver stage

malaria in mice with a single low dose of 60 mg/kg (human equivalent dose of 300 mg).

A dose of 20 mg/kg in mice over three days, which is equivalent to a 100 mg single

dose in humans, was also curative. Importantly, the two molecular constituents of

cethromycin—erythomycin and quinoline—when given alone or together as separate

molecules, did not cure any mice. We compared the same dose of a related macrolide

in widespread clinical use, azithromycin at 60 mg/kg, and this was not curative.

Primaquine was curative at the single dose of 40 mg/kg. The single dose of cethromycin

given on day 2 or 3 was not curative, showing that its primary effect is on the liver stage

and not subsequent blood stages.

Cethromycin 60 mg/kg, which is allometrically equivalent to the 300 mg single

daily dose in humans, achieved maximum liver concentrations of nearly 3 mM and

remained above 1 μM for 24 hours(21). Azithromycin liver concentrations dosed at 200

mg/kg were 36 mcg/mL (48 mM) a day after dosing(22), and were 200 times that of

serum. Another publication noted a peak liver azithromycin concentration of 73 mcg/mL

(97 mM) after 50 mg/kg dose with four times higher concentrations in liver than lung

(23). Here we noted 2,063 mcg/mL as the observed peak liver concentration after the

60 mg/kg oral dose, which was measured 2.5 hours post-dosing. These data support

other findings which show greatest cethromycin deposition in the liver. Macrolides are

ideally suited by pharmacokinetics to work on liver infections.

In separate experiments initiated by blood stage infection and bypassing the

liver, 7 daily oral doses were curative of a high blood stage parasite density. This is

comparable to the seven days necessary to cure with artesunate—a first-line agent for

severe malaria— in the murine mouse model(24). However, the slow parasiticidal

activity of macrolides which kill by delayed generational death precludes their use as

monotherapy for severe malaria, where rapid parasite reduction is necessary(25). They

do have near equal curative activity in the mouse model to artesunate when dosed on a

daily basis. Artesunate is superior in rapid parasite clearance activity in humans

compared to the antibacterial malaria drugs(26).

The demonstrated cure and an excellent therapeutic product profile of the 3-keto

macrolide quinoline cethromycin suggest further investigation into the niche area of

dormant liver stage malaria. Recent work noted disruption of the apicoplast in both

dormant *P. vivax* and the primate equivalent species *P. cynomolgi*(14) with the

speculation that azithromycin might work in relapsing malaria. Azithromycin in a

triangular human clinical test design of 5 patients was not active in prevention of

dormant *P. vivax* with 3 relapses stopping the trial (27). In the *P. cynomolgi* model

clindamycin by itself or with subcurative primaquine did not stop relapse but

azithromycin with subcurative primaquine prevented relapse in one of two monkeys, but

could not be distinguished from radical cure with primaquine alone in this model(28).

We hypothesize that the quinoline ring might give more liver stage killing than

azithromycin. The two current anti-hypnozoite 8-aminoquinolines, primaquine(29) and

tafenoquine(30, 31), require glucose-6-phosphate dehydrogenase deficiency testing

which is an impediment (time, extra clinical visit, cost and accuracy of point of care test)

to widespread drug implementation(32-35). The target indication for cethromycin is

prevention of dormant *P. vivax* malaria after curative chloroquine or artemisinin

combination treatment for primary *P. vivax* parasitemia. The INSPECTOR trial noted

antagonism of tafenoquine with dihydroartemesinin-piperaquine for dormant *P.*

vivax(36), which might be avoided with cethromycin's different mechanism of action.

Thus, cethromycin might be superior to daily doxycycline prophylaxis with shorter

duration after malaria endemic country exit. Advantages to cethromycin are single daily

doses, high liver tissue concentrations and broad antimicrobial activity. Next steps or

validation of efficacy for dormant malaria in the *P. cynomolgi* model and good

manufacturing practice and investigational new drug enabling preclinical studies for

human malaria testing.

MATERIALS AND METHODS

Animals and parasites. Inbred, female 6-7-week-old wild-type BALB/c and C57Bl/6 mice were purchased from The Jackson Laboratory (Bar Harbor, ME), and kept in the animal facility at the Johns Hopkins Bloomberg School of Public Health. All animal experiments were approved by the Johns Hopkins Animal Care and Use Committee (ACUC), under protocol MO15H319. Sporozoite infected mCherry *P. berghei* mosquitoes were used for liver stage infections (37). *P. berghei* MRA-868 ANKA (*Pb-ANKA-Luc*) transgenic strain was used for all murine blood stage infections. The *Pb-ANKA-Luc* parasite strain expresses a GFP-luciferase fusion gene [19]. NF54 *P. falciparum* parasites at 4% hematocrit were cultured as described(38).

Drug preparation and dosing

Primaquine, quinoline, and erythromycin were from Sigma. Azithromycin (Astatech cat # N460720) and cethromycin (BDG Synthesis, lot 16822, Wellington, New Zealand) were solubilized in either DMSO for in vitro assays or PBS/10% EtOH/20 mM HCl (875 μL PBS + 100 μL ethanol + 50 μL 1 N HCl per mL) for oral dosing in mice. For all the experimental studies done, each drug solution was dosed at a volume of 200 µL per mouse.

Drug Measurements

Cethromycin concentrations in plasma and tissue were determined via liquid chromatography-tandem mass spectrometry by the Clinical Pharmacology Analytical Laboratory at Johns Hopkins. Briefly, cethromycin was isolated from plasma and methanol-homogenized tissue lysate via protein precipitation using a 0.45 µm Captiva filter plate (Agilent Technologies, Wilmington, DE). Cethromycin ($C_{42}H_{59}N_3O_{10}$) peak

area ratios were normalized to the structural analog telithromycin ($C_{43}H_{65}N_5O_{11}$), which

served as an internal standard. Cethromycin and its internal standard were quantified

on an API 4000 (plasma) or API 4500 (tissue homogenates) (SCIEX, Foster City, CA)

operated in positive ionization and selective reaction monitoring modes. Ion transitions

monitored were *m/z* 766.5→158.2 for cethromycin and *m/z* 813.6→656.2 for

telithromycin. The primary analytical measuring ranges for plasma and tissue assays

were 0.1-2500 ng/mL and 10-2500 ng/sample, respectively. Tissue results were

normalized to the weight of tissue analyzed. Assays were validated in accordance with

regulatory recommendations.

Liver stage in vitro Assay

Hepa 1-6 (ATCC CRL-1830) were grown to confluence in DMEM with Fungizone/10%

Pencillin/Streptomycin and plated onto Permanox LAB-TEK slides. Approximately

50,000 *P. berghei* sporozoites were added per well. After 2 hours medium was washed

with fresh medium containing drugs at 20 microM. At 24 hours after sporozoite infection,

cells were washed with fresh medium alone or fresh medium with drugs. At 48 hours

medium was removed and cells fixed with 4% paraformaldehyde for 15 minutes,

washed 3 times with PBS for GFP-luciferase assays or in cold methanol for 15 minutes,

dried and stored at -20C overnight for immunofluorescence. Antibodies were to mouse

anti-*P. berghei* HSP70 (2E6) provided by Fidel Zavala and rabbit anti-Pb acyl carrier

protein (ACP) provided by Dr S Prigge (Johns Hopkins University). Secondary FITC

goat anti-mouse and Alexa594 goat anti-rabbit. Liver stage assay experiment was

performed in biologic duplicate at two different dates.

Model of sporozoite infection

Female *Anopheles stephensi* mosquitoes were fed with *mcherry P. berghei*-infected

blood for sporozoite development, as previously described(37, 39). Twenty days later

sort cage of 200 mosquitoes with fluorescent microscope, Infect mice with 5-7

mosquitoes, examine patency by Giemsa blood film microscopy starting from day 4 to

day 16 post infection.

Plasmodium falciparum culture

Low passage 3D7 or NF54 *Plasmodium falciparum* parasites were cultured and

synchronized as described(40). Cultures were incubated with RPMI-1640 with L-

glutamine, 25 mM HEPES, 40 µM hypoxanthine, 25 mM sodium bicarbonate, and 10%

O+ human serum (Interstate Blood Bank). All cultures, unless otherwise noted, are in

2% hematocrit with O+ red blood cells. Cultures were gassed with 5% O_2/ 5% CO_2 /90%

Nitrogen and placed in 37°C incubators.

In vivo cytocidal model of murine malaria

The rodent model used for all the experiments were Balb/cj female mice from the

Jackson Laboratories aged at least 6 weeks old and weighed approximately 20 grams

each. For each study done, replicate of three mice were used for each drug dose

regimen tested. *P. berghei* ANKA, 676m1cl1, Green fluorescence protein-luciferase

(PbANKA GFP-Luc) obtained from ATCC (catalog # MRA-868) constitutively expresses

the luciferase at all stages in the life cycle. For each experiment Balb/cj female mice

were infected via intraperitoneal (i.p) with approximately 100,000 erythrocytes infected

with PbANKA GFP-Luc from donor mouse between 5 to 10 % parasitemia (in first or

second passage). For malaria blood stage infections and drug responses, there are not

sex differences in Balb/c mice. As male and female Balb/c mice display differences in

tissue iron levels, female mice were used for consistency throughout all

experiments(41).

Blood stage murine luciferase assay and analysis

During the drug treatment, 5 µL of blood was collected from the tail of each mouse at

regular intervals and deposited into 45 µL of lysis buffer in a 96 well plate (29). Samples

were stored at -80°C until processed. A total of 5 µL of blood/lysis buffer (whole blood

equivalent of 0.5 µL) was transferred to a black, opaque 96-well plate and 95 µl of

luciferase buffer (20 mM Tricine, 100 µM EDTA, 1.07 mM K_2CO_3, 2.67 mM $MgSO_4$, 17

mM DTT, 250 µM ATP, 250 µM D-Luciferin) was added. Luciferase activity was

measured in the IVIS Spectrum In Vivo Imaging System and analyzed using Living

Image v. 4.4 software. The raw luciferase activity is expressed as radiant flux in

photons/second. Total radiant flux was compared to parasites per well using GraphPad

Prism 5 software

Mosquito transmission

Gametocytes were cultured as previously described(42). Gametocytes (mature stages

IV-V) were counted by blood film at day 15 to 16. Gametocyte cultures were then

incubated with fresh medium alone or fresh medium with 10 µM drug for 48 hours, with

medium replaced (with or without drug) after 24 hours exposure. Cultures were washed

three times with fresh medium, and then diluted withRBCs and O+ human serum to a

gametocytemia of 0.05-0.07%, 50% hematocrit, and kept warm at 37°C. Cultures were

fed to 4-6 hour starved *Anopheles stephensi,* using the standard membrane feeding

assay(42). After feeding, blood-fed mosquitoes were sorted and kept. Mosquitoes were

maintained for 8 days, fed with 10% sucrose solution. At day 8, mosquitoes were

dissected and midguts were stained with 1% mercurochrome to count oocysts.

Pharmacokinetics

Pharmacokinetic (PK) data were collected at multiple time points from mice dosed orally

with cethromycin at 30, 60, or 120 mg/kg from plasma, lung, and liver. For cross-tissue

comparisons tissue concentrations in ng/mg were converted to ng/mL by multiplying by

the approximate tissue density (lung ~ 1000 mg/mL; liver ~ 1050 mg/mL).

Noncompartmental analyses were performed using the *PKNCA* (version 0.11.0)

package in R (version 4.4.1). Mean tissue concentration-time profiles were generated

for each dose group and tissue, and PKNCA was used to compute C_{max} and area under

the concentration-time curve (AUC_{0-72}) via the linear-up/log-down trapezoidal method.

To estimate variability, bootstrapped NCA analyses (n=1000 resamples) were

performed for each tissue-dose combination. To capture active elimination kinetics, and

given that concentration-time profiles showed two phases—a rapid initial decline over

24 hours (alpha half-life, $t_{1/2\alpha}$) and a very slow terminal phase >100 hours, likely

reflecting tissue redistribution—we calculated $t_{1/2\alpha}$ by fitting a linear regression of the

log-transformed data for the first 24 hours.

ACKNOWLEDGMENTS

We thank the Johns Hopkins Malaria Research Institute and The Bloomberg Family

Foundation (D.J.S). This work was supported by NCATS R21 TR001737 titled

"Quantum model repurposing of cethromycin for liver stage malaria" (DJS) and

Maryland TEDCO funding (D.J.S. and N.K.). M.M.I. was supported by NIAID K23

AI139343 titled "Clinical pharmacology of malaria elimination" and by the Johns Hopkins

Malaria Research Institute and The Bloomberg Family Foundation. We also thank

Pamela Hummert, MS, in support of the drug concentration analyses.

Author contributions

G.K. and D.S. designed experiments. M.M. designed and implemented pharmacokinetic

results. M. M. and M.I. analyzed PK data. G.K., R.W. and K.P. implemented

immunofluorescent hepatic assays. G.K. and B.B implemented liver stage transmission

assays. R. W. implemented *P. falciparum* drug inhibition assays. G.K., B.B. and D.S.

implemented murine blood stage assays. R.W., N.K. and D.S. implemented mosquito

transmission assays. All authors analyzed the data. D.S. drafted manuscript. All authors

provided critical review.

Declaration of competing interests

DS is Founder, Board Member, and stock/option owner of AliquantumRx(macrolide for

antimicrobial and malaria use); coinventor on USP 7,270,948 (Detection of malaria

parasites by laser desorption mass spectrometry), USP9,568,471 (Malaria diagnosis in

urine), USP 9,642,865 (New angiogenesis inhibitors), and PCT/US2015/046665 (Salts

and polymorphs of cethromycin for the treatment of disease); and has received royalties

from Binax Inc/D/B/A Inverness Medical for plasmids for HRP aldolase for malaria

diagnostic kit. All other authors declare no competing interests. NK is Founder, Board

Member, and stock/option owner of AliquantumRx(macrolide for antimicrobial and

malaria use) and coinventor PCT/US2015/046665 (Salts and polymorphs of

cethromycin for the treatment of disease). Other authors have no competing interests to

declare.

REFERENCES

1. Mansour H, Chahine EB, Karaoui LR, El-Lababidi RM. 2013. Cethromycin: a new ketolide antibiotic. Ann Pharmacother 47:368-79.

2. English ML, Fredericks CE, Milanesio NA, Rohowsky N, Xu ZQ, Jenta TR, Flavin MT, Eiznhamer DA. 2012. Cethromycin versus clarithromycin for community-acquired pneumonia: comparative efficacy and safety outcomes from two double-blinded, randomized, parallel-group, multicenter, multinational noninferiority studies. Antimicrob Agents Chemother 56:2037-47.

3. Georgopapadakou NH. 2014. The wobbly status of ketolides: where do we stand? Expert Opin Investig Drugs 23:1313-9.

4. Vázquez-Laslop N, Mankin AS. 2018. How Macrolide Antibiotics Work. Trends Biochem Sci 43:668-684.

5. Shortridge VD, Zhong P, Cao Z, Beyer JM, Almer LS, Ramer NC, Doktor SZ, Flamm RK. 2002. Comparison of in vitro activities of ABT-773 and telithromycin against macrolide-susceptible and -resistant streptococci and staphylococci. Antimicrob Agents Chemother 46:783-6.

6. Champney WS, Mentens N, Zurawick K. 2004. An examination of the differential sensitivity to ketolide antibiotics in ermB strains of Streptococcus pyogenes and Streptococcus pneumoniae. Curr Microbiol 49:239-47.

7. Davies TA, Ednie LM, Hoellman DM, Pankuch GA, Jacobs MR, Appelbaum PC. 2000. Antipneumococcal activity of ABT-773 compared to those of 10 other agents. Antimicrob Agents Chemother 44:1894-9.

8.   Jacobsson S, Alirol E, Unemo M. 2019. In vitro activity of the ketolide

cethromycin in multidrug-resistant clinical Neisseria gonorrhoeae isolates and

international reference strains. J Chemother 31:246-251.

9.   Khan AA, Araujo FG, Craft JC, Remington JS. 2000. Ketolide ABT-773 is active

against Toxoplasma gondii. J Antimicrob Chemother 46:489-92.

10.   Low LM, Ssemaganda A, Liu XQ, Ho MF, Ozberk V, Fink J, Sundac L, Alcorn K,

Morrison A, O'Callaghan K, Gerrard J, Stanisic DI, Good MF. 2019. Controlled

Infection Immunization Using Delayed Death Drug Treatment Elicits Protective

Immune Responses to Blood-Stage Malaria Parasites. Infect Immun 87.

11.   Wittlin S, Ekland E, Craft JC, Lotharius J, Bathurst I, Fidock DA, Fernandes P.

2012. In vitro and in vivo activity of solithromycin (CEM-101) against Plasmodium

species. Antimicrob Agents Chemother 56:703-7.

12.   Dahl EL, Rosenthal PJ. 2008. Apicoplast translation, transcription and genome

replication: targets for antimalarial antibiotics. Trends Parasitol 24:279-84.

13.   Yeh E, DeRisi JL. 2011. Chemical rescue of malaria parasites lacking an

apicoplast defines organelle function in blood-stage Plasmodium falciparum.

PLoS Biol 9:e1001138.

14.   Amanzougaghene N, Tajeri S, Franetich JF, Ashraf K, Soulard V, Bigeard P,

Guindo CO, Bouillier C, Lemaitre J, Relouzat F, Legrand R, Kocken CHM,

Zeeman AM, Roobsoong W, Sattabongkot J, Yang Z, Snounou G, Mazier D.

2024. Azithromycin disrupts apicoplast biogenesis in replicating and dormant

liver stages of the relapsing malaria parasites Plasmodium vivax and

Plasmodium cynomolgi. Int J Antimicrob Agents 63:107112.

15.   Friesen J, Silvie O, Putrianti ED, Hafalla JC, Matuschewski K, Borrmann S. 2010.

379   Natural immunization against malaria: causal prophylaxis with antibiotics. Sci

380   Transl Med 2:40ra49.

381   16.   Andersen SL, Ager AL, McGreevy P, Schuster BG, Ellis W, Berman J. 1994.

382   Efficacy of azithromycin as a causal prophylactic agent against murine malaria.

383   Antimicrob Agents Chemother 38:1862-3.

384   17.   Anderson SL, Berman J, Kuschner R, Wesche D, Magill A, Wellde B, Schneider

385   I, Dunne M, Schuster BG. 1995. Prophylaxis of Plasmodium falciparum malaria

386   with azithromycin administered to volunteers. Ann Intern Med 123:771-3.

387   18.   Shmuklarsky MJ, Boudreau EF, Pang LW, Smith JI, Schneider I, Fleckenstein L,

388   Abdelrahim MM, Canfield CJ, Schuster B. 1994. Failure of doxycycline as a

389   causal prophylactic agent against Plasmodium falciparum malaria in healthy

390   nonimmune volunteers. Ann Intern Med 120:294-9.

391   19.   Sullivan DJ, Liu Y, Mott BT, Kaludov N, Martinov MN. 2015. Discovery of Novel

392   Liver-Stage Antimalarials through Quantum Similarity. PLoS One 10:e0125593.

393   20.   Fichera ME, Roos DS. 1997. A plastid organelle as a drug target in

394   apicomplexan parasites. Nature 390:407-9.

395   21.   Freireich EJ, Gehan EA, Rall DP, Schmidt LH, Skipper HE. 1966. Quantitative

396   comparison of toxicity of anticancer agents in mouse, rat, hamster, dog, monkey,

397   and man. Cancer Chemother Rep 50:219-44.

398   22.   Araujo FG, Shepard RM, Remington JS. 1991. In vivo activity of the macrolide

399   antibiotics azithromycin, roxithromycin and spiramycin against Toxoplasma

400   gondii. Eur J Clin Microbiol Infect Dis 10:519-24.

23.   Girard AE, Girard D, English AR, Gootz TD, Cimochowski CR, Faiella JA,

Haskell SL, Retsema JA. 1987. Pharmacokinetic and in vivo studies with

azithromycin (CP-62,993), a new macrolide with an extended half-life and

excellent tissue distribution. Antimicrob Agents Chemother 31:1948-54.

24.   Walker LA, Sullivan DJ, Jr. 2017. Impact of Extended Duration of Artesunate

Treatment on Parasitological Outcome in a Cytocidal Murine Malaria Model.

Antimicrob Agents Chemother 61.

25.   Lell B, Kremsner PG. 2002. Clindamycin as an antimalarial drug: review of

clinical trials. Antimicrob Agents Chemother 46:2315-20.

26.   Pukrittayakamee S, Imwong M, Looareesuwan S, White NJ. 2004. Therapeutic

responses to antimalarial and antibacterial drugs in vivax malaria. Acta Trop

89:351-6.

27.   Ranque S, Badiaga S, Delmont J, Brouqui P. 2002. Triangular test applied to the

clinical trial of azithromycin against relapses in Plasmodium vivax infections.

Malar J 1:13.

28.   Deye GA, Gettayacamin M, Hansukjariya P, Im-erbsin R, Sattabongkot J,

Rothstein Y, Macareo L, Fracisco S, Bennett K, Magill AJ, Ohrt C. 2012. Use of a

rhesus Plasmodium cynomolgi model to screen for anti-hypnozoite activity of

pharmaceutical substances. Am J Trop Med Hyg 86:931-5.

29.   Thomas D, Tazerouni H, Sundararaj KG, Cooper JC. 2016. Therapeutic failure of

primaquine and need for new medicines in radical cure of Plasmodium vivax.

Acta Trop 160:35-8.

30. Lu KY, Derbyshire ER. 2020. Tafenoquine: A Step toward Malaria Elimination.

Biochemistry 59:911-920.

31. Hounkpatin AB, Kreidenweiss A, Held J. 2019. Clinical utility of tafenoquine in the

prevention of relapse of Plasmodium vivax malaria: a review on the mode of

action and emerging trial data. Infect Drug Resist 12:553-570.

32. Dean L, Kane M. 2012. Tafenoquine Therapy and G6PD Genotype. *In* Pratt VM,

Scott SA, Pirmohamed M, Esquivel B, Kattman BL, Malheiro AJ (ed), Medical

Genetics Summaries. National Center for Biotechnology Information (US),

Bethesda (MD).

33. Commons RJ, McCarthy JS, Price RN. 2020. Tafenoquine for the radical cure

and prevention of malaria: the importance of testing for G6PD deficiency. Med J

Aust 212:152-153.e1.

34. Santos A, Brito M, Silva E, Rocha F, Oliveira A, Dávila R, Gama H, Albuquerque

J, Paiva M, Baía-Silva D, Sampaio V, Balieiro P, Rufatto R, Grewal Daumerie P,

Peterka C, Edilson Lima F, Jr., Monteiro W, Arcanjo A, Silva R, Batista Pereira D,

Lacerda M, Murta F. 2024. Perspectives of healthcare professionals on training

for quantitative G6PD testing during implementation of tafenoquine in Brazil

(QualiTRuST Study). PLoS Negl Trop Dis 18:e0012197.

35. Adissu W, Brito M, Garbin E, Macedo M, Monteiro W, Mukherjee SK, Myburg J,

Alam MS, Bancone G, Bansil P, Pal S, Sharma A, Zobrist S, Bryan A, Chu CS,

Das S, Domingo GJ, Hann A, Kublin J, Lacerda MVG, Layton M, Ley B, Murphy

SC, Nosten F, Pereira D, Price RN, Talukdar A, Yilma D, Gerth-Guyette E. 2023.

Clinical performance validation of the STANDARD G6PD test: A multi-country

pooled analysis. PLoS Negl Trop Dis 17:e0011652.

36.   Sutanto I, Soebandrio A, Ekawati LL, Chand K, Noviyanti R, Satyagraha AW,

Subekti D, Santy YW, Crenna-Darusallam C, Instiaty I, Budiman W, Prasetya

CB, Lardo S, Elyazar I, Duparc S, Cedar E, Rolfe K, Fernando D, Berni A, Jones

S, Kleim JP, Fletcher K, Sharma H, Martin A, Taylor M, Goyal N, Green JA, Tan

LK, Baird JK. 2023. Tafenoquine co-administered with dihydroartemisinin-

piperaquine for the radical cure of Plasmodium vivax malaria (INSPECTOR): a

randomised, placebo-controlled, efficacy and safety study. Lancet Infect Dis

23:1153-1163.

37.   Hopp CS, Chiou K, Ragheb DR, Salman AM, Khan SM, Liu AJ, Sinnis P. 2015.

Longitudinal analysis of Plasmodium sporozoite motility in the dermis reveals

component of blood vessel recognition. Elife 4.

38.   Trager W, Jensen JB. 1976. Human malaria parasites in continuous culture.

Science 193:673-5.

39.   Flores-Garcia Y, Herrera SM, Jhun H, Pérez-Ramos DW, King CR, Locke E,

Raghunandan R, Zavala F. 2019. Optimization of an in vivo model to study

immunity to Plasmodium falciparum pre-erythrocytic stages. Malar J 18:426.

40.   Lambros C, Vanderberg JP. 1979. Synchronization of Plasmodium falciparum

erythrocytic stages in culture. J Parasitol 65:418-20.

41.   Hahn P, Song Y, Ying GS, He X, Beard J, Dunaief JL. 2009. Age-dependent and

gender-specific changes in mouse tissue iron by strain. Exp Gerontol 44:594-

600.

42.   Tripathi AK, Mlambo G, Kanatani S, Sinnis P, Dimopoulos G. 2020. Plasmodium

falciparum Gametocyte Culture and Mosquito Infection Through Artificial

Membrane Feeding. J Vis Exp doi:10.3791/61426.

Figure legends

Fig 1 **Liver, lung blood pharmacokinetics-** Single oral cethromycin doses were given

to groups of three mice with anticoagulated blood, liver and lung harvested, and frozen

on dry ice after separation of plasma. A) 30 mg/kg B) 60 mg/kg C) 120 mg/kg. Graphs

are mean and standard deviation. Dotted lines are 1 and 10 μM at 765 and 7650 ng/mL

respectively

Fig 2 **Hepatocyte *Plasmodium berghei* apicoplast imaging after drug treatment-**

Drugs at 20 μM were incubated 2 hours after sporozoite Hepa-1 hepatocyte invasion for

48 hours at which time cells were fixed for immunofluorescence. A) Representative

immunofluorescent images at 48 hours stained for nuclei with DAPI, cytosol with *P.*

berghei heat shock protein 70 (HSP70) and the apicoplast with acyl carrier protein

(ACP). B) Segregation of number size and branching of the apicoplast in hepatocytes

under experimental conditions with the addition of cethromycin incubated for the first 24

hours and last 24 hours in addition to 48 hour drug incubations. Experiments were done

in biologic replicates.

Fig 3. **Schema for liver stage dosing schedule with drugs dosed before during and**

after 48 hours *P. berghei* liver stage. Infected mosquitos were sorted with fluorescent

microscopy for mcherry sporozoites and 5-7 mosquitoes were fed on anesthetized mice.

Patency was measured by Giemsa blood film for 16 day after sporozoite infection. Red

labeled drugs indicate partial or all mice patent and green labeled drugs indicate zero

patent mice after liver stage infection.

Fig 4 **Blood stage inhibition** A) *P. falciparum* inhibition- Continuous 72 and 144 hour

P. falciparum 3D7 drug incubations with cethromycin and clindamycin. Assays

performed in biologic duplicates with technical triplicates. B) *P. berghei* blood stage 4

day suppression test with drugs starting 24 hours after inoculation for 4 days. C) High

parasitemia cytocidal test with 2 and 4 days of oral cethromycin and azithromycin D)

High parasitemia test model with 4, 5, 6 and 7 days of cethromycin. One of three mice

recrudesced with 6 days of dosing and none of the 7 day dosing

Female Balb/cj mice (n=3) were inoculated with approximately 1 million infected

erythrocytes from a donor mouse. 4-5 days before drug dosing at about 1 million

parasites per μL. Data is represented as mean ± SEM.

Fig 5 **Late stage gametocyte to oocyst test**- A) Schema for late stage IV and V

gametocytes on day 15 were dose with 10 μM drug for 48 hours, cells washed and

immediately placed in membrane feeding apparatus, after which fed mosquitoes were

sorted and number of oocysts per mosquito counted. At least 20 mosquitoes were

counted.

Table 1 Tissue distribution of cethromycin after single oral doses in mice (n=3 per

group)

Tissue	Dose (mg/kg)	C_{max} (ng/ml)	AUC_{0-72} (ng·h /ml)	$t_{1/2\alpha}$ (h)
Plasma	30	4,767 ± 754	18,682 ± 2,216	1.77 (1.56-2.03)
Plasma	60	5,994 ± 700	34,429 ± 3,572	2.32 (2.40-2.69)
Plasma	120	8,297 ± 604	117,134 ± 9,894	7.53 (5.82-10.66)
Lung	30	481,883 ± 45,681	1,502,310 ± 122,398	1.90 (1.58-2.39)
Lung	60	1,291,870 ± 189,700	4,856,050 ± 616,430	1.70 (1.48-2.00)
Lung	120	2,322,030 ± 83,339	17,565,300 ± 2,061,010	4.21 (3.31-5.80)
Liver	30	1,098,800 ± 253,711	3,585,590 ± 596,799	1.85 (1.61-2.17)
Liver	60	2,063,840 ± 74,153	8,573,030 ± 451,875	2.02 (1.85-2.23)
Liver	120	6,221,820 ± 1,501,440	37,067,800 ± 4,292,660	4.73 (3.64-6.74)

512   Mean and s.d. maximum concentration (C_{max}) and area-under-the-curve (AUC_{0-72}) are

513   shown. The 24-hour half-life ($t_{1/2\alpha}$) was estimated from linear regression of the log-

514   transformed data with 95% confidence intervals shown in parentheses.

516   Table 2 Liver stage patency

Mouse	# Mosq	Drug Groups	Dose mg/kg	day of dose	number pos/total	average day to patency	number positive	total number
BALB/c	7	Cethromycin	30	-1	3/3			
BALB/c	7	Cethromycin	60	-1	3/3			
BALB/c	7	Azithromycin	45	0	3/3	8	3	3
BALB/c	7	Azithromycin	60	0	0/6		0	6
BALB/c	7	Azithromycin	120	0	0/3		0	3
BALB/c	7	Cethromycin	45	0	1/3	8	1	3
BALB/c	7	Cethromycin	60	0	0/3		0	3
BALB/c	7	Cethromycin	120	0	0/2		0	2
BALB/c	7	Azithromycin	15	1	2/3	7	2	3
BALB/c	7	Azithromycin	30	1	3/6	9	3	6
BALB/c	7	Azithromycin	45	1	1/3	13	1	3
BALB/c	5	Azithromycin	60	1	3/6	9	3	6
BALB/c	7	Cethromycin	10	1	3/3			
BALB/c	7	Cethromycin	15	1	5/12	8	5	12
BALB/c	7	Cethromycin	30	1	1/9	10	1	9
BALB/c	7	Cethromycin	45	1	0/3		0	3
BALB/c	5	Cethromycin	60	1	1/9	13	1	9
BALB/c	5	Cethromycin	120	1	0/3		0	3
BALB/c	5	Primaquine	40	1	0/3		0	3
BALB/c	5	Primaquine	60	1	0/3		0	3
BALB/c	7	Cethromycin	60	2	3/3	6		
BALB/c	7	Cethromycin	60	3	3/3	6		
BALB/c	7	Cethromycin	20	-1, 0, 1	0/3			
BALB/c	5	Cethromycin	120	0,1	0/3		0	3
BALB/c	5	Erythromycin Ethyl Succinate	120	0,1	2/2	6	2	2
BALB/c	5	Quinoline	120	0,1	3/3	4	3	3
BALB/c	5	Primaquine	60	1,1	0/3		0	3
BALB/c	7	Untreated			24/24	6	24	24
C57Bl6	7	Azithromycin	45	1	1/3	13	1	3
C57Bl6	7	Cethromycin	45	1	0/3		0	3
C57Bl6	8	Cethromycin	60	1	0/9		0	9
C57Bl6	8	Untreated	n/a		3/3	5	3	3

Fig 1 **Liver, lung blood pharmacokinetics-** Single oral cethromycin doses were given

to groups of three mice with anticoagulated blood, liver and lung harvested, and frozen

on dry ice after separation of plasma. A) 30 mg/kg B) 60 mg/kg C) 120 mg/kg. Graphs

are mean and standard deviation. Dotted lines are 1 and 10 μM at 765 and 7650 ng/mL

respectively.

Fig 2 **Hepatocyte *Plasmodium berghei* apicoplast imaging after drug treatment-**

Drugs at 20 μM were incubated 2 hours after sporozoite Hepa-1 hepatocyte invasion for

48 hours at which time cells were fixed for immunofluorescence. A) Representative

immunofluorescent images at 48 hours stained for nuclei with DAPI, cytosol with *P.*

berghei heat shock protein 70 (HSP70) and the apicoplast with acyl carrier protein

(ACP). B) Segregation of number size and branching of the apicoplast in hepatocytes

under experimental conditions with the addition of cethromycin incubated for the first 24

hours and last 24 hours in addition to 48 hour drug incubations. Experiments were done

in biologic replicates.

Fig 3. **Schema for liver stage dosing schedule with drugs dosed before during and**

after 48 hours *P. berghei* liver stage. Infected mosquitos were sorted with fluorescent

microscopy for mCherry sporozoites and 5-7 mosquitoes were fed on anesthetized

mice. Patency was measured by Giemsa blood film for 16 day after sporozoite infection.

Red labeled drugs indicate partial or all mice patent and green labeled drugs indicate

zero patent mice after liver stage infection.

Fig 4 **Blood stage inhibition** A) *P. falciparum* inhibition- Continuous 72 and 144 hour

P. falciparum 3D7 drug incubations with cethromycin and clindamycin. Assays

performed in biologic duplicates with technical triplicates. B) *P. berghei* blood stage 4

day suppression test with drugs starting 24 hours after inoculation for 4 days. C) High

parasitemia cytocidal test with 2 and 4 days of oral cethromycin and azithromycin D)

High parasitemia test model with 4, 5, 6 and 7 days of cethromycin. One of three mice

recrudesced with 6 days of dosing and none of the 7 day dosing. Female Balb/cj mice

(n=3) were inoculated with approximately 1 million infected erythrocytes from a donor

mouse. 4-5 days before drug dosing at about 1 million parasites per μL. Data is

represented as mean ± SEM.

Fig 5 **Late stage gametocyte to oocyst test**- A) Schema for late stage IV and V

gametocytes on day 15 were dose with 10 µM drug for 48 hours, cells washed and

immediately placed in membrane feeding apparatus, after which fed mosquitoes were

sorted and number of oocysts per mosquito counted. At least 20 mosquitoes were

counted.