(Article)

Phytoplankton recruitment of specific microbial assemblages and

phylosymbiotic patterns

Joo-Hwan Kim[1,a,*], Jin Ho Kim[2,*], Zhun Li[3,*], Hyeon Ho Shin[4], Young Kyun Lim[5], Seung Ho Baek[6],

Pengbin Wang[7], Ro Young Park[1], Soo Hwan Jin[1], Myung-Soo Han[1,b], and Bum Soo Park[1,8,9,10,†]

[1]Department of Life Science, College of Natural Sciences, Hanyang University, Seoul 04763, Republic

of Korea

[2]Department of Earth and Marine Science, College of Ocean Sciences, Jeju National University, Jeju

63243, Republic of Korea

[3]Biological Resource Center/Korean Collection for Type Cultures, Korea Research Institute of

Bioscience and Biotechnology, Jeongeup, 56212, Republic of Korea

[4]Division of Fisheries Life Science, Pukyong National University, Busan 48574, Republic of Korea

[5]Ocean Climate Response & Ecosystem Research Department, Korea Institute of Ocean Science &

Technology, Busan 49111, Republic of Korea

[6]Ecological Risk Research Department, Korea Institute of Ocean Science & Technology, Geoje 53201,

Republic of Korea

[7]Key Laboratory of Marine Ecosystem Dynamics, Second Institute of Oceanography, Ministry of

Natural Resources, Hangzhou, 310012, China

[8]Hanyang Institute of Bioscience and Biotechnology, Hanyang University, Seoul 04763, Republic of

Korea

[9]Research Institute for Convergence of Basic Science, Hanyang University, Seoul 04763, Republic of

Korea

[10]Hanyang Institute of Advanced BioConvergence, Hanyang University, Seoul 04763, Republic of

Korea

* Co-first authors

† Corresponding author contact information:

Department of Life Science, College of Natural Sciences, Hanyang University, Seoul 04763, Republic

of Korea, Tel: +82-2-2220-0961, E-mail: parkbs@hanyang.ac.kr

Present address:

[a]Ministry of Environment, Government Complex-Sejong, Sejong 30103, Republic of Korea

[b]Mirae Shangtae (Co., Ltd.), R&D Headquarters, 10, songi-ro 24-gil, Songpa-gu, Seoul 05829,

Republic of Korea

Abstract

Phytoplankton and bacteria represent two major pillars of the carbon cycle in marine ecosystems.

While their interactions are known to be tightly linked, the specific mechanisms underlying these

interactions remain largely unexplored. Evidence of host-specific microbial assemblages could serve

as a foundation for studies of detailed host-microbe interactions, yet such research remains limited in

phytoplankton. Here, we not only investigate the microbial assemblages of six phytoplankton species,

including multiple strains of dinoflagellates and diatoms, but also samples of phytoplankton blooms

from the field. Our results reveal the presence of host-specific microbial assemblages in

phytoplankton, with members of the core microbial lineages (MCGs) playing pivotal roles in shaping

host-specific microbial assemblages and contributing to network structures. Deterministic processes,

particularly host genotypes, were the dominant factors shaping microbial assemblages, overriding

environmental influences. Consequently, microbial composition reflected the evolutionary

relationships of the host species, demonstrating phylosymbiotic patterns. These findings suggest that

studying MCGs will be a crucial foundation for investigating specific phytoplankton-microbe

interactions and highlight the ecological and evolutionary importance of host-microbial specificity in

phytoplankton.

Main

Microbial communities in ecosystems play crucial roles in host nutrition, metabolism, protection from pathogens, and regulation of physiological processes such as immunity and development[1, 2, 3, 4, 5]. Host-specific microbial assemblages have been documented in a wide range of organisms, such as vertebrates[6, 7, 8, 9, 10, 11], invertebrates[9, 12, 13], and plants[14, 15], providing clear evidence of host-microbiota coevolution and mutual benefits. Research into host-specific microbiomes is critical for understanding how these microbial partnerships influence the fitness, ecological dynamics, and persistence of host species. In some cases, host specificity is observed as phylosymbiosis, in which microbial assemblages reflect the evolutionary relationships of their hosts[15, 16, 17, 18]. However, this pattern is not universal, as not all marine invertebrates and terrestrial mammals exhibit phylosymbiosis[19, 20].

As key primary producers in aquatic ecosystems, phytoplankton are responsible for approximately half of Earth's oxygen production and are crucial for the biogeochemical cycling of carbon, nitrogen, sulphur, and iron[21, 22, 23]. Simultaneously, bacteria play central roles in the decomposition and remineralization of organic matter, forming the functional backbone of these ecosystems[24, 25]. Phytoplankton and bacteria are closely intertwined, with phytoplankton providing dissolved organic carbon, and bacteria supplying essential nutrients such as bioavailable nitrogen and phosphorus, as well as growth-promoting compounds such as vitamins[25, 26, 27]. These reciprocal interactions are particularly evident during algal blooms, during which bacterial communities actively support phytoplankton growth and shape bloom dynamics[27, 28, 29, 30, 31].

Recent findings have shown that bacteria can promote the growth of specific phytoplankton by regulating their aging physiology, revealing intricate mechanisms underlying these interactions[32]. However, the details of specific interactions between phytoplankton and bacteria remain largely unexplored. A promising starting point for studying these intricate relationships is to examine whether phytoplankton hosts harbour distinct host-specific microbial assemblages.

Although some studies provide evidence for host-specific microbial assemblages in phytoplankton[29, 33, 34, 35, 36, 37, 38], in part because many analyses were limited to comparisons within a single species or between species using only one strain per species, most research focuses on diatoms, despite the critical ecological roles of dinoflagellates in aquatic systems[39, 40]. Many studies relied on laboratory

strains, leaving host-specific microbial assemblages in natural environments largely unexplored.

In this study, we address these gaps by investigating host-specific microbial assemblage patterns in

both laboratory-cultured strains and samples of phytoplankton blooms collected from the field. We

examined six microalgal species, including diatoms and dinoflagellates, across multiple strains and

growth stages, analysing core microbial lineages. Our results provide strong evidence for host-

specific microbial assemblages in phytoplankton, even in highly mixed aquatic environments, and

identify potential phylosymbiotic patterns between phytoplankton hosts and their microbial

assemblages.

Results

Host-specificity of microbial assemblages across phytoplankton species

Microbial assemblages associated with six phytoplankton species exhibited distinct compositions

depending on the host species (Fig. 1a). Among dinoflagellates, the dominant bacterial taxa were

Rhodobacterales (Alphaproteobacteria) and Alteromonadales (Gammaproteobacteria). Sub-dominant

taxa varied by species: *Flavobacteriales* (Flavobacteriia) were more abundant in *Alexandrium*

catenella (Group I) and *Gymnodinium catenatum*, while members of Cytophagales (Cytophagia)

showed higher relative abundance in *A. catenella* and *A. pacificum* (Group IV). In diatoms,

Rhodobacterales consistently dominated in *Cylindrotheca closterium* and *Coscinodiscuss granii*,

whereas *Alteromonadales* dominated in *Pseudo-nitzschia pungens*. Sub-dominant taxa also varied by

species, including Bacillales (Bacilli) in *Cylindrotheca closterium*, Vibrionales (Gammaproteobacteria)

in *P. pungens*, and a mix of Alteromonadales and Flavobacteriales in *Coscinodiscuss granii*.

Dinoflagellates were significantly enriched in nine bacterial families, including Alteromonadaceae

(Gammaproteobacteria), Flavobacteriaceae (Flavobacteriia), and Flammeovirgaceae (Cytophagia),

compared with diatoms (linear discriminant analysis [LDA] score > 3.12, adjusted p-value < 0.05). In

contrast, Bacillaceae (Bacilli) was the only significantly enriched family in diatoms (Extended Data Fig.

1a). Alpha diversity was significantly higher in dinoflagellates than in diatoms, with dinoflagellates

associated with a greater Shannon index (2.54 vs. 1.46) and Chao1 values (125.6 vs. 65.18) (p <

0.001) (Extended Data Fig. 1b and c). This indicates that dinoflagellates harbour a more diverse array

of microbial lineages.

Beta diversity analysis revealed host-specific microbial assemblages which involve significant
differences in the microbial assemblages of each phytoplankton host based on Bray–Curtis
dissimilarity and confirmed by principal coordinates analysis (PCoA) and permutational multivariate
analysis of variance (PERMANOVA, R^2 = 0.794, p < 0.001) (Fig. 1b). Most microbial assemblages
were clustered with the microbial assemblages of the same phytoplankton species isolated from
different spatiotemporal contexts (Extended Data Table 1, Extended Data Fig. 2b). For example,
although the two strains of *Cylindrotheca Closterium* were separated spatially by 375 km and
temporally by 347 days, they were clustered into the same clade in the hierarchical clustering analysis,
indicating that they share a conserved microbial assemblage and are distinct from other
phytoplankton species (Extended Data Fig. 3). Host-specificity in microbial assemblages was evident
not only in Bray–Curtis dissimilarity results but also across other beta diversity metrics, consistently
showing statistically significant distinctions (Extended Data Table 2).

Distinct microbial compositions were observed across different growth stages of the host species,
highlighting growth-specificity of microbial assemblages. For example, in *A. catenella*, the relative
abundance of Rhodobacterales increased, whereas Kiloniellales decreased during the stationary
phase compared with the lag phase. This pattern was consistent across all strains and replicates
examined (Fig. 1c). PERMANOVA analysis supported these observations, showing that the microbial
assemblage during the stationary phase of *A. catenella* strain AC-1 was more similar to the stationary
phase microbial assemblage of strain AC-2 than to its own assemblage during the lag or log phases
(R^2 = 0.82, p < 0.001). No significant difference between strains was evident (R^2 = 0.05, p > 0.05),
indicating that the growth stage, rather than strain, plays a dominant role in shaping microbial
assemblages. These growth phase–dependent variations were more prominent in dinoflagellates but
were only observed in *Cylindrotheca closterium* among diatoms. While the microbiomes of
dinoflagellates were highly similar between strains, several strain-specific differences were noted in *P.*
pungens and *Coscinodiscuss granii* among diatoms (Fig. 1c). In *P. pungens* and *Coscinodiscuss*
granii, no significant differences in microbiome assembly were observed between growth stages,
although statistically significant strain-level differences were seen (R^2 = 0.24, p < 0.05; R^2 = 0.95, p <
0.001, respectively).

Host-specific microbial assemblages were evident in phytoplankton bloom samples taken from the

field, with microbial communities clustering distinctly according to the host species of the bloom (Fig.

1d, Extended Data Fig. 4). For example, microbial assemblages associated with blooms caused by

Margalefidinium polykrikoides (found in both Korean coastal waters and those of the United States),

Akashiwo sanguinea (Korea), *Alexandrium monilatum* (United States), and *Phaeocystis globosa*

(China) formed tight clusters distinct from one another in Bray–Curtis dissimilarity-based PCoA

analysis (PERMANOVA, R^2 = 0.36, $p < 0.05$) (Extended Data Fig. 2, Extended Data Table 3).

Microbial assemblages were more strongly differentiated by the host phytoplankton species (R^2 = 0.36)

than by the geographical location of the bloom (R^2 = 0.27), highlighting the dominant influence of host

species on community structure ($p < 0.05$).

Core and host-specific microbial lineages across dinoflagellates and diatoms

In laboratory-cultured strains, the number of core microbial lineages (core MLs) varied by host

species, ranging from 5 to 33, excluding *Coscinodiscuss granii*, which lacked observable core MLs

(Fig. 2a). Host-specific microbial lineages (HS-MLs) which were statistically significantly different and

uniquely associated with individual hosts ranged from 1 to 16 (Fig. 2a, 2b). Notably, nearly all HS-MLs

(45 of 47 microbial lineages) overlapped with core MLs, highlighting their strong association (Fig. 2a).

Among the top 200 microbial lineages for each host species, the proportion of core MLs and HS-MLs,

collectively termed the members of the core MLs group (MCG), ranged from 3% to 17%, indicating

that MCGs were numerically fewer than non-MCGs (Fig. 2c). However, MCGs accounted for a

substantial portion of the total rarefied abundance for each host species, contributing 80.8% to 96.2%,

suggesting that MCGs, while limited in diversity, dominate the microbial community quantitatively.

MCGs shared between different hosts were also observed. The number of MCGs shared between

host pairs ranged from 1 to 3, with nearly all shared MCGs found exclusively within dinoflagellates

(Fig. 2d). Additionally, four MCGs were identified in the dinoflagellates *A. catenella*, *A. pacificum*, and

G. catenatum, three of which belonged to the family Alteromonadaceae, genus *Marinobacter* (the

fourth was from an unidentified genus within Alteromonadaceae). In contrast, no shared MCGs were

found among diatom hosts. Dinoflagellates consistently harboured more MCGs (17–34 per species)

compared with diatoms (0–11 per species).

Among dinoflagellates, the bacterial taxa associated with MCGs primarily included Rhodobacterales

(Alphaproteobacteria), Alteromonadales (Gammaproteobacteria), and Flavobacteriales

(Flavobacteriia), closely reflecting the taxonomic composition of all the microbial lineages detected in

dinoflagellates (Fig. 2e). In contrast, diatoms exhibited a more restricted and simplified taxonomic

composition within their MCGs. For example, while Bacilli and Flavobacteriia constituted 9.1% and

8.5%, respectively, of the total microbial lineages observed in diatoms, neither was represented

among MCGs. Instead, MCGs in diatoms were dominated by Rhodobacterales (50.0%) and

Alteromonadales (43.8%), suggesting that the core microbiome of diatoms is not only smaller but

more taxonomically constrained compared with that of dinoflagellates.

MCGs played a major role in driving host-specific microbial assemblages in phytoplankton species.

Although MCGs collectively accounted for only 77 of the top 242 MLs observed in this study, a PCoA

and PERMANOVA based solely on these MLs found stronger differentiation among host species

compared with analyses of the remaining 165 MLs (non-MCGs) (Fig. 2f). Specifically, weighted

UniFrac-based PERMANOVA results for MCG ($R^2 = 0.877$, $p < 0.0001$) showed significantly greater

intergroup differences compared with those observed for the remaining MLs ($R^2 = 0.516$, $p < 0.0001$)

or the full dataset of 242 MLs ($R^2 = 0.794$, $p < 0.001$). Consistent trends were also observed across

beta diversity metrics, such as Bray–Curtis dissimilarity, further supporting the robustness of these

findings (Table 1). The difference in Bray–Curtis dissimilarity between species and within species was

larger when calculated using MCG compared with calculations using non-MCGs or the total

microbiome, highlighting the high contribution of MCGs to host specificity (Fig. 2g and h).

Functional group differences between dinoflagellate and diatom microbiomes and between

core MLs group and other MLs

Functional annotation of prokaryotic taxa (FAPROTAX) analysis revealed significant differences in

bacterial functional groups between dinoflagellates and diatoms (Fig. 2i, Extended Data Fig. 5).

Notably, all functional groups showing significant differences were more enriched in dinoflagellates

compared with diatoms. The functional group with the greatest difference between these two

microalgal groups, based on LDA effect size, was "hydrocarbon degradation". Other notable functional

groups enriched in dinoflagellates included those related to carbon metabolism, such as

"methylotrophy" and "methanol oxidation", as well as groups associated with nitrogen and sulphur

recycling and biopolymer degradation.

Core MLs exhibited distinct functional profiles compared with other MLs, with chemoheterotrophic

functions being significantly enriched across all host species (Fig. 2j). Additionally, hydrocarbon

degradation was a prominent function observed in the core MLs of most hosts. Some functional

differences among core MLs were also noted across hosts. For example, core MLs in *A. catenella*

were enriched in "xylanolysis", while those in *A. pacificum* exhibited functions associated with

"intracellular parasites". In *G. catenatum*, "methylotrophy"-related functions were prominent, whereas

core MLs in *P. pungens* were enriched in functions related to "fermentation" and "nitrate reduction".

Functions observed in HS-MLs overlapped significantly with those in core MLs, including key

functions enriched in specific hosts, such as hydrocarbon degradation and chemoheterotrophy.

Network analysis of microbiomes by microalgal host species

Co-occurrence network analysis of the microbiomes associated with each microalgal host revealed

that bacterial network connectivity within dinoflagellate microbiomes was generally weaker compared

with diatom microbiomes (Fig. 3a). Specifically, the clustering coefficient was significantly lower in

dinoflagellates (0.57) than in diatoms (0.73) (Fig. 3b). Although the average degree (31.68 in

dinoflagellates versus 65.46 in diatoms) and connectance (0.16 in dinoflagellates versus 0.33 in

diatoms) also differed, these differences were not statistically significant.

Within dinoflagellate microbiome networks, MCG nodes exhibited significantly higher degree,

closeness, and betweenness centrality compared with other nodes, emphasizing their critical role in

network structure (Fig. 3c). In contrast, in diatom microbiomes, other nodes showed higher degrees

and closeness centrality compared with MCG nodes, indicating that the relative importance of MCG

nodes in diatom networks was lower than in dinoflagellates.

Deterministic influences and phylosymbiotic patterns in microalgal microbiome assembly

To evaluate the relative contributions of stochastic and deterministic processes in shaping the host-

specific microbiome assembly, we analysed the normalized stochasticity ratio (NST) of the Jaccard

matrix (NST_{jac}). This metric distinguishes between stochastic (NST > 50%) and deterministic (NST <

50%) influences on community structure. The results indicate that microbiome structures in both

dinoflagellates and diatoms were predominantly shaped by deterministic factors (Fig. 4a).

Dinoflagellates exhibited a mean NST value of 16.8 ± 3.1%, with a maximum value of 32.3%,

reflecting a strong deterministic control. In contrast, diatoms showed a higher mean NST value of 32.4

± 13.2% and a maximum of 90.1%, suggesting that stochastic influences were more prominent in

diatom microbiomes compared with dinoflagellates.

Correlation analysis to identify deterministic factors influencing microbial assemblages revealed that

the dissimilarity between microalgal-associated microbiomes was not significantly correlated with

physicochemical factors, including temperature, salinity, pH, dissolved oxygen (DO), total nitrogen

(TN), total phosphorus (TP), chemical oxygen demand (COD), chlorophyll-a (Chl-a), or spatiotemporal

distance at the time of host strain isolation. However, a strong positive correlation was observed with

the phylogenetic distance between hosts (R = 0.61, p < 0.001) (Fig. 4b). Specifically, microbiome

similarity increased with decreasing evolutionary distance between hosts. This relationship was

notably stronger in dinoflagellates (R = 0.80, p < 0.001) compared with diatoms (R = 0.33, p < 0.001),

indicating a more pronounced influence from host genetic relatedness on microbiome assembly in

dinoflagellates.

To further investigate phylosymbiotic patterns, we performed Robinson–Foulds and matching cluster

analyses, which showed complete congruence (normalized distance = 0.0) between the molecular

phylogeny of the host phytoplankton and the dendrogram topology of their microbiome dissimilarity

across both evaluation methods (nRF = 0.00, p < 0.01; nMC = 0.00, p < 0.01) (Fig. 4c). The

microbiome dendrograms could be divided into two major groups: dinoflagellates and diatoms. Within

these groups, the microbiomes reflected the phylogenetic traits of their hosts, with dinoflagellate

microbiomes clustering into unarmoured and armoured subgroups, while diatom microbiomes

separated into pennate and centric subgroups. Additionally, congruence was also observed when

using unweighted UniFrac to calculate microbiota distances, although no significant congruence was

detected with weighted UniFrac (Table 2).

Discussion

In this study, through comparative analyses of both laboratory-cultured strains and field-collected

bloom samples, we found evidence suggesting that microbial assemblage in phytoplankton exhibited

a degree of host-specificity. We found that strains of the same microalgal species, isolated from

spatiotemporally distant locations, exhibited highly similar microbiome assemblies. Even

phytoplankton strains isolated from the same time and location, such as *A. catenella* (AC-2), *A.*

pacificum (AP-2), and *G. catenatum* (GC-1), shared only 4 MLs, and these MLs were consistent

detected in MLs of strains isolated from different times and locations, such as *A. catenella* (AC-1), *A.*

pacificum (AP-1), and *G. catenatum* (GC-2), indicating microbial assemblages in phytoplankton are

highly preserved and have their own specificity (Extended Data Fig. 6). The region surrounding

phytoplankton cells, known as the phycosphere, is rich in organic matter, which is known to support

active interactions with bacteria[27, 41]. Our findings suggest that the phytoplankton phycosphere is

remarkably robust in maintaining its microbial assemblages, likely facilitated by active host-microbe

interactions within this microenvironment.

While previous studies have explored host-specific microbial assemblages in phytoplankton, they

primarily focused on single species or limited geographical regions[28, 29, 36, 38, 42, 43, 44]. By analysing

samples of bloom samples collected from multiple geographically distinct locations in the field, our

study provides novel insights into host-specificity across multiple phytoplankton species under natural

conditions. Notably, while the similarity of microbial assemblages within the same species was

moderately lower in samples of phytoplankton blooms compared with lab-cultured strains, the degree

of similarity remained high in natural environments. This difference can be attributed to the

methodologies used for sample preparation. Lab-cultured strains, established through single-cell

isolation, are expected to retain microbes that are physically closely associated with the host cells. In

contrast, field-collected bloom samples, analysed by filtering entire seawater samples, likely included

microbes unrelated to the host. As a result, host specificity in microbial assemblages was moderately

reduced in field samples. Despite this reduction, microbial assemblages associated with the same

phytoplankton species exhibited remarkable similarity across field samples, even when collected from

geographically and temporally distant locations. This underscores the robustness of host-specific

microbial assemblages in phytoplankton, even under the dynamic conditions of natural environments.

The observation that phytoplankton, which inhabit a homogenous water column, maintain host-

specific microbial assemblages is particularly intriguing. Most studies reporting host-specific microbial

assemblages focused on heterogeneous environments, such as the gut microbiomes of vertebrates

and invertebrates, or root-associated microbiomes of terrestrial plants[6, 13, 14, 15, 18, 45, 46, 47]. Our findings

indicate that even phytoplankton, which lack organs such as a gut, can exert a significant influence on

the structure of their associated microbiomes.

A key factor underlying the host specificity observed in phytoplankton microbiomes is the role of

MCGs, which dominate host-associated microbial assemblages and contribute significantly to their

distinct composition. Despite representing a smaller fraction of microbial diversity compared with non-

MCGs, MCGs dominate host microbiomes and contribute significantly to their host specificity. The

majority of MCGs identified in our study belong to the orders Rhodobacterales, Alteromonadales, and

Flavobacteriales, which have been frequently associated with phytoplankton cultures and blooms in

previous studies[25, 48, 49, 50]. Among these, members of the family Roseobacteraceae (Order

Rhodobacterales) dominated the Alphaproteobacteria class, accounting for 86 of 122 microbial

lineages. Roseobacteraceae species are known to utilize various organic compounds released by

phytoplankton, such as dimethylsulphoniopropionate and amino acids, and to participate in key

biogeochemical processes involving carbon, nitrogen, sulphur, and phosphorus cycling. Additionally,

they produce vitamins and growth-promoting compounds that benefit phytoplankton[51, 52].

Alteromonadaceae (Order Alteromonadales) break down diverse polysaccharides, using

carbohydrate-active enzymes encoded in polysaccharide utilization loci[53]. This group is particularly

adept at degrading complex biopolymers, such as laminarin, alginate, xylan, and mannans, facilitated

by their large genomes and extensive enzymatic repertoire. Similarly, Flavobacteriia use

polysaccharides and proteins as primary sources of carbon and energy, efficiently recycling organic

material and nutrients from phytoplankton blooms by processing photosynthetic byproducts or

degrading algal cell walls[30, 54, 55, 56]. Together, these MCGs are likely to play complementary roles in

the remineralization of organic matter within the phytoplankton microenvironment, potentially

contributing to localized nutrient cycling and energy flow.

Beyond the dominance of MCGs in host microbiomes, their ecological significance lies in their diverse

functional capabilities, which can be characterized using FAPROTAX predictions and reveal key

metabolic traits crucial for interactions with phytoplankton hosts. Across all host species, MCGs

exhibited distinct functional profiles compared with non-MCGs, with chemoheterotrophic functions

being significantly enriched. This indicates that MCGs are likely highly dependent on host-derived

organic matter, and suggests a commensal or even mutualistic relationship between phytoplankton

hosts and their associated MCGs. Hydrocarbon degradation was a prominent function consistently

observed in the MCGs of most hosts, far exceeding its representation in non-MCGs. Among

dinoflagellates, the genus *Marinobacter* (family Alteromonadaceae) was commonly detected in the

MCGs of all host species. *Marinobacter* is a well-known hydrocarbon-degrading bacterium, and one

lineage, OTU22, was found to share 99.57% of its gene sequence with *Marinobacter algicola*, a

hydrocarbon-degrading bacterium previously isolated from *Alexandrium tamarense* and *Gymnodinium*

catenatum[57]. This highlights the inclusion of hydrocarbon-degrading bacteria in the core microbiome

of dinoflagellates, a result consistent with previous studies[42, 58]. Such associations suggest a

particularly close ecological relationship between these microbial lineages and dinoflagellates. Some

phytoplankton species, including dinoflagellates, are known to produce hydrocarbons[59, 60, 61, 62], which

may explain the frequent presence of bacteria capable of hydrocarbon degradation. However, it

remains unclear whether such bacteria are associated with all phytoplankton because they

metabolize host-derived hydrocarbons or because they provide essential benefits critical for host

survival. Previous studies have shown that hydrocarbon-degrading bacteria can promote

phytoplankton growth by supplying essential nutrients or growth-promoting compounds[63, 64]. Further

research is needed to clarify whether these interactions constitute a mutualistic relationship and

elucidate the underlying mechanisms driving these associations.

As revealed by a microbiome co-occurrence network analysis, the MCG nodes in dinoflagellates

played more significant roles in network connectivity compared with other nodes. This suggests that

MCGs may act as ecological bridges, not only mediating interactions with the host but facilitating

connections among other bacterial taxa. MCGs appeared to contribute substantially to maintaining the

overall microbial assemblages of their hosts. Meanwhile, the co-occurrence network results for

dinoflagellates, in which bacterial networks were weaker than those observed in diatoms, imply that

dinoflagellate-associated bacterial communities are more tightly regulated by the host. This pattern

resembles the weakening of microbial network stability seen under environmental stress, in which

microbial assemblages across samples exhibit minimal variation and high similarity [65].

The weaker network connectivity observed in dinoflagellate microbiomes can be attributed to their

host-specific microbial assemblages, which are more pronounced compared with those in diatoms.

This is supported by the larger number of MCGs found in dinoflagellates and the higher similarity

values within species than between species. The within-species similarity observed in our study aligns

with that reported previously in research on microalgal microbiomes. The bacterial community

structures associated with dinoflagellates were largely consistent at the order level, while diatoms

exhibited greater variability. For example, in *A. catenella*, both our study and previous research

identified Rhodobacterales, Cytophagales, and Flavobacteriales as dominant taxa[44]. In contrast,

during the exponential phase of *Cylindrotheca closterium*, Rhodobacterales consistently emerged as

the dominant taxon, aligning with our findings. However, unlike *A. catenella*, the composition of

secondary dominant taxa in *Cylindrotheca closterium* varied between studies[43]. In *P. pungens*, earlier

studies reported Rhodobacterales was the dominant taxon, with minimal representation of

Alteromonadales. In our study, however, Alteromonadales was dominant, with Rhodobacterales being

subdominant[33]. This suggests that diatom microbiomes are less conserved and more dynamic

compared with dinoflagellate microbiomes.

These findings align with previous reports indicating that diatom microbiomes, while generally

conserved, can vary depending on environmental conditions at the time of strain isolation[43, 66]. This

flexibility in microbiome composition was most pronounced in *Coscinodiscuss granii*, which lacked

observable core MLs, whereas the microbiomes of the other two diatom species were more stable.

This observation does not entirely align with studies reporting a highly conserved microbiome

structure in diatoms[33, 34, 35]. However, the taxonomic diversity within diatoms, categorized into three

major classes — Coscinodiscophyceae, Mediophyceae, and Bacillariophyceae — provides a

plausible explanation[67, 68]. In this study, *Coscinodiscuss granii* belongs to Coscinodiscophyceae, while

the other two species are members of Bacillariophyceae. The variation in microbiome conservation

observed across diatom species and studies may reflect evolutionary and coevolutionary dynamics

between hosts and their microbiomes, underscoring the need for further research to elucidate these

relationships[47, 69, 70].

The reason for the greater microbial assemblage similarity among dinoflagellate hosts compared to

diatom hosts is uncertain. Many animals and plants regulate their microbiomes to benefit the host by

producing metabolites, such as adhesive molecules, antimicrobial peptides, and hormones[15, 71, 72, 73].

Similarly, host-specific microbial assemblages in phytoplankton are thought to be strongly shaped by

host-derived metabolites. For example, diatoms selectively modulate surrounding bacteria through the

production of secondary metabolites[74]. Other studies demonstrated that manipulating the ratios of

representative metabolites produced by two phytoplankton species can not only alter the bacterial

community composition in natural seawater consortia but also produce predictable outcomes[75, 76].

This serves as an example of how microbial assemblages can be regulated by phytoplankton hosts.

The relatively strong host specificity observed in the microbiomes of dinoflagellates compared with

that in diatoms may suggest that dinoflagellate-derived metabolites exert stronger selective pressures

and that dinoflagellates may rely more heavily on their microbiomes. Results of a FAPROTAX analysis

showed that dinoflagellates harbour a much richer array of bacterial functional groups compared with

diatoms. Although the specific benefits these functional groups provide to the host remain unclear,

dinoflagellates may depend more heavily on their microbiome's functions and may apply stricter

filtering to recruit beneficial bacteria[77].

The microbial assemblage is influenced by various factors, including selection, dispersal,

diversification, and drift [78, 79, 80, 81]. These factors are typically categorized into deterministic and

stochastic processes. While traditional niche-based ecological perspectives emphasize that

environmental factors play a significant role in shaping communities through deterministic processes[82],

recent studies suggest that stochastic factors may better predict variations in community structure[83, 84,

[85]. In this study, microbial assemblages observed in both dinoflagellate and diatom microbiomes were

predominantly governed by deterministic processes. However, NST analysis of diatom microbiomes

revealed a greater prevalence of stochastic influences, with values exceeding the 50% threshold. This

suggests that diatom microbiomes may be shaped by priority effects and other stochastic factors[38, 86].

In our analysis, deterministic environmental factors such as temperature, salinity, pH, DO, COD, TN,

TP, and Chl-*a*, as well as the spatiotemporal location of strain isolation, were considered. However,

the factor most strongly correlated with microbiome dissimilarity (Bray–Curtis) between samples was

not the physicochemical environment or spatiotemporal distance, but the evolutionary distance

between hosts. This aligns with the results of previous studies showing that a host's genotype can

significantly influence its microbiome[38]. It also suggests that microalgal microbiomes are shaped

primarily by host filtering, with selection acting as the dominant deterministic factor in microbiome

assembly.

The proposition that microbiome similarity increases as evolutionary relationships between hosts

converge is central to the emerging idea of phylosymbiosis, which has attracted significant attention

from ecologists over the past decade[70, 86, 87]. Phylosymbiosis has been observed in various organisms,

including insects, mammals, and marine invertebrates, although some taxa do not exhibit this pattern

[9, 20, 88]. To our knowledge, this is the first study to suggest that phytoplankton associated microbiomes

may exhibit phylosymbiotic characteristics. We found evidence of phylosymbiosis in phytoplankton

microbiomes despite their unicellular structure and microscopic size. This contrasts with findings

reported for marine microscopic invertebrates, where such features are absent[20]. However, the

congruence between microbiota dendrograms and host phylogenies varied depending on the beta

diversity metrics used, highlighting the need for further validation. Given that this study examined only

six host species and two strains per species, additional research with a broader range of host taxa

and multiple strains is required to confirm the phylosymbiotic nature of phytoplankton-associated

microbiomes. Moreover, future studies should explore whether phylosymbiosis is a universal trait

among phytoplankton and identify the mechanisms driving this pattern, such as host-derived

metabolites or ecological interactions shaping microbial community assembly.

Conclusion

This study identifies MCGs in phytoplankton microbiomes and describes their contribution to host-

specific microbial assemblages. Investigating the interactions between host-specific MCGs and their

phytoplankton hosts will be critical for elucidating previously unknown mechanisms of phytoplankton-

bacteria interactions. Additionally, phylosymbiotic patterns were observed between phytoplankton and

their microbiomes, suggesting a more interdependent co-evolutionary relationship than previously

recognized. These findings underscore the necessity of jointly considering phytoplankton and

bacterial communities to comprehensively understand their ecological dynamics and evolutionary

interplay in marine environments.

Methods

Microalgal strain collection and bloom sampling

To analyse the microbiomes of phytoplankton, three dinoflagellate species and three diatom species

were selected (hereafter referred to as "lab microbiomes"). To minimize the impact of environmental

factors on microbiome similarity, two strains with different spatiotemporal origins were used for each

microalgal host species. For example, two strains of *Cylindrotheca closterium* were isolated from

locations 375 km and 347 days apart. All algal strains were obtained from the Library of Marine

Samples of the Korea Institute of Ocean Science and Technology in Geoje, South Korea. Information

on the environmental conditions at the time of strain isolation (e.g., temperature, salinity, pH,

dissolved oxygen, COD, TN, TP, and Chl-a) was provided (Extended Data Table 1). All strains were

cultured in duplicate at 20°C in an F/2 growth medium [89] with a salinity of 31–33 under cool-white

fluorescent lamps (photon flux of 100 μE m^{-2} s^{-1}) on a 12-h light/12-h dark cycle.

To observe microbiome composition changes across growth stages for each host, samples of 300 mL

were taken during the lag, log, and stationary phases based on cell density. The samples were size-

fractioned using 47 mm diameter Isopore membrane filters (Millipore, Cork, Ireland) with a pore size

of 0.22–3.0 µm, and the filters were stored in an EX buffer (100 mM Tris-HCl, 100 mM Na2-EDTA,

100 mM sodium phosphate, 1.5 M NaCl, 1% CTAB) at –80°C until DNA extraction[90].

To compare microbiomes across field samples ("field microbiomes"), blooms of *M. polykrikoides*, a

major harmful algal bloom dinoflagellate species in Northeast Asia, were collected from four different

locations along the Korean coast in 2012 and 2015 (Extended Data Table 4). Additionally, samples of

Akashiwo sanguinea (Dinophyceae) blooms from 2011 were included for comparison. Seawater

samples were collected at a depth of <1 m using a 4.2 L Van Dorn water sampler (Wildlife Supply

Company, MI, USA). Volumes of 50–300 mL of seawater were filtered by gravity onto Isopore filters

(Millipore, Cork, Ireland) with a pore size of 5 µm, and the filters were submerged in the EX buffer and

stored at –80°C until further analysis.

In addition to the field samples, previously published microbiome data from other microalgal blooms

were included for comparison. Microbiome data from *M. polykrikoides*, *Alexandrium monilatum*, *A.*

sanguinea, and *Phaeocystis globosa* (Prymnesiophyceae) blooms were retrieved from the National

Center for Biotechnology Information (NCBI) Sequence Read Archive (PRJNA731462,

PRJNA1061360, PRJNA685631, PRJNA678802)[28, 91, 92, 93] (Extended Data Table 3).

DNA extraction and sequencing of 28S rRNA gene

Extraction of host algal DNA was performed on cell pellets from the cultured isolates following a

protocol supplied with the DNeasy Plant Kit (Qiagen, Valencia, CA, USA). Polymerase chain reaction

(PCR) amplification was performed with 50 µL reaction mixtures containing 23 µL of sterile distilled

water, 10 of µL of 5× Phusion HF buffer (New England Biolabs, Beverly, MA, USA), 1 µL of a dNTP

mixture (2.5 mM each), 2.5 µL of each primer (10 pmol), 1 µL of Phusion DNA polymerase (2.0

units/50 µL), and 10 µL of template DNA. Partial large subunit (LSU) rDNA sequences were amplified

using primers described in Scholin et al. (1994). The ITS-5.8S-ITS2 rDNA sequences were

determined using primers described in Nézan et al. (2012). PCR cycling for partial LSU and ITS-5.8S-

ITS2 rDNA was carried out in an iCycler Thermal cycler (Bio-Rad, Hercules, CA, USA) using the

following conditions: ITS-5.8S-ITS2; pre-denaturation 94°C for 2 min, 37 cycles of 94°C for 30 s, 52°C

for 30 s, 72°C for 2 min, and a final extension at 72°C for 5 min, LSU; pre-denaturation 94°C for 3.5

min, 36 cycles of 94°C for 50 s, 45°C for 50 s, 72°C for 80 s, and a final extension at 72°C for 10 min.

The PCR products were purified using a QIAquick PCR Purification Kit (Qiagen, Hilden, Germany)

according to the manufacturer's instructions and sequenced (MCLAB, San Francisco, USA). Editing

and contig assembly of rDNA sequence fragments were carried out using Bioedit v7.2.5 (Hall, 1999).

DNA extraction and sequencing of 16S rRNA gene

DNA Extraction

For both the lab and field microbiomes, DNA was extracted from filters using a modified CTAB

protocol[94]. Briefly, 2 mL microtubes containing membrane filters and an extraction buffer were

subjected to three cycles of freezing in liquid nitrogen followed by thawing in a 65°C water bath.

Afterward, 8 µL of proteinase K (10 mg mL^{-1} in a TE buffer) was added, and the samples were

incubated at 37°C for 30 min. Next, 80 µL of 20% sodium dodecyl sulphate was added, and the

samples were incubated at 65°C for 2 h. The solution was then gently mixed with an equal volume of

chloroform-isoamyl alcohol (24:1) and centrifuged at 10,000 g for 5 min. The aqueous phase was

transferred to a fresh 2 mL tube containing 88.8 µL of 3 M sodium acetate (pH 5.1), followed by the

addition of 587 µL of isopropanol (≥99%). After centrifugation at 14,000 g for 20 min, the supernatant

was discarded, and 1 mL of cold 70% ethanol was added. The samples were then centrifuged at

14,000 g for 15 min, after which the pellets were air-dried at room temperature and dissolved in 100

µL of TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8).

16S rRNA Gene Sequencing

Lab microbiome

Sequencing libraries were prepared according to Illumina 16S Metagenomic Sequencing Library

protocols to amplify the V3 and V4 regions of the 16S rRNA gene. Genomic DNA (2 ng/10 ng input)

was amplified by PCR using a 5× reaction buffer, 1 mM dNTP mix, 500 nM each of the universal

forward/reverse primers, and Herculase II Fusion DNA Polymerase (Agilent Technologies, Santa

Clara, CA, USA). The conditions for the first PCR cycle were 95°C for 3 min for initial heat activation,

followed by 25 cycles of 30 s at 95°C, 30 s at 55°C, and 30 s at 72°C, with a final extension at 72°C

for 5 min.

The universal primer pair with Illumina adapter overhang sequences used for the initial PCR

amplifications were as follows: forward primer (5′-

TCGTCGGCAGCGTCAGATGTGTATAAGAGACAGCCTACGGGNGGCWGCAG-3′) and reverse

primer (5′-GTCTCGTGGGCTCGGAGATGTGTATAAGAGACAGGACTACHVGGGTATCTAATCC-3′).

The first PCR product was purified using AMPure XP beads (Agencourt Bioscience, Beverly, MA,

USA). Next, 10 µL of the first PCR product was used for a second round of PCR to construct the final

library, incorporating NexteraXT indexed primers. The conditions for the second PCR cycle were

identical to the first, except for 10 cycles. The PCR products were again purified with AMPure beads,

and the final purified product was quantified using quantitative PCR (KAPA Library Quantification Kit

for Illumina Sequencing Platforms) and validated using TapeStation D1000 ScreenTape (Agilent

Technologies, Waldbronn, Germany). Sequencing was performed on the Illumina HiSeq platform

(Illumina, San Diego, CA, USA).

Field microbiome

The extracted DNA samples were processed to generate amplicons for variable region 4 (V4) of the

nuclear small subunit (SSU) rRNA gene using the universal bacterial/archaeal primers 515F (5′-

GTGCCAGCMGCCGCGGTAA-3′) and 806R (5′-GGACTACHVGGGTWTCTAAT-3′), which contain a 6

bp error-correcting barcode unique to each sample. The amplified samples were sequenced on an

Illumina MiSeq platform with the 515F and 806R primers.

Operational taxon unit (microbial lineage) analysis

Lab microbiome

The raw data were demultiplexed by index sequences to generate FASTQ files for each sample.

Adapter sequences were trimmed using the fastp tool[95], and error correction was performed for

overlapping reads. The paired-end data for each sample were merged into single sequences using

FLASH (v1.2.11)[96]. Sequences shorter than 400 bp or longer than 500 bp were discarded. The

assembled sequences were processed using QIIME (v1.9)[97] and UCLUST[98] for microbial lineage

picking in a closed-reference approach. The NCBI 16S Microbial DB was used as the reference.

Finally, the microbial lineage abundance table was rarefied to the minimum number of sequences per

sample.

Field microbiome

The raw sequencing data were processed in R using the EasyAmplicon pipeline package and

incorporating various bioinformatics tools[99]. After removing barcodes and primers, paired-end

sequences were merged, and quality filtering was applied. Dereplication of full-length sequences was

performed, and sequences with a minimum unique size of 10 were retained. MLs were clustered at 97%

similarity using de novo chimera removal. Taxonomic classification of the representative microbial

lineages was performed against the SILVA 138.2 database[100] with a confidence threshold of 60%.

Finally, the microbial lineage abundance table was rarefied to the minimum number of sequences per

sample. The raw sequencing reads were deposited at the NCBI Short Read Archive under BioProject

ID PRJNA1180708.

Statistical analysis

All statistical analyses were conducted in RStudio (v2024.04.2+764) using R (v.4.3.2). Microbial

community comparisons were performed using the phyloseq (v.1.46.0)[101], Microbiotaprocess

(v.1.17.1)[102], and vegan (v.2.6.6.1)[103] packages based on ML abundance, taxonomy, and phylogenetic

tree data. Visualizations were generated using the ggplot2 package (v.3.5.1) [104] as needed. Prior to all

analyses, chloroplast- and mitochondria-related microbial lineages were filtered out, and microbial

lineage abundance was rarefied. For all analyses except co-occurrence networks, FAPROTAX, and

phylosymbiosis, MLs with an average relative abundance of less than 0.005% were excluded. All p-

values indicating the significance of microbiome differences between groups were adjusted using the

Bonferroni or Benjamini-Hochberg correction. All PERMANOVA permutation tests were conducted

with 9,999 permutations. To evaluate the statistical power of detecting group differences in beta

diversity via PERMANOVA, a post hoc power analysis was performed using G*Power (v.3.1.9.7)[105].

The analysis was based on a total sample size of 72, with 6 groups, each consisting of 2 strains, and

6 repeated measurements per strain. An effect size (f) of 0.35, an α error probability of 0.05, and a

correlation among repeated measures of 0.5 were applied. The power analysis yielded a noncentrality

parameter λ of 15.12 and a critical F value of 2.3538, resulting in an achieved power (1-β) of 0.836

(83.6%). These results indicate that the study had adequate statistical power to detect significant

between-group differences at a 5% significance level.

Network analysis was carried out using the ggClusterNet R package[106]. Microbial functional prediction

was performed using FAPROTAX based on taxonomic data using Python script[107]. To assess the

impact of stochastic processes on the microbiome, we applied the normalized stochastic ratio on an

NST_{jac} using the NST R package[108], with host taxonomy (dinoflagellate or diatom) as the grouping

variable and strain origin as the meta-group. A correlation analysis of Bray–Curtis dissimilarity

between phytoplankton strain microbiomes and the strain's environmental and spatiotemporal

information (Extended Data Table 1) was performed after log-transforming the data and normalizing

Euclidean distances between samples to a maximum value of 1. Data for microbiome dissimilarity

within the same strain across growth phases and replicates were excluded from the analysis.

Evolutionary distance and phylosymbiosis analysis

Evolutionary distance and host phylogeny

Large subunit ribosomal DNA sequences (total length 1031 bp) were used to calculate evolutionary

distances and construct phylogenetic trees among phytoplankton hosts (Extended Data Table 1). For

dinoflagellates, LSU sequences from six strains were analysed in this study, with one representative

sequence per species chosen to maximize the length of the comparison region. For diatoms, LSU

sequences were downloaded from the NCBI, except for *Coscinodiscus granii*, for which no LSU

sequences were available; those of the closely related *Coscinodiscus wailesii* were used instead. The

evolutionary history was inferred using the maximum likelihood method and the Tamura-Nei model[109].

A bootstrap consensus tree based on 500 replicates[110] was generated to represent the evolutionary

history of the analysed taxa. A discrete gamma distribution was used to model evolutionary rate

variation among sites (5 categories (+G, parameter = 0.6128)). All evolutionary analyses, including

model testing, were performed in MEGA11 [111].

Microbiota dendrograms

To obtain representative microbiota profiles for each species, the microbial lineage abundance data

for each strain and growth phase of the six microalgal hosts were collapsed into a single profile for

each species. Beta diversity distances between the six species' microbiomes were calculated using

the Bray–Curtis method. Distance matrices were then clustered using the UPGMA method to

generate dendrograms reflecting interspecific microbiome relatedness. To assess the impact of

different beta diversity measures on congruence with host phylogeny, dendrograms were generated

using weighted UniFrac and unweighted UniFrac distance methods.

Robinson–Foulds and matching cluster congruency analyses

Congruence between host phylogeny and microbiota dendrograms was quantified using normalized

Robinson–Foulds and normalized matching cluster methods. The analyses were performed using a

Python script, with 100,000 random trees used as the comparison baseline[9].

References

1. Douglas AE. Multiorganismal insects: diversity and function of resident microorganisms. *Annual review of entomology* **60**, 17-34 (2015).

2. Douglas AE. *Fundamentals of microbiome science: how microbes shape animal biology*. Princeton University Press (2018).

3. Flórez LV, Biedermann PH, Engl T, Kaltenpoth M. Defensive symbioses of animals with prokaryotic and eukaryotic microorganisms. *Natural Product Reports* **32**, 904-936 (2015).

4. Kissoyan KA*, et al.* Natural C. elegans microbiota protects against infection via production of a cyclic lipopeptide of the viscosin group. *Curr Biol* **29**, 1030-1037. e1035 (2019).

5. Obeng N, Bansept F, Sieber M, Traulsen A, Schulenburg H. Evolution of Microbiota–Host Associations: The Microbe's Perspective. *Trends in Microbiology* **29**, 779-787 (2021).

6. Ochman H*, et al.* Evolutionary relationships of wild hominids recapitulated by gut microbial communities. *PLoS biology* **8**, e1000546 (2010).

7. Phillips CD*, et al.* Microbiome analysis among bats describes influences of host phylogeny, life history, physiology and geography. *Molecular ecology* **21**, 2617-2627 (2012).

8. Moeller AH*, et al.* Rapid changes in the gut microbiome during human evolution. *Proc Natl Acad Sci USA* **111**, 16431-16435 (2014).

9. Brooks AW, Kohl KD, Brucker RM, van Opstal EJ, Bordenstein SR. Phylosymbiosis: relationships and functional effects of microbial communities across host evolutionary history. *PLoS biology* **14**, e2000225 (2016).

10. Pratte ZA, Besson M, Hollman RD, Stewart FJ. The gills of reef fish support a distinct microbiome influenced by host-specific factors. *Applied and environmental microbiology* **84**, e00063-00018 (2018).

11. Sylvain F-É*, et al.* Fish skin and gut microbiomes show contrasting signatures of host species and habitat. *Applied and environmental microbiology* **86**, e00789-00720 (2020).

12. Easson CG, Thacker RW. Phylogenetic signal in the community structure of host-specific microbiomes of tropical marine sponges. *Frontiers in microbiology* **5**, 532 (2014).

13. Sanders JG, Powell S, Kronauer DJ, Vasconcelos HL, Frederickson ME, Pierce NE. Stability and phylogenetic correlation in gut microbiota: lessons from ants and apes. *Molecular Ecology* **23**, 1268-1283 (2014).

14. Ofek M, Voronov-Goldman M, Hadar Y, Minz D. Host signature effect on plant root-associated microbiomes revealed through analyses of resident vs. active communities. *Environmental microbiology* **16**, 2157-2167 (2014).

635 15. Jones P, Garcia BJ, Furches A, Tuskan GA, Jacobson D. Plant host-associated mechanisms
636   for microbial selection. *Frontiers in Plant Science* **10**, 862 (2019).

637
638 16. Dubilier N, Bergin C, Lott C. Symbiotic diversity in marine animals: the art of harnessing
639   chemosynthesis. *Nature Reviews Microbiology* **6**, 725-740 (2008).

640
641 17. Venn AA, Loram JE, Douglas AE. Photosynthetic symbioses in animals. *Journal of
642   experimental botany* **59**, 1069-1080 (2008).

643
644 18. Visconti A*, et al.* Interplay between the human gut microbiome and host metabolism. *Nature
645   communications* **10**, 1-10 (2019).

646
647 19. Chandler JA, Morgan Lang J, Bhatnagar S, Eisen JA, Kopp A. Bacterial communities of
648   diverse Drosophila species: ecological context of a host–microbe model system. *PLoS
649   genetics* **7**, e1002272 (2011).

650
651 20. Boscaro V*, et al.* Microbiomes of microscopic marine invertebrates do not reveal signatures of
652   phylosymbiosis. *Nature Microbiology* **7**, 810-819 (2022).

653
654 21. Falkowski PG*, et al.* The evolution of modern eukaryotic phytoplankton. *science* **305**, 354-360
655   (2004).

656
657 22. Field CB, Behrenfeld MJ, Randerson JT, Falkowski P. Primary production of the biosphere:
658   integrating terrestrial and oceanic components. *science* **281**, 237-240 (1998).

659
660 23. Landa M*, et al.* Sulfur metabolites that facilitate oceanic phytoplankton–bacteria carbon flux.
661   *The ISME journal* **13**, 2536-2550 (2019).

662
663 24. Bar-On YM, Milo R. The Biomass Composition of the Oceans: A Blueprint of Our Blue Planet.
664   *Cell* **179**, 1451-1454 (2019).

665
666 25. Buchan A, LeCleir GR, Gulvik CA, González JM. Master recyclers: features and functions of
667   bacteria associated with phytoplankton blooms. *Nature Reviews Microbiology* **12**, 686-698
668   (2014).

669
670 26. Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. Algae acquire vitamin B12
671   through a symbiotic relationship with bacteria. *Nature* **438**, 90-93 (2005).

672
673 27. Seymour JR, Amin SA, Raina J-B, Stocker R. Zooming in on the phycosphere: the ecological
674   interface for phytoplankton–bacteria relationships. *Nature microbiology* **2**, 1-12 (2017).

675
676 28. Fortin SG, Song B, Anderson IC, Reece KS. Blooms of the harmful algae Margalefidinium
677   polykrikoides and Alexandrium monilatum alter the York River Estuary microbiome. *Harmful
678   Algae* **114**, 102216 (2022).

679
680 29. Jackrel SL, Yang JW, Schmidt KC, Denef VJ. Host specificity of microbiome assembly and its
681   fitness effects in phytoplankton. *The ISME Journal* **15**, 774-788 (2021).

683 30. Teeling H*, et al.* Substrate-controlled succession of marine bacterioplankton populations
684   induced by a phytoplankton bloom. *Science* **336**, 608-611 (2012).

685
686 31. Kuhlisch C, Shemi A, Barak-Gavish N, Schatz D, Vardi A. Algal blooms in the ocean: hot
687   spots for chemically mediated microbial interactions. *Nature Reviews Microbiology* **22**, 138-
688   154 (2024).

689
690 32. Deng Y*, et al.* Bacteria modulate microalgal aging physiology through the induction of
691   extracellular vesicle production to remove harmful metabolites. *Nature Microbiology* **9**, 2356-
692   2368 (2024).

693
694 33. Sison-Mangus MP, Jiang S, Tran KN, Kudela RM. Host-specific adaptation governs the
695   interaction of the marine diatom, Pseudo-nitzschia and their microbiota. *The ISME journal* **8**,
696   63-76 (2014).

697
698 34. Behringer G, Ochsenkühn MA, Fei C, Fanning J, Koester JA, Amin SA. Bacterial
699   Communities of Diatoms Display Strong Conservation Across Strains and Time. *Frontiers in*
700   *Microbiology* **9**,   (2018).

701
702 35. Mönnich J, Tebben J, Bergemann J, Case R, Wohlrab S, Harder T. Niche-based assembly of
703   bacterial consortia on the diatom Thalassiosira rotula is stable and reproducible. *The ISME*
704   *Journal* **14**, 1614-1625 (2020).

705
706 36. Mars Brisbin M, Mitarai S, Saito MA, Alexander H. Microbiomes of bloom-forming Phaeocystis
707   algae are stable and consistently recruited, with both symbiotic and opportunistic modes. *The*
708   *ISME Journal* **16**, 2255-2264 (2022).

709
710 37. Husnik F, Tashyreva D, Boscaro V, George EE, Lukeš J, Keeling PJ. Bacterial and archaeal
711   symbioses with protists. *Curr Biol* **31**, R862-R877 (2021).

712
713 38. Ahern OM, Whittaker KA, Williams TC, Hunt DE, Rynearson TA. Host genotype structures the
714   microbiome of a globally dispersed marine phytoplankton. *Proc Natl Acad Sci USA* **118**,
715   e2105207118 (2021).

716
717 39. Cohen NR*, et al.* Dinoflagellates alter their carbon and nutrient metabolic strategies across
718   environmental gradients in the central Pacific Ocean. *Nature Microbiology* **6**, 173-186 (2021).

719
720 40. Zheng B*, et al.* Dinoflagellate vertical migration fuels an intense red tide. *Proc Natl Acad Sci*
721   *USA* **120**, e2304590120 (2023).

722
723 41. Bell W, Mitchell R. Chemotactic and growth responses of marine bacteria to algal extracellular
724   products. *The Biological Bulletin* **143**, 265-277 (1972).

725
726 42. Deng Y, Wang K, Hu Z, Hu Q, Tang YZ. Identification and implications of a core bacterial
727   microbiome in 19 clonal cultures laboratory-reared for months to years of the cosmopolitan
728   dinoflagellate Karlodinium veneficum. *Frontiers in Microbiology* **13**,   (2022).

729
730 43. Stock W, Willems A, Mangelinckx S, Vyverman W, Sabbe K. Selection constrains lottery
731   assembly in the microbiomes of closely related diatom species. *ISME communications* **2**, 11
732   (2022).

733
734 44. Zou S, Yu X, Sun T, Wei L, Wu X. Unveiling bacterial communities linked with Alexandrium
735 catenella, their impact and influential mechanism on paralytic shellfish toxin production. *Algal*
736 *Research* **83**, 103698 (2024).

737
738 45. Bledsoe JW, Waldbieser GC, Swanson KS, Peterson BC, Small BC. Comparison of Channel
739 Catfish and Blue Catfish Gut Microbiota Assemblages Shows Minimal Effects of Host
740 Genetics on Microbial Structure and Inferred Function. *Frontiers in Microbiology* **9**,   (2018).

741
742 46. Grieneisen L*, et al.* Gut microbiome heritability is nearly universal but environmentally
743 contingent. *Science* **373**, 181-186 (2021).

744
745 47. Mallott EK, Amato KR. Host specificity of the gut microbiome. *Nature Reviews Microbiology*
746 **19**, 639-653 (2021).

747
748 48. Park BS, Kim J-H, Kim JH, Gobler CJ, Baek SH, Han M-S. Dynamics of bacterial community
749 structure during blooms of Cochlodinium polykrikoides (Gymnodiniales, Dinophyceae) in
750 Korean coastal waters. *Harmful Algae* **48**, 44-54 (2015).

751
752 49. Hattenrath-Lehmann TK, Gobler CJ. Identification of unique microbiomes associated with
753 harmful algal blooms caused by Alexandrium fundyense and Dinophysis acuminata. *Harmful*
754 *Algae* **68**, 17-30 (2017).

755
756 50. Zhang X*, et al.* Seasonal dynamics of bacterial community and co-occurrence with eukaryotic
757 phytoplankton in the Pearl River Estuary. *Marine Environmental Research* **192**, 106193
758 (2023).

759
760 51. Moran MA*, et al.* Ecological genomics of marine Roseobacters. *Applied and environmental*
761 *microbiology* **73**, 4559-4569 (2007).

762
763 52. Newton RJ*, et al.* Genome characteristics of a generalist marine bacterial lineage. *The ISME*
764 *journal* **4**, 784-798 (2010).

765
766 53. Francis B, Urich T, Mikolasch A, Teeling H, Amann R. North Sea spring bloom-associated
767 Gammaproteobacteria fill diverse heterotrophic niches. *Environmental Microbiome* **16**, 15
768 (2021).

769
770 54. Williams TJ*, et al.* The role of planktonic F lavobacteria in processing algal organic matter in
771 coastal E ast A ntarctica revealed using metagenomics and metaproteomics. *Environmental*
772 *microbiology* **15**, 1302-1317 (2013).

773
774 55. Reeves AR, Wang G, Salyers AA. Characterization of four outer membrane proteins that play
775 a role in utilization of starch by Bacteroides thetaiotaomicron. *Journal of bacteriology* **179**,
776 643-649 (1997).

777
778 56. González JM*, et al.* Genome analysis of the proteorhodopsin-containing marine bacterium
779 Polaribacter sp. MED152 (Flavobacteria). *Proc Natl Acad Sci USA* **105**, 8724-8729 (2008).

780
781 57. Green DH, Bowman JP, Smith EA, Gutierrez T, Bolch CJ. Marinobacter algicola sp. nov.,
782 isolated from laboratory cultures of paralytic shellfish toxin-producing dinoflagellates.

783   *International journal of systematic and evolutionary microbiology* **56**, 523-527 (2006).

58.   Lawson CA, Raina JB, Kahlke T, Seymour JR, Suggett DJ. Defining the core microbiome of the symbiotic dinoflagellate, Symbiodinium. *Environmental Microbiology Reports* **10**, 7-11 (2018).

59.   Andelman JB, Suess MJ. Polynuclear aromatic hydrocarbons in the water environment. *Bulletin of the World Health Organization* **43**, 479 (1970).

60.   Gunnison D, Alexander M. Basis for the resistance of several algae to microbial decomposition. *Applied Microbiology* **29**, 729-738 (1975).

61.   Marlowe I, Green JC, Neal A, Brassell S, Eglinton G, Course P. Long chain (n-C37–C39) alkenones in the Prymnesiophyceae. Distribution of alkenones and other lipids and their taxonomic significance. *British Phycological Journal* **19**, 203-216 (1984).

62.   McKay WA, Turner MF, Jones BMR, Halliwell CM. Emissions of hydrocarbons from marine phytoplankton—Some results from controlled laboratory experiments. *Atmospheric Environment* **30**, 2583-2593 (1996).

63.   Park BS, Erdner DL, Bacosa HP, Liu Z, Buskey EJ. Potential effects of bacterial communities on the formation of blooms of the harmful dinoflagellate Prorocentrum after the 2014 Texas City "Y" oil spill (USA). *Harmful Algae* **95**, 101802 (2020).

64.   Yang Q*, et al.* Marinobacter alexandrii sp. nov., a novel yellow-pigmented and algae growth-promoting bacterium isolated from marine phycosphere microbiota. *Antonie Van Leeuwenhoek* **114**, 709-718 (2021).

65.   Hernandez DJ, David AS, Menges ES, Searcy CA, Afkhami ME. Environmental stress destabilizes microbial networks. *The ISME Journal* **15**, 1722-1734 (2021).

66.   Ajani PA, Kahlke T, Siboni N, Carney R, Murray SA, Seymour JR. The microbiome of the cosmopolitan diatom Leptocylindrus reveals significant spatial and temporal variability. *Frontiers in microbiology* **9**, 2758 (2018).

67.   Medlin LK. Evolution of the diatoms: major steps in their evolution and a review of the supporting molecular and morphological evidence. *Phycologia* **55**, 79-103 (2016).

68.   Medlin LK, Desdevises Y. Phylogenetic reconstruction of diatoms using a seven-gene dataset, multiple outgroups, and morphological data for a total evidence approach. *Phycologia* **59**, 422-436 (2020).

69.   Lewin-Epstein O, Hadany L. Host–microbiome coevolution can promote cooperation in a rock–paper–scissors dynamics. *Proceedings of the Royal Society B* **287**, 20192754 (2020).

70.   Richardson LA. Evolving as a holobiont. *PLoS biology* **15**, e2002168 (2017).

71.   Franzenburg S*, et al.* Distinct antimicrobial peptide expression determines host species-specific bacterial associations. *Proc Natl Acad Sci USA* **110**, E3730-E3738 (2013).

832
833 72. McLoughlin K, Schluter J, Rakoff-Nahoum S, Smith AL, Foster KR. Host selection of
834 microbiota via differential adhesion. *Cell host & microbe* **19**, 550-559 (2016).

835
836 73. Schluter J, Foster KR. The evolution of mutualism in gut microbiota via host epithelial
837 selection. *PLoS biology* **10**, e1001424 (2012).

838
839 74. Shibl AA*, et al.* Diatom modulation of select bacteria through use of two unique secondary
840 metabolites. *Proc Natl Acad Sci USA* **117**, 27445-27455 (2020).

841
842 75. Fu H, Uchimiya M, Gore J, Moran MA. Ecological drivers of bacterial community assembly in
843 synthetic phycospheres. *Proc Natl Acad Sci USA* **117**, 3656-3662 (2020).

844
845 76. Landa M, Burns AS, Roth SJ, Moran MA. Bacterial transcriptome remodeling during
846 sequential co-culture with a marine dinoflagellate and diatom. *The ISME Journal* **11**, 2677-
847 2690 (2017).

848
849 77. Bolch CJ, Subramanian TA, Green DH. The toxic dinoflagellate Gymnodinium catenatum
850 (Dinophyceae) requires marine bacteria for growth 1. *J Phycol* **47**, 1009-1022 (2011).

851
852 78. Zhou J, Ning D. Stochastic Community Assembly: Does It Matter in Microbial Ecology?
853 *Microbiology and Molecular Biology Reviews* **81**, 10.1128/mmbr.00002-00017 (2017).

854
855 79. Hubbell SP. *The unified neutral theory of biodiversity and biogeography (MPB-32).* Princeton
856 University Press (2011).

857
858 80. Stegen JC, Lin X, Konopka AE, Fredrickson JK. Stochastic and deterministic assembly
859 processes in subsurface microbial communities. *The ISME journal* **6**, 1653-1664 (2012).

860
861 81. Stegen JC*, et al.* Quantifying community assembly processes and identifying features that
862 impose them. *The ISME journal* **7**, 2069-2079 (2013).

863
864 82. Fargione J, Brown CS, Tilman D. Community assembly and invasion: an experimental test of
865 neutral versus niche processes. *Proc Natl Acad Sci USA* **100**, 8916-8920 (2003).

866
867 83. Chave J. Neutral theory and community ecology. *Ecology letters* **7**, 241-253 (2004).

868
869 84. Chase JM. Stochastic community assembly causes higher biodiversity in more productive
870 environments. *Science* **328**, 1388-1391 (2010).

871
872 85. Dini-Andreote F, Stegen JC, Van Elsas JD, Salles JF. Disentangling mechanisms that mediate
873 the balance between stochastic and deterministic processes in microbial succession. *Proc
874 Natl Acad Sci USA* **112**, E1326-E1332 (2015).

875
876 86. Kohl KD. Ecological and evolutionary mechanisms underlying patterns of phylosymbiosis in
877 host-associated microbial communities. *Philosophical Transactions of the Royal Society B:
878 Biological Sciences* **375**, 20190251 (2020).

880   87.   Lim SJ, Bordenstein SR. An introduction to phylosymbiosis. *Proceedings of the Royal Society*
881   *B* **287**, 20192900 (2020).

882
883   88.   O'Brien PA*, et al.* Diverse coral reef invertebrates exhibit patterns of phylosymbiosis. *The*
884   *ISME Journal* **14**, 2211-2222 (2020).

885
886   89.   Guillard RR. Culture of phytoplankton for feeding marine invertebrates. In: *Culture of marine*
887   *invertebrate animals* (ed Smith WL, Chanley, M.H.). Springer US (1975).

888
889   90.   Park BS, Kim S, Kim J-H, Ho Kim J, Han M-S. Dynamics of Amoebophrya parasites during
890   recurrent blooms of the ichthyotoxic dinoflagellate Cochlodinium polykrikoides in Korean
891   coastal waters. *Harmful Algae* **84**, 119-126 (2019).

892
893   91.   Kim H-J*, et al.* Co-occurrence between key HAB species and particle-attached bacteria and
894   substrate specificity of attached bacteria in the coastal ecosystem. *Harmful Algae* **138**,
895   102700 (2024).

896
897   92.   Xu S, He C, Song S, Li C. Spatiotemporal dynamics of marine microbial communities
898   following a Phaeocystis bloom: biogeography and co‐occurrence patterns. *Environmental*
899   *Microbiology Reports* **13**, 294-308 (2021).

900
901   93.   Xu S, He C, Kang Z, Song S, Li C. The ecological responses of bacterioplankton during a
902   Phaeocystis globosa bloom in Beibu Gulf, China highlighted by integrated metagenomics and
903   metatranscriptomics. *Mar Biol* **169**, 36 (2022).

904
905   94.   Park BS, Baek SH, Ki J-S, Cattolico RA, Han M-S. Assessment of EvaGreen-based
906   quantitative real-time PCR assay for enumeration of the microalgae *Heterosigma* and
907   *Chattonella* (Raphidophyceae). *J Appl Phycol* **24**, 1555-1567 (2012).

908
909   95.   Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.
910   Bioinformatics 34: i884–i890.) (2018).

911
912   96.   Magoč T, Salzberg SL. FLASH: fast length adjustment of short reads to improve genome
913   assemblies. *Bioinformatics* **27**, 2957-2963 (2011).

914
915   97.   Caporaso JG*, et al.* Ultra-high-throughput microbial community analysis on the Illumina HiSeq
916   and MiSeq platforms. *The ISME journal* **6**, 1621-1624 (2012).

917
918   98.   Edgar RC. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**,
919   2460-2461 (2010).

920
921   99.   Liu YX*, et al.* EasyAmplicon: An easy‐to‐use, open‐source, reproducible, and
922   community‐based pipeline for amplicon data analysis in microbiome research. *IMeta* **2**, e83
923   (2023).

924
925   100.   Quast C*, et al.* The SILVA ribosomal RNA gene database project: improved data processing
926   and web-based tools. *Nucleic Acids Res* **41**, D590-D596 (2012).

927
928   101.   McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and

929   graphics of microbiome census data. *PloS one* **8**, e61217 (2013).

930
931   102.   Xu S*, et al.* MicrobiotaProcess: A comprehensive R package for deep mining microbiome. *The*
932   *Innovation* **4**,   (2023).

933
934   103.   Oksanen J*, et al.* Package 'vegan'. *Community ecology package, version* **2**, 1-295 (2013).

935
936   104.   Wickham H. ggplot2. *Wiley interdisciplinary reviews: computational statistics* **3**, 180-185
937   (2011).

938
939   105.   Faul F, Erdfelder E, Lang A-G, Buchner A. G* Power 3: A flexible statistical power analysis
940   program for the social, behavioral, and biomedical sciences. *Behavior research methods* **39**,
941   175-191 (2007).

942
943   106.   Wen T*, et al.* ggClusterNet: An R package for microbiome network analysis and
944   modularity-based multiple network layouts. *Imeta* **1**, e32 (2022).

945
946   107.   Louca S, Parfrey LW, Doebeli M. Decoupling function and taxonomy in the global ocean
947   microbiome. *Science* **353**, 1272-1277 (2016).

948
949   108.   Ning D, Deng Y, Tiedje JM, Zhou J. A general framework for quantitatively assessing
950   ecological stochasticity. *Proc Natl Acad Sci USA* **116**, 16892-16898 (2019).

951
952   109.   Tamura K, Nei M. Estimation of the number of nucleotide substitutions in the control region of
953   mitochondrial DNA in humans and chimpanzees. *Mol Biol Evol* **10**, 512-526 (1993).

954
955   110.   Felsenstein J. Confidence limits on phylogenies: an approach using the bootstrap. *evolution*
956   **39**, 783-791 (1985).

957
958   111.   Tamura K, Stecher G, Kumar S. MEGA11: molecular evolutionary genetics analysis version
959   11. *Mol Biol Evol* **38**, 3022-3027 (2021).

Acknowledgements

964 This research was supported by the National Research Foundation of Korea (NRF) grant funded by
965 the Korea Government (MSIT) (No. RS-2023-00209356 and No. 2022R1C1C1003582), and the
966 Korea Institute of Marine Science & Technology (KIMST) funded by the Ministry of Oceans and
967 Fisheries (RS-2023-00256330, Development of risk managing technology tackling ocean and
968 fisheries crisis around the Korean Peninsula by Kuroshio Current).

Author contributions

B.S.P. and J.-H.K. conceptualized the project. J.-H.K., J.H.K., Z.L., and Y.K.L. performed the wet
laboratory experiments. H.H.S. collected phytoplankton strains. B.S.P., J.-H.K., and J.H.K. collected
field phytoplankton bloom samples. J.-H.K. and P.W. conducted the metagenomic analyses. J.-H.K.,
R.Y.P., and S.H.J. curated the metadata of public samples. J.-H.K., J.H.K., and Z.L. designed the
statistical analyses. B.S.P., H.H.S., S.H.B., and M.-S.H. acquired funding. J.-H.K., J.H.K., and Z.L.
wrote the manuscript. B.S.P. supervised the work. All authors read and approved the manuscript.

Competing interests

979 The authors declare no competing interests.

980   # Tables

981   **Table 1.** PERMANOVA results comparing microbial community structure across groups and distance metrics.

Group	Number of MLs	Distance metrics	Pseudo F	R^2	P-value
Total MLs	242	Bray-Curtis	50.884	0.794	p < 0.0001
Total MLs	242	Weighted UniFrac	49.862	0.791	p < 0.0001
Total MLs	242	Unifrac	22.225	0.627	p < 0.0001
MCGs	77	Bray-Curtis	61.228	0.823	p < 0.0001
MCGs	77	Weighted UniFrac	93.756	0.877	p < 0.0001
MCGs	77	UniFrac	21.098	0.615	p < 0.0001
non-MCGs	165	Bray-Curtis	18.053	0.578	p < 0.0001
non-MCGs	165	Weighted UniFrac	14.088	0.516	p < 0.0001
non-MCGs	165	UniFrac	28.860	0.686	p < 0.0001

984   **Table 2**. The normalised Robinson–Foulds (nRF) and the normalized Matching Cluster metric analysis across Bray–Curtis, weighted and unweighted

985   UniFrac beta-diversity metrics.

Metric	Bray-Curtis	Weighted UniFrac	Unweighted UniFrac
nRF	RF = 0.00, p = 0.0031	RF = 0.50, p = 0.0735	RF = 0.00, p = 0.0031
nMC	MC = 0.00, p = 0.0031	MC = 0.83, p = 0.9637	MC = 0.00, p = 0.0031

 Figure legends

Fig. 1: Host-specificity of microbial assemblages in six phytoplankton species, confirmed in
both laboratory-cultured strains and field-collected phytoplankton blooms. a: Order-level
composition of microbiomes across strains and growth stages of the six phytoplankton hosts. **b:** PCoA
of microbiomes associated with the six phytoplankton species, based on Bray–Curtis dissimilarity.
Ellipses represent 95% confidence intervals for each host species. PERMANOVA results indicate
significant clustering by host species. **c:** PCoA highlighting microbiome differences among six
phytoplankton species by strain and growth phase. Dashed ellipses indicate growth-phase separation,
while solid ellipses represent strain differences. PERMANOVA significance is shown with *, $P < 0.05$;
, $P < 0.01$; ***, $P < 0.001$; ns, not significant. **d: PCoA of microbiomes in field-collected blooms of
four phytoplankton species (*Akashiwo sanguinea, Alexandrium monilatum*, *Margalefidinium*
polykrikoides, *Phaeocystis globosa*) sampled across Korea, China, and the US. Plots display
microbiomes along PCoA axes 1 and 2 (left) and axes 1 and 3 (right). Data for samples from China,
the US, and some Korean coastal regions were retrieved from the NCBI (Extended Data Table 3). To
minimize variability from different sequencing hypervariable regions, microbiome abundance data
were collapsed at the genus level.

1002

Fig. 2: Members of core microbial lineages (MCGs) play a key role in shaping host-specific
microbial assemblages across phytoplankton species. a: Number of core MLs and HS-MLs at the
bacterial order level for each host species. ND indicates not detected. The central Venn diagram
shows the overlap between Core MLs and HS-MLs. **b:** Microbiome taxa tree highlighting host-specific
taxa. The outermost points of the tree represent the relative abundance of each microbial lineage.
Nodes are colour-coded to indicate statistically enriched taxa for each host species, with node size
reflecting the adjusted p-value (Bonferroni correction). The external bar plot shows linear discriminant
analysis scores of differentially abundant taxa, with an effect size threshold of 3.2. Differential
abundance analysis was conducted using a two-step approach: a Kruskal–Wallis test to detect taxa
with overall group differences, followed by a Wilcoxon rank-sum test for pairwise comparisons. **c:**
Proportion of core MLs and HS-MLs among the top 200 MLs for each host species by number of MLs

and relative abundance. Core MLs shown here exclude HS-MLs within the MCGs. **d:** Venn diagrams

of Core MLs for each host species. Circle sizes are proportional to the numbers indicated.

Coscinodiscuss granii, which lacks core MLs, is not represented. **e:** Bacterial composition by lineage

group (ML types) within dinoflagellates and diatoms. Percentages were calculated based on the

number of microbial lineages. **f:** Bray–Curtis dissimilarity-based PCoA plots of MCGs (left) and non-

MCGs (right) datasets. PERMANOVA results indicate significant clustering by host species. **g:**

Distribution of Bray–Curtis dissimilarity values between and within species for the total microbiome,

MCGs, and non-MCGs datasets. Boxplot boundaries represent the first and third quartiles, with the

median indicated. Whiskers extend to the furthest data points within 1.5× the interquartile range.

Coloured points inside the boxes denote mean values (each calculated from n=630). **h:** Comparison

of between-species and within-species Bray–Curtis dissimilarity values for the total microbiome,

MCGs, and non-MCGs datasets. The endpoints of the dumbbells represent the mean values from

panel g. The values at the midpoints of the dumbbells represent the differences between between-

species and within-species dissimilarities. **i:** Bacterial functional group abundances across

phytoplankton species, as predicted by FAPROTAX analysis. Of the 19 functional groups that differed

significantly between dinoflagellates and diatoms, all were more abundant in dinoflagellates on

average. The left panel depicts the differences in functional group abundance, while the right panel

presents linear discriminant analysis scores alongside adjusted p-values (Benjamini-Hochberg

correction). Differential abundance was determined using a two-step approach: first, a Kruskal–Wallis

test to identify group-level differences, followed by pairwise comparisons using a Wilcoxon rank-sum

test. Boxplot boundaries represent the first and third quartiles, with the median indicated, and

whiskers extend to the furthest data points within 1.5× the interquartile range. **j:** Differential

abundance analysis of bacterial functional groups between ML groups (left: MCGs vs. non-MCGs;

right: HS-MLs vs. other) using FAPROTAX predictions. Bubble colours indicate significantly enriched

functional groups (adjusted p-value ≤ 0.05). To reduce visual complexity, nonsignificant results (p-

value > 0.05) and groups with small effect sizes (LDA score < 6.28) are not shown.

1040

1041

Fig. 3: Properties of co-occurrence networks and the significance of MCG nodes in

phytoplankton-associated microbiomes. a: Co-occurrence networks constructed from the top 200

microbial lineages by abundance for each of the six phytoplankton hosts. The upper row represents

microbiomes associated with dinoflagellates, while the lower row depicts those of diatoms. Nodes are

colour-coded to distinguish MCGs from non-MCGs lineages. *Coscinodiscuss granii* contains only non-

MCG nodes as no MCGs were identified. **b:** Comparison of three network properties (clustering

coefficient, connectance, and degree) between dinoflagellate- and diatom-associated microbiomes.

Data represent the mean values for each group (n = 3). Error bars indicate standard errors. Numbers

above the bars denote p-values. Statistical significance was calculated using a two-sided Student's t-

test. **c:** Distribution of three node properties (degree, closeness, and betweenness) for MCG and non-

MCG nodes within each host's co-occurrence network. Boxplot boundaries represent the first and

third quartiles, with the median indicated. Whiskers extend to the furthest data points within 1.5× the

interquartile range. To evaluate statistical significance and address differences in sample sizes

between groups, a permutation test with 1,000 resamples was performed. P-values were adjusted

using the Benjamini–Hochberg method.

1057

Fig. 4: Microbiome distance in phytoplankton is most strongly correlated with the evolutionary

distance between hosts, supporting the observation of phylosymbiotic patterns. a: Comparison

of the normalized stochasticity ratio (NST) of the Jaccard matrix between dinoflagellates and diatoms.

Values greater than 0.5 indicate that stochastic factors play a dominant role in microbiome assembly,

while values below 0.5 indicate a stronger influence of deterministic factors. **b:** Correlation of Bray–

Curtis dissimilarity between host microbiomes with phylogenetic, physicochemical, and

spatiotemporal distances. The phylogenetic distance between hosts was calculated using pairwise

distances based on LSU rDNA sequences (1,031 bp). All distance types, except for Bray–Curtis

dissimilarity and phylogenetic distance, were measured as Euclidean distances and normalized to a

maximum value of 1. The R^2 values for each distance type's correlation with Bray–Curtis dissimilarity

are displayed in radar plots, categorized by dinoflagellates, diatoms, and total data. All significant

correlations had P < 0.001. Lines show loess regression fits between Bray–Curtis dissimilarity and

each type of distance, with shaded areas represent the 95% confidence intervals. **c:** Phylosymbiotic

analysis comparing topological congruence between the rooted host phylogeny and microbiota

dendrogram. The host phylogenetic tree was constructed using LSU rDNA sequences (1,031 bp), and

the microbiota dendrogram was generated by UPGMA clustering based on Bray–Curtis dissimilarity.

Topological congruences were quantified using the normalized Robinson–Foulds (nRF) and

normalized matching cluster (nMC) metrics, with values ranging from 0 (complete congruence) to 1

(complete incongruence). *Note: LSU rDNA sequence data for* Coscinodiscuss granii *was unavailable*

in NCBI, so the sequence from its closely related species, Coscinodiscuss wailesii*, was used instead.*

1078

1079 **Extended Data**

1080

Extended Data Fig. 1. Significant differences in major bacterial taxa and higher diversity in

dinoflagellate microbiomes compared to diatom microbiomes. a: Microbiome taxa tree illustrating

differentially abundant taxa between dinoflagellate and diatom microbiomes. Tree nodes and

highlights are colour-coded to indicate the group (dinoflagellate or diatom) in which the corresponding

bacterial taxon is significantly more abundant. Node size represents the adjusted p-value (Bonferroni

correction). Differential abundance analysis involved a two-step approach: first, a Kruskal–Wallis test

was used to identify taxa with overall group differences, followed by a Wilcoxon rank-sum test for

pairwise comparisons. Taxa with an absolute linear discriminant analysis) score > 3.12 are shown. **b:**

Alpha diversity comparison between dinoflagellate (*n*=36) and diatom (*n*=36) microbiomes. Statistical

significance was assessed using a Wilcoxon rank-sum test. **c**: Alpha diversity comparison across six

phytoplankton species (*n*=12 per species). Statistical significance was also determined using a

Wilcoxon rank-sum test. Significance levels: *$P < 0.05$, ***$P < 0.001$; NS: not significant. Boxplot

boundaries indicate the first and third quartiles, with the median shown as a line. Whiskers extend to

the furthest data points within 1.5× the interquartile range.

1095

Extended Data Fig. 2. Geographic locations of host phytoplankton strain isolation and

phytoplankton bloom collection analysed in this study. a: Map showing study regions in Korea,

China, and the US. The red rectangle (b, d) represents the area shown in panels b and d, while c and

e indicate the regions depicted in panels c and e, respectively. **b:** Isolation sites of phytoplankton

strains along the Korean coastline. The numbers shown between the two shapes in the inset (n/n)

represent the spatial (km) and temporal (days) distances between the two strains isolated from each

location. Extended Data Table 1 provides more details. **c:** Collection sites of phytoplankton blooms in

the Beibu Gulf, China. Detailed information can be found in Extended Data Table 3. **d:** Collection sites

of phytoplankton blooms along the Korean coastline. **e:** Collection sites of phytoplankton blooms in

the York River, eastern US. In panels c to e, the numbers within the symbols indicate the number of

samples collected at each site.

1107

1108   **Extended Data Fig. 3. Heatmap and hierarchical cluster analysis of Bray–Curtis dissimilarity**

1109   **between phytoplankton-associated microbiomes.** The dendrogram was constructed using the

1110   UPGMA method. Bray–Curtis dissimilarity is represented by both colour gradation and circle size.

1111

1112   **Extended Data Fig. 4. Order-level composition of microbiomes from four microalgal blooms**

1113   **collected across various spatial and temporal scales.** Detailed sample information can be found in

1114   Extended Data Table 3.

1115

Extended Data Fig. 5. Distribution of bacterial functional group abundances across different

microalgal host species, as predicted by FAPROTAX analysis. A total of 19 functional groups

indicate significant differences between dinoflagellate and diatom hosts, with all groups more

abundant in dinoflagellates on average. Panels are arranged in descending order based on linear

discriminate analysis scores. Boxplot boundaries indicate the first and third quartiles, with the median

shown as a line. Whiskers extend to the furthest data points within 1.5× the interquartile range.

1122

Extended Data Fig. 6. Limited overlap in core bacterial taxa among strains of different species,

even when isolated from the same spatiotemporal conditions. Venn diagrams compare the

number of core bacteria (operational taxonomic unit level) shared among three strains of

dinoflagellates isolated from the same site on the same day (left) and those isolated from different

times and locations (right; details in Extended Data Table 1). Strain names are shown inside the Venn

diagram, and species names are labelled outside. The numbers in the intersections represent the

shared core bacteria. Arrows between the circles indicate the spatial (km) and temporal (days)

distances between the isolation sites of each strain.

1131

1132 **Extended Data Table 1.** Information on phytoplankton strains used in this study, including time, location, and environmental factors at the time of isolation.

Species	Group	Strain	Sample ID	Collection date	Latitude	Longitude	Temperature (℃)	Salinity	pH	DO (mg / L)	COD (mg / L)	T-N (µg / L)	T-P (µg / L)	Chl-a (µg / L)
Alexandrium catenella	Dinoflagellate	AC-1	ACA1	2019-03-05	35°3'5.4"N	128°47'7.8"E	13.24	34.23	8.24	9.37	0.8	129.8	15.9	0.53
Alexandrium catenella	Dinoflagellate	AC-2	ACA2	2017-08-29	34°45'9.2"N	128°22'46.7"E	27.56	32.19	8.16	7.93	1.5	250.9	28.7	6
Alexandrium pacificum	Dinoflagellate	AP-1	APA1	2018-08-03	34°59'39.1"N	128°40'28.8"E	28.56	32.52	7.76	4.6	1.87	152.2	26.5	4.28
Alexandrium pacificum	Dinoflagellate	AP-2	APA2	2017-08-29	34°45'9.2"N	128°22'46.7"E	27.56	32.19	8.16	7.93	1.5	250.9	28.7	6
Gymnodinium catenatum	Dinoflagellate	GC-1	GYM1	2017-08-29	34°45'9.2"N	128°22'46.7"E	27.56	32.19	8.16	7.93	1.5	250.9	28.7	6
Gymnodinium catenatum	Dinoflagellate	GC-2	GYM2	2016-07-12	34°42'43.21"N	127°40'0.45"E	26.01	32.01	7.93	3.48	1.92	222.1	63.6	5.28
Cylindrotheca closterium	Diatom	LIMS-PS-3013	CYL1	2016-06-13	37°46'25.69"N	126°19'2.27"E	17.36	31.73	8.29	8.99	1.43	271.8	22.2	6
Cylindrotheca closterium	Diatom	LIMS-PS-3150	CYL2	2017-05-26	34°59'36.71"N	128°40'27.73"E	17.59	33.95	8.17	8.5	1.41	97.4	13.9	1.81
Pseudo-nitzschia pungens	Diatom	LIMS-PS-3189	PSE1	2017-08-03	34°59'36.71"N	128°40'27.73"E	27.4	32.56	8.16	7.2	1.01	171.8	23.7	4.16
Pseudo-nitzschia pungens	Diatom	LIMS-PS-3248	PSE2	2017-05-18	34°41'34.50"N	127°50'27.60"E	15.97	34.2	8.24	8.62	1.96	125.4	16.8	1.8
Coscinodiscus granii	Diatom	LIMS-PS-3151	COS1	2017-06-06	34°59'36.71"N	128°40'27.73"E	18.58	33.9	8.2	8.75	1.37	86.4	13.5	2
Coscinodiscus granii	Diatom	LIMS-PS-3273	COS2	2018-06-25	35°36'33.37"N	129°36'25.31"E	25.77	32.55	7.86	8.73	1.23	109.7	9	0.41

1133

1134

1135 **Extended Data Table 2.** Comparison of beta diversity metrics for microbial assemblages across phytoplankton hosts.

Distance method	Pseudo F	R^2	P-value
Bray-Curtis	50.884	0.794	0.0001
Weighted UniFrac	49.862	0.791	0.0001
Jaccard	32.809	0.713	0.0001
Unifrac	22.225	0.627	0.0001

1136

1137 **Extended Data Table 3.** Information on phytoplankton bloom microbiome samples and DNA sequence sources used in this study.

SampleID	Original sample name	Bloom species	Collection date	Location	Latitude	Longitude	Instrument	Region	Primer_F	Primer_R	Study group
Marg_KOR1_1	P_CP_B_PH	*Margalefidinium polykrikoides*	2015-08-28	Pohang, South Korea	35°59'03.58"N	129°27'27.97"E	Illumina MiSeq	V4	515F (5-GTGCCAGCMGCCGCGGTAA-3)	806R (5-GGACTACHVGGGTWTCTAAT-3)	This study
Marg_KOR1_2	P_CP_LateB_PH	*Margalefidinium polykrikoides*	2015-09-03	Pohang, South Korea	35°59'03.58"N	129°27'27.97"E	Illumina MiSeq	V4	515F (5-GTGCCAGCMGCCGCGGTAA-3)	806R (5-GGACTACHVGGGTWTCTAAT-3)	This study
Marg_KOR1_3	P_CP_B_MG	*Margalefidinium polykrikoides*	2012-10-05	Mulgun, South Korea	34°47'38.19"N	128°03'07.69"E	Illumina MiSeq	V4	515F (5-GTGCCAGCMGCCGCGGTAA-3)	806R (5-GGACTACHVGGGTWTCTAAT-3)	This study
Marg_KOR1_4	P_CP_B_MJ	*Margalefidinium polykrikoides*	2015-09-08	Mijo, South Korea	34°42'35.66"N	128°02'50.53"E	Illumina MiSeq	V4	515F (5-GTGCCAGCMGCCGCGGTAA-3)	806R (5-GGACTACHVGGGTWTCTAAT-3)	This study
Marg_KOR1_5	P_CP_B_WD	*Margalefidinium polykrikoides*	2015-09-08	Wando, South Korea	34°23'34.07"N	126°38'23.22"E	Illumina MiSeq	V4	515F (5-GTGCCAGCMGCCGCGGTAA-3)	806R (5-GGACTACHVGGGTWTCTAAT-3)	This study
Marg_US_1	SF105	*Margalefidinium polykrikoides*	2017-08-09	York River Estuary, VA, USA	37°15'36.0"N	76°31'48.0"W	Illumina MiSeq	V4-V5	515F-Y (5-GTGYCAGCMGCCGCGGTAA)	926R (5-CCGYCAATTYMTTTRAGTTT)	[28]
Marg_US_2	SF104	*Margalefidinium polykrikoides*	2017-08-09	York River Estuary, VA, USA	37°14'24.0"N	76°30'36.0"W	Illumina MiSeq	V4-V5	515F-Y (5-GTGYCAGCMGCCGCGGTAA)	926R (5-CCGYCAATTYMTTTRAGTTT)	[28]
Marg_US_3	SF102	*Margalefidinium polykrikoides*	2017-08-09	York River Estuary, VA, USA	37°13'48.0"N	76°28'48.0"W	Illumina MiSeq	V4-V5	515F-Y (5-GTGYCAGCMGCCGCGGTAA)	926R (5-CCGYCAATTYMTTTRAGTTT)	[28]
Akas_KOR2_1	Bacteria 16S V3-V4 in Jangmok b	*Akashiwo sanguinea*	2016-11-14	Geoje, South Korea	34°59'38.6"N	128°40'27.9"E	Illumina MiSeq	V3-V4	341F (5- CCTACGGGNGGCWGCAG-3)	800R (5- TACCAGGGTATCTAATCC-3)	[91]
Akas_KOR2_2	Bacteria 16S V3-V4 in Jangmok b	*Akashiwo sanguinea*	2016-11-21	Geoje, South Korea	34°59'38.6"N	128°40'27.9"E	Illumina MiSeq	V3-V4	341F (5- CCTACGGGNGGCWGCAG-3)	800R (5- TACCAGGGTATCTAATCC-3)	[91]
Akas_KOR1_1	P_AS_B_JM	*Akashiwo sanguinea*	2011-10-28	Geoje, South Korea	34°59'37.47"N	128°40'28.46"E	Illumina MiSeq	V4	515F (5-GTGCCAGCMGCCGCGGTAA-3)	806R (5-GGACTACHVGGGTWTCTAAT-3)	This study
Alex_US_1	SF10	*Alexandrium monilatum*	2017-08-22	York River Estuary, VA, USA	37°15'00.0"N	76°29'24.0"W	Illumina MiSeq	V4-V5	515F-Y (5-GTGYCAGCMGCCGCGGTAA)	926R (5-CCGYCAATTYMTTTRAGTTT)	[28]
Alex_US_2	SF9	*Alexandrium monilatum*	2017-08-22	York River Estuary, VA, USA	37°13'48.0"N	76°27'00.0"W	Illumina MiSeq	V4-V5	515F-Y (5-GTGYCAGCMGCCGCGGTAA)	926R (5-CCGYCAATTYMTTTRAGTTT)	[28]
Phae_CHN1_h1	SFZZN3M1	*Phaeocystis globosa*	2017-03-23	Beibu Gulf, China	20°49'05.9"N	109°19'44.4"E	Illumina HiSeq	V3-V4	341F (5'-CCTAYGGGRBGCASCAG-3)	806R (5'-GGACTACHVGGGTWTCTAAT-3')	[92]
Phae_CHN1_h2	SFZZN3M2	*Phaeocystis globosa*	2017-03-23	Beibu Gulf, China	20°49'05.9"N	109°19'44.4"E	Illumina HiSeq	V3-V4	341F (5'-CCTAYGGGRBGCASCAG-3)	806R (5'-GGACTACHVGGGTWTCTAAT-3')	[92]
Phae_CHN1_h3	SFZZN3M3	*Phaeocystis globosa*	2017-03-23	Beibu Gulf, China	20°49'05.9"N	109°19'44.4"E	Illumina HiSeq	V3-V4	341F (5'-CCTAYGGGRBGCASCAG-3)	806R (5'-GGACTACHVGGGTWTCTAAT-3')	[92]
Phae_CHN2_h1	FLp1.1a.fastq	*Phaeocystis globosa*	2018-01-14	Beibu Gulf, China	21°42'35.0"N	108°35'12.0"E	Illumina HiSeq	V3-V4	341F (5'-CCTAYGGGRBGCASCAG-3)	806R (5'-GGACTACHVGGGTWTCTAAT-3')	[93]
Phae_CHN2_h2	FLp7.2a.fastq	*Phaeocystis globosa*	2018-01-14	Beibu Gulf, China	21°38'31.0"N	108°38'12.0"E	Illumina HiSeq	V3-V4	341F (5'-CCTAYGGGRBGCASCAG-3)	806R (5'-GGACTACHVGGGTWTCTAAT-3')	[93]

1138

a

A. catenella   *A. pacificum*

G. catenatum   *C. closteriun*

P. pungens   *C. granii*

Relative abundance (%)

lag   log   stn   lag   log   stn

Others, Thiotrichales, Saprospirales, Cytophagales, Alteromonadales, Sphingomonadales, Oceanospirillales, Kiloniellales, Vibrionales, Rhodobacterales, Pirellulales, Planctomycetales, Bacillales, Flavobacteriales

b

C. granii, *P. pungens*, *C. closterium*, *G. catenatum*, *A. pacificum*, *A. catenella*

PCo2 (15.54%)

A. pacificum, *A. catenella*, *G. catenatum*, *C. granii*, *C. closterium*, *P. pungens*

PERMANOVA :
$R^2 = 0.794$
$P < 0.0001$

PCo1 (22.28%)

c

A. catenella

lag, stn, log

PCo2 (33.76%)

PERMANOVA :
$R^2 = 0.82$ (Hg***)
$R^2 = 0.05$ (Hs^ns)

PCo1 (36.49%)

A. pacificum

PERMANOVA :
$R^2 = 0.76$ (Hg***)
$R^2 = 0.03$ (Hs^ns)

lag, stn, log

PCo2 (13.3%)

PCo1 (60.32%)

G. catenatum

stn, log, lag

PCo2 (16.61%)

PERMANOVA :
$R^2 = 0.84$ (Hg***)
$R^2 = 0.02$ (Hs^ns)

PCo1 (66.1%)

C. closterium

lag, log, stn

PCo2 (23.83%)

PERMANOVA :
$R^2 = 0.76$ (Hg**)
$R^2 = 0.02$ (Hs^ns)

PCo1 (61.59%)

P. pungens

strain 1, strain 2

PCo2 (30.39%)

PERMANOVA :
$R^2 = 0.21$ (Hg^ns)
$R^2 = 0.24$ (Hs*)

PCo1 (40.89%)

C. granii

strain 2, strain 1

PCo2 (1.55%)

PERMANOVA :
$R^2 = 0.02$ (Hg^ns)
$R^2 = 0.95$ (Hs**)

PCo1 (96.15%)

d

P. globosa, *M. polykrikoides*, *A. monilatum*, *A. sanguinea*

PCo2 (18.08%)

PERMANOVA :
$R^2 = 0.361$
$P < 0.0001$

A. monilatum, *P. globosa*, *M. polykrikoides*, *A. sanguinea*

PCo1 (21.28%)

P. globosa, *M. polykrikoides*, *A. monilatum*, *A. sanguinea*

PCo3 (12.23%)

PCo1 (21.28%)

a

b

Distance type
- Phylogenetic (*n*=2376)
- Environment (*n*=2376)
- Spacetime (*n*=2376)
- Space (*n*=2376)
- Time (*n*=2376)

Correlation with BCD
- Dinoflagellates
- Diatoms
- Total
- Significance (p < 0.001)

c

Host phylogeny

Microbiota Dendrogram – Bray-Curtis

Dinoflagellates

- *A. catenella*
- *A. pacifcum*
- *G. catenatum*

nRF = 0.00   nMC = 0.00
P = 0.0031   P = 0.0031

Diatoms

- *C. closterium*
- *P. pungens*
- *C. granii ***

a

Host group
- Dinoflagellate
- Diatom

Host group specific taxa
- a: f__Flammeovirgaceae
- b: f__Cryomorphaceae
- c: f__Flavobacteriaceae
- d: f__Bacillaceae
- e: f__Kiloniellaceae
- f: f__Rhodospirillaceae
- g: f__Erythrobacteraceae
- h: f__Methylophilaceae
- i: f__Alteromonadaceae
- j: f__Piscirickettsiaceae
- k: g__Reichenbachiella
- l: g__Crocinitomix
- m: g__Arenibacter
- n: g__Bacillus
- o: g__Thalassospira
- p: g__Erythrobacter
- q: g__Alteromonas
- r: g__Marinobacter
- s: g__Methylophaga

-log10 (P-value)
- 6
- 8
- 10

b Shannon / Chao1

c Shannon / Chao1

Predicted functional group

Same spatiotemporal

Different spatiotemporal