**Phyllosphere microbiome-based biocontrol solution against the filamentous**

**pathogens *Botrytis cinerea* and *Plasmopara viticola* in grapevine**

Nicolas-David Rappo[1], Sylvain Schnée[2], Clara Chevalley[2], Émilie Michellod[2], Katia Gindro[2], Laure

Weisskopf[1], Sébastien Bruisson[1*]

[1]Department of Biology, University of Fribourg, CH-1700 Fribourg, Switzerland.

[2]Mycology Group, Agroscope of Changins, 1260 Nyon, Switzerland

[*]Corresponding author: S. Bruisson; E-mail: sebastien.bruisson@unifr.ch

Chemin du musée 10, CH-1700 Fribourg, Switzerland

**Abstract**

Plants harbor different microbial communities in their different organs. This difference is due to the various conditions to which the different parts of the plant are subjected. As a result, the phyllosphere microbiome may be drastically different than the root microbiome. Since each microbiome is supposed to be well adapted to its environment, we hypothesize that bacterial communities from the leaves should be more effective in protecting the plants against the pathogen attacking the aerial part of the plant than the ones from the soil commonly used as biocontrol agents. In a previous study, we demonstrated that many phyllosphere bacteria isolated from the grapevine phyllosphere were indeed very effective against such plant pathogens under *in vitro* conditions, inhibiting their mycelial and spore development. In this project, we investigated the biocontrol ability of these phyllosphere bacteria *in planta* against two filamentous pathogens colonizing the aerial part of the plant: *Botrytis cinerea*, the ascomycete responsible for gray mold and *Plasmopara viticola* the oomycete responsible for downy mildew. Our results show that more than 40 strains were able to inhibit at least one pathogen in their spore physiology or infection progress. Among these strains, 27 strains were able to impact both pathogens and 20 were also capable of stimulating plant defenses and. Finally, when used in consortia, bacteria showed synergistic effects, resulting in a much stronger protection than that obtained by the application of the strains alone. These data suggest that phyllosphere bacteria could be a promising tool to help wine grower to fight against grapevine diseases by providing a strong long-lasting protection on their crops.

Keywords: *Vitis vinifera*, *Botrytis cinerea*, *Plasmopara viticola*, plant protection, biocontrol, plant

defense,

Funding: co-financed by Innosuisse; co-financed by the Swiss National Foundation (grant 310030

207917 to LW).

**Introduction**

Plant microbiota form a rich and complex network that is organized into different communities.

Each part of the plant provides a specific environment with conditions favorable to the

development of different species of bacteria (Compant et al., 2019). Thus, within the same

plant, bacterial communities found in roots are different from those found in leaves and flowers

(Zarraonaindia et al., 2015). Furthermore, because the composition of each community is

determined by the sum of all conditions they have to face, the bacteria found at the surface or

inside the plant of the same organ are also different (Dastogeer et al., 2020). Many recent

studies have shown that the plant microbiota contributes to its host's health by modulating its

growth, inhibiting pathogen development and promoting plant defenses (Brader et al., 2017; Rai

et al., 2023) and such bacteria are good candidates for developing biocontrol solution.

However, it appears that current biocontrol solutions only rely on organisms isolated from the

soil or roots, even when the aim is to control pathogens affecting the aerial part of the plant

(Syed Ab Rahman et al., 2018; Pascale et al., 2020, EU Pesticide database)

In this study, we therefore wanted to test the potential of bacteria isolated from the phyllosphere (the leaves and their surrounding environment) to protect plants against leaf-borne pathogens. We selected grapevine as a model plant to test such biocontrol solutions, for several reasons : i) it is a culture with a strong economic interest worldwide; ii) it is a perennial plant, and such plants are expected to harbor more adapted phyllosphere microbiota compared to annual plants, since perennial plants have longer-lasting interactions with their associated bacterial communities (Primieri et al., 2022); iii) most grapevine cultivars are highly susceptible to leaf-borne pathogens which makes this culture one the most fungicide consuming crops in the world, requiring several applications of copper, sulfur and other chemical compounds which can exceed the amount of one hundred kg per hectare each year (W. J. Li et al., 2011). Biocontrol is of particular interest for such cultures, that need alternative and novel effective solutions, to limit the toxicity linked to the excessive use of phytosanitary products.

Biocontrol phyllosphere bacteria have a wide range of modes of actions and can act either directly on the pathogens or indirectly, through the plant. Thus, they can have a direct inhibitory effect through the production of metabolites that can disturb the pathogen's growth and physiology, impair their virulence or they can also compete for space and nutrient, e.g. by trapping vital resources such as iron through the secretion of siderophores. They can also have an indirect effect on the pathogen by improving plant health through the production of plant growth promoting compounds or the induction of a local or systemic defense response (Legein et al., 2020). In grapevine, this defense response may involve the production of stilbenic

compounds, a group of phytoalexins of major importance for this crop due to their high toxicity

for fungal pathogens. This family includes a large number of inducible molecules that are

produced in several forms from monomers to tetramers. Their distribution and concentration

are cultivar and organ dependent and the resistant cultivars have been shown to synthetize high

amounts of the more active forms such as viniferins (Waffo-Teguo et al., 2008; Viret et al., 2018;

Guerrero et al., 2020).

In view of these multiple modes of action, combining different strains into a mixed microbe-

based biocontrol solution could provide a multi-layer protection by combining several

mechanisms that may lead to synergies and increased plant protection. Several studies already

highlighted that consortia of beneficial microbes can help to improve the effectiveness of the

protection or at least decrease its variability (Guetsky et al., 2001; Yu et al., 2022; Wang et al.,

2023).

In a previous study, we demonstrated that the use of the grapevine microbiota isolated from the

phyllosphere contained many powerful strains that could effectively inhibit the development of

leaf pathogens when co-cultivated in *in vitro* experiments (Bruisson et al., 2019). Thus, several

bacteria were able to strongly inhibit the mycelial growth of the fungus *B. cinerea* and the

oomycete *Phytophthora infestans* in *in vitro* conditions. They were also able to disrupt the

physiology of these pathogens by affecting their spores. We concluded that the use of such

bacteria was a promising strategy to increase the effectiveness of microbe-based biocontrol

solutions, but that more advance testing using plants was required.

In this work, we show that grapevine phyllosphere bacteria used as single strains or consortia

can provide a very effective protection *in planta* against two of the most detrimental grapevine

diseases caused by filamentous organisms, downy mildew caused by the oomycete *Plasmopara*

*viticola*, and gray mold caused by the ascomycete *Botrytis cinerea*. Our results suggest that this

protection is achieved by combining direct effects on the pathogens, affecting their spores and

their mycelium development, as well as indirect effects linked to their ability to increase the

expression of defense-related genes and the production of specialized metabolites involved in

plant defense.

**Materials and methods**

**Biological material and culture media**

Bacterial strains isolated from grapevine phyllosphere as described in (Vionnet et al., 2018) and

screened for their biocontrol capacity in (Bruisson et al., 2019) were grown on LB (Lysogeny

Broth) medium in the dark and at room temperature. LB medium was prepared by mixing 12.5

g/L of LB Broth Miller and 10 g/L of LB Broth Lennox (Fisher Bioreagents) in distilled water. For

solid medium, 15 g/L of Agar-Agar Kobe I (Roth) were added.

*Botrytis cinerea* strain 2905 was provided by the Mycology group at the Agroscope of Changins

and was grown in potato dextrose agar (PDA) (Roth). PDA was prepared by dissolving 39 g/L of

PDA powder (Sigma-Aldrich) in distilled water.

*Plasmopara viticola* sporangia were provided by the Agroscope of Changins. They were

harvested from infected leaves in vineyards each year to reflect the genotypes currently

infecting the grapevines at the time of the experiments.

Cuttings of *Vitis vinifera* cultivar Gamay, Gamaret and the interspecific hybrid Divico (Gamaret x

Bronner). After 2 months, plants were transferred into growth chambers with a 16h/8h

photoperiod, 21/20 °C day/night temperature and 40 % relative humidity (RH). Two Phillips

master TL-D super 80 36W/840 neon bulbs were used to provide light.

**Effect of bacterial strains on *B. cinerea* spore germination**

Spores of *B. cinerea* grown on PDA plate were harvested by watering the plate with 5 mL of

sterile water and filtration through glass wool to remove hyphae. The spore solution was then

adjusted to $5 \times 10^4$ spores/mL and 12 μL of this solution was mixed in an Eppendorf tube with 36

μL of bacterial suspension adjusted at $OD_{600}$=1 in V8 medium, clarified using a 0.2 μm Millex®-FG

filter (Merck). The mixture was transferred into a 24-well plate and the plates were incubated at

21 °C at light for 8 hours. Three technical replicates were performed for each modality and the

control consisted in spore solutions mixed with filtered V8 without bacteria. After 8 h pictures

were taken using an imaging plate reader (Cytation5, Biotek). Spores were counted according to

their development stage (not germinated, normal germination, delayed germination) and the

results were analyzed with a chi-square test using the package fifer on R with * = $p < 0.05$, ** = $p$

$< 0.01$ and *** = $p < 0.001$.

**Effect of bacterial strains on *P. viticola* zoospore motility**

Zoospores of *P. viticola* were released from sporangia harvested by suction from sporulating

lesions with a filter-containing pipette tip connected to a vacuum device. The sporangia were

resuspended in sterile water and their concentration was adjusted to 5 x $10^4$ sporangia/mL in a

falcon tube. The tube was placed in a shaker at 25 rpm at room temperature and in the dark for

2 hours for zoospore release. After 1 h, the zoospore solution concentration was adjusted to 5 x

$10^4$ zoospores/mL. The effect of bacteria on zoospore motility was assessed by mixing 500 µL of

a solution of bacteria diluted at $OD_{600nm}$ = 2 in sterile water with 500 µL of the zoospore solution

in a 1.5 mL Eppendorf. The Eppendorf was then incubated at room temperature in a shaker at 25

rpm for 30 minutes. After incubation, the solution was transferred into a counting chamber and

observed under a microscope. After counting, percentage of motile and non-motile zoospores

were computed and compared to the control (zoospore mixed with distilled water only).

Differences between bacterial treatments and the control were assessed by using an ANOVA

followed by a Dunnett's test with $* = p < 0.05$, $** = p < 0.01$, and $*** = p < 0.001$. Zoospores

were then used for leaf disc infection following the procedure described below.

**Electron microscope observation of zoospores**

Pellets of P. viticola zoospores, bacteria, or bacteria in contact with zoospores released using the

method described above were prepared according to Roland and Vian (1991). Briefly, the pellets

were centrifuged for 5 minutes at 2000 rpm and resuspended in a solution of 3%

glutaraldehyde–2% paraformaldehyde in 0.14 M PIPES buffer (pH 7) for prefixation. Samples

were then centrifuged for 20 minutes at 2000 rpm, the supernatant was removed, and the

samples were washed twice with 0.07 M PIPES buffer. They were embedded in 2% agarose and

postfixed with a solution of 1% OsO4. The samples were then dehydrated in a graded series of

ethanol solutions (30–50–70–95–100% v/v) and infiltrated in a solution of ethanol mixed with

propylene oxide (1:1) and pure propylene oxide, followed by a series of epoxy resin mixed with

propylene oxide (1:3, 1:1) for 2 hours each. Finally, they were infiltrated with epoxy resin for 17

hours. After polymerization for 48 hours at 60°C, a staining step was performed by floating on a

50°C solution of 1% methylene blue, 1% sodium borate, and 1% azure II for 10 seconds for semi-

thin sections. Thin sections were cut and stained with 2% uranyl acetate followed by lead citrate,

according to Reynolds (1963). The thin sections were observed with a Tecnai G2 Spirit BioTWIN

transmission electron microscope (TEM) equipped with a Gatan UltraScan High-resolution CCD

camera.

**Leaf discs infection assays**

Leaf discs were cut from 2-month-old grapevine leaves, taken from leaf stages 4 to 6. Leaves

were taken randomly from at least ten plants and were rinsed with tap water and surface

sterilized by immersion in hypochlorite calcium (40 g/L) for 7 minutes followed by rinsing in

sterile water prior to cutting. Leaves were then dried with sterile filter paper, used to cut leaf

discs using a 1.7 cm diameter punch and placed in sterile water. Ten leaf discs were taken

randomly and placed on top of a filter paper moistened with sterile water placed inside a 9 mm

Petri dish with the abaxial side facing upward and placed in a growth chamber with a 16h/8h

photoperiod, 21/20 °C day/night temperature and 40 % RH. For each modality, three Petri dishes

containing ten leaf discs each were used. After 24 h each plate was inoculated with 500 µL of

bacterial solution diluted in sterile water and adjusted to $OD_{600nm}$ = 1, using a sterile glass

atomizer and placed back in the growth chamber. Two days after the inoculation, the lead discs

were infected with *P. viticola* or *B. cinerea*. For *P. viticola*, the infection was performed using 500

μL of solution adjusted at $5 \times 10^4$ zoospores/mL for each plate. For *B. cinerea*, the infection was

performed by depositing a 20 μL droplet of a spore solution adjusted to $1 \times 10^5$ spores/mL in

PDB$^{1/4}$ on a wound made by a sterile syringe at the center of the lamina. The same procedure

was used with sterile water only for the negative control. For positive controls, plants were

treated with a 2 g/L solution of copper hydroxide (Kocide® Opti, Bayer) against *P. viticola* and a

0.375 % solution of brown algae (Alginure, Andermatt) against *B. cinerea*. After infection, Petri

dishes were sealed with Parafilm® and placed back in the growth chamber. For each modality,

one plate containing 10 leaf discs was used to collect material for gene expression analysis by

qPCR and stilbene production by HPLC. Five leaf discs were collected, dried and snap frozen in

liquid nitrogen after 6 h for qPCR analysis and after 3 days for HPLC analysis. Pictures of each

remaining plate were taken one week after infection to assess the development of symptoms.

**Whole plant infection with *P. viticola***

Half of the leaves from foliar stage 4 to 7 of two-month-old grapevine plants (4 plants per

treatment) were sprayed with a solution of 4 mL of bacteria adjusted to $OD_{600nm} = 1$ on their

adaxial face, while the other half was protected with plastic bags to prevent unwanted

inoculation. Once the inoculation finished, plastic bags were removed and uninoculated leaves

were sprayed with sterile water. Two days after the bacterial inoculation, all leaves from foliar

stage 4 to 7 were sprayed with 200 μL of a solution of *P. viticola* zoospores adjusted to $5 \times 10^4$

zoospores/mL. After a week, leaves were collected, the level of symptoms was quantified and

compared to the control (plants inoculated with water only) in search of local and systemic

protection from the bacteria.

**Symptom assessment**

For symptom measurement, pictures of leaves were analyzed in an automated and quantitative

manner, a macroinstruction was developed in the freeware ImageJ to measure the necrosis

development of *B. cinerea* and the formation of sporangiophores of *P. viticola* as described in

(Guyer et al., 2015). The level of inhibition was calculated as a percentage of the level of

infection observed in control.

**Gene expression analysis**

The ability of phyllosphere bacteria to promote defense gene expression on grapevine was

analyzed by qPCR in 8-week-old plants. Five leaf discs from infection assays described above

were sampled 6 h after infection, snap-frozen in liquid nitrogen and stored at −80 °C. RNA was

extracted with the NucleoSpin RNA Plant from Macherey-Nagel with a modified procedure

(Schellenbaum et al., 2008). Reverse transcription was performed using the SensiFAST cDNA

Synthesis Kit from Bioline. Quantitative PCR reactions were performed using the SensiFAST SYBR

Hi-ROX Kit from Bioline. Each reaction was carried out with 5 μL of cDNA (5 ng/μL), 7.5 μL of

SYBR Hi-ROX mix, 0.5 μL of each primer and 1.5 μL of sterile water. For amplification, an initial

denaturation step at 95 °C for 15 min was done, followed by 45 amplification cycles (95 °C for 15

s, 60 °C for 15 s, 72 °C for 30 s). Each reaction was run in duplicate and the experiment was

repeated twice. All results were analyzed using the double delta Cq method with uninfected

samples treated with water as reference and normalization was performed relative to the

expression of the two reference genes Actine and 60SRP (60S ribosomal subunit). Primers were

used to monitor the expression of the following defense-related genes, PR-1 (pathogenesis-

related protein 1), PR-4 (pathogenesis-related protein 4), STS (stilbene synthase), ROMT

(resveratrol O-methyltransferase), PAL (phenylalanine ammonia-lyase) and LOX (lipoxygenase)

(Supp. Table 1). A two-way ANOVA followed by Tukey-HSD post-hoc test was performed for

statistical analysis with $* = p < 0.05$, $** = p < 0.01$, and $*** = p < 0.001$.

**Stilbenic compound measurement**

Leaf disc fragments were weighed and placed into microcentrifuge tubes. Samples were

immediately snap-frozen and immersed in 300 µL of methanol. The tubes were then heated at

60°C for 15 minutes with agitation at 800 rpm, followed by rapid cooling in an ice bath for 5

minutes. After cooling, the samples were centrifuged to separate the supernatant. The

methanolic extracts were analyzed using Ultra-Performance Liquid Chromatography (UPLC) to

quantify the presence of various stilbenic compounds, including piceid, resveratrol, α-viniferin,

δ-viniferin, ε-viniferin, isohopeaphenol, and pterostilbene, following the method described by

Pezet et al. (2003). The analysis was performed using a Vanquish UHPLC system (Thermo

Scientific) with a mobile phase consisting of $H_2O$ with 0.1% formic acid (Phase A) and

acetonitrile with 0.1% formic acid (Phase B). The gradient elution program started with 5% B and

increased to 40% over 17 minutes, followed by an increase to 50% B between 17 and 20

minutes. From 20 to 25 minutes, the proportion of B increased to 100%, which was maintained

until 29 minutes. The flow rate was set at 0.3 mL/min. Separation was carried out using a

Hypersil Gold column (1.9 µm, 100 x 2.1 mm, Thermo Scientific) equipped with a 0.2 µm UHPLC

filter (Thermo Scientific), maintained at 30°C. The injection volume was 2 µL, and detection was

performed using a UV detector at 280 nm and 310 nm. This method ensures accurate and

reproducible quantification of stilbenic compounds in leaf tissues.

**Results**

**Motility disruption of *P. viticola* zoospores after contact with phyllosphere bacteria**

We tested all the strains from our collection of bacteria regarding their ability to disturb *P.*

*viticola* zoospores *in vitro*.  Confrontation assays between bacteria and *P. viticola* zoospores

showed that the contact with bacteria can strongly affect zoospore motility. A significant

reduction was measured with 28 strains out of the 53 tested. Visual assessment revealed that

zoospores either completely lost their motility, or kept motile as in the non-treated controls, but

we did not observe slower movement or lesser dispersal. However, the proportion of affected

zoospores was different depending on the bacterial strain used. Thus, a group of 16 strains

caused very strong reduction, affecting 90%-100% of the zoospores and reaching the level of

inhibition achieved with the copper used as a positive control, while a group of 12 strains caused

a significant but lower reduction (40%-90 %) (Table 1). A large predominance of bacteria from

the *Bacillus* genus were found among the active strains, since 19 of the 22 *Bacillus* strains tested

belonged to this group. However, the two most effective strains affecting both 100 % of the

zoospores (PIP10 and SOP5) were a *Pantoea agglomerans* and a *Microbacterium hominis* strain,

respectively. Interestingly the active strains were composed of a balanced proportion of

epiphytic (13) and endophytic (16) strains.

After being in contact with the bacteria, the zoospores were observed using a transmission

electron microscope (TEM). The zoospores that were in contact with non-active bacteria and

had not lost their motility displayed a standard oval shape comparable to the control zoospores.

However, the zoospores that had been in contact with the active strain PID6 (*Bacillus cereus*)

showed an irregular shape, suggesting that the whole cell had lost its integrity and had been

strongly affected by the presence of the bacteria (Fig. 1).

After addition of the bacteria, all the solutions containing the zoospores were used to infect leaf

discs cut from the susceptible cultivar Gamaret to monitor pathogen development. The

observations showed that in the case of 18 strains, the infections performed with the treated

zoospores resulted in a significant reduction in symptoms (from ca. 30% to 80 % of the control)

compared with the infection performed with non-treated zoospores (Table 1). From these 18

strains, only 13 were also significantly impairing motility, suggesting that the loss of motility was

not always correlated with symptom reduction and that other mechanisms are involved. Indeed,

five strains that had not impaired zoospore motility were nevertheless efficiently reducing the

symptoms caused by the zoospores with which they had been co-incubated.

Interestingly, four of these five strains belong to different genera than those highlighted

previously, namely *Rhodopseudomonas*, *Cupriavidus*, *Variovorax* and *Herbaspirillum.*

It seemed that when bacteria isolated from the cultivar Pinot noir were co-inoculated with

zoospores, this almost systematically resulted in significant symptom reduction, whereas the

two other cultivars harbored a lesser proportion of protective strains in this leaf disc

experiment.

**Phyllosphere bacteria inhibit the germination of *B. cinerea* spores**

The ability of phyllosphere bacteria to disrupt the germination of *B. cinerea* spores was assessed

by confrontation tests. After being in contact with the bacteria for 8 h, the spores displayed four

different morphologies: i) normal germination, with expected formation of a long germ tube, ii)

partially inhibited germination, with only the tip of the germ tube visible, iii) complete inhibition

of germination, with full inability for the spore to form a germ tube, iv) in a few rare cases,

spores could also form a swollen germ tube, revealing an abnormal germination in which the

germ tip swelled, stopping its elongation (Fig. 2 B-E).

Among the 53 bacteria that were tested, 34 strains were able to strongly disturb the

germination of more than 50% of *B. cinerea* spores, either by strongly delaying the formation of

the germ tube or by fully preventing spore germination. The six most effective strains SOD14,

PIP6, PIP4, PID6, PID13 and PID1 were even capable of fully inhibiting the germination, leaving

no spore able to germinate properly. These strains belonged mainly to the *Bacillus* or

*Paenibacillus* genera, with the exception of PIP4 (*Cupriavidus*) and of the most effective one,

PIP6 (*Herbaspirillum*).

**The inhibition of *B. cinerea* spore germination is higher with consortia than with bacteria**

**alone**

Consortia containing two to three active strains impairing spore germination were used to

assess the ability of the bacteria to generate synergies with higher protection abilities. The

solution containing the combination of bacteria contained the same final cell concentration than

the one used in the previous experiment, so any increase in effectiveness could not be simply

due to the increased bacterial density in the mixture. This combination resulted in an overall

increase of the strains' germination inhibition abilities. All consortia were able to affect all or

almost all spores, with the least effective one (CHP14, PIP6 and SOD2) inhibiting 98 % of the

spores and 20 out of the 23 tested consortia inhibiting 100 % of the spores (Fig. 3). The

difference between the consortia consisted in the proportion of spores that were fully vs.

partially inhibited in their germination. Interestingly, the presence of some strains such as PID6

(*Bacillus cereus*) and SOP8 (*Cupriavidus metallidurans*) seemed to increase the effect of all

consortia containing them and the use of these two strains alone prevented almost all spores

from germinating. On the contrary, the strains SOD2 (*Bacillus halotolerans*) and CHP14 (*Bacillus*

*subtilis*) slightly reduced the effectiveness of the consortia which contained them.

**Phyllosphere bacteria can inhibit symptom development *in planta* against *P. viticola* and *B.***

***cinerea***

The protective abilities of the 53 phyllosphere bacteria were assessed *in planta* on grapevine

leaf discs. Bacteria were sprayed onto grapevine leaf discs two days prior to infection, to let

them time to adapt and develop in their new environment. Against *P. viticola*, 14 strains were

able to significantly reduce the symptoms of the pathogen, to a level of from 24 to 59 % on

average (Table 2, Fig. 4), while all other strains did not result in any significant reduction in any

significant symptom reduction. These 14 strains included 10 strains already identified as active

in the first infection assay carried out with exposed zoospores described above (Table 1), while

four other strains that were not active in the previous experiment, SOD6 (*Bacillus cereus)*, CHP2

(*Cupriavidus metallidurans*), CHP20 (*Bacillus cereus*) and CHP29 (*Microbacterium proteolyticum*),

revealed their antioomycete abilities when pre-inoculated on the leaf discs (Table 2).

Higher levels of inhibition were achieved against *B. cinerea* with 15 strains allowing a significant

reduction of the necrosis going from 38 to 81 % on average (Table 3). In our conditions, seven

strains were able to achieve a better reduction in symptoms (more than 46 %) than the Botector

used as positive control: CHP29 (*Microbacterium proteolyticum*), SOP47 (*Variovorax paradoxus*),

CHP2 (*Cupriavidus metallidurans*), PIP6 (*Herbaspirillum huttiense*) PIP36 (*Bacillus cereus*), SOP8

(*Cupriavidus metallidurans*) and PID2 (*Bacillus simplex*) (Fig.5).

**Phyllosphere bacteria can stimulate plant defenses against *P. viticola* and *B. cinerea***

Infected leaf discs from the previous experiment were also collected in search for effect on plant

defenses. We monitored defense genes expression and stilbenic compound production. A set of

defense gene markers linked to the SAR (PAL), ISR (LOX), stilbene biosynthesis pathway (STS and

ROMT) and two pathogenesis related protein encoding genes (PR-1 and PR-4) were used for the

qPCR analysis. For each pathogen, we tested all strains that were able to significantly reduce symptom

development for their ability to induce plant defenses. Among the 14 strains that were effective

against *P. viticola*, 11 significantly upregulated the expression of at least three defense genes and

three strains had no positive impact on the defense gene expression (Fig. 6). These latter strains

(CHP20, CHP29 and SOD31), showed the lowest level of inhibition of symptoms on leaf discs and

mainly downregulated gene expression or had no significant effect on the monitored genes.

Among the strains that significantly upregulated the expression of several defense-gene, we

observed a tendency for upregulation of the genes from the SAR and the stilbene synthesis

pathways (PR-1, PAL, STS and ROMT). In contrast, the expression of the genes LOX and PR4 were

induced by fewer strains. Furthermore, it seems that when these two genes were upregulated,

the fold change was globally lower compared to the genes from the SAR or the stilbene pathway.

Interestingly, the strains PIP7, PIP17, SOD13, SOD2 and CHP2 were able to stimulate all or almost

all genes tested without downregulating any single gene. Finally, regarding the two best

performing strains in term of symptom reduction, PIP27 and CHP15, they had an opposite effect

on gene expression: PIP27 upregulated ROMT, PR-1 and PR-4 and slightly downregulated LOX,

PAL and STS, while CHP15 caused the opposite effect.

For *B. cinerea*, we tested the seven most effective strains that triggered an inhibition higher than

the one observed in the positive control. In contrast to *P. viticola* infection, the strains causing

the highest inhibition against *B. cinerea* mainly triggered a downregulation of the defense-

related genes (Fig. 6). Except for the strain PIP36 (*Bacillus cereus*), which was able to promote

the expression of almost all genes and SOP8, which only promoted the expression of PR-4 and

had moderate to no effect on the other genes, all the other strains had no significant or

negatively affected the expression of the tested set of defense-related genes.

Regarding the production of stilbenic compounds, differences were only observed in plants

infected with *P. viticola* (Fig. 7). Thus, in infected plants inoculated with protective bacteria, the

production of piceid and resveratrol (an inactive storage form and a precursor of more active

forms respectively) was stimulated by most of the bacteria. On the contrary, the production of

the active compounds, isohopeaphenol, α-viniferin and ε-viniferin was negatively impacted by

the bacterial inoculation with most strains. However, few strains were able to stimulate the

production of the different active forms. Thus, CHP15 (*Micrococcus yunnanensis*), SOD13

(*Paenibacillus xylanexedens*), SOD2 (*Bacillus* subtilis), SOD6 and PID6 (two *Bacillus cereus*) were

able to promote the synthesis of at least three of the active forms in infected leaves. Except for

CHP15 that triggered an inhibition of symptoms of 46 %, these strains were responsible for an

inhibition of 36, 39 and 32 % respectively corresponding to an moderate level among our active

strains.

**Strains combinations are more efficient than single strains for whole plant protection**

The ability of the protective strains to inhibit the symptoms of *P. viticola* on whole plants was

assessed using combinations containing three of the best performing strains. The strains used in

the consortia were selected based on their ability to affect spore physiology, to inhibit symptom

development, to stimulate defenses and to survive when exposed to at least the recommended

dose of antifungal compounds commonly used in the vineyard for crop protection such as

wettable sulfur and potassium hydrogenocarbonate (Supp. Table 2). Thus, random combinations

of CHP2 (*Cupriavidus metallidurans*), PID6 (*Bacillus cereus*), PIP6 (*Herbaspirillum huttiense*),

CHP14 (*Bacillus subtilis*), PID13 (*Bacillus endophyticus*), SOP8 (*Cupriavidus metallidurans*) and

SOD2 (*Bacillus halotolerans*) were assembled and sprayed on half the leaves of grapevine plants

while the other half was sprayed with water prior to the whole plant infection with *P. viticola*

zoospores. After a week, symptoms were measured on bacterial-inoculated and uninoculated

leaves in search of local and systemic protective effects of the bacteria. Result showed that the

strains CHP14, CHP2 and PID13 alone provided the lowest reduction of symptoms along with the

highest variability (38, 34 and 31 % respectively), the reduction triggered by CHP2 and PID13

being not even significant. However, all combinations were able to significantly reduce the level

of symptoms (Fig. 8). For all plants and for all treatments, the severity of symptoms was similar

on bacteria-inoculated leaves and in leaves that were spread with only water, suggesting that

the presence of bacteria can trigger a systemic response. However, the level of symptom

reduction was different between the different treatments. The three best combinations

(CHP2/PID6/PIP6; CHP14/PID13/PIP6 and CHP2/CHP14/PID13) led to a symptom reduction

ranging from 88.1 to 94.2 % on average, reaching an almost complete protection, however still

lower than the positive control (Table 4). The last combination (CHP2/CHP14/SOD2) provided a

reduction of 57.6 % which, although significant, was not very consistent (standard deviation of

39.2) and remained lower than the protection generated by the strains PID6 and SOP8 alone (82

and 68 % respectively).

**Discussion**

The results of this study show that phyllosphere bacteria are powerful agent to fight against

grapevine diseases that could be able to provide several layers of protection effective against

different kinds of pathogens.

**Spore physiology disruption against *P. viticola* and *B. cinerea***

The first layer of protection consists in affecting the reproduction of the pathogens trying to

colonize the plant. In past studies, bacteria such as streptomyces strains have been showed to

negatively affect the motility of *P. viticola* zoospores through the production of oligomycin and

pamemycin (Dame et al., 2016). In our study, among the 30 strains able to impair the motility of

*P. viticola* zoospores, almost 20 strains belonged to the genus *Bacillus*. This genus is well known

for containing many biocontrol species and several studies have highlighted the effect of *Bacillus*

strains on zoospores. *Bacillus* sp. 109GG020 is able to affect *Phytophthora capsici* zoospore

motility by producing gageopeptins, cyclic lipopeptides resulting in a loss of motility and an

abnormal helical instead of straight swimming (Tareq et al., 2015). *B. licheniformis* has also been

shown to negatively impair *P. capsici* zoospores by affecting their motility in the same way, as

well as causing zoospore lysis (Y. Li et al., 2020). Comparable effects have been described using

*Bacillus amyloliquefaciens* against *Phytophthora sojae* (D. Liu et al., 2019) and *Bacillus cereus*

against the pathogen *Pythium torulosum* (Shang et al., 1999). However, in our study, the most

active strains which, after co-incubation with the zoospores, induced a reduction in the

subsequent development of symptoms, were *Pantoea agglomerans*, *Microbacterium hominis*

and *Rhodopseudomonas boonkerdii*, species which have so far not been reported to harbor

strains affecting zoospores of oomycetes.  The origin of the bacteria does not seem to explain

their level of inhibition, since the proportion of endophytic and epiphytic strains was

comparable among active strains. One might have expected to find more active bacteria among

epiphytes, because due to their location they are in direct contact with the zoospores before

these enter the leaves. However, previous reports have already shown that other endophytic

microorganisms such as beneficial fungi like the ascomycete *Curvularia* sp. M12, were also able

to affect *P. capsici* zoospores (Mondol et al., 2017). In our work, after contact with bacteria, no

abnormal swimming was observed and the number of zoospores was comparable with the

control. The mechanism involved in the loss of motility was not investigated, but it seems that

the gageopeptins or cell lysis are not at play since abnormal motility was not observed and there

was no apparent sign of cell lysis as seen in Shang et al., 1999 and Tareq et al., 2015. However,

the TEM observations suggest that zoospores might be strongly affected by some bacteria since

they displayed a complete loss of their oval shape after contact with one of our tested bacteria.

It seems probable that the zoospores may have been killed although not lysed. Despite the strong

effects of several bacteria on zoospore motility, the zoospores were still able to infect grapevine

leaf discs, although with a significant reduction in symptoms compared to the control. Motility is

an essential function of the zoospore for plant infection in oomycetes (Chepsergon et al., 2020).

It is therefore expected that zoospores unable to swim can no longer infect plants. However,

several zoospores were still able to cause infection despite this loss of function. It seems that

they were still able to reach an infection site. We can hypothesize that the infection method

(spraying of a zoospore solution dissolved in water) could have led to the direct deposition of

living zoospores on a stomata or at least close enough for the zoospore to be transported inside

the plant with water as they entered the plant by absorption. In all cases it is important to note

that even with a complete loss of motility, spores were still able to infect the plant and that a

high loss in motility was not always correlated with the highest symptom reduction. Thus,

zoospores treated with strains like SOD2 and CHP14 despite an almost full inhibition of motility

were still able to successfully infect the plants after subsequent infection. In contrast, some

zoospores treated with strains such as PIP27 and PIP4 that were not able to inhibit the spore

motility showed a limited capacity to infect plants, suggesting that loss of motility alone is not

enough to fully hinder the virulence of *P. viticola* zoospores' and that the contact with bacteria

not only affects the motility but also other key functions in the zoospores.

The phyllosphere bacteria not only affected the spore motility of *P. viticola* but also the spore

germination of *B. cinerea*. The most effective strains regarding this effect belonged to the genera

*Herbaspirillum*, *Cupriavidus* and *Bacillus.* The biocontrol abilities for these two first genera have not been

well studied so far, but the effect of *Bacillus* strains on fungal pathogens including *B. cinerea* is well

known. *Bacillus subtilis* has been reported to affect *B. cinerea* hyphal morphology and to inhibit

spore germination through the release of antifungal volatiles (Chen et al., 2008). The use of a

broth from *B. cereus* strains also caused comparable effects (Li et al., 2012), as did the

application of *B. mycoides* (from the *B. cereus* group) as a biocontrol agent against *B. cinerea* in

strawberry (Guetsky et al., 2001a). Interestingly, one strain of *B. thuringiensis* was found among

the four best performing bacteria. This species is also a member of the *B. cereus* group and a

source of powerful biocontrol agents against insects mainly due to its production of Cry and Cyt

toxins (Bravo et al., 2011), but also because of its production of chitinase (Honda et al., 2017).

Since *B. cinerea* hyphal and spore cell are also composed of chitin (Chardonnet et al., 1999), this

chitinase could be responsible for this germination inhibition and it is possible that *B.*

*thuringiensis* could be a very effective biocontrol agent against fungal pathogens and not only

insects as suggested by a recent study (Martínez-Zavala et al., 2024). However, this strain was

not found in the best performing strain consortia because it was not synergizing well with other

bacteria unlike the others aforementioned *Bacillus*, *Herbaspirillum* and *Cupriavidus*. The

combination of bacteria from these genera increased tremendously the inhibitory effect on

sporulation, especially when different genera were mixed in the same consortia. Thus, consortia

of two bacteria from different genera were able to significantly increase the effectiveness of the

strains, some of which almost reached a complete inhibition of the germination. The fact that a

full of inhibition was reached with two combinations containing bacteria from three different

genera supports the hypothesis that the best synergies would be achieved when different

genera, supposedly using different mechanisms would be combined. Higher effectiveness of

biocontrol solutions using mixtures of bacteria instead of single strains has already been shown

for pathogen spore germination (Guetsky et al., 2001). In this case, the mechanisms involved are

also unknown, but the authors suggested that when consortia are used, the reliability of the

biocontrol is increased while the variability is decreased. This is what we observed in this study

as well, but in certain conditions, some associations were less effective. Thus, all combinations

containing the strain SOD2 (*B. halotolerans*) and CHP14 (*B. subtilis*) showed a lower effect

compared to the same combinations in which this strain was absent or replaced by another one.

The mechanism underlying this decreased efficacy is not known but we can rule out antagonism

between the strains regarding their growth, since compatibility had been tested for all strains

prior to designing the consortia. However, such effectiveness reduction when using

combinations of microorganisms in a plant biocontrol context has already been described in

banana treated with beneficial fungi (Sukhada et al., 2011)

**Preventive inoculation of phyllosphere bacteria prior to infection led to significant protection**

**against *P. viticola* and *B. cinerea***

Fewer bacterial strains caused a significant reduction of symptoms against *P. viticola* compared

with the previous test. Despite the two days between the bacterial inoculation and the

pathogen infection to give plants and bacteria time to adapt to each other, it seems that a direct

contact with the zoospores was more effective than a plant inoculation. However, the conditions of a

bacterial preinoculation are closer to natural conditions or conditions that would be used in the field

when applying the strains. If we compare the list of effective strains from the first assay, that were at the

same time able to affect motility and to reduce symptom development, with the strains effective in the

bacterial preinoculation experiment, only three strains, SOP5, PID6 and PIP17 (*Microbacterium hominis*,

*Bacillus cereus* and *Staphylococcus arlettae* respectively) proved  effective in both experimental set up,

suggesting that the weakening of spores does not play such an important role in natural or field

conditions and that other mechanisms are at play. Regarding the type of bacteria that showed activity in

this set up, we found once again the genera from the previous experiment *i.e.*: *Cupriavidus, Paenibacillus*

*Rhodopseudomonas* and a large representation of *Bacillus* species, but we also found three different

strains of *Microbacterium*. So far, this genus has not been often mentioned in the literature as effective

against *P. viticola,* but some strains have been shown to be effective at stimulating plant defense and at

inhibiting the growth of oomycetes and several fungal pathogens (Barnett et al., 2006; Freitas et al.,

2019; Ray et al., 2021; Suárez-Estrella et al., 2023).

The same observations have been made against *B. cinerea*, less bacteria caused a reduction in symptoms

compared to the test performed with spores, suggesting that for both oomycetes and fungi, activities on

spores, although helpful, are not essential for achieving effective biocontrol, it is also possible that the

ability of bacteria to disturb the pathogens' spores *in planta* is weaker than *in vitro*. Almost all genera

found previously were also found to be active against *B. cinerea* in these conditions, with the addition of

one *Variovorax* and one *Plantibacter.* These two genera are known for their plant growth promoting

abilities (Giannelli et al., 2024; Han et al., 2013; Mayer et al., 2019; Sun et al., 2018), but their

biocontrol abilities are poorly studied. *Variovorax* has shown moderate biocontrol activities in

tomatoes against *B. cinerea* and in wheat against *Fusarium* (Besset-Manzoni et al., 2019; Köhl et al.,

2020), but to our knowledge, no study has yet reported any biocontrol effects of *Plantibacter*

strains.

**Several phyllosphere bacteria contribute to enhance grapevine defenses**

Bacterial strains successfully inhibiting the development of symptoms when they were

inoculated *in planta* prior to infection were tested for their ability to stimulate the expression of

several defense genes belonging to different defense pathways. Most of the strains effective

against *P. viticola* were able to upregulate several defense genes, but several bacteria such as

CHP20, CHP29 and SOD31 show few to no significant effect regarding defense gene upregulation

and seemed to even slightly downregulate plant defenses. Most of the strains shared the

downregulation of the PR-4 gene a gene encoding a protein with chitinase and chitin-binding

activities (Enoki & Suzuki, 2016; M. Y. Li et al., 2020; G. T. Liu et al., 2021). The expression of this

protein has been shown to be upregulated in grapevine resistant cultivars against *P. viticola* or in

susceptible cultivar upon transgenic overexpression, but not in susceptible cultivars (Enoki & Suzuki,

2016; M. Y. Li et al., 2020; G. T. Liu et al., 2021). Other studies mentioned that the expression of PR-4 is

upregulated in association with the JA-ethylene pathway against necrotrophic fungi like *B. cinerea* while

SA pathway is recruited against biotrophic pathogens such as *P. viticola* (Trusov et al., 2009; Van Loon

et al., 2006). This could explain that in our case, the expression of this protein-encoding gene

was not promoted since we are using a susceptible cultivar and because the SA pathway,

antagonistic to the JA pathway (Vlot et al., 2021), seems to be mainly recruited after infection in

our conditions. Thus, for most bacteria stimulating plant defenses against *P. viticola*, PAL, STS,

ROMT and PR1 were among the genes that were most frequently upregulated. These data

suggest that our active bacteria use different pathways to protect the plant against the pathogen

and that several of them do not rely on plant defense promotion. Interestingly, it seems that the

most effective strains against *P. viticola*, PIP27 and CHP15, were not the best at stimulating plant

defenses, so we may assume that they have a direct and strong effect on the pathogen.

When testing induction of defense genes upon infection with *B. cinerea*, we observed that

except for PIP36 and SOP8, that triggered the upregulation of LOX and PR-4 (consistent with

Trusov et al., 2009 and Van Loon et al., 2006), the other strains negatively affected the

expression of the monitored defense related genes. If we compare this observation with the one

we made in the case of *P. viticola*, we observe that *Cupriavidus metallidurans* CHP2, stimulated

the expression of defense-related genes in presence of *P. viticola*, but mostly inhibited the

expression of defense-related genes in presence of *B. cinerea*. This suggests that the effect of

this strain is specific to the type of pathogen, whether biotrophic or necrotrophic. But the

reduction in plant defense does not appear to be detrimental to the plant, since despite the lack

of plant defense stimulation, the bacteria were able to successfully protect the plant against the

pathogen suggesting that their effect is more likely acting on the pathogen rather than on plant

defenses. However, it is still possible that a different set of defense related genes are involved

for the protection against *B. cinerea*. Thus, according to a recent study, the genetic of grapevine

resistance is still poorly known, but different defense gene markers such as Ethylene Responsive

Factors (ERF) could be involved in the defense against this pathogen (Zhu et al., 2019; Rahman et

al., 2024).

The inoculation of bacteria also affected the production of stilbenes in leaves. These compounds

are phytoalexins with different levels of antifungal properties depending on their structure.

Against *P. viticola*, several strains were able to stimulate the production of several of these

compounds. Most of the strains promoted the production of resveratrol, synthetized by the

stilbene synthase (STS), an active form of stilbene and precursor of several more active

compounds such as viniferins and pterostilbenes (Coutos-Thévenot et al., 2001). CHP15, SOD13,

CHP2, SOD6 and PID6 (*M. yunnanensis*, *B. xylanexedens*, *C. metallidurans* and two *B. cereus*, and

*Cupriavidus metallidurans*) were able to strongly increase the production of at least two

different forms of viniferins, one the most active stilbenic compounds against *P. viticola* (Alonso-

Villaverde et al., 2011; Pezet et al., 2003; Van Leeuwen et al., 2013). The correlation between the

increase of the production of resveratrol and viniferin and the higher expression of The STS

genes in several active strains suggest that they indeed positively affect this pathway enough to

influence the development of the pathogen. However, no significant amounts of pterostilbenes,

a compound with higher antifungal effects than viniferins have been found in any sample,

despite the increase in the expression of the ROMT gene encoding the enzyme responsible for

its synthesis from resveratrol. This compound is produced in small amounts by grapevine even

after being infected by downy mildew (Godard et al., 2009) and a previous study showed that

even with higher gene expression and upon infection by a pathogen, levels of pterostilbenes

may not change significantly and remain low (Kong et al., 2020; Rahman et al., 2024). It is

therefore possible that despite the upregulation of the ROMT gene, the level of pterostilbene

production was too low for detection in our conditions.

On the contrary, no significant increase of any stilbenic compounds were found against *B.*

*cinerea*. These compounds have yet been shown to be effective against *B. cinerea* dormant

conidia (Pezet & Pont, 1990; Pont & Pezet, 1990), but when not in its quiescent form, this

pathogen produces lytic enzymes able to degrade stilbenic compounds (Pezet et al., 1991; Goetz

et al., 1999). The production of such enzymes could explain why no significant changes in

stilbene production were observed in our samples after infection with *B cinerea*, both in the

presence or absence of phyllosphere bacteria.

**Combinations of phyllosphere bacteria are effective *in planta* against *P. viticola***

The combination of strains we used on whole plants were all able to significantly protect the

plant both locally and systematically with a comparable level of symptom inhibition (Table 4).

For each combination, the strains were selected based not only on their abilities to inhibit

pathogen growth or to stimulate plant defenses, but also on their abilities to grow well together

and to survive in the presence of several phytosanitary products used in the field. Thus, each

selected strain was able to develop properly in culture media in presence of potassium

hydrogenocarbonate and wettable sulfur. Interestingly, as observed in experiments targeting *B.*

*cinerea* spore germination, the combination of strains containing three different genera was the

most effective against *P. viticola* compared to the other combinations containing at least two

*Bacillus* strains. Thus, the combination CHP2/PID6/PIP6 (*Cupriavidus metallidurans*, *Bacillus*

*cereus* and *Herbaspirillum huttiense*) was the most effective one and was able to reach an

almost full inhibition of symptoms (94 %). This level of protection was only surpassed by the

positive control Alginure®, a product certified for field use against *P. viticola* in Switzerland

containing a combination of brown algae, potassium phosphonate and amino acids, acting as a

direct fungicide and as a plant defense enhancer. Our best combination of bacteria, despite its

lack of formulation, was therefore able to match the protection conferred by a mixture of

biocontrol and chemical agents. This result suggests that strains greatly differing in genetic

makeup and potentially using different mode of actions are more effective than combinations of

very effective but genetically close strains that are more likely to use similar strategies of

inhibition. Consistent with the results observed on *B. cinerea* spores, the presence of the strain

SOD2 (*Bacillus halotolerans*) had a negative impact on the performance of the consortia in

which it was used. Thus, this consortium was the least successful at *P. viticola* inhibition and

resulted in lesser inhibition than SOP8 and PID6 alone. This combination also showed the

highest variability compared to the other consortia, making it particularly unreliable.

The use of combinations of bacteria with other bacterial strains, microorganisms from different

kingdoms or even chemical compounds has been reported as successful against several plant

pathogens in the past. This solution has several advantages, it allows the addition of several

effects such as direct pathogen inhibition, accumulation of more diverse antipathogenic

compounds, plant defense stimulation and higher competition for space and resources (Niu et

al., 2020). In tomatoes, combination of a *Bacillus* strain with several bactericides and a mixture

of two streptomyces was found more effective than the use of each strain or compound applied

separately against tomato wilt (Wang et al., 2023; Yu et al., 2022). Similarly, strains of *Bacillus*

*mycoides* and *Pichia guilermondii* were more effective at inhibiting *B. cinerea* spore germination

and reducing necrotic lesion size in strawberry leaves when used in combination compared to

their use alone, showing that organisms from different kingdoms can synergize and also provide

more consistent protection (Guetsky et al., 2001). On the contrary, a synthetic microbial

community composed of *Pseudomonas*, *Bacillus*, *Trichoderma* and *Rhizophagus* was shown to

not synergize against the two fungal pathogens *Fusarium oxysporum* and *B. cinerea* in tomato

(Minchev et al., 2021). In this study, the level of protection conferred by the consortium was

always equivalent to the best performing organism alone, suggesting that strains assembled in a

consortium do not always display synergistic effects in terms of protection against diseases. Even

if the use of microorganism combinations as biocontrol agents against plant pathogens has

mainly been studied against root pathogens, literature data show that some strains can be more

effective than others in term of plant protection. Thus, for bacteria, the genera *Bacillus* and

*Pseudomonas* are the most commonly found (Niu et al., 2020). In our study, several different

*Bacillus* species were proven effective but no *Pseudomonas* were present. Instead, other genera

such as *Cupriavidus*, *Herbaspirillum*, *Microbacterium*, and other less commonly studied species

showed high potential. We can speculate that to achieve the best protection possible with

bacteria-based biocontrol solutions, the use of combination including such genera is key and

that strains showing only moderate effects when used alone could be of high importance to

stabilize or increase the effectiveness of a consortium.

**Conclusion**

This study reports that the use of phyllosphere bacteria as biocontrol agents could be an

efficient strategy to fight against two of the most detrimental filamentous pathogens for

grapevine cultivation, *P. viticola* and *B. cinerea*. The plant natural microbial communities are a

good source of potent biocontrol strains and the use of microorganisms already adapted to the

environment and the organ to protect. After optimization, their use can probably show higher

effectiveness compared to solutions based on the application of exogenous organisms not fit to

live in the aerial part of the plant and consequently less likely to reach their full potential against

plant diseases. These bacteria were even more effective when they were used in consortia

combining diverse bacterial genera. They were shown to disturb the pathogens' spore

physiology, the mycelium development and also to stimulate plant defenses. Thus, combinations

of different species that show high effectiveness in these three aspects were able to almost fully

inhibit *P. viticola* infection in grapevine susceptible cultivars. Whether such level of protection

could also be conferred under field conditions is the next question we will need to address. To

this end, it will be important to develop a formulation recipe, since in our experiment, bacteria

were used only dissolved in sterile water, without any UV or dessication protection. Our results

suggest that a solution based on consortia from grapevine phyllosphere isolates could be a

powerful tool to control diseases in open vineyards or at least under plant production conditions

in nurseries. We believe that such biocontrol solution, providing several layers of defense against

pathogen and based on bacteria already adapted to the organ to protect could bring a long-term

protection with no harmful side effects on the plant or the environment. This solution could help

the farmer to reduce their use of pesticide without compromising the health of their plant and

the safety of their products.

**Acknowledgement**

The authors are grateful to Alisson Gillon and Gaëlle Ruffieux for their help in the experimental

work with grapevine infection and bacterial inoculum preparation.

**References**

Alonso-Villaverde, V., Voinesco, F., Viret, O., Spring, J. L., & Gindro, K. (2011). The effectiveness of

stilbenes in resistant Vitaceae: Ultrastructural and biochemical events during Plasmopara viticola

infection process. *Plant Physiology and Biochemistry*, *49*(3), 265–274.

https://doi.org/10.1016/j.plaphy.2010.12.010

Barnett, S. J., Roget, D. K., & Ryder, M. H. (2006). Suppression of Rhizoctonia solani AG-8 induced disease

on wheat by the interaction between Pantoea, Exiguobacterium, and Microbacteria. *Australian*

*Journal of Soil Research*, *44*(4), 331–342. https://doi.org/10.1071/SR05113

Besset-Manzoni, Y., Joly, P., Brutel, A., Gerin, F., Soudière, O., Langin, T., & Prigent-Combaret, C. (2019).

Does in vitro selection of biocontrol agents guarantee success in planta? A study case of wheat

protection against Fusarium seedling blight by soil bacteria. *PLoS ONE*, *14*(12).

https://doi.org/10.1371/journal.pone.0225655

Brader, G., Compant, S., Vescio, K., Mitter, B., Trognitz, F., Ma, L.-J., & Sessitsch, A. (2017). Ecology and

Genomic Insights into Plant-Pathogenic and Plant-Nonpathogenic Endophytes. *Annual Review of*

*Phytopathology*, *55*(1), 61–83. https://doi.org/10.1146/annurev-phyto-080516-035641

Bravo, A., Likitvivatanavong, S., Gill, S. S., & Soberón, M. (2011). Bacillus thuringiensis: A story of a

successful bioinsecticide. In *Insect Biochemistry and Molecular Biology* (Vol. 41, Issue 7, pp. 423–

431). https://doi.org/10.1016/j.ibmb.2011.02.006

Bruisson, S., Zufferey, M., L'Haridon, F., Trutmann, E., Anand, A., Dutartre, A., De Vrieze, M., & Weisskopf,

L. (2019). Endophytes and Epiphytes From the Grapevine Leaf Microbiome as Potential Biocontrol

Agents Against Phytopathogens. *Frontiers in Microbiology*, *10*(November), 1–17.

https://doi.org/10.3389/fmicb.2019.02726

Chardonnet, C. O., Sams, C. E., & Conway, W. S. (1999). Calcium effect on the mycelial cell walls of

Botrytis cinerea. *Phytochemistry*, *52*(6), 967–973. https://doi.org/10.1016/S0031-9422(99)00315-5

Chen, H., Xiao, X., Wang, J., Wu, L., Zheng, Z., & Yu, Z. (2008). Antagonistic effects of volatiles generated

by Bacillus subtilis on spore germination and hyphal growth of the plant pathogen, Botrytis cinerea.

*Biotechnology Letters*, *30*(5), 919–923. https://doi.org/10.1007/s10529-007-9626-9

Chepsergon, J., Motaung, T. E., Bellieny-Rabelo, D., & Moleleki, L. N. (2020). Organize, don't agonize:

Strategic success of phytophthora species. In *Microorganisms* (Vol. 8, Issue 6, pp. 1–21). MDPI AG.

https://doi.org/10.3390/microorganisms8060917

Compant, S., Samad, A., Faist, H., & Sessitsch, A. (2019). A review on the plant microbiome: Ecology,

functions, and emerging trends in microbial application. In *Journal of Advanced Research* (Vol. 19,

pp. 29–37). Elsevier B.V. https://doi.org/10.1016/j.jare.2019.03.004

Coutos-Thévenot, P., Poinssot, B., Bonomelli, A., Yean, H., Breda, C., Buffard, D., Esnault, R., Hain, R., &

Boulay, M. (2001). In vitro tolerance to Botrytis cinerea of grapevine 41B rootstock in transgenic

plants expressing the stilbene synthase Vst1 gene under the control of a pathogen-inducible PR 10

promoter. *Journal of Experimental Botany*, *52*(358), 901–910.

https://doi.org/10.1093/jexbot/52.358.901

Dame, Z. T., Islam, M. T., Helmke, E., von Tiedemann, A., & Laatsch, H. (2016). Oligomycins and

pamamycin homologs impair motility and induce lysis of zoospores of the grapevine downy mildew

pathogen, Plasmopara viticola. *FEMS Microbiology Letters*, *363*(16).

https://doi.org/10.1093/femsle/fnw167

Dastogeer, K. M. G., Tumpa, F. H., Sultana, A., Akter, M. A., & Chakraborty, A. (2020). Plant microbiome–

an account of the factors that shape community composition and diversity. In *Current Plant Biology*

(Vol. 23). Elsevier B.V. https://doi.org/10.1016/j.cpb.2020.100161

Enoki, S., & Suzuki, S. (2016). Pathogenesis-Related Proteins in Grape. In *Grape and Wine Biotechnology*.

InTech. https://doi.org/10.5772/64873

Freitas, M. A., Medeiros, F. H. V., Melo, I. S., Pereira, P. F., Peñaflor, M. F. G. V., Bento, J. M. S., & Paré, P. W.

(2019). Stem inoculation with bacterial strains Bacillus amyloliquefaciens (GB03) and

Microbacterium imperiale (MAIIF2a) mitigates Fusarium root rot in cassava. *Phytoparasitica*, *47*(1),

135–142. https://doi.org/10.1007/s12600-018-0706-2

Giannelli, G., Del Vecchio, L., Cirlini, M., Gozzi, M., Gazza, L., Galaverna, G., Potestio, S., & Visioli, G.

(2024). Exploring the rhizosphere of perennial wheat: potential for plant growth promotion and

biocontrol applications. *Scientific Reports*, *14*(1), 22792. https://doi.org/10.1038/s41598-024-

73818-6

Godard, S., Slacanin, I., Viret, O., & Gindro, K. (2009). Induction of defence mechanisms in grapevine

leaves by emodin- and anthraquinone-rich plant extracts and their conferred resistance to downy

mildew. *Plant Physiology and Biochemistry*, *47*(9), 827–837.

https://doi.org/10.1016/j.plaphy.2009.04.003

Goetz, G., Fkyerat, A., Métais, N., Kunz, M., Tabacchi, R., Pezet, R., & Pont, V. (1999). Resistance factors to

grey mould in grape berries: identification of some phenolics inhibitors of Botrytis cinerea stilbene

oxidase. *Phytochemistry*, *52*(5), 759–767. https://doi.org/10.1016/S0031-9422(99)00351-9

Guerrero, R. F., Aliaño-González, M. J., Puertas, B., Richard, T., & Cantos-Villar, E. (2020). Comparative

analysis of stilbene concentration in grapevine shoots of thirteen Vitis during a three-year study.

*Industrial Crops and Products*, *156*. https://doi.org/10.1016/j.indcrop.2020.112852

Guetsky, R., Shtienberg, D., Elad, Y., & Dinoor, A. (2001). Combining biocontrol agents to reduce the

variability of biological control. *Phytopathology*, *91*(7), 621–627.

https://doi.org/10.1094/PHYTO.2001.91.7.621

Guyer, A., De Vrieze, M., Bönisch, D., Gloor, R., Musa, T., Bodenhausen, N., Bailly, A., & Weisskopf, L.

(2015). The anti-phytophthora effect of selected potato-associated Pseudomonas strains: From the

laboratory to the field. *Frontiers in Microbiology*, *6*(NOV).

https://doi.org/10.3389/fmicb.2015.01309

Han, J.-I., Spain, J. C., Leadbetter, J. R., Ovchinnikova, G., Goodwin, L. A., Han, C. S., Woyke, T., Davenport,

K. W., & Orwin, P. M. (2013). Genome of the Root-Associated Plant Growth-Promoting Bacterium

Variovorax paradoxus Strain EPS. *Genome Announcements*, *1*(5).

https://doi.org/10.1128/genomea.00843-13

Honda, S., Kunii, T., Nohara, K., Wakita, S., Sugahara, Y., Kawakita, M., Oyama, F., & Sakaguchi, M. (2017).

Characterization of a Bacillus thuringiensis chitinase that binds to cellulose and chitin. *AMB Express*,

*7*(1). https://doi.org/10.1186/s13568-017-0352-y

Khatun, A., Farhana, T., Sabir, A. A., Islam, S. M. N., West, H. M., Rahman, M., & Islam, T. (2018).

Pseudomonas and Burkholderia inhibit growth and asexual development of Phytophthora capsici.

*Zeitschrift Fur Naturforschung - Section C Journal of Biosciences*, *73*(3–4), 123–135.

https://doi.org/10.1515/znc-2017-0065

Köhl, J., Medeiros, F. H. V., Lombaers-van der Plas, C., Groenenboom-de Haas, L., & van den Bosch, T.

(2020). Efficacies of bacterial and fungal isolates in biocontrol of Botrytis cinerea and Pseudomonas

syringae pv. tomato and growth promotion in tomato do not correlate. *Biological Control*, *150*.

https://doi.org/10.1016/j.biocontrol.2020.104375

Kong, Q., An, P., Xu, Z., Zhang, R., Qi, J., & Ren, X. (2020). New insights into the alleviating role of

Melaleuca alternifolia oil on metabolites pathway disorder of grapes caused by Aspergillus niger,

verified by corresponding key genes expression. *Food Chemistry*, *327*.

https://doi.org/10.1016/j.foodchem.2020.127083

Legein, M., Smets, W., Vandenheuvel, D., Eilers, T., Muyshondt, B., Prinsen, E., Samson, R., & Lebeer, S.

(2020). Modes of Action of Microbial Biocontrol in the Phyllosphere. In *Frontiers in Microbiology*

(Vol. 11). Frontiers Media S.A. https://doi.org/10.3389/fmicb.2020.01619

LI, F.-X., MA, H.-Q., LIU, J., & ZHANG, C. (2012). Antagonistic Effects of Bacillus cereus Strain B-02 on

Morphology, Ultrastructure and Cytophysiology of Botrytis cinerea. *Polish Journal of Microbiology*,

*61*(2), 119–128. https://doi.org/10.33073/pjm-2012-015

Li, M. Y., Jiao, Y. T., Wang, Y. T., Zhang, N., Wang, B. B., Liu, R. Q., Yin, X., Xu, Y., & Liu, G. T. (2020).

CRISPR/Cas9-mediated VvPR4b editing decreases downy mildew resistance in grapevine (Vitis

vinifera L.). *Horticulture Research*, *7*(1). https://doi.org/10.1038/s41438-020-00371-4

Li, W. J., Qin, Z. H., Zhang, M. H., & Browde, J. (2011). An index method to evaluate growers' pesticide use

for identifying on-farm innovations and effective alternative pest management strategies: A case

study of winegrape in Madera County, California. *Journal of Zhejiang University: Science B*, *12*(3),

226–246. https://doi.org/10.1631/jzus.B0900264

Li, Y., Feng, X., Wang, X., Zheng, L., & Liu, H. (2020). Inhibitory effects of Bacillus licheniformis BL06 on

Phytophthora capsici in pepper by multiple modes of action. *Biological Control*, *144*.

https://doi.org/10.1016/j.biocontrol.2020.104210

Liu, D., Li, K., Hu, J., Wang, W., Liu, X., & Gao, Z. (2019). Biocontrol and action mechanism of bacillus

amyloliquefaciens and bacillus subtilis in soybean phytophthora blight. *International Journal of*

*Molecular Sciences*, *20*(12). https://doi.org/10.3390/ijms20122908

Liu, G. T., Wang, B. B., Lecourieux, D., Li, M. J., Liu, M. B., Liu, R. Q., Shang, B. X., Yin, X., Wang, L. J.,

Lecourieux, F., & Xu, Y. (2021). Proteomic analysis of early-stage incompatible and compatible

interactions between grapevine and P. viticola. *Horticulture Research*, *8*(1).

https://doi.org/10.1038/s41438-021-00533-y

Martínez-Zavala, S. A., Ortiz-Rodríguez, T., Salcedo-Hernández, R., Casados-Vázquez, L. E., Del Rincón-

Castro, M. C., Bideshi, D. K., & Barboza-Corona, J. E. (2024). The chitin-binding domain of Bacillus

thuringiensis ChiA74 inhibits gram-negative bacterial and fungal pathogens of humans and plants.

*International Journal of Biological Macromolecules*, *254*.

https://doi.org/10.1016/j.ijbiomac.2023.128049

Mayer, E., de Quadros, P. D., & Fulthorpe, R. (2019). Plantibacter flavus, Curtobacterium herbarum,

Paenibacillus taichungensis, and Rhizobium selenitireducens endophytes provide host-specific

growth promotion of Arabidopsis thaliana, basil, lettuce, and bok choy plants. *Applied and*

*Environmental Microbiology*, *85*(19). https://doi.org/10.1128/AEM.00383-19

Minchev, Z., Kostenko, O., Soler, R., & Pozo, M. J. (2021). Microbial Consortia for Effective Biocontrol of

Root and Foliar Diseases in Tomato. *Frontiers in Plant Science*, *12*.

https://doi.org/10.3389/fpls.2021.756368

Mondol, M. A. M., Farthouse, J., Islam, M. T., Schüffler, A., & Laatsch, H. (2017). Metabolites from the

Endophytic Fungus Curvularia sp. M12 Act as Motility Inhibitors against Phytophthora capsici

Zoospores. *Journal of Natural Products*, *80*(2), 347–355.

https://doi.org/10.1021/acs.jnatprod.6b00785

Niu, B., Wang, W., Yuan, Z., Sederoff, R. R., Sederoff, H., Chiang, V. L., & Borriss, R. (2020). Microbial

Interactions Within Multiple-Strain Biological Control Agents Impact Soil-Borne Plant Disease. In

*Frontiers in Microbiology* (Vol. 11). Frontiers Media S.A. https://doi.org/10.3389/fmicb.2020.585404

*osafune1992*. (n.d.).

Pascale, A., Proietti, S., Pantelides, I. S., & Stringlis, I. A. (2020). Modulation of the Root Microbiome by

Plant Molecules: The Basis for Targeted Disease Suppression and Plant Growth Promotion. *Frontiers*

*in Plant Science*, *10*(January), 1–23. https://doi.org/10.3389/fpls.2019.01741

Pezet, R., Perret, C., Jean-Denis, J. B., Tabacchi, R., Gindro, K., & Viret, O. (2003). δ-viniferin, a resveratrol

dehydrodimer: One of the major stilbenes synthesized by stressed grapevine leaves. *Journal of*

*Agricultural and Food Chemistry*, *51*(18), 5488–5492. https://doi.org/10.1021/jf030227o

Pezet, R., & Pont, V. (1990). Ultrastructural Observations of Pterostilbene Fungitoxicity in Dormant

Conidia of *Botrytis cinerea* Pers. *Journal of Phytopathology*, *129*(1), 19–30.

https://doi.org/10.1111/j.1439-0434.1990.tb04286.x

Pezet, R., Pont, V., & Hoang-Van, K. (1991). Evidence for oxidative detoxication of pterostilbene and

resveratrol by a laccase-like stilbene oxidase produced by Botrytis cinerea. *Physiological and*

*Molecular Plant Pathology*, *39*(6), 441–450. https://doi.org/10.1016/0885-5765(91)90010-F

Pont, V., & Pezet, R. (1990). Relation Between the Chemical Structure and the Biological Activity of

Hydroxystilbenes Against *Botrytis cinerea*. *Journal of Phytopathology*, *130*(1), 1–8.

https://doi.org/10.1111/j.1439-0434.1990.tb01147.x

Primieri, S., Magnoli, S. M., Koffel, T., Stürmer, S. L., & Bever, J. D. (2022). Perennial, but not annual

legumes synergistically benefit from infection with arbuscular mycorrhizal fungi and rhizobia: a

meta-analysis. *New Phytologist*, *233*(1), 505–514. https://doi.org/10.1111/nph.17787

Rahman, M. U., Liu, X., Wang, X., & Fan, B. (2024). Grapevine gray mold disease: Infection, defense and

management. In *Horticulture Research* (Vol. 11, Issue 9). Oxford University Press.

https://doi.org/10.1093/hr/uhae182

Rai, S., Omar, A. F., Rehan, M., Al-Turki, A., Sagar, A., Ilyas, N., Sayyed, R. Z., & Hasanuzzaman, M. (2023).

Crop microbiome: their role and advances in molecular and omic techniques for the sustenance of

agriculture. In *Planta* (Vol. 257, Issue 2). Springer Science and Business Media Deutschland GmbH.

https://doi.org/10.1007/s00425-022-04052-5

Ray, T., Pandey, A., Pandey, S. S., Singh, S., Shanker, K., & Kalra, A. (2021). Molecular insights into

enhanced resistance of Papaver somniferum against downy mildew by application of endophyte

bacteria Microbacterium sp. SMR1. *Physiologia Plantarum*, *173*(4), 1862–1881.

https://doi.org/10.1111/ppl.13528

Reynolds, E. S. (1963). THE USE OF LEAD CITRATE AT HIGH pH AS AN ELECTRON-OPAQUE STAIN IN

ELECTRON MICROSCOPY. *The Journal of Cell Biology*, *17*(1), 208–212.

https://doi.org/10.1083/jcb.17.1.208

Shang, H., Chen, J., Handelsman, J., & Goodman, R. M. (1999). Behavior of Pythium torulosum Zoospores

During Their Interaction with Tobacco Roots and Bacillus cereus. *Current Microbiology*, *38*(4), 199–

204. https://doi.org/10.1007/PL00006787

Suárez-Estrella, F., Jurado, M. M., López-González, J. A., Toribio, A., Martínez-Gallardo, M. R., Estrella-

González, M. J., & López, M. J. (2023). Seed priming by application of Microbacterium spp. strains

for control of Botrytis cinerea and growth promotion of lettuce plants. *Scientia Horticulturae*, *313*.

https://doi.org/10.1016/j.scienta.2023.111901

Sukhada, M., Manjula, R., & Rawal, R. D. (2011). Evaluation of arbuscular mycorrhiza and other biocontrol

agents against Phytophthora parasitica var. nicotianae infecting papaya (Carica papaya cv. Surya)

and enumeration of pathogen population using immunotechniques. *Biological Control*, *58*(1), 22–

29. https://doi.org/10.1016/j.biocontrol.2011.03.013

Sun, S.-L., Yang, W.-L., Fang, W.-W., Zhao, Y.-X., Guo, L., Dai, Y.-J., & Cann, I. (2018). *The Plant Growth-*

*Promoting Rhizobacterium Variovorax boronicumulans CGMCC 4969 Regulates the Level of Indole-3-*

*Acetic Acid Synthesized from Indole-3-Acetonitrile*. https://doi.org/10.1128/AEM

Syed Ab Rahman, S. F., Singh, E., Pieterse, C. M. J., & Schenk, P. M. (2018). Emerging microbial biocontrol

strategies for plant pathogens. *Plant Science*, *267*(November 2017), 102–111.

https://doi.org/10.1016/j.plantsci.2017.11.012

Tareq, F. S., Hasan, C. M., Lee, H. S., Lee, Y. J., Lee, J. S., Surovy, M. Z., Islam, M. T., & Shin, H. J. (2015).

Gageopeptins A and B, new inhibitors of zoospore motility of the phytopathogen Phytophthora

capsici from a marine-derived bacterium Bacillus sp. 109GGC020. *Bioorganic and Medicinal*

*Chemistry Letters*, *25*(16), 3325–3329. https://doi.org/10.1016/j.bmcl.2015.05.070

Trusov, Y., Sewelam, N., Rookes, J. E., Kunkel, M., Nowak, E., Schenk, P. M., & Botella, J. R. (2009).

Heterotrimeric G proteins-mediated resistance to necrotrophic pathogens includes mechanisms

independent of salicylic acid-, jasmonic acid/ethylene- and abscisic acid-mediated defense signaling.

*Plant Journal*, *58*(1), 69–81. https://doi.org/10.1111/j.1365-313X.2008.03755.x

Van Leeuwen, C., Roby, J. P., Alonso-Villaverde, V., & Gindro, K. (2013). Impact of clonal variability in Vitis

vinifera cabernet franc on grape composition, wine quality, leaf blade stilbene content, and downy

mildew resistance. *Journal of Agricultural and Food Chemistry*, *61*(1), 19–24.

https://doi.org/10.1021/jf304687c

Van Loon, L. C., Rep, M., & Pieterse, C. M. J. (2006). Significance of inducible defense-related proteins in

infected plants. In *Annual Review of Phytopathology* (Vol. 44, pp. 135–162).

https://doi.org/10.1146/annurev.phyto.44.070505.143425

Vionnet, L., De Vrieze, M., Dutartre, A., Gfeller, A., Lüthi, A., L'Haridon, F., & Weisskopf, L. (2018).

Microbial life in the grapevine: What can we expect from the leaf microbiome? *Oeno One*, *52*(3),

205–210. https://doi.org/10.20870/oeno-one.2018.52.3.2120

Viret, O., Spring, J. L., & Gindro, K. (2018). Stilbenes: Biomarkers of grapevine resistance to fungal

diseases. *Oeno One*, *52*(3). https://doi.org/10.20870/oeno-one.2018.52.3.2033

Vlot, A. C., Sales, J. H., Lenk, M., Bauer, K., Brambilla, A., Sommer, A., Chen, Y., Wenig, M., & Nayem, S.

(2021). Systemic propagation of immunity in plants. In *New Phytologist* (Vol. 229, Issue 3, pp. 1234–

1250). Blackwell Publishing Ltd. https://doi.org/10.1111/nph.16953

Waffo-Teguo, P., Krisa, S., Richard, T., & Mérillon, J.-M. (2008). Grapevine Stilbenes and Their Biological

Effects. In *Bioactive Molecules and Medicinal Plants* (pp. 25–54). Springer Berlin Heidelberg.

https://doi.org/10.1007/978-3-540-74603-4_2

Wang, J., Yue, T., He, C., Zhou, Y., Bai, Y., Li, Q., Jiang, W., Huang, Y., & Liu, X. (2023). Biocontrol of tomato

bacterial wilt by a combination of Bacillus subtilis GSJB-1210 and ningnanmycin. *Scientia*

*Horticulturae*, *321*. https://doi.org/10.1016/j.scienta.2023.112296

Yu, X., Zhang, X., Zhang, J., Zhang, L., Jiao, Y., Guo, L., Wang, J., Wang, X., Zhao, J., & Xiang, W. (2022).

Mixtures of suppressive bacteria enhance biological control of tomato bacterial wilt. *Biological*

*Control*, *170*. https://doi.org/10.1016/j.biocontrol.2022.104937

Zarraonaindia, I., Owens, S. M., Weisenhorn, P., West, K., Hampton-Marcell, J., Lax, S., Bokulich, N. A.,

Mills, D. A., Martin, G., Taghavi, S., van der Lelie, D., & Gilbert, J. A. (2015). The soil microbiome

influences grapevine-associated microbiota. *MBio*, *6*(2). https://doi.org/10.1128/mBio.02527-14

Zhu, Y., Li, Y., Zhang, S., Zhang, X., Yao, J., Luo, Q., Sun, F., & Wang, X. (2019). Genome-wide identification

and expression analysis reveal the potential function of ethylene responsive factor gene family in

response to Botrytis cinerea infection and ovule development in grapes (Vitis vinifera L.). *Plant*

*Biology*, *21*(4), 571–584. https://doi.org/10.1111/plb.12943

**Table 1**: Inhibition of *P. viticola* zoospore motility after 30 minutes contact with phyllosphere bacteria and level of symptom development 1 week after leaf disc infection.

| strain | organism / treatment | motility inhibition (%) | SD | significance | symptoms inhibition (%) | SD | significance |
|---|---|---|---|---|---|---|---|
| Cu | copper hydroxide | 100 | 0 | *** | 92 | 3.8 | *** |
| PIP10 | *Pantoea agglomerans* | 100 | 0 | *** | 64 | 21.9 | *** |
| SOP5 | *Microbacterium hominis* | 100 | 0 | ** | 61.8 | 29.3 | *** |
| PIP7 | *Microbacterium hominis* | 99.4 | 1.1 | *** | 66.2 | 29.9 | *** |
| PID6 | *Bacillus cereus* | 99.0 | 1.7 | *** | 83.3 | 10.8 | *** |
| SOD2 | *Bacillus halotolerans* | 99.0 | 1.7 | *** | 22.7 | 39.0 | - |
| CHP14 | *Bacillus subtilis* | 98.8 | 2.1 | *** | 38.5 | 39.2 | ** |
| PID5 | *Bacillus altitudinis* | 98.7 | 1.3 | *** | 18.5 | 61.3 | - |
| SOP51 | *Bacillus pumilus* | 98.7 | 1.2 | *** | 49.4 | 34.4 | *** |
| CHD1 | *Bacillus cereus* | 98.5 | 2.6 | *** | 30.1 | 58.7 | - |
| PIP36 | *Bacillus cereus* | 98.2 | 3.2 | *** | 46.7 | 42.2 | *** |
| SOD14 | *Bacillus thuringiensis* | 97.5 | 2.1 | *** | 4.4 | 40.5 | - |
| CHD4 | *Bacillus subtilis* | 97.1 | 5.0 | *** | 25.4 | 36.1 | - |
| SOD31 | *Bacillus zhangzhouensis* | 95.7 | 4.6 | *** | 26.0 | 46.7 | - |
| CHP20 | *Bacillus cereus* | 95.3 | 7.3 | *** | 18.1 | 51.1 | - |
| CHD2 | *Bacillus cereus* | 95.2 | 4.4 | *** | 33.5 | 36.9 | - |
| SOD6 | *Bacillus cereus* | 92.2 | 4.7 | *** | 39.4 | 47.6 | - |
| PIP2 | *Sphingomonas zeae* | 84.0 | 27.7 | ** | 50.6 | 36.5 | ** |
| PIP17 | *Staphylococcus arlettae* | 81.4 | 12.3 | ** | 39.1 | 42.8 | ** |
| SOD5 | *Bacillus licheniformis* | 79.9 | 28.3 | ** | 10.3 | 48.6 | - |
| PID2 | *Bacillus simplex* | 77.9 | 13.6 | *** | 74.5 | 28.4 | *** |
| CHP29 | *Microbacterium proteolyticum* | 76.3 | 30.7 | ** | 5.0 | 55.0 | - |
| SOD20 | *Bacillus stratosphericus* | 75.5 | 3.3 | * | 19.3 | 50.2 | - |
| PID1 | *Paenibacillus amylolyticus* | 64.1 | 22.9 | *** | 8.2 | 48.8 | - |
| CHP35 | *Bacillus aryabhattai* | 60.2 | 17.5 | * | 35.0 | 58.6 | - |
| CHP6 | *Rhodopseudomonas boonkerdii* | 56.3 | 13.8 | *** | 13.4 | 55.6 | - |
| PID13 | *Bacillus endophyticus* | 51.9 | 16.4 | ** | 52.3 | 41.1 | *** |
| SOP16 | *Bacillus butanolivorans* | 43.1 | 11.1 | ** | 14.9 | 49.8 | - |
| CHD6 | *Micrococcus aloeverae* | 39.7 | 11.1 | ** | 23.1 | 46.7 | - |
| PIP27 | *Rhodopseudomonas boonkerdii* | 36.3 | 31.2 | - | 63.7 | 24.5 | *** |
| PIP4 | *Cupriavidus metallidurans* | 28.0 | 18.3 | - | 60.3 | 32.3 | *** |
| SOD13 | *Paenibacillus xylanexedens* | 35.6 | 13.9 | - | 47.3 | 31.6 | *** |
| PIP3 | *Variovorax paradoxus* | 7.7 | 28.0 | - | 50.6 | 62.6 | *** |
| PIP6 | *Herbaspirillum huttiense* | 3.8 | 22.6 | - | 31.8 | 36.5 | * |

∗ = p < 0.05; ∗∗ = p < 0.01; ∗∗∗ = p < 0.001; - = not significant

**Table 2**: *P. viticola* symptom inhibition 1 week after infection in grapevine leaf discs inoculated with phyllosphere bacteria relative to uninfected control. Only strains that generated significant inhibition are displayed.

| strain | organism / treatment | inhibition (%) | SD | significance |
|---|---|---|---|---|
| Cu | copper hydroxide | 89.9 | 4.9 | *** |
| PIP27 | *Rhodopseudomonas boonkerdii* | 58.6 | 41.8 | ** |
| CHP15 | *Micrococcus yunnanensis* | 46.1 | 49.1 | * |
| PIP7 | *Microbacterium hominis* | 41.6 | 34.1 | ** |
| CHP14 | *Bacillus subtilis* | 39.7 | 32.8 | ** |
| PIP17 | *Staphylococcus arlettae* | 39.1 | 47.3 | * |
| SOD6 | *Bacillus cereus* | 39 | 41.7 | ** |
| SOD13 | *Paenibacillus xylanexedens* | 37.9 | 52.5 | * |
| SOD2 | *Bacillus halotolerans* | 36.6 | 45.5 | ** |
| CHP2 | *Cupriavidus metallidurans* | 35.5 | 47.2 | * |
| PID6 | *Bacillus cereus* | 32.4 | 42.7 | * |
| SOP5 | *Microbacterium hominis* | 28.9 | 24.3 | ** |
| CHP20 | *Bacillus cereus* | 24.8 | 42.4 | * |
| CHP29 | *Microbacterium proteolyticum* | 24 | 28.3 | * |
| SOD31 | *Bacillus zhangzhouensis* | 23.8 | 31.5 | * |

**Table 3**: *B. cinerea* symptom inhibition 1 week after infection in grapevine leaf discs inoculated with phyllosphere bacteria relative to uninoculated control. Only strains that generated significant inhibition are displayed.

| strain | organism / treatment | inhibition (%) | SD | significance |
|---|---|---|---|---|
| CHP29 | *Microbacterium proteolyticum* | 81.4 | 10.5 | *** |
| SOP47 | *Variovorax paradoxus* | 75.4 | 9.7 | *** |
| CHP2 | *Cupriavidus metallidurans* | 70.8 | 19.1 | *** |
| PIP6 | *Herbaspirillum huttiense* | 69.3 | 19 | *** |
| PIP36 | *Bacillus cereus* | 64.8 | 16.1 | ** |
| SOP8 | *Cupriavidus metallidurans* | 63.8 | 19.4 | * |
| PID2 | *Bacillus simplex* | 62.4 | 11.6 | * |
| Botector | *Aureobasidium pullulans* | 61.1 | 14.8 | *** |
| PIP26 | *Plantibacter flavus* | 59.2 | 25.2 | * |
| SOP16 | *Bacillus butanolivorans* | 55.8 | 25.4 | ** |
| SOD13 | *Paenibacillus xylanexedens* | 54.2 | 25.5 | ** |
| SOP48 | *Staphylococcus xylosus* | 52.1 | 25.3 | * |
| CHP26 | *Bacillus drentensis* | 50 | 40.6 | ** |
| CHP20 | *Bacillus cereus* | 49.8 | 43.7 | ** |
| CHD1 | *Bacillus cereus* | 39.4 | 22.2 | * |
| SOD7 | *Bacillus psychrosaccharolyticus* | 38.6 | 33.6 | * |

**Table 4**: Local and systemic *P. viticola* symptom inhibition 1 week after infection in whole plants inoculated with phyllosphere bacteria relative to uninoculated control

| Strain | organism / treatment | local inhibition (%) | SD | significance | systemic inhibition (%) | SD | significance |
|---|---|---|---|---|---|---|---|
| Alginure | Alginure (brown algae) | 98.2 | 0.9 | *** | 98.2 | 0.9 | *** |
| PID6 | *Bacillus cereus* | 82.4 | 21.7 | *** | 68.5 | 38.5 | *** |
| SOP8 | *Cupriavidus metallidurans* | 68.3 | 29.5 | *** | 74 | 19.8 | *** |
| PIP6 | *Herbaspirillum huttiense* | 63.4 | 35.9 | *** | 29.6 | 55.5 | N.S. |
| SOD2 | *Bacillus halotolerans* | 52.6 | 31.9 | *** | 58.3 | 18.1 | *** |
| CHP14 | *Bacillus subtilis* | 38 | 64.4 | * | 31.1 | 73.5 | * |
| CHP2 | *Cupriavidus metallidurans* | 33.8 | 46.6 | N.S. | 25.8 | 44.6 | N.S. |
| PID13 | *Bacillus endophyticus* | 31.2 | 70.8 | N.S. | 32.2 | 53.4 | N.S. |
| CHP2 / PID6 / PIP6 | *Cupriavidus metallidurans / Bacillus cereus / Herbaspirillum huttiense* | 94.2 | 5.5 | *** | 94 | 5.2 | *** |
| CHP14 / PID13 / PIP6 | *Bacillus subtilis / Bacillus endophyticus / Herbaspirillum huttiense* | 88.1 | 8.6 | *** | 90 | 16.2 | *** |
| CHP2 / CHP14 / PID13 | *Cupriavidus metallidurans / Bacillus subtilis / Bacillus endophyticus* | 84.9 | 13.1 | *** | 87.7 | 15.7 | *** |
| CHP2 / CHP14 / SOD2 | *Cupriavidus metallidurans / Bacillus subtilis / Bacillus halotolerans* | 57.6 | 39.2 | ** | 67 | 26.8 | ** |

∗ = p < 0.05; ∗∗ = p < 0.01; ∗∗∗ = p < 0.001; - = not significant

908    **Figure 1:** Transmission electron microscopy image of *P. viticola* zoospores placed for 30 min in

909    contact with water (control), or a solution of the *Bacillus cereus* strain PID6 $OD_{600}$ = 1.

**Figure 2:** Effect of phyllosphere bacteria on *B. cinerea* spore germination. A: Each colored bar

represents the proportion of spores showing the corresponding morphology. Green, normal

germination; yellow, partial inhibition (only the tip of germ tube is visible); orange, swelling; and

red, no germination. Each treatment with bacteria significantly affected the physiology of the

spores, with *p* < 0.05 according to a Chi-square test with Yate's correction. The different types of

germination morphologies are shown in (B to E); B: normal germination; C; absence of

germination; D: partial germination and E: swelling.

**Figure 3**: Effect of consortia of phyllosphere bacteria on *B. cinerea* spore germination. A: Each

colored bar represents the proportion of spores showing the corresponding morphology. green:

normal germination; yellow; partial inhibition (only the tip of germ tube is visible); orange:

swelling and red: no germination. Each treatment with bacteria significantly affected the

physiology of the spores, with $p < 0.05$ according to a Chi-square test with Yate's correction. The

different types of germination morphologies are shown in (B to E); B: normal germination; C;

absence of germination; D: partial germination and E: swelling.

**Figure 4:** Representative pictures of grapevine leaf discs inoculated with phyllosphere bacteria

prior to infection with *P. viticola*. Control: leaf disc not inoculated with bacteria and infected with

*P. viticola zoospores*; PIP27, SOD13 and SOD31: leaf discs inoculated with *Rhodopseudomonas*

*boonkerdii, Paenibacillus xylanexedens* and *Bacillus zhangzhouensis* respectively prior to

infection. Picture were taken 1 week after infection.

949 **Figure 5:** Representative pictures of grapevine leaf discs inoculated with phyllosphere bacteria

950 prior to infection with *B. cinerea*. Control: leaf disc not inoculated with bacteria and infected

951 with *B. cinerea* spores; CHP29, PID2 and SOD7: leaf discs inoculated with *Microbacterium*

952 *proteolyticum*, *Bacillus simplex* and *Bacillus psychrosaccharolyticus* respectively prior to

953 infection. Picture were taken 1 week after infection.

963 **Figure 6:** Level of expression of defense-related gene in grapevine leaf discs inoculated with

964 phyllosphere bacteria that significantly protected the plant against *P. viticola* or *B. cinerea* in the

965 leaf disc assay. Levels of expression were normalized with the two reference genes Actine and

966 60SRP and expressed relative to infected control leaf discs that were not inoculated with

967 bacteria.

| pathogen | strain | organism | LOX | PAL | STS | ROMT | PR-1 | PR-4 |
|---|---|---|---|---|---|---|---|---|
| *P. viticola* | PIP27 | *Rhodopseudomonas boonkerdii* | | | | | | |
| | CHP15 | *Micrococcus yunnanensis* | | | | | | |
| | PIP7 | *Microbacterium hominis* | | | | | | |
| | CHP14 | *Bacillus subtilis* | | | | | | |
| | PIP17 | *Staphylococcus arlettae* | | | | | | |
| | SOD6 | *Bacillus cereus* | | | | | | |
| | SOD13 | *Paenibacillus xylanexedens* | | | | | | |
| | SOD2 | *Bacillus halotolerans* | | | | | | |
| | CHP2 | *Cupriavidus metallidurans* | | | | | | |
| | PID6 | *Bacillus cereus* | | | | | | |
| | SOP5 | *Microbacterium hominis* | | | | | | |
| | CHP20 | *Bacillus cereus* | | | | | | |
| | CHP29 | *Microbacterium proteolyticum* | | | | | | |
| | SOD31 | *Bacillus zhangzhouensis* | | | | | | |
| *B. cinerea* | CHP29 | *Microbacterium proteolyticum* | | | | | | |
| | SOP47 | *Variovorax paradoxus* | | | | | | |
| | CHP2 | *Cupriavidus metallidurans* | | | | | | |
| | PIP6 | *Herbaspirillum huttiense* | | | | | | |
| | PIP36 | *Bacillus cereus* | | | | | | |
| | SOP8 | *Cupriavidus metallidurans* | | | | | | |
| | PID2 | *Bacillus simplex* | | | | | | |

log2 fold change

-2        2

972    **Figure 7:** Stilbene production levels in grapevine leaf discs inoculated with phyllosphere bacteria

973    3 days after infection with *P. viticola* zoospores. Levels of production based on dry weight,

974    relative to infected leaves that were not inoculated with bacteria.

**Figure 8:** Representative pictures of grapevine leaves from whole grapevine plants inoculated

with phyllosphere bacteria prior to infection with *P. viticola*. Leaves were inoculated with

different mixtures of the following six strains: *Cupriavidus metallidurans* (CHP2), *Bacillus cereus*

(PID6), *Herbaspirillum huttiense* (PIP6), *Bacillus subtilis* (CHP14), *Bacillus endophyticus* (PID13)

and *Bacillus halotolerans* (SOD2). Control: non-inoculated leaf infected with *P. viticola zoospores*.

Picture were taken 1 week after infection.

| CONTROL | CHP2/PID6/PIP6 | CHP14/PID13/PIP6 | CHP2/PID6/PIP6 | CHP2/CHP14/SOD2 |