1
2
3
4

Specific cell states underlie complex tissue regeneration in spiny mice

Emilio Oviedo Rivadeneira[1], Robyn Allen[1], Mike Adam[2], and Ashley W. Seifert[1,3,4*]

[1]*Department of Biology, University of Kentucky, Lexington, KY 40506, USA*
[2]*Division of Experimental Hematology and Cancer Biology, Cancer and Blood Diseases Institute, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA*
[3]*Department of Veterinary Anatomy and Physiology, University of Nairobi, Nairobi, Kenya*
[4]*Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD 20815, USA*

19
20
21
22

ORCID: Ashley W. Seifert, 0000-0001-6576-3664

24
25
26
27

Key words: Cell proliferation, regeneration, cell cycle, cell state, fibroblasts, chondroblasts

[*]To whom correspondence should be addressed: Ashley W. Seifert, e-mail: awseifert@uky.edu

41 **SUMMARY STATEMENT**

42 Comparing regenerative vs. fibrotic healing, we identify injury-induced cell states associated

43 with persistent cell cycle progression and complex tissue regeneration in mammals.

44
45
ABSTRACT

Cell proliferation is an elemental feature of epimorphic regeneration in vertebrate taxa. We

previously reported that in contrast to fibrotic repair observed in laboratory mouse (*Mus*) strains,

highly regenerative spiny mice (*Acomys* spp.) exhibit cell cycle progression and cell proliferation

to faithfully replace missing tissue. However, little is known about proliferation dynamics, and

specific cell types and states that may contribute to complex tissue regeneration in mammals.

Using temporal pulse-chase experiments, we show that stromal cells in *Acomys dimidiatus*

rapidly re-enter the cell cycle in response to injury and maintain tight spatiotemporal control of

cell cycle progression to restrict the proliferative population to a distal area relative to the injury.

Conversely, *Mus* stromal cells incorporate thymidine analogs without cell division supporting an

S-phase arrest after D10. Deploying immunostaining and scRNA-seq, we identify several key

cell types (CRABP1+, αSMA+) differentially associated with regenerating versus scar tissue.

Importantly, our single cell data revealed distinct gene expression profiles for cross-species

stromal cell types, identifying cell states specific for regenerative or fibrotic healing. While

CRABP1+ fibroblasts are enriched in *Acomys* ears before and after injury, similar fibroblasts

enriched in young, postnatal *Mus* ears remain unable to promote regeneration. Our data

underscore the finely regulated dynamics of proliferating cells during regeneration and

emphasize that regeneration depends on multiple factors including the presence of specific cell

types and the ability of cells to acquire specific states.

Key Conclusions:

-Differentiated cells in *Acomys*, *Mus* and *Danio* re-enter the cell cycle in response to injury,

while homeostatic cycling cells contribute to blastema formation in *Ambystoma*

-Pulse-chase thymidine analog labeling shows tight spatiotemporal control of proliferating

stromal cells during regeneration in *Acomys*.

-Following injury, CRABP1 and αSMA are expressed in distinct stromal cell populations in

Acomys but are co-expressed in *Mus* stromal cell populations.

-Species-specific cell states underlie regenerative and fibrotic repair

-CRABP1+ cells are lost during embryonic development in *Mus* ear pinna but are retained in

Acomys to adulthood.

-Young neonatal *Mus* with abundant CRABP1+ cells still fail to execute regenerative healing

INTRODUCTION

Epimorphic regeneration is mediated by the accumulation of local progenitor cells that proliferate to form a blastema and subsequently differentiate to replace missing or damaged tissue (Morgan, 1901). With few exceptions (e.g. hydra, colonial animals, etc.), most regeneration-competent invertebrate and vertebrate species use epimorphic regeneration to rebuild missing tissue. Seminal studies using urodeles, zebrafish, and planarians explored the dynamics of proliferating cells to understand their origin, proliferative rates, migration patterns, and responses to different injury types (Hays & Fischman, 1961; Poleo et al., 2001; Wallace, 1981; Wenemoser & Reddien, 2010). For example, thymidine analogs applied to amputated newt limbs revealed that cells start to synthesize DNA in tissues adjacent to the wound site (e.g. muscle, connective tissue, periosteum, etc.) and that these proliferating cells are then liberated and maintained distally from the injury in the blastema (Hays & Fischman, 1961). Similar studies in axolotls investigated cell cycle length, factors required to maintain proliferation and the impact of inhibiting proliferation on regeneration (Maden & Wallace, 1976; Wallace, 1981; Wallace & Maden, 1976). Different types of injury trigger distinct mitotic responses in planarians. An amputation, for example, can cause both local and systemic responses, while any other injury that does not remove tissue triggers only the first systemic response without triggering a local one (Wenemoser & Reddien, 2010). Similarly, studies in zebrafish have improved our understanding of the spatiotemporal distribution of proliferating cells in the blastema (Nechiporuk & Keating, 2002; Poleo et al., 2001). All these studies have advanced our understanding of the dynamics of the proliferative population in highly regenerative animals. However, the spatiotemporal dynamics of proliferating cells in regenerative mammals remains poorly understood.

Complex tissue regeneration in vertebrates has been primarily studied in fishes, amphibians and invertebrates. However, some mammals exhibit bona fide regeneration (Billingham et al., 1959; Borgens, 1982; Carlson, 2005; Gawriluk et al., 2016; Goss & Grimes, 1972; Neeham, 1953; Seifert et al., 2012; Vorontsova et al., 1960). For instance, patterned musculoskeletal tissue is reconstituted after removal with a full thickness hole punch through the ear pinna in some lagomorphs and deomyine rodents (e.g., *Acomys* and *Lophuromys*) (Gawriluk et al., 2016; Goss, 1980; Goss & Grimes, 1972; Seifert et al., 2012). Comparative studies

between regenerative and non-regenerative rodents showed that spiny mice (*Acomys*) can

regenerate ear pinna tissue after full thickness excisional injury; a response mediated by cell

proliferation and blastema formation. In response to an identical injury, however, laboratory

mice (*Mus*) cannot regenerate missing tissue and while cells are activated to re-enter the cell

cycle in response to injury, these cells are arrested shortly afterwards (Gawriluk et al., 2016;

Seifert et al., 2012).

Generally speaking, complex tissue cell proliferation in non-regenerative mammals

occurs exclusively during embryonic development, while cell cycle progression and cell division

become increasingly restricted to stem and progenitor cells in tissues that require constant

turnover during postembryonic development and adulthood (Cheung & Rando, 2013; Liu et al.,

2019; Orford & Scadden, 2008). This raises a question about the identity of cells capable of

maintaining cell cycle progression in regenerative mammals. Said another way, are there specific

cell types or cell states needed for regeneration to occur? In highly regenerative vertebrates,

multiple studies have pinpointed that blastema formation is supported by cellular re-

programming (termed dedifferentiation by most investigators) of lineage-restricted progenitors

located near the wound site (Echeverri et al., 2001; Gerber et al., 2018; Jopling et al., 2010;

Leigh et al., 2018). Interestingly, while lineage-restricted progenitors have been shown to

support digit tip regeneration in *Mus* (G. L. Johnson et al., 2020; Lehoczky et al., 2011;

Rinkevich et al., 2011), subsequent research showed that progenitor cell activation is important

but also the capacity of mesenchymal cells to maintain a specific transcriptomic state associated

with regeneration (Storer et al., 2020). Similar observations come from research investigating

deer antler velvet regeneration. Research comparing injured velvet versus dorsal back skin found

that while both skin types have *Acta2*+ myofibroblast cells, specific transcriptomic profiles were

associated with location. Importantly, this work showed that all velvet skin fibroblasts can access

regenerative regulatory networks that skew them to an immunosuppressive, pro-regenerative

fetal phenotype, while back skin fibroblasts express genes related to an inflammatory, pro-

fibrotic healing outcome (Sinha et al., 2022). Other wound healing studies, suggest that large

skin wounds capable of pro-regenerative healing are biased to use a specific fibroblast subtype

that is inhibited in fibrotic responses; a finding that is also associated with better healing

outcomes in neonatal *Mus* and human skin (Guerrero-Juarez et al., 2019; Mine et al., 2008; Phan

et al., 2021a). Together, these studies support that access to regenerative cell states ***and*** the

presence of specific cell types is essential to promote a regenerative outcome.

Here we use the spiny mouse ear pinna regeneration model to explore controlled cell

proliferation in adult mammals. Leveraging a comparative approach to fibrotic repair in

laboratory mice, we explore the spatiotemporal dynamics of proliferating cells and begin to

identify cell types that participate in regeneration. First, we use a pulse-chase method to quantify

cell cycle activation in regenerative vertebrates with determinate and indeterminate growth and

find that growth mode is not strictly associated with regenerative ability or the lack thereof.

Additionally, this analysis found no evidence of long-range cell activation in response to injury.

Using a pulse-chase labeling strategy, we show that proliferating cells maintain a distal position

relative to the wound site, while cells in the proximal compartment exit the cell cycle and

differentiate. Finally, using a combination of immunostaining and scRNA-sequencing, we

identify similar cell types across species that express distinct transcriptomic states associated

with regenerative or fibrotic repair supporting that species-specific cell states underlie

regeneration and scaring.

METHODS

Animal care. Sexually mature, spiny mice (*Acomys dimidiatus* - aged six months to one year

old) and *Mus musculus*, ~three months old) were obtained from our in-house breeding colonies

at the University of Kentucky. Animals were maintained on a 12:12 h L:D light cycle. Both

species were fed 14% mouse chow (Teklad Global 2014, Harlan Laboratories) and *Acomys* diets

were supplemented with black sunflower seeds. All animal experiments were approved by

University of Kentucky Institutional Animal Care and Use Committee (IACUC) under protocol

2013–3254.

4mm ear punch assay regeneration assay. A 4mm full thickness excisional punch through the

ear pinna serves as a tissue regeneration assay (Gawriluk et al., 2016). Animals were

anesthetized with 3% vaporized isoflurane (v/v) (Henry Schein Animal Health, Dublin, OH) at 1

psi oxygen flow rate. A 4mm biopsy punch (Sklar Instruments, West Chester, PA) was used to

create a through-and-through hole in the right and left ear pinna. Ear tissue was collected at

specified time points using an 8mm biopsy punch (Sklar Instruments, West Chester, PA)

circumscribing the original injury. The published protocol can be found here:

dx.doi.org/10.17504/protocols.io.bp2l6dx65vqe/v1.

Pulse-chase experiments. We injected Ethynyl deoxyuridine (EdU) (10 ug/g) and

Bromodeoxyuridine (BrdU) (1 ug/g) diluted in normal saline, for pulse-chase experiments to

label and track cycling cells (the published protocol can be found here:

dx.doi.org/10.17504/protocols.io.261gerdmol47/v1). Both solutions were injected

intraperitoneally with a difference of five days between the EdU and BrdU injections. EdU and

BrdU injections were given as three separate injections, one every three hours over a period of

nine hours. The left ear pinna was collected after the EdU pulse to assess baseline cell numbers

in S-phase, and the right pinna was collected after the BrdU pulse to track single (EdU+ or

BrdU+) and double positive cells. The injection scheme was as follows: the first group of

animals was injected with EdU at D0 and BrdU at D5, the next group was injected with EdU at

D5 and BrdU at D10 and so on until D30. Hence, tissue samples were collected at 0, 5, 10, 15,

20, 25, and 30-days post injury.

Our experimental scheme allowed us to distinguish cells that (i) were in S-phase at the

time of the EdU injections and stopped proliferating within next five days (Edu+ alone); (ii) were

in S-phase at the time of the EdU injections, divided multiple times over five days diluting the

EdU, and remained in the cell cycle acquiring BrdU (weak EdU+/ strong BrdU+) - because each

cell division dilutes the EdU by half, a scenario where EdU is not detectable is possible, and

these cells would contribute to the BrdU+ alone counts); (iii) cells not in S-phase during the EdU

pulse but were cycling during the BrdU injections (BrdU+ alone); (iv) cells that acquired EdU,

divided only a few times, and remained in the cell cycle during the BrdU chase (strong EdU+/

strong Brdu+).

Histology, EdU detection and Immunohistochemistry. Harvested tissue was fixed

overnight in 10% neutral buffered formalin (NBF) (Thermo Fisher Scientific Inc. Epredia

5701) at 4˚C on a shaker. After 16-20 hrs the tissue was washed 3x with PBS, 3 minutes

per wash, followed by 3x 70% ethanol washes for dehydration and storage at 4˚C. Stored

tissue was processed for paraffin (Leica Biosystems, Buffalo Grove, IL, Formula "R"

REF: 3801450) embedding using a Histo5 tissue processor machine (Milestone). Samples

were sectioned at 5μm. EdU was detected using click chemistry as follows: Tissue

sections were de-paraffinized and rehydrated. Antigen retrieval was performed if co-

stained with antibodies. Tissues were washed two times for 10 minutes in Tris-buffered

saline (TBS) to incubate in the prepared reaction cocktail (see table below) for 30

minutes at RT protected from light. Tissues were washed two times in TBS for 5 minutes

and if co-stained with antibodies were blocked following the fluorescent IHC protocol for

BrdU and cell type specific antibodies tissue sections were deparaffinized and re-

hydrated. Tissues were then exposed to high heat antigen retrieval in a steamer in antigen

unmasking TRIS or citric acid-based solutions (Vector Laboratories, Inc.). After a 5-

minute wash, tissues were blocked in blocking serum (see table below) for 30 minutes at

RT. Tissues were then incubated with primary antibody diluted in blocking serum at 4 ˚C

overnight. After a 5 minutes wash, tissues were incubated in secondary antibody diluted

in blocking serum for 30 min at RT. Tissue samples were incubated with Hoechst 33342

for 5 minutes at RT and then dried and mounted in Prolong. Finally, protein similarity

was compared to assess antibody binding efficacy (Supplementary table S1). For

complete protocol details regarding paraffin sections

(dx.doi.org/10.17504/protocols.io.6qpvr9353vmk/v1), immunohistochemistry

(dx.doi.org/10.17504/protocols.io.n92ldrmd8g5b/v1), and EdU detection on sections

(dx.doi.org/10.17504/protocols.io.e6nvwbdrzvmk/v1).

EdU reaction cocktail	
Components	Volume (ul)
2 M Tris, pH 8.5	5
50 mM CuSO4	2
0.5 M ascorbic acid	20
Alexa fluor azide (0.25 mg/ml)	0.2
ddH2O	73
Total	100

Primary Antibodies					
Antibody	Company, Cat#	RRID	Host species	Dilution	Antigen retrieval
Anti-BrdU (Bromodeoxyuridine)	Abcam, Ab6326	AB_305426	Rat	1:500	Citric Acid

Anti-wide spectrum Cytokeratin	Abcam, Ab9377	AB_307222	Rabbit	1:500	None
Anti-ionized calcium-binding adapter molecule 1 (IBA1)	Novus Biologicals, NB100-10288s	AB_521594	Goat	1:400	Tris
Anti-retinoic acid-binding protein 1 (CRABP1)	Cell Signaling Technology, Inc, 13163	AB_2750569	Rabbit	1:500	Tris
Anti- alpha-Smooth Muscle Actin (alpha-SMA)	Novus Biologicals, NB300-978	AB_2273628	Goat	1:200	Tris
Anti-SRY Box Transcription Factor 9 (SOX9)	Abcam, ab185230	AB_2715497	Rabbit	1:1000	None
Anti-macrophage mannose receptor (MMR) (CD206)	R&D Systems, AF2535	AB_2063012	Goat	1:500	Tris

Secondary Antibodies				
Antibody	**Company, Cat#**	**RRID**	**Host species**	**Dilution**
Alexafluor 594 anti-Rabbit	Thermo Fisher Scientific, A21207	AB_141637	Donkey	1:400
Alexafluor 647 anti-goat	Thermo Fisher Scientific, A21447	AB_2535864	Donkey	1:400
Alexafluor 647 anti-rat	Thermo Fisher Scientific, A21247	AB_141778	Goat	1:400

Western Blotting. Protein extraction and quantification was performed as indicated

(dx.doi.org/10.17504/protocols.io.rm7vzkx12vx1/v1). Western blotting was performed as

indicated (dx.doi.org/10.17504/protocols.io.kqdg3qxz1v25/v1).

Imaging and Cell Quantification. Fluorescent images were collected using Cells Sens from

Olympus in Olympus inverted microscope (Olympus America Inc). Whole tissue sections were

captured in 20X and stitched automatically. Cell counting was manually performed in defined

areas of the images separately in each channel using ImageJ. Percentages of positive cells were

calculated by quantifying nuclei (Hoechst 33342) using the threshold tool in ImageJ.

Statistics. Statistical analyses were performed using JMP (version Pro 15.0, SAS Institute Inc)

and SPSS (version 28.0, IBM) where alpha was set at 0.05 for all statistical analyses. Graphs

were made in excel and JMP. To analyze our pulse chase data across species, we performed

Two-way ANOVAs with either EdU+, BrdU+ or double positive cells (dependent variable), and

days post injury (independent variable) followed by Tukey HSD multiple comparisons to test for

differences in positive cells as a function of time (Supplementary table S2).

To analyze proliferation during regeneration in *Acomys*, we used a linear mixed model to

assess differences in percentage of positive cells between three positions (A, B, C) over time

(D5-D30) (Supplementary table S3). Likelihood ratio testing and Akaike Information Criterion

(AIC) were used to select an appropriate covariance structure. A Kenward-Roger adjustment was

used to correct for negative bias in the 170 standard errors and degrees of freedom calculations

induced by the small sample size. Relevant pairwise comparisons were performed and controlled

for multiple comparisons using Fisher's Least Significant Differences (LSD). Mixed modeling

analysis was performed using SAS 9.4 (SAS Institute, Inc.; Cary, NC, USA).

Single-cell analysis. scRNA-seq data used in this study were download and available through

the Gene Expression Omnibus (GEO) (RRID:SCR_005012) under the accession number

GSE182141. The raw fastqs were processed through the CellRanger pipeline v8.0.1

(RRID:SCR_017344) using 10X Genomics' mm10-2020-A reference genome or a custom

annotated *Acomys dimidiatus* genome (GCA_907164435.1 (mAcoDim1_REL_1905), with the

'include introns' set to true to obtain gene expression matrices.

Background reads were removed using decontX from the celda package using the filtered

barcodes as the cells to keep and the remaining cells in the raw barcode matrix as the background

(Campbell et al., 2024). Doublets were minimized using DoubletFinder (RRID:SCR_018771)

(McGinnis et al., 2019). The R v4.4 (RRID:SCR_001905) library Seurat v5.1

(RRID:SCR_016341) (Hao et al., 2021; Hao et al., 2024)) was used for cell type clustering and

marker gene identification. Cells expressing >500 genes were retained for downstream analysis.

Each sample was normalized SCTransform v2 (RRID:SCR_022146), using the glmGamPoi

method and the number of RNA molecules per cell were regressed out. Samples were integrated

with common anchor genes using the CCA method to minimize sample to sample variation. Cell

clusters were determined by the Louvain algorithm using a resolution of 0.3. UMAP dimension

reduction was done using the first thirty principal components. Marker genes for each cell type

were calculated using the Wilcoxon Rank Sum test returning only genes that are present in a

minimum of 25% of the analyzed cluster. Cross species markers were done as above in cell types

of interest, but the pct.2 score was filtered to be > 0 to eliminate uniquely annotated genes from

dominating the lists. Volcano plots were created with the Enhanced Volcano (Blighe et al., 2024)

R library using the avg_log2FC and p_val_adj values as inputs.

Gene ontology (GO) analyses were performed in the Enrichr platform

(https://maayanlab.cloud/Enrichr/). Only differentially expressed genes with a two-fold change

were included in the analysis of GO Biological Process.

RESULTS

Vertebrate stromal cells rapidly re-enter the cell cycle in response to injury

Complex tissue regeneration in vertebrates requires proliferation to expand cell

populations at the injury site which then undergo expansion and differentiation. Notably, few

papers using salamanders or fish to study regeneration have indicated whether stromal cells are

proliferating during normal homeostasis. To explore proliferation in uninjured tissue, we used

EdU to label cells in S-phase in two highly regenerative amniotes (*Danio rerio* - zebrafish,

Ambystoma mexicanum - axolotl), in an adult mammal with enhanced regenerative ability

(*Acomys dimidiatus* – spiny mouse) and in a poorly regenerating mammal (*Mus musculus* –

outbred ND4 laboratory mouse strain) (Fig. 1A). Examining the ear pinna (spiny mouse and lab

mouse), limb (axolotl) and caudal fin (zebrafish), we found cycling cells in the epidermis from

all four species prior to injury; a finding that was expected given the presence of slow cycling

stem cells in the epidermis (Fig. 1B). In uninjured *Danio* and *Ambystoma* tissue, we also found a

small percentage of cycling cells in the stroma (1% and 0.5% respectively) supporting previous

reports for these species (Fig. 1B-C) (K. Johnson et al., 2018; Simkin & Seifert, 2018). In

contrast, we found only one or two cycling stromal cells across entire tissue sections in *Acomys*

or *Mus* prior to tissue removal (Fig. 1B-C). These data support that stromal cell proliferation is

essentially non-existent prior to injury in mammals, at least in the ear and skin.

A longstanding hypothesis is that cells in non-regenerative vertebrates are refractory to

cell cycle re-entry and progression whereas cells in highly regenerative vertebrates easily re-

enter the cell cycle and proliferate to replace missing tissue (Pajcini et al., 2010; Simkin &

Seifert, 2018; Tanaka et al., 1997; Tanaka & Brockes, 1998). To test the capacity for cell cycle

re-entry in response to injury, we used a pulse chase strategy whereby we injected animals with

EdU, performed the specified injury, injected BrdU at either 15 or 63hrs post injury (hpi) and

then harvested tissue 9hrs later at 24 and 72 hpi (see Methods). We quantified stromal cells that

were EdU+, or BrdU+, or EdU+/BrdU+ (double positive) (Fig. 1B, D). 24 hpa spiny mice and

zebrafish showed the highest percentage of newly cycling cells in response to injury (BrdU+).

Mus cells also re-entered the cell cycle, albeit later at 72 hpa. Axolotl limb tissue, on the other

hand, exhibited very few singly BrdU+ cells but presented the highest number of double positive

cells (Fig. 1D). Importantly, our data demonstrate that stromal cells in *Acomys*, *Mus* and

zebrafish re-enter the cell cycle *de novo* in response to injury with a 48hr lag observed for *Mus*

cells.

To assess cell activation arising from the injury but independent of the local repair

response (i.e., systemic cell activation), we collected the contralateral ear pinna in *Mus* and

Acomys, the contralateral limb in axolotls, and the pelvic fin in zebrafish 24 hpa and 72 hpa after

tissue amputation. Using a pulse chase strategy as described above, we looked for cell cycle re-

entry in uninjured tissues at 24 hpa and 72 hpa. EdU+ cells were cycling prior to injury, BrdU+

cells were those activated systemically and EdU+/BrdU+ were those cycling prior to injury that

were activated systemically. Despite reports in lab mice of long-range cell activation in skeletal

muscle stem cells (Rodgers et al., 2014), in *Acomys* and *Mus*, we did not find a noticeable

response in the contralateral ear pinnae where the number of BrdU+ cells did not exceed 0.5% at

either time point (Fig. S1A-B). Similarly, in axolotl and zebrafish, the number of EdU+ cells at

24 hpa and 72 hpa was similar to the number of positive cells found in uninjured tissue at D0

(see above) (Fig. S1A-B). These data indicate that a systemic response to injury is negligible at

24 hpa and 72hpa in all four species.

Proliferating cells are maintained in a distal position relative to the wound site over time

Acomys species can regenerate the ear pinna after tissue removal with a 4 mm (or larger)

ear hole punch whereas identical tissue excisions in wild or laboratory *Mus* resulted in scar tissue

formation (Seifert et al., 2012). Deeper examination revealed cycling cells at the injury site

during regeneration and failure to maintain a proliferative population during fibrotic healing

(Gawriluk et al., 2016). To more thoroughly understand the kinetics of proliferating cells during

regeneration and fibrotic repair, we again employed a pulse (EdU) chase (BrdU) scheme where

our pulse and chase were separated by a five-day window (see Methods). First, we analyzed

BrdU+ cells to estimate the proliferative rate from injury until D30 and then analyzed the spatial

distribution of cycling cells by quantifying cells adjacent to the injury (aka proximal) and at two

distal locations (i.e., closest to the new epidermis); distal dorsal and distal ventral (Fig. 2A-E). In

addition to the proliferative rate, we also quantified the total area of newly generated tissue from

D10-30 which supported continued addition of new tissue during regeneration (Fig. 2B). In

contrast, after a proliferative burst, no new fibrotic tissue was added in *Mus* ears after D10 (Fig.

2C).

Spatially analyzing BrdU+ cells from D10-30 during regeneration revealed that from D15

to D30, the number of cycling cells was significantly higher in the distal areas compared to the

proximal areas (injury site) while there was no difference between the distal-dorsal and distal-

ventral areas (Fig. 2D and Supplementary Table 1). The exception to this pattern was D20 when

the new tissue areas were small or flat and thus distal and proximal areas were close (Fig. 2D).

Importantly, the proliferative rate captured in the distal compartment remained >10% from D10-

30 which contributed to the steady production of new tissue while cell proliferation in the

proximal compartment decreased over time. (Fig. 2D). When we quantified BrdU+ cells in *Mus*

ears, we observed a narrow extension of new tissue (Fig. 2A). While we observed a proliferative

rate of >25% at D5 in *Mus,* the percent of BrdU+ cells precipitously declined from D10-30

which coincided with no increase in tissue area over this time period (Fig. 2C, E).

To assess the spatial distribution of cycling cells in the stromal compartment during the

proliferative phase of regeneration (D10-30), we first analyzed the epidermis as a simpler tissue

where stem cell localization and organization are well known. Basal stem cells are located in the

deeper layer of the epidermis and show symmetrical and asymmetrical cell division depending

on whether they produce progeny or more stem cells (Blanpain & Fuchs, 2006). Using our pulse-

chase scheme and based on localization, we observed stem, progenitor, and differentiating cells

in the epidermis (Fig. S2A-B). The stratum basale contained slow-cycling stem cells (strong

EdU+) and infrequently dividing stem cells (EdU+/BrdU+) while also indicating rapidly-cycling

progenitor cells (strong BrdU+/diluted EdU+), and recently-cycling progenitor cells (strong

BrdU+ Fig. S2A-B). In addition, we observed progenitor cells having acquired EdU that

migrated apically withdrawing from the cell cycle and undergoing differentiation as per the

absence of BrdU (Fig. S2B, yellow arrow). These cell behaviors demonstrated that our labeling

scheme could effectively capture predicted stem cell behavior in a tissue with known

proliferative properties.

Next we used this scheme to similarly characterize proliferative behavior in the stromal

compartment by quantifying cells that re-entered the cell cycle after injury but arrested within

five days (EdU+ only) possibly to differentiate or re-enter the cell cycle later (slow-cycling

progenitors), and cells that re-entered the cell cycle but maintained their proliferative state over

the same period (EdU+/BrdU+ and BrdU+ only) in both distal and proximal areas relative to the

wound site (Fig. 2F-I). We hypothesized that cell proliferation could either be clonal, with

multiple daughter cells near proliferative progenitors in rosettes or could expand from the wound

site. In the latter case, we predicted to capture progenitors cycling near the wound margin and

daughter cells building up the blastema distally. At D10, we found a random, stochastic pattern

of proliferation with a mixture of EdU+ cells, BrdU+ cells, and double-positive cells in distal and

proximal compartments, where the EdU+ cell population represented the biggest population

(>25%) (Fig. 2F-G). At D25, however, we were able to observe a pattern where rapid-cycling

(BrdU+) and infrequently-dividing (double positive) cells represented half of the proliferative

population (>9%) in the distal compartment with the other half (~8%) being represented by

potentially slow cycling progenitors (EdU+) (Fig. 2H-I). In comparison, the proximal area was

represented mostly by EdU+ cells (5%) that may have started differentiation and a very small

percentage of rapid-cycling and infrequently-dividing (1%) cells that could support expansion of

some differentiating cell types (Fig. 2H-I). Supporting this pattern, we observed differentiation

of new hair follicles and cartilage near the wound site. *In toto*, these results show that the

proliferative progenitor population is maintained in a distal position relative to the wound site.

Furthermore, our pulse chase labeling supports cell cycle progression and asynchronous cell

proliferation during regeneration in *Acomys*, whereas thymidine analog incorporation without

cell division was observed in *Mus*, suggesting an S-phase arrest after D10.

The proliferative population comprises diverse cell types in Acomys

As noted earlier, cell proliferation is required during epimorphic regeneration to supply

cells for new tissue generation. Having shown that cell cycle progression and division occur

during ear pinna regeneration in spiny mice, we next began characterizing the cellular

composition of the proliferative population (PP). To determine cell state or identity among the

PP, we co-labeled with cell type specific markers and EdU (see Methods). As previously shown

in uninjured tissue, we did not find any mesenchymal cell type in a proliferative state. We,

therefore, characterized the distribution of macrophages (IBA1+), chondroblasts (SOX9+) and

two fibroblast subtypes (CRABP1+ or αSMA+) in comparison with *Mus* to identify any obvious

differences in presence and location in various uninjured tissue compartments (Fig. S3A-J). We

found relatively few CD206+ and IBA1+ macrophages in both species (Fig. S3A, B, F, G).

Similarly, αSMA+ cells were few and mostly marking pericytes (Fig. S3E, J). We observed two

populations of SOX9+ cells in both species: one associated with the perichondrium along the

elastic cartilage, and another associated with the sebaceous gland and hair follicle (Fig. S3C, H).

Interestingly, we observed a population of CRABP1+ fibroblasts scattered throughout the

papillary dermis in *Acomy*s while these cells were not detectable in uninjured *Mus* skin (Fig.

S3D, I).

We next examined these same cell types at D5 (Fig. 3A-J). First, we observed that

multiple cell types re-entered the cell cycle in both species. By counting all the EdU+ cells in a

counting frame and then quantifying double positive cells we estimated the fraction of cycling

cells represented by each cell type. Because inflammation peaks at D5 (Gawriluk et al., 2020),

we hypothesized that a large fraction of proliferating cells would be macrophages. Instead,

IBA1+ and CD206+ cells represented only 5.3% and 4.95% of the PP respectively in *Acomys*

and 5.3% and 0.96% in *Mus* (Fig 3A-D). In *Acomys*, the two fibroblast subtypes (CRABP1+ and

αSMA+ cells) represented ~14% and 18.65% respectively while chondroblasts (SOX9+ cells)

represented ~16% of the PP in *Acomys* (Fig. 3E, G, H). Surprisingly, although totally absent

prior to injury in *Mus*, an emergent CRABP1 population represented 28.9% of the PP at D5 (Fig.

3I). SOX9+ and αSMA+ cells represented 7.96%, and 33% of the PP in *Mus* (Fig. 3F, J).

Together, this data supports that various cell types re-enter the cell cycle during both

regeneration and fibrotic repair.

As described earlier, by day 20, only *Acomys* maintains their proliferative population

during regeneration whereas activated cells did not appear to maintain cell cycle progression or

appreciably divide during fibrotic repair. Co-staining with EdU, we confirmed all the cell types

mentioned above remained part of the PP at D20 in *Acomys* (Fig. 4A-C and Fig. S4A, B). Of

note, we observed that CRABP1+, αSMA+, and SOX9+ cells were segregated spatially

throughout the blastema (Fig. 4A-C, G, H and Fig. S5A). At D0, the CRABP1+ population

seemed restricted to the papillary dermis (Fig. S3D and Fig. S5A). However, this population

expanded to other regions deeper in the dermis at D5 (Fig. 3G and Fig. S5A) and occupied the

dorsal part of the blastema with another, smaller population in the ventral papillary zone at D20

(Fig. 4A-A', G, H and Fig. S5A). In contrast, the αSMA+ fibroblast population was absent at D0

(Fig. S3E and Fig. S5A) but was found after injury in a distal and central position where reticular

fibroblasts localize (Fig. 3H-H' and Fig. S5A). At D20, this population was comparatively much

smaller in size and largely non-overlapping with CRABP1+ cells, and located ventrodistally near

the wound epidermis (Fig. 4B-B', G, H and Fig. S5A). Interestingly, while the blastema expands

and both populations maintain cell cycle progression, the αSMA+ fibroblasts seem to maintain a

small population at the very distal tip of the blastema and ends up disappearing in proximal

tissue. While the CRABP1+ population seems much more abundant occupying more space over

time in the blastema at D25 and D30 (Fig. 4G and H). The SOX9+ population was present in the

cartilage before injury (Fig. S3C and Fig. S5A) and distributed in the middle of the blastema

perfectly aligned with the uninjured cartilage after injury (Fig. 4C and Fig. S5A). Finally, IBA1+

and CD206+ cells were spread evenly across the blastema (Fig. S4A-B and Fig. S5A). Assessing

the spatial segregation of these cell types in *Mus*, we noted a much different configuration of

these populations (Fig. 4D-F and Fig. S5B). We found that the CRABP1+ population in *Mus* was

absent at D0 (Fig. S3I and Fig. S5B) and then expanded throughout the injury zone at D5 (Fig. 3I

and Fig. S5B). However, this population became restricted to the very distal tip of the ear and the

staining was significantly fainter at D20 compared to cells we observed at D5 (Fig. 4D and Fig.

S5B). Although, the αSMA population was absent in *Mus* at D0 like in *Acomys* (Fig. S3E, J and

Fig. S5B), it became the dominant population in the distal part of the scar after injury (Fig. 4E

and Fig. S5B). We did observe some Sox9+ cells in the middle of the outgrowth (Fig. 4F and

Fig. S5B), and IBA1+ and CD206+ cells distributed widely throughout the scar tissue (Fig. S4C,

D and Fig. S5B).

These results show that the same cell types re-enter the cell cycle after injury in both

species, but only *Acomys* maintains cells in a proliferative state during regeneration.

Furthermore, we show a distinct spatial segregation of cell types in regeneration and scarring and

demonstrate that the CRABP1 population is present in *Acomys* before and after injury, while

there is only a transient expansion of this population in *Mus* immediately after injury. To further

support the expression pattern of CRABP1 observed in *Acomys* and *Mus*, we performed a

western blot and found high expression before and after injury in *Acomys* and little to no

expression in *Mus* before injury with an increase in expression at D10 and decrease at D20 (Fig.

S5C). Together, these data support that an expansive and persistent CRABP1+ population is

unique to regeneration.

Species-specific cell states separate scaring and regeneration phenotypes

To support and extend our proliferation and cell studies, we interrogated a single-cell

RNA sequencing (scRNA-seq) dataset harvested from uninjured ear pinna (D0) and healing

tissue at D3, 5, 10 and 15 from *Acomys* and *Mus* (Simkin et al., 2024). Unbiased clustering of

batch corrected data for all time points (see Methods) revealed the same cell populations in both

species (Fig. S7). Next, we extracted the four main stromal cell populations, fibroblasts (cluster

18), chondroblasts (cluster 19), lymphatic endothelial cells (cluster 16) and vascular endothelial

cells (cluster 9), and generated a new UMAP (Fig. 5A). This re-clustering revealed eight

subpopulations among the four original stromal populations: two populations each of vascular

and lymphatic endothelial cells, three fibroblast subtypes, and a single chondroblast population

(Fig. 5B). To investigate markers from our immunohistochemical analysis, we plotted *Crabp1*,

Acta2 (αSMA), and *Sox9* across cells on this UMAP (Fig. 5B). Interestingly, we found that

Acta2 transcripts were strongly associated with one fibroblast subtype (cluster 7) while *Crabp1*

was totally excluded from this cluster (Fig. 5B-D). *Sox9* marked chondroblasts (cluster two) in

both species (Fig. 5E). Since we noticed cells expressing *Acta2* in clusters four and six in both

species, we plotted *CRABP1 and Acta2* together and found very few double positive cells in

Acomys (Fig. S8A). In contrast, fibroblast cluster six was mostly double-positive for *Crabp1* and

Acta2 in *Mus* (Fig. S8B). In addition, fibroblast cluster four and lymphatic endothelial cluster

five were heavily enriched for *Acta2* in *Mus* (Fig. S8B). Together with our immunohistochemical

results, these data support a distinct fibroblast population that is *Acta2*+ in both species (cluster

7). However, they also support that a *Crabp1+* fibroblast subtype exists in *Acomys* that does not

express *Acta2* whereas in *Mus*, *Crabp1* appears to mark only a transitional cell state to finally

express *Acta2*. Moreover, increased *Acta2* positivity among additional fibroblasts and lymphatic

endothelial cells in *Mus* support a bias in the mouse healing response towards fibrosis.

To further understand intrinsic differences between the fibroblast subpopulations in both

species, we ran differential gene expression (DGE) analysis comparing the three *Acomys*

fibroblast clusters against the three *Mus* clusters (Fig. 5F-K). This analysis revealed 1497 genes

(488 in *Acomys*, 1009 in *Mus*) differentially upregulated in cluster four, 1228 (510 in *Acomys*,

718 in *Mus*) in cluster six, and 743 (248 in *Acomys*, 495 in *Mus*) in cluster seven among both

species (Fig 5F, H, J). Gene ontology of each cluster separated by species revealed a clear

segregation in fibroblast responses to amputation (Supplementary Table S4). The three

fibroblasts clusters in *Mus* were associated with fibrotic pathways such as: cellular response to

Transforming Growth Factor Beta stimulus (GO:0071560), cellular response to hypoxia

(GO:0071456), and negative regulation of epithelial cell proliferation (GO:0050680), among

others (Fig 5G, I, K). In contrast, *Acomys* subpopulations were associated with telomere

maintenance and cell proliferation pathways. Clusters four and six, for example, were associated

with positive regulation of establishment of protein localization to telomere (GO:1904851),

positive regulation of telomerase RNA localization to Cajal Body (GO:1904874), and response

to Endoplasmic Reticulum Stress (ERS) (GO:0034976) (Fig. 5G, I). Interestingly, *Acta2* (marker

associated with myofibroblasts) expressing cluster seven was associated with regulation of fat

cell differentiation (GO:0045598), positive regulation of translation (GO:0045727) and

regulation of membrane protein ectodomain proteolysis (GO:0051043) (Fig 5K). Chondroblasts

(cluster two) also followed the same pattern with 2234 genes (786 in *Acomys*, 1448 in *Mus*)

differentially upregulated. *Acomys* chondroblasts upregulated genes associated with translation

(GO:0006412), response to hydrogen peroxide (GO:0042542) and gene expression

(GO:0010467), while *Mus* chondroblasts upregulated genes associated with cellular response to

Transforming Growth Factor Beta stimulus (GO:0071560) and regulation of Transforming

Growth Factor Beta Receptor signaling pathway (GO:0017015) (Fig. S9B). Of note, all four

clusters (two, four, six and seven) in *Acomys* highly upregulated *Zc3h10* and *Tssc4* (Fig. 5F, H, J

and fig. S9A). The first gene has been shown to boost mitochondrial function (Audano et al.,

2018), while the second gene promotes cell survival and proliferation in breast cancer tumors

(Chen et al., 2022).

Together, these data show that all four *Mus* subpopulations activate fibrosis-related

pathways, while the four subpopulations in *Acomys* are biased toward pathways associated with

cell proliferation and growth. Importantly, while the same markers may be able to identify

fibroblast subtypes in both species, the intrinsic ability of each subtype to respond to injury

differs.

CRABP1+ fibroblasts do not induce regeneration in neonatal Mus.

Previous studies using very large wounds in lab mice observed hair follicle cells in the

center of the wound bed, an observation that has been attributed to the presence of *Crabp1+*

fibroblasts (the so-called WIHN model) (Jiang & Rinkevich, 2021; Phan et al., 2021b). Since we

did not observe CRABP1+ cells in uninjured *Mus* ear pinna tissue, but did observe them in

Acomys ear dermis prior to injury and during regeneration, we hypothesized these cells might

disappear during *Mus* ear development. To test this hypothesis, we assessed CRABP1 reactivity

in embryonic, neonatal, juvenile and adult ear tissue (Fig. 6A). In *Acomys,* we found CRABP1+

reactivity to cells across all stages (embryonic and neonatal) and into adulthood (8 months old)

(Fig. 6A). CRABP1 reactivity was strongest at embryonic stages but always restricted to the

developing dermis where it persisted into adulthood, mainly in the papillary dermis (Fig. 6A). In

contrast, we found that *Mus* fibroblasts started to lose CRABP1 expression around P20 and it

was totally absent from the ear pinna in three month old adults (Fig. 6A).

Based on these observations, we hypothesized that challenging *Mus* ear pinnae using our

regeneration assay in the presence of CRABP1+ fibroblasts might lead to enhanced ear hole

regeneration in *Mus*. Therefore, we performed a 4 mm ear punch in P20 animals and found that

neonatal ear holes produce tissue till D15 (Fig. 6B-C). Similarly, we performed 2 mm punches in

P23 animals and reported tissue growth till D20 (Fig. S10A-B), although we did not observe

regeneration. We went back and performed another assay in P15 *Mus* and collected tissue at D5

and D20. Although P15 uninjured tissue also has CRABP1+ fibroblasts, CRABP1 reactivity

after injury was similar to that observed in adult *Mus* with increased expression at D5 and then

disappearing by D20 (Fig 6D). Together, these data show the presence of a CRABP1+ fibroblast

population in neonatal *Mus* is not sufficient to promote a regenerative response and support an

intrinsic difference in the capacity for this population to respond to injury across species.

DISCUSSION

We previously reported that in response to an ear pinna injury, *Acomys* cells progress

through the cell cycle and maintain proliferation during regeneration. In response to an identical

injury in laboratory mouse strains (*Mus*), cell proliferation in the stromal compartment was

minimal, where cells initiated proliferation only to arrest shortly afterwards (Gawriluk et al.,

2016). Here, we conducted a comprehensive spatiotemporal analysis of cell cycle re-entry,

progression, and proliferation during regeneration and found cycling cells maintained a distal

position relative to the original injury, while proximal cells exited the cell cycle and

differentiated. Importantly, these data revealed finely controlled regulation of the cell cycle in

response to injury such that while the injury served as an activation trigger in both species, cells

in tissue destined to regenerate maintained cell cycle progression while those participating in

fibrotic repair arrested. Importantly, proliferation in *Acomys* did not lead to tissue overgrowth

demonstrating regulated shutdown of cell cycle progression. Examples of cell proliferation in

organ stroma where dedicated stem cell populations are not present are rarely observed except

for neoplasms where cell proliferation is uncontrolled and caused by the accumulation of genetic

mutations (Hanahan & Weinberg, 2011). Tightly controlled cell proliferation is a hallmark of

regenerating tissues, occurring in amputated zebrafish caudal fins (Nechiporuk & Keating, 2002;

Poleo et al., 2001) and newt limbs (Hays & Fischman, 1961; Wallace & Maden, 1976).

Although the evolutionary basis for the presence of regenerative ability among some

vertebrate clades remains controversial, indeterminate growth has been hypothesized as a

contributing factor because animals with determinate growth tend not to regenerate as adults

(Froehlich et al., 2013; Sebens, 1987). This longstanding hypothesis suggests that indeterminate

growers maintain populations of somatic stem or progenitor cells that behave like slow-cycling

stem cells to support growth. There are two arms to this hypothesis. The first is the presence of

cycling cells during normal homeostasis and growth lowers the barrier to cell cycle re-entry. The

second is that once cycling, cells do not activate tumor suppressor machinery during S-phase

progression to initiate cell cycle exit. These ideas stem from observations using newt myotubes

which easily re-enter the cell cycle in culture after serum stimulation in contrast to mouse

myotubes formed from C2C12 cells which are refractory to cell cycle re-entry after the same

treatment (Tanaka et al., 1997; Tanaka & Brockes, 1998). *Mus* myotubes can be forced to re-

enter through the cell cycle by deleting or inactivating tumor suppressors such us the

retinoblastoma protein (pRB) or p16^{ink4a} (Gu et al., 1993; Schneider et al., 1994). The question

arises, do cycling cells in uninjured tissue contribute to regeneration?

579   To examine this question, we compared cell cycle re-entry and progression prior to and

after injury between determinate (*Acomys*, regeneration-competent; *Mus*, regeneration-

incompetent) and indeterminate growers (*Ambystoma* and *Danio* – regeneration-competent). In

line with previous observations (K. Johnson et al., 2018; Simkin & Seifert, 2018), we found a

small population of slow-cycling stromal cells present during homeostasis in indeterminate

growers while we failed to detect these cells in determinate growers. However, the cycling

population observed pre-injury in zebrafish scarcely contributed to blastema formation. Instead,

cells that re-entered the cell cycle in response to injury were the major contributors to

regeneration. This result supports a previous study in zebrafish where slow-cycling cells marked

for eight weeks before injury did not contribute to blastema formation (Nechiporuk & Keating,

2002). In our experiments, axolotls had the highest number of double positive cells after injury

suggesting that slow-cycling cells substantially contribute to the blastema. However, a relatively

long S-phase time (~53hrs) for axolotl cells (Wallace & Maden, 1976) may explain this result.

Importantly, our data show that cells in both *Acomys* and *Mus* re-enter the cell cycle in response

to injury independent of their regenerative ability. An open question is how cell cycle

progression is controlled for almost forty days and then downregulated so as not to create tissue

overgrowth. Considered *in toto*, our cross-species analysis rejects the notion that a determinate

growth mode prevents cell cycle re-entry and supports that the contribution of slow-cycling cells

to the blastema is minimal, at least in fish.

598   Another question we examined with our cross-species study was the extent to which

injury induced a systemic activation of cells distant from the injury site. Work in lab mice found

that acute muscle injury elicits a systemic response in which contralateral-located stem cells

(CSCs) re-enter the cell cycle and access a priming state named G$_{Alert}$ (Rodgers et al., 2014). In

contrast to quiescent stem cells (QSCs), CSCs in mouse show an increased propensity to cycle,

although their numbers remain low in comparison with stem cells located at the injury site

(Rodgers et al., 2014). A similar systemic response was observed in axolotl limbs contralateral to

amputated limbs (Johnson et al., 2018). Using contralateral control tissue collected in our

experiments, we looked for primed cells *in vivo* following injury. In contrast to muscle stem cells

in *Mus*, we found a negligible stromal response in *Acomys* and *Mus* with less than 0.5% of

contralateral ear pinna cells responding to a systemic signal. This difference may be explained by

our methodology used to assess proliferation and by the specific cell types analyzed. While we

used immunofluorescence to quantify active or primed mesenchymal cells in tissue sections,

previous studies in mouse muscle used FACS sorted satellite stem cells to assess *in vivo* BrdU

incorporation. Supporting our observations in *Acomys* and *Mus*, we could not find a significant

response in contralateral axolotl limbs or zebrafish pelvic fins, at least in the first 72hrs after

injury. While these results are at odds with a single axolotl study, those data included epidermal

cells in their cell counts and later time points (Johnson et al., 2018). Together, our data do not

support significant cell cycle re-entry in uninjured contralateral tissue from the four species

analyzed.

Contemporary regeneration studies have revisited cell type contributions to the blastema

with many studies exploring lineage contributions using transgenics in regenerative models. The

general consensus is that the blastema particular to an injured tissue is formed from a

heterogeneous population of cells at the amputation plane (Seifert & Muneoka, 2018). Although

various stem cell reservoirs have been discovered in different mammalian tissues, except for the

digit tip, exploring cell type contributions during complex tissue regeneration in mammals has

been impossible. While we could not identify stem cells with our methods, our data support that

Acomys expands pre-existing cell types to participate in regeneration via cell proliferation. One

exception in our analysis were αSMA+ fibroblasts which appeared after injury but did not

maintain αSMA identity in newly differentiated tissue. Interestingly, our scRNA-seq data

indicated that an *Acta2* expressing population expresses genes related to the regulation of fat cell

differentiation suggesting that some myofibroblasts may differentiate into adipocytes in a

regenerative context. Similarly, αSMA+ myofibroblasts were shown to differentiate into

adipocytes when exposed to bone morphogenetic protein (BMP) triggered by neogenic hair

follicles in large skin wounds (Plikus et al., 2017). Further supporting our observations, recent

data from another *Acomys* ear pinna regeneration study reported that αSMA fibroblasts

disappear after wound closure with only ACTA2+ vascular smooth muscle cells remaining

(Brewer et al., 2021). In addition, we observed high expression of TGFβ pathway components in

Mus, but not *Acomys* fibroblasts. Taken with the previously observed insensitivity of *Acomys*

fibroblasts to TGFβ induced myofibroblast differentiation (Brewer et al., 2021), *Acomys* appears

to support reduced myofibroblast activation. While these data suggest a possible differentiation

trajectory for myofibroblasts, the exact dynamics of αSMA+ cells await further study in a

regenerative context.

In contrast to their role during regeneration, the role of myofibroblasts during fibrotic

skin repair is well known (rev. in Plikus et al., 2021). Previous studies in skin have described

first wound responders as a large proportion of reticular fibroblasts including αSMA fibroblasts

(Driskell et al., 2013) which rapidly seal the wound area promoting scaring and inhibiting hair

follicle formation (Correa-Gallegos et al., 2019). These data support our observations in *Mus*

where αSMA+ fibroblasts are the main cell population in newly generated scar tissue. Together,

these data show that αSMA fibroblasts are dominant in scaring but seem to be transient in

regeneration.

CRABP1+ papillary fibroblasts abundantly participate in the regeneration of neonatal

skin and deer antler velvet and thus we were interested to interrogate their behavior in our ear

pinna assay (Phan et al., 2021b; Sinha et al., 2022). In contrast to the ephemeral nature of the

αSMA+ population in *Acomys*, we observed a stark difference in CRABP1+ cells during

regeneration and fibrotic repair. While we were unable to detect CRABP1+ cells prior to injury

in *Mus*, these cells appeared at D5 as a subset of cycling cells only to disappear by D20. In

Acomys, however, we detected CRABP1+ cells prior to injury and this population expanded to

participate in blastema formation. CRABP1 has been found in a variety of regeneration contexts.

For instance, this protein is expressed in *Mus* papillary fibroblasts that expand in large,

somewhat regenerative wounds (Guerrero-Juarez et al., 2019). Other examples are the

regenerating wounds in newborn (P2) mice that have a higher proportion of CRABP1+

fibroblasts (Phan et al., 2021a) and the velvet skin on deer antlers which have regeneration-

primed fibroblasts expressing genes putatively associated with a regenerative phenotype (e.g.,

CRABP1, RUNX1, and *PRSS35*) (Sinha et al., 2022).

Rather than the presence or lack of a specific cell type, our scRNA-seq data suggests that

it may also be one of cell state. Although all four stromal subclusters express some of the same

markers in *Acomys* and *Mus*, our DGE and GO analyses revealed that the four *Mus* subclusters

express biological processes related to fibrosis. In contrast, all four subclusters in *Acomys* seem to use biological processes related with cell proliferation; high transcription and telomere maintenance and genes that boost mitochondrial function and cell survival. Specifically, a subpopulation in both species that highly expressed *Acta2* exhibited divergent transcriptomic responses. Future studies are necessary to address whether *Acomys* αSMA+ cells transiently become myofibroblast-like only to later differentiate into adipocytes as suggested by our GO analysis. Our finding that two subpopulations expressed high telomerase and endoplasmic reticulum stress (ERS) related activity supported the persistence of proliferation in these cells. Telomerase activity is low in most non-proliferating somatic cells in humans and mice, while high activity has been reported in highly proliferating immortal and cancer cells (Kim et al., 1994; Sha & Bacchetti, 1997). Moreover, inhibition of telomerase results in death of tumor cells *in vitro* and *in vivo* showing the importance of this enzyme for tumor growth (Hahn et al., 1999). Similarly, various ERS responses have been shown to promote cell proliferation and survival in various types of cancer (Zhang et al., 2024). These data suggest that *Acomys* can access and use pathways other adult mammals do not outside of tumorigenesis.

As a functional test of cell type versus cell state as a driver of regeneration, we assessed the role of CRABP1+ fibroblasts after injury in neonatal mice before this population disappears. Previous studies have shown that papillary CRABP1+ fibroblasts are enriched in regenerative neonatal human and mouse skin but are lost early during postnatal development (Mine et al., 2008; Sinha et al., 2022). Examining neonatal and early postnatal stages, we found CRABP1+ fibroblasts in the *Mus* ear pinna which persisted until approximately P20. Using our ear punch assay, we did not find a significant improvement in ear hole closure or evidence of regeneration in animals injured at P15 or P20. These results support that the presence of a cell population alone is less important that the ability for cells to acquire a specific cell state needed to facilitate regeneration. A recent examination of regeneration and scarring phenotypes across eleven murid rodents, underscored the binary nature of a regenerative phenotype in that species appear to either completely regenerate or not (i.e., no partial regeneration) (REF). This supports that complex tissue regeneration requires multiple features across stereotypical healing processes such as intrinsic cell cycle licensing (Gawriluk et al., 2016; Tomasso et al., 2023), goldilocks regulation of reactive oxygen species (ROS) (Saxena et al., 2019) and a regenerative skewed immune cell response (Gawriluk et al., 2020; Simkin et al., 2017, 2024). Future work

investigating specific cell states across more regenerative and non-regenerative mammals will

undoubtably shed light on a blueprint for complex tissue regeneration in mammals.

ACKNOWLEDGEMENTS

We would like to thank Josh Sarli, Brennan Riddell and Ava Musarra for animal husbandry. We

thank all members of the Seifert lab for helpful discussions in developing the manuscript and

Brennan Riddle for helping to create the GO graphs. This research was funded in part by grants

from the NIH (R01AR070313) and Aligning Science Across Parkinson's (ASAP-020495) through

the Michael J. Fox Foundation for Parkinson's Research (MJFF) to AWS. For the purpose of open

access, the author has applied a CC BY public copyright license to all Author Accepted

Manuscripts arising from this submission.

AUTHOR CONTRIBUTIONS

EOR, RA, and AWS designed research. EOR, MA and RA performed research. EOR, MA and

AWS analyzed data. EOR and AWS wrote the paper, and all authors commented on and edited the

paper.

COMPETING INTERESTS

The authors declare no competing interests.

AVAILABILITY STATEMENT

The data, code, protocols, and key lab materials used and generated in this study are listed in a

Key Resource Table alongside their persistent identifiers (Supplementary Tables).

REFERENCES

Audano, M., Pedretti, S., Cermenati, G., Brioschi, E., Diaferia, G. R., Ghisletti, S., Cuomo, A.,
Bonaldi, T., Salerno, F., Mora, M., Grigore, L., Garlaschelli, K., Baragetti, A., Bonacina, F.,
Catapano, A. L., Norata, G. D., Crestani, M., Caruso, D., Saez, E., … Mitro, N. (2018).
Zc3h10 is a novel mitochondrial regulator. *EMBO Reports*, *19*(4).
https://doi.org/10.15252/embr.201745531

Billingham, R. E., Mangold, R., & Silvers, W. K. (1959). THE NEOGENESIS OF SKIN IN THE ANTLERS OF DEER*. *Annals of the New York Academy of Sciences*, *83*(3), 491–498. https://doi.org/10.1111/J.1749-6632.1960.TB40922.X

Blanpain, C., & Fuchs, E. (2006). *Epidermal Stem Cells of the Skin*.

Blighe K, Rana S, Lewis M (2024). *EnhancedVolcano: Publication-ready volcano plots with enhanced colouring and labeling*. R package version 1.24.0, https://github.com/kevinblighe/EnhancedVolcano.

Borgens, R. B. (1982). Mice Regrow the Tips of Their Foretoes. *Science*, *217*(4561), 747–750. https://doi.org/10.1126/SCIENCE.7100922

Brewer, C. M., Nelson, B. R., Wakenight, P., Collins, S. J., Okamura, D. M., Dong, X. R., Mahoney, W. M., McKenna, A., Shendure, J., Timms, A., Millen, K. J., & Majesky, M. W. (2021). Adaptations in Hippo-Yap signaling and myofibroblast fate underlie scar-free ear appendage wound healing in spiny mice. *Developmental Cell*, *56*(19), 2722–2740. https://doi.org/10.1016/j.devcel.2021.09.008

Carlson, B. M. (2005). Some principles of regeneration in mammalian systems. *Anatomical Record. Part B, New Anatomist*, *287*(1), 4–13. https://doi.org/10.1002/AR.B.20079

Chen, Y., Zhang, Z., Henson, E. S., Cuddihy, A., Haigh, K., Wang, R., Haigh, J. J., & Gibson, S. B. (2022). Autophagy inhibition by TSSC4 (tumor suppressing subtransferable candidate 4) contributes to sustainable cancer cell growth. *Autophagy*, *18*(6), 1274–1296. https://doi.org/10.1080/15548627.2021.1973338

Cheung, T. H., & Rando, T. A. (2013). Molecular regulation of stem cell quiescence. In *Nature Reviews Molecular Cell Biology* (Vol. 14, Issue 6, pp. 329–340). https://doi.org/10.1038/nrm3591

Correa-Gallegos, D., Jiang, D., Christ, S., Ramesh, P., Ye, H., Wannemacher, J., Kalgudde Gopal, S., Yu, Q., Aichler, M., Walch, A., Mirastschijski, U., Volz, T., & Rinkevich, Y. (2019). Patch repair of deep wounds by mobilized fascia. *Nature*, *576*(7786), 287–292. https://doi.org/10.1038/s41586-019-1794-y

Driskell, R. R., Lichtenberger, B. M., Hoste, E., Kretzschmar, K., Simons, B. D., Charalambous, M., Ferron, S. R., Herault, Y., Pavlovic, G., Ferguson-Smith, A. C., & Watt, F. M. (2013). Distinct fibroblast lineages determine dermal architecture in skin development and repair. *Nature*, *504*(7479), 277–281. https://doi.org/10.1038/nature12783

Echeverri, K., Clarke, J. D. W., & Tanaka, E. M. (2001). In vivo imaging indicates muscle fiber dedifferentiation is a major contributor to the regenerating tail blastema. *Developmental Biology*, *236*(1), 151–164. https://doi.org/10.1006/dbio.2001.0312

Froehlich, J. M., Fowler, Z. G., Galt, N. J., Smith, D. L., & Biga, P. R. (2013). Sarcopenia and piscines: The case for indeterminate-growing fish as unique genetic model organisms in aging

and longevity research. *Frontiers in Genetics*, *4*(AUG). https://doi.org/10.3389/fgene.2013.00159

Gawriluk, T. R., Simkin, J., Hacker, C. K., Kimani, J. M., Kiama, S. G., Ezenwa, V. O., & Seifert, A. W. (2020). Complex Tissue Regeneration in Mammals Is Associated With Reduced Inflammatory Cytokines and an Influx of T Cells. *Frontiers in Immunology*, *11*. https://doi.org/10.3389/FIMMU.2020.01695/FULL

Gawriluk, T. R., Simkin, J., Thompson, K. L., Biswas, S. K., Clare-Salzler, Z., Kimani, J. M., Kiama, S. G., Smith, J. J., Ezenwa, V. O., & Seifert, A. W. (2016). Comparative analysis of ear-hole closure identifies epimorphic regeneration as a discrete trait in mammals. *Nature Communications*, *7*. https://doi.org/10.1038/ncomms11164

Gerber, T., Murawala, P., Knapp, D., Masselink, W., Schuez, M., Hermann, S., Gac-Santel, M., Nowoshilow, S., Kageyama, J., Khattak, S., Currie, J. D., Camp, J. G., Tanaka, E. M., & Treutlein, B. (2018). Single-cell analysis uncovers convergence of cell identities during axolotl limb regeneration. *Science*, *362*(6413). https://doi.org/10.1126/science.aaq0681

Goss, R. J. (1980). Prospects for Regeneration in Man. In *SECTION III BASIC SCIENCE AND PATHOLOGY*.

Goss, R. J., & Grimes, L. N. (1972). Tissue Interactions in the Regeneration of Rabbit Ear Holes. In *AM. Zooi.or.isr* (Vol. 12). https://academic.oup.com/icb/article/12/1/151/2107413

Gu, W., Schneider, J. W., Condorelli, G., Kaushai, S., Yahdavi, V., & Nadai-Ginard, B. (1993). Interaction of Myogenic Factors and the Retinoblastoma Protein Mediates Muscle Cell Commitment and Differentiation. In *Cell* (Vol. 72).

Guerrero-Juarez, C. F., Dedhia, P. H., Jin, S., Ruiz-Vega, R., Ma, D., Liu, Y., Yamaga, K., Shestova, O., Gay, D. L., Yang, Z., Kessenbrock, K., Nie, Q., Pear, W. S., Cotsarelis, G., & Plikus, M. V. (2019). Single-cell analysis reveals fibroblast heterogeneity and myeloid-derived adipocyte progenitors in murine skin wounds. *Nature Communications*, *10*(1). https://doi.org/10.1038/s41467-018-08247-x

HAHN, W. C., STEWART, S. A., BROOKS, M. W., YORK, S. G., EATON, E., KURACHI, A., BEIJERSBERGEN, R. L., KNOLL, J. H. M., MEYERSON, M., & WEINBERG, R. A. (1999). Inhibition of telomerase limits the growth of human cancer cells. *Nature*.

Hanahan, D., & Weinberg, R. A. (2011). Hallmarks of cancer: The next generation. In *Cell* (Vol. 144, Issue 5, pp. 646–674). https://doi.org/10.1016/j.cell.2011.02.013

Hao, Y., Stuart, T., Kowalski, M. H., Choudhary, S., Hoffman, P., Hartman, A., Srivastava, A., Molla, G., Madad, S., Fernandez-Granda, C., & Satija, R. (2024). Dictionary learning for integrative, multimodal and scalable single-cell analysis. *Nature Biotechnology*, *42*(2), 293–304. https://doi.org/10.1038/s41587-023-01767-y

Hao Y, Hao S, Andersen-Nissen E, Mauck WM 3rd, Zheng S, Butler A, Lee MJ, Wilk AJ, Darby C, Zager M, Hoffman P, Stoeckius M, Papalexi E, Mimitou EP, Jain J, Srivastava A, Stuart

T, Fleming LM, Yeung B, Rogers AJ, McElrath JM, Blish CA, Gottardo R, Smibert P, Satija
R. (2021). Integrated analysis of multimodal single-cell data. Cell. Jun 24;184(13):3573-
3587.e29. doi: 10.1016/j.cell.2021.04.048. Epub 2021 May 31. PMID: 34062119; PMCID:
PMC8238499.

Hays, E. D., & Fischman, D. A. (1961). Origin of the Blastema in Regenerating Limbs of the Newt
Triturus viridescens An Autoradiographic Study Using Tritiated Thymidine to Follow Cell
Proliferation and Migration'. In *DEVELOPMENTAL BIOLOGY* (Vol. 3).

Jiang, D., & Rinkevich, Y. (2021). Distinct fibroblasts in scars and regeneration. In *Current
Opinion in Genetics and Development* (Vol. 70, pp. 7–14). Elsevier Ltd.
https://doi.org/10.1016/j.gde.2021.04.005

Johnson, G. L., Masias, E. J., & Lehoczky, J. A. (2020). Cellular Heterogeneity and Lineage
Restriction during Mouse Digit Tip Regeneration at Single-Cell Resolution. *Developmental
Cell*, *52*(4), 525-540.e5. https://doi.org/10.1016/j.devcel.2020.01.026

Johnson, K., Bateman, J., DiTommaso, T., Wong, A. Y., & Whited, J. L. (2018). Systemic cell
cycle activation is induced following complex tissue injury in axolotl. *Developmental
Biology*, *433*(2), 461–472. https://doi.org/10.1016/j.ydbio.2017.07.010

Jopling, C., Sleep, E., Raya, M., Martí, M., Raya, A., & Belmonte, J. C. I. (2010). Zebrafish heart
regeneration occurs by cardiomyocyte dedifferentiation and proliferation. *Nature*, *464*(7288),
606–609. https://doi.org/10.1038/nature08899

Kim, Nam. W., Piatyszek, Miezyslaw. A., Prowse, Karen. R., Harley, Calvin. B., West, Michael.
D., Ho, Peter. L. C., Coviello, Gina. M., Wright, Woodring. E., Weinrich, Scott. L., & Shay,
Jerry. W. (1994). Specific association of human telomerase activity with immortal cells and
cancer. *Science*.

Lehoczky, J. A., Robert, B., & Tabin, C. J. (2011). Mouse digit tip regeneration is mediated by
fate-restricted progenitor cells. *Proceedings of the National Academy of Sciences of the
United States of America*, *108*(51), 20609–20614. https://doi.org/10.1073/pnas.1118017108

Leigh, N. D., Dunlap, G. S., Johnson, K., Mariano, R., Oshiro, R., Wong, A. Y., Bryant, D. M.,
Miller, B. M., Ratner, A., Chen, A., Ye, W. W., Haas, B. J., & Whited, J. L. (2018).
Transcriptomic landscape of the blastema niche in regenerating adult axolotl limbs at single-
cell resolution. *Nature Communications*, *9*(1). https://doi.org/10.1038/s41467-018-07604-0

Liu, L., Michowski, W., Kolodziejczyk, A., & Sicinski, P. (2019). The cell cycle in stem cell
proliferation, pluripotency and differentiation. In *Nature Cell Biology* (Vol. 21, Issue 9, pp.
1060–1067). Nature Publishing Group. https://doi.org/10.1038/s41556-019-0384-4

Maden, M., & Wallace, H. (1976). *How X-Rays inhibit Amphibian Limb Regeneration*.

Mine, S., Fortunel, N. O., Pageon, H., & Asselineau, D. (2008). Aging alters functionally human
dermal papillary fibroblasts but not reticular fibroblasts: A new view of skin morphogenesis
and aging. *PLoS ONE*, *3*(12). https://doi.org/10.1371/journal.pone.0004066

Morgan, T. H. (1901). *Regeneration /*. New York : http://hdl.handle.net/2027/mdp.39015001913774

Nechiporuk, A., & Keating, M. T. (2002). A proliferation gradient between proximal and msxb-expressing distal blastema directs zebrafish fin regeneration. *Development*.

Neeham, A. (1953). Regeneration and Wound-healing. *BioScience*, *3*(1), 16–16. https://doi.org/10.1093/AIBSBULLETIN/3.1.16-F

Orford, K. W., & Scadden, D. T. (2008). Deconstructing stem cell self-renewal: Genetic insights into cell-cycle regulation. In *Nature Reviews Genetics* (Vol. 9, Issue 2, pp. 115–128). https://doi.org/10.1038/nrg2269

Pajcini, K. V., Corbel, S. Y., Sage, J., Pomerantz, J. H., & Blau, H. M. (2010). Transient inactivation of Rb and ARF yields regenerative cells from postmitotic mammalian muscle. *Cell Stem Cell*, *7*(2), 198–213. https://doi.org/10.1016/j.stem.2010.05.022

Phan, Q. M., Sinha, S., Biernaskie, J., & Driskell, R. R. (2021a). Single-cell transcriptomic analysis of small and large wounds reveals the distinct spatial organization of regenerative fibroblasts. *Experimental Dermatology*, *30*(1), 92–101. https://doi.org/10.1111/exd.14244

Phan, Q. M., Sinha, S., Biernaskie, J., & Driskell, R. R. (2021b). Single-cell transcriptomic analysis of small and large wounds reveals the distinct spatial organization of regenerative fibroblasts. *Experimental Dermatology*, *30*(1), 92–101. https://doi.org/10.1111/exd.14244

Plikus, M. V, Guerrero-Juarez, C. F., Ito, M., Li, Y. R., Dedhia, P. H., Zheng, Y., Shao, M., Gay, D. L., Ramos, R., Hsi, T.-C., Oh, J. W., Wang, X., Ramirez, A., Konopelski, S. E., Elzein, A., Wang, A., Supapannachart, R. J., Lee, H.-L., Lim, C. H., … Cotsarelis, G. (2017). *Regeneration of fat cells from myofibroblasts during wound healing*. https://www.science.org

Plikus, M. V., Wang, X., Sinha, S., Forte, E., Thompson, S. M., Herzog, E. L., Driskell, R. R., Rosenthal, N., Biernaskie, J., & Horsley, V. (2021). Fibroblasts: Origins, definitions, and functions in health and disease. In *Cell* (Vol. 184, Issue 15, pp. 3852–3872). Elsevier B.V. https://doi.org/10.1016/j.cell.2021.06.024

Poleo, G., Brown, C. W., Laforest, L., & Akimenko, M. A. (2001). Cell proliferation and movement during early fin regeneration in zebrafish. In *Developmental Dynamics* (Vol. 221, Issue 4, pp. 380–390). https://doi.org/10.1002/dvdy.1152

Rinkevich, Y., Lindau, P., Ueno, H., Longaker, M. T., & Weissman, I. L. (2011). Germ-layer and lineage-restricted stem/progenitors regenerate the mouse digit tip. *Nature*, *476*(7361), 409–414. https://doi.org/10.1038/nature10346

Rodgers, J. T., King, K. Y., Brett, J. O., Cromie, M. J., Charville, G. W., Maguire, K. K., Brunson, C., Mastey, N., Liu, L., Tsai, C. R., Goodell, M. A., & Rando, T. A. (2014). MTORC1 controls the adaptive transition of quiescent stem cells from G 0 to GAlert. *Nature*, *510*(7505), 393–396. https://doi.org/10.1038/nature13255

Saxena, S., Vekaria, H., Sullivan, P. G., & Seifert, A. W. (2019). Connective tissue fibroblasts
from highly regenerative mammals are refractory to ROS-induced cellular senescence.
Nature Communications, *10*(1). https://doi.org/10.1038/s41467-019-12398-w

Schneider, J. W., Gu, W., Zhu, L., Mahdavi, V., & Nadal-Ginard, B. (1994). Reversal of Terminal
Differentiation Mediated by p107 in Rb-/- Muscle Cells. In *New Series* (Vol. 264, Issue
5164).

Sebens, K. P. (1987). The ecology of indeterminate growth in animals. *Ann. Rev. Ecol. Syst.*

Seifert, A. W., Kiama, S. G., Seifert, M. G., Goheen, J. R., Palmer, T. M., & Maden, M. (2012).
Skin shedding and tissue regeneration in African spiny mice (Acomys). *Nature*, *489*(7417),
561–565. https://doi.org/10.1038/nature11499

Seifert, A. W., & Muneoka, K. (2018). The blastema and epimorphic regeneration in mammals. In
Developmental Biology (Vol. 433, Issue 2, pp. 190–199). Elsevier Inc.
https://doi.org/10.1016/j.ydbio.2017.08.007

Sha, J. W., & Bacchetti, S. (1997). A Survey of Telomerase Activity in Human Cancer. In *Eur J*
Cancer (Vol. 33, Issue 5).

Simkin, J., Aloysius, A., Adam, M., Safaee, F., Donahue, R. R., Biswas, S., Lakhani, Z., Gensel,
J. C., Thybert, D., Potter, S., & Seifert, A. W. (2024). Tissue-resident macrophages
specifically express Lactotransferrin and Vegfc during ear pinna regeneration in spiny mice.
Developmental Cell, *59*(4), 496-516.e6. https://doi.org/10.1016/j.devcel.2023.12.017

Simkin, J., Gawriluk, T. R., Gensel, J. C., & Seifert, A. W. (2017). *Macrophages are necessary*
for epimorphic regeneration in African spiny mice. https://doi.org/10.7554/eLife.24623.001

Simkin, J., & Seifert, A. W. (2018). Concise Review: Translating Regenerative Biology into
Clinically Relevant Therapies: Are We on the Right Path? In *Stem Cells Translational*
Medicine (Vol. 7, Issue 2, pp. 220–231). John Wiley and Sons Ltd.
https://doi.org/10.1002/sctm.17-0213

Sinha, S., Sparks, H. D., Labit, E., Robbins, H. N., Gowing, K., Jaffer, A., Kutluberk, E., Arora,
R., Raredon, M. S. B., Cao, L., Swanson, S., Jiang, P., Hee, O., Pope, H., Workentine, M.,
Todkar, K., Sharma, N., Bharadia, S., Chockalingam, K., … Biernaskie, J. (2022). Fibroblast
inflammatory priming determines regenerative versus fibrotic skin repair in reindeer. *Cell*,
185(25), 4717-4736.e25. https://doi.org/10.1016/j.cell.2022.11.004

Storer, M. A., Mahmud, N., Karamboulas, K., Borrett, M. J., Yuzwa, S. A., Gont, A., Androschuk,
A., Sefton, M. V., Kaplan, D. R., & Miller, F. D. (2020). Acquisition of a Unique
Mesenchymal Precursor-like Blastema State Underlies Successful Adult Mammalian Digit
Tip Regeneration. *Developmental Cell*, *52*(4), 509-524.e9.
https://doi.org/10.1016/j.devcel.2019.12.004

Tanaka, E. M., & Brockes, J. P. (1998). A target of thrombin activation promotes cell cycle re-
entry by urodele muscle cells. *Wound Repair and Regeneration*, *6*(4), S-371-S-381.
https://doi.org/10.1046/J.1524-475X.1998.60413.X/FORMAT/PDF

Tanaka, E. M., Gann, A. A. F., Gates, P. B., & Brockes, J. P. (1997). Newt Myotubes Reenter the
Cell Cycle by Phosphorylation of the Retinoblastoma Protein. In *The Journal of Cell Biology*
(Vol. 136, Issue 1).

Tomasso, A., Koopmans, T., Lijnzaad, P., Bartscherer, K., & Seifert, A. W. (2023). *An ERK-*
dependent molecular switch antagonizes fibrosis and promotes regeneration in spiny mice
(Acomys). https://www.science.org

Vorontsova, M. A., Liosner, L. D., Allen, P. M., & Billett, F. (1960). *Asexual propagation and*
regeneration.

Wallace, H. (1981). *Vertebrate Limb Regeneration* (John Wiley & Sons, Ed.; First edition).

Wallace, H., & Maden, M. (1976). THE CELL CYCLE DURING AMPHIBIAN LIMB
REGENERATION. In *J. CeUSci.zo*.

Wenemoser, D., & Reddien, P. W. (2010). Planarian regeneration involves distinct stem cell
responses to wounds and tissue absence. *Developmental Biology*, *344*(2), 979–991.
https://doi.org/10.1016/j.ydbio.2010.06.017

Zhang, W., Shi, Y., Oyang, L., Cui, S., Li, S., Li, J., Liu, L., Li, Y., Peng, M., Tan, S., Xia, L., Lin,
J., Xu, X., Wu, N., Peng, Q., Tang, Y., Luo, X., Liao, Q., Jiang, X., & Zhou, Y. (2024).
Endoplasmic reticulum stress—a key guardian in cancer. In *Cell Death Discovery* (Vol. 10,
Issue 1). Springer Nature. https://doi.org/10.1038/s41420-024-02110-3

Figure 1. Cell cycle re-entry before and after injury in *Acomys*, *Mus*, *Danio* and

***Ambystoma*.** A) Tissue section planes and EdU/BrdU pulse-chase scheme indicating injections

and harvest times; D = distal, P = proximal. B) Representative images of EdU+ and BrdU+ cells

in uninjured tissue (D0), D1 and D3. In uninjured tissue, white arrows = positive cells in the

epidermis and green arrows = positive cells in the stromal compartment. In injured tissue, red

arrows = EdU+ cells, light blue arrows = BrdU+ cells, and white arrows = double positive cells

(EdU+/BrdU+). C) EdU+ cell quantification as a percentage of the total number of cells in

uninjured tissue from all four species. Autofluorescent tissue, including red blood cells, stains in

multiple channels (yellow). D) EdU+, Brdu+ and EdU+/BrdU+ cell quantification as a

percentage of the total number of cells at D1 and D3 post-injury. Images represent n = 3

biological samples/time point in *Acomys* and *Mus*. See Methods for cell counting quantification.

In *Danio* (zebrafish), n = 9 at D0, n = 5 at D1 and n = 4 at D3. In *Ambystoma* (axolotl), n = 12 at

D0, n = 6 at D1, n = 6 at D3. (* $p<0.05$, ** $p<0.01$).

Figure 2. Spatiotemporal distribution of proliferating cells in *Acomys* and *Mus*. A)

Representative images of blastema and scar tissue changes over time (D5-D30), Ca = cartilage,

WM = wound margin, HF = hair follicle. B-C) Blastema or scar tissue area in mm^2 from D10 to

D30 based on tissue sections. D-E) Quantification of BrdU+ cells as a percentage of the total

number of cells in three areas (A = blue, B = red, C = green) of the blastema in *Acomys* or the

scar in *Mus* from D5 to D30. F-H) Representative images of D10 (F) and D25 (H) blastema

tissue with EdU+, BrdU+ and double positive cells. Red arrows = EdU+ cells, light blue arrows

= BrdU+ cells, and white arrows = double positive cells (EdU+/BrdU+). G-I) Quantification of

EdU+, BrdU+ and EdU+/BrdU+ cells as a percentage of the total number of cells in distal and

proximal areas relative to the wound site at D10 (G) and D25 (I). n = 5 per time point per

species. (* *p*<0.05).

Figure 3. S-phase quantification of fibroblasts, chondroblasts and macrophages after injury at D5 in *Acomys* and *Mus*. A-D) Cycling macrophages double labeled with IBA1 and EdU in *Acomys* (A-A') and *Mus* (C-C') or with CD206 and EdU in *Acomys* (B-B') and *Mus* (D-

D'). E-F) Chondroblasts double labeled with Sox9 and EdU in *Acomys* (E-E') and *Mus* (F-F').

G-J) Fibroblasts double labeled with CRABP1 and EdU in *Acomys* (G-G') and *Mus* (I-I'). or

with αSMA and EdU in *Acomys* (H-H') and *Mus* (J-J'). White arrows indicate double positive

cells (EdU+/cell type marker). Pie charts indicate percentage double positive cells of the total

number of counted EdU+ cells. n = 5/species. RBC = red blood cells.

Figure 4. Spatial distribution of cell types within blastema or scar at D20. A-C) CRABP1+

(A-A') or αSMA+ (B-B') fibroblasts, and Sox9+ chondroblasts (C-C') in *Acomys* at D20. D-F)

CRABP1+ (D-D') or αSMA+ (E-E') fibroblasts, and Sox9+ chondroblasts (F-F') in *Mus* at D20.

G) CRABP1 and αSMA double staining in *Acomys* over time (D5-D30). H) Cartoon showing

general distribution of CRABP1+ and αSMA+ fibroblasts in *Acomys*. White arrows indicate

double positive cells (EdU+/cell type marker). Pie charts indicate percentage double positive of

the total number of counted EdU+ cells. n = 5 per species. RBC = red blood cells.

Figure 5. Single-cell analysis reveals different fibroblast cell states in *Acomys* and *Mus*. Cell

suspensions collected from healing tissue preps representing D0, 3, 5, 10 and 15 were batch-

corrected and plotted together (Fig. S7) (see Methods). A) UMAP projection of stromal cell

types re-clustered from all cells (Fig. S7) from both species. B) UMAP projection of stromal

subpopulations separated by species. Clusters represent subpopulations of major subtypes

indicated in A. C-E) Feature plots for *Crabp1* (C), *Acta2* (D), and *Sox9* (E) separated by species.

F, H, J) Volcano plots showing differentially upregulated genes between *Acomys* and *Mus* from

clusters four (F), six (H), and seven (J) in B. Gene ontology Biological Process based on

differentially upregulated genes between *Acomys* and *Mus* clusters four (G), six (I), and seven

(K). Dot size represents the relative number of genes in each plot. Upregulated genes associated

with each pathway are in color boxes. Refer to supplementary table S4 to see full list of genes.

1007

1008

Figure 6. Tissue injury in neonatal mouse ears when CRABP1+ fibroblasts are present

does not facilitate regeneration. A) Immunostaining for CRABP1 in *Acomys* and *Mus*. Age in

days indicated in upper right corner of each image. B) Images showing representative ear hole

closure from P20 injured *Mus* after a 4 mm ear punch. C) Quantification of ear hole closure area

from D5 to D40. D) CRABP1 staining in P15 *Mus* before and after injury. n = 3 per species per

time point. Green channel shows autofluorescence including red blood cells (RBC).

1016

1017