1

Specific cell states underlie complex tissue regeneration in spiny mice

Emilio Oviedo Rivadeneira[1], Robyn Allen[1], Mike Adam[2], and Ashley W. Seifert[1,3,4*]

[1]Department of Biology, University of Kentucky, Lexington, KY 40506, USA
[2]Division of Experimental Hematology and Cancer Biology, Cancer and Blood Diseases Institute, Cincinnati Children's Hospital Medical Center, Cincinnati, OH, USA
[3]Department of Veterinary Anatomy and Physiology, University of Nairobi, Nairobi, Kenya
[4]Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD 20815, USA

ORCID: Ashley W. Seifert, 0000-0001-6576-3664

Key words: Cell proliferation, regeneration, cell cycle, cell state, fibroblasts, chondroblasts

*To whom correspondence should be addressed: Ashley W. Seifert, e-mail: awseifert@uky.edu

SUMMARY STATEMENT

Comparing regenerative vs. fibrotic healing, we identify injury-induced cell states associated with persistent cell cycle progression and complex tissue regeneration in mammals.

ABSTRACT

Cell proliferation is an elemental feature of epimorphic regeneration in vertebrate taxa. We previously reported that in contrast to fibrotic repair observed in laboratory mouse (*Mus*) strains, highly regenerative spiny mice (*Acomys* spp.) exhibit cell cycle progression and cell proliferation to faithfully replace missing tissue. However, little is known about proliferation dynamics, and specific cell types and states that may contribute to complex tissue regeneration in mammals. Using temporal pulse-chase experiments, we show that stromal cells in *Acomys dimidiatus* rapidly re-enter the cell cycle in response to injury and maintain tight spatiotemporal control of cell cycle progression to restrict the proliferative population to a distal area relative to the injury. Conversely, *Mus* stromal cells incorporate thymidine analogs without cell division supporting an S-phase arrest after D10. Deploying immunostaining and scRNA-seq, we identify several key cell types (CRABP1+, αSMA+) differentially associated with regenerating versus scar tissue. Importantly, our single cell data revealed distinct gene expression profiles for cross-species stromal cell types, identifying cell states specific for regenerative or fibrotic healing. While CRABP1+ fibroblasts are enriched in *Acomys* ears before and after injury, similar fibroblasts enriched in young, postnatal *Mus* ears remain unable to promote regeneration. Our data underscore the finely regulated dynamics of proliferating cells during regeneration and emphasize that regeneration depends on multiple factors including the presence of specific cell types and the ability of cells to acquire specific states.

68 **Key Conclusions:**

69 -Differentiated cells in *Acomys*, *Mus* and *Danio* re-enter the cell cycle in response to injury,

70 while homeostatic cycling cells contribute to blastema formation in *Ambystoma*

71 -Pulse-chase thymidine analog labeling shows tight spatiotemporal control of proliferating

72 stromal cells during regeneration in *Acomys*.

73 -Following injury, CRABP1 and αSMA are expressed in distinct stromal cell populations in

74 *Acomys* but are co-expressed in *Mus* stromal cell populations.

75 -Species-specific cell states underlie regenerative and fibrotic repair

76 -CRABP1+ cells are lost during embryonic development in *Mus* ear pinna but are retained in

77 *Acomys* to adulthood.

78 -Young neonatal *Mus* with abundant CRABP1+ cells still fail to execute regenerative healing

INTRODUCTION

Epimorphic regeneration is mediated by proliferating cells that form a blastema and subsequently differentiate to replace missing tissue (Morgan, 1901). Nearly all regenerative species use epimorphic regeneration to rebuild missing tissue. Seminal studies using urodeles, zebrafish, and planarians explored the dynamics of proliferating cells to understand their origin, proliferative rates, migration patterns, and responses to different injury types (Hays & Fischman, 1961; Poleo et al., 2001; Wallace, 1981; Wenemoser & Reddien, 2010). For example, thymidine analogs applied to amputated newt limbs revealed that cells start to synthesize DNA in differentiated tissues adjacent to the wound site (Hays & Fischman, 1961). These cells are then liberated and displaced distally to form the blastema. Similar studies in axolotls quantified cell cycle length, identified factors that maintain proliferation, and blocked proliferation to study regeneration (Maden & Wallace, 1976; Wallace, 1981; Wallace & Maden, 1976). In planarians, different injury types trigger distinct mitotic responses. For example, amputation causes local and systemic responses, while injuries that do not remove tissue only trigger a systemic response (Wenemoser & Reddien, 2010). Similarly, zebrafish regeneration studies have characterized the spatiotemporal distribution of proliferating cells across the blastema (Nechiporuk & Keating, 2002; Poleo et al., 2001). Considering these studies, an open question is the degree to which patterns and processes governing cell proliferation may hold during instances of epimorphic regeneration in mammals.

Complex tissue regeneration in vertebrates has been primarily studied in fishes, amphibians and invertebrates. However, some mammals exhibit bona fide regeneration (Billingham et al., 1959; Borgens, 1982; Carlson, 2005; Gawriluk et al., 2016; Goss & Grimes, 1972; Neeham, 1953; Seifert et al., 2012; Vorontsova et al., 1960). For instance, patterned musculoskeletal tissue is reconstituted after removal with a full thickness hole punch through the ear pinna in some lagomorphs and deomyine rodents (e.g., *Acomys* and *Lophuromys*) (Gawriluk et al., 2016; Goss, 1980; Goss & Grimes, 1972; Joseph & Dyson, 1966; Riddell et al., 2025; Seifert et al., 2012). Comparative studies between regenerative and non-regenerative rodents showed that spiny mice (*Acomys*) can regenerate ear pinna tissue after full thickness excisional injury; a response mediated by cell proliferation and blastema formation. In response to an identical injury, however, laboratory mice (*Mus*) cannot regenerate missing tissue and while cells

are activated to re-enter the cell cycle in response to injury, these cells are arrested shortly

afterwards (Gawriluk et al., 2016; Seifert et al., 2012).

In adult mammals, cell division is generally restricted to stem and progenitor cells in

tissues that require constant turnover during homeostasis (Cheung & Rando, 2013; Liu et al.,

2019; Orford & Scadden, 2008). In highly regenerative mammals, are there certain cell types or

states that can specifically re-enter the cell cycle and proliferate? In highly regenerative

vertebrates, blastema formation appears to be supported by cellular re-programming (aka

dedifferentiation) of lineage-restricted progenitors located near the wound site (Echeverri et al.,

2001; Gerber et al., 2018; Jopling et al., 2010; Leigh et al., 2018). Interestingly, while lineage-

restricted progenitors undoubtably support mouse digit tip regeneration (G. L. Johnson et al.,

2020; Lehoczky et al., 2011; Rinkevich et al., 2011), the capacity of mesenchymal cells to

maintain a specific transcriptomic state is also associated with regeneration (Storer et al., 2020).

Similar observations come from research comparing injured deer antler velvet (regenerative) to

dorsal back skin (non-regenerative) where both skin types have *Acta2+* myofibroblast cells, but

specific transcriptomic profiles in these cells are associated with location. Importantly, velvet

fibroblasts access regulatory networks skewing them towards an immunosuppressive, pro-

regenerative fetal phenotype, while back skin fibroblasts express networks related to an

inflammatory, pro-fibrotic outcome (Sinha et al., 2022). Wound healing studies in large skin

wounds that regenerate hair follicles are biased to use a specific fibroblast subtype that is

inhibited in fibrotic responses; a finding that is also associated with better healing outcomes in

neonatal *Mus* and human skin (Guerrero-Juarez et al., 2019; Mine et al., 2008; Phan et al.,

2021a). Together, these studies support that access to regenerative cell states ***and*** the presence of

specific cell types is essential to promote a regenerative outcome.

Here we use the spiny mouse ear pinna regeneration model compared to fibrotic repair of

identical injuries in laboratory mice to explore the spatiotemporal dynamics of proliferating cells

and begin identifying cell types that participate in regeneration. First, we use a pulse-chase

method to quantify cell cycle activation in regenerative vertebrates with determinate and

indeterminate growth and find that growth mode is not strictly associated with regenerative

ability or the lack thereof. Additionally, we find no evidence of long-range cell activation in

response to injury. Using a pulse-chase labeling strategy in *Acomys*, we show that proliferating

cells maintain a distal position relative to the wound site, while cells in the proximal

143 compartment exit the cell cycle and differentiate. Finally, using a combination of

144 immunostaining and scRNA-sequencing, we identify similar cell types across species that

145 express distinct transcriptomic states associated with regenerative or fibrotic repair supporting

146 that species-specific cell states underlie regeneration and scaring.

148 **METHODS**

150 **Animal care**. Sexually mature, spiny mice (*Acomys dimidiatus* - aged six months to one year

151 old) and *Mus musculus*, ~three months old) were obtained from our in-house breeding colonies

152 at the University of Kentucky. Animals were maintained on a 12:12 h L:D light cycle. Both

153 species were fed 14% mouse chow (Teklad Global 2014, Harlan Laboratories) and *Acomys* diets

154 were supplemented with black sunflower seeds. All animal experiments were approved by

155 University of Kentucky Institutional Animal Care and Use Committee (IACUC) under protocol

156 2013–3254.

158 **4mm ear punch assay regeneration assay**. A 4mm full thickness excisional punch through the

159 ear pinna serves as a tissue regeneration assay (Gawriluk et al., 2016). Animals were

160 anesthetized with 3% vaporized isoflurane (v/v) (Henry Schein Animal Health, Dublin, OH) at 1

161 psi oxygen flow rate. A 4mm biopsy punch (Sklar Instruments, West Chester, PA) was used to

162 create a through-and-through hole in the right and left ear pinna. Ear tissue was collected at

163 specified time points using an 8mm biopsy punch (Sklar Instruments, West Chester, PA)

164 circumscribing the original injury. The published protocol can be found here:

165 dx.doi.org/10.17504/protocols.io.bp2l6dx65vqe/v1.

167 **Pulse-chase experiments**. We injected Ethynyl deoxyuridine (EdU) (10 ug/g) and

168 Bromodeoxyuridine (BrdU) (1 ug/g) diluted in normal saline, for pulse-chase experiments to

169 label and track cycling cells (the published protocol can be found here:

170 dx.doi.org/10.17504/protocols.io.261gerdmol47/v1). Both solutions were injected

171 intraperitoneally with a difference of five days between the EdU and BrdU injections. EdU and

172 BrdU injections were given as three separate injections, one every three hours over a period of

173 nine hours. The left ear pinna was collected after the EdU pulse to assess baseline cell numbers

in S-phase, and the right pinna was collected after the BrdU pulse to track single (EdU+ or

BrdU+) and double positive cells. The injection scheme was as follows: the first group of

animals was injected with EdU at D0 and BrdU at D5, the next group was injected with EdU at

D5 and BrdU at D10 and so on until D30. Hence, tissue samples were collected at 0, 5, 10, 15,

20, 25, and 30-days post injury.

Our experimental scheme allowed us to distinguish cells that (i) were in S-phase at the

time of the EdU injections and stopped proliferating within next five days (Edu+ alone); (ii) were

in S-phase at the time of the EdU injections, divided multiple times over five days diluting the

EdU, and remained in the cell cycle acquiring BrdU (weak EdU+/ strong BrdU+) - because each

cell division dilutes the EdU by half, a scenario where EdU is not detectable is possible, and

these cells would contribute to the BrdU+ alone counts); (iii) cells not in S-phase during the EdU

pulse but were cycling during the BrdU injections (BrdU+ alone); (iv) cells that acquired EdU,

divided only a few times, and remained in the cell cycle during the BrdU chase (strong EdU+/

strong Brdu+).

Histology, EdU detection and Immunohistochemistry. Harvested tissue was fixed

overnight in 10% neutral buffered formalin (NBF) (Thermo Fisher Scientific Inc. Epredia

5701) at 4˚C on a shaker. After 16-20 hrs the tissue was washed 3x with PBS, 3 minutes

per wash, followed by 3x 70% ethanol washes for dehydration and storage at 4˚C. Stored

tissue was processed for paraffin (Leica Biosystems, Buffalo Grove, IL, Formula "R"

REF: 3801450) embedding using a Histo5 tissue processor machine (Milestone). Samples

were sectioned at 5μm. EdU was detected using click chemistry as follows: Tissue

sections were de-paraffinized and rehydrated. Antigen retrieval was performed if co-

stained with antibodies. Tissues were washed two times for 10 minutes in Tris-buffered

saline (TBS) to incubate in the prepared reaction cocktail (see table below) for 30

minutes at RT protected from light. Tissues were washed two times in TBS for 5 minutes

and if co-stained with antibodies were blocked following the fluorescent IHC protocol for

BrdU and cell type specific antibodies tissue sections were deparaffinized and re-

hydrated. Tissues were then exposed to high heat antigen retrieval in a steamer in antigen

unmasking TRIS or citric acid-based solutions (Vector Laboratories, Inc.). After a 5-

minute wash, tissues were blocked in blocking serum (see table below) for 30 minutes at

RT. Tissues were then incubated with primary antibody diluted in blocking serum at 4 ˚C

overnight. After a 5 minutes wash, tissues were incubated in secondary antibody diluted

in blocking serum for 30 min at RT. Tissue samples were incubated with Hoechst 33342

for 5 minutes at RT and then dried and mounted in Prolong. Finally, protein similarity

was compared to assess antibody binding efficacy (Supplementary table S1). For

complete protocol details regarding paraffin sections

(dx.doi.org/10.17504/protocols.io.6qpvr9353vmk/v1), immunohistochemistry

(dx.doi.org/10.17504/protocols.io.n92ldrmd8g5b/v1), and EdU detection on sections

(dx.doi.org/10.17504/protocols.io.e6nvwbdrzvmk/v1).

EdU reaction cocktail	
Components	Volume (ul)
2 M Tris, pH 8.5	5
50 mM CuSO4	2
0.5 M ascorbic acid	20
Alexa fluor azide (0.25 mg/ml)	0.2
ddH2O	73
Total	100

Primary Antibodies					
Antibody	Company, Cat#	RRID	Host species	Dilution	Antigen retrieval
Anti-BrdU (Bromodeoxyuridine)	Abcam, Ab6326	AB_305426	Rat	1:500	Citric Acid
Anti-wide spectrum Cytokeratin	Abcam, Ab9377	AB_307222	Rabbit	1:500	None
Anti-ionized calcium-binding adapter molecule 1 (IBA1)	Novus Biologicals, NB100-10288s	AB_521594	Goat	1:400	Tris
Anti-retinoic acid-binding protein 1 (CRABP1)	Cell Signaling Technology, Inc, 13163	AB_2750569	Rabbit	1:500	Tris
Anti- alpha-Smooth Muscle Actin (alpha-SMA)	Novus Biologicals, NB300-978	AB_2273628	Goat	1:200	Tris

Anti-SRY Box Transcription Factor 9 (SOX9)	Abcam, ab185230	AB_2715497	Rabbit	1:1000	None
Anti-macrophage mannose receptor (MMR) (CD206)	R&D Systems, AF2535	AB_2063012	Goat	1:500	Tris

Secondary Antibodies				
Antibody	Company, Cat#	RRID	Host species	Dilution
Alexafluor 594 anti-Rabbit	Thermo Fisher Scientific, A21207	AB_141637	Donkey	1:400
Alexafluor 647 anti-goat	Thermo Fisher Scientific, A21447	AB_2535864	Donkey	1:400
Alexafluor 647 anti-rat	Thermo Fisher Scientific, A21247	AB_141778	Goat	1:400

Western Blotting. Protein extraction and quantification was performed as indicated

(dx.doi.org/10.17504/protocols.io.rm7vzkx12vx1/v1). Western blotting was performed as

indicated (dx.doi.org/10.17504/protocols.io.kqdg3qxz1v25/v1).

Imaging and Cell Quantification. Fluorescent images were collected using Cells Sens from

Olympus in Olympus inverted microscope (Olympus America Inc). Whole tissue sections were

captured in 20X and stitched automatically. Cell counting was manually performed in defined

areas of the images separately in each channel using ImageJ. Percentages of positive cells were

calculated by quantifying nuclei (Hoechst 33342) using the threshold tool in ImageJ.

Statistics. Statistical analyses were performed using JMP (version Pro 15.0, SAS Institute Inc)

and SPSS (version 28.0, IBM) where alpha was set at 0.05 for all statistical analyses. Graphs

were made in excel and JMP. To analyze our pulse chase data across species, we performed

Two-way ANOVAs with either EdU+, BrdU+ or double positive cells (dependent variable), and

days post injury (independent variable) followed by Tukey HSD multiple comparisons to test for

differences in positive cells as a function of time (Supplementary table S2).

To analyze proliferation during regeneration in *Acomys*, we used a linear mixed model to

assess differences in percentage of positive cells between three positions (A, B, C) over time

(D5-D30) (Supplementary table S3). Likelihood ratio testing and Akaike Information Criterion
(AIC) were used to select an appropriate covariance structure. A Kenward-Roger adjustment was
used to correct for negative bias in the 170 standard errors and degrees of freedom calculations
induced by the small sample size. Relevant pairwise comparisons were performed and controlled
for multiple comparisons using Fisher's Least Significant Differences (LSD). Mixed modeling
analysis was performed using SAS 9.4 (SAS Institute, Inc.; Cary, NC, USA).
242
Single-cell analysis. scRNA-seq data used in this study were download and available through
the Gene Expression Omnibus (GEO) (RRID:SCR_005012) under the accession number
GSE182141. The raw fastqs were processed through the CellRanger pipeline v8.0.1
(RRID:SCR_017344) using 10X Genomics' mm10-2020-A reference genome or a custom
annotated *Acomys dimidiatus* genome (GCA_907164435.1 (mAcoDim1_REL_1905), with the
'include introns' set to true to obtain gene expression matrices.
Background reads were removed using decontX from the celda package using the filtered
barcodes as the cells to keep and the remaining cells in the raw barcode matrix as the background
(Campbell et al., 2024). Doublets were minimized using DoubletFinder (RRID:SCR_018771)
(McGinnis et al., 2019). The R v4.4 (RRID:SCR_001905) library Seurat v5.1
(RRID:SCR_016341) (Hao et al., 2021; Hao et al., 2024)) was used for cell type clustering and
marker gene identification. Cells expressing >500 genes were retained for downstream analysis.
Each sample was normalized SCTransform v2 (RRID:SCR_022146), using the glmGamPoi
method and the number of RNA molecules per cell were regressed out. Samples were integrated
with common anchor genes using the CCA method to minimize sample to sample variation. Cell
clusters were determined by the Louvain algorithm using a resolution of 0.3. UMAP dimension
reduction was done using the first thirty principal components. Marker genes for each cell type
were calculated using the Wilcoxon Rank Sum test returning only genes that are present in a
minimum of 25% of the analyzed cluster. Cross species markers were done as above in cell types
of interest, but the pct.2 score was filtered to be > 0 to eliminate uniquely annotated genes from
dominating the lists. Volcano plots were created with the Enhanced Volcano (Blighe et al., 2024)
R library using the avg_log2FC and p_val_adj values as inputs.

Gene ontology (GO) analyses were performed in the Enrichr platform

(https://maayanlab.cloud/Enrichr/). Only differentially expressed genes with a two-fold change

were included in the analysis of GO Biological Process.

RESULTS

Vertebrate stromal cells rapidly re-enter the cell cycle in response to injury

Notably few regeneration papers using salamanders or fish have indicated whether

stromal cells are proliferating during normal homeostasis. To explore proliferation in uninjured

tissue, we used EdU to label cells in S-phase in two highly regenerative amniotes (*Danio rerio* -

zebrafish, *Ambystoma mexicanum* - axolotl), in an adult mammal with enhanced regenerative

ability (*Acomys dimidiatus* – spiny mouse) and in a poorly regenerating mammal (*Mus musculus*

– outbred ND4 laboratory mouse strain) (Fig. 1A). Examining the ear pinna (*Acomys* and *Mus*),

limb (*Ambystoma*) and caudal fin (*Danio*), we found cycling cells in the epidermis from all four

species prior to injury; a finding that was expected given the presence of slow cycling stem cells

in the epidermis (Fig. 1B). In uninjured *Danio* and *Ambystoma* tissue, we also found a small

percentage of cycling cells in the stroma (1% and 0.5% respectively) supporting previous reports

for these species (Fig. 1B-C) (K. Johnson et al., 2018; Simkin & Seifert, 2018). In contrast, we

found only one or two cycling stromal cells across entire tissue sections in *Acomys* or *Mus* prior

to tissue removal (Fig. 1B-C). These data support that stromal cell proliferation is essentially

non-existent prior to injury in mammals, at least in the ear and skin.

A longstanding hypothesis is that cells in mammals are refractory to cell cycle re-entry

and progression whereas cells in highly regenerative vertebrates easily re-enter the cell cycle and

proliferate to replace missing tissue (Pajcini et al., 2010; Simkin & Seifert, 2018; Tanaka et al.,

1997; Tanaka & Brockes, 1998). To test this hypothesis, we used a pulse chase strategy whereby

we injected animals with EdU, performed the specified injury, injected BrdU at either 15 or

63hrs post injury (hpi) and then harvested tissue 9hrs later at 24 and 72 hpi (see Methods). We

quantified stromal cells that were EdU+, or BrdU+, or EdU+/BrdU+ (double positive) (Fig. 1B,

D). 24 hpi spiny mice and zebrafish showed the highest percentage of newly cycling cells in

response to injury (BrdU+). *Mus* cells also re-entered the cell cycle, albeit later at 72 hpi. Axolotl

limb tissue, on the other hand, exhibited very few singly BrdU+ cells but presented the highest

number of double positive cells (Fig. 1D). These data demonstrate that stromal cells in *Acomys*,

Mus and zebrafish re-enter the cell cycle *de novo* in response to injury with a 48hr lag observed

for *Mus* cells.

299   To assess cell activation arising from the injury but independent of the local repair

response (i.e., systemic cell activation), we collected the contralateral ear pinna in *Mus* and

Acomys, the contralateral limb in axolotls, and the pelvic fin in zebrafish 24 hpi and 72 hpi after

tissue amputation. Using a pulse chase strategy as described above, we looked for cell cycle re-

entry in uninjured tissues at 24 hpi and 72 hpi. EdU+ cells were cycling prior to injury, BrdU+

cells were those activated systemically and EdU+/BrdU+ were those cycling prior to injury that

were activated systemically. Despite reports in lab mice of long-range cell activation in skeletal

muscle stem cells (Rodgers et al., 2014), in *Acomys* and *Mus*, we did not find a noticeable

response in the contralateral ear pinna where the number of BrdU+ cells did not exceed 0.5% at

either time point (Fig. S1A-B). Similarly, in axolotl and zebrafish, the number of EdU+ cells at

24 and 72 hpi was similar to the number of positive cells found in uninjured tissue at D0 (see

above) (Fig. S1A-B). These data indicate that a systemic response to injury is negligible at 24

and 72hpi in all four species.

Proliferating cells are maintained in a distal position relative to the wound site over time

314   *Acomys* species regenerate ear pinna tissue after removal with a 4 mm (or larger) biopsy

punch whereas identical tissue excisions in wild or laboratory *Mus* result in scar tissue formation

(Seifert et al., 2012). To more thoroughly understand the kinetics of proliferating cells during

regeneration and fibrotic repair, we again employed a pulse (EdU) chase (BrdU) scheme where

our pulse and chase were separated by a five-day window (see Methods). First, we analyzed

BrdU+ cells to estimate the proliferative rate from injury until D30 and then analyzed the spatial

distribution of cycling cells by quantifying cells adjacent to the injury (aka proximal) and at two

distal locations (i.e., closest to the new epidermis); distal dorsal and distal ventral (Fig. 2A-E).

We quantified healing tissue area from D10-30 which supported continued addition of new tissue

during regeneration (Fig. 2B). In contrast, after a proliferative burst, no new fibrotic tissue was

added in *Mus* ears after D10 (Fig. 2C).

325   Analyzing BrdU+ cells during regeneration revealed that from D15 to D30, the number

of cycling cells was significantly higher in the distal areas compared to the proximal areas

(injury site) while there was no difference between the distal-dorsal and distal-ventral areas (Fig.

2D and Supplementary Table 1). An exception was D20 when the new tissue areas were small or

flat and thus distal and proximal areas were close (Fig. 2D). While cell proliferation in the

proximal compartment decreased over time, proliferation in the distal compartment remained

>10% from D10-30 contributing to the steady production of new tissue. (Fig. 2D). Quantifying

BrdU+ cells in *Mus* ears, we observed a narrow extension of new tissue (Fig. 2A). After a burst

of DNA replication (>25%) in *Mus* at D5, the percent of BrdU+ cells precipitously declined from

D10-30 coinciding with no increase in tissue area over this time period (Fig. 2C, E).

To assess the spatial distribution of cycling cells during regeneration (D10-30), we first

analyzed the epidermis where stem cell localization and proliferation are well known. Basal stem

cells in the deeper layer of the epidermis show symmetrical and asymmetrical cell division

depending on whether they produce progeny or more stem cells (Blanpain & Fuchs, 2006).

Analyzing our pulse-chase data, we observed stem, progenitor, and differentiating cells in the

epidermis (Fig. S2A-B). The stratum basale contained slow-cycling stem cells (strong EdU+)

and infrequently dividing stem cells (EdU+/BrdU+) while also indicating rapidly-cycling

progenitor cells (strong BrdU+/diluted EdU+), and recently-cycling progenitor cells (strong

BrdU+ Fig. S2A-B). In addition, we observed progenitor cells having acquired EdU that

migrated apically withdrawing from the cell cycle and undergoing differentiation as per the

absence of BrdU (Fig. S2B, yellow arrow). These cell behaviors demonstrated that our labeling

scheme could effectively capture predicted stem cell behavior in a tissue with known

proliferative properties.

Next we characterized proliferative behavior in the stromal compartment by quantifying

cells that re-entered the cell cycle after injury but arrested within five days (EdU+ only) possibly

to differentiate or re-enter the cell cycle later (slow-cycling progenitors), and cells that re-entered

the cell cycle but maintained their proliferative state over the same period (EdU+/BrdU+ and

BrdU+ only) in both distal and proximal areas relative to the wound site (Fig. 2F-I). We

hypothesized that cell proliferation could either be clonal, with multiple daughter cells near

proliferative progenitors in rosettes or could expand from the wound site. In the latter case, we

predicted to capture progenitors cycling near the wound margin and daughter cells building up

the blastema distally. At D10, we found a stochastic pattern of proliferation with a mixture of

EdU+ cells, BrdU+ cells, and double-positive cells in distal and proximal compartments, where

the EdU+ cell population represented the biggest population (>25%) (Fig. 2F-G). At D25,

however, we were able to observe a pattern where rapid-cycling (BrdU+) and infrequently-

dividing (double positive) cells represented half of the proliferative population (>9%) in the

distal compartment with the other half (~8%) being represented by potentially slow cycling

progenitors (EdU+) (Fig. 2H-I). In comparison, the proximal area was represented mostly by

EdU+ cells (5%) that may have started differentiation and a very small percentage of rapid-

cycling and infrequently-dividing (1%) cells that could support expansion of some differentiating

cell types (Fig. 2H-I). Supporting this pattern, we observed differentiation of new hair follicles

and cartilage near the wound site. *In toto*, these results show that the proliferative progenitor

population is maintained in a distal position relative to the wound site. Furthermore, our pulse

chase labeling supports cell cycle progression and asynchronous cell proliferation during

regeneration in *Acomys*, whereas thymidine analog incorporation without cell division was

observed in *Mus*, suggesting an S-phase arrest after D10.

The proliferative population comprises diverse cell types in Acomys

To determine cell state or identity among the proliferative population (PP), we co-labeled

with cell type specific markers and EdU (see Methods). Although we found no mesenchymal cell

types proliferating in uninjured tissue, we characterized the distribution of macrophages

(IBA1+), chondroblasts (SOX9+) and two fibroblast subtypes (CRABP1+ or αSMA+) in

comparison with *Mus* to identify any obvious differences in presence and location across tissue

compartments (Fig. S3A-J). First, we found relatively few CD206+ and IBA1+ macrophages

(Fig. S3A, B, F, G) nor αSMA+ cells which were sparse (Fig. S3E, J). We observed two

populations of SOX9+ cells in both species: one associated with the perichondrium, and another

associated with the sebaceous gland and hair follicle (Fig. S3C, H). Importantly, we observed a

population of CRABP1+ fibroblasts scattered throughout the papillary dermis in *Acomy*s while

these cells were not detectable in uninjured *Mus* skin (Fig. S3D, I).

Examining these same cell types at D5, we observed cell-cycle re-entry among many of

them (Fig. 3A-J). By counting all the EdU+ cells in a counting frame and then quantifying

double positive cells we estimated the fraction of cycling cells represented by each cell type.

Because inflammation peaks at D5 (Gawriluk et al., 2020), we hypothesized that a large fraction

of proliferating cells would be macrophages. Instead, IBA1+ and CD206+ cells represented only

5.3% and 4.95% of the PP respectively in *Acomys* and 5.3% and 0.96% in *Mus* (Fig 3A-D). In

Acomys, the two fibroblast subtypes (CRABP1+ and αSMA+ cells) represented ~14% and

18.65% respectively while chondroblasts (SOX9+ cells) represented ~16% of the PP in *Acomys*

(Fig. 3E, G, H). Surprisingly, although totally absent prior to injury in *Mus*, an emergent

CRABP1+ population represented 28.9% of the PP at D5 (Fig. 3I). SOX9+ and αSMA+ cells

represented 7.96%, and 33% of the PP in *Mus* (Fig. 3F, J). These data support that various cell

types re-enter the cell cycle during regeneration *and* fibrotic repair.

396   Only *Acomys* maintains a proliferative population during regeneration whereas activated

cells did not appear to maintain cell cycle progression or appreciably divide during fibrotic

repair. Co-staining with EdU, we confirmed the above-mentioned cell types remained part of the

PP at D20 in *Acomys* (Fig. 4A-C and Fig. S4A-B). Of note, we observed that CRABP1+,

αSMA+, and SOX9+ cells were segregated spatially throughout the blastema (Fig. 4A-C, G, H

and Fig. S5A). While the CRABP1+ population was restricted to the papillary dermis at D0, this

population expanded to deeper dermal areas at D5 (Fig. 3D, G and Fig. S5A) and occupied the

dorsal blastema with another, smaller population in the ventral papillary zone at D20 (Fig. 4A-

A', G-H and Fig. S5A). In contrast, the αSMA+ fibroblast population appeared after injury in a

distal and central position where reticular fibroblasts localize (Fig. 3H-H' and Figs. S3E, S5A).

This population shrunk at D20 and was largely non-overlapping with CRABP1+ cells (Fig. 4B-

B', G-H and Fig. S5A). Thus, as the blastema expanded, the αSMA+ fibroblasts maintained a

small, distal population that disappeared in proximal tissue, while the CRABP1+ population

expanded over time (Fig. 4G-H). In response to injury, the SOX9+ population that labeled the

perichondrium appeared in the blastema perfectly aligned with the cartilage (Fig. 4C and Fig.

S5A). IBA1+ and CD206+ cells were spread evenly across the blastema (Fig. S4A-B and Fig.

S5A).

413   The configuration of these populations was notably different in *Mus* during healing (Fig.

4D-F and Fig. S5B). Absent at D0, a CRABP1+ population emerged throughout the injury zone

at D5, only to become restricted to the distal tip of the ear at D20 with distinctly fainter antibody

reactivity (Figs. 3I & 4D and Fig. S5B). αSMA+ cells, absent in *Mus* at D0, became the

dominant population in distal scar tissue (Fig. 4E and Fig. S5B). We observed some Sox9+ cells

in the middle of the outgrowth (Fig. 4F and Fig. S5B), and IBA1+ and CD206+ cells distributed

widely throughout the scar tissue (Fig. S4C, D and Fig. S5B). Western blotting for CRABP1

supported our observations using immunohistochemistry (Fig. S5C). Together, these results

uncovered a distinct spatial segregation of cell types with a prominent CRABP1+ population

present in *Acomys* before and after injury, that was merely transient in *Mus*.

Species-specific cell states separate scaring and regeneration phenotypes

To extend our cellular studies, we next interrogated a scRNA-seq dataset harvested from

uninjured (D0) and healing ear tissue at D3, 5, 10 and 15 from *Acomys* and *Mus* (Simkin et al.,

2024). Unbiased clustering of batch corrected data for all time points (see Methods) revealed the

same cell populations across species (Fig. S7). Thus, we extracted the four main stromal cell

populations, fibroblasts (cluster 18), chondroblasts (cluster 19), lymphatic endothelial cells

(cluster 16) and vascular endothelial cells (cluster 9), and generated a new UMAP (Fig. 5A). Re-

clustering revealed eight subpopulations among the four original stromal populations: two

populations each of vascular and lymphatic endothelial cells, three fibroblast subtypes, and a

single chondroblast population (Fig. 5B). To investigate markers from our immunohistochemical

analysis, we plotted *Crabp1*, *Acta2* (αSMA), and *Sox9* across cells on this UMAP (Fig. 5B).

Interestingly, we found that *Acta2* transcripts were strongly associated with one fibroblast

subtype (cluster 7) while *Crabp1* was totally excluded from this cluster (Fig. 5B-D). *Sox9*

marked chondroblasts in both species (Fig. 5E). To assess co-localization, we plotted *CRABP1*

and *Acta2* together and found few double positive cells in *Acomys* (Fig. S8A). In contrast,

fibroblast cluster six was mostly double-positive for *Crabp1* and *Acta2* in *Mus* (Fig. S8B). In

addition, fibroblast cluster four and lymphatic endothelial cluster five were heavily enriched for

Acta2 in *Mus* (Fig. S8B). These data support a distinct *Acta2*+ fibroblast population in both

species (cluster 7) and a *Crabp1*+ fibroblast subtype in *Acomys* that is *Acta2*-. In *Mus*, *Crabp1*

appears to only mark a transitional cell state that ultimately expresses *Acta2* while *Acta2*

generally increases among fibroblasts and lymphatic endothelial cells supporting a fibrotic

phenotype in *Mus*.

To better understand intrinsic differences between the fibroblast subpopulations, we ran

differential gene expression (DGE) analysis comparing the three *Acomys* fibroblast clusters

against the three *Mus* clusters (Fig. 5F-K). This analysis revealed 1497 genes (488 in *Acomys*,

1009 in *Mus*) differentially upregulated in cluster four, 1228 (510 in *Acomys*, 718 in *Mus*) in

cluster six, and 743 (248 in *Acomys*, 495 in *Mus*) in cluster seven among both species (Fig 5F, H,

J). Gene ontology of each cluster separated by species revealed a clear segregation in the

fibroblast response to injury (Supplementary Table S4). The three fibroblast populations in *Mus*

were associated with fibrotic pathways: response to TGFß stimulus, cellular response to hypoxia,

and negative regulation of epithelial cell proliferation, among others (Fig 5G, I, K). In contrast,

Acomys subpopulations were associated with telomere maintenance and cell proliferation

pathways (Fig 5G, I, K). Interestingly, *Acta2*-expressing cluster seven was associated with

regulation of fat cell differentiation, positive regulation of translation and regulation of

membrane protein ectodomain proteolysis (Fig 5K). Chondroblasts (cluster two) followed a

similar pattern with 2234 genes (786 in *Acomys*, 1448 in *Mus*) differentially upregulated. *Acomys*

chondroblasts upregulated genes associated with translation, response to hydrogen peroxide and

gene expression, while *Mus* chondroblasts upregulated genes associated with TGFß stimulus and

regulation of TGFß-Receptor signaling (Fig. S9B). Of note, all four clusters (two, four, six and

seven) in *Acomys* highly upregulated *Zc3h10* and *Tssc4* (Fig. 5F, H, J and fig. S9A). The first

gene has been shown to boost mitochondrial function (Audano et al., 2018), while the second

gene promotes cell survival and proliferation in breast cancer tumors (Chen et al., 2022).

Together, these data show that all four *Mus* subpopulations activate fibrosis-related

pathways, while the four subpopulations in *Acomys* are biased toward pathways associated with

cell proliferation and growth. Importantly, while the same markers may be able to identify

fibroblast subtypes in both species, the intrinsic ability of each subtype to respond to injury

differs.

CRABP1+ fibroblasts do not induce regeneration in neonatal Mus.

Previous studies using very large wounds in lab mice observed hair follicle regeneration

in the wound center, an observation that has been attributed to the presence of *Crabp1+*

fibroblasts (the so-called WIHN model) (Jiang & Rinkevich, 2021; Phan et al., 2021b). Since we

did not observe CRABP1+ cells in uninjured *Mus* tissue, but did observe them in *Acomys* ear

dermis prior to injury and during regeneration, we hypothesized these cells disappear during *Mus*

ear development. To test this hypothesis, we assessed CRABP1 reactivity in embryonic,

neonatal, juvenile and adult ear tissue (Fig. 6A). In *Acomys,* we found CRABP1+ reactivity

across all stages (embryonic and neonatal) and into adulthood (8 months old) (Fig. 6A).

CRABP1 reactivity was strongest at embryonic stages and restricted to the dermis where it

persisted into adulthood; mainly the papillary dermis (Fig. 6A). In contrast, we found *Mus*

fibroblasts started to lose CRABP1 expression around P20 and it was totally absent from the ear

pinna in three month old adults (Fig. 6A).

Based on these observations, we hypothesized that challenging *Mus* ear pinnae using our

regeneration assay in the presence of CRABP1+ fibroblasts might enhance ear hole regeneration.

Performing a 4 mm ear punch in P20 animals, we found neonatal ear holes produced tissue until

D15 (Fig. 6B-C), but did not close. Similarly, we performed 2 mm punches in P23 animals and

did not observe regeneration (Fig. S10A-B). Next, we injured P15 *Mus* which have CRABP1+

fibroblasts and collected tissue at D5 and D20. CRABP1 reactivity after injury was similar to

that observed in adult *Mus* with increased expression at D5 which disappeared by D20 (Fig 6D).

Again, we did not observe regeneration. Together, these data show the presence of a CRABP1+

fibroblast population is not sufficient to promote a regenerative response and instead supports

species-intrinsic adoption of cells states that support regeneration.

DISCUSSION

Here we conducted a comprehensive analysis of cell proliferation during regeneration

and found cycling cells maintained a distal position relative to the original injury, while proximal

cells exited the cell cycle and differentiated. These data revealed controlled regulation of the cell

cycle in response to injury such that while amputation served as an activation trigger in both

species, regenerating tissue maintained cell cycle progression while cells participating in fibrotic

repair arrested. Importantly, proliferation in *Acomys* did not lead to tissue overgrowth

demonstrating regulated shutdown of cell cycle progression. This unites *Acomys* with other

highly regenerative vertebrates where tightly controlled cell proliferation is a hallmark of

epimorphic regeneration in zebrafish caudal fins (Nechiporuk & Keating, 2002; Poleo et al.,

2001) and newt limbs (Hays & Fischman, 1961; Wallace & Maden, 1976). Cell proliferation in

injured mammalian organ stroma, apart from dedicated stem cell populations, is rarely observed

except during tumorigenesis where cell proliferation is uncontrolled (Hanahan & Weinberg,

2011).

Although evolutionary explanations for the presence of regenerative ability among some

vertebrate clades remains poorly supported, the presence of indeterminate growth has been

offered as an underlying prerequisite (Froehlich et al., 2013; Sebens, 1987). This hypothesis

claims that indeterminate growers maintain populations of somatic stem or progenitor cells that

behave like slow-cycling stem cells to support growth. There are two arms to this hypothesis.

First is the presence of cycling somatic cells during homeostasis lowers the barrier to cell cycle

re-entry upon injury. Second is that once cycling, cells do not activate tumor suppressor

machinery during S-phase progression to initiate cell cycle arrest. These ideas partly stem from

culture experiments with newt myotubes that easily re-enter the cell cycle after serum

stimulation in contrast to mouse myotubes formed from C2C12 cells which are refractory to cell

cycle re-entry (Tanaka et al., 1997; Tanaka & Brockes, 1998). Although *Mus* myotubes can be

forced to re-enter the cell cycle, this requires deletion or inactivation of tumor suppressors

retinoblastoma protein (pRB) and p16^{ink4a} (Gu et al., 1993; Schneider et al., 1994).

To examine if cycling somatic cells exist in highly regenerative vertebrates, we analyzed

cell cycle re-entry and progression prior to and after injury between determinate (*Acomys*, *Mus*)

and indeterminate growers (*Ambystoma* and *Danio*). In line with previous observations (K.

Johnson et al., 2018; Simkin & Seifert, 2018), we found a small population of slow-cycling

stromal cells present during homeostasis in indeterminate growers while we failed to detect these

cells in determinate growers. This cycling population, however, scarcely contributed to blastema

formation in zebrafish. Instead, cells that re-entered the cell cycle in response to injury were the

major contributors to regeneration. This result supports a previous study in zebrafish where slow-

cycling cells marked for eight weeks before injury did not contribute to blastema formation

(Nechiporuk & Keating, 2002). Importantly, our data show that cells in *Acomys* and *Mus* re-enter

the cell cycle in response to injury independent of their regenerative ability. An open question is

how cell cycle progression is controlled for almost forty days and then downregulated so as not

to create tissue overgrowth. Considered *in toto*, our cross-species analysis rejects the notion that

a determinate growth mode prevents cell cycle re-entry and supports that the contribution of

slow-cycling cells to the blastema is minimal, at least in fish.

Our cross-species proliferation experiments also assessed whether cells on the other side

of the animal were activated by injury (so-called systemic cell activation). Work in lab mice

found that acute muscle injury elicits a systemic response in which contralateral-located stem

cells (CSCs) re-enter the cell cycle and access a priming state named G_{Alert} (Rodgers et al.,

2014). In contrast to quiescent stem cells (QSCs), CSCs in mouse show an increased propensity

to cycle, although their numbers remain low in comparison with stem cells located at the injury

site (Rodgers et al., 2014). A similar systemic response was observed in axolotl limbs

contralateral to amputated limbs (Johnson et al., 2018). Using contralateral control tissue

collected in our experiments, we looked for primed cells *in vivo* following injury. In contrast to

muscle stem cells in *Mus*, we found a negligible stromal response in *Acomys* and *Mus* with less

than 0.5% of contralateral ear pinna cells responding to a systemic signal. While we used

immunofluorescence to quantify active or primed mesenchymal cells in tissue sections, previous

studies in mouse muscle used FACS sorted satellite stem cells to assess *in vivo* BrdU

incorporation. Supporting our observations in *Acomys* and *Mus*, we could not find a significant

response in contralateral axolotl limbs or zebrafish pelvic fins, at least in the first 72hrs after

injury. While these results are at odds with a single axolotl study, those data included epidermal

cells in their cell counts and later time points (Johnson et al., 2018). Thus, our data do not

support significant stromal cell cycle re-entry in uninjured contralateral tissue from the four

species analyzed.

Contemporary vertebrate regeneration studies using transgenics have concluded that

blastemas particular to an injured tissue are formed from a heterogeneous population of cells at

the amputation plane (rev. in Seifert & Muneoka, 2018). Although various stem cell reservoirs

have been discovered in different mammalian tissues, except for the digit tip, exploring cell type

contributions during complex tissue regeneration in mammals has been impossible. While we

could not identify stem cells with our methods, our data support that *Acomys* expands pre-

existing cell types to participate in regeneration via cell proliferation. One exception in our

analysis were αSMA+ fibroblasts which appeared after injury but did not maintain αSMA

identity in newly differentiated tissue. Interestingly, our scRNA-seq data indicated that an *Acta2*-

expressing population expressed genes related to the regulation of fat cell differentiation

suggesting that some myofibroblasts may differentiate into adipocytes in a regenerative context.

Similarly, αSMA+ myofibroblasts were shown to differentiate into adipocytes when exposed to

bone morphogenetic protein (BMP) triggered by neogenic hair follicles in large skin wounds

(Plikus et al., 2017). Further supporting our observations, recent data from another *Acomys* ear

pinna regeneration study reported that αSMA fibroblasts disappear after wound closure with

only ACTA2+ vascular smooth muscle cells remaining (Brewer et al., 2021). We too observed

high expression of TGFβ pathway components in *Mus*, but not *Acomys* fibroblasts. Taken with

the previously observed insensitivity of *Acomys* fibroblasts to TGFβ induced myofibroblast

differentiation (Brewer et al., 2021), *Acomys* appears to support reduced myofibroblast

activation. While these data suggest a possible differentiation trajectory for myofibroblasts, the

exact dynamics of αSMA+ cells await further study in a regenerative context.

In contrast to their role during regeneration, the role of myofibroblasts during fibrotic

skin repair is well known (rev. in Plikus et al., 2021). Previous studies in skin have described

first wound responders as a large proportion of reticular fibroblasts including αSMA+ fibroblasts

(Driskell et al., 2013) which rapidly seal the wound area promoting scaring and inhibiting hair

follicle formation (Correa-Gallegos et al., 2019). These data support our observations in *Mus*

where αSMA+ fibroblasts are the main cell population in newly generated scar tissue. Together,

these data show that αSMA fibroblasts are dominant in scaring but seem to be transient in

regeneration.

CRABP1+ papillary fibroblasts abundantly participate in the regeneration of neonatal

skin and deer antler velvet and thus we interrogated their behavior in our ear pinna assay (Phan

et al., 2021b; Sinha et al., 2022). In contrast to the ephemeral nature of the αSMA+ population in

Acomys, we observed a stark difference in CRABP1+ cells during regeneration and fibrotic

repair. While we were unable to detect CRABP1+ cells prior to injury in *Mus*, these cells

appeared at D5 as a subset of cycling cells only to disappear by D20. In *Acomys*, however, we

detected CRABP1+ cells prior to injury and this population expanded to participate in blastema

formation. CRABP1 has been found in a variety of regeneration contexts. For instance, this

protein is expressed in *Mus* papillary fibroblasts that expand in large, somewhat regenerative

wounds (Guerrero-Juarez et al., 2019). Other examples are the regenerating wounds in newborn

(P2) mice that have a higher proportion of CRABP1+ fibroblasts (Phan et al., 2021a) and the

velvet skin on deer antlers which have regeneration-primed fibroblasts expressing genes

putatively associated with a regenerative phenotype (e.g., *CRABP1, RUNX1,* and *PRSS35*)

(Sinha et al., 2022).

Rather than the presence or lack of a specific cell type, our scRNA-seq data suggests that

it may also be one of cell state. Although all four stromal subclusters express some of the same

markers in *Acomys* and *Mus*, our DGE and GO analyses revealed that the four *Mus* subclusters

express biological processes related to fibrosis. In contrast, all four subclusters in *Acomys* seem

to use biological processes related with cell proliferation; high transcription and telomere

maintenance and genes that boost mitochondrial function and cell survival. Specifically, a

subpopulation in both species that highly expressed *Acta2* exhibited divergent transcriptomic

responses. Future studies are necessary to address whether *Acomys* αSMA+ cells transiently

become myofibroblast-like only to later differentiate into adipocytes as suggested by our GO

analysis. Our finding that two subpopulations expressed high telomerase and endoplasmic

reticulum stress (ERS) related activity supported the persistence of proliferation in these cells.

Telomerase activity is low in most non-proliferating somatic cells in humans and mice, while

high activity has been reported in highly proliferating immortal and cancer cells (Kim et al.,

1994; Sha & Bacchetti, 1997). Moreover, inhibition of telomerase results in death of tumor cells

in vitro and *in vivo* showing the importance of this enzyme for tumor growth (Hahn et al., 1999).

Similarly, various ERS responses have been shown to promote cell proliferation and survival in

various types of cancer (Zhang et al., 2024). These data suggest that *Acomys* can access and use

pathways other adult mammals do not outside of tumorigenesis.

As a functional test of cell type versus cell state as a driver of regeneration, we assessed

the role of CRABP1+ fibroblasts after injury in neonatal mice before this population disappears.

Previous studies have shown that papillary CRABP1+ fibroblasts are enriched in regenerative

neonatal human and mouse skin but are lost early during postnatal development (Mine et al.,

2008; Sinha et al., 2022). Examining neonatal and early postnatal stages, we found CRABP1+

fibroblasts in the *Mus* ear pinna which persisted until approximately P20. Using our ear punch

assay when CRABP1+ fibroblasts are present in the *Mus* ear pinna, did not improve ear hole

closure or stimulate regeneration. These results support that the presence of a cell population

alone may be less important that the ability for cells to acquire a specific cell state needed to

facilitate regeneration. A recent examination of regeneration and scarring phenotypes across

eleven murid rodents, underscored the binary nature of a regenerative phenotype in that species

appear to either completely regenerate or not (i.e., no partial regeneration) (Riddell et al., 2025).

This supports that complex tissue regeneration requires multiple features across stereotypical

healing processes such as intrinsic cell cycle licensing (Gawriluk et al., 2016; Tomasso et al.,

2023), goldilocks regulation of reactive oxygen species (ROS) (Saxena et al., 2019) and a

regenerative skewed immune cell response (Gawriluk et al., 2020; Simkin et al., 2017, 2024).

Future work investigating specific cell states across more regenerative and non-regenerative

mammals will undoubtably shed light on a blueprint for complex tissue regeneration in
mammals.
638

ACKNOWLEDGEMENTS

We would like to thank Josh Sarli, Brennan Riddell and Ava Musarra for animal husbandry. We
thank all members of the Seifert lab for helpful discussions in developing the manuscript and
Brennan Riddle for helping to create the GO graphs. This research was funded in part by grants
from the NIH (R01AR070313) and Aligning Science Across Parkinson's (ASAP-020495) through
the Michael J. Fox Foundation for Parkinson's Research (MJFF) to AWS. For the purpose of open
access, the author has applied a CC BY public copyright license to all Author Accepted
Manuscripts arising from this submission.

AUTHOR CONTRIBUTIONS
EOR, RA, and AWS designed research. EOR, MA and RA performed research. EOR, MA and
AWS analyzed data. EOR and AWS wrote the paper, and all authors commented on and edited the
paper.
652

653 **COMPETING INTERESTS**
654 The authors declare no competing interests.
655

656 **AVAILABILITY STATEMENT**
657 The data, code, protocols, and key lab materials used and generated in this study are listed in a
658 Key Resource Table alongside their persistent identifiers (Supplementary Tables).
659

660 **REFERENCES**
661

662 Audano, M., Pedretti, S., Cermenati, G., Brioschi, E., Diaferia, G. R., Ghisletti, S., Cuomo, A.,
663   Bonaldi, T., Salerno, F., Mora, M., Grigore, L., Garlaschelli, K., Baragetti, A., Bonacina, F.,
664   Catapano, A. L., Norata, G. D., Crestani, M., Caruso, D., Saez, E., … Mitro, N. (2018).
665   Zc3h10 is a novel mitochondrial regulator. *EMBO Reports*, *19*(4).
666   https://doi.org/10.15252/embr.201745531

Billingham, R. E., Mangold, R., & Silvers, W. K. (1959). THE NEOGENESIS OF SKIN IN THE ANTLERS OF DEER*. *Annals of the New York Academy of Sciences*, *83*(3), 491–498. https://doi.org/10.1111/J.1749-6632.1960.TB40922.X

Blanpain, C., & Fuchs, E. (2006). *Epidermal Stem Cells of the Skin*.

Borgens, R. B. (1982). Mice Regrow the Tips of Their Foretoes. *Science*, *217*(4561), 747–750. https://doi.org/10.1126/SCIENCE.7100922

Brewer, C. M., Nelson, B. R., Wakenight, P., Collins, S. J., Okamura, D. M., Dong, X. R., Mahoney, W. M., McKenna, A., Shendure, J., Timms, A., Millen, K. J., & Majesky, M. W. (2021). Adaptations in Hippo-Yap signaling and myofibroblast fate underlie scar-free ear appendage wound healing in spiny mice. *Developmental Cell*, *56*(19), 2722–2740. https://doi.org/10.1016/j.devcel.2021.09.008

Carlson, B. M. (2005). Some principles of regeneration in mammalian systems. *Anatomical Record. Part B, New Anatomist*, *287*(1), 4–13. https://doi.org/10.1002/AR.B.20079

Chen, Y., Zhang, Z., Henson, E. S., Cuddihy, A., Haigh, K., Wang, R., Haigh, J. J., & Gibson, S. B. (2022). Autophagy inhibition by TSSC4 (tumor suppressing subtransferable candidate 4) contributes to sustainable cancer cell growth. *Autophagy*, *18*(6), 1274–1296. https://doi.org/10.1080/15548627.2021.1973338

Cheung, T. H., & Rando, T. A. (2013). Molecular regulation of stem cell quiescence. In *Nature Reviews Molecular Cell Biology* (Vol. 14, Issue 6, pp. 329–340). https://doi.org/10.1038/nrm3591

Correa-Gallegos, D., Jiang, D., Christ, S., Ramesh, P., Ye, H., Wannemacher, J., Kalgudde Gopal, S., Yu, Q., Aichler, M., Walch, A., Mirastschijski, U., Volz, T., & Rinkevich, Y. (2019). Patch repair of deep wounds by mobilized fascia. *Nature*, *576*(7786), 287–292. https://doi.org/10.1038/s41586-019-1794-y

Driskell, R. R., Lichtenberger, B. M., Hoste, E., Kretzschmar, K., Simons, B. D., Charalambous, M., Ferron, S. R., Herault, Y., Pavlovic, G., Ferguson-Smith, A. C., & Watt, F. M. (2013). Distinct fibroblast lineages determine dermal architecture in skin development and repair. *Nature*, *504*(7479), 277–281. https://doi.org/10.1038/nature12783

Echeverri, K., Clarke, J. D. W., & Tanaka, E. M. (2001). In vivo imaging indicates muscle fiber dedifferentiation is a major contributor to the regenerating tail blastema. *Developmental Biology*, *236*(1), 151–164. https://doi.org/10.1006/dbio.2001.0312

Froehlich, J. M., Fowler, Z. G., Galt, N. J., Smith, D. L., & Biga, P. R. (2013). Sarcopenia and piscines: The case for indeterminate-growing fish as unique genetic model organisms in aging and longevity research. *Frontiers in Genetics*, *4*(AUG). https://doi.org/10.3389/fgene.2013.00159

Gawriluk, T. R., Simkin, J., Hacker, C. K., Kimani, J. M., Kiama, S. G., Ezenwa, V. O., & Seifert, A. W. (2020). Complex Tissue Regeneration in Mammals Is Associated With Reduced

Inflammatory Cytokines and an Influx of T Cells. *Frontiers in Immunology*, *11*.
https://doi.org/10.3389/FIMMU.2020.01695/FULL

Gawriluk, T. R., Simkin, J., Thompson, K. L., Biswas, S. K., Clare-Salzler, Z., Kimani, J. M.,
Kiama, S. G., Smith, J. J., Ezenwa, V. O., & Seifert, A. W. (2016). Comparative analysis of
ear-hole closure identifies epimorphic regeneration as a discrete trait in mammals. *Nature*
Communications, *7*. https://doi.org/10.1038/ncomms11164

Gerber, T., Murawala, P., Knapp, D., Masselink, W., Schuez, M., Hermann, S., Gac-Santel, M.,
Nowoshilow, S., Kageyama, J., Khattak, S., Currie, J. D., Camp, J. G., Tanaka, E. M., &
Treutlein, B. (2018). Single-cell analysis uncovers convergence of cell identities during
axolotl limb regeneration. *Science*, *362*(6413). https://doi.org/10.1126/science.aaq0681

Goss, R. J. (1980). Prospects for Regeneration in Man. In *SECTION I11 BASIC SCIENCE AND*
PATHOLOGY.

Goss, R. J., & Grimes, L. N. (1972). Tissue Interactions in the Regeneration of Rabbit Ear Holes.
In *AM. Zooi.or.isr* (Vol. 12). https://academic.oup.com/icb/article/12/1/151/2107413

Gu, W., Schneider, J. W., Condorelli, G., Kaushai, S., Yahdavi, V., & Nadai-Ginard, B. (1993).
Intmion of Myogenic Factors and the l%tinobl ma Protein Muscle Cell Commitment and
Dmeren#ation. In *Cell* (Vol. 72).

Guerrero-Juarez, C. F., Dedhia, P. H., Jin, S., Ruiz-Vega, R., Ma, D., Liu, Y., Yamaga, K.,
Shestova, O., Gay, D. L., Yang, Z., Kessenbrock, K., Nie, Q., Pear, W. S., Cotsarelis, G., &
Plikus, M. V. (2019). Single-cell analysis reveals fibroblast heterogeneity and myeloid-
derived adipocyte progenitors in murine skin wounds. *Nature Communications*, *10*(1).
https://doi.org/10.1038/s41467-018-08247-x

HAHN, W. C., STEWART, S. A., BROOKS, M. W., YORK, S. G., EATON, E., KURACHI, A.,
BEIJERSBERGEN, R. L., KNOLL, J. H. M., MEYERSON, M., & WEINBERG, R. A.
(1999). Inhibition of telomerase limits the growth of human cancer cells. *Nature*.

Hanahan, D., & Weinberg, R. A. (2011). Hallmarks of cancer: The next generation. In *Cell* (Vol.
144, Issue 5, pp. 646–674). https://doi.org/10.1016/j.cell.2011.02.013

Hao, Y., Stuart, T., Kowalski, M. H., Choudhary, S., Hoffman, P., Hartman, A., Srivastava, A.,
Molla, G., Madad, S., Fernandez-Granda, C., & Satija, R. (2024). Dictionary learning for
integrative, multimodal and scalable single-cell analysis. *Nature Biotechnology*, *42*(2), 293–
304. https://doi.org/10.1038/s41587-023-01767-y

Hays, E. D., & Fischman, D. A. (1961). Origin of the Blastema in Regenerating Limbs of the Newt
Triturus viridescens An Autoradiographic Study Using Tritiated Thymidine to Follow Cell
Proliferation and MFgration'. In *DEVELOPMENTAL BIOLOGY* (Vol. 3).

Jiang, D., & Rinkevich, Y. (2021). Distinct fibroblasts in scars and regeneration. In *Current*
Opinion in Genetics and Development (Vol. 70, pp. 7–14). Elsevier Ltd.
https://doi.org/10.1016/j.gde.2021.04.005

Johnson, G. L., Masias, E. J., & Lehoczky, J. A. (2020). Cellular Heterogeneity and Lineage
Restriction during Mouse Digit Tip Regeneration at Single-Cell Resolution. *Developmental*
Cell, *52*(4), 525-540.e5. https://doi.org/10.1016/j.devcel.2020.01.026

Johnson, K., Bateman, J., DiTommaso, T., Wong, A. Y., & Whited, J. L. (2018). Systemic cell
cycle activation is induced following complex tissue injury in axolotl. *Developmental*
Biology, *433*(2), 461–472. https://doi.org/10.1016/j.ydbio.2017.07.010

Jopling, C., Sleep, E., Raya, M., Martí, M., Raya, A., & Belmonte, J. C. I. (2010). Zebrafish heart
regeneration occurs by cardiomyocyte dedifferentiation and proliferation. *Nature*, *464*(7288),
606–609. https://doi.org/10.1038/nature08899

Joseph, J., & Dyson, M. (1966). Tissue replacement in the rabbit's ear. *The British Journal of*
Surgery, *53*(4), 372–380. https://doi.org/10.1002/BJS.1800530415

Kim, Nam. W., Piatyszek, Miezyslaw. A., Prowse, Karen. R., Harley, Calvin. B., West, Michael.
D., Ho, Peter. L. C., Coviello, Gina. M., Wright, Woodring. E., Weinrich, Scott. L., & Shay,
Jerry. W. (1994). Specific association of human telomerase activity with immortal cells and
cancer. *Science*.

Lehoczky, J. A., Robert, B., & Tabin, C. J. (2011). Mouse digit tip regeneration is mediated by
fate-restricted progenitor cells. *Proceedings of the National Academy of Sciences of the*
United States of America, *108*(51), 20609–20614. https://doi.org/10.1073/pnas.1118017108

Leigh, N. D., Dunlap, G. S., Johnson, K., Mariano, R., Oshiro, R., Wong, A. Y., Bryant, D. M.,
Miller, B. M., Ratner, A., Chen, A., Ye, W. W., Haas, B. J., & Whited, J. L. (2018).
Transcriptomic landscape of the blastema niche in regenerating adult axolotl limbs at single-
cell resolution. *Nature Communications*, *9*(1). https://doi.org/10.1038/s41467-018-07604-0

Liu, L., Michowski, W., Kolodziejczyk, A., & Sicinski, P. (2019). The cell cycle in stem cell
proliferation, pluripotency and differentiation. In *Nature Cell Biology* (Vol. 21, Issue 9, pp.
1060–1067). Nature Publishing Group. https://doi.org/10.1038/s41556-019-0384-4

Maden, M., & Wallace, H. (1976). *How X-Rays inhibit Amphibian Limb Regeneration*.

Mine, S., Fortunel, N. O., Pageon, H., & Asselineau, D. (2008). Aging alters functionally human
dermal papillary fibroblasts but not reticular fibroblasts: A new view of skin morphogenesis
and aging. *PLoS ONE*, *3*(12). https://doi.org/10.1371/journal.pone.0004066

Morgan, T. H. (1901). *Regeneration /*. New York:
http://hdl.handle.net/2027/mdp.39015001913774

Nechiporuk, A., & Keating, M. T. (2002). A proliferation gradient between proximal and msxb-
expressing distal blastema directs zebrafish fin regeneration. *Development*.

Neeham, A. (1953). Regeneration and Wound-healing. *BioScience*, *3*(1), 16–16.
https://doi.org/10.1093/AIBSBULLETIN/3.1.16-F

Orford, K. W., & Scadden, D. T. (2008). Deconstructing stem cell self-renewal: Genetic insights into cell-cycle regulation. In *Nature Reviews Genetics* (Vol. 9, Issue 2, pp. 115–128). https://doi.org/10.1038/nrg2269

Pajcini, K. V., Corbel, S. Y., Sage, J., Pomerantz, J. H., & Blau, H. M. (2010). Transient inactivation of Rb and ARF yields regenerative cells from postmitotic mammalian muscle. *Cell Stem Cell*, 7(2), 198–213. https://doi.org/10.1016/j.stem.2010.05.022

Phan, Q. M., Sinha, S., Biernaskie, J., & Driskell, R. R. (2021a). Single-cell transcriptomic analysis of small and large wounds reveals the distinct spatial organization of regenerative fibroblasts. *Experimental Dermatology*, 30(1), 92–101. https://doi.org/10.1111/exd.14244

Phan, Q. M., Sinha, S., Biernaskie, J., & Driskell, R. R. (2021b). Single-cell transcriptomic analysis of small and large wounds reveals the distinct spatial organization of regenerative fibroblasts. *Experimental Dermatology*, 30(1), 92–101. https://doi.org/10.1111/exd.14244

Plikus, M. V, Guerrero-Juarez, C. F., Ito, M., Li, Y. R., Dedhia, P. H., Zheng, Y., Shao, M., Gay, D. L., Ramos, R., Hsi, T.-C., Oh, J. W., Wang, X., Ramirez, A., Konopelski, S. E., Elzein, A., Wang, A., Supapannachart, R. J., Lee, H.-L., Lim, C. H., … Cotsarelis, G. (2017). *Regeneration of fat cells from myofibroblasts during wound healing*. https://www.science.org

Plikus, M. V., Wang, X., Sinha, S., Forte, E., Thompson, S. M., Herzog, E. L., Driskell, R. R., Rosenthal, N., Biernaskie, J., & Horsley, V. (2021). Fibroblasts: Origins, definitions, and functions in health and disease. In *Cell* (Vol. 184, Issue 15, pp. 3852–3872). Elsevier B.V. https://doi.org/10.1016/j.cell.2021.06.024

Poleo, G., Brown, C. W., Laforest, L., & Akimenko, M. A. (2001). Cell proliferation and movement during early fin regeneration in zebrafish. In *Developmental Dynamics* (Vol. 221, Issue 4, pp. 380–390). https://doi.org/10.1002/dvdy.1152

Riddell, B., McDonough, M., Ferguson, A., Kimani, J. M., Gawriluk, T. R., Peng, C., Kiama, S. G., Ezenwa, V. O., & Seifert, A. W. (2025). Complex tissue regeneration in Lophuromys reveals a phylogenetic signal for enhanced regenerative ability in deomyine rodents. *Proceedings of the National Academy of Sciences of the United States of America*, 122(1). https://doi.org/10.1073/PNAS.2420726122

Rinkevich, Y., Lindau, P., Ueno, H., Longaker, M. T., & Weissman, I. L. (2011). Germ-layer and lineage-restricted stem/progenitors regenerate the mouse digit tip. *Nature*, 476(7361), 409–414. https://doi.org/10.1038/nature10346

Rodgers, J. T., King, K. Y., Brett, J. O., Cromie, M. J., Charville, G. W., Maguire, K. K., Brunson, C., Mastey, N., Liu, L., Tsai, C. R., Goodell, M. A., & Rando, T. A. (2014). MTORC1 controls the adaptive transition of quiescent stem cells from G 0 to GAlert. *Nature*, 510(7505), 393–396. https://doi.org/10.1038/nature13255

Saxena, S., Vekaria, H., Sullivan, P. G., & Seifert, A. W. (2019). Connective tissue fibroblasts from highly regenerative mammals are refractory to ROS-induced cellular senescence. *Nature Communications*, 10(1). https://doi.org/10.1038/s41467-019-12398-w

Schneider, J. W., Gu, W., Zhu, L., Mahdavi, V., & Nadal-Ginard, B. (1994). Reversal of Terminal
Differentiation Mediated by p107 in Rb-/- Muscle Cells. In *New Series* (Vol. 264, Issue
5164).

Sebens, K. P. (1987). The ecology of indeterminate growth in animals. *Ann. Rev. Ecol. Syst.*

Seifert, A. W., Kiama, S. G., Seifert, M. G., Goheen, J. R., Palmer, T. M., & Maden, M. (2012).
Skin shedding and tissue regeneration in African spiny mice (Acomys). *Nature*, *489*(7417),
561–565. https://doi.org/10.1038/nature11499

Seifert, A. W., & Muneoka, K. (2018). The blastema and epimorphic regeneration in mammals. In
Developmental Biology (Vol. 433, Issue 2, pp. 190–199). Elsevier Inc.
https://doi.org/10.1016/j.ydbio.2017.08.007

Sha, J. W., & Bacchetti, S. (1997). A Survey of Telomerase Activity in Human Cancer. In *Eur J*
Cancer (Vol. 33, Issue 5).

Simkin, J., Aloysius, A., Adam, M., Safaee, F., Donahue, R. R., Biswas, S., Lakhani, Z., Gensel,
J. C., Thybert, D., Potter, S., & Seifert, A. W. (2024). Tissue-resident macrophages
specifically express Lactotransferrin and Vegfc during ear pinna regeneration in spiny mice.
Developmental Cell, *59*(4), 496-516.e6. https://doi.org/10.1016/j.devcel.2023.12.017

Simkin, J., Gawriluk, T. R., Gensel, J. C., & Seifert, A. W. (2017). *Macrophages are necessary*
for epimorphic regeneration in African spiny mice. https://doi.org/10.7554/eLife.24623.001

Simkin, J., & Seifert, A. W. (2018). Concise Review: Translating Regenerative Biology into
Clinically Relevant Therapies: Are We on the Right Path? In *Stem Cells Translational*
Medicine (Vol. 7, Issue 2, pp. 220–231). John Wiley and Sons Ltd.
https://doi.org/10.1002/sctm.17-0213

Sinha, S., Sparks, H. D., Labit, E., Robbins, H. N., Gowing, K., Jaffer, A., Kutluberk, E., Arora,
R., Raredon, M. S. B., Cao, L., Swanson, S., Jiang, P., Hee, O., Pope, H., Workentine, M.,
Todkar, K., Sharma, N., Bharadia, S., Chockalingam, K., … Biernaskie, J. (2022). Fibroblast
inflammatory priming determines regenerative versus fibrotic skin repair in reindeer. *Cell*,
185(25), 4717-4736.e25. https://doi.org/10.1016/j.cell.2022.11.004

Storer, M. A., Mahmud, N., Karamboulas, K., Borrett, M. J., Yuzwa, S. A., Gont, A., Androschuk,
A., Sefton, M. V., Kaplan, D. R., & Miller, F. D. (2020). Acquisition of a Unique
Mesenchymal Precursor-like Blastema State Underlies Successful Adult Mammalian Digit
Tip Regeneration. *Developmental Cell*, *52*(4), 509-524.e9.
https://doi.org/10.1016/j.devcel.2019.12.004

Tanaka, E. M., & Brockes, J. P. (1998). A target of thrombin activation promotes cell cycle re-
entry by urodele muscle cells. *Wound Repair and Regeneration*, *6*(4), S-371-S-381.
https://doi.org/10.1046/J.1524-475X.1998.60413.X/FORMAT/PDF

Tanaka, E. M., Gann, A. A. F., Gates, P. B., & Brockes, J. P. (1997). Newt Myotubes Reenter the
Cell Cycle by Phosphorylation of the Retinoblastoma Protein. In *The Journal of Cell Biology*
(Vol. 136, Issue 1).

Tomasso, A., Koopmans, T., Lijnzaad, P., Bartscherer, K., & Seifert, A. W. (2023). *An ERK-*
dependent molecular switch antagonizes fibrosis and promotes regeneration in spiny mice
(Acomys). https://www.science.org

Vorontsova, M. A., Liosner, L. D., Allen, P. M., & Billett, F. (1960). *Asexual propagation and*
regeneration.

Wallace, H. (1981). *Vertebrate Limb Regeneration* (John Wiley & Sons, Ed.; First edition).

Wallace, H., & Maden, M. (1976). THE CELL CYCLE DURING AMPHIBIAN LIMB
REGENERATION. In *J. CeUSci.zo*.

Wenemoser, D., & Reddien, P. W. (2010). Planarian regeneration involves distinct stem cell
responses to wounds and tissue absence. *Developmental Biology*, *344*(2), 979–991.
https://doi.org/10.1016/j.ydbio.2010.06.017

Zhang, W., Shi, Y., Oyang, L., Cui, S., Li, S., Li, J., Liu, L., Li, Y., Peng, M., Tan, S., Xia, L., Lin,
J., Xu, X., Wu, N., Peng, Q., Tang, Y., Luo, X., Liao, Q., Jiang, X., & Zhou, Y. (2024).
Endoplasmic reticulum stress—a key guardian in cancer. In *Cell Death Discovery* (Vol. 10,
Issue 1). Springer Nature. https://doi.org/10.1038/s41420-024-02110-3

FIGURES AND LEGENDS

Figure 1. Cell cycle re-entry before and after injury in *Acomys*, *Mus*, *Danio* and

***Ambystoma*.** A) Tissue section planes and EdU/BrdU pulse-chase scheme indicating injections

and harvest times; D = distal, P = proximal. B) Representative images of EdU+ and BrdU+ cells

in uninjured tissue (D0), D1 and D3. In uninjured tissue, white arrows = positive cells in the

epidermis and green arrows = positive cells in the stromal compartment. In injured tissue, red

arrows = EdU+ cells, light blue arrows = BrdU+ cells, and white arrows = double positive cells

(EdU+/BrdU+). C) EdU+ cell quantification as a percentage of the total number of cells in

uninjured tissue from all four species. Autofluorescent tissue, including red blood cells, stains in

multiple channels (yellow). D) EdU+, Brdu+ and EdU+/BrdU+ cell quantification as a

percentage of the total number of cells at D1 and D3 post-injury. Images represent n = 3

biological samples/time point in *Acomys* and *Mus*. See Methods for cell counting quantification.

In *Danio* (zebrafish), n = 9 at D0, n = 5 at D1 and n = 4 at D3. In *Ambystoma* (axolotl), n = 12 at

D0, n = 6 at D1, n = 6 at D3. (* $p < 0.05$, ** $p < 0.01$).

Figure 2. Spatiotemporal distribution of proliferating cells in *Acomys* and *Mus*. A)

Representative images of blastema and scar tissue changes over time (D5-D30), Ca = cartilage,

WM = wound margin, HF = hair follicle. B-C) Blastema or scar tissue area in mm^2 from D10 to

D30 based on tissue sections. D-E) Quantification of BrdU+ cells as a percentage of the total

number of cells in three areas (A = blue, B = red, C = green) of the blastema in *Acomys* or the

scar in *Mus* from D5 to D30. F-H) Representative images of D10 (F) and D25 (H) blastema

tissue with EdU+, BrdU+ and double positive cells. Red arrows = EdU+ cells, light blue arrows

= BrdU+ cells, and white arrows = double positive cells (EdU+/BrdU+). G-I) Quantification of

EdU+, BrdU+ and EdU+/BrdU+ cells as a percentage of the total number of cells in distal and

proximal areas relative to the wound site at D10 (G) and D25 (I). n = 5 per time point per

species. (* $p<0.05$).

Figure 3. S-phase quantification of fibroblasts, chondroblasts and macrophages after

injury at D5 in *Acomys* and *Mus*. A-D) Cycling macrophages double labeled with IBA1 and

EdU in *Acomys* (A-A') and *Mus* (C-C') or with CD206 and EdU in *Acomys* (B-B') and *Mus* (D-

D'). E-F) Chondroblasts double labeled with Sox9 and EdU in *Acomys* (E-E') and *Mus* (F-F').

G-J) Fibroblasts double labeled with CRABP1 and EdU in *Acomys* (G-G') and *Mus* (I-I'). or

with αSMA and EdU in *Acomys* (H-H') and *Mus* (J-J'). White arrows indicate double positive

cells (EdU+/cell type marker). Pie charts indicate percentage double positive cells of the total

number of counted EdU+ cells. n = 5/species. RBC = red blood cells.

Figure 4. Spatial distribution of cell types within blastema or scar at D20. A-C) CRABP1+

(A-A') or αSMA+ (B-B') fibroblasts, and Sox9+ chondroblasts (C-C') in *Acomys* at D20. D-F)

CRABP1+ (D-D') or αSMA+ (E-E') fibroblasts, and Sox9+ chondroblasts (F-F') in *Mus* at D20.

G) CRABP1 and αSMA double staining in *Acomys* over time (D5-D30). H) Cartoon showing

general distribution of CRABP1+ and αSMA+ fibroblasts in *Acomys.* White arrows indicate

double positive cells (EdU+/cell type marker). Pie charts indicate percentage double positive of

the total number of counted EdU+ cells. n = 5 per species. RBC = red blood cells.

Figure 5. Single-cell analysis reveals different fibroblast cell states in *Acomys* and *Mus*. Cell

suspensions collected from healing tissue preps representing D0, 3, 5, 10 and 15 were batch-

corrected and plotted together (Fig. S7) (see Methods). A) UMAP projection of stromal cell

types re-clustered from all cells (Fig. S7) from both species. B) UMAP projection of stromal

subpopulations separated by species. Clusters represent subpopulations of major subtypes

indicated in A. C-E) Feature plots for *Crabp1* (C), *Acta2* (D), and *Sox9* (E) separated by species.

F, H, J) Volcano plots showing differentially upregulated genes between *Acomys* and *Mus* from

clusters four (F), six (H), and seven (J) in B. Gene ontology Biological Process based on

differentially upregulated genes between *Acomys* and *Mus* clusters four (G), six (I), and seven

(K). Dot size represents the relative number of genes in each plot. Upregulated genes associated

with each pathway are in color boxes. Refer to supplementary table S4 to see full list of genes.

Figure 6. Tissue injury in neonatal mouse ears when CRABP1+ fibroblasts are present

does not facilitate regeneration. A) Immunostaining for CRABP1 in *Acomys* and *Mus*. Age in

days indicated in upper right corner of each image. B) Images showing representative ear hole

closure from P20 injured *Mus* after a 4 mm ear punch. C) Quantification of ear hole closure area

from D5 to D40. D) CRABP1 staining in P15 *Mus* before and after injury. n = 3 per species per

time point. Green channel shows autofluorescence including red blood cells (RBC).

954