Nanoprotrusion enlargement conducts cytocapsular tube elongation along the path of least resistance

Tingfang Yi[1,2*] and Gerhard Wagner[1,2,3*]

1. Cytocapsula Research Institute, 245 First Street, Cambridge, MA, 02142 USA
2. Centiver Ltd., 245 First Street, Cambridge, MA, 02142 USA
3. Department of Biological Chemistry and Molecular Pharmacology of Harvard Medical School, 240 Longwood, Boston, MA, 02115 USA

*To whom correspondence may be addressed. E-mail:tingfang_yi@centiver.com, or Gerhard_Wagner@hms.harvard.edu.

Abstract

Cancer metastasis caused multiple tissue and organ biological function failure is a major resource of cancer lethality. Recently, we identified membrane protein Ca^{2+} ion channel Ca2+-ATPase 2 (PMCA2) as a biomarker of malignant tumors. Cytocapsular tumors at primary and secondary cancer niches are interconnected by three-dimension (3D) cytocapsular tube networks and compose dynamic and integrated cytocapsular tumor network systems (CTNSs) in cancer patients. Cytocapsular tubes (CCTs) provide membrane-enclosed freeways for protected, bi-directional and efficient cancer metastasis. It is obscure how CCTs elongate in heterogeneous human tissues and organs harboring heterogeneous resistances. Here, we report that CCT nanoprotrusions (NPs) with least resistance enlarge and grow into enlarged NPs. Subsequently, enlarged NPs develop into initial CCTs followed by new CCT formation and CCT elongation.

CCTs elongate in the path along least resistance in cancers in 34 kinds of human tissues and organs. CCT branching morphogenesis employs CCT side NP enlargement and bifid branching format, which increases CCT numbers and expands 3D CCT networks. CCT regeneration drives repeated cancer metastasis and promotes cancer relapse. CCT superstructures facilitate complete cancer metastasis in acytocapsular oncocell mass-CC/CCT complex (AMCC) tissues. This study demonstrated nanoprotrusion-conducted CCT elongation along the path with least resistance and branching morphogenesis in bifid branching style. The cytocapsular membrane system expansion mechanics of CCTs and CCT networks described here may open new avenues for native cancer research and CTNS-targeted effective cancer cure. (227 words)

Significance statement

The recent identification of PMCA2 as a biomarker for malignant tumors, allows imaging very early native cancer growths on culture plates *in vitro* and in human bodies as revealed from snapshot images of numerous cancer biopsies *in vivo*. Aggressive cancer metastasis, cancer progression, repeated cancer relapse, limited cancer therapy outcomes, pan-cancer drug resistance, immune cell attack escape, and immunotherapy non-responsive "cold" tumors are unmet challenges in clinical cancer therapy. Moreover, the cytocapsular tube (CCT) elongation mechanisms in human tissues/organs with heterogeneous resistances is unclear. Here, we report that CCT nanoprotrusions (NPs) with least resistance preferably grow up into enlarged NPs, which subsequently develop into initial CCTs. Intracytocapsular oncocells released cytocapsulasomes fuse into initial CCTs, and increase cytocapsular membrane areas. Oncocell push and deform initial CCT membranes and generate new CCTs with new nanoprotrusion layers. CCT

elongate along the path with least resistances in heterogeneous tissues/organs. CCT branching morphogenesis develops in a bifid format, significantly increasing CCT number and expanding 3D CCT networks in all kinds of human tissues. CCT superstructures promote complete cancer metastasis in AMCC complexes. Repeated CCT regeneration drives repeated cancer relapse. This study reveals the CCT elongation laws, and elucidates that CCT elongation and branching morphogenesis drives cytocapsular membrane protected cancer metastasis, and that repeated CCT regeneration promotes repeated tumor relapse *in vivo*.

Introduction

Recently, we identified membrane protein Ca^{2+} ion channel Ca2+-ATPase 2 (PMCA2, or ATP2B2) as a biomarker of malignant tumors of 293 subtypes of cancers in 34 kinds of human tissues and organs[1-3]. The anti-PMCA2 antibodies can be used in immunohistochemistry (IHC) florescence staining assays for fluorescence microscope imaging analysis of both solid cancers and hematology cancers in tissues[1-3]. Now, we can initiate native malignant tumor growth *in vitro* with our optimized culture kits of CC/CCT culture kits, and identify basic mechanisms for initial growth of native malignant tumors with bright field and IHC fluorescence microscope imaging technologies *in vitro*. By accessing more than 15,000 pieces of specimens of more than 14,000 cancer patients from clinically annotated cancer tissue banks, we can also follow the growth/elongation of cancers in human bodies *in vivo*.

Cancer metastases are the major source of cancer deaths[4-12]. New insights into mechanisms of metastases have come from the discovery of cytocapsulas and cytocapsular tubes (CCTs) which provide the major pathways for cancer cell dissemination. The understanding of native cancer progression mechanics at the level of cytocapsular tumor network systems (CTNSs) in cancer patients is a prerequisite for effective and efficient cancer research, therapy and treatments[1-3]. The Cytocapsula and the cytocapsular tube (CCT), two newly discovered large, cancer cell specific organelles enclosing oncocells or oncocell masses, provide the keys for understanding CTNS progression and cytocapsular membrane-sheltered tumor evolution in tissues *in vivo*. In particular, native CCT networks provide 3D, membrane-enclosed freeway systems for protected and efficient cancer cell migration and dissemination in human tissues and dominate cancer metastasis in cancer patients *in vivo*[1-3]. Human tissues and organs in cancer micro-environments harbor extremely heterogeneous materials/factors, including all kinds of cells, extracellular matrix (ECM), cell-cell and cell-ECM interactions, solid stresses, interstitial fluid pressure, physical microarchitectures, which are resistances of CCT elongation[13-24]. Interestingly, we find that progressing tubes generate wave-like pattens and larger triangular like progression paths. It is obscure how CCTs elongate and expand in heterogeneous human tissues and organs.

There are growth lifecycle of native malignant tumors and cytocapsular tumor network system (CTNS), and native cancer metastases in our previous report, but little about metastasis pathways, which is focused here. There are important open questions: 1) How do cytocapsular tumor growth from single cancer cell to cytocapsular tumorspheres and cancers? 2) Whether and how do cytocapsular nanoprotrusions develop into cytocapsular tubes (CCTs)? 3) What determines the direction of cytocapsular tube tumor growth? Here, we investigated CCT elongation in

cancers in 34 kinds of human tissues and organs, and found that: (i) CCT nanoprotrusion enlargement conducts CCT elongation along the path with least resistance, (ii) CCT bifid branching morphogenesis increases CCT numbers and expands CCT networks. (iii) CCT regeneration in AMCC associates with new cycles of CCT-conducted cancer metastasis, and (iv) CCT superstructures and mega superstructures promote complete cancer metastasis in AMCC. These discoveries may facilitate cancer researches, cancer drug research and development, and clinical cancer applications in cancer diagnosis, surgery, therapy, prognosis and treatments.

Results

Cytocapsular nanoprotrusions grow into initial cytocapsular tubes followed by new CCT formation and CCT elongation

At 24hour (h) after cell implantation, single Bxpc3 pancreas cancer cells in CC/CCT culture kits generate cytocapsulas (CCs) with many nanoprotrusions (NPs) and NP bunches, which interconnect and form NP networks surrounding CCs (**Extended Data Fig. 1A**, n = 316 tested cytocapsular oncocells). In the ecellulated CCs (ECCs) of single cytocapsular oncocells, NPs and NP bunches formed NP networks interconnect with and surround ECCs (**Extended Data Fig. 1B**, n=56 ECCs). These results show that single cytocapsular oncocells' cytocapsular membranes extend and generate many NPs and NP bunches, which interconnect and form NP networks surrounding cytocapsular oncocells (**Extended Data Figs. 1A-1B**). After about 6h, ecellulated CCs and their NP networks degrade, followed by disappearance (**Extended Data Fig. 1B**), suggesting that intracytocapsular oncocells drive NP generation and maintenance. At 36h, Bxpc3 pancreas intracytocapsular oncocells proliferate and generate cytocapsular tumorspheres wrapped in the enlarged CCs, which engender larger NP networks surrounding enlarged CCs

(**Extended Data Fig. 1C**, n= 98 cytocapsular tumorspheres). Transmission electron microscope (TEM) analyses of breast cancer stem cell (BCSC) HMLER (CD44^{high}/CD24^{low})^{FA} cell NPs show that the NPs are coated with glycocalyx (a mixture of glycoproteins and glycolipids) in variable thickness (up to 78nm in thickness, **Extended Data Fig. 1D**). At 36h, the NP sizes are variable (21nm ~ 82nm in diameter/width, 300-2700nm in length). Further TEM analyses of these BCSC cytocapsular oncocell NP bunches show that two or more straight or curved NPs align together tightly/loosely and form NP bunches, coated with glycocalyx layers in wider thickness range (2nm~1000nm in thickness, **Extended Data Fig. 1E**). In NP bunches, single NP sizes are 18nm ~ 32nm in diameter/width. The NP bunches sizes are variable (40nm ~ 152nm in width, **Extended Data Fig. 1E**). NP generate branches, indicating NP branching morphogenesis is a resource of NP number increase and NP network formation surrounding CCs (**Extended Data Fig. 1E**). Glycocalyx may cause the blur/smear effects of NP networks in bright filed and immunohistochemistry fluorescence microscope imaging analyses. Subsequently, at 48h, one or a few single Bxpc3 cytocapsular tumorspheres' cytocapsular NPs elongate in length (up to 100μm in length, n=46 NPs) and enlarge in size (up to 0.1μm in diameter/width, n=46 enlarged NPs), and link CCs of cytocapsular tumorspheres (CTs, **Extended Data Fig. 2A**). There is no oncocell in migration in these early enlarged NPs (82nm<NP diameter/width <0.1μm) directly linking two CTs (**Extended Data Fig. 2A,** n=126 enlarged NPs). Not all NPs are equally to have chances to enlarge, and only the shortest NPs between CTs are preferred to grow up and enlarge (**Extended Data Fig. 2A**, n=126 enlarged NPs), indicating that NPs networks sense distances between CTs and the NPs with shortest distance and least resistance of molecular communications between CTs stand out, grow up and enlarge in sizes.

At 56h, single Bxpc3 cytocapsular tumorsphere enlarged NPs linking CTs continue to grow up in sizes and generate cytocapsular tubes (CCTs) in various sizes of 0.1μm ~ 3μm in diameter/width (**Extended Data Fig. 2B,** n=65 NPs) *in vitro*. In the NP enlargement generated early CCTs with sizes (0.1μm ~ 0.5μm in diameter/width), there are oncocells in migration inside, indicating these early CCTs are propertied with biological functions for CCT-conduced oncocell migration inside. These NP-enlargement engendered early CCTs are termed as initial CCTs (ICCTs). Established CCTs with sizes (>0.5μm in diameter/width, **Extended Data Fig. 2C**) allow bi-directional oncocell migration between cytocapsular tumorspheres *in vitro*. There are many NPs surrounding ICCTs and CCTs, anchoring ICCTs and CCTs in place. The above data demonstrated that a few cytocapsular NPs with shortest distance and least resistance of molecular communications between CTs are selectively to grow into ICCTs and CCTs for oncocell migration inside *in vitro*. NP enlargement and CCT development are terminated in ecellulated cytocapsulas (from cytocapsular oncocells/tumorspheres, **Extended Data Figs. 1 and 2**), demonstrating that NP enlargement, ICCT development and CCT elongation are depended on incytocapsular oncocell driving (cytocapsulasomes and other molecules).

Consistently, in the nuperphase tumor of invasive breast carcinoma, there are many initial CCTs *in vivo* (ICCTs, 0.2μm ~ 0.5μm in diameter/width, 10μm ~ 230μm in length in sectioned specimens, n=215 ICCTs, from 112 breast cancer patient; Fig. S3A). There are oncocells in migration in some initial CCTs (**Extended Data Fig.S3A**). These observations of initial CCTs in clinical invasive breast carcinoma tissues evidenced that clinical breast cancer cytocapsular nanoprotrusions (NPs) grow into initial CCTs in tissues *in vivo*. The initial CCTs elongate in tiny pores or small cavities with least resistance (**Extended Data Fig.S3A**), indicating the NPs sense

resistance in tissues, and NPs with least resistances grow up and enlarge, and engender initial CCTs. Next, we examined initial CCTs in cancers of 34 kinds of human tissues/organs (with 289 subtypes). Indeed, initial CCTs are presented in all the investigated cancers of 34 kinds of human tissues/organs (289 subtypes of cancers, **Extended Data Fig. 4**). In some instances, the sectioned thin initial CCT fragments are straight and crossing tissues. Most sectioned initial CCT fragments are in curved or irregular morphologies in tissues. Later, the breast cancer initial CCTs grow up into CCTs (0.8μm ~ 6μm in diameter/width) and with oncocells in dissemination inside. Both initial CCTs and CCTs are anchored in local positions in tissues, and CCT morphologies and locations largely reflect the initial CCTs' morphologies and locations from which they grow up in tissues (**Extended Data Fig. 3A-B**). In summary, cytocapsular NPs grow into CCTs for oncocell migration and cancer metastasis *in vitro* and in tissues *in vivo*, and CCTs anchored in place in tissues reflect/record the locations and paths of initial CCTs and the selected NPs (**Extended Data Fig.3C**).

Cytocapsular tubes elongate in the path with least resistance

Human tissues and organs harbors extremely heterogeneous materials/factors, including all kinds of cells, extracellular matrix (ECM), cell-cell and cell-ECM interactions, solid stresses, interstitial fluid pressure, physical microarchitectures, which are resistances of CCT elongation[25-27]. However, it is obscure how CCTs elongate and expand in heterogeneous human tissues and organs.

Next, we investigated elongation of super-long CCTs in tissues *in vivo*. In some instances, single pancreas adenocarcinoma cytocapsular oncocells generate very long, bent and curved CCTs (>7,000μm in length in sectioned cancer tissue specimens, >700-folds of average single cell

diameter of 10μm) in the same plane in heterogeneous tissues, and being contained in a macro bent space (**Figs. 1A-B**). The super-long, super-large, oval-shaped and bent CCT suggests that the long pancreas cancer CCT elongates in a path with least resistance, which is sandwiched and bent by two layers of substantial resistances (the above big oval-shaped CCT bunch belt and the below big oval-shaped dense CCT strand belt, **Fig 1A**). The super-large, oval-shaped, bent CCT superstructure harbors a lot of small, near periodic or aperiodic waves, or irregular curves. At the crest and trough positions, CCTs meet physical resistances from extracytocapsular tissues, and choose to change CCT elongation directions and extend along the paths with least resistance. In near periodic waves, the average wave heights are 52 μm ± 6.2 μm, and average wavelengths (λ) are 61 μm ± 3.4 μm （n=122 near periodic waves in tissues). These evidences demonstrated that the CCT experienced variable resistance at micro local position and always elongate forward in a path with least resistance (**Fig.1B**). Furthermore, single pancreas adenocarcinoma cytocapsular oncocell generate super-long, (>8,200 μm in length in sectioned cancer tissues >820-folds of average single cell diameter of 10μm), super-large, bent, flat ring-shaped CCT in the same plane in heterogeneous tissues (**Extended Data Fig. 5**). This long pancreas cancer CCT elongates in a path with least resistance sandwiched and bent by two layers of substantial resistances (the above super-large, flat ring-shaped and dense tissue cell layer and the below big flat ring-shaped dense CCT strand belt, **Extended Data Fig. 5**). Moreover, single prostate adenocarcinoma cytocapsular oncocells generate super-long, (>20,500μm in length in sectioned cancer tissues, >2,050-folds of average single cell diameter of 10μm), super-large, irregular-shaped, continuous, and highly bent CCT in superstructures in the same plane in heterogeneous tissues (**Extended Data Fig. 6**). The very-long single prostate cancer CCT with many small or big curves consistently elongates along the path with least resistance in the heterogeneous tissues

(**Extended Data Fig. 6**). In colon adenocarcinoma tissues, single colon cytocapsular oncocells generate very long, highly curved CCTs in heterogenous tissues (**Extended Data Fig. 7**). The above data evidenced that native single cytocapsular oncocells can generate super-long CCTs with variable superstructures, and that CCTs elongate along the path with least resistance in heterogeneous tissues *in vivo* (**Figs. 1 and Extended Data Fig. 5-7**).

In most instances, in invasive breast carcinoma tissues, there are large quantities of sectioned, short and curved CCT fragments. The high percentages of short CCT fragments with curves/turns/coils/ spirals suggest the high frequency of CCT elongation direction changes (**Fig. 1C-D**). The CCT elongation direction changes at CCT turns, curves, or crests/troughs of CCT wave fragments are caused by heterogeneous obstacles from tissue cells, ECMs, microarchitectures, and extend along the path with pores, microenvironment space, less cell density, less ECM density, less CCT density, where have least physical resistances (**Fig. 1C-D**).

In summary, the above *in vitro* and *in vivo* data on CCT elongation demonstrated that CCT elongation procedure/cycle contains several successive steps: 1) generation of enlarged NP via elongation and enlargement of selective nanoprotrusions (NPs) with least resistance, 2) enlarged NP grows into initial cytocapsular tube (ICCT), 3) with cytocapsulasome fusion, ICCT increase cytocapsular membrane areas, oncocell push and deform CCT membranes, and elongate CCT (**Extended Data Fig. 8**).

Consequently, CCT elongation, cancer cell metastasis via CCTs, and CCT degradation in tissues usually occur ahead of CCT invasion into blood vessels, as blood vessel walls have tight and dense cell-cell interactions and strong physical resistances (**Extended Data Fig. 9**).

CCT branching morphogenesis increases CCT number and drives cancer metastasis

In some instances, at 72h after Bxpc3 pancreas cancer cell implantation, nanoprotrusions (NPs) at the side of CCTs enlarge and grow into branch enlarged NPs (branch ENPs, **Fig.S10A-B**). Subsequently, the branch ENPs develop into a branch initial CCTs (branch ICCTs) and branch CCTs (**Extended Data Fig. 10A-B**). In the examined 75 cytocapsular oncocell CCT with branches *in vitro* (21 of Bxpc3 pancreas cytocapsular oncocells, 18 of A549 lung cytocapsular oncocells, 36 of MDA-MB-231 breast cytocapsular oncocells), all these CCT branches employ bifid branching format.

Consistently, in clinical cancer tissues, CCT branching morphogenesis utilizes CCT side NP enlargement and bifid branching format (**Fig. 2A**). In the examined 126 CCTs with branch CCTs (55 CCT branches from breast carcinoma, 26 CCT branches from colon carcinoma, 17 CCT branches from prostate carcinoma, 28 CCT branches from lung carcinoma) employ side NP enlargement and bifid branching format. One stem CCT can perform multiple branching in a short CCT distance (**Fig.2A4**). After CCT remodeling, the branch CCT sizes in diameter/width are almost similar as that of stem CCTs. Oncocells bi-directionally migrate in CCT networks containing stem and branch CCTs (**Figs. 2A-B, and Extended Data Fig. 10C**).

With the >120,000 pieces of CCT images of 13,000 pieces of cancer tissues in AMCC, we suggest that AMCC can be divided into 4 stages based on ICCTs/CCTs/CCT strand occupied area percentages: Stage I, ICCTs/CCTs/CCT strands occupied area is 0-10%; Stage II, 10%-40%; Stage III, 40%-70%; and Stage IV, 70%-100%. In AMCC stage in clinical breast and colon carcinoma tissues, CCT branching morphogenesis significantly increase CCT numbers from AMCC phase I to AMCC phase IV (**Fig.2**). CCT branching morphogenesis contributes significant increase in both CCT number and density in AMCC stage in breast cancer, prostate cancer and colon cancer (**Extended Data Fig. 11**). CCT branching morphogenesis is universally presented in all the checked 288 subtypes of cancers in 35 kinds of human tissues and organs (**Extended Data Fig. 4**). In addition, increased CCT occupied areas associate with increased cancer cell metastasis, and decreased left cancer cells (**Fig. 2C-D, and Extended Data Fig. 11**). These data demonstrated that CCT branching morphogenesis substantially contributes to CCT number and density increase, cancer cell metastasis progression, and cancer evolution.

CCT regeneration and elongation drive continuous cancer metastasis in AMCC

In acytocapsular oncocell mass-CC/CCT complex (AMCC) stage, acytocapsular oncocell continuously perform uncontrolled oncocell proliferation, and one generation of CCTs can only reach metastasis of parts of acytocapsular oncocells, and massive cancer cells remain (**Figs.3 and Extended Data Fig. 12**). Interestingly, after old CCTs degrade, decompose, and disappear, leaving CCT cavity strips formed 3D CCT cavity networks in acytocapsular oncocell mass (**Figs.3 and Extended Data Fig. 12**). Subsequently, new CCTs are regenerated. With CCT branching morphogenesis, CCT number and density significantly increase, and extend along CCT cavity strips formed 3D CCT cavity networks left by previous generation CCT network (**Figs.3 and Extended Data Fig. 12**). The new generation CCT networks continuously let more

oncocells dissemination and leave away, and therefore drive cancer relapse (**Figs.3 and Extended Data Fig. 12**). The phenomena of repeated CCT regeneration and elongation universally present in all checked 289 subtypes of cancers in 34 kinds of human tissues and organs (**Extended Data Fig. 4**). These observations demonstrated that repeated CCT regeneration and elongation in AMCC drive repeated cancer metastasis and cancer relapse**.**

CCT superstructures promote complete cancer metastasis in AMCC stage

Single, super-long, and highly curved/coiled CCTs generate big, near triangle CCT super-structures (**Fig.4**). Multiple big, near triangle CCT super-structures compose multi-layered CCT super-structure complex, wrapping multiple irregular shaped cytocapsular tumors in variable sizes (**Fig.4**). Similar multiple-layered CCT superstructure complexes in diverse morphologies present in breast carcinoma, lung carcinoma, pancreas carcinoma (**Extended Data Fig. S13-14**). The average acytocapsular oncocell densities in the CCT super-structures are close to or almost zero, or much lower than that in the neighboring acytocapsular oncocell mass area (**Fig.4, and Extended Data Figs. 13-14**), suggesting that CCT superstructure complexes significantly enhance CCT density, dramatically increase CCT membrane surface areas for acytocapsular oncocell alloentry into CCTs, and therefore promote cancer metastasis via CCTs. CCT superstructure complexes allow acytocapsular oncocell masses to completely leave away via CCT conducted cancer disseminate with no acytocapsular oncocell left (**Fig.4, and Extended Data Figs. 13-14**). Therefore, CCT superstructure complexes advance in reaching complete cancer metastasis of acytocapsular oncocell masses.

Discussion

The discovery of cytocapsula, cytocapsular tube (CCT), CCT networks, cytocapsular oncocells/tumors, and cytocapsular tumor network systems (CTNSs) in our two initial reports on CCT-cancer have unveiled a previously unnoticed principle of native cancer *in vivo*. Here, we show that CCT elongation in the least resistance path and CCT branching morphogenesis in bifid format universally present in cancers *in vivo*. This was achieved via a 7-year effort to collect >300,000 pieces of CCT fluorescence microscope images of clinical cancer tissue specimens from >15,000 cancer patients from tissue banks worldwide, with a goal to investigate CCT development in 288 subtypes of human cancers in diverse stages. The CCT-elongation mechanisms that emerged as principles for native cancer progression research, and clinical cancer diagnosis and therapies.

The CCT-elongation mechanisms elucidate the principles of CCT elongation, CCT network expansion, CCT freeway system conducted and cytocapsular membrane protected cancer metastasis, and intro-cytocapsular oncocell proliferation and migration driven cancer progression in heterogeneous human tissues and organs. The leading intro-cytocapsular oncocells drive CCT elongation and CCT network expansion in multiple aspects: 1) generate and release cytocapsulasomes and other membrane-enclosed tiny bodies as CCT elongation membrane building blocks; 2) release proteases and provide proteolysis to remove resistances from ECMs (including bone matrices), cell-cell interactions (including brain-blood barriers) and cell-ECM interactions; 3) receive signals from outside nanoprotrusions and decide the preferable NPs with least resistance to enlarge and grow up; 4) generate blebs to sense extracytocapsular microenvironmental resistances and push neighboring cells/ECMs back to make space for NP enlargement and initial CCT expansion in diameter/width and length; 5) push and deform cytocapsular membranes into tube-shaped morphologies, and generate new NPs anchoring CCT

in place; 6) drive side NPs to develop into branch CCTs and promote CCT branching morphogenesis, and therefore provide expanded 3D CCT freeway systems for oncocell metastasis in a membrane barrier protected cell dissemination format, escape from immune cell attack, achieve pan-cancer drug resistance; 7) Drive repeated CCT regeneration, and new generations of CCTs and CCT networks promote new cycle of cancer metastasis, resulted in repeated tumor relapse.

The advantages of CCT elongation via nanoprotrusion (NP) enlargement conducted new CCT formation are: 1) the single NPs are in tiny sizes of 21nm ~ 82nm in diameter/width, much smaller than CCTs in sizes (3-10 μm in diameter/width), which help NPs to extend into large quantities of micropores (<100nm in width) in microenvironments with minimum resistance; 2) the large quantities of NPs surrounding CCTs and CCT leads help CCTs to sense resistance in all 3D directions in micro positions in local microenvironments, and choose the NPs with least resistances to enlarge and develop into new CCTs; 3) NP membranes are extended from CCT membranes and share similar cytocapsular membrane structural components (membrane lipids, membrane proteins, beneath microfilament networks, and so on), and relatively easy to enlarge in sizes; 4) the open connection of NPs at/in CCT ultrastructural complexes facilitate NPs to access and obtain intracytocapsular oncocell released membrane building materials and proteases, without physical barriers or obstacles; 5) NPs in the leading release proteases and proteolyze to remove resistances from ECMs (including bone matrices), cell-cell interactions (even brain-blood barriers) and cell-ECM interactions in tiny scale initially, which is an efficient way to find out a path with least resistance. The above multiple unique advantages significantly facilitate NP-conducted CCT elongation in the least resistance path to overcome/remove heterogeneous macro and micro resistances in heterogeneous human tissues/organs.

CCT branching morphogenesis in side NP enlargement and bifid format is propertied with multiple characterizations: 1) branch CCTs are developed from CCT side and enlarged NPs with least resistance, which interconnect with stem CCTs with open connection, sharing with similar cytocapsular membrane components and shared the similar/same materials in stem CCT lumens and NP lumens, 2) intracytocapsular oncocells (in long, thin and spindle-shaped mesenchymal morphologies in initial CCTs) release cytocapsulasomes, which integrate into initial CCTs, and increase CCT membrane areas; push and deform cytocapsular membranes to form bigger tube-shaped CCTs; 3) stem CCT sides have large quantities of CCTs already extend into ECMs anchoring CCTs in place, which provide substantial opportunities to develop multiple or many CCT branches via multiple/many times of bifid branching; 4) CCT branching morphogenesis significantly increase CCT numbers and expand CCT networks in network sizes and cytocapsular membrane area sizes for outside acytocapsular oncocells to invade into CCT network systems via alloentry format. Therefore, more CCTs promote cancer cell metastasis to leave away from the stressful inner tumor microenvironments (uncontrolled cancer cell proliferation caused nutrient deprivation, hypoxia, low pH, elevated metabolism waste molecules, insufficient growth factors, high competition from neighboring cancer cells, toxic microenvironments) for better survival conditions.

In summary, the CCT elongation laws and mechanisms of CCT elongation and CCT branching morphogenesis described here may facilitate native cancer research in a CTNS level and help clinical cancer applications, including cancer diagnosis, surgery, caner drug research and development, and CTNS targeted effective cancer therapy.

Materials and Methods

Reagent, antibodies and devices

CC/CCT culture kits (Celldevi, www.celldevi.com , Cat. CD0104, 6-well plates; Cat. CD0105, 12-well plates, and Cat. CD 0106, 24-well plates) and kits with glass cover slips in the well bottom and embedded by CC/CCT culture matrix layer (Cat. CD 0112, 6-well plate) were ordered from Celldevi Inc. CC/CCT culture kit fixation kit (Cat. CD0201) were ordered from Celldevi Inc. Cancer cell lines of pancreas cancer cell Bxpc3, breast cancer cell MCF-7 and colon cancer cell SK-CO-1 and cell culture media were ordered from ATCC. $^{13}C_6$, $^{15}N_2$-L-Lysine (Cat. 88209) and $^{13}C_6$, $^{15}N_4$-L-Arginine (Cat. 89990) for SILAC labeling were ordered from Thermo Fisher Scientific. Unipick^{TM} and capillary units were ordered from NeuroInDx Inc. Rabbit anti-PMCA2 antibody (polyclonal, ab3529; 1:200 dilution), mouse anti-γ-actin antibody (monoclonal, ab123034; 1:200 dilution) for immunofluorescence assay were ordered from Abcam. DAPI (1:1,000 dilution in the immunofluorescence assay) was ordered from KPL. Human normal and cancer tissue specimens were ordered from TissueArray and US Biolab (or gifted by local hospitals).

References:

1. Yi T, Wagner G, Cytocapsular tubes conduct cell translocation. *Proc Natl Acad Sci USA*. **115**, E1137-E1146 (2018).
2. Yi T, Wagner G, Malignant tumor cells engender second membrane-lined organelles for self-protection and tumor progression. *Proc Natl Acad Sci USA*. **121**, e2317141121 (2024).
3. Yi T, Wagner G, Discovery of the Cytocapsular Membrane as Hallmark of Malignant Tumors. *Biochemistry*. 64:555-562 (2025).
4. Cassetta L, Pollard J, A timeline of tumor-associated macrophage biology. *Nat Rev Cancer*. **23**, 238-257 (2023)
5. Chaffer C, Weinberg RA, A Perspective on Cancer Cell Metastasis. *Science*. **331**, 1559-64 (2011).
6. Hanahan D, Weinberg RA, Hallmarks of cancer: the next generation. Cell. **144**, 646-74 (2011).
7. Sung H, et al, Global cancer statistics 2020: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. Online ahead of print (2021).
8. Chiang A, Massagué J, Molecular basis of metastasis. *N Engl J Med*. **359**, 2814-23(2008).
9. Dienstmann R, et al, Personalizing colon cancer adjuvant therapy: selecting optimal treatments for individual patients. *J Clin Oncol*. **33**, 1787-96 (2015).
10. Mody K, Bekaii-Saab T, Clinical Trials and Progress in Metastatic Colon Cancer. *Surg Oncol Clin N Am*. **27**, 349-365 (2018).
11. Patras L, et al, Immune determinants of the pre-metastatic niche. *Cancer Cell*. **41**, 546-572(2023).
12. Turajlic S, Swanton C, Metastasis as an evolutionary process. *Science*. **352**, 169-75 (2016).
13. Vendramin R, et al, Cancer evolution: Darwin and beyond. *EMBO J*. **40**, e108389 (2021).
14. Valastyan S, Weinberg RA, Tumor metastasis: molecular insights and evolving paradigms. *Cell*. **147**, 275-292 (2011).
15. Erler J, et al, Lysyl oxidase is essential for hypoxia-induced metastasis. *Nature*. **440**, 1222-6 (2006).
16. Kannarkatt J, et al, Adjuvant Chemotherapy for Stage II Colon Cancer: A Clinical Dilemma. *J Oncol Pract*. **13**, 233-241 (2017).
17. Koo M, et al, Conceptual Framework to Guide Early Diagnosis Programs for Symptomatic Cancer as Part of Global Cancer Control, *JCO Glob Oncol.* **7**, 35-45 (2021).
18. Nowell P, The clonal evolution of tumor cell populations. *Science*. **194**, 23-28 (1976).
19. Ring A, et al, Biology, vulnerabilities and clinical applications of circulating tumour cells. *Nat Rev Cancer*. **23**, 95-111(2023).
20. Seferbekova Z, et al, Spatial biology of cancer evolution. *Nat Rev Genet*. **24**, 295-313 (2023).
21. Friedl P, Wolf K, Tumour-cell invasion and migration: Diversity and escape mechanisms. *Nat Rev Cancer*. **3**, 362-374 (2003).
22. Huerta S, et al, Colon cancer and apoptosis. *Am J Surg*. **191**, 517-26 (2006).
23. Labianca R, et al, Colon cancer. *Crit Rev Oncol Hematol*. **74**, 106-33 (2010).
24. Visser K, et al, The evolving tumor microenvironment: From cancer initiation to metastatic outgrowth. *Cancer Cell*. **41**, 374-403 (2023).
25. Gerstberger S, et al, Metastasis. *Cell*. **186**, 564-1579 (2023).
26. Iqbal A, George TJ, Randomized Clinical Trials in Colon and Rectal Cancer. *Surg Oncol Clin N Am.* **26**, 689-704 (2017).
27. Schiffman J, et al, Early detection of cancer: past, present, and future. *Am Soc Clin Oncol Educ Book*. 57-65 (2015).

28. Jeong J, et al, PMCA2 regulates HER2 protein kinase localization and signaling and promotes HER2-mediated breast cancer. *Proc Natl Acad Sci USA*. **113**, E282-90 (2016).
29. Street A, et al, Mutations in a plasma membrane Ca2+-ATPase gene cause deafness in deafwaddler mice. *Nat Genet*.**19**, 390-4 (1998).
30. VanHouten J, et al (2010). PMCA2 regulates apoptosis during mammary gland involution and predicts outcome in breast cancer. *Proc Natl Acad Sci USA*. **107**, 11405-11410 (2010).
31. FDA: https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfRL/rl.cfm?lid=740305&lpcd=MVU (2020)
32. Gerstberger S, et al, Metastasis. *Cell*. **186**, 564-1579 (2023).
33. Hebert D, et al, Dissecting metastasis using preclinical models and methods. *Nat Rev Cancer*. **23**, 391-407 (2023).
34. Lawrence R, et al, Circulating tumour cells for early detection of clinically relevant cancer. *Nat Rev Clin Oncol*. **20**, 487-500 (2023).

Nolan E, et al, Deciphering breast cancer: from biology to the clinic. *Cell.* **186**, 708-1728 (2023).

Acknowledgements and funding sources:
This work was funded by: Cytocapsula Research Institute Fund (T. Y.) and Centiver Ltd Fund (T. Y. and G.W.). This work was supported by Cellmig Biolabs Inc. All data produced in the present work are contained in the manuscript. We thank Centiver Ltd team and Cellmig Biolabs Inc. team for the help on CCT analyses. We thank Dr. Steven Gygi of Harvard Medical School for the help of proteome analyses. We greatly acknowledge Dr. Ed Harlow of Harvard Medical School (USA) and Cancer Institute of University of Cambridge (UK), Dr. Nahum Sonenberg of McGill University of Canada, Dr. Michael Roehrl of Beth Israel Deaconess Medical Center (BIDMC), Dr. Stephen Ganshirt of Northwest University Medicine Hospital, Dr. Joan Brugge of Harvard Medical School, for their help and meaningful discussion in the study.

Author contributions: T.Y. designed research; T.Y. Resources; T.Y. performed research; T.Y. supervision and project administration; T.Y. and G.W. analyzed data; T.Y. and G. W. wrote the manuscript.

Conflict of interest statement: The authors are cofounders of and hold equity in Centiver Ltd and Cellmig Biolabs Inc. to support future basic research on cytocapsular oncocells and cytocapsular tumor network systems.

Figure legends:

Fig. 1. Cytocapsular tubes elongate along the path with least resistances in tissues *in vivo*.

(**A**) Representative immunohistochemistry (IHC) fluorescence microscope image of a single pancreas cytocapsular oncocell generated, a part of a single, super-long CCT with many short, near waves/curves and super-large CCT superstructure in a superlarge, bent, oval-shaped morphology in heterogeneous pancreas ductal adenocarcinoma tissue in the same plane. The macro resistances from the above and lower dense tissues force the super-long pancreas CCT to form a bent, oval-shaped superlarge morphology. The framed area is enlarged and shown in (B). CCT (white arrow) is shown. (**B**) An enlarged CCT fragment from (A). This CCT fragment has many CCT near waves and irregular curves with short wave/curve height. Each pancreas CCT turn (orange arrows show crest/upper turns, white arrows show trough/bottom turns) is caused by

the local micro resistances from the dense tissues along the pancreas CCT elongation direction. Then, the pancreas CCT nanoprotrusion (NP) with least resistance enlarge and form pancreas enlarged NP, followed by pancreas initial CCT generation in the least resistance, and established pancreas CCT formation in the least resistance. Therefore, form a pancreas CCT turn. An inserted enlarged near CCT wave panel shows the crest, trough, wave height and wave length (λ) of CCT near wave. (**C**) Representative IHC fluorescence microscope image of invasive breast carcinoma in acytocapsular mass-CC/CCT complex (AMCC) with many sectioned CCT fragments with CCT turns. (**D**) Quantitative statistical analysis of number of sectioned CCT fragments with CCT turns/curves of clinical breast carcinoma tissues. Scale bar, 10μm.

Fig.2. Cytocapsular tube bifid branching morphogenesis significantly increase CCT number and promote CCT conducted cancer metastasis. (**A**) Representative IHC fluorescence microscope images of bifid CCT branching morphogenesis format in breast and pancreas cancer tissues in vivo. Panel A1, a breast cancer stem CCT (SC) with two branch CCTs (branch 1 (B1) and branch 2 (B2)). Panel A2, two breast cancer stem CCTs (CSs) and each CS with two branch CCTs. Panel 3, a pancreas stem CCT with two branch CCTs. Panel 4, a breast cancer stem CCT (SC) with two branch CCTs (B1 and B2), and branch 1 (B1) continuously develops into two CCT branches (B1a and B1b). (**B**) Representative IHC fluorescence microscope images of invasive breast carcinoma in 4 AMCC stages: stage I, II, III and IV: panel B1 (AMCC stage I), panel B2 (AMCC stage II), panel B3 (AMCC stage III), and panel B4 (AMCC stage IV). CCT branching morphogenesis significantly increase CCT numbers, and promote acytocapsular oncocells invade into CCTs. via alloentry followed by CCT-conducted cancer ell metastasis. Scale bar, 10μm. (**C**) Quantitative statistical analysis of occupied areas of

breast cancer CCT/CCT strands, acytocapsular oncocells, intercellular spaces/others (such as micro blood vessels) in the 4 AMCC stages. (**D**) Quantitative statistical analysis of occupied areas of colon cancer CCT/CCT strands, acytocapsular oncocells, intercellular spaces/others (such as micro blood vessels) in the 4 AMCC stages.

Fig. 3. CCT regeneration in AMCC promotes new cycles of CCT-conducted cancer metastasis.

(**A**) Representative breast invasive carcinoma in AMCC with CCT Degradation left cavity Strip (CDS, white arrows) interconnection formed CDS networks. There are many CCT degradation left cavity strips (CDSs) in variable widths crossing in the dense acytocapsular oncocell masses in AMCC, and separate dense acytocapsular oncocell (AO) mass into many dense AO mass islands. The CCTs degrade into cloud-like CCT strands (smear green cloud in the image) followed by disappearance. The framed areas (1, 2, and 3) are enlarged and shown in the panel (B, C, D), respectively. (**B**) Enlarged framed area 1 in (A). Many newly regenerated ICCTs (purple arrows) and CCTs (orange arrows) elongate along and fill the previous CCT degradation left cavity strips (CDSs, white arrows). (**C**) Enlarged area 2 in (A). Massive newly regenerated CCTs fill CDSs (white arrows) and many cytocapsular oncocells invade into regenerated CCTs and leave aways. There are some regenerated CCTs in degradation into CCT strands (CTSs, cyan arrows). (**D**) Enlarged area 3 in (A). Massive newly regenerated CCTs fill CDSs, in which regenerated CCTs tightly contact with dense AO islands, and allow AOs invade into CCTs via alloentry, followed by continuous cancer metastasis in AMCC. Scale bar, 10μm.

Fig. 4. CCT mega superstructures promote CCT-conducted cancer metastasis in AMCC stage *in vivo*. (**A**) Representative IHC fluorescence image of a super-large breast CCT mega superstructure complex with two-layered, irregular triangle-shaped, and highly curved CCT superstructures with many and dense CCT curves/coils of each CCT superstructure. There are two irregular-shaped cytocapsular tumors (CTs, white arrows) in the CCT mega superstructure. Many acytocapsular oncocells (AOs) are left via mega CCT superstructure conducted metastasis. Some CCTs (orange arrows) degrade into CCT strands (CTSs, purple arrows). (**B**) Representative image of super-large, many-layered, highly dense CCT formed mega CCT superstructures in the late breast carcinoma AMCC stage. Most acytocapsular oncocells (AOs) are left via mega CCT (orange arrows) superstructure conducted metastasis. CCT mega superstructures promote complete cancer metastasis of AMCCs. Scale bar, 10μm.

Nanoprotrusion enlargement conducts cytocapsular tube elongation along the path of least resistance

Tingfang Yi[1,2]* and Gerhard Wagner[1,2,3]*

4. Cytocapsula Research Institute, 245 First Street, Cambridge, MA, 02142 USA
5. Centiver Ltd., 245 First Street, Cambridge, MA, 02142 USA
6. Department of Biological Chemistry and Molecular Pharmacology of Harvard Medical School, 240 Longwood, Boston, MA, 02115 USA

*To whom correspondence may be addressed. E-mail:tingfang_yi@centiver.com, or

Gerhard_Wagner@hms.harvard.edu.

Materials and Methods

Reagent, Antibodies, and Devices.

CC/CCT culture kits (Celldevi, https://www. celldevi.com, Cat. CD0104, six-well plates; Cat. CD0105, 12-well plates, and Cat. CD 0106, 24-well plates) and kits with glass cover slips in the well bottom and embedded by CC/CCT culture matrix layer (Cat. CD 0112, six-well plate) were ordered from Celldevi Inc. CC/CCT culture kit fixation kit (Cat. CD0201) were ordered from Celldevi Inc. CT-M2 fluorescence microscope (with high resolution, high signal/noise ratio, low background noise) and CCT analysis software were ordered from Celldevi. Cancer cell lines of pancreas cancer cell Bxpc3, breast cancer cell MCF-7 and colon cancer cell SK-CO-1 and cell culture media were ordered from ATCC. 13C6, 15N2-L-Lysine (Cat. 88209) and 13C6, 15N4-L-Arginine (Cat. 89990) for SILAC labeling were ordered from Thermo Fisher Scientific.

UnipickTM and capillary units were ordered from NeuroInDx Inc. Rabbit anti-PMCA2 antibody (polyclonal, ab3529; 1:200 dilution), mouse anti-γ-actin antibody (monoclonal, ab123034; 1:200 dilution) for immunofluorescence assay were ordered from Abcam. DAPI (1:1,000 dilution in the immunofluorescence assay) was ordered from KPL. Human normal and cancer tissue specimens described here were ordered from Tissue Array and US Biolabs and were de-identified prior to use.

Cytocapsular tumorsphere and cytocapsula growth *in vitro*, immunohistochemistry staining and imaging.

Pancreas cancer Bxpc3 cells were implanted in CC/CCT culture kit (Cat. CD 0112, Celldevi) following the kit manual. At 36h, Bxpc3 cells generated cytocapsulas (CCs). Some cytocapsular oncocells performed ecellulation. Cytocapsular oncocells and ecellulated CCs were performed fixation kit and immunohistochemistry staining. At different time of 48h, 72h, 68h, 74h, 78h, 84h, 96h, 108h after cell implantation, Bxpc3 cancer cells engender cytocapsular oncocells, and grow into cytocapsular tumorspheres in different sizes with CC tightly wrapping oncocell mass or with wide cytocapsular lumens, and ecellulation of cytocapsular tumorspheres. These cytocapsular tumorspheres and ecellulated cytocapsular tumorspheres were fixed by Celldevi Inc. CC/CCT fixation kit (Celldevi, CD0201) in the 6-well plate, and then taken out and put onto slides, followed by immunohistochemistry (IHC) staining.

Cytocapsular tumorsphere and cytocapsula growth *in vitro*, immunohistochemistry staining and imaging

IHC staining was performed with rabbit anti-PMCA2 polyclonal primary antibodies (1:200 dilution), mouse anti-γ-actin monoclonal primary antibodies (1:200 dilution), Goat anti-Mouse

IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 555 (Thermo Fisher), and Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 488, Thermo Fisher), and DAPI staining (1:000 dilution). Fluorescence images were taken with a Nikon 80i upright microscope with a 20× or 40× lens. All images were obtained using MetaMorph image acquisition software and were analyzed with ImageJ software.

CCT Histology and Immunohistochemical Staining Analysis

The 9972 formalin-fixed, paraffin-embedded (FFPE) human cancer tissue specimens (4-5μm in thickness) from 9784 cancer patients, 14 human normal tissue FFPE specimens form 14 patients, and 126 human benign tumor tissue FFPE specimens from 126 patients were processed immunohistochemistry and hematoxylin and eosin (H&E) staining. Immunohistochemical fluorescence tests were performed to stain cytocapsular tubes using rabbit anti-PMCA2 polyclonal primary antibodies (1:200 dilution), mouse anti-γ-actin monoclonal primary antibodies (1:200 dilution), Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 555 (Thermo Fisher), and Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 488, Thermo Fisher), and DAPI staining (1:000 dilution). Fluorescence images were taken with a Nikon 80i upright microscope with a 20× or 40× lens. All images were obtained using MetaMorph image acquisition software and were analyzed with ImageJ software.

Imaging Acquisition.

DIC and fluorescence images of fixed cells (with or without cytocapsulae) were taken with an 80i upright microscope and a digital Hamamatsu ORCA-ER cooled CCD camera with a 20× or

40× lens. The bright-field phase-contrast image was taken using a Nikon digital camera. The cytocapsula initiation ratio per high-performance field (HPF; 200×) and the number of elongated cytocapsulae per high-performance field were quantified. All images were obtained using MetaMorph image acquisition software and were analyzed with ImageJ software.

Transmission Electron Microscope.

Matrigel matrix layers were prepared on plastic discs in six- well plates. HMEC cultures with cytocapsulae in Matrigel matrix (>40 μm in depth) were fixed with 1:1 mixtures of formaldehyde- glutaraldehyde-picric acid fixative (2.5% paraformaldehyde, 5.0% glutaraldehyde, 0.06% picric acid in 0.2 M cacodylate buffer):cell culture medium. The cells and cytocapsulae were then postfixed for 30 min in 1% osmium tetroxide (OsO4)/1.5% potassium fer- rocyanide (KFeCN6), washed three times in water, and incubated in 1% aqueous uranyl acetate for 30 min followed by two washes in water and subsequent dehydration in grades of alcohol (50, 70, 95, and 2 × 100%; 5 min each) (1). Cells and cytocapsulae were infiltrated for 2 h to overnight in a 1:1 mixture of propylene oxide and TAAB Epon (Marivac Canada Inc.). The samples were subsequently embedded in TAAB Epon and polymerized at 60 °C for 48 h. Ultrathin sections (∼60 nm) were cut on a Reichert Ultracut S microtome, picked up on copper grids, stained with lead citrate, and examined in a JEOL 1200EX Transmission electron micro- scope or a Tecnai G2 Spirit BioTWIN microscope, and images were recorded with an AMT 2k CCD camera.

Nanoprotrusion bunches in bright filed and fluorescence microscope images *in vitro*
Nanoprotrusions (NPs, 21nm ~ 82nm in diameter/width) of CCs/CCTs are in visible in bright field and fluorescence microscope due to the limited resolution of these microscopes. The NP

bunches with big sizes (100nm ~ 152nm in width) are visible in high resolution bright filed microscope images. The NP bunches with big sizes (100nm ~ 152nm in width) and coated with thin or no glycocalyx layers are visible in CT-M2 fluorescence microscope images with high resolution, high signal/noise ratio and low background (but invisible in common fluorescence microscope).

Initial cytocapsular tube (ICCT) and CCT analysis in cancer tissues *in vivo*

In cancer tissues, initial cytocapsular tubes (ICCTs) are small (0.2μm ~ 0.5μm in diameter/width) straight/curved cytocapsular tubes without degradation, with or without long and thin cancer cells in migration inside. Cytocapsular tubes (CCTs) are established cytocapsular membrane enclosed tubes (0.5μm ~ 10μm in diameter/width). CCT degraded strands show phenomena of: networks of very thin strands, dot array collapse membrane degradation, cloud line-status.

Data collection

Cytocapsular tubes (CTs, not sectioned, longitudinally sectioned, and cross sectioned) without degradation (3~6μm in measured diameter) were counted using a fluorescence microscope and ImageJ. The presence of CTs degrading into thick strands (1~2μm in measured diameter), thin strands (0.2~1μm in measured diameter), or the disintegration state were reported without quantification. The patients providing formalin-fixed paraffin-embedded tissue samples gave informed consent that they understood that the biopsies (needle biopsy or surgical biopsy, from US Biomax) were performed for *in vitro* research purposes only. Comparative deidentified samples of normal tissues, benign tissues, carcinoma *in situ*, cancer, paracancer, metastatic tissues with their cancer stages identified according to the tumor (T), node (N), and metastasis

(M) TNM system (cancer stages: 0, I, II, III, IV) were obtained from archival materials (Tables S3-4). The cancer, paracancer and metastatic cancer tissues, in which the original cancer niches were identified by indicated cancer specific molecular markers, were identified by hospital pathology laboratories and obtained from archival material. Autopsy tissues samples have many post-life CC/CCT degradation and CCs/CCTs will not be quantified. Biopsy samples from FFPE with fresh tissues present high fidelity of CC/CCT status and CCs/CCTs are quantified and reported.

Quantification and Statistical Analysis.

The statistical methods used for comparisons are indicated in the relevant Fig. legends and in the sections below. The diameters, widths, and lengths of cytocapsulae and cytocapsular tubes were measured with MetaMorph or ImageJ. The time of individual cytocapsulae and cytocapsular tubes was counted from cytocapsula generation to acellular cytocapsula (or cytocapsular tube) decomposition. For lifetime of cytocapsular oncocell and cytocapsular tumorsphere assays, at least 20 cytocapsular oncocell or cytocapsular tumorsphere were measured per condition, and two-tailed Student's test was used to determine statistical significance. In cytocapsular tube quantitation assays, for each specimen, the number of fully intact cytocapsular tubes was counted in 5 areas (0.35mm x 0.35mm, length x width) of the sample (top, bottom, left, right, and center), and the cytocapsular tube density (CCT/mm2) was calculated and determined for each area. The average CCT density across the 5 sites was treated as the specimen's overall CCT density and round up to digits.

Extended Data Figures and Legend:

Extended data Fig.1. Cytocapsular oncocells and tumorspheres generate cytocapsular nanoprotrusions (NPs), and ultrastructure of cytocapsular nanoprotrusions (NPs) and NP bunches. (A) Representative bright field (BF) and immunohistochemistry (IHC) fluorescence microscope images of a single cytocapsular oncocell (CO) with cytocapsular nanoprotrusions (NPs) at 24h after Bxpc3 pancreas cancer cell implantation in 3D CC/CCT culture kits. A schematic diagram of single COs with NPs and NP bunches. **(B)** Representative bright field and IHC fluorescence microscope images of ecellulated cytocapsulas (ECs) with NPs and NP bunches of single Bxpc3 cytocapsular oncocells in 3D CC/CCT culture kits. The ECs collapse, and form deflated, concave discs. NP networks (NNs) interconnect with and surround the acellular cytocapsulas. A schematic diagram of single ecellulated cytocapsula with NPs and NP bunches. **(C)** Representative bright field and IHC fluorescence microscope image of a Bxpc3 pancreas cancer cytocapsula tumorsphere with NPs and CNP bunches. A schematic diagram of single cytocapsular tumorsphere with CNPs and CNP bunches. The cytocapsula (CC, orange arrows), cytocapsular nanoprotrusion (NP), cytocapsular nanoprotrusion bunches (NB, white arrows), cytocapsular nanoprotrusion network (NN, purple arrows), incytocapsular oncocell (ICO, purple arrows) are shown. Scale bar, 10μm. **(D)** Representative transmission electron microscope (TEM) image of longitudinal section of a single cytocapsular nanoprotrusion (NP) fragment at 42h after breast cancer stem cell (BCSC) HMLER(CD24[low]/CD44[high])[FA] subpopulation cell implantation in 3D CC/CCT culture kits. The NPs are coated with glycocalyx (GC, black arrows) layers in variable thickness. Cytocapsular nanoprotrusion (NP, red arrows), cytocapsular nanoprotrusion membrane (NPM, blue arrows) are shown. Scale bar, 250nm. **(E)** Representative TEM image of longitudinal section of cytocapsular nanoprotrusion bunches (NBs,

red arrows). Multiple NPs align together tightly or loosely and form NBs in curved morphologies in thick glycocalyx layers. Scale bar, 100nm. The enlarged NBs are show in the right two panels: Panel E1, longitudinal section of a stem NP with two branches (B1 and B2) with an open interconnection between the stem CNP and the two NP branches; Panel E2, longitudinal sections of two NPs align together closely with the 4 sectioned and exposed NP membranes, which are embedded in glycocalyx layer. Asterisks in cyan and black colors: longitudinal sectioned lumens of the two NP lumens. Scale bar, 50nm.

Extended data Fig.2. CCT nanoprotrusions (NPs) grow into enlarged NPs, and develop into initial CCTs in vitro. (**A**) Representative bright field and IHC fluorescence microscope images show that cytocapsular nanoprotrusions (NPs) with shortest distance and least resistance between ecellulated cytocapsulas (ECCs) of cytocapsular tumorspheres (CTs) enlarge, and grow into enlarged cytocapsular nanoprotrusions (ECNs, white arrows). (**B**) Representative bright field (BF) and IHC fluorescence microscope images show that in initial cytocapsular tubes (ICCTs), single oncocells migrate inside of ICCT, and make enlarged CCT fragments behind. A single oncocell ecellulated, and left an enlarged and oval-shaped CCT membrane bulge. The enlarged CCT membrane bulge is enlarged in panel B1' and B2'. Purple arrows show the open connection between the CCT bulge front and ICCT. (**C**) Representative bright field (BF) and IHC fluorescence microscope images show that a long, straight CCTs interconnect cytocapsular tumorspheres (CTs) with shorted distance and least resistance. There are many NPs surrounding the CCTs. Scale bar, 10μm.

Extended data Fig.3. Initial CCTs present in cancer tissues in vivo.

(**A**) Panel A1, a representative IHC fluorescence microscope image of invasive breast cancer post nuperphase tumor with many regenerated initial CCTs. The framed area is enlarged and shown in panel A2. Panel A2, there are many long or short, straight or highly-curved initial CCTs are regenerated in this area. The framed two areas of 2a and 2b are enlarged and show in the panels A3 and A4, respectively. Panel A3, there are several long, straight or curved, regenerated initial CCTs (ICCTs, white arrows), and some breast cancer cells are in migration in ICCTs (orange arrows). Panel 4, there are many regenerated initial CCTs (ICCTs, white arrows), and some breast cancer cells are in migration in ICCTs (orange arrows). There are a few established cytocapsular tubes (CCTs, cyan arrow). (**B**) In a representative image of in situ breast carcinoma, there already are large quantities of initial CCTs in high ICCT (white arrows) density. (**C**) In a representative image of clinical colon carcinoma, there are some ICCs (white arrows) in colon cancer tissues. Scale bar, 10μm.

Extended data Fig.4. Characterization of initial cytocapsular tubes, curved cytocapsular tube fragments, CCT branching morphogenesis, CCT regeneration, and CCT superstructures in 289 types/subtypes of cancers in human organs and tissues.

Extended data Fig.5. Single super-long pancreas cytocapsular tubes elongate along the path with least resistance in heterogeneous pancreas tissues and form superstructures with many curves *in vivo*. (**A**) Representative IHC fluorescence microscope image of a single pancreas cytocapsular oncocell generated, single, super-long CCT with many short/middle length curves and super-large CCT superstructure in a superlarge bent morphology in heterogeneous pancreas ductal adenocarcinoma tissue in the same plane. The macro resistances from the above

and lower dense tissues force the super-long CCT to form a superlarge bent morphology. The framed area is enlarged and shown in (B). (**B**) An enlarged CCT fragment from (A). This CCT fragment has many irregular-shaped, short/middle length curves. Each pancreas CCT curve turn (white arrows) is caused by the local micro resistances from the dense tissues along the pancreas CCT elongation direction. Then, the pancreas CCT nanoprotrusion (NP) with least resistance enlarge and form enlarged NP, followed by pancreas initial CCT generation in the least resistance, and established pancreas CCT formation in the least resistance. Therefore, form a pancreas CCT turn. The orange asterisks show less or no tissue (with less or no tissue resistance) below the CCT turns. Scale bar, 10μm.

Extended data Fig.6. Single super-long prostate cytocapsular tubes elongate along the path with least resistance in heterogeneous prostate tissues and form superstructures with many long curves *in vivo*. (**A**) Representative IHC fluorescence microscope image of a single prostate cytocapsular oncocell generated, single, super-long CCT with many short and long curves and super-large CCT superstructure in a superlarge and irregular bent morphology in heterogeneous prostate adenocarcinoma tissue in the same plane. The macro resistances from the above and lower dense tissues force the super-long CCT to form a bent superlarge morphology. The framed area is enlarged and shown in (B). CCT (white arrows) are shown. (**B**) An enlarged CCT fragment from (A). This CCT fragment has many irregular-shaped, long curves. Each prostate CCT curve turn (white arrows) is caused by the local micro resistances from the dense tissues along the prostate CCT elongation direction. Then, the prostate CCT nanoprotrusion (NP) with least resistance enlarge and form prostate enlarged NP, followed by prostate initial CCT generation in the least resistance, and established prostate CCT formation in the least resistance.

Therefore, form a prostate CCT turn. The orange asterisks show less or no tissue (with less or no tissue resistance) below the CCT turns. CCT turns (white arrows), and CCT strand (CTS, orange arrows) are shown. Scale bar, 10μm.

Extended data Fig.7. Single super-long colon cytocapsular tubes elongate along the path with least resistance in heterogeneous colon tissues and form superstructures with many curves *in vivo*. (**A**) Representative IHC fluorescence microscope image of a single colon cytocapsular oncocell generated, single, super-long colon CCT with many middle-length curves and super-large CCT superstructure in a superlarge and irregular waved morphology in heterogeneous colon adenocarcinoma tissue in the same plane. The macro resistances from the above and lower dense tissues force the super-long colon CCT to form highly-curved superlarge morphologies. The framed area is enlarged and shown in (B). CCT (white arrows) are shown. (**B**) An enlarged colon CCT fragment from (A). This colon CCT fragment has many irregular-shaped, middle-length curves. Each colon CCT curve turn (white arrows) is caused by the local micro resistances from the dense tissues along the colon CCT elongation direction. Then, the colon CCT nanoprotrusion (NP) with least resistance enlarge and form colon enlarged NP, followed by colon initial CCT generation in the least resistance, and established colon CCT formation in the least resistance. Therefore, form a colon CCT turn. CCT turns (white arrows) are shown. Scale bar, 10μm.

Extended data Fig.8. Schematic diagram of nanoprotrusion enlargement conducted cytocapsular tube elongate along the path with least resistance.

There are 6 successive major steps in cytocapsular tube elongation in heterogeneous tissues *in vivo*: (**1**) Transformation: normal cell transformation generates acytocapsular oncocell (CO);

(**2**) Devolution: acytocapsular oncocell devolution engenders cytocapsular oncocell with extracellular cytocapsula (CC) wrapping AO inside. The CC isolates AO from external stressful or toxic microenvironments and help AO for survival;

(**3**) Nanoprotrusion (NP) generation: cytocapsular oncocells (COs) generate many nanoprotrusions (NPs, 21nm ~ 82nm in diameter/width, 300-2700nm in length), NP bunches (40nm ~ 152nm in width), which interconnect and form NP networks surrounding the CO. Intracytocapsular oncocells proliferate in CCs and generate cytocapsular tumors (CTs). CTs generate NP networks surrounding CTs. NPs are coated with glycocalyx in variable thickness;

(**4**) Enlarged nanoprotrusion (ENP) generation: the NPs in least resistance enlarge and generate enlarged NPs (ENPs, 82nm ~ 0.1μm);

(**5**) Initial CCT (ICCT) generation: the ENPs continue to growth in size and length and generate initial CCTs (ICCTs, 0.1μm ~ 0.5μm in diameter/width).

(**6**) CCT generation and elongation: Single intracytocapsular oncocell squeezes into ICCTs, and promote ICCTs to continue to grow up in sizes and length and engender established CCTs (>0.5μm in diameter/width), therefore, CCT is elongated. CCT elongation is conducted by CCT nanoprotrusion enlargement and along the path with least resistance.

Extended data Fig.9. Cytocapsular tube elongate in the least resistance path lead to CCT conducted cancer metastasis occurrence much prior to CCT invasion into blood vessel and oncocell release into blood. (**A**) Representative image of clinical breast cancer tissues with many CCTs (white arrows) in the late CCT lifecycle stage of CCT degradation into CCT strands

(CTSs, orange arrows). Most of breast acytocapsular oncocells (AOs) invade into breast CCTs and leave away, and there are only a few AOs left in this area. There are multiple micro blood vessels (MBVs, red arrows) in intact status without CCT invasion caused oncocell release into the blood in this area. (**B**) Representative image of clinical colon cancer tissues with many colon CCTs (white arrows) with many colon oncocells in migration in CCTs (purple arrows). Most of colon acytocapsular oncocells (AOs) invade into colon CCTs and leave away, and there are almost no colon AOs left in this area. There are multiple micro blood vessels (MBVs, red arrows) in intact status without CCT invasion caused oncocell release into the blood in this area. (**C**) Representative image of clinical prostate cancer tissues with many prostate CCTs (white arrows) in the late CCT lifecycle stage of CCT degradation into CCT strands (CTSs, orange arrows). Most of prostate acytocapsular oncocells (AOs) invade into prostate CCTs and leave away, and there are almost no prostate AOs left in this area. There are blood vessels (BV) and micro blood vessel (MBV, red arrow) in intact status without CCT invasion caused oncocell release into the blood in this area. (**D**) Representative image of clinical breast cancer tissues with large CCTs (white arrows) networks in the late stage. Most of breast acytocapsular oncocells (AOs) invade into breast CCT networks and leave away, and there is no breast AOs left in this area. There are a micro blood vessel (MBV, red arrows) in intact status without CCT invasion caused oncocell release into the blood in this area. Scale bar, 10μm.

Extended data Fig.10. CCT branching morphogenesis in side NP enlargement conducted bifid branching morphogenesis format in vitro. (**A**) Representative bright field (BF) and fluorescence microscope images of CCT branching morphogenesis CC/CCT culture kits *in vitro*. At 72h after Bxpc3 pancreas cancer cell implantation in CC/CCT culture kits, pancreas

cytocapsular tumorspheres (CTs), multiple CCTs interconnect CTs, and CCT branches are generated. The framed area in BF and fluorescence microscope images are enlarged and shown in (B). Scale bar, 10μm. (**B**) Enlarged BF and fluorescence microscope images of framed areas in (A). The stem CCT (SC) side nanoprotrusions in least resistance generate enlarged NPs, followed by initial CCT (ICCT) and established CCT generation, and engender a branch CCT. After CCT membrane remodeling, the branch CCT and stem CCT are remodeled into two near similar CCT branches (branch 1 (B1) and branch 2 (B2), orange arrows). This kind of CCT branching morphogenesis format is named as bifid branching morphogenesis format. Scale bar, 10μm. (**C**) Schematic diagram of CCT bifid branching morphogenesis format, which contains 4 successive major steps: (1) Side NP enlargement: CCT side NPs in least resistance enlarge and generate branch enlarged NPs (branch ENPs); (2) Branch initial CCT (ICCT) generation: Branch ENPs continue to grow up into branch ICCT; (3) Branch CCT formation: With intracytocapsular oncocell riven ICCT growth in size and length, branch ICCTs continue to grow up into branch CCTs. (4) CCT remodeling: the connected cytocapsular membranes of stem CCT and branch CCT are remodeled into two similar or near similar CCTs (in diameter/width) and interconnecting with the stem CCT, and form two branch CCTs.

Extended data Fig.11. CCT bifid branching morphogenesis significantly increase CCT numbers and promote cancer metastasis in tissues *in vivo*. (**A**) Representative IHC fluorescence image of breast invasive carcinoma in the early CCT regeneration stage I in acytocapsular oncocell mass-CC/CCT complex (AMCC stage I). The framed area is enlarged and shown in (B). (**B**) Enlarged framed area in (A). In the dense acytocapsular oncocell mass,

regenerated initial CCTs (ICCTs) and CCTs perform bifid branching morphogenesis and increase CCT numbers. In AMCC stage I, ICCTs/CCTs occupied areas are 0-10% in areas in the AMCC. Multiple CCT branches (CBs, orange arrows) and ICCTs (white arrows) are shown. The two framed areas (area 1 and area 2) with CCT branches are enlarged and shown in Figure 2A1 and 2A2 panels, respectively. (**C**) Representative IHC fluorescence image of breast invasive carcinoma in AMCC stage II. The framed area is enlarged and shown in (D). (**D**) Enlarged framed area in (C). In AMCC stage II, ICCTs/CCTs/CCT strands occupied areas are 10%-40% in areas in the AMCC. (**E**) Representative IHC fluorescence image pancreas adenocarcinoma in the AMCC stage III. The framed area is enlarged and shown in (F). (**F**) Enlarged framed area in (E). In AMCC stage III, ICCTs/CCTs/CCT strands occupied areas are 40%-70% in areas in the AMCC III. The framed area with CCT branch is enlarged and shown in Fig2A3 panel. (**G**) Representative IHC fluorescence image of breast invasive carcinoma in the AMCC stage IV. The framed area is enlarged and shown in (H). (**H**) Enlarged framed area in (G). In AMCC stage IV, ICCTs/CCTs/CCT strands occupied areas are 70%-100% in areas in the AMCC III. CCTs (white arrows) are shown. With CCT number significantly increase (from A to C, to E, to G, from AMCC stage I to stage II, stage III, and stage IV), more and more acytocapsular oncocells (AOs) invade into CCTs via alloentry, and leave away, and less and less AOs left. Scale bar, 10μm.

Extended data Fig.12. CCT regeneration associates with new cycle of cancer metastasis in AMCC.

(**A**) Representative invasive ductal breast carcinoma in AMCC with many CCT degradation left cavity strips (CDSs) interconnected CDS networks. There are many CCT degradation left cavity

strips (CDSs) in variable widths crossing in the dense acytocapsular oncocell masses, and separate dense acytocapsular oncocell (AO) mass into many dense AO mass islands. The CCTs degrade into cloud-like CCT strands (smear cloud in green color in the image) followed by disappearance. The framed areas (1, 2, and 3) are enlarged and shown in the panel (B, C, D), respectively. (**B**) Enlarged area 1 in (A). Many new CCTs are regenerated and elongate in the previous CCT degradation left cavity strips (CDSs). Most regenerated CCTs elongate in CDSs with less resistances, a few CCTs invade into dense AO mass islands. (**C**) Enlarged area 2 in (A). There are multiple CDSs interconnect and form CDS network for regenerated new CCTs to elongate. (**D**) Enlarged area 2 in (A). In the CDSs, there are many regenerated new CCTs, which tightly contact with dense AO islands, and allow AOs invade into CCTs via alloentry, followed by new cycles of cancer metastasis in AMCC. Scale bar, 10μm.

Extended data Fig.13. CCT mega superstructures in pancreas, lung and prostate cancers in vivo.

(**A**) Representative IHC fluorescence image of a part of a super-large pancreas CCT mega superstructures with three-layered, irregular triangle-shaped, and highly curved CCT superstructures with many CCT curves/coils of each CCT superstructure. (**B**) Representative image of a super-large lung CCT mega superstructures with five-layered, irregular-shaped CCT superstructures with many CCT curves/coils of each CCT superstructure. Some CCTs are in degradation into CCT strands (CTSs, orange arrows). (**C**) Representative image of a super-large prostate CCT mega superstructures with more than seven-layered, irregular-shaped CCT superstructures with many CCT curves of each CCT superstructure.

Extended data Fig.14. CCT mega superstructures promote complete cancer metastasis in AMCC stage *in vivo*. (**A**) Representative image of super-large CCT bunch interconnection formed mega CCT network superstructures in the late breast cancer AMCC stage. All acytocapsular oncocells (AOs) are left via mega CCT superstructure conducted metastasis. (**B**) Representative image of super-large, many-layered, highly curved and coiled CCT formed mega CCT superstructures in the late breast cancer AMCC stage. All acytocapsular oncocells (AOs) are left via mega CCT superstructure conducted metastasis. (**C**) Representative image of super-large, many-layered, highly dense CCT formed mega CCT superstructures in the late breast carcinoma AMCC stage. All acytocapsular oncocells (AOs) are left via mega CCT superstructure conducted metastasis. Some CCTs degrade into CCT strands (CTSs, orange arrows). CCTs (white arrows) are shown. Scale bar, 10μm.

Fig.1

A Pancreas ductal adenocarcinoma

DAPI: PMCA2: γ-actin

CCT

B Pancreas ductal adenocarcinoma

DAPI: PMCA2: γ-actin

CCT turn

CCT turn

Crest

Crest

λ

Wave height

Trough

C Invasive breast carcinoma in AMCC

DAPI: PMCA2: γ-actin

D

N= 514

Number of sectioned CCT fragments with CCT turn/curve at one or two ends of 5 specimens of clinical breast carcinoma tissues

Length (μm)

<20   20~40   40~60   60~80   >80

Fig.2

A

DAPI: PMCA2: γ-actin

1 Invasive breast carcinoma AMCC — B1, SC, B2

2 Invasive breast carcinoma AMCC — B2, B1, B1, B2, SC, SC

3 Pancreas carcinoma AMCC — B1, B2, SC

4 Invasive breast carcinoma AMCC — B1a, B1b, B1, B2, SC

B

Invasive breast carcinoma

1 DAPI: PMCA2: γ-actin

2 DAPI: PMCA2: γ-actin

3 DAPI: PMCA2: γ-actin

4 DAPI: PMCA2: γ-actin

C

- Breast cancer CCT/CCT strands
- Acytocapsular oncocell
- Intercellular space/others

Average percentage of occupied areas of acytocapsular oncocells, CCTs, and intercellular space in clinical invasive breast carcinoma of 5 pieces of tissue specimens

AMCC phases: Phase I, Phase II, Phase III, Phase IV

D

- Colon cancer CCT/CCT strands
- Acytocapsular oncocell
- Intercellular space/others

Average percentage of occupied areas of acytocapsular oncocells, CCTs, and intercellular space in clinical colon cancer of 5 pieces of tissue specimens

AMCC phases: Phase I, Phase II, Phase III, Phase IV

Fig.3

Fig.4

A

Clinical breast carcinoma

B

Clinical breast carcinoma

Extended data Fig.1

Extended data Fig.2

A Bxpc3

BF — 48h — ECC of CT, ENP, NB, ECC of CT

DAPI: PMCA2: γ-actin — ENP, NB

Schematic: ECC of CT, NP, CC, ENP, ECC of CT, NP

B Bxpc3

1 BF — 56h — CCT, ICCT, CT, ECC, ENP, CT

1' BF

2 DAPI: PMCA2: γ-actin — CT, CT

2' DAPI: PMCA2: γ-actin

Schematic: ECC of CT, NP, CC, ICCT, CC, ECC of CT, NP

C Bxpc3

BF — 72h — CCT

DAPI: PMCA2: γ-actin

Schematic: NP, CC, CT, CCT, CT, CC, NP

Extended data Fig.3

Extended data Fig.4

Characterization of initial cytocapsular tubes, curved cytocapsular tube fragments, CCT branching morphogenesis, CCT regeneration, and CCT superstructures in 289 types/subtypes of cancers in human organs and tissues.

	Cancer types by organs/tissues	*Cancer subtypes	Tissue specimen number	Patient number	With initial CCTs in all or parts of tested subtypes	With curved CCT fragments in each subtype	With CCT elongation in least resistance path in each subtype	CCT branching morphogenesis in all or parts of tested subtypes	With CCT degradation and regeneration	With CCT super-structures
1	Adrenal Gland	3	23	22	+	+	+	+	+	+
2	Bladder	7	46	46	+	+	+	+	+	+
3	Blood (Leukemia in tissues)	2	14	14		+	+		+	
4	Blood (Multiple myeloma in tissues)	2	35	35		+			+	
5	Bone	12	212	210		+	+		+	
6	Bone marrow	5	20	20		+			+	+
7	Brain	8	124	122	+	+	+	+	+	+
8	Breast	49	3960	3890	+	+	+	+	+	+
					+	+	+	+	+	+
					+	+	+	+	+	+
					+	+	+	+	+	+
					+	+	+	+	+	+
					+	+	+	+	+	+
9	Cervix	7	53	51	+	+	+	+	+	+
10	Colon	6	1586	1586	+	+	+	+	+	+
11	Esophagus	7	63	62	+	+	+	+	+	+
12	Fibrous	7	25	25	+	+	+	+	+	+
13	Gallbladder	4	26	26	+	+	+	+	+	+
14	Head/neck	5	12	12	+	+	+	+	+	+
15	Intestine	3	15	15	+	+	+	+	+	+
16	Kidney	11	123	120	+	+	+	+	+	+
17	Liver	8	362	358	+	+	+	+	+	+
18	Lung	30	683	676	+	+	+	+	+	+
19	Lymph (in tissues)	13	56	56	+	+	+	+	+	+
20	Oral cavity	6	17	17	+	+	+	+	+	+
21	Ovary	6	83	83	+	+	+	+	+	+
22	Pancreas	16	222	218	+	+	+	+		+
23	Penis	3	3	3	+	+		+		+
24	Prostate	7	1606	1528	+	+	+	+	+	+
25	Rectum	7	5	5	+	+	+	+	+	+
26	Skin	10	122	122	+	+	+	+	+	+
27	Spleen	3	6	6	+	+	+	+	+	+
28	Smooth muscle	3	20	20	+	+	+	+	+	+
29	(other) Soft tissues	12	120	106	+	+	+	+	+	+
30	Stomach	10	245	245	+	+	+	+	+	+
31	Testis	5	24	24	+	+	+	+	+	+
32	Thymus	8	5	5	+	+	+	+	+	+
33	Thyroid	3	17	17	+	+	+	+	+	+
34	Uterus	1	26	26	+	+	+	+	+	+
35	Vulva	3	13	13	+	+	+	+	+	+
	Total	**289**	**9972**	**9784**						

(* Checked cancer subtypes are listed in Table S1.)

Extended data Fig.6

A   Prostate adenocarcinoma

B   Prostate adenocarcinoma

A

Colon adenocarcinoma

B

Colon adenocarcinoma

Extended data Fig.8

Extended data Fig.9

Extended data Fig.10

Extended data Fig.11

Extended data Fig.12

Extended data Fig.13

Extended data Fig.14

A   **Clinical breast cancer in late AMCC stage**

DAPI: PMCA2: γ-actin

CCT

B   **Clinical breast cancer in late AMCC stage**

DAPI: PMCA2: γ-actin

CCT

C   **Breast carcinoma in late AMCC stage**

DAPI: PMCA2: γ-actin

CCT

CTS