Title

Identification and classification of ion-channels across the tree of life provide functional insights

into understudied CALHM channels

Authors and affiliations

Rahil Taujale[1,4†], Sung Jin Park[2†], Nathan Gravel[1], Saber Soleymani[4], Rayna Carter[4], Kennady

Boyd[4], Sarah Keuning[2], Zheng Ruan[3], Wei Lü[2,5,6*], Natarajan Kannan[1,4*]

[1]Institute of Bioinformatics, University of Georgia, Athens, GA 30606

[2]Department of Molecular Biosciences, Northwestern University, Evanston, IL 60208

[3]Department of Biochemistry & Molecular Biology, Thomas Jefferson University, Philadelphia,

PA 19107

[4]Department of Biochemistry and Molecular Biology, University of Georgia, Athens, GA 30606

[5]Department of Pharmacology, Northwestern University, Chicago, IL 60611

[6]Chemistry of Life Processes Institute, Northwestern University, Evanston, IL 60208

* Corresponding authors: nkannan@uga.edu, wei.lu@northwestern.edu

[†] These authors contributed equally to the presented work.

Abstract

The ion channel (IC) genes encoded in the human genome play fundamental roles in cellular functions and disease and are one of the largest classes of druggable proteins. However, limited knowledge of the diverse molecular and cellular functions carried out by ICs presents a major bottleneck in developing selective chemical probes for modulating their functions in disease states. The wealth of sequence data available on ICs from diverse organisms provides a valuable source of untapped information for illuminating the unique modes of channel regulation and functional specialization. However, the extensive diversification of IC sequences and the lack of a unified resource present a challenge in effectively using existing data for IC research. Here, we perform integrative mining of available sequence, structure, and functional data on 419 human ICs across disparate sources, including extensive literature mining by leveraging advances in large language models to annotate and curate the full complement of the "channelome". We employ a well-established orthology inference approach to identify and extend the IC orthologs across diverse organisms to above 48,000. We show that the depth of conservation and taxonomic representation of IC sequences can further be translated to functional similarities by clustering them into functionally relevant groups, which can be used for downstream functional prediction on understudied members. We demonstrate this by delineating co-conserved patterns characteristic of the understudied family of the Calcium Homeostasis Modulator (CALHM) family of ICs. Through mutational analysis of co-conserved residues altered in human diseases and electrophysiological studies, we show that these evolutionarily-constrained residues play an important role in channel gating functions. Thus, by providing new tools and resources for performing large comparative analyses on ICs, this study addresses the unique needs of the IC community and provides the groundwork for accelerating the functional characterization of dark channels for therapeutic intervention.

44   Keywords: ion channel classification, LLM, RAG, human ion channel orthology, CALHM1,

45   CALHM6, electrophysiology

Introduction

Ion Channels (ICs) are membrane-bound proteins critical to many physiological processes in the human body, including regulating cell volume, neurotransmitter release, muscle contraction, and glandular secretion [1-4]. Abnormal channel functions have been causally associated with "channelopathies" such as Parkinson's disease, epilepsy, cardiac arrhythmia, cancer, and cystic fibrosis, to name but a few [5-8]. The development of selective chemical probes for channels in disease states is currently hindered by the limited knowledge of the diverse gating mechanisms, ion selectivity, and pathway associations displayed by the various channels encoded in the human genome [8, 9]. While IC sequences from diverse organisms encode critical information regarding underlying functions and effective mining of sequence data can provide important context for predicting and testing understudied IC functions, traditional bioinformatic approaches have had limited success due to the challenges in consistently defining the full IC complement [10, 11], and accurately aligning and mining large sequence datasets [12]. Online resources like the collaborative platform Channelpedia [13], the Guide to Pharmacology (GtoP) database [14], structure centric ChanFAD [15] and ChannelsDB [16], and several other efforts have previously catalogued and curated ICs and provide extensive information on a number of ion channel sequences, but do not include all the human encoded ICs, nor provide information on ICs from other organisms and taxa [17-20]. Most IC studies across taxa are largely confined to closely related families or limited taxonomic groups [21, 22].

The extensive diversification and widespread abundance of ICs across cell types and their similarities to other transmembrane protein families, such as transporters, have led to an inconsistent definition of the human IC complement. For example, previous literature has reported around 230 human ICs [17]. The Kyoto Encyclopedia of Genes and Genomes (KEGG) database has cataloged around 316 human ICs [23, 24], the GtoP database lists 278 ICs [14],

while Pharos [25] and The Human Genome Organization (HUGO) [26] list around 344 and 330

proteins, respectively. Moreover, these resources classify the human ICs based on their

function, which is not known for many. Furthermore, current classification rarely accounts for the

pore-forming or auxiliary roles of ICs. Channel proteins often form large macromolecular

complexes in association with other IC and auxiliary subunits. Separate studies have highlighted

the myriad of roles that the non-conducting auxiliary subunits play in an ion channel complex

that control channel trafficking, gating and pharmacology [27-29], making them suitable

therapeutic targets for drugs to regulate ion channel activity. Unfortunately, many of the existing

resources fail to identify, include, and annotate such auxiliary channels, leading to their

exclusion from large scale analyses for clinical outcomes. Due to these inconsistencies,

downstream analyses have been hindered and limited to certain families or groups of closely

related channels. Consequently, our current knowledge of IC functions is skewed towards a

subset of well-studied "light" channels, while a significant portion of the channelome remains

understudied and is referred to as "dark" channels by the Illuminating Druggable Genome (IDG)

consortium [30, 31]. Concerted efforts such as IDG, and the Structural Genomics Consortium

[29] help address these gaps in knowledge by systematically identifying and prioritizing such

understudied members. However, it is imperative to develop a functional and an evolutionary

classification framework that provides tools for researchers to extend knowledge from light

members to their dark counterparts enabling them to make meaningful biological inferences and

shed light on their critical roles in function and disease states.

The current classification of channel genes is largely based on experimentally determined

electrophysiological parameters such as the type of ions they transport (Na^+, K^+, Ca^{2+}, Cl^-) and

the channel gating mechanism such as gating by voltage potential (voltage-gated) or ligand

binding (ligand-gated), which has been quite helpful in placing ICs in a functional context [14].

One of the first comprehensive classification of ICs was done by Jegla et. al. [17], where they

described 235 human ICs distributed in 24 families, along with their conservation across select

metazoan species, and family specific phylogeny and evolutionary analyses. A more recent

collection of ICs along with their classification can be obtained from the GtoP database [14], a

resource dedicated to providing detailed overviews of over 1900 human drug targets, spanning

six major pharmacological targets – ICs being one of them. GtoP classifies 278 human ICs into

3 major groups (Voltage-Gated Ion Channels (VGICs), Ligand-Gated Ion Channels (LGICs),

and Other) and 31 families, and provides detailed overview of these channels in a friendly

interface. However, it does not include a lot of now well-known ion channel families such as

Calcium Homeostasis Modulator channels (CALHMs) or Bestrophins (Best). Additionally, the

resource being a general drug target resource, is not IC-centric, and lacks evolutionary and

residue level functional insights that might help hypotheses generation. The role or annotation

for auxiliary channels, that play critical roles in function and regulation of ICs [27, 32] have not

been addressed in these resources either.

On the other hand, cryo-electron microscopy (cryo-EM) studies and analysis of IC structures

have shown that despite extensive variation in primary sequences, several channels adopt

common structural folds and mechanisms of action [33]. For example, several ICs - Potassium

(K+), Sodium (Na+), Calcium (Ca2+), Transient Receptor Potential (TRP), and

Hyperpolarization-activated Cyclic Nucleotide-gated (HCN), grouped together as voltage-gated-

like (VGL), are proposed to have evolved from a common ancestor because they share key

sequence and structural features [34, 35]. Thus, quantitative comparisons of these commonly

conserved regions at the sequence and structural level can provide new residue-level insights

into IC function and a stronger basis for functional classification, much like in other large protein

families such as kinases and glycosyltransferases [36-39]. This is imperative since selective

targeting of channels in diseases and a mechanistic understanding of disease-associated

mutations in channelopathies will require a residue-level understanding of function-determining

features in both well-studied "light" and understudied "dark" channels. Thus, there is a critical

need for an updated classification that places ICs in functionally, and evolutionarily related

groups, enabling a deeper residue level analysis of light and dark channels alike.

To develop a comprehensive classification of ICs, we first mined literature and sequence data

sources to collect information on ICs stored across various databases and in disparate data

sources and formats to build a well-curated knowledgebase towards defining the human ion

"channelome." A comprehensive list encompassing established and putative IC sequences in

the human proteome is provided as a unified table alongside other contextual data for functional

annotations such as regulatory interactions, ion-selectivity, and pore-lining residues. Leveraging

the identification of a pore region and pore-lining residues, we make a clear distinction between

pore-containing versus auxiliary ICs. Based on this curation, we then perform a comprehensive

classification of ICs into well-defined groups and families and identify the extent of understudied

channels in each family, providing a premise for prioritizing studies in IC families with more

understudied members.

We further performed a comprehensive evolutionary analysis to define and collect more than

48,000 well-defined IC orthologous relationships from diverse taxonomic groups spanning

metazoans, fungi, protists, bacteria, and archaea. We demonstrate the application of these

sequences in annotating understudied CALHMs at residue-level resolution using Bayesian

statistical approaches [40] and in predicting and experimentally validating the structural and

functional impact of disease-associated mutations. These studies support our working premise

that integrative mining of available sequence, structure, and functional data on ICs from diverse

organisms (well-studied and understudied) will provide an important context for defining

sequence and structural features associated with understudied channel functions.

Results

Defining the IC complement of the human genome using informatics approaches

We first sought to collect and curate the full IC complement across the human proteome by

systematically mining all the existing data sources, collating information from literature, and

running various informatics tools (see methods) to identify related sequences based on overall

similarity and pore-defining regions. In brief, we first collected all the sequences listed as ICs in

the comprehensive UniProt database [41], as well as Pharos [25], KEGG [23, 24], GtoP [14],

and HUGO Gene Nomenclature Committee (HGNC) [26] databases and subjected them to a

series of annotations as described in Table 1. The complete annotation table for all human ICs

is available in Supplementary Table 1. Primarily, we mined the literature for functional and

structural annotations on these sequences. Broadly, we have included 6 annotation categories.

1) The identifier labels include the UniProt ID, name, and Target Development Level (TDL)

designation from Pharos (as of 3 July 2025). The IDG consortium, through Pharos, assigns

TDL, which can be Tclin, Tchem, Tbio or Tdark, based on the literature data available for

proteins in terms of their potential as drug targets. Including this designation in the annotation is

especially relevant for identifying and prioritizing understudied ICs for further investigation. 2)

The classification labels describe the Group, Class, and Family the IC falls into. This information

has been mined from literature including UniProt, Pharos, GtoP, and HGNC, and the field

"Family designation" provides a consensus family label for the IC. 3) The functional labels have

been manually mined from various sources, relying mostly on previous literature in the "Lit

Resource" column. An important column in this section is the "Unit" column, which describes

whether the IC is directly involved in forming the pore region, where it can be pore-forming

(directly involved in forming an ion-conducting pore), 2-pore (contains 2 tandem pore-

conducting regions), or auxiliary (does not have its own pore domain but interacts with other

pore conducting IC subunits as part of a complex) (please see below and Methods for further

definition of an auxiliary IC). We also provide the ions, gating mechanism, and UniProt

functional description in this annotation category. To verify the ion and gating mechanism

annotations, we mined the literature using Large Language Models (LLM) and Retrieval

Augmented Generation (RAG) systems (Methods, Supplementary Figure 1). The RAG system

was able to successfully verify about 40% of the ion and gating mechanism annotations, while

for others, it was unable to find supporting evidence, either because no evidence was available

in the existing literature or no relevant references were found. The annotations that were not

verified by the RAG system are indicated with an asterisk in Supplementary Table 1. 4) The

structure-related category includes the PDB ID of any experimentally generated crystal structure

or an Alpha-Fold ID for a computationally predicted structure of the IC. 5) The

Complex/Interaction label provides information on the types of complex the IC is involved in

forming, along with the interactors. 6) The last section for transmembrane (TM) and pore

domain-related labels has been curated extensively using various sources and predictors. The

TM regions and pore containing functional domains are annotated based on literature

references and prediction tools, as outlined in the Methods section. TM predictions by TMHMM

[42] and Phobius [43] are provided alongside the TM annotations provided in UniProt. First, we

supplemented this information by adding TM information based on the literature review and the

source. Then, we further analyzed any ICs with an experimentally determined structure using

the MOLE software [44] to identify and annotate the pore region and pore-lining residues. The

MOLE software starts by defining the membrane region of the protein, followed by the

identification of cavities and computation of the pore boundaries to predict the pore lining

residues. These predictions are used as additional evidence for defining the pore containing

functional domain region and a pore containing IC sequence. Sequences without an identified

pore region were classified as auxiliary, as defined by Gurnett et al. 1996 [45], to provide

functional context for this classification. A snapshot of the list of annotations we collected

through this process is shown in Table 1, using Aquaporin 1 as an example.

In the final phase, we conducted sequence similarity searches to detect sequences exhibiting

significant homology with any curated sequences, subsequently subjecting these candidates to

all annotation procedures to determine their eligibility as ICs. By combining the results from this

annotation process, we were able to achieve the following: 1) compile a list of human ICs based

on the presence of distinct transmembrane regions, a detectable pore, and evidence of ion

conductivity, 2) distinguish between a pore containing IC vs an auxiliary IC, 3) identify novel

putative IC sequences and 4) curate and classify the full IC complement in the human

proteome. These results are summarized in Figure 1 and Supplementary Table 1.

419 human ion channel (IC) sequences were curated using this approach, representing an

increase of 75 sequences compared to the previous consensus of 344 ICs in humans [8[25]]. A

comparison of the curated IC sequences in this study against the list of ICs in other resources –

KEGG, GtoP, and Pharos, is provided in Supplementary Figure 2 and Supplementary Table 2.

These sequences were classified into four major groups—VGICs, LGICs, Chloride Channels,

and Others—defined previously [12], and 55 families. VGICs constitute the largest group,

comprising 186 sequences distributed across 21 families, followed by LGICs with 82 sequences

in 10 families. Of the 419 sequences, 62 could not be assigned to the four major groups and

were categorized into outlier families. Among these, 28 are pore-containing ICs, with 19 of the

28 distributed across four families (CALHM, otopetrin (OTOP), transmembrane channel-like

(TMC), and tweety homolog (TTYH)). These families, collectively referred to as "Unclassified"

families, are labeled as such in Figure 1. The remaining 9 pore-containing sequences could not

be assigned to a distinct family and were grouped together under a single "Unclassified" family.

Based on the evaluation of the pore-containing regions, 343 out of 419 ICs were annotated as

pore-containing ICs, while the remaining 76 were classified as auxiliary and fell into 17 different

families. 23 of these auxiliary ICs are soluble with no detectable transmembrane domains. Any

auxiliary ICs that did not belong to a distinct family were collectively classified into an "Auxiliary

unclassified" family.

Next, we sought to use the curated set to define similarities across IC families. Traditional

sequence-based bioinformatics approaches, such as pairwise sequence alignment (e.g.,

BLASTp) [46] or profile hidden Markov models (e.g., HMMER) [47], are limited in their ability to

detect relationships among human ICs due to the extensive divergence in their primary

sequences. These methods often fail to identify homologous ICs across different families, as the

sequence similarity falls below detectable thresholds. Consequently, such approaches are

inadequate for comprehensive classification or comparison of ion channel families in humans.

Instead, we relied on representations (embeddings) derived from evolutionary scale protein

language models [48] to capture sequence, structure, and evolutionary information and use

them to generate pairwise sequence alignment. Specifically, for the 343 pore-containing ICs, we

passed their pore-containing functional domains to a protein sequence embedding model called

DEDAL [49] to perform a pairwise sequence alignment. Given that sequence similarity is largely

restricted to the pore-containing functional domains of ICs, we computed protein embeddings

using only these regions. As auxiliary ICs lack such domains, they were excluded from this part

of the analysis. The resulting all-vs-all sequence similarity scores were used to generate uniform

manifold approximation and projection (UMAP) embedding scores (Supplementary Figure 3).

An average of all IC sequences within a family was calculated to place the IC family bubbles in

Figure 1. Thus, families placed close to each other in Figure 1 indicate their similarities in

sequence embeddings, which capture learned representations of sequence, structure,

evolutionary conservation, and functional properties. Since the auxiliary IC families did not have

UMAP embedding scores, they are arbitrarily placed at the bottom of the figure.

Most of the VGIC families group together on the top right except for inward rectifier potassium

channels (K_{IR}), two-pore domain potassium channels (K_{2P}), and plasmolipin (PLLP), where K_{2P}

and PLLP are more centrally dispersed, while K_{IR} falls closer to LGIC families acid-sensing ion

channels (ASICs), epithelial sodium channel (ENAC), and P2X purinoceptors, all of which share

similar transmembrane topology with only two transmembrane helices. The inositol 1,4,5-

trisphosphate (IP3) and ionotropic glutamate receptors are also centrally located, while the Cys

loop receptor LGIC families group separately from others towards the bottom right, indicating

their shared functional and structural similarities. Interestingly, pannexins, volume-regulated

anion channels (VRAC), connexins, chloride channel CLIC-like (CLCC), and CALHM families

group closer together centrally, indicating shared features.

We also mapped TDL annotations defined by the IDG [30] in Pharos [25] to highlight the depth

of understudied ICs within each group, and hopefully, spark additional studies of those

members. Briefly, IDG describes Tclin as targets that have an approved drug; Tchem has

known high potency small molecule binding targets; Tbio has experimentally supported Gene

Ontology [50] annotations based on published literature; while Tdark is manually curated at the

primary sequence level in UniProt, but do not meet any of the Tclin, Tchem or Tbio criteria [25].

There are 19 Tdark, 185 Tbio, 88 Tchem and 127 Tclin ICs spread across all families. The

proportion of these levels for each family is shown as pie charts in Figure 1. CALHM family has

the highest proportion of Tdark ICs, with OTOP, TMC, Connexins, calcium-activated chloride

channels (CaCC), calcium voltage-gated channel (CACN), $K_{2P,}$ and the unclassified families

accounting for the remaining Tdark ICs, emphasizing the need for additional research on these

families. On the other hand, VGICs voltage-gated potassium channel (K_V), voltage-gated

sodium channel (Na_V), voltage-gated calcium channel (Ca_V), ryanodine receptor (RyR), and

LGICs Gamma-Aminobutyric acid type A ($GABA_A$) receptor, nicotinic acetylcholine (nACh)

receptor, ENAC have the highest proportion of Tclin ICs, solidifying their status as the well-

studied families.

Mining and cataloging IC orthologs across the tree of life

Next, we sought to extend this annotation and collect related ICs from other organisms across

different taxonomic lineages from the tree of life [41] using the full complement of the annotated

human IC sequences. To this end, we used a graph-based orthology inference approach [51]

starting from the full length and the pore-containing functional domains of the human IC

sequences to define orthologous relationships across more than 1,500 proteomes from the

UniProt Proteomes database [41]. These selected proteomes dataset includes 353 Archaea,

696 Bacteria, and 547 Eukaryota organisms. This method has previously been successfully

employed to define orthology across such a large collection of proteomes for protein kinases

[51], and since it relies on both the full-length sequence and the annotated pore-containing

domains, it can more accurately define true orthologs that are indeed evolutionarily related

across organisms. Since this method relies on the pore containing domain annotations for a

domain-based orthology inference, we only used the 343 pore-containing IC sequences for this

analysis, and did not include the auxiliary ICs that lack a pore containing domain.

□

By combining domain and full-length searches, we could identify more than 48,000 IC

orthologous relationships across diverse organisms. Figure 2A shows a heatmap depicting the

phylogenetic profiling of human IC orthologs along the horizontal axis across the taxonomic

lineages shown by the tree along the vertical axis. The color intensity indicates the percentage

of orthologs detected for all the sequences in the IC family for a given taxonomic group. The

highest number of orthologs were found for Aquaporins, including bacterial and archaeal

orthologs, indicating that this family of ICs is the most conserved across lineages. Along with

Aquaporins, the two paralogs of Golgi pH regulator ICs (GPHRs), both indicated as Tbio

channels, were found to have the largest number of orthologs across metazoans. Along with

GPHRs, several other ICs, notably members of the K_V family and $GABA_A$ receptors, display

some of the most widespread orthologs across organisms. Some dark channels, such as

members of the TMC (TMC7 and TMC3) and CALHM (CALHM5 and CALHM3) families, are

well conserved across metazoans and vertebrates, respectively, yet very little is known about

their functions. On the other hand, some Tdark channels, such as Potassium channel K

member 7 (KCNK7) and TMC4 have a lineage-specific set of orthologs that extend only up to

mammals. A full list of all the orthologs detected through this analysis is provided in

Supplementary Table 3.

To use this evolutionary conservation for functional inference of dark ICs, we first used

hierarchical clustering to group ICs with similar orthology profiles into related clusters, which led

to the definition of 9 clusters, as shown in Figure 2B. Since the orthology profiles in prokaryotic

lineages were very sparse, only the eukaryotic orthology profiles were retained for this analysis.

Within eukaryotes, each cluster has a well-defined signature of orthology conservation. For

example, ICs in cluster 2 have most of their orthologs only in mammals, whereas clusters 4, 5,

and 6 have orthologs spanning other vertebrates. Similarly, clusters 7 and 8 have orthologs

extending to other metazoans, while seven ICs in cluster 9, including the 2 Golgi pH regulators

A and B and the sodium leak channel NALCN, have detectable orthologs in Fungi, Plants,

Protists, and other eukaryotic lineages. To translate these patterns of orthology profiles to

function, we performed a functional Gene Ontology (GO) term enrichment analysis for the

human ICs in each cluster (Figure 2C). As expected, the most significant GO terms were related

to ion channel function (colored green in Supplementary Table 4). However, beyond channel

function, the enriched functional GO terms obtained for each cluster correlate well with the

physiological functions present in the orthologous organismal groups. For example, clusters 4,

5, and 6 are conserved in vertebrates and have six dark IC sequences, including unclassified

ICs such as CALHM 3 and 5, and connexins GJA10, GJD4, and GJE1, which had functional

enrichment for regulation of heart contraction and heart rate, traits that could be closely related

to a closed blood circulatory system with an endothelium, present in vertebrates [52]. On the

other hand, cluster 2 with orthologs in mammals was conserved in mammalian-specific

reproductive functions such as sperm capacitation.

Using the orthology information to identify evolutionary constraints in CALHMs

Once the orthologs have been identified across different organisms, they can be leveraged to

find evolutionarily conserved signals that could point to functional similarities within related

groups of sequences. To this end, we classified subsets of orthologous IC sequences into

evolutionarily related clusters using a Bayesian Partitioning with Pattern Selection (BPPS)

algorithm, which classifies sequences based on patterns of amino acid conservation and

variation in a large multiple sequence alignment (see methods) [40]. For this analysis, we

focused on the orthologs of CALHM family of IC sequences. The CALHM proteins constitute a

family of large pore channels, forming oligomeric assemblies of different sizes [53]. It has 6

member sequences in humans (CALHM1-6), 3 labeled as dark channels, constituting one of the

families with the highest prevalence of dark ICs. It has been well established that CALHM1 is

activated by removing extracellular Ca^{2+} and membrane depolarization and that the heteromeric

CALHM1 and CALHM3 channels are implicated in the ATP release during taste perception [54].

CALHM1 has also been shown to regulate cortical neuron excitability [55], locomotion and

induces neurodegeneration in *C. elegans* [56]. On the other hand, CALHM6 has been shown to

be important for immune system functions by facilitating induction of immune cells during

infection [57]. In contrast, the activation stimuli and physiological roles of other family members

remain largely unexplored. All CALHM family members share a common arrangement in the

transmembrane domain, with each subunit consisting of four transmembrane segments (S1–4),

forming a large cylindrical pore lined by the first transmembrane segment S1 along with a short

N-terminal helix preceding S1. The exact gating mechanism of the CALHM channels is still

unclear, partly since all their cryo-EM structures have been in an open state. However, previous

studies have indicated that the N-terminal region, called the amino-terminal helix (NTH) plays a

crucial role in modulating voltage dependence and stabilizing the closed channel state [58],

while a more recent study has suggested that the voltage dependent gate is formed by the

proximal regions of S1 [59]. Collectively, these studies indicate the role of the amino termini –

the NTH and S1 regions, to play a crucial role in the gating mechanism and the determination of

pore size [60] of CALHMs.

Thus, we performed a phylogenetic analysis to find evolutionarily conserved residues that might

shed more light on these critical mechanisms that govern CALHM function. Figure 3A shows a

phylogenetic tree depicting the evolutionary relationship across the 6 members of the CALHM

family across diverse taxa. CALHM1 and 3 form a distinct clade from CALHM 2, 4, 5 and 6. We

first analyzed 5805 CALHM homologs to identify pattern positions conserved across all these

sequences with the hypothesis that such conserved positions could point to shared functional

features across all CALHMs. The Bayesian analysis identified 13 aligned positions conserved

across all 6 CALHM homologs, which we will refer to as CALHM shared patterns (Figure 3B,C).

Most of these conserved positions were hydrophobic amino acid residues, and 5 conserved

Cystine residues, 4 of which are involved in forming inter-molecular disulfide bridges

(C46=C130, C48=C162). Residue position numbers reflect the numbering based on human

CALHM2 (PDB id: 6uiv). Interestingly, five of the conserved positions (F44, Y56, I61, P64,

W117) are located close to the amino termini in a functionally important linker region connecting

S1 and S2 (S1-S2 linker). Specifically, this linker region could regulate the dynamic

conformational changes of S1, where the S1 could adopt either a vertical conformation relative

to the membrane plane, resulting in an enlarged pore size, or a lifted conformation, leading to a

reduced pore size (Figure 3D) [60]. Therefore, mutations in this linker are expected to affect

channel functions. Based on the positioning of these conserved residues, and previous studies

that highlight the importance of this region in gating functions, we hypothesized that these

conserved pattern positions play a role in the gating mechanism of CALHM. The conservation of

these residues across orthologs of all 6 CALHM sequences further suggests that all CALHM

paralogs could share this gating mechanism.

To test our hypothesis and determine the functional importance of the CALHM shared patterns,

we sought to perform a series of mutational experiments to determine the functional implications

of perturbations at these positions. To achieve this, we methodically determined target

mutations for each position. We first scanned the Genome Aggregation Database (gnomAD)

[61] to check for any prevalent variations at these conserved positions within the sampled

population to use as our mutational targets. We found several disease variants at these

positions that are listed in Figure 3E that were used to prioritize target mutations. For positions

where a variation was not found, we tested their significance by performing alanine mutations,

causing a deletion of the side chain at the β-carbon.

Targeted mutational and electrophysiological studies of CALHM conserved residues

To assess the functional roles of the predicted conserved residues in CALHM channels, we

performed targeted mutational and electrophysiological analyses in two representative human

CALHMs: the well-studied human CALHM1 and the relatively understudied human CALHM6.

Building on the insights from previous structural and functional studies [59, 60], which implicated

the NTH/S1 region in channel gating, we prioritized the constraints located in the S1-S2 linker

and at the interface between S1 and the transmembrane domain (Figure 3B,C,E). These two

regions were hypothesized to play distinct roles in gating: the S1-S2 linker as a flexible hinge

and the S1-TMD interface as a stabilizing contact for S1 movement.

Both total protein Western blotting and surface biotinylation assays showed that the mutants

have either comparable or substantially higher expression levels than wild type, confirming that

neither protein expression nor trafficking to the plasma membrane was impaired

(Supplementary Figure 4). We then established electrophysiological measurements for wild-type

CALHM1 and CALHM6 following protocols described in the methods. Our results were

consistent with previous findings [55] (Figure 4). In addition to room temperature, which is

commonly used for patch-clamp studies of CALHM channels, we also conducted measurements

at physiological temperatures (37°C). We observed that CALHM1 activity was significantly

higher at physiological temperature than at room temperature, as reported previously [62, 63],

while the CALHM6 currents showed only a small increase (Figure 4). Notably, both CALHM1

and CALHM6 currents were inhibited by extracellular Gd^{3+} (Figure 5C,I), a commonly used

inhibitor for the CALHM family, consistent with previous studies [55, 57, 59]. The robust currents

at 37°C, particularly those of CALHM1, provide a solid basis to interpret the phenotypes of the

mutants tested, as detailed below.

Mutations of a predicted conserved residue in the S1–S2 linker (F40 in CALHM1; F39 in

CALHM6) either abolished or markedly reduced channel activity in both CALHM1 and CALHM6,

presumably by impeding the conformational dynamics of S1 required for channel gating (Figure

5P,Q; Supplementary Figure 5). Similarly, mutation of a conserved tyrosine residue on S2 (Y52

and Y51 in CALHM1 and CALHM6, respectively), whose side chain directly contacts the S1-S2

linker, also resulted in strong phenotypic changes: Y52A in CALHM1 abolished channel

activation (Figure 5P,Q; Supplementary Figure 5), while Y51A in CALHM6 converted the

channel from voltage-independent to voltage-dependent gating, resulting in outward rectification

(Figure 5J-L,P,Q; Supplementary Figure 5). These residues are likely key determinants

controlling the conformational dynamics during gating.

Interestingly, mutations in other conserved residues, near – but not within – the S1–S2 linker

(W114, L57, and P60 in CALHM1; W113 and A127 in CALHM6), also abolished or markedly

reduced channel activity (Figure 5P,Q; Supplementary Figure 5). Among these, W114 in

CALHM1 and W113 in CALHM6 directly contact S1. Thus, substituting these bulky hydrophobic

residues with smaller (cysteine or alanine) or positively charged (arginine) residues is expected

to alter the conformational dynamics of S1 and therefore impair channel gating. Notably,

however, the CALHM6 W113A was an exception, retaining wild-type like currents at 37°C and

exhibiting voltage dependence and inhibition by extracellular Gd^{3+} similar to wild-type CALHM6

(Figure 5M-O), suggesting that this mutation does not impair fundamental gating properties.

Finally, since most mutants showed either reduced or completely abolished activity, we included

a positive control in the electrophysiological experiments: I109W on CALHM1, which has been

previously documented to increase channel activity. As expected, we reproduced this gain-of-

function phenotype (Figure 5). To further explore channel-specific differences, we also

examined the corresponding mutant in CALHM6 (L108W). Interestingly, L108W decreased

CALHM6 channel activity (Figure 5), possibly reflecting inherent differences between the two

channels.

Discussion

Here, for the first time, we have computationally defined the full IC complement of the human

genome – the "channelome". We provide this comprehensive list and a rich annotation of

functional, structural, and sequence features, mapping them to 4 widely accepted IC groups and

further classifying them into 55 families. We also highlight unclassified outlier groups that

contain most of the understudied "dark" and potentially novel unclassified IC sequences. As part

of our curation, we also provide annotation for the pore-containing domain that is based on

multiple sources, including an extensive literature review, further strengthened by the

application of literature mining using LLMs and structural analysis tools. We use the pore-

containing domain to distinguish pore-containing from auxiliary ICs, providing additional

functional context. Our annotations can serve as a reference for comparing ICs within or across

families, designing experiments for functional studies, and identifying potential targets to

illuminate the functional relevance of understudied ICs further.

The application of LLMs, specifically RAG systems, in our research demonstrates significant

potential in automating literature mining and correcting certain inaccuracies. However, the

system currently faces limitations related to diverse data presentation formats, complex channel

subunit relationships, and variant gene nomenclature, which hinder its effectiveness. The high

frequency of responses where evidence was not found underscores the need for enhanced

entity recognition capabilities, particularly for accurately identifying gene and protein names

across different nomenclatures. Additionally, the RAG system's ability to critically engage with

data by identifying and amending presumed inaccuracies can be widely used to verify and

correct annotation entries, though this also raises concerns about potential overcorrections or

misinterpretations without adequate validation. By more data cleaning and standardizations,

LLMs and RAG systems can become more reliable and integral tools for curating and

expanding bioinformatics databases, ultimately enhancing the accuracy and

comprehensiveness of our IC knowledgebase.

To the best of our knowledge, identifying true orthologs across the tree of life performed here is

the first large-scale effort to consolidate ion channel orthologs across diverse organisms and

provides a comprehensive view of the evolutionary conservation of individual ICs. Based on this

analysis, we could pinpoint the Aquaporins family of ICs as the most ancient family of ICs with

orthologous sequences present back to bacterial and archeal species. More importantly, the

clustering of human ICs based on their depth of conservation placed them into functionally

meaningful clusters with many enriched functions exclusive to their orthologous taxonomic

group. The placement of understudied ICs in such groups allows for meaningful functional

predictions, which can be further elucidated using experimental approaches.

The orthologous sequences identified and presented here can serve as a valuable resource for

performing evolutionary and functional analysis using statistical and machine learning

approaches. We demonstrate one such use by performing a Bayesian pattern-based

classification on the CALHM subset of ICs to identify amino acid positions significantly

conserved across all CALHM sequences. These features reside on a functionally important S1-

S2 linker region that potentially governs their gating mechanism. Because these features were

conserved and shared across all CALHM sequences, we also hypothesize that all CALHM

sequences share this gating mechanism. By performing targeted mutations on these conserved

residues, we show that a mutation in any of these evolutionarily conserved residues results in a

dramatic loss of gating function, thus highlighting the functional importance of the identified

pattern positions.

Among the conserved positions targeted for mutations, mutations of residues in the S1-S2 linker

(F40C in CALHM1; F39L in CALHM6 and Y52A in CALHM1; Y51A in CALHM6) resulted in

strong phenotypic changes, most likely due to their direct involvement in the conformational

dynamics of S1 for channel gating. Mutations in W114 in CALHM1,which is not directly on the

S1-S2 linker, also resulted in abolished activity, however mutations in an analogous position

W113 in CALHM6 did not impair channel function, presenting an exception compared to other

residues across CALHM1 and 6. Among other residues that are near – but not within – the S1-

S2 linker, the positioning of P60 on the S2 helix of CALHM1 is particularly intriguing. In the cryo-

EM structure, P60 resides approximately midway along S2, inducing a distortion or bending of

the helix due to the rigid ring structure of proline. This bent configuration allows extensive

contact between the extracellular portion of S2 and the S1–S2 linker, while the intracellular

portion of S2 interacts extensively with S1. Furthermore, we hypothesize that this bent

conformation of S2 contributes to the flexibility of the S1–S2 linker, as if S2 were straightened, it

would consequently straighten the S1–S2 linker, potentially reducing its flexibility. Replacing the

proline with glycine eliminates the structural constraint imposed by proline's cyclic side chain,

allowing S2 to adopt a more regular, uninterrupted helical structure. Consequently, a

straightened S2 may lead to reduced flexibility of the S1–S2 linker, ultimately impacting channel

gating. Similarly, L57 in CALHM1 is positioned near P60 on the protrusion of the bend of S2,

with its sidechain facing residues on the adjacent S3. We hypothesize that the larger sidechain

of the L57R mutant would force S2 to straighten due to steric collision between the bulky

arginine sidechain and residues on S2, again reducing the flexibility of the S1-S2 linker and alter

channel gating. A127 in CALHM6 resides in a short alpha helix within the extracellular domain.

While A127 does not directly interact with the S1–S2 linker, it is adjacent to the highly

conserved disulfide bond (between C41 and C126), linking the helix containing A127 to the S1–

S2 linker. Thus, it is conceivable that the A127G mutant may indirectly affect the conformation

of the S1–S2 linker through this disulfide bond.

These findings complement prior studies and provide additional insights into CALHM gating

mechanisms. Several previous studies examined conserved residues in CALHM channels and

proposed roles in gating [56-59, 62, 64]. These studies were based on comparisons of a limited

set of homologs from model organisms such as *C. elegans* or mouse, and identified residues in

various parts of the protein that influence gating, providing important clues towards their

functioning. Here, we have extended the analysis to encompass thousands of CALHM

sequences collected from the entire tree of life, allowing us to identify residues conserved

across all family members through evolution. Guided by a hypothesis from previous structural

and functional studies [59, 60] which highlighted the NTH/S1 region as a key element in channel

gating, we focused on evolutionarily conserved residues in the S1–S2 linker and at the interface

of S1 with the rest of the TMD. We reasoned that if S1 movement is critical for gating, then

these two structural elements—the S1–S2 linker, acting as a hinge, and the S1 interface with

the TMD, serving as a stabilizing contact—would be key determinants of the conformational

dynamics of S1.

Together, our experimental data indicate that the conserved residues at the S1–S2 linker and its

immediate surroundings play an important role in CALHM channel gating, supporting a model in

which S1 dynamics are central to the gating mechanism [59, 60]. Furthermore, the contrasting

phenotypes of Y52A in CALHM1 versus Y51A in CALHM6, W114A in CALHM1 versus W113A

of CALHM6, and I109W in CALHM1 versus L108W in CALHM6 highlight differences in gating

properties specific to each family member. Overall, the complementary results from this study

and the published literature highlight the complexity of CALHM gating and suggest that distinct

conserved elements contribute to both shared and lineage-specific gating mechanisms in this

unique family of large-pore channels. Importantly, this was demonstrated not only for the well-

studied CALHM1 but also for the relatively understudied human CALHM6, providing valuable

clues into shared gating features across light and dark CALHM sequence sets.

In addition to the CALHM shared patterns, we also present a second subset of conserved

positions shared only within CALHM 2,4,5 and 6, paralogs that are evolutionarily closer

compared to CALHM1 and 3 (Supplementary Figure 6). These subsets of conserved residues

fall in the intracellular helical segment that is involved in oligomerization and formation of the

pore complex governing CALHM function. Previous studies have indicated that CALHM2 adopts

a unique undecameric oligomer different from the CALHM1 octamer [65]. Thus, with the

presence of shared patterns within the subset of CALHM2 and the understudied CALHM4,5 and

6 in this intracellular helical region, we hypothesize that these conserved positions help maintain

a similar mode of oligomerization between these evolutionarily related subsets of CALHMs.

Because many disease mutations map to these conserved positions, we believe the datasets

and approaches offer a powerful avenue for elucidating the functional and clinical relevance of

the understudied ICs.

Methods

Identification and annotation of human ICs

The annotation of human ICs was performed using a semi-automated pipeline. First, all the

protein sequences that were labeled as ICs were collected from the UniProt [41], KEGG [23,

24], Pharos [25], GtoP [14], and HGNC [26] databases. The annotations described in Table 1

were then collected from literature sources manually and compiled together. The UniProt

specific labels and the complex formation information were extracted from the UniProt

database. The sequences were run through TMHMM [42] and Phobius [43] to predict

transmembrane and helical regions. The predictions were cross-referenced and confirmed

against the CDD [66], Pfam [67], PrositePattern , PrositeProfiles [68] and Simple Modular

Architecture Research Tool (SMART) [69] databases, where a reference was available. Finally,

the prediction of the pore region and pore-lining residues was supplemented using the MOLE

software [44]. The prediction of pore region and the information available in previous literature

was combined to annotate the final set of auxiliary IC sequences. An auxiliary IC is defined as

any sequence that itself does not have a pore domain, but has experimental evidence of being

part of an IC complex. Thus, any IC where a pore domain could not be found was subjected to

additional literature review to find evidence for their interactions with the pore containing ICs,

and if the evidence was found, they were included as auxiliary ICs, otherwise they were

removed from our curated IC list.

Retrieval-Augmented Generation (RAG) system for verifying ion-selectivity and gating-

mechanism annotations

To systematically evaluate the ion-selectivity and gating-mechanism fields in the curated human

IC data set, we built a RAG pipeline comprising three sequential stages: corpus construction

and vectorization, query formulation and similarity search, and evidence synthesis with a LLM.

All scripts and configuration files are archived in the project repository

(https://github.com/esbgkannan/ionchannels-final-pdf).

Corpus construction and vectorization: PubMed identifiers (PMIDs) linked to each IC were

obtained from our annotation pipeline. Full-text PDFs were downloaded, converted to plain text

with the *pdfminer.six* Python library, and segmented into overlapping fragments of

approximately 1000 tokens (50-token overlap) using the *CharacterTextSplitter* module of

LangChain. Each fragment was embedded with the OpenAI *text-embedding-3-large* model

(3072-dimensional vectors). Embeddings and fragment metadata (PMID, page number,

fragment index) were stored in a local *Qdrant* vector database. Prior to embedding, boiler-plate

headers, footers and reference lists were removed to retain only article body text.

Query formulation and similarity search: For every IC, two natural-language questions were

generated automatically: 1. "Is there any evidence that *<ION>* is the ion selectivity of the *<IC-*

NAME> ion channel?", and 2."Does this article provide evidence for *<GATING>* as the gating

mechanism for the *<IC-NAME>* ion channel?".

Here, *<IC-NAME>* encompasses the UniProt primary name and all recognized synonyms

(produced by extract_alternative_names.py script in

https://github.com/esbgkannan/ionchannels-final-pdf). Each query was submitted to Qdrant to

retrieve relevant fragments from the database. When PMIDs were available in the dataset, an

initial PMID-filtered search was performed; filtered and unfiltered results were subsequently

merged and fragments with the highest cosine similarity retained.

Evidence synthesis with an LLM: The retrieved fragments were inserted into a fixed prompt

and supplied to GPT-4o (temperature = 0). The model was instructed to return a JSON object

having three keys and values of answer (Found, or Not Found), confidence (a number between

0 to 1), and evidence (an explanation of LLM's decision for the answer). Outputs were parsed

with LangChain's *JsonOutputParser*, combined with fragment metadata, and compiled to a

single JSON file for further analysis. Predictions with confidence ≥ 0.80 were accepted

automatically; lower-confidence or *Not Found* results triggered manual review. The structured

records were merged with the annotation table, enabling automatic comparison with pre-existing

entries and flagging of discrepancies for curator review.

Entries for which no supporting evidence could be located (model answer 'Not Found' confirmed

on manual inspection) are marked with an asterisk in Supplementary Table 1. An illustrative

exchange is provided in Supplementary Figure 1.

Defining a pore containing functional domain

We aimed to define the pore-containing functional domain so that it spanned the pore region

and included all the TM domains present in an IC. This ensured that our domain definition

included the pore region and any other functionally important TMs, while excluding any

accessory domains on the flanking sequences. To ensure we did not miss any TM regions, four

different sources of annotation information were combined to define the pore containing

functional domain for the 343 ICs. 1) The TM predictions from TMHMM [42] and Phobius [43]

was first used to identify the TM regions. 2) The UniProt-based TM annotations were matched

with the predictions to verify the TM regions further. 3) The literature-based TM annotations

were then used to verify the TM positions and organization, where available manually. 4) Where

experimentally resolved crystal structure coordinates were available, the MOLE software was

used to identify the pore lining residues.

Pairwise sequence alignment using sequence embeddings

The pore containing functional domains for the 343 pore containing ICs were passed to DEDAL

[49] which was run using standard parameters and resulted in homology logit scores based on

an all-vs-all pairwise sequence alignment using their sequence embeddings. These scores were

used to generate a sequence similarity matrix passed to the umap function of the umap_learn

package v0.5.7 [70] in python 3.9 to generate 2D UMAP embeddings. This was used to

generate a 2D scatterplot that defined the placement of individual ICs shown in Supplementary

Figure 3. Finally, an average position of ICs within a family was used to define the placement of

that family in Figure 1.

Orthology detection and analysis

The KinOrtho pipeline [51] was used for defining the orthologs and co-orthologs of human ICs

across the tree of life. KinOrtho employs a graph-based orthology inference approach using

both the full-length and the pore domain regions for inferring orthologous relationships primarily

relying on sequence similarity within these regions. The pipeline began with pairwise sequence

similarity searches using both full-length and pore-containing domain sequences of human ICs

against the target proteomes database. Since a pore domain needs to be defined for this

analysis, we only used the 343 pore containing IC sequences to perform orthology detection.

They were used as query for running KinOrtho against a reference database of curated

canonical proteomes from the UniProt Proteomes Release 2022_05 that consisted of 343

Archaea, 696 Bacteria and 548 Eukaryota proteomes. The initial sequence similarity search was

conducted using BLASTp [46] with default parameters and an e-value cutoff of 1e-5 to retain

high-confidence hits. The top hit from this forward search was then used as query in a reciprocal

BLASTp search against the human proteome, applying a more stringent e-value threshold of

1e-200 to retain only the top and high confidence hits. Pairs of sequences were retained only if

each was the top hit for the other in this reciprocal search. Such a reciprocal best-hit (RBH)

strategy minimizes false positives by ensuring that the candidate orthologs are each other's

most similar sequence in the respective proteomes, which is particularly important when

distinguishing true orthologs from paralogs or other homologs that may share partial similarity

but have diverged functionally [71].

All the retained hit pairs were then passed to cluster analysis using OrthoMCL [72] and filtering

of relationships to keep only relationships within the same cluster. This process was conducted

separately using both the full-length human IC sequences and their pore domain sequences.

Only sequences that were identified as orthologs in both the full-length and domain-based

analyses were retained as high-confidence orthologs. This ensured that both overall sequence

similarity and conservation of functionally critical domains were satisfied.

Finally, the obtained ortholog sequence sets were subjected to a series of validation tests that

provide more guardrails and ensure we avoid duplicate, fragments and extraneous hits ensuring

that only sequences with high quality annotation were retained as true orthologous

relationships. These validation tests followed the following steps:

Step 1: Check UniProt Entry Type: We first check the "entryType" flag in UniProt to check

whether it is: "Reviewed" or "Unreviewed". "Reviewed" sequences are manually annotated and

reviewed, thus are of the highest quality and safe to include. Any sequences with this status are

passed. Sequences with "Unreviewed" status proceed to Step 2.

Step 2: Check for Protein Existence Evidence: For the "Unreviewed" sequences, check their

"proteinExistence" flag. This flag can have one of five values: 1. Evidence at protein level 2.

Evidence at Transcript level 3. Inferred from homology 4. Predicted 5. Uncertain. Flags 1,2, and

3 are dependable evidence of protein existence, thus are passed, whereas 4 and 5 are low

confidence and subjected to further verification in Step 3.

Step 3: For the proteins with low confidence (4 or 5) protein existence values, we next check

five different measures/flags to ensure its integrity:

1. Check their sequence version for unusual patterns. Multiple updates with high

sequence versions or a very old version update indicate unstable or deprecated sequences,

respectively.

2. Perform sequence level checks.

691   A. Check for compositional bias: Sequences with compositional bias might be

692   low complexity regions or repeats.

693   B. Check for proportion of non-standard amino acids: Presence of large number

694   of non-standard amino acids could also indicate poor protein annotation.

695   C. Check sequence length: Unusually short (<30 aa) or long (>5000 aa)

sequences could indicate fragments or fusion errors and misannotation.

3. Check for cross-references: If the protein completely lacks annotated domains, or

additional cross referenced metadata, it might suggest poorly characterized or erroneous

proteins. For any sequence that is subjected to Step 3, it should pass all the checks in this step

to be included as an orthologous hit.

48694 unique orthologous relationships from 36846 sequences passed the orthology pipeline

and validation checks and were included in the final list of orthologous sequences. The list of all

the orthologus sequences that passed the validation check are provided in Supplementary

Table 3.

To illustrate the extent of these orthologous sequences across different taxonomic lineages,

they were used to create a presence/absence matrix of orthologs, with rows representing each

human IC and columns representing the different organisms. To reduce individual granularity

and get estimates at the family and lineage level for visualization, first, the orthologs were

grouped by IC families and then, by their defined taxonomic lineages. For each group

representing one IC family and one taxonomic lineage, a percentage value representing the

proportion of detected orthologs was calculated using the following: (total number of orthologs

found for all ICs in a family)/(total number of organisms queried in the taxonomic lineage *

number of human IC sequences in the family). These percentages are depicted in the heatmap

in Figure 2A where each cell represents an IC family and its proportion of orthologs in a given

taxonomic lineage.

Next, the presence/absence matrix of individual ICs was used to perform an orthology profiling

clustering and enrichment analysis. Since this analysis is human IC-centric, only the orthologs

from eukaryotic lineages were selected. Hierarchical clustering was performed with the Ward

method of clustering and Euclidean distance metric using the scipy package [73] in Python. The

resulting dendrogram was used to define 9 clusters that group ICs with similar

presence/absence patterns. Human ICs falling in these 9 clusters were then subjected to a

functional enrichment analysis using the Gene Ontology resource GO enrichment tool [74]. GO

terms with a corrected FDR <0.01 were retained as significantly enriched terms.

CALHM evolutionary analysis

BPPS was used to perform a pattern-based classification of the CALHM homologs. First,

orthologs for all 6 human CALHMs were collected. This ortholog dataset was supplemented with

more hits from the UniProt database using MAPGAPS, a multiply-aligned profile for global

alignment of protein sequences [12]. Along with finding the best hits for the human CALHMs,

MAPGAPS also aligns those hits to the template profile alignment to generate a large multiple-

sequence alignment of the resulting 5805 CALHM. This large alignment was then subjected to

734 BPPS, which performs a hierarchical classification of the sequence sets based on conserved

735 pattern positions shared by subsets of sequences using a Bayesian statistical procedure [40].

736 This generates a hierarchical cluster where sequences within each cluster are defined by

737 distinct conserved patterns.

739 **Cell lines**

740 HEK293T cells are purchased from Sigma-Aldrich (Catalogue Number: 96121229). Neuro2A

741 cells are purchased from ATCC (Catalogue Number: CCL-131). The cells are authenticated and

742 are mycoplasma contamination tested negative by the vendor.

744 **CALHM plasmid construction and expression by transient transfection**

745 Full-length human CALHM1 and CALHM6 in the pEGC Bacmam vector was used. The

746 translated product contains the human CALHM1 or CALHM6 protein, a thrombin digestion site

747 (LVPRGS), an enhanced GFP protein, and an 8x His tag. Primers for site-directed mutagenesis

748 were designed using Snapgene and synthesized by Eurofins Genomics. The QuikChange

749 mutagenesis protocol was used to generate all the mutants of the study. Sanger sequencing

750 was performed to identify positive clones.

752 Adherent HEK293T (ECACC, Catalogue Number: 96121229) cells were grown in DMEM media

753 supplemented with 10% fetal bovine serum. Transient transfection was conducted using

754 lipofectamine-2000 by following the manufacturer's protocol. Specifically, the cells were cultured

755 in 60 mm Petri dishes until 80% confluency. Transfection solution was made by mixing 500 ng

756 of plasmid DNA, 4 uL of lipofectamine-2000 reagent, and 100 uL Opti-MEM media. After 20 min

757 incubation at room temperature, the DNA-lipid complexes were added to the cell culture and

758 incubated at 37°C. The next day, 10 mM sodium butyrate was added to the cells to boost

protein expression. The cell culture was then grown at 30°C for another day before harvesting.

The cell pellet was flash-frozen with liquid nitrogen and stored at -80°C. Each CALHM1 and

CALHM6 mutant were transfected in triplicate as biological replicates.

Expression analysis of CALHM1, CALHM6, and their mutants

To analyze total expression levels of wild-type CALHM1, wild-type CALHM6, and their mutants,

cells were lysed in TBS buffer (20 mM Tris, pH 8.0, 150 mM NaCl) supplemented with 10%

lauryl maltose neopentyl glycol and cholesterol hemisuccinate detergents on ice. Lysates were

solubilized at 4°C for 1 hour and clarified by centrifugation at 13,000 rpm for 5 min. The

supernatant was mixed with 4× SDS loading buffer containing 5% 2-mercaptoethanol and

resolved on a 4–20% gradient SDS-PAGE gel.

In-gel fluorescence imaging of the C-terminal GFP tag was performed immediately after

electrophoresis using a Chemidoc system to visualize CALHM1 and CALHM6 proteins.

Following imaging, proteins were transferred to a nitrocellulose membrane using semi-dry

transfer buffer (48 mM Tris base, 39 mM glycine, 20% methanol). Membranes were blocked

with TBST (20 mM Tris, pH 8.0, 150 mM NaCl, 0.1% Tween-80) containing 4% non-fat milk for

1 hour at room temperature. β-actin was detected as a loading control by incubating

membranes with HRP-conjugated anti-β-actin antibody (Proteintech, cat. no. HRP-60008;

1:2000 dilution) for 1 hour at 4°C. After four washes with TBST (15 min each),

chemiluminescence signals were developed using Pierce ECL substrate and imaged on a

Chemidoc system. A brightfield image was overlaid to visualize protein marker positions.

Each mutant was analyzed in triplicate from independently transfected cell pellets. GFP signal

intensities for CALHM1 and CALHM6 were quantified using ImageJ and normalized to β-actin

chemiluminescence signals. Mean values and SEM were calculated for each mutant, and

relative expression levels were compared to wild-type proteins using bar graphs generated in

Microsoft Excel.

Surface expression analysis

Surface biotinylation of CALHM1, CALHM6, and their mutants was performed using the Pierce™ Cell Surface Biotinylation and Isolation Kit (Thermo Fisher Scientific) following the manufacturer's protocol. Cells were cultured and harvested 48 hours post-transfection as described above. Surface-isolated proteins were analyzed by SDS-PAGE, and in-gel fluorescence imaging of the C-terminal GFP tag was performed using a Chemidoc system.

Electrophysiology

TsA201 cells expressing plasmids encoding N-terminal GFP tagged human CALHM1 or CALHM6 were used. After 1-day post-transfection with plasmid DNA (100 ng/mL) and Lipofectamine 2000 (Invitrogen, 11668019), the cells were trypsinized and replated onto poly-L-lysine-coated (Sigma P4707) glass coverslips After cell attachment, the coverslip was transferred to a recording chamber. Whole-cell patch-clamp recordings were performed at room temperature (21-23°C) or body temperature (36-38°C). Signals were amplified using a Multiclamp 700B amplifier and digitized using a Digidata 1550B A/D converter (Molecular Devices, Sunnyvale CA). The whole-cell current was measured on the cells with an access resistance of less than 10 MΩ after the whole-cell configuration was obtained. The amplifier circuitry compensated the whole-cell capacitance. The two-step pulse from 120 to -80 mV for 50 ms was continuously applied to the cell membrane every 5 sec to monitor the activation of the CALHM current. The step pulse from -100 to 140 mV (or –80 to 120 mV) for 200 msec with a holding potential of 0 mV was applied to plot the current-voltage relationship. Electrical signals were digitized at 10 kHz and filtered at 2 kHz. Recordings were analyzed using Clampfit 11.3 (Axon Instruments Inc), GraphPad Prism 10 (La Jolla, CA), and OriginPro 2024 (OriginLab, Northampton, MA). The standard bath solution contains (in mM): 150 NaCl, 5 KCl, 1 $MgCl_2$, 2 $CaCl_2$, 12 Mannitol, 10 HEPES, pH=7.4 with NaOH. For a whole-cell recording, the extracellular solution contains (in mM): 150 NaCl, 10 HEPES, 1 $MgCl_2$, 5 $CaCl_2$. To establish a zero Ca^{2+}

condition, 5 mM $CaCl_2$ was replaced with 5 mM EGTA. For the zero Ca^{2+} condition with 100 μM

Gd^{3+}, 5 mM $CaCl_2$ was omitted entirely without adding additional EGTA. The intracellular

solution contains (in mM): 150 NaCl, 10 HEPES, 1 $MgCl_2$, 5 EGTA.

All data are expressed as mean±SEM. Multiple comparisons were performed by one-way or

two-way ANOVA with Bonferroni's post hoc test. n indicates the number of cells. Significance

was defined as: $*P<0.05$, $**P<0.01$, $***P<0.001$. The absence of an asterisk indicates non-

significance.

821 Data and Code Availability

The human IC annotation table with all the curation information is made available with the

manuscript as Supplementary Table 1. The fasta sequences for the human ICs (both full-length

sequences and the pore domain sequences) are available through Zenodo at doi

10.5281/zenodo.16232527. The full length sequences for all the identified orthologs are also

available through the Zenodo doi link. The code and results related to the RAG annotation

pipeline are available at github: https://github.com/esbgkannan/ionchannels-final-pdf

829 Acknowledgments

830 We thank members of the Kannan Lab for feedback and rotation student William N Lantz for the

831 manual evaluation of RAG LLM-based annotations. Maya Salcedo is acknowledged for

832 onboarding the undergraduates involved in the project. Funding for NK from NIH Common Fund

833 (IDG) U01CA271376 is acknowledged. ZR is supported by 5R00NS128258 from NINDS.

835 Author Contributions

836 Conceptualization: RT, NK, WL. Methodology: RT, NK, SJP, SK, NG, SS, WL. Formal analysis

837 and investigation: RT, SJP, SK, SS, RC, KB, NK. Data curation: RT, RC, KB, SS. Validation:

838 RT, NG, RC, SS, WL, ZR. Writing – original draft: RT, SJP, SS, NK. Writing - review and

839 editing: RT, SS, ZR, WL, NK. Supervision, project administration, and funding acquisition: NK,

840 WL.

Competing Interests Statement

843 The authors declare no competing interests.

Reference:

1.   Como M, Koppala BR, Hasan MN, Han VL, Arora I, Sun D. Cell Volume Regulation in
Immune Cell Function, Activation and Survival. Cell Physiol Biochem. 2021;55(S1):71-88. doi:
10.33594/000000331. PubMed PMID: 33611867.
2.   Meir A, Ginsburg S, Butkevich A, Kachalsky SG, Kaiserman I, Ahdut R, et al. Ion channels
in presynaptic nerve terminals and control of transmitter release. Physiol Rev. 1999;79(3):1019-
88. doi: 10.1152/physrev.1999.79.3.1019. PubMed PMID: 10390521.
3.   Thorneloe KS, Nelson MT. Ion channels in smooth muscle: regulators of intracellular
calcium and contractility. Can J Physiol Pharmacol. 2005;83(3):215-42. doi: 10.1139/y05-016.
PubMed PMID: 15870837.
4.   Rajan AS, Aguilar-Bryan L, Nelson DA, Yaney GC, Hsu WH, Kunze DL, Boyd AE, 3rd. Ion
channels and insulin secretion. Diabetes Care. 1990;13(3):340-63. doi:
10.2337/diacare.13.3.340. PubMed PMID: 1689632.
5.   Edelman A, Saussereau E. [Cystic fibrosis and other channelopathies]. Arch Pediatr.
2012;19 Suppl 1:S13-6. doi: 10.1016/s0929-693x(12)71101-6. PubMed PMID: 22682482.
6.   Kim JB. Channelopathies. Korean J Pediatr. 2014;57(1):1-18. Epub 20140131. doi:
10.3345/kjp.2014.57.1.1. PubMed PMID: 24578711; PubMed Central PMCID:
PMCPMC3935107.
7.   Rivolta I, Binda A, Masi A, DiFrancesco JC. Cardiac and neuronal HCN channelopathies.
Pflugers Arch. 2020;472(7):931-51. Epub 20200518. doi: 10.1007/s00424-020-02384-3. PubMed
PMID: 32424620.
8.   Bagal SK, Brown AD, Cox PJ, Omoto K, Owen RM, Pryde DC, et al. Ion Channels as
Therapeutic Targets: A Drug Discovery Perspective. Journal of Medicinal Chemistry.
2013;56(3):593-624. doi: 10.1021/jm3011433.

9.   Braun N, Sheikh ZP, Pless SA. The current chemical biology tool box for studying ion
channels. J Physiol. 2020;598(20):4455-71. Epub 20200817. doi: 10.1113/jp276695. PubMed
PMID: 32715480.
10.   Wickenden A, Priest B, Erdemli G. Ion Channel Drug Discovery: Challenges And Future
Directions. Future Medicinal Chemistry. 2012;4(5):661-79. doi: 10.4155/fmc.12.4.
11.   Fodor AA, Aldrich RW. Statistical limits to the identification of ion channel domains by
sequence similarity. J Gen Physiol. 2006;127(6):755-66. doi: 10.1085/jgp.200509419. PubMed
PMID: 16735758; PubMed Central PMCID: PMCPMC2151544.
12.   Neuwald AF. Rapid detection, classification and accurate alignment of up to a million or
more related protein sequences. Bioinformatics. 2009;25(15):1869-75. Epub 20090608. doi:
10.1093/bioinformatics/btp342. PubMed PMID: 19505947; PubMed Central PMCID:
PMCPMC2732367.
13.   Ranjan R, Khazen G, Gambazzi L, Ramaswamy S, Hill SL, Schürmann F, Markram H.
Channelpedia: an integrative and interactive database for ion channels. Front Neuroinform.
2011;5:36. Epub 20111230. doi: 10.3389/fninf.2011.00036. PubMed PMID: 22232598; PubMed
Central PMCID: PMCPMC3248699.
14.   Alexander SPH, Mathie AA, Peters JA, Veale EL, Striessnig J, Kelly E, et al. The Concise
Guide to PHARMACOLOGY 2023/24: Ion channels. Br J Pharmacol. 2023;180 Suppl 2(Suppl
2):S145-s222. doi: 10.1111/bph.16178. PubMed PMID: 38123150; PubMed Central PMCID:
PMCPMC11339754.
15.   Castro EV, Shepherd JW, Guggenheim RS, Sengvoravong M, Hall BC, Chappell MK, et al.
ChanFAD: A Functional Annotation Database for Ion Channels. Frontiers in Bioinformatics.
2022;Volume 2 - 2022. doi: 10.3389/fbinf.2022.835805.
16.   Špačková A, Vávra O, Raček T, Bazgier V, Sehnal D, Damborský J, et al. ChannelsDB 2.0: a
comprehensive database of protein tunnels and pores in AlphaFold era. Nucleic Acids Research.
2023;52(D1):D413-D8. doi: 10.1093/nar/gkad1012.
17.   Jegla TJ, Zmasek CM, Batalov S, Nayak SK. Evolution of the human ion channel set. Comb
Chem High Throughput Screen. 2009;12(1):2-23. doi: 10.2174/138620709787047957. PubMed
PMID: 19149488.
18.   Jianzhao G, Zhen M, Zhaopeng Z, Hong W, Lukasz K. Prediction of Ion Channels and their
Types from Protein Sequences: Comprehensive Review and Comparative Assessment. Current
Drug Targets. 2019;20(5):579-92. doi:
http://dx.doi.org/10.2174/1389450119666181022153942.
19.   Li B, Gallin WJ. VKCDB: Voltage-gated potassium channel database. BMC Bioinformatics.
2004;5(1):3. doi: 10.1186/1471-2105-5-3.
20.   Moran Y, Barzilai MG, Liebeskind BJ, Zakon HH. Evolution of voltage-gated ion channels
at the emergence of Metazoa. Journal of Experimental Biology. 2015;218(4):515-25. doi:
10.1242/jeb.110270.
21.   Lara A, Simonson BT, Ryan JF, Jegla T. Genome-Scale Analysis Reveals Extensive
Diversification of Voltage-Gated K+ Channels in Stem Cnidarians. Genome Biology and
Evolution. 2023;15(3). doi: 10.1093/gbe/evad009.
22.   Uribe C, Nery MF, Zavala K, Mardones GA, Riadi G, Opazo JC. Evolution of ion channels in
cetaceans: a natural experiment in the tree of life. Scientific Reports. 2024;14(1):17024. doi:
10.1038/s41598-024-66082-1.

23.   Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids
Res. 2000;28(1):27-30. doi: 10.1093/nar/28.1.27. PubMed PMID: 10592173; PubMed Central
PMCID: PMCPMC102409.
24.   Kanehisa M, Furumichi M, Sato Y, Kawashima M, Ishiguro-Watanabe M. KEGG for
taxonomy-based analysis of pathways and genomes. Nucleic Acids Res. 2023;51(D1):D587-d92.
doi: 10.1093/nar/gkac963. PubMed PMID: 36300620; PubMed Central PMCID:
PMCPMC9825424.
25.   Kelleher KJ, Sheils TK, Mathias SL, Yang JJ, Metzger Vincent T, Siramshetty Vishal B, et al.
Pharos 2023: an integrated resource for the understudied human proteome. Nucleic Acids
Research. 2022;51(D1):D1405-D16. doi: 10.1093/nar/gkac1033.
26.   Seal RL, Braschi B, Gray K, Jones TEM, Tweedie S, Haim-Vilmovsky L, Bruford EA.
Genenames.org: the HGNC resources in 2023. Nucleic Acids Res. 2023;51(D1):D1003-d9. doi:
10.1093/nar/gkac888. PubMed PMID: 36243972; PubMed Central PMCID: PMCPMC9825485.
27.   Kato AS, Bredt DS. Pharmacological regulation of ion channels by auxiliary subunits. Curr
Opin Drug Discov Devel. 2007;10(5):565-72. PubMed PMID: 17786855.
28.   Li Y, Um SY, McDonald TV. Voltage-gated potassium channels: regulation by accessory
subunits. Neuroscientist. 2006;12(3):199-210. doi: 10.1177/1073858406287717. PubMed
PMID: 16684966.
29.   Abbott GW. Kv Channel Ancillary Subunits: Where Do We Go from Here? Physiology
(Bethesda). 2022;37(5):0. doi: 10.1152/physiol.00005.2022. PubMed PMID: 35797055; PubMed
Central PMCID: PMCPMC9394777.
30.   Sharma K, Nadler L. Identifying New Drug Targets by Illuminating the Druggable
Genome. The FASEB Journal. 2021;35(S1). doi:
https://doi.org/10.1096/fasebj.2021.35.S1.01799.
31.   Sheils T, Mathias SL, Siramshetty VB, Bocci G, Bologa CG, Yang JJ, et al. How to
Illuminate the Druggable Genome Using Pharos. Curr Protoc Bioinformatics. 2020;69(1):e92.
doi: 10.1002/cpbi.92. PubMed PMID: 31898878; PubMed Central PMCID: PMCPMC7818358.
32.   Vacher H, Trimmer JS. Diverse roles for auxiliary subunits in phosphorylation-dependent
regulation of mammalian brain voltage-gated potassium channels. Pflügers Archiv - European
Journal of Physiology. 2011;462(5):631-43. doi: 10.1007/s00424-011-1004-8.
33.   Chen GL, Li J, Zhang J, Zeng B. To Be or Not to Be an Ion Channel: Cryo-EM Structures
Have a Say. Cells. 2023;12(14). Epub 20230717. doi: 10.3390/cells12141870. PubMed PMID:
37508534; PubMed Central PMCID: PMCPMC10378246.
34.   Yu FH, Yarov-Yarovoy V, Gutman GA, Catterall WA. Overview of molecular relationships
in the voltage-gated ion channel superfamily. Pharmacol Rev. 2005;57(4):387-95. doi:
10.1124/pr.57.4.13. PubMed PMID: 16382097.
35.   Huffer KE, Aleksandrova AA, Jara-Oseguera A, Forrest LR, Swartz KJ. Global alignment
and assessment of TRP channel transmembrane domain structures to explore functional
mechanisms. eLife. 2020;9:e58660. doi: 10.7554/eLife.58660.
36.   Shrestha S, Byrne DP, Harris JA, Kannan N, Eyers PA. Cataloguing the dead: breathing
new life into pseudokinase research. Febs j. 2020;287(19):4150-69. Epub 20200310. doi:
10.1111/febs.15246. PubMed PMID: 32053275; PubMed Central PMCID: PMCPMC7586955.
37.   Picado A, Chaikuad A, Wells CI, Shrestha S, Zuercher WJ, Pickett JE, et al. A Chemical
Probe for Dark Kinase STK17B Derives Its Potency and High Selectivity through a Unique P-Loop

Conformation. J Med Chem. 2020;63(23):14626-46. Epub 20201120. doi: 10.1021/acs.jmedchem.0c01174. PubMed PMID: 33215924; PubMed Central PMCID: PMCPMC7816213.

38.   Taujale R, Venkat A, Huang L-C, Zhou Z, Yeung W, Rasheed KM, et al. Deep evolutionary analysis reveals the design principles of fold A glycosyltransferases. eLife. 2020;9:e54532. doi: 10.7554/eLife.54532.

39.   Taujale R, Zhou Z, Yeung W, Moremen KW, Li S, Kannan N. Mapping the glycosyltransferase fold landscape using interpretable deep learning. Nature Communications. 2021;12(1):5656. doi: 10.1038/s41467-021-25975-9.

40.   Neuwald AF. A Bayesian sampler for optimization of protein domain hierarchies. J Comput Biol. 2014;21(3):269-86. Epub 20140204. doi: 10.1089/cmb.2013.0099. PubMed PMID: 24494927; PubMed Central PMCID: PMCPMC3948484.

41.   UniProt Consortium T. UniProt: the universal protein knowledgebase. Nucleic Acids Res. 2018;46(5):2699. doi: 10.1093/nar/gky092. PubMed PMID: 29425356; PubMed Central PMCID: PMCPMC5861450.

42.   Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. J Mol Biol. 2001;305(3):567-80. doi: 10.1006/jmbi.2000.4315. PubMed PMID: 11152613.

43.   Käll L, Krogh A, Sonnhammer EL. Advantages of combined transmembrane topology and signal peptide prediction--the Phobius web server. Nucleic Acids Res. 2007;35(Web Server issue):W429-32. Epub 20070505. doi: 10.1093/nar/gkm256. PubMed PMID: 17483518; PubMed Central PMCID: PMCPMC1933244.

44.   Sehnal D, Svobodová Vařeková R, Berka K, Pravda L, Navrátilová V, Banáš P, et al. MOLE 2.0: advanced approach for analysis of biomacromolecular channels. J Cheminform. 2013;5(1):39. Epub 20130816. doi: 10.1186/1758-2946-5-39. PubMed PMID: 23953065; PubMed Central PMCID: PMCPMC3765717.

45.   Gurnett CA, Campbell KP. Transmembrane auxiliary subunits of voltage-dependent ion channels. J Biol Chem. 1996;271(45):27975-8. doi: 10.1074/jbc.271.45.27975. PubMed PMID: 8910401.

46.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, Madden TL. BLAST+: architecture and applications. BMC Bioinformatics. 2009;10(1):421. doi: 10.1186/1471-2105-10-421.

47.   Eddy SR. Profile hidden Markov models. Bioinformatics. 1998;14(9):755-63. doi: 10.1093/bioinformatics/14.9.755. PubMed PMID: 9918945.

48.   Lin Z, Akin H, Rao R, Hie B, Zhu Z, Lu W, et al. Evolutionary-scale prediction of atomic-level protein structure with a language model. Science. 2023;379(6637):1123-30. Epub 20230316. doi: 10.1126/science.ade2574. PubMed PMID: 36927031.

49.   Llinares-López F, Berthet Q, Blondel M, Teboul O, Vert J-P. Deep embedding and alignment of protein sequences. Nature Methods. 2023;20(1):104-11. doi: 10.1038/s41592-022-01700-2.

50.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. Nat Genet. 2000;25(1):25-9. doi: 10.1038/75556. PubMed PMID: 10802651; PubMed Central PMCID: PMCPMC3037419.

1001 51. Huang LC, Taujale R, Gravel N, Venkat A, Yeung W, Byrne DP, et al. KinOrtho: a method
1002 for mapping human kinase orthologs across the tree of life and illuminating understudied
1003 kinases. BMC Bioinformatics. 2021;22(1):446. Epub 20210918. doi: 10.1186/s12859-021-04358-
1004 3. PubMed PMID: 34537014; PubMed Central PMCID: PMCPMC8449880.
1005 52. Monahan-Earley R, Dvorak AM, Aird WC. Evolutionary origins of the blood vascular
1006 system and endothelium. J Thromb Haemost. 2013;11 Suppl 1(Suppl 1):46-66. doi:
1007 10.1111/jth.12253. PubMed PMID: 23809110; PubMed Central PMCID: PMCPMC5378490.
1008 53. Ma Z, Tanis JE, Taruno A, Foskett JK. Calcium homeostasis modulator (CALHM) ion
1009 channels. Pflugers Arch. 2016;468(3):395-403. Epub 20151125. doi: 10.1007/s00424-015-1757-
1010 6. PubMed PMID: 26603282; PubMed Central PMCID: PMCPMC5308871.
1011 54. Ma Z, Taruno A, Ohmoto M, Jyotaki M, Lim JC, Miyazaki H, et al. CALHM3 Is Essential for
1012 Rapid Ion Channel-Mediated Purinergic Neurotransmission of GPCR-Mediated Tastes. Neuron.
1013 2018;98(3):547-61.e10. Epub 20180419. doi: 10.1016/j.neuron.2018.03.043. PubMed PMID:
1014 29681531; PubMed Central PMCID: PMCPMC5934295.
1015 55. Ma Z, Siebert AP, Cheung KH, Lee RJ, Johnson B, Cohen AS, et al. Calcium homeostasis
1016 modulator 1 (CALHM1) is the pore-forming subunit of an ion channel that mediates
1017 extracellular Ca2+ regulation of neuronal excitability. Proc Natl Acad Sci U S A.
1018 2012;109(28):E1963-71. Epub 20120618. doi: 10.1073/pnas.1204023109. PubMed PMID:
1019 22711817; PubMed Central PMCID: PMCPMC3396471.
1020 56. Tanis JE, Ma Z, Krajacic P, He L, Foskett JK, Lamitina T. CLHM-1 is a functionally
1021 conserved and conditionally toxic Ca2+-permeable ion channel in Caenorhabditis elegans. J
1022 Neurosci. 2013;33(30):12275-86. doi: 10.1523/jneurosci.5919-12.2013. PubMed PMID:
1023 23884934; PubMed Central PMCID: PMCPMC3721838.
1024 57. Danielli S, Ma Z, Pantazi E, Kumar A, Demarco B, Fischer FA, et al. The ion channel
1025 CALHM6 controls bacterial infection-induced cellular cross-talk at the immunological synapse.
1026 Embo j. 2023;42(7):e111450. Epub 20230302. doi: 10.15252/embj.2022111450. PubMed PMID:
1027 36861806; PubMed Central PMCID: PMCPMC10068325.
1028 58. Tanis JE, Ma Z, Foskett JK. The NH(2) terminus regulates voltage-dependent gating of
1029 CALHM ion channels. Am J Physiol Cell Physiol. 2017;313(2):C173-c86. Epub 20170517. doi:
1030 10.1152/ajpcell.00318.2016. PubMed PMID: 28515089; PubMed Central PMCID:
1031 PMCPMC5582874.
1032 59. Ma Z, Paudel U, Wang M, Foskett JK. A mechanism of CALHM1 ion channel gating. Am J
1033 Physiol Cell Physiol. 2025;328(4):C1109-c24. Epub 20250221. doi: 10.1152/ajpcell.00925.2024.
1034 PubMed PMID: 39981825; PubMed Central PMCID: PMCPMC12164053.
1035 60. Choi W, Clemente N, Sun W, Du J, Lü W. The structures and gating mechanism of human
1036 calcium homeostasis modulator 2. Nature. 2019;576(7785):163-7. doi: 10.1038/s41586-019-
1037 1781-3.
1038 61. Gudmundsson S, Singer-Berk M, Watts NA, Phu W, Goodrich JK, Solomonson M, et al.
1039 Variant interpretation using population databases: Lessons from gnomAD. Human Mutation.
1040 2022;43(8):1012-30. doi: https://doi.org/10.1002/humu.24309.
1041 62. Kwon JW, Jeon YK, Kim J, Kim SJ, Kim SJ. Intramolecular Disulfide Bonds for Biogenesis of
1042 CALHM1 Ion Channel Are Dispensable for Voltage-Dependent Activation. Mol Cells.
1043 2021;44(10):758-69. doi: 10.14348/molcells.2021.0131. PubMed PMID: 34711692; PubMed
1044 Central PMCID: PMCPMC8560582.

63.   Jeon YK, Choi SW, Kwon JW, Woo J, Choi SW, Kim SJ, Kim SJ. Thermosensitivity of the
voltage-dependent activation of calcium homeostasis modulator 1 (calhm1) ion channel.
Biochem Biophys Res Commun. 2021;534:590-6. Epub 20201114. doi:
10.1016/j.bbrc.2020.11.035. PubMed PMID: 33199024.
64.   Syrjänen JL, Epstein M, Gómez R, Furukawa H. Structure of human CALHM1 reveals key
locations for channel regulation and blockade by ruthenium red. Nat Commun.
2023;14(1):3821. Epub 20230628. doi: 10.1038/s41467-023-39388-3. PubMed PMID:
37380652; PubMed Central PMCID: PMCPMC10307800.
65.   Syrjanen JL, Michalski K, Chou TH, Grant T, Rao S, Simorowski N, et al. Structure and
assembly of calcium homeostasis modulator proteins. Nat Struct Mol Biol. 2020;27(2):150-9.
Epub 20200127. doi: 10.1038/s41594-019-0369-9. PubMed PMID: 31988524; PubMed Central
PMCID: PMCPMC7015811.
66.   Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, et al. CDD/SPARCLE: the
conserved domain database in 2020. Nucleic Acids Res. 2020;48(D1):D265-d8. doi:
10.1093/nar/gkz991. PubMed PMID: 31777944; PubMed Central PMCID: PMCPMC6943070.
67.   Mistry J, Chuguransky S, Williams L, Qureshi M, Salazar GA, Sonnhammer ELL, et al.
Pfam: The protein families database in 2021. Nucleic Acids Res. 2021;49(D1):D412-d9. doi:
10.1093/nar/gkaa913. PubMed PMID: 33125078; PubMed Central PMCID: PMCPMC7779014.
68.   Henschel A, Winter C, Kim WK, Schroeder M. Using structural motif descriptors for
sequence-based binding site prediction. BMC Bioinformatics. 2007;8 Suppl 4(Suppl 4):S5. Epub
20070522. doi: 10.1186/1471-2105-8-s4-s5. PubMed PMID: 17570148; PubMed Central PMCID:
PMCPMC1892084.
69.   Letunic I, Khedkar S, Bork P. SMART: recent updates, new developments and status in
2020. Nucleic Acids Research. 2020;49(D1):D458-D60. doi: 10.1093/nar/gkaa937.
70.   Großberger LMaJHaNSaL. UMAP: Uniform Manifold Approximation and Projection.
Journal of Open Source Software. 2018;3(29):861.
71.   Hernández-Salmerón JE, Moreno-Hagelsieb G. Progress in quickly finding orthologs as
reciprocal best hits: comparing blast, last, diamond and MMseqs2. BMC Genomics.
2020;21(1):741. doi: 10.1186/s12864-020-07132-6.
72.   Li L, Stoeckert CJ, Jr., Roos DS. OrthoMCL: identification of ortholog groups for
eukaryotic genomes. Genome Res. 2003;13(9):2178-89. doi: 10.1101/gr.1224503. PubMed
PMID: 12952885; PubMed Central PMCID: PMCPMC403725.
73.   Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy
1.0: fundamental algorithms for scientific computing in Python. Nature Methods.
2020;17(3):261-72. doi: 10.1038/s41592-019-0686-2.
74.   The Gene Ontology resource: enriching a GOld mine. Nucleic Acids Res.
2021;49(D1):D325-d34. doi: 10.1093/nar/gkaa1113. PubMed PMID: 33290552; PubMed Central
PMCID: PMCPMC7779012.
1083
1084
1085

Figures and Tables

Identifier labels		Structure related labels			UniProt TM		
Uniprot	P29972	PDB ID	8CT2		# of TMs	6	
Name	Aquaporin-1	Alphafold ID			# of Transmembranes (TMs) + Intramembranes (IMs)	10	
Symbol	AQP1 (CHIP28)	**Complex/Interaction related labels**			TM Start (UniProt)	8	
Target Development level (Pharos)	Tbio	Auxiliary	No		TM End (UniProt)	228	
Length	269	Characterized domains	MIP		TMsList	T:8-36,T:49-66,I:71-76,I:77-84,T:95-	
Classification labels		Auxiliary domain	no		TM Lit resource	https://scholar.google.co	
Family designation	Aquaporin	Auxiliary protein			Lit based TMstart_1	8	
Group	Other	Notable interactors	EPHB2		Lit based TMend _1	36	
Class		**TM and pore domain related labels**			Lit based TMstart_2	49	
Family	Aquaporin	Pore Domain start	8		Lit based TMend_2	66	
Sub Family		Pore Domain end	228		Lit based TMstart_3	95	
Functional labels		Domain length	220		Lit based TMend_3	115	
Unit	pore-containing	Does it pass through membrane at least once	Yes		Lit based TMstart_4	137	
Ion	water	# of TM domains (predicted by TMHMM)	6		Lit based TMend_4	155	
Gate Mechanism	ligand-gated (cGMP)	# of TM domain (predicted by Phobius)	6		Lit based TMstart_5	167	
Lit Resource	PMID: 26365508, PMID: 16962972	TM organization	1|2|3|4|5|6		Lit based TMend_5	183	
Uniprot Function	Form water-specific channel / plasma membranes of red cells and kidney proximal tubules	MOLE pore residue first	35		Lit based TMstart_6	208	
		MOLE pore residue last	185		Lit based TMend_6	228	

(Right sections labeled: "UniProt TM" and "Literature search based TM labels"; lower-middle section labeled "MOLE".)

Table 1: List of features annotated for the collected IC sequences. The labels are colored by their annotation category. Labels for Aquaporin 1 are shown as examples of each annotation label.

Figure 1: Distribution of human ICs across different families. Each circle represents a human IC family, with the symbol at the center indicating its Group. The size of the circles is proportional to the number of sequences in that family, and the colored pies indicate the proportion of their

1098 TDL status as designated by IDG. The placement of the bubbles is based on the average co-

1099 ordinates of all the members within that family in a distribution of UMAP embeddings generated

1100 using protein embedding-based pairwise sequence alignments. An embedding-based sequence

1101 alignment approach was used to overcome the vast divergence of IC sequences with minimal

1102 sequence similarity. A family level abstraction was done to provide an intuitive view undeterred

1103 by the relationships of individual ICs across families. A detailed view of the full UMAP plot

1104 showing the placement of individual ICs is provided in Supplementary Figure 3 for reference.

1105 Auxiliary IC families at the bottom were not part of the embedding-based analysis due to the

1106 lack of a pore-containing domain, and thus placed arbitrarily at the bottom of the figure.

1107

1108

Figure 2: Orthology profiling of human ICs. (A) Heatmap showing the percent of orthologs

detected for each IC family within a given taxonomic lineage. The taxonomic groups are shown

in the vertical axis with a tree on the left. Darker color represents a higher percentage of

orthologs detected. Percentages were calculated as (total number of orthologs found for all ICs

in a family)/(total number of organisms queried in the taxonomic lineage * number of sequences

in the family) (B) Clustergram depicting the presence/absence of orthologous sequences of ICs

across eukaryotic taxonomic lineages. ICs are clustered along the horizontal axis into 9 distinct

clusters. Taxonomic groups are shown on the vertical axis. Each square in the heatmap is

colored based on the orthology relationship found for a specific IC in a specific organism (black:

one-to-one ortholog present, red: co-ortholog detected, brown: no orthology detected). (C)

Results from the enrichment analysis performed on human ICs of each cluster. The x-axis

1120 shows the number of ICs in the cluster enriched for the GO term shown on the y-axis. The bars

1121 are colored based on their FDR values for the enriched term. For a full list of enriched terms,

1122 please refer to Supplementary Table 4.

1123

1124

Figure 3: Evolutionary analysis of CALHM reveals conserved pattern positions. (A) A

phylogenetic tree depicting the evolutionary relationships across all 6 CALHM members with

orthologs across different taxa. (B) The conserved pattern positions conserved across all 6

CALHM members are shown as a weblogo with the red bars indicating the significance (longer

bars indicate higher significance) of conservation and are mapped into a representative

structure of human CALHM2. The four transmembrane helices are labeled S1-S4. Conserved

residues that are targeted for mutation are highlighted using a asterisk symbol in the labels. (C)

Schematic representing the location of transmembrane regions and identified conserved pattern

positions in a representative human CALHM2 sequence. The residues targeted for mutations

are labelled with their corresponding positions for CALHM2, 1 and 6 shown in the labels. (D)

Cartoon representation of CALHM2 structure (PDB ID: 6uiv and 6uiw) in open and closed

conformation, respectively. (E) List of disease variants and mutations performed in the

conserved pattern positions for functional studies.

1138

Figure 4. Electrophysiological studies of human CALHM1 and CALHM6. Whole-cell voltage-

clamp recordings were performed in tsA cells overexpressing wild-type CALHM1 (A–D) and

wild-type CALHM6 (E–H), as well as in non-transfected cells (I–L). Currents were measured

under three conditions sequentially from the same cell: 5 mM Ca^{2+} at 22 °C (a, e, i; black), 0 mM

Ca^{2+} at 22 °C (B,F,J; blue), and 0 mM Ca^{2+} at 37 °C (C,G,K; red). Voltage steps ranged from

–100 mV to +140 mV, followed by a final tail pulse at –100 mV, with a holding potential of 0 mV

(protocol illustrated in the box on the right). Current-voltage (I–V) relationships were plotted in D,

H, and L using mean current amplitudes (averaged across independent cells) measured at the

end of the voltage steps. The arrow indicates the time point at which current amplitudes were

measured. Error bars represent SEM (D, n=5; H, n=5; L, n=5)

Figure 5: Functional characterization of CALHM1 and CALHM6 mutants at conserved residues

at 37 °C. Whole-cell voltage-clamp recordings were performed in tsA cells overexpressing wild-

type CALHM1 (A-C), CALHM1 mutant (I109W) (D-F), wild-type CALHM6 (G-I), and CALHM6

mutants (Y51A (J-L)and W113A(M-O)). Currents were measured under two conditions: 0 mM

Ca^2 at 37 °C (A,D,G,J,M; red) and 0 mM Ca^2 at 37°C plus 100 µM Gd^3 (B,E,H,K,N; blue),

with both conditions recorded sequentially from the same cell. Voltage steps ranged from

−80 mV to +120 mV, followed by a final tail pulse at −80 mV, with a holding potential of 0 mV

(protocol shown in the box on the right). Current-voltage (I–V) relationships were plotted in

C,F,I,L, and O using mean current amplitudes (averaged across independent cells) measured at

the end of the voltage steps. The arrow indicates the time point at which the current was

measured. The number of independent measurements (cells) were: n = 5 (C); n = 5 (F); n = 5

(I); n = 5 (L); n=5 (O). Error bars represent SEM. (P,Q) Current amplitudes obtained using a

two-step voltage protocol (from +120 mV to −80 mV; protocol shown in the box on the right) are

compared between wild-type CALHM1 and its mutants (P) and between wild-type CALHM6 and

its mutants (Q). Each dot represents an independent measurement (cell), and bar represents

the mean current amplitude across cells. The number of independent measurements (cells) for

each bar in P and Q are shown from left to right: 5, 8, 5, 8, 6, 7, 5, 5, 5, 6, 7, 7 (P); 5, 6, 5, 7, 7,

5, 6, 6 (Q). Statistical analysis was performed using one-way ANOVA with Bonferroni's post hoc

test, comparing each mutant to wild type (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$).

1169

1170 **Supplementary Figures**

1171
1172 **Supplementary Figure 1:** RAG annotation pipeline used for validating the annotation of ion
1173 specificity and gating mechanism

1174

1175
1176 **Supplementary Figure 2:** Upset plot showing the overlap of IC sequences based on their
1177 UniProt IDs across the current study, the KEGG database, GtoP and Pharos.
1178
1179

1180
1181
Supplementary Figure 3: UMAP embeddings plot of all human IC sequences. The UMAP
embeddings were generated based on the pairwise similarity scores of the protein embedding
alignment generated for all pairs of human ICs. Shapes of the markers indicate the different IC
groups (circle: VGIC, square: LGIC, diamond: Chloride channel, triangle: Other, plus:
Unclassified) and the colors indicate different IC families within each group as shown in the
legend above.

1188
1189

1190
1191
Supplementary Figure 4. Expression analysis of wild-type CALHM1, wild-type CALHM6, and
their mutants. (A,B) Representative gels showing CALHM1 and its mutants (A), and CALHM6
and its mutants (B). CALHM1 and CALHM6 signals were detected using in-gel fluorescence of

the C-terminal GFP tag, while β-actin was detected by Western blotting as a loading control.
(C,D) Quantification of total protein expression levels of wild-type CALHM1 and its mutants (C),
and wild-type CALHM6 and its mutants (D). Each dot represents an independent measurement
(transfection) and error bars represent SEM (c, n = 3; d, n = 3). Statistical analysis was
performed using one-way ANOVA with Bonferroni's post hoc test, comparing each mutant to
wild type (*p < 0.05; **p < 0.01; ***p < 0.001). (E,F) Surface biotinylation assays of wild-type
CALHM1 and its mutants (E), and wild-type CALHM6 and its mutants (F), detected using the C-
terminal GFP tag by in-gel fluorescence.
1203

1205
1206
Supplementary Figure 5: Functional characterization of CALHM1 and CALHM6 mutants at
conserved residues at 22 °C. Current amplitudes obtained using a two-step voltage protocol
(from +120 mV to –80 mV; protocol shown in the box on the right) are compared between wild-
type CALHM1 and its mutants (A, B) and between wild-type CALHM6 and its mutants (C, D).
The cells analyzed here are the same as those in Figure 5P, Q. Briefly, for each cell, currents
were measured sequentially under three conditions: 5 mM Ca^{2+} at 22 °C, 0 mM Ca^{2+} at 22 °C,
and 0 mM Ca^{2+} at 37 °C. The currents from the first two conditions are plotted here, while
currents at 0 mM Ca^{2+} at 37 °C are shown in Figure 5R, S. Each dot represents an independent
measurement (cell), and bar represents the mean current amplitude across cells. The number of
independent measurements (cells) for each bar in a–d are shown from left to right: 5, 8, 5, 8, 6,
7, 5, 5, 5, 6, 7, 7 (A, B); 5, 6, 5, 7, 7, 6, 6, 5 (C, D). Statistical analysis was performed using one-
way ANOVA with Bonferroni's post hoc test, comparing each mutant to wild type (*$p < 0.05$;
**$p < 0.01$; ***$p < 0.001$).
1220
1221

1222
1223
Supplementary Figure 6: Conserved pattern positions identified within the clade for
CALHM2,4,5 and 6. A) Phylogenetic tree of CALHM sequences where the orange star indicates
the clade for CALHM2,4,5 and 6. B) The identified pattern positions are mapped into a
representative structure of human CALHM2 (PDB: 6uiw). C) Weblogo showing the conserved
pattern positions. The red bar indicates the significance of conservation where a taller bar
indicates higher significance.
1230