Spatial dynamics of the tumor microenvironment in emerging resistance to

targeted therapy in *EGFR*-mutated NSCLC

Satoshi Nakamura[1] Daisuke Shibahara[1,*], Kentaro Tanaka[1,2], Yasuyuki Kishikawa[1],

Mikiko Hashisako[3], Keita Nakatomi[4], Noriaki Nakagaki[5], Mikihiro Kohno[6], Koichi

Azuma[7], Ritsu Ibusuki[1], Kohei Otsubo[1], Yasuto Yoneshima[1], Eiji Iwama[1], Yoshinao

Oda[3], Isamu Okamoto[1]

[1]Department of Respiratory Medicine, Graduate School of Medical Sciences, Kyushu

University, Fukuoka, Japan

[2]Department of Pulmonary Medicine, Graduate School of Medical and Dental Sciences,

Kagoshima University, Kagoshima, Japan

[3]Department of Anatomic Pathology, Graduate School of Medical Sciences, Kyushu

University, Fukuoka, Japan

[4]Department of Respiratory Medicine, Kyushu Central Hospital, Fukuoka, Japan

[5]Department of Respiratory Medicine, Fukuoka Red Cross Hospital, Fukuoka, Japan

[6]Department of Surgery and Science, Graduate School of Medical Sciences, Kyushu

University, Fukuoka, Japan

[7]Division of Respirology, Neurology, and Rheumatology, Department of Internal

Medicine, Kurume University School of Medicine, Fukuoka, Japan

*Corresponding Author: Daisuke Shibahara

Email address: shibahara.daisuke.805@m.kyushu-u.ac.jp

Postal address: 3-1-1 Maidashi, Higashi-ku, Fukuoka 812-8582, Japan

Disclosure

Daisuke Shibahara has received research funding from Eli Lilly Japan K.K. Koichi

Azuma has received honoraria from AstraZeneca, MSD, Chugai Pharmaceutical, Ono

Pharmaceutical, Bristol-Myers Squibb, Takeda Pharmaceutical, Taiho Pharmaceutical,

and Amgen. Yasuto Yoneshima has received honoraria from AstraZeneca. Kentaro

Tanaka has received honoraria from Chugai Pharmaceutical, AstraZeneca, Ono

Pharmaceutical, Bristol-Myers Squibb, Eli Lilly, Takeda Pharmaceutical, Novartis

Pharma K.K., Merck Biopharma, Kyowa Kirin, Daiichi-Sankyo, Pfizer, Amgen,

Janssen Pharmaceutical K.K., and MSD, and is an advisory board member for Pfizer

and Novartis Pharma K.K. Eiji Iwama has received honoraria from AstraZeneca and

Chugai Pharmaceutical. Isamu Okamoto has received honoraria and research funding

from Daiichi Sankyo, Chugai Pharmaceutical, Eli Lilly Japan K.K., AstraZeneca, Taiho

Pharmaceutical, Boehringer Ingelheim, and Ono Pharmaceutical; honoraria from

Takeda Pharmaceutical and Novartis Pharma K.K.; and research funding from Bristol-

Meyers Squibb and MSD Oncology. Other authors declare no potential conflicts of

interest.

__**Abstract**__

Epidermal growth factor receptor tyrosine kinase inhibitors (EGFR-TKIs) provide

therapeutic benefit in *EGFR*-mutated non-small cell lung cancer (NSCLC); however,

early resistance develops in some patients. To explore the underlying mechanisms, we

performed spatial transcriptomics using Visium HD and StarDist processing on paired

pre- and post-treatment samples from the same patients. In early resistance samples,

fibroblasts increased in proportion and exhibited enhanced activation following EGFR-

TKI treatment, while a distinct tumor cell cluster showed TNF-α/NF-kB pathway

activation even prior to treatment. Furthermore, in the pre-treatment sample with early

resistance, specific tumor clusters interacted with immune and stromal cells through

inflammatory pathways, including MIF/CD74 and PDGF signaling. Immature tertiary

lymphoid structures (TLSs) were enriched in early resistance, whereas mature TLSs

were observed in long-term responders. These findings highlight the role of tumor

heterogeneity and an inflammatory tumor microenvironment in early EGFR-TKI

resistance, providing insights into potential therapeutic strategies to improve treatment

outcomes.

__Introduction__

Epidermal growth factor receptor (EGFR) tyrosine kinase inhibitors (EGFR-TKIs)

demonstrate durable therapeutic efficacy in the majority of patients with *EGFR*-mutated

non-small cell lung cancer (NSCLC)[1]. Nevertheless, resistance to EGFR-TKIs

inevitably emerges over time after treatment initiation[2,3]. While several resistance

mechanisms to EGFR-TKIs have been identified, the underlying causes of resistance

remain unknown in 40-50% of cases[4–6]. Furthermore, some patients develop early

resistance during treatment, whereas others achieve long-term responses, yet the reasons

for these divergent outcomes remain unclear. Therefore, there is a critical need to

elucidate the precise mechanisms of resistance not only to develop effective therapies

for overcoming resistance but also to extend the duration of response to EGFR-TKIs.

Increasing evidence suggests that the tumor microenvironment (TME) plays a pivotal

role in shaping the response to EGFR-TKIs[7–12]; however, spatial analyses of TME using

paired pre- and post-treatment samples from the same patients with *EGFR*-mutated

NSCLC have been highly limited[13].

Spatial transcriptomics has emerged as a powerful approach to unravel the

TME[13–18]. As an array-based method, Visium HD allows comprehensive mRNA capture

without limiting the analysis to specific target genes, as well as allowing analysis of the

entire sample, compared to in situ hybridization and sequencing methods. In this study,

we conducted a detailed analysis of the TME in paired pre- and post-treatment samples

from patients with *EGFR*-mutated NSCLC using Visium HD. Furthermore, we

converted Visium HD spot data into single cell data using StarDist, a deep learning-

based method for nuclei/cell detection and segmentation. By integrating the spatial

coordinates derived from StarDist into the single-cell data, we conducted spatial

transcriptomic analysis at the single-cell level while preserving the actual spatial

relationships between cells. Our findings elucidate the spatial dynamics of TME,

highlighting the contribution of an inflammatory TME and tumor heterogeneity to the

early onset of resistance mechanisms in *EGFR*-mutated NSCLC.

__Result__

Integrative analysis of spatial transcriptomics and nuclear segmentation

To investigate differences in the duration of response to osimertinib, a third-generation

EGFR-TKI, we collected and analyzed paired pre- and post-treatment samples from

four patients with *EGFR*-mutated NSCLC (Fig. 1a and Extended Data Fig. 1). Clinical

characteristics of patients are listed in Supplementary Table 1. Based on each patient's

progression-free survival duration[2], the samples were categorized into one early

resistance (ER), one standard response (SR), and two long-term response (LR) groups.

All samples were processed using Visium HD platform, which provides a spatial

resolution of 2 μm per spot. To reconstruct single cells from the data, we employed

StarDist[19], a deep learning-based tool, which enables the detection and segmentation of

nuclei from hematoxylin and eosin (H&E) stained images generated alongside the

spatial transcriptomics data (Fig 1b and Extended Data Fig. 2). Using StarDist, we

aggregated 2 μm spots that overlapped with the polygonal coordinates corresponding to

the detected nuclei, thereby reconstructing spatial transcriptomics data mapped to

individual cell positions (Fig. 1c). Additionally, we incorporated the centroid

coordinates of each polygon as new spot coordinates into the data structure to ensure

compatibility with existing tools for spatial transcriptomics analysis (Extended Data Fig.

3a-c).

A total of 814,636,020 reads was obtained from Slide 1 (ER, SR), and

730,477,318 reads were obtained from Slide 2 (LR1, LR2). StarDist identified 141,324

nuclei in Slide 1 and 99,658 nuclei in Slide 2. Samples were partitioned based on

coordinates from their location on the image, and the sample names were linked to the

data (Extended Data Fig. 3d). The integrated data for these eight samples were then

processed for quality control, filtering, normalization, dimensionality reduction, batch

correction, nearest neighbor graph building, and Leiden clustering, and visualized using

Uniform Manifold Approximation and Projection (UMAP). The cell population was

broadly categorized into three main lineages: epithelial cells, stromal cells, and immune

cells, with a small number of blood cells (Fig. 1d and Extended Data Fig. 3e). Within

the epithelial cell population, tumor cells, basal cells, club cells, alveolar cell type 2 and

goblet/ciliated cells were further divided (Fig. 1e and Extended Data Fig. 4a). Immune

and stromal cells were clustered into lymphocytes, plasmablasts, macrophages,

fibroblasts, endothelial cells, and smooth muscle cells (Fig. 1f,g and Extended Data Fig.

4b-c). Additionally, lymphocytes were subdivided, plasmablasts were classified by the

immunoglobin they produced, and macrophages were distinguished from mixed cells

(Extended Data Fig. 4d-i). The results of these clustering analyses were integrated with

the polygonal coordinate data obtained from StarDist, enabling the visualization of

distinct cell populations on spatial coordinates (Fig. 1h-k and Extended Data Fig. 5,6).

Changes of TME between pre- and post-treatment samples

To investigate changes in tumor cell characteristics before and after treatment, we

performed differentially gene expression (DGE) analysis following pseudo-bulk

processing, comparing tumor cells from pre- and post-treatment samples (Fig. 2a).

Genes with significantly higher expression in the pre-treatment group demonstrated a

tendency toward favorable prognosis (Extended Data Fig. 7a-d), whereas genes with

higher expression in the post-treatment group were associated with poorer prognosis

(Extended Data Fig. 7e-h). Gene Set Enrichment Analysis (GSEA) revealed significant

enrichment in gene sets associated with cell cycle and proliferation in post-treatment

tumor cells (Fig. 2b,c and Extended Data Fig. 7i,j). These results indicate that tumor

cells exhibit enhanced proliferative activity after EGFR-TKI resistance.

We next compared the proportion of stromal and immune cell clusters between

pre- and post-treatment samples. The proportion of stromal cells, including fibroblasts

and endothelial cells, was elevated in the ER_post sample compared to the ER_pre

sample (Fig. 2d). DGE analysis of fibroblasts from ER samples revealed increased

expression of collagen-related genes following treatment. (Fig. 2e). GSEA further

demonstrated that a gene set related to extracellular matrix production was significantly

enriched in fibroblasts from the ER_post sample (Fig. 2f). Additionally, collagen gene

expressions were significantly elevated in ER_post sample compared to the other

samples (Fig 2g and Extended Data Fig. 7k). These results suggest that fibroblasts

exhibit increased activity following EGFR-TKI treatment, contributing to changes in the

TME associated with early resistance.

In the T cell cluster, CD4-expressing T cells were present in slightly higher

numbers, while the low expression levels of CD4, CD8A, and FOXP3 impeded their

classification (Fig. 2h and Extended Data Fig. 7l). This could be attributed to the

filtering effect of the analytical methods employed. In the macrophage cluster, tumor-

associated macrophage markers (SPP1, C1QC) were highly expressed in accordance

with the distribution of M2-like markers (CD163, MSR1)[20–22], suggesting an immune-

suppressive phenotype that was consistently observed across all samples. (Fig. 2i and

Extended Data Fig. 7m).

Spatial distribution of tumor cell heterogeneity between early resistance and long-

term response samples

To explore factors contributing to early resistance, tumor cells from all pre-treatment

samples were categorized into two groups: ER_pre and SLR_pre (comprising SR_pre,

LR1_pre, and LR2_pre) for DGE analysis (Fig. 3a). GSEA revealed significant

enrichment in the inflammatory response, TNF-alpha signaling via NF-κB pathway, IL-

6 pathway, and chemokine signaling pathways in ER_pre tumor cells (Fig. 3b,c and

Extended Data Fig. 8a,b), indicating that the TME of ER_pre was characterized by

active inflammation.

To better understand the factors that contribute the differences between samples,

we performed a comparative analysis of tumor cell clusters. Based on differentially

expressed genes, we classified and labeled each tumor cell cluster using cancer-related

genes[23–28] as discriminatory markers (Fig. 3d,e and Extended Data Fig.8c,d). Applying

inferCNV, copy number variation (CNV) analysis revealed a tendency for consistent

CNV alterations in each tumor cell cluster when compared to epithelial cell clustering

as a reference group (Extended Data Fig. 8e). To quantify the distribution, we

visualized the tumor cluster across samples and mapped it spatially for each sample (Fig.

3f,g). The tc_S100A6 and tc_AQP1 clusters were present in all pre- and post-treatment

samples, accounting for a high proportion in nearly all samples. The tc_S100A6 cluster

exhibited significantly higher expressions of *CCND1* and *ERBB2* compared to the other

clusters (Extended Data Fig. 8f,g), while *MET* and *CD74* expressions were markedly

elevated in the tc_AQP1 and tc_TMSB4X cluster, respectively (Extended Data Fig.

8h,i). GSEA based on DGE analysis of the tc_S100A6 cluster revealed significant

enrichment of gene sets related to cell proliferation, including E2F targets and G2-M

checkpoint, compared to the others (Fig. 4a,b). In contrast, gene sets linked to

inflammatory response, epithelial mesenchymal transition (EMT), and the KRAS

signaling were not enriched (Fig. 4c). DGE analysis and GSEA of the tc_AQP1 cluster

demonstrated upregulation of pathways associated with mTORC1 and IL6/JAK/STAT3

signaling, both of which have been implicated in EGFR-TKI resistance[4,29–31], in

addition to EMT and inflammatory responses (Fig. 4d-f).

Next, we analyzed distinct clusters that showed differences compared to the

other samples. The tc_TMSB4X cluster was predominantly observed in ER_pre, with a

lower frequency in other samples, suggesting its potential involvement in early

resistance (Fig. 3f). GSEA followed by DGE analysis revealed significant enrichment of

genes associated with metabolic pathways, while genes related to cell proliferation

exhibited relatively low activity (Fig. 4g-i). The tc_MMP7 cluster was detected in both

pre- and post-treatment ER samples, yet was nearly absent in other samples, supporting

its potential role in early resistance (Fig. 3f). Additionally, DGE analysis and GSEA of

the tc_MMP7 demonstrated significant enrichment of pathways related to EMT,

inflammatory response, IL-6/JAK/STAT3 signaling, KRAS signaling, and TNF-alpha

signaling via NF-kB (Fig. 4j-l). The tc_SCGB3A1 cluster was the most abundant in

LR1_pre (Fig. 3f), possibly indicating a role in the long-term response. DGE analysis

and GSEA showed a decrease in E2F targets and G2-M checkpoint, suggesting a

relatively slow cell cycle (Extended Data Fig. 9a-c). The tc_AZGP1 cluster was

characteristic of SR_pre (Fig. 3f), although no significant enrichment was observed

(Extended Data Fig. 9d-f). This cluster may have been affected by the extremely low

cell number.

To investigate differences in the TME across tumor clusters, we quantified the

spatial distribution of surrounding cell types within the ER_pre samples. For each tumor

cluster, we calculated the proportion of cell types within concentric annular regions

extending up to 1000 μm from each tumor cell (Extended Data Fig. 9g). This analysis

revealed that the tc_S100A6 and tc_AQP1 clusters were diffusely distributed, whereas

the tc_TMSB4X and tc_MMP7 clusters were each localized in close proximity, a

pattern also observed in the spatial mapping of ER_pre (Fig. 3g). Notably, the

tc_MMP7 cluster displayed the highest proportion of immune cells within its

surrounding microenvironment. As shown in Figure 4l, GSEA of the tc_MMP7 cluster

revealed an enrichment of genes associated with several inflammatory responses. Taken

together, these findings suggest that this cluster is characterized by an "immune hot"

state.

Maturity of tertiary lymphoid structures in TME

To examine the spatial relationships of each cluster within the TME, we performed a

neighborhood enrichment analysis using the spatial map coordinates annotated with cell

types. This analysis revealed a close spatial relationship between T cells, B cells, and

follicular dendritic cells (FDCs), particularly in the ER_pre and LR1_pre samples (Fig.

5a). By visualizing these cell populations on the spatial map, they were distributed

across several regions in ER_pre sample, whereas only one region was observed in

LR1_pre sample (Extended Data Fig. 10a,b). In ER_pre sample, the clustering regions

exhibited a structure where B cells and FDCs were intermixed, sparsely surrounded by

T cells (Fig. 5b and Extended Data Fig. 10c). In contrast, in the LR1_pre sample, the

FDCs were positioned around the B cells, with T cells surrounding the outer perimeter

(Fig. 5c and Extended Data Fig. 10d). Moreover, CXCL13 expression was elevated

within these structures, indicating the presence of tertiary lymphoid structures (TLS).

According to previous reports, the poorly organized clustering region in the ER_pre

sample corresponded to an immature TLS, whereas the well-structured regions in the

LR1_pre sample were identified as a mature TLS[32].

Enrichment of MIF- and PDGF- mediated cell-cell interactions in the early

resistant TME

To further elucidate the characteristics of TME, we analyzed intercellular interactions

using COMMOT. This Python library calculates cell-cell interactions based on ligand-

receptor competition and spatial proximity. We extracted ligand-receptor gene pairs

with sufficient expression levels from the CellChat database of known ligand-receptor

interactions (Supplementary Table 2). Pathway comparisons across samples revealed

elevated expression of genes associated with the macrophage migration inhibitory factor

(MIF) pathway in ER and SR (Fig. 6a,b). MIF is a ligand of the CD74/CD44 or

CD74/CXCR4 complex and regulates the inflammatory response[33]. To quantify ligand-

receptor interactions, we calculated signal strength based on the spatial proximity of

ligand- and receptor-expressing cells and their expression levels. Comparing the ER_pre

and LR1_pre spatial signal vectors, the signals on the spatial map were directed to

regions containing TLS and inflammatory cell infiltration (Fig. 6c,d). Signal intensity

and direction was further analyzed by cell type (Fig. 6e-g). In both ER_pre and LR1_pre,

signals showed bidirectional signaling, predominantly among T cells, follicular

1   dendritic cells, and B cells. Tumor cells exhibited limited signals in the LR1_pre sample,

2   whereas tumor cells in the ER_pre sample displayed signals directed toward

3   macrophages and TLS-related cell types. Some tumor cell clusters activated this

4   pathway via an autocrine mechanism. These findings suggest that tumor cells actively

5   stimulated macrophages and TLS through the MIF pathway in the ER_pre sample.

6   As shown in Figure 2d, the proportion of fibroblasts and endothelial cells

increased post-treatment in the ER sample. Therefore, we focused on the platelet-

derived growth factor (PDGF) pathway, which is associated with fibroblast

proliferation[34]. The signal was not localized to specific regions within the ER_pre and

LR1_pre samples (Fig. 6h,i). Bidirectional signaling was observed between fibroblasts

and endothelial cells in both samples (Fig. 6j-l). In the ER_pre sample, tumor cells,

including tc_TMSB4x and tc_MMP7 clusters, exhibited signals directed toward stromal

cells. In contrast, tumor cell clusters in LR1_pre sample showed limited signal directed

at stromal cells. These results suggest that tumor cells interact with stromal cells via the

PDGF pathway, possibly contributing to the formation of an early-resistant TME.

__Discussion__

Although various mechanisms of resistance to EGFR-TKIs in *EGFR*-mutated NSCLC

have been reported, there are limited studies comparing early resistance cases with long-

term responders[3,35,36]. In this study, we showed that the cell proliferation of residual

tumors was enhanced following osimertinib treatment. Genes significantly upregulated

after treatment were associated with poor prognosis[37–39], emphasizing the need to target

the remaining proliferative tumor cells. Tumor cell clusters, including the tc_S100A6

and tc_AQP1, persisted before and after osimertinib treatment, suggesting that

resistance has developed through the activation of downstream or alternative pathways.

In fact, genes involved in resistance mechanisms such as CCND1, HER2, or MET were

significantly high in these clusters, with their amplifications being associated with

EGFR-TKI resistance [4]. Compared to tc_S100A6 and tc_AQP1 clusters, the

tc_TMSB4X cluster, predominantly observed in ER_pre, demonstrated a relatively slow

cell cycle, suggesting a quiescent state associated with EGFR-TKIs resistance[40]. We

showed that the MIF-CD74 pathway was enhanced in the early resistance sample, with

the tc_TMSB4X cluster activating this pathway via an autocrine mechanism. Notably,

this cluster exhibited significantly higher CD74 expression compared to the other

clusters. CD74 has been previously associated with resistance to EGFR-TKIs[41],

suggesting that the tc_TMSB4X cluster acts as a reservoir for resistant cells.

Furthermore, this cluster showed elevated bile acid metabolism, potentially reflecting

suppression of immune cell responses[42]. In contrast, the tc_SCGB3A1 cluster, which

also exhibited a slow cell cycle, was mainly present in LR1_pre and disappeared after

osimertinib treatment, implying characteristics of low proliferation and low malignancy.

The tc_MMP7 cluster had already exhibited an abundant TNF-alpha signaling via NF-

kB even before the initiation of osimertinib treatment. The activation of TNF-alpha

signaling induced by osimertinib has been shown to contribute EGFR-TKI resistance[43–

45], indicating it as an early resistance mechanism. Furthermore, in spatial distribution

analysis, the tc_MMP7 cluster was in an immunologically hot state due to a large

amount of plasmablasts infiltrating the surrounding area. A recent report has shown that

CXCR1$^+$ neutrophils contribute to early resistance to EGFR-TKI through TNF signaling

via NF-kB[13]. However, in our analysis, the number of neutrophils was extremely low

and thus not included in the assessment. Taken together, our findings reveal that tumor

heterogeneity contributes to resistance to EGFR-TKIs.

Increasing evidence suggests that interactions between tumor cells and

surrounding stromal or immune cells involved in mechanisms of EGFR-TKI

resistance[9,10,46]. Cancer associated fibroblasts (CAFs) have been reported as key

regulators of the TME[47]. Our comparison analysis revealed an increase in the proportion

of stromal cells in early resistance samples after osimertinib treatment, while it was no

change in the SR samples, and it decreased in the LR samples. Notably, fibroblasts

exhibited a marked increase, along with enhanced extracellular matrix production and

upregulation of collagens such as COL1A1 and COL3A1, suggesting a potential

correlation with early resistance[31]. In addition, PDGFRA and PDGFRB, which are

receptors for PDGF signaling, are known markers of tumor-associated fibroblasts and

regulate angiogenesis[47]. In the ER_pre sample, this PDGF signaling was active between

tumor cells and stomal cells, suggesting that tumor cells interact with stromal cells via

the PDGF pathway, thereby inducing an early resistant TME.

We demonstrated that inflammatory response was significantly upregulated in

early resistance tissues compared to long-term response tissues before osimertinib

treatment. We further showed that TME in early resistance was characterized by

immature TLSs and infiltration of B cells and plasmablasts, whereas TME in long-term

response exhibited mature TLSs with reduced immune cell infiltration. Previous reports

have shown that immature TLSs were implicated in tumor-promoting effects, while

mature TLSs were tumor-suppressive[32,48,49]. Our findings suggest that the degree of

TLS maturation is associated with early resistance mechanisms. *EGFR*-mutated NSCLC

is generally characterized by low immunogenicity, often described as an "immune cold"

state, which contributes to resistance to immunotherapy[8,9]. In contrast, an "immune hot"

state, defined by TLS formation and high immune cell infiltration, has been associated

with enhanced responsiveness to immunotherapy[50–54]. Notably, patients with a short

response to EGFR-TKIs have shown better responses to immunotherapy[55], suggesting

that their TME exhibits an "immune hot" state. Consistently, we showed that

inflammatory responses were enriched in ER_pre samples. Taken together, these results

indicate that an "immune hot" TME serves as a predictor of early resistance in *EGFR*-

mutated NSCLC (Fig. 7).

Visium HD allows for high-resolution analysis, but it is difficult to distinguish

between noise genes and differentially expressed genes. We addressed this by detecting

and segmenting cell nuclei, reconstructing read count data and cell coordinate data to

improve data reliability. While this approach minimized false positives, some clusters

still contained mixed cell types, which we subsequently excluded. As a result of these

processes, comprehensive gene expression analysis and spatial analysis has become

possible while improving the accuracy of single-cell data and spatial coordinates. For

further improvement in precision, multi-omics approaches are required.

This study has several limitations. First, the size of the bronchoscopic lung

biopsy specimen was small compared to the size of a Visium HD slide, limiting the

amount of data that could be obtained. The number of tumor cells may have been

insufficient to analyze the changes pre- and post-treatment. Second, enhancing single-

cell resolution resulted in reduced count data, potentially affecting the analysis of

differentially expressed genes. Third, we cannot rule out the possibility that our findings

were affected by sample variability. Future studies should collect additional clinical data

to validate these findings.

In conclusion, our study suggests that an inflammatory TME and tumor

heterogeneity contribute to early resistance mechanisms in *EGFR*-mutated NSCLC.

Notably, even before treatment, certain tumor cell clusters already exhibit upregulated

resistance-related pathways.

__Method__

Sample preparation

We collected formalin fixed paraffin embedded (FFPE) block specimens from four

EGFR-mutated NSCLC patients pre- and post-treatment with osimertinib from multiple

institutions. The biopsy methods for the specimens were surgical biopsy, bronchoscopic

biopsy, surgical pleural biopsy, and single hole thoracoscopic pleural biopsy. Patient

clinical data are listed in Supplementary Table 1. Patients were classified into early

resistance (ER, n=1), standard response (SR, n=1), and long-term response (LR, n=2)

based on the duration of progression free survival. These specimens were originally

performed to determine treatment. We obtained written consent from all patients for

specimen collection and research use. This study was carried out in accordance with the

Declaration of Helsinki and was approved by the Institutional Review Board of Kyushu

University Hospital (approval no. 23297-01)

10X Genomics Visium HD data generation

In this study, two capture areas on a Visium HD slide were utilized, with four samples

assigned to each capture area. A total of eight samples consisting of paired pre- and

post-treatment samples from four patients (ER, SR, LR1 and LR2) were analyzed, and

each sample was differentiated from (ER_pre, ER_post, SR_pre, SR_post, LR1_pre,

LR1_post, LR2_pre, LR2_post). Specifically, ER and SR samples were assigned to one

capture area, while LR1 and LR2 samples were assigned to the other capture area.

Initially, RNA was extracted from FFPE tissue sections prepared for RNA quality

assessment, and the DV200 value was measured using the 4200 Tape Station. The

DV200 value for all samples was greater than or equal to 30%. The FFPE tissue

sections were then subjected to deparaffinization and H&E staining following the

Visium HD FFPE Tissue Preparation protocol provided by 10x Genomics (Visium HD

FFPE Tissue Preparation Handbook, CG000684 Rev A). After staining, tissue sections

were imaged using the EVIDENT VS200 microscope in accordance with the 10X

Genomics protocol (Visium HD Spatial Applications Image Guidelines CG000688 Rev

A). Gene expression libraries were prepared from the H&E-stained sections following

the 10X Genomics protocol (Visium HD Spatial Gene Expression Reagent Kits User

Guide CG000685 Rev A). The quality of the prepared libraries was evaluated using the

4200 Tape Station, and all libraries were confirmed to have sufficient concentrations.

The Visium libraries were sequenced using the NovaSeq X Plus platform with a paired-

end configuration (read1: 150 bp, read2: 150 bp). Sequencing data (FASTQ files) were

processed and analyzed using Space Ranger (version 3.0.0) software to assess data

quality and generate spatial gene expression profiles.

Analysis Environment

The analysis was conducted using Python (version 3.10) on either Ubuntu (22.04.5

LTS) with JupyterLab (4.2.5) or Windows (Windows 11 Home edition) with PyCharm

(Community Edition 2024.1.1). The primary Python libraries and their versions

included: AnnData (0.10.8), AnnData2ri (1.3.2), Commot (0.0.3), GeoPandas (1.0.1),

GSEApy (1.1.3), HarmonyPy (0.0.10), inferCNVpy (0.5.0), Matplotlib (3.8.4), NumPy

(1.26.4), Pandas (2.2.2), Rpy2 (3.5.16), Scanpy (1.10.3), SciPy (1.11.4), Seaborn

(0.13.2), Squidpy (1.6.1) and StarDist (0.9.1). In some analyses, R Script was executed

in Python. The primary R packages and their versions used were: tradeSeq (1.8.0),

edgeR (3.36.0), and clusterExperiment (2.14.0). While the libraries were generally

applied as described in their documentation, modifications were made to Commot to

improve visual clarity in plots.

Nuclei detection and segmentation

Although Visium HD allows for the use of 8 µm square spots as single cells, these spots

do not accurately reflect the true cellular coordinates. To preserve cellular coordinates

as much as possible, nuclei detection and segmentation were performed using StarDist[19].

Specifically, Python code was developed based on Analysis Guide provided by 10x

Genomics (https://www.10xgenomics.com/jp/analysis-guides/segmentation-visium-hd).

Running this code generated coordinate data for individual nuclei represented as

polygons. Each 2 µm spot count data within the polygon area was relabeled as a single

cell. Additionally, the centroid coordinates of each polygon were added to the count

data, along with metadata content, creating a data structure compatible with libraries

commonly used in spatial transcriptomics analysis.

Sample labeling on spatial coordinates

In typical spatial transcriptomics analyses, all samples within a single capture area are

assumed to be the same. However, in this study, different four samples were present

within the same capture area. Since this situation is not typically supported by existing

tools, it was necessary to manually label samples based on spatial coordinates. Using

the spatial coordinate data within the data structure, samples were divided into eight

groups by referencing images and assigning coordinates manually. Since there was an

overlapping area between LR1_pre and LR1_post, a blank area was created to avoid

sample contamination.

Major clustering

The newly generated data from nuclei segmentation were analyzed using a pipeline

similar to single-cell RNA-seq analysis with Scanpy package. Cells with fewer than 20

counts per cell or containing more than 10% mitochondrial gene, genes with fewer than

20 total counts were excluded from the analysis. Cells with a cross-sectional area of 900

μm2 or more on the image were excluded in consideration of the possibility that they

were composed of multiple cells. After filtering, the data were calculated for highly

variable genes (HVGs) and normalized using Pearson residuals, which adjusts for

library size and overdispersion. Principal component analysis (PCA) using the top 4000

HVGs calculated the principal components up to the top 35. To account for batch

effects across samples, Harmony integration was applied using the HarmonyPy library

through the Scanpy with the sample labels. Subsequently, the nearest neighbors distance

matrix and the neighborhood graph of the observations were calculated with a local

neighborhood size of 100 and 35 principal components. A UMAP embedding was

generated with a minimum distance parameter set to 0.5. Leiden clustering was

performed at a resolution of 0.4 and identified 18 clusters. However, due to the high

noise level in cells, it was possible that the differences between cells and noise could

not be distinguished. Instead, we analyzed HVGs in the expression of each cluster and

compared the expression levels of known cell markers to classify the clusters into three

major cell types: epithelial, stromal, and immune cell types (Supplementary Table 3).

Sub-cluster analysis

Sub-clustering was performed to further explore the epithelial, stromal, and immune cell

populations identified in the initial clustering. For each cell type cluster, the analysis

was start again from the raw count data. The data were filtered to the same threshold as

the major clustering. HVGs were calculated within some sub-cluster, PCA was

calculated, followed by the construction of neighborhood graphs, UMAP embedding,

and clustering. The distinction between tumor cells and normal cells was evaluated

using inferCNVpy, which is based on inferCNV, in addition to the pathological images.

Normal airway epithelial cells obtained from the clustering results were set as non-

malignant cells and analyzed. Further, for tumor cells, additional clustering was

performed. For immune cells, three cell types—lymphocytes, plasmablasts, and

macrophages—were identified, and each type was subsequently analyzed with

individual clustering steps. Detailed parameters for each clustering step and cell

annotation markers are provided in the supplementary materials (Supplementary Table

3,4).

Differential Gene Expression Analysis

We performed differential gene expression (DGE) analysis using created pseudo-bulk

samples combined with the edgeR package[56,57]. Single-cell RNA-seq data were

aggregated at the sample level for each cell type of interest. Samples with fewer than 30

cells of the target cell type were excluded. For comparisons between ER and SLR

groups, as well as pre- and post-treatment comparisons of the tc_AZGP1 cluster, two

random replicates were generated for each group because the edgeR library requires

biological replicates for statistical modeling. In the others, pseudo-bulk aggregation was

performed without creating additional replicates.

After pseudo-bulk aggregation, DGE analysis was performed using the edgeR

library[57]. Low-expressed genes were first filtered out, retaining only those with

sufficient counts across experimental groups to improve statistical power. Library size

normalization was carried out using the trimmed mean of M-values method, which

adjusts for sequencing depth and compositional biases across samples. A design matrix

was constructed to account for sample conditions and randomly generated replicates,

enabling the estimation of both biological variation and group-specific effects. The

quasi-likelihood variance estimation method was employed to generate a gene list for

DGE. Adjusted p-values were calculated using the Benjamini-Hochberg method to

control the false discovery rate. Results for each cell type were integrated back into the

annotated data object to facilitate further interpretation. The aggregated gene expression

values were subsequently used for downstream analyses, including the creation of

volcano plots and GSEA.

The Kaplan-Meier Plotter database

We used data from lung cancer patients in the Kaplan-Meier Plotter database

(http://kmplot.com/analysis/) to evaluate differentially expressed genes in tumor cells

compared before and after treatment, respectively[58]. Patients on the database were

divided into high and low expression groups by median expression values, and survival

curves were plotted with no follow-up restrictions.

Integration of cell type and spatial coordinate data

The dataset used in this analysis was generated by integrating single-cell analysis for

cell type identification with spatial coordinate data derived from StarDist. The StarDist-

based coordinate data, which represents new spot coordinates, was transformed based

on the ratio of the actual distance to the pixel coordinates, as described in the metadata.

This integration resulted in a unified data structure compatible with analysis tools such

as Scanpy and Squidpy.

Neighborhood enrichment analysis

Neighborhood enrichment analysis was performed on the spatial transcriptomics data to

assess the enrichment of different cell clusters within their spatial neighborhoods. For

each of the 8 sample datasets, spatial neighbors function within Squidpy was used to

calculate the spatial neighbors of each spot. This was followed by the application of the

neighborhood enrichment function to calculate the z-scores, using predefined cell

cluster labels. To standardize the enrichment scores across all samples, the minimum

and maximum values were first calculated across all datasets. Each z-score matrix was

then normalized by subtracting the global mean and dividing by the standard deviation,

ensuring that enrichment scores were comparable across samples. This normalization

allowed for the unified analysis of neighborhood enrichment across different

experimental conditions.

Spatial distribution and neighborhood analysis of tumor cells

1 The spatial distribution and neighborhood relationships of tumor cells were analyzed by

2 integrating single-cell cluster information with spatial coordinate data. For each of the 8

3 sample datasets, the spatial positions of tumor cells were identified. To quantify the

4 surrounding cell composition, we calculated the pairwise distances between each tumor

5 cell and all other cell types within the tissue. To analyze the spatial distribution, we

6 defined concentric annular regions extending up to 1000 µm from each tumor cell, with

7 each region increasing in 20 µm increments. These regions served as distance bins,

8 allowing us to categorize surrounding cells based on their proximity to tumor cells. We

9 then calculated the proportion of each cell type within these bins to assess their spatial

10 distribution relative to tumor clusters. This method enabled a quantitative evaluation of

11 neighborhood enrichment patterns, providing insights into the TME and its cellular

12 composition.

14 **Cell-cell communication analysis**

15 Cell-cell interactions were analyzed using COMMOT, a tool that enables

16 comprehensive modeling of both direct and indirect interactions between different cell

17 types while incorporating the spatial organization of cells within tissues[59]. For this

18 analysis, smooth muscle cells, normal airway epithelial cells, and blood cells were

19 excluded to focus on the TME. Additionally, genes expressed in more than 0.5% of all

20 cells were first filtered. This gene list was further refined by extracting known ligand-

21 receptor pairs from the CellChat database. Only ligand-receptor pairs co-occurring

22 within a distance of 500 µm were considered for subsequent analysis. For the identified

23 ligand-receptor pairs and their associated pathways, signal strength and vectors were

calculated based on the distance between sending and receiving cells as well as gene

expression intensity.

Gene expression analysis and statistics

Gene expression scores were calculated using marker gene-based scoring method

implemented in Scanpy. The "MIF Score" was derived from the MIF pathway-related

genes: *MIF*, *CD74*, *CD44* and *CXCR4*. Background correction was performed using

control gene sets of 5000 randomly selected genes to ensure robust scoring. Pairwise

comparisons of each score between the Comparisons of MIF scores between samples

and gene expression in tumor cells were performed using a pairwise comparison using a

unilateral Mann-Whitney U test. Visualization was conducted with Seaborn for violin

and strip plots   annotated with the value of the p-value. The significance level of

GSEA after DGE analysis was set as p value less than 0.05 and false discovery rate as

0.25.

__Data availability__

The Visium HD dataset from this study is available on the GEO repository under

accession numbers GSE288758.

_

_Reference__

1. Ramalingam SS, Vansteenkiste J, Planchard D, et al. Overall Survival with
Osimertinib in Untreated, EGFR-Mutated Advanced NSCLC. *New England Journal*
of Medicine. 2020;382(1):41-50. doi:10.1056/NEJMoa1913662

2. Soria JC, Ohe Y, Vansteenkiste J, et al. Osimertinib in Untreated EGFR-Mutated
Advanced Non–Small-Cell Lung Cancer. *New England Journal of Medicine*.
2018;378(2):113-125. doi:10.1056/NEJMoa1713137

3. Oxnard GR, Hu Y, Mileham KF, et al. Assessment of Resistance Mechanisms and
Clinical Implications in Patients With EGFR T790M–Positive Lung Cancer and
Acquired Resistance to Osimertinib. *JAMA Oncol*. 2018;4(11).
doi:10.1001/jamaoncol.2018.2969

4. Leonetti A, Sharma S, Minari R, Perego P, Giovannetti E, Tiseo M. Resistance
mechanisms to osimertinib in EGFR-mutated non-small cell lung cancer. *Br J*
Cancer. 2019;121(9):725-737. doi:10.1038/s41416-019-0573-8

5. Maynard A, McCoach CE, Rotow JK, et al. Therapy-Induced Evolution of Human
Lung Cancer Revealed by Single-Cell RNA Sequencing. *Cell*. 2020;182(5):1232-
1251.e22. doi:10.1016/j.cell.2020.07.017

6. Oh S, Koh J, Kim TM, et al. Transcriptomic Heterogeneity of EGFR-Mutant Non–
Small Cell Lung Cancer Evolution Toward Small-Cell Lung Cancer. *Clinical Cancer*
Research. 2024;30(20):4729-4742. doi:10.1158/1078-0432.CCR-24-0160

7. Madeddu C, Donisi C, Liscia N, Lai E, Scartozzi M, Macciò A. EGFR-Mutated Non-
Small Cell Lung Cancer and Resistance to Immunotherapy: Role of the Tumor
Microenvironment. *Int J Mol Sci*. 2022;23(12):6489. doi:10.3390/ijms23126489

8. Chen S, Tang J, Liu F, et al. Changes of tumor microenvironment in non-small cell
lung cancer after TKI treatments. *Front Immunol*. 2023;14.
doi:10.3389/fimmu.2023.1094764

9. Yang L, He YT, Dong S, et al. Single-cell transcriptome analysis revealed a
suppressive tumor immune microenvironment in EGFR mutant lung adenocarcinoma.
J Immunother Cancer. 2022;10(2):e003534. doi:10.1136/jitc-2021-003534

10.   Matsumoto Y, Sawa K, Fukui M, et al. Impact of tumor microenvironment on
the efficacy of epidermal growth factor receptor-tyrosine kinase inhibitors in patients
with -mutant non-small cell lung cancer. *Cancer Science*. 2019;110(10):3244-3254.
doi:10.1111/cas.14156

11.   Yang CY, Liao WY, Ho CC, et al. Association between programmed death-
ligand 1 expression, immune microenvironments, and clinical outcomes in epidermal
growth factor receptor mutant lung adenocarcinoma patients treated with tyrosine

kinase inhibitors. *European Journal of Cancer*. 2020;124:110-122. doi:10.1016/j.ejca.2019.10.019

12. Hsu KH, Tseng JS, Yang TY, et al. PD-L1 strong expressions affect the clinical outcomes of osimertinib in treatment naïve advanced EGFR-mutant non-small cell lung cancer patients. *Sci Rep*. 2022;12(1):9753. doi:10.1038/s41598-022-13102-7

13. Wang H, Xiong A, Chen X, et al. CXCR1+ neutrophil infiltration orchestrates response to third-generation EGFR-TKI in EGFR mutant non-small-cell lung cancer. *Sig Transduct Target Ther*. 2024;9(1):1-15. doi:10.1038/s41392-024-02045-2

14. Zhang H, AbdulJabbar K, Moore DA, et al. Spatial Positioning of Immune Hotspots Reflects the Interplay between B and T Cells in Lung Squamous Cell Carcinoma. *Cancer Res*. 2023;83(9):1410-1425. doi:10.1158/0008-5472.CAN-22-2589

15. De Zuani M, Xue H, Park JS, et al. Single-cell and spatial transcriptomics analysis of non-small cell lung cancer. *Nat Commun*. 2024;15(1):4388. doi:10.1038/s41467-024-48700-8

16. Peng H, Wu X, Liu S, et al. Multiplex immunofluorescence and single‐cell transcriptomic profiling reveal the spatial cell interaction networks in the non‐small cell lung cancer microenvironment. *Clin Transl Med*. 2023;13(1):e1155. doi:10.1002/ctm2.1155

17. Wang S, Rong R, Yang DM, et al. Features of tumor-microenvironment images predict targeted therapy survival benefit in patients with EGFR-mutant lung cancer. *J Clin Invest*. 133(2):e160330. doi:10.1172/JCI160330

18. Dagogo-Jack I, Valiev I, Kotlov N, et al. B-Cell Infiltrate in the Tumor Microenvironment Is Associated With Improved Survival in Resected Lung Adenocarcinoma. *JTO Clinical and Research Reports*. 2023;4(7). doi:10.1016/j.jtocrr.2023.100527

19. Schmidt U, Weigert M, Broaddus C, Myers G. Cell Detection with Star-Convex Polygons. In: Frangi AF, Schnabel JA, Davatzikos C, Alberola-López C, Fichtinger G, eds. *Medical Image Computing and Computer Assisted Intervention – MICCAI 2018*. Springer International Publishing; 2018:265-273. doi:10.1007/978-3-030-00934-2_30

20. Mantovani A, Sica A. Macrophages, innate immunity and cancer: balance, tolerance, and diversity. *Current Opinion in Immunology*. 2010;22(2):231-237. doi:10.1016/j.coi.2010.01.009

21. Matsubara E, Yano H, Pan C, et al. The Significance of SPP1 in Lung Cancers and Its Impact as a Marker for Protumor Tumor-Associated Macrophages. *Cancers (Basel)*. 2023;15(8):2250. doi:10.3390/cancers15082250

22. Song Y, Fu Y, Wang J, et al. Complement C1q induces the M2-polarization of tumor-associated macrophages in lung adenocarcinoma. *Genes Dis*. 2023;11(4):101093. doi:10.1016/j.gendis.2023.101093

23. Wei X, Dong J. Aquaporin 1 promotes the proliferation and migration of lung cancer cell in vitro. *Oncology Reports*. 2015;34(3):1440-1448. doi:10.3892/or.2015.4107

24. Li P, Lv X, Zhang Z, Xie S. S100A6/miR193a regulates the proliferation, invasion, migration and angiogenesis of lung cancer cells through the P53 acetylation. *Am J Transl Res*. 2019;11(8):4634-4649.

25. Yang Z, Luo J, Zhang M, et al. TMSB4X: A novel prognostic marker for non-small cell lung cancer. *Heliyon*. 2023;9(11):e21505. doi:10.1016/j.heliyon.2023.e21505

26. Yamada K, Kadota K, Fujimoto S, et al. MMP-7 expression is associated with a higher rate of tumor spread through air spaces in resected lung adenocarcinomas. *Lung Cancer*. 2023;175:125-130. doi:10.1016/j.lungcan.2022.12.001

27. Mazumdar J, Hickey MM, Pant DK, et al. HIF-2α deletion promotes Kras-driven lung tumor development. *Proc Natl Acad Sci U S A*. 2010;107(32):14182-14187. doi:10.1073/pnas.1001296107

28. Albertus DL, Seder CW, Chen G, et al. AZGP1 Autoantibody Predicts Survival and Histone Deacetylase Inhibitors Increase Expression in Lung Adenocarcinoma. *Journal of Thoracic Oncology*. 2008;3(11):1236-1244. doi:10.1097/JTO.0b013e318189f5ec

29. Li L, Li Z, Lu C, et al. Ibrutinib reverses IL-6-induced osimertinib resistance through inhibition of Laminin α5/FAK signaling. *Commun Biol*. 2022;5(1):1-14. doi:10.1038/s42003-022-03111-7

30. Fang W, Huang Y, Gu W, et al. PI3K-AKT-mTOR pathway alterations in advanced NSCLC patients after progression on EGFR-TKI and clinical response to EGFR-TKI plus everolimus combination therapy. *Transl Lung Cancer Res*. 2020;9(4):1258-1267. doi:10.21037/tlcr-20-141

31. Yamazaki S, Higuchi Y, Ishibashi M, et al. Collagen type I induces EGFR☐TKI resistance in EGFR☐mutated cancer cells by mTOR activation through Akt☐independent pathway. *Cancer Sci*. 2018;109(6):2063-2073. doi:10.1111/cas.13624

32. Bruno TC. B cells to the forefront of immunotherapy. *Nature*. 2020;577(7791):474-476. doi:10.1038/d41586-019-03943-0

33. Calandra T, Roger T. Macrophage migration inhibitory factor: a regulator of innate immunity. *Nat Rev Immunol*. 2003;3(10):791-800. doi:10.1038/nri1200

34. Yao L, Rathnakar BH, Kwon HR, et al. Temporal control of PDGFRα regulates the fibroblast-to-myofibroblast transition in wound healing. *Cell Rep*. 2022;40(7):111192. doi:10.1016/j.celrep.2022.111192

35. Thress KS, Jacobs V, Angell HK, et al. Modulation of Biomarker Expression by Osimertinib: Results of the Paired Tumor Biopsy Cohorts of the AURA Phase I Trial. *Journal of Thoracic Oncology*. 2017;12(10):1588-1594. doi:10.1016/j.jtho.2017.07.011

36. Isomoto K, Haratani K, Hayashi H, et al. Impact of EGFR-TKI Treatment on the Tumor Immune Microenvironment in *EGFR* Mutation–Positive Non–Small Cell Lung Cancer. *Clinical Cancer Research*. 2020;26(8):2037-2046. doi:10.1158/1078-0432.CCR-19-2027

37. Jiang C, Zhang Y, Li Y, et al. High CEP55 expression is associated with poor prognosis in non-small-cell lung cancer. *Onco Targets Ther*. 2018;11:4979-4990. doi:10.2147/OTT.S165750

38. Luo A, Yin Y, Li X, Xu H, Mei Q, Feng D. The clinical significance of FSCN1 in non-small cell lung cancer. *Biomedicine & Pharmacotherapy*. 2015;73:75-79. doi:10.1016/j.biopha.2015.05.014

39. Li Y, Leng Y, Dong Y, et al. Cyclin B1 expression as an independent prognostic factor for lung adenocarcinoma and its potential pathways. *Oncol Lett*. 2022;24(6):441. doi:10.3892/ol.2022.13561

40. Lindell E, Zhong L, Zhang X. Quiescent Cancer Cells—A Potential Therapeutic Target to Overcome Tumor Resistance and Relapse. *Int J Mol Sci*. 2023;24(4):3762. doi:10.3390/ijms24043762

41. Kashima Y, Shibahara D, Suzuki A, et al. Single-cell analyses reveal diverse mechanisms of resistance to EGFR tyrosine kinase inhibitors in lung cancer. *Cancer Res*. 2021;81(18):4835-4848. doi:10.1158/0008-5472.CAN-20-2811

42. Varanasi SK, Chen D, Liu Y, et al. Bile acid synthesis impedes tumor-specific T cell responses during liver cancer. *Science*. 2025;387(6730):192-201. doi:10.1126/science.adl4100

43. Gong K, Guo G, Gerber DE, et al. TNF-driven adaptive response mediates resistance to EGFR inhibition in lung cancer. *J Clin Invest*. 128(6):2500-2518. doi:10.1172/JCI96148

44. Nam YW, Shin JH, Kim S, et al. EGFR inhibits TNF-α-mediated pathway by phosphorylating TNFR1 at tyrosine 360 and 401. *Cell Death Differ*. 2024;31(10):1318-1332. doi:10.1038/s41418-024-01316-3

45. Ibusuki R, Iwama E, Shimauchi A, et al. TP53 gain-of-function mutations promote osimertinib resistance via TNF-α–NF-κB signaling in EGFR-mutated lung cancer. *NPJ Precis Oncol*. 2024;8:60. doi:10.1038/s41698-024-00557-2

46. Alsaed B, Lin L, Son J, et al. Intratumor heterogeneity of EGFR expression mediates targeted therapy resistance and formation of drug tolerant microenvironment. *Nat Commun*. 2025;16(1):28. doi:10.1038/s41467-024-55378-5

47. Luo H, Xia X, Huang LB, et al. Pan-cancer single-cell analysis reveals the heterogeneity and plasticity of cancer-associated fibroblasts in the tumor microenvironment. *Nat Commun*. 2022;13(1):6619. doi:10.1038/s41467-022-34395-2

48. Helmink BA, Reddy SM, Gao J, et al. B cells and tertiary lymphoid structures promote immunotherapy response. *Nature*. 2020;577(7791):549-555. doi:10.1038/s41586-019-1922-8

49. Petitprez F, de Reyniès A, Keung EZ, et al. B cells are associated with survival and immunotherapy response in sarcoma. *Nature*. 2020;577(7791):556-560. doi:10.1038/s41586-019-1906-8

50. Ukita M, Hamanishi J, Yoshitomi H, et al. CXCL13-producing CD4+ T cells accumulate in the early phase of tertiary lymphoid structures in ovarian cancer. *JCI Insight*. 7(12):e157215. doi:10.1172/jci.insight.157215

51. Germain C, Gnjatic S, Tamzalit F, et al. Presence of B Cells in Tertiary Lymphoid Structures Is Associated with a Protective Immunity in Patients with Lung Cancer. *Am J Respir Crit Care Med*. 2014;189(7):832-844. doi:10.1164/rccm.201309-1611OC

52. Cabrita R, Lauss M, Sanna A, et al. Tertiary lymphoid structures improve immunotherapy and survival in melanoma. *Nature*. 2020;577(7791):561-565. doi:10.1038/s41586-019-1914-8

53. Laumont CM, Banville AC, Gilardi M, Hollern DP, Nelson BH. Tumour-infiltrating B cells: immunological mechanisms, clinical impact and therapeutic opportunities. *Nat Rev Cancer*. 2022;22(7):414-430. doi:10.1038/s41568-022-00466-1

54. Leong TL, Bryant VL. B cells in lung cancer—not just a bystander cell: a literature review. *Transl Lung Cancer Res*. 2021;10(6):2830-2841. doi:10.21037/tlcr-20-788

55. Yoshida H, Kim YH, Ozasa H, et al. Nivolumab in non-small-cell lung cancer with EGFR mutation. *Annals of Oncology*. 2018;29(3):777-778. doi:10.1093/annonc/mdx745

56. Murphy AE, Skene NG. A balanced measure shows superior performance of pseudobulk methods in single-cell RNA-sequencing analysis. *Nat Commun*. 2022;13(1):7851. doi:10.1038/s41467-022-35519-4

57.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics*.
2010;26(1):139-140. doi:10.1093/bioinformatics/btp616

58.   Győrffy B. Transcriptome-level discovery of survival-associated biomarkers and
therapy targets in non-small-cell lung cancer. *British Journal of Pharmacology*.
2024;181(3):362-374. doi:10.1111/bph.16257

59.   Cang Z, Zhao Y, Almet AA, et al. Screening cell–cell communication in spatial
transcriptomics via collective optimal transport. *Nat Methods*. 2023;20(2):218-228.
doi:10.1038/s41592-022-01728-4

 __**Figure legends**__

 Fig. 1 | Analysis overview and clustering, visualization of cell type and spatial

 location. a, Biopsy interval pre- and post-osimertinib treatment and patient

 classification. **b**, Detecting and segmenting cell nuclei using StarDist on high-resolution

 tissue images loaded with Visium HD protocol. **c**, Visualization of detection and

 segmentation by StarDist and 2 µm spot extraction of identified cell nuclei positions. **d**,

 UMAP of annotated cell types by single cell RNA-seq based on cell nuclei identified by

 StarDist. **e**, UMAP with epithelial cells re-clustered and labeled for cell type. **f**, UMAP

 with Immune cells re-clustered and labeled for cell type. **g**, UMAP with stromal cells

 re-clustered and labeled for cell type. **h,i**, H&E-stained pathological image of a certain

 area of ER_pre (h), and image of the same area with cell type identification reflected in

 polygon coordinate data obtained by StarDist (i). **j,k**, H&E-stained pathological images

 of magnified regions of LR2_pre (j), and images of cell type-identified same regions

 reflected in polygonal coordinate data obtained by StarDist (k). Normal epithelial cells

 and tumor cells were distinguished, and surrounding mesenchymal cells and immune

 cells were identified with high accuracy.

 Fig. 2 | Comparative analysis before and after treatment. a, Volcano plot showing

 the DGE analysis results by extracting tumor cell clusters from all samples and dividing

 them into two groups before and after treatment (All_pre vs All_post). **b**, Dot plot of

 GSEA with Hallmarker gene sets based on DGE analysis of All_pre vs All_post. **c**,

 GSEA plot with high absolute value of normalized enrichment score between All_pre

 and All_post. **d**, Stacked bar chart showing the proportion of cell types within each

 sample. **e**, Volcano plot showing DGE analysis results comparing fibroblasts extracted

from ER_pre and ER_post. **f**, GSEA plot of the only significant gene set in Hallmark

gene sets. **g**, Violin plot and strip plot showing raw counts of *COL1A1* gene for each

sample. One-sided Mann-Whitney U tests were performed to compare the other samples

with ER_post, and the test results were described in the graph. ***p < 0.0005. **h,**

Expression levels of *CD3D*, *CD4*, *CD8A*, and *FOXP3* related to T cell subtypes

displayed on UMAP of T cell parts extracted from lymphocyte clusters. **i,** Expression

levels of genes related to tumor-associated macrophage markers (SPP1, C1QC) and M2

macrophage markers (CD163, MSR1) on UMAP of Macrophage..

Fig. 3 | Analysis exploring mechanisms of tumor heterogeneity and early resistance.

a, Volcano plot showing the DGE analysis results by extracting tumor cell clusters from

pre-treatment samples and dividing them into two groups: ER_pre and SLR_pre. **b**, Dot

plot of GSEA with Hallmarker gene sets based on DGE analysis of ER_pre vs SLR_pre.

c, GSEA plot with high absolute value of normalized enrichment score between ER_pre

and SLR_pre. **d**, Matrix plot showing clustering of tumor cells based on differentially

expressed genes. **e**, UMAP with Tumor cells re-clustered and labeled for differentially

expressed genes. **f**, Violin plot showing the distribution of tumor cell clusters per

sample. **g**, Image of tumor cell cluster positions plotted on HE-stained tissue image of

Slide 1 and Slide 2.

Fig. 4 | Individual analysis of tumor cell clusters. a, Volcano plot showing DGE

analysis results between the tc_S100A6 cluster and the rest. **b**, Dot plot of GSEA with

Hallmarker gene sets based on DGE analysis of the tc_S100A6 cluster vs the rest. **c**,

GSEA plot with high absolute value of normalized enrichment score between the

tc_S100A6 cluster and the rest. **d**, Volcano plot showing DGE analysis results between

the tc_AQP1 cluster and the rest. **e**, Dot plot of GSEA with Hallmarker gene sets based

on DGE analysis of the tc_AQP1 cluster vs the rest. **f**, GSEA plot with high absolute

value of normalized enrichment score between the tc_AQP1 cluster and the rest. **g**,

Volcano plot showing DGE analysis results between the tc_TMSB4X cluster and the

rest. **h**, Dot plot of GSEA with Hallmarker gene sets based on DGE analysis of the

tc_TMSB4X cluster vs the rest. **i**, GSEA plot with high absolute value of normalized

enrichment score between the tc_TMSB4X cluster and the rest. **j**, Volcano plot

showing DGE analysis results between the tc_MMP7 cluster and the rest. **k**, Dot plot of

GSEA with Hallmarker gene sets based on DGE analysis of the tc_MMP7 cluster vs the

rest. **l**, GSEA plot with high absolute value of normalized enrichment score between the

tc_MMP7 cluster and the rest.

Fig. 5 | Spatial neighborhood enrichment analysis and TLS identification. a,

Heatmap representing spatial neighborhood enrichment analysis results. 5000

surrounding cells were counted from each cell, and the proportion of that cell type was

expressed as z-scores normalized to all samples. **b**, Enlarged images of a TLS in the

ER_pre sample. The area surrounded by the white dotted line showed images of HE-

stained tissue, cell types, and the expression level of CXCL13. **c**, Enlarged images of a

TLS in the LR1_pre sample. The area surrounded by the white dotted line showed

images of HE-stained tissue, cell types, and the expression level of CXCL13.

Fig. 6 | Cell-cell interaction analysis using COMMOT. a, Violin plot and strip plot

showing MIF scores for each sample. One-sided Mann-Whitney U tests were performed

to compare the other samples with the ER_pre sample, and the test results were

described in the graph. ns p ≥ 0.05; ***p < 0.0005. **b**, Images of the expression levels of

MIF, *CD74*, *CD44*, and *CXCR4*, which constitute the MIF pathway, and were expressed

in 0.5% or more of all cells in each sample. **c**, An image of spatial directions of cell-cell

communication of MIF pathway in the ER_pre sample. **d**, An image of spatial

directions of cell-cell communication of MIF pathway in the LR1_pre sample. **e**,

Visualization of significant MIF pathway signal networks summarized by cell type in

ER_pre and LR1_pre samples. **f**, Dot plot of significant MIF pathway signal networks

summarized by cell type in the ER_pre sample. **g**, Dot plot of significant MIF pathway

signal networks summarized by cell type in the LR1_pre sample. **h**, An image of spatial

directions of cell-cell communication of PDGF pathway in the ER_pre sample. **i**, An

image of spatial directions of cell-cell communication of PDGF pathway in the

LR1_pre sample. **j**, Visualization of significant PDGF pathway signal networks

summarized by cell type in ER_pre and LR1_pre samples. **k**, Dot plot of significant

PDGF pathway signal networks summarized by cell type in the ER_pre sample. **l**, Dot

plot of significant PDGF pathway signal networks summarized by cell type in the

LR1_pre sample.

Fig. 7 | Proposed model of early resistance to EGFR-TKIs in the tumor

microenvironment. Schematic diagram summarizing findings from the analysis of this

study

Neighborhood enrichment

Early Resistance Mechanisms to EGFR-TKIs in the Tumor Microenvironment

MIF/CD74 pathway

PDGF pathway

Immature TLS

Mature TLS

Fibroblast activity

TNFα/NF-κB signaling

TNFα

NF-κB

Tumor cell   B cell   FDC   Macrophage   T cell   Fibroblast   plasmablast