Duplications in the Chlorophyll *a* oxygenase (CAO) gene across the Viridiplantae

Mackenzie C. Poirier[1], Roberta Wright[1], Marina Cvetkovska[1]*

[1]Department of Biology, University of Ottawa, Ottawa, ON, Canada

*Corresponding author:

Marina Cvetkovska
Department of Biology, University of Ottawa, 30 Marie-Curie Pr., Ottawa, ON, Canada, K1N 6N5
Email: mcvetkov@uottawa.ca
ORCID: 0000-0002-7080-8203

Statements and Declarations

Competing Interests: The authors report that there are no competing interests to declare

Acknowledgements: This project was supported by Natural Sciences and Engineering Research Council of Canada Discovery Grants (NSERC DG) awarded to M.C. The authors are grateful for the support from the Canada Foundation for Innovation (CFI) and University of Ottawa start-up funding. M.P. was supported by Ontario Graduate Scholarship (OGS), NSERC Graduate Scholarship, and Polar Knowledge Canada Antarctic Doctoral Scholarship.

Author's contributions: M. Poirier and M. Cvetkovska conceptualized the work and designed the experiments. Material preparation, data collection and analysis were performed by M. Poirier, R. Wright and M. Cvetkovska. The first draft of the manuscript was written by M. Poirier and all authors commented on all versions of the manuscript. All authors read and approved the final manuscript.

Abstract (150-250 words)

Viridiplantae, a diverse group of green plants and alga that have evolved from a common ancestor,
are unified in their ability to produce and use two types of chlorophyll (chlorophyll *a* and chlorophyll
b) to capture light energy. In addition to its role in light harvesting, appropriate levels of chlorophyll
b are required for accumulation, assembly, and stability of light harvesting complexes within the
photosynthetic apparatus. Chlorophyll *b* is synthesized from chlorophyll *a* by the enzyme
chlorophyllide *a* oxygenase (CAO), a Rieske-type mononuclear non-heme iron oxygenase. A
regulatory degron sequence, described in detail only in land plants, has been demonstrated to
regulate the stability of CAO proteins based on the availability of chlorophyll *b*. Recent
identification of CAO gene duplication in bryophyte and green algal species, combined with
expanded availability of sequenced genomes within the Viridiplantae, prompted further
investigation into the role of gene duplication in the evolution of chlorophyll *b* biosynthesis.
Examination of genomes from 246 plant and algae species revealed independently occurring CAO
duplications throughout the Viridiplantae, with a higher prevalence of duplication in land plants
compared to their algal relatives. Additionally, we demonstrate that the degron sequence is absent
in in chlorophytes, but first appears as a conserved sequence in charophytes, and is very highly
conserved among the embryophytes. The evolutionary history of CAO throughout the Viridiplantae
lineage is discussed based on these key observations, adding to our understanding of chlorophyll *b*
biosynthesis and the role of CAO in photosynthetic species.

Keywords (4-6 keywords)

Introduction

Photosynthetic organisms form the base of all food chains and provide the organic carbon compounds that support life in our biosphere. The capture of light energy by chlorophyll is the first step in the process of photosynthesis that fixes of atmospheric CO_2 into stable organic compounds. Overwhelming evidence indicates that oxygenic eukaryotic photosynthesis that occurs in the chloroplasts of modern-day plants and algae traces its origins from an endosymbiotic event with cyanobacteria-like prokaryotes (Blankenship 2010; Cardona 2019; Sánchez-Baracaldo and Cardona 2020). Viridiplantae (or green plants), are primary endosymbionts that encompass two major clades: the chlorophytes and the streptophytes. Chlorophytes are a monophyletic group of marine, freshwater, and terrestrial green algae. With ~8,000 described species (Guiry 2024), this group encompasses a large diversity of adaptations, morphologies, and life histories (Leliaert et al. 2012). Streptophyta includes the charophytes, ~5,500 species of largely freshwater green algae (Guiry 2024) and the embryophytes, or land plants. Land plants are thought to have diverged from a single clade within the streptophytes (de Vries and Archibald 2018) and number ~430,000 species (Ruggiero et al. 2015). One characteristic that unifies members of the Viridiplantae is their ability to synthesize and use two types of chlorophylls: chlorophyll a (Chl *a*) and chlorophyll b (Chl *b*).

The photosynthetic apparatus in the Viridiplatae is derived from the ancestral cyanobacterial endosymbiont, which used chlorophyll as the main pigment in their photosystems (Xiong and Bauer 2002; Sánchez-Baracaldo and Cardona 2020). Chl *a* is ubiquitously present in all oxygenic photosynthetic eukaryotes, and functions in both energy capture in the antenna light harvesting complexes (LHCs) and in driving electron transfer in the photosystem II (PSII) and photosystem I (PSI) reaction centers. Chl *b* is specific to the Viridiplantae although it has also been detected in prochlorophytes and *Acaryochloris*, unique photosynthetic prokaryotic groups that lack typical cyanobacterial phycobilin light-harvesting pigments (Palenik and Haselkorn 1992; Roche et al. 1996; Tomitani et al. 1999; Partensky et al. 2018). Chl *b* resides predominantly in the peripheral LHCs (Neilson and Durnford 2010) although it has been detected in the core complexes of some deep water marine chlorophytes (Kunugi et al. 2016).

Strict Chl *a/b* stoichiometry is required for optimal energy transfer during photosynthesis. Maintaining the correct Chl *a/b* ratios is a dynamic process and indicative of adaptation to different light conditions (Tanaka and Tanaka 2007). In addition to its role in light harvesting, appropriate levels of Chl *b* are required for accumulation, assembly, and stability of LHCs in algae (Bujaldon et al. 2017), bryophytes (Zhang et al. 2023), and angiosperms (Król et al. 1995; Reinbothe et al. 2006; Kim et al. 2009; Nick et al. 2013). Under low light conditions, the amount of Chl *b* relative to Chl *a* increases, leading to larger LHC antenna size to maximizes the surface area for light absorption (Kunugi et al. 2016; Kume et al. 2018; Ueno et al. 2019). Thus, the amount of Chl *b* is a major regulator of light-harvesting capacity.

Chlorophyll biosynthesis and turnover occurs through a series of complex multistep pathways (reviewed in Brzezowski et al. 2015; Willows 2020; Schumacher et al. 2022). Chl *b* is synthesized from Chl *a*, via the intermediate 7-hydromethyl chlorophyll *a*, by the action of a single enzyme: chlorophyllide *a* oxygenase (CAO) (Tanaka et al. 1998; Espineda et al. 1999; Oster et al. 2000; Mueller et al. 2012). Chl *b* can be converted back into Chl *a* through the action of chlorophyll *b* reductase (CBR) and 7-hydroxymethyl reductase (HCAR) to complete the cycle (Figure 1; reviewed in detail in Tanaka and Tanaka, 2019). This chlorophyll cycle plays an essential role in many

biological processes, including biogenesis of LHCs, antenna size regulation during light
acclimation, and chlorophyll degradation during senescence. While many of the intricacies of how
this pathway is regulated are still not fully understood (Tanaka and Tanaka 2019), it has been
suggested that CAO levels and activity are key determinants of Chl *b* accumulation.

Figure 1. The chlorophyll cycle in land plants. Chlorophyll *a* is converted to chlorophyll *b* via the
intermediate 7-hydromethyl chlorophyll *a*, by the action of the enzyme chlorophyllide *a* oxygenase
(CAO). In a negative feedback loop, high levels of chlorophyll *b* trigger the binding of a stromal Clp
protease to the degron motif (QDLLTIMILH) and the subsequent destabilization of CAO.
Chlorophyll b is necessary for the stabilization of Light Harvesting Complexes (LHC) in plants.
Excessive amounts of LHC or accumulation of energetically uncoupled non-functional LHCs (e.g.,
during senescence), triggers the accumulation of Chlorophyll b reductase (CBR) and the
conversion of chlorophyll *b* to chlorophyll *a* via the action of CBR and 7-hydroxymethyl reductase
(HCAR). This regulatory mechanism of CAO and CBR enables the fine- tuning and optimization of
chlorophyll *b* and LHC levels.

CAO catalyzes the conversion of the methyl group of Chl *a* into a formyl group to form Chl *b* through
two oxygenation reactions, likely using ferredoxin as a reductant (Oster et al. 2000). CAO is part of
the Rieske-type mononuclear non-heme iron oxygenase group of proteins (Gray et al. 2004). CAO
has three main regions: an N-terminal regulatory domain and a C-terminal catalytic domain, which
are connected by a more variable linker region (Nagata et al. 2004). The catalytic domain contains a
Rieske cluster and a mononuclear iron-binding domain, both of which are necessary for the
enzymatic activity of CAO (Yamasato et al. 2005; Kunugi et al. 2013). The regulatory domain has

been characterized in detail only in land plants. In *Arabidopsis*, a conserved sequence
(QDLLTIMILH) termed a degron was demonstrated to regulate Chl *b* accumulation by affecting CAO
protein turnover. In a negative feedback loop, high levels of Chl *b* trigger the binding of a stromal
Clp protease to the degron motif leading to destabilization of the CAO protein; this, in turn, halts
Chl *b* biosynthesis (Sakuraba et al., 2009; Nakagawara et al., 2007).

The evolutionary origins of CAO can be traced back to prochlorophytes, which encode a CAO gene
homolog involved in Chl biosynthesis (Tomitani et al. 1999; Nagata et al. 2004). Notably, the
prokaryotic CAO-like enzymes lack the regulatory N-terminal domain, suggesting that the Chl *b*-
dependant mechanism for controlling CAO stability was acquired in eukaryotes. Recent work on
several chlorophyte, charophyte, and embryophyte representative species suggested that the
degron recognition sequence first appeared in charophytes as an adaptation to high-light
environments in shallow waters (Kunugi et al. 2016). These conclusions, however, were based on
the presence of a degron in the CAO sequence of a single charophyte species (*Klebsormidinium*
flaccidum) and require further validation.

It has been postulated that gene duplication is the driver of plant evolution (Rensing 2014), but
previous reports suggested that the CAO protein is encoded by a single gene in plants and algae
(Kunugi et al. 2016; Schumacher et al. 2022). Two CAO genes were detected in rice (Lee et al. 2005)
although only one of these homologs is necessary for Chl *b* synthesis (Jung et al. 2021). Recent
work revealed CAO duplication in the bryophyte *Physcomitrium patens,* where both homologs
contribute to Chl *b* accumulation (Zhang et al. 2023). CAO duplication was also reported in two
Antarctic *Chlamydomonas* species , were the CAO duplicates appeared to be paralogous
suggesting an independent, rather than ancestral, duplication event (Cvetkovska et al. 2019; Poirier
et al. 2025).

These insights, combined with advances in plant and algal genome sequencing, prompted us to
further examine the *CAO* gene content and the presence of the regulatory domain across the
diverse plant and algal groups. In this work we identified and examined CAO sequence in diverse
species across Viridiplantae and demonstrate a widespread occurrence of CAO duplicates in this
group. Furthermore, we demonstrate that the degron sequence is absent in chlorophytes, but
present in charophyte and embryophytes, suggesting a different mechanism for Chl *b* regulation in
early aquatic lineages.

Materials and Methods

CAO sequence identification and analyses

To identify *CAO* sequences and duplication events with high accuracy, we used publicly available,
high quality, and fully annotated genomes. Chlorophyte and streptophyte genomes were retrieved
from Phycosom (Grigoriev et al. 2021), while embryophyte genomes were obtained from
Phytozome v13 (Goodstein et al. 2012). Putative *CAO* genes were identified with a tBLASTn search
with the *Chlamydomonas reinhardtii* (CrCAO; Cre01.g043350) and the *Arabidopsis thaliana*
(AtCAO; AT1G44446) full-length CAO peptides (Merchant et al. 2007; Lamesch et al. 2012). A strict
e-value cutoff of ≤e-30 was taken as a threshold, to avoid many unspecific hits due to the
conserved nature of the Fe-binding and Rieske cluster domains present among diverse plant and
algal proteins (Schmidt and Shaw 2001; Ferraro et al. 2005; Przybyla-Toscano et al. 2021). Putative

CAO sequences were manually curated, and truncated or low-quality sequences were excluded
from downstream analyses. Curated *CAO* sequences were aligned using MUSCLE (Edgar 2004).
The degron sequence (QDLLTUMILH) (Sakuraba et al. 2009) was identified by multiple sequence
alignment of the full-length CAO amino acid sequences and the regulatory N-terminal domain.
Sequence logos for each major group were generated by Geneious Prime 2024.11 (Dotmatics),
where the size of the letter reflects its frequence.

Phylogenetic Inference

The analyses described above resulted in 374 individual CAO sequences (Supplementary Table 1).
To avoid inaccurate inference and over-representation, we manually removed heterodimeric CAO
proteins, where the Rieske domain and the Fe-binding domain are located on different genes
(Kunugi et al. 2013) (Supplementary Table S1). We also removed identical CAO sequences isolated
from algal cultures identified as the same species but belonging to different strains or culture
collections (*Scenedesmus obliquus*, *Auxenochlorella prototothecoides*, *Chlorella sorokiniana*,
Ostreococcus tauri, *Mesostigma viride*, *Zygnema circumcarinatum*). This resulted in a 280 amino
acid alignment of 326 unique CAO sequences across 246 species.

These manually curated CAO sequences were aligned with MUSCLE and trimmed to remove gaps
and ambiguously aligned regions with Gblock (Castresana 2000; Talavera and Castresana 2007).
Maximum likelihood trees were inferred in the CIPRES Science Gateway (Miller et al. 2015) using
RAxML v8.0 (Stamatakis 2014) with 1000 bootstraps with the Whelan Goldman matrix for globular
proteins (WAG), a gamma shape parameter, and empirical estimation of invariable sites. Trees
were visualized in iTOL v6.8.2 (Letunic and Bork 2021). Previously inferred taxonomic information
for chlorophytes and streptophytes was obtained from AlgaeBase, University of Galway
(www.algaebase.org), and for land plants from the Plants Of the World Online, Royal Botanic
Gardens, Kew (https://powo.science.kew.org).

Results and Discussion

Occurrences of multiple CAO genes are widespread among the Viridiplantae

Our analysis demonstrates a widespread occurrence of *CAO* gene duplication across the
Viridiplantae. Multiple gene copies are more prevalent among the embryophytes, where ~41% of
examined species encode for more than one *CAO* gene. In comparison, we detected two CAO gene
copies in only ~11% of chlorophyte and ~20% of charophyte species (Table 1; Table 2,
Supplementary Table 1; Supplementary Table 2). All 23 species where we detect more than two
CAO genes are members of the Embryophyta. The modern-day cultivar of sugarcane (*Saccharum*
sp. R570) encodes for 6 *CAO* copies, the largest number we report. This species is a result of inter-
specific hybridization (*Saccharum officinarum* x *spontaneum;* Dumont et al. 2022), which has
resulted in a very large and highly redundant genome (~0.3 Gb, ~48,000 genes) (Healey et al. 2024).

We also detected multiple occurrences of heterodimeric *CAO* gene, where the functional CAO
protein is encoded by two genes (Table 1, Supplementary Table 1). Previous work in the
prasinophyte *Micromonas pusila* demonstrated that in this species CAO was encoded by two non-
homologous genes, one that possesses the mononuclear iron-binding motif (*MpCAO1*) and a
second one that encodes for the Rieske cluster (*MpCAO2*). It was shown that a combination of
both gene products was necessary to form an active heterodimeric CAO protein (Kunugi et al.

2013; Dey et al. 2023). We expand on this work by showing the presence of a heterodimeric CAO
encoded by two separate genes across the Mammiellophycea, a group within the prasinophytes
that contains *M. pusila* and several of its relatives (Worden et al. 2009) , the ecologically important
picoalgae *Ostreococcus tauri*, *Ostreococcus lucimarinus* and *Bathycoccus prasinos* (Palenik et al.
2007; Moreau et al. 2012; Blanc-Mathieu et al. 2014)*,* as well as the related *Chloropicon primus*
(Chloropicophyceae) (Lemieux et al. 2019).

We also show that the heterodimeric CAO is not restricted only within the Prasinophytes. We
detect a similar *CAO* gene arrangement in the genome of the multinucleate green alga *Caulerpa*
lentillifera (Bryopsidales) (Arimoto et al. 2019) , as well as in in the genome of the secondary
endosymbiont *Bigelowiella natans* (Chlorarachniophyceae) (Curtis et al. 2012). Phylogenetic
analysis based on plastid-encoded protein sequences revealed that chlorarachiniophytes acquired
their secondary plastids by the uptake of a filamentous green alga in the Bryopsidales (Suzuki et al.
2016), suggesting that a heterodimeric CAO has been inherited from the green algal endosymbiont.
Considering that CAO in prochlorophytes is encoded by a single gene that contains both the Rieske
and iron-binding domain (Nagata et al. 2004), it is likely that the occurrence of two separate genes
is not an ancestral condition but rather occurred after the primary endosymbiotic event. Indeed, it
has been suggested that acquisition of two separate genes encoding for key functional domains
may not be an evolutionarily difficult process. CAO likely functions as a trimer (Kunugi et al. 2013;
Liu et al. 2022), and the electron transfer from the Rieske to the mononuclear iron occurs between
two neighbouring CAO proteins. A functionally similar process likely occurs in the CAO enzyme
encoded by two separate genes.

Table 1: Number of species across the Viridiplantae with fully sequenced and annotated genomes
available through Phytozome (Total), compared to the number of species that encode for multiple
CAO genes (Multiple CAO) and number of species that encode for two *CAO* genes that form
heterodimeric complex (Heterodimeric CAO). The number in brackets represents the percentage of
species within the group that encode multiple CAO genes or heterodimeric *CAO* genes
239

Group	Total	Heterodimeric CAO	Multiple CAO
Chlorophytes*	70	11 (15.7%)	8 (11.5%)
Charophytes	10	0	2 (20%)
Non-vascular plants	6	0	3 (50%)
Lycophytes	3	0	2 (66%)
Gymnosperms	1	0	0
Angiosperms	151	0	62 (41.1%)

240   *The Chlorophyte group also includes the green algal secondary endosymbiont within the
241   Chlorarachniophyceae (*Bigelowiella natans*)
242

All examined streptophytes encode for at least one *CAO* gene (Supplementary Table 1) signifying
the importance of Chl *b* for plant physiology, but we were not able to identify a homolog of *CAO* in
several chlorophyte algal species. Not surprisingly, we did not detect a *CAO* homolog in the
invertebrate parasite *Helicosporidium* sp., a colorless protist that belongs within the
trebouxiophycean green algal clade (Tartar et al. 2003). As a consequence of its parasitic lifestyle,
Helicosporidium has lost nearly all genes associated with light harvesting, photosynthesis and
chlorophyll biogenesis (Pombert et al. 2014), including CAO.

We were also not able to conclusively identify a *CAO* gene in the genomes of the endolithic coral
holobiont *Ostreobium quekettii* (Bryopsidales) (Iha et al. 2021), the aquaculture species
Tetraselmis striata (Chlorodendrophyceae) (Steadman Tyler et al. 2019) and the halotolerant
Picocystis sp. ML (Picocystophyceae) (Junkins et al. 2019) . All identified sequences within the
genomes of these species had e-value higher than the cut-off (≥e-15) used in this work and high
similarity to pheophorbide *a* oxygenase (PAO), a Rieske-type protein involved in chlorophyll
degradation (Pružinská et al. 2003). The lack of *CAO* genes in these algae is surprising and could be
an artifact of genome assembly, as Chl *b* has been experimentally detected in all three species
(Roesler et al. 2002; Massé et al. 2020; Conlon et al. 2024).

It is tempting to speculate that these species may have a different mechanism for Chl *b*
biosynthesis or a CAO sequence significantly different than most algae and plants. For instance,
Ostreobium is an endolithic coral holobiont adapted to extreme shading, and has lost many genes
associated with photoprotection and light perception due to evolution in a light-limited
environment (Iha et al. 2021). This alga is also the only known eukaryote that has lost the gene
encoding light-dependent protochlorophyllide oxidoreductase (LPOR), a key enzyme that catalyzes
the conversion of protochlorophyllide to chlorophyllide *a*, a necessary step towards the synthesis
of Chl *a* (Reinbothe et al. 2010). Instead, this alga fully depends on dark operative POR (DPOR) to
synthesize Chl *a,* likely as an adaptation to extreme shading (Iha et al. 2021). The mechanism of
chlorophyll biosynthesis has not been examined in the species in detail, but it is clear that the
endolithic lifestyle has resulted in a unique physiology. Further work will shed light on the role of
CAO in this and other shade-tolerant species.

271 *CAO duplications are not ancestral within the Viridiplantae*

To determine whether the *CAO* gene duplication was an ancestral event, we preformed a
phylogenetic analysis of all high-quality CAO amino acid sequences (with the exception of
heterodimeric *CAO* genes). Phylogenetic patterns typically match previously shown relationship
among major Viridiplantae groups (Figure 2), although some relationships are poorly resolved
(particularly among the Rosids; Supplementary Figure 1). The presence of two *CAO* genes has been
previously reported in rice (*OsCAO1* and *OsCAO2*) where the two genes were positioned in tandem
on Chromosome 10, taken as evidence for a recent gene duplication (Lee et al. 2005) but this
appears to be the exception rather than the rule. In addition to rice, tandem *CAO* duplicates are
only found in a handful of species (Table 2). In most cases, *CAO* duplicates are located on different
chromosomes and are paralogous, hinting to independent non-recent duplication events (Figure 2;
Supplementary Figure 1). Thus, we surmise that encoding for multiple *CAO* genes is not related to
plant evolutionary patterns; rather, it may be that *CAO* duplication occurred independently in
different species.

There are several exceptions to this pattern, mainly in very closely related species where the
duplication may have originated in a recent shared ancestor. There are two possible cases in the
Trebouxiophyceae class within the Chlorophyte green algae, where *CAO* genes detected in closely
related *Chlorella* species share a close phylogenetic relationship (*Chlorella sorokiniana* UTEX1230
and *Chlorella variabilis*; *Chlorellaceae* sp. 4966 and *Chlorellaceae* sp. 4402) (Supplementary
Figure 1A). Within the bryophytes, *Spaghnum magellanicum* and *Sphagnum falax* both encode for
homologous *CAO* genes. These two species share a very close evolutionary relationship

(Piatkowski and Shaw 2019), and it is likely that a *CAO* duplication occurred in a recent shared
ancestor (Supplementary Figure 1B).

Figure 2: A phylogenetic tree of CAO genes inferred from using maximum likelihood analysis, using
326 unique CAO sequences across 246 species color-coded to represent the major Viridiplantae
groups. Scale bar indicates amino acid substitutions per site. Species with more than one detected
CAO gene are highlighted in red

Species that encode for multiple *CAO* genes within their genomes are more prevalent among the
Angiosperms. Within the monocots, *CAO* duplication arising from a shared ancestor is likely
present in the Asparagaceae, specifically among the closely related *Agave tequiliana*, *Yucca*
filamentosa and *Yucca aloifolia* (Ji et al. 2023), although we were only able to detect a single *CAO*
copy in *Asparagus officinalis* (Supplementary Figure 1C). We detected two *CAO* gene copies in two
Brachypodium species (*B. mexicanum* and *B. hybridum*) but not in their relatives from the same
genus (*B. arbuscula*, *B. stacei*, *B. sylvaticum* and *B. distachyon*). The two *CAO* genes detected in *B.*
hybridum appear to me more closely related to their counterparts from other species within their
genome (Supplementary Figure 1C), raising the possibility of an ancestral duplication within this
genus combined with gene loss in some species. All phylogenetic studies in *Brachypodium* support
a rapid and recent divergence from a common ancestor, but their precise relationship is often
unresolved or weakly supported (Catalan et al. 2016) making it difficult to accurately determine
evolutionary relationships between them.

CAO duplication may be ancestral in the two closely related Saxifragales *Kalanchoe laxiflora* and
Kalanchoe fedtschenkoi (Han et al. 2024). We detected two *CAO* genes in *K. fedtschenkoi* and four
in *K. laxiflora*, suggesting a duplication in a recent ancestor, followed by a second duplication
within *K. laxiflora* (Supplementary Figure 1E). Within the Fabids, four *Populus* species and their
relative *Salix purpurea*, each encode for two *CAO* gene copies that may be a result of an ancestral
duplication. The cultivated soybean *Glycine max* and its wild relative *Glycine soja* each encode for
four *CAO* gene copies (Supplementary Figure 1E) that more closely related to their homologs
between the two species. The phylogenetic separation of *CAO1/2* and *CAO3/4* is puzzling and
could simply be an artifact of phylogenetic uncertainty. Finally, all *Gossypium* species examined
here encode for four *CAO* genes, except *G. raimondii* which encodes for two copies
(Supplementary Figure 1E). The only other member of the Malvaceae examined, *Theobroma cacao,*
encodes for a single *CAO* gene, and although it clusters with *CAO* from *Gossypium* sp., the
relationship is poorly supported.

We show that *CAO* gene duplication is not a rarity among the Viridiplantae, but do all *CAO* genes
encode for an active protein? It is tempting to speculate that species that encode for multiple *CAO*
genes have a more robust capacity for Chl *b* biosynthesis through the gene dosage mechanism or a
finer control over Chl *b* accumulation though gene subfunctionalization (Kondrashov 2012; Panchy
et al. 2016) , but experimental evidence supporting these hypotheses is largely lacking.

Current work on species that encode for multiple CAO genes suggest that the mechanisms behind
the control of Chl *b* synthesis may be species specific. Mutational analysis in rice revealed that
only *OsCAO1* was necessary for Chl *b* biosynthesis. While *OsCAO1* knockout mutants exhibited
low Chl *b* levels, disordered thylakoid membrane arrangements, membrane damage due to
excessive ROS accumulation and decreased chilling tolerance (Lee et al. 2005; Jung et al. 2021;
Xiong et al. 2024), these effects were absent in *OsCAO2* knockouts (Lee et al. 2005). In contrast,
knocking out either of the two *CAO* homologs in *P. patens* lead to significantly reduced Chl *b* levels
suggesting a dependency on two active CAO enzymes for the appropriate control of Chl *b* synthesis
in this species (Zhang et al. 2023).

Table 2: Distribution of *CAO* duplicate genes across Viridiplantae. Only genes with confirmed Riske conserved cluster, Fe-binding domain (all species), and degron sequence (Streptophytes only) are listed. Accession numbers can be found in Supplementary Table 1 (continued on next page).

Taxon	Species	#CAO genes	Tandem
Chlorophyta			
Chlorophyceae	*Chlamydomonas priscui*	2	x
Chlorophyceae	*Volvox reticulferus*	2	✓
Chlorophyceae	*Tetradesmus deserticola*	2	x
Trebouxiophyceae	*Chlorella sp. A99*	2	x
Trebouxiophyceae	*Chlorella sorokiniana*	2	x
Trebouxiophyceae	*Chlorellaceae sp.*	2	x
Trebouxiophyceae	*Micractinium conductrix*	2	x
Pedinophyceae	*Pedinomonas minor*	2	x
Charophyta			
Zygnematophyceae	*Mesotaenium endlicherianum*	2	x
Zygnematophyceae	*Spirogloea muscicola*	2	x
Bryophyta			
Bryopsida	*Physcomitrium patens*	2	x
Sphagnopsida	*Sphagnum fallax*	2	x
Sphagnopsida	*Sphagnum magellanicum*	3	x
Lycophyta			
Equisetopsida	*Diphasiastrum complanatum*	4	✓✓
Equisetopsida	*Ceratopteris richardii*	2	x
Angiosperms			
Laurales	*Cinnamomum kanehirae*	2	✓
Nymphaeales	*Nymphaea colorata*	2	x
Asparagales	*Agave tequilana*	2	x
Asparagales	*Yucca aloifolia*	3	x
Asparagales	*Yucca filamentosa*	3	x
Zingiberales	*Musa acuminata v1*	2	x
Alismatales	*Spirodela polyrhiza v2*	2	x
Poales	*Typha latifolia*	2	x
Poales	*Eleusine coracana*	2	x
Poales	*Oryza sativa*	2	✓
Poales	*Thinopyrum intermedium*	3	x
Poales	*Brachypodium hybridum*	2	x
Poales	*Brachypodium mexicanum*	2	x
Poales	*Zea mays*	2	x
Poales	*Andropogon gerardi*	3	x
Poales	*Miscanthus sinensis*	2	x
Poales	*Panicum virgatum*	3	x
Poales	*Saccharum officinarum*	6	x
Caryophyllales	*Chenopodium quinoa*	2	x
Gentianales	*Coffea arabica geisha*	2	x

Table 2:(continued)

Taxon	Species	#CAO genes	Tandem
Apiales	*Hydrocotyle leucocephala*	2	x
Boraginales	*Ehretia anacua*	2	x
Asterales	*Helianthus annuus*	2	x
Lamiales	*Olea europaea*	2	x
Saxifragales	*Kalanchoe fedtschenkoi*	2	x
Saxifragales	*Kalanchoe laxiflora*	4	x
Fabales	*Arachis hypogaea*	2	x
Fabales	*Chamaecrista fasciculata*	2	x
Fabales	*Glycine max*	4	x
Fabales	*Glycine soja*	4	x
Fabales	*Lupinus albus*	3	x
Fabales	*Phaseolus vulgaris*	2	x
Fagales	*Carya illinoinensis*	2	x
Rosales	*Fragaria x ananassa*	3	x
Rosales	*Malus domestica*	2	x
Malpighiales	*Linum usitatissimum*	4	x
Malpighiales	*Populus deltoides*	2	x
Malpighiales	*Populus nigra x maximowiczii*	2	x
Malpighiales	*Populus tremula x alba*	2	x
Malpighiales	*Populus trichocarpa*	2	x
Malpighiales	*Salix purpurea*	2	x
Sapindales	*Anacardium occidentale*	3	x
Malvales	*Gossypium barbadense*	4	x
Malvales	*Gossypium darwinii*	4	x
Malvales	*Gossypium hirsutum*	4	x
Malvales	*Gossypium mustelinum*	4	x
Malvales	*Gossypium raimondii*	2	x
Malvales	*Gossypium tomentosum*	4	x
Brassicales	*Alyssum linifolium*	2	x
Brassicales	*Caulanthus amplexicaulis*	2	x
Brassicales	*Iberis amara*	2	x
Brassicales	*Lepidium sativum*	2	x
Brassicales	*Stanleya pinnata*	2	x
Brassicales	*Brassica juncea*	3	x
Brassicales	*Brassica oleracea*	2	x
Brassicales	*Cakile maritima*	2	x
Brassicales	*Crambe hispanica*	2	x
Brassicales	*Eruca vesicaria*	2	x
Brassicales	*Sinapis alba*	4	x
Brassicales	*Camelina sativa*	3	x
Brassicales	*Isatis tinctoria*	4	x

Significantly, all experimental studies to date suggest that *CAO* duplicates may be differently
regulated at the level of gene expression. In rice, the expression of *OsCAO1* is light-inducible, while
OsCAO2 accumulates during periods of darkness and is expressed mainly in non-photosynthetic
tissues (Lee et al. 2005). In *P. patens*, both *PpCAO1* and *PpCAO2* are expressed in both light and
dark conditions, but only *PpCAO1* is upregulated with prolonged light exposure. Similarly, in two
Chlamydomonas species that harbour *CAO* duplications (*Chlamydomonas priscui* and
Chlamydomonas sp. ICE-MDV), only one *CAO* homolog is upregulated in response to increases in
light intensity, while the second one is constitutively expressed (Poirier et al. 2025). The role of light
in regulating *CAO* expression has been demonstrated previously in *Arabidopsis* and tobacco
(Pattanayak et al. 2005; Tanaka and Tanaka 2005; Biswal et al. 2012, 2024), species with a single
CAO gene copy, but the implication of light regulation is not known in species that harbour multiple
CAO genes. Two gene copies with different expression patterns could provide a fine-tuned control
of Chl *b* levels and photosynthetic performance under varying light conditions, but careful
experimental work is needed to substantiate this hypothesis.

Evolutionary implications of the CAO degron sequence

In addition to light-dependant regulation at the level of transcription, the main regulation of CAO
occurs post-translationally. Early work in *A. thaliana* demonstrated that this regulation is governed
by the presence of a degron sequence (QDLLTIMILH) in the protein's N-terminal A domain that acts
as a docking site for a chloroplast Clp protease (Yamasato et al. 2005; Nakagawara et al. 2007;
Sakuraba et al. 2009). It has been hypothesized that the degron sequence is "hidden" within the A
domain when there are insufficient amounts of Chl *b*, but accumulation of this product causes a
conformational change in the CAO protein, exposing the degron, and giving access to the protease
(Tanaka and Tanaka 2011). The evolution of this negative-feedback regulatory loop has not been
examined in detail, but it has been suggested that the degron was acquired early in the
Viridiplantae evolution as a mechanism for high light adaptation (Kunugi et al. 2016).

Here we show that the degron sequence is highly conserved among all land plants (86% identity),
less conserved among charophytes and non-vascular plants (68% and 66% identity respectively)
and very poorly conserved or absent from chlorophytes (49% identity) (Figure 3; Supplementary
Figure 2). Land plants (both vascular and non-vascular) have several highly conserved sites across
all species (Asparagine at position 2, Isoleucine at position 6, Histidine at position 10), whereas
these positions are only moderately conserved in charophytes, and poorly conserved or absent in
chlorophytes. In contrast, the Rieske cluster [CXH(X)15-17CXXH] exhibits an 84% pairwise identity
among all species, where the two histidine (positions 3 and 23) and cysteines (positions 1 and 20)
that coordinate the metallocentre [2Fe-2S] (Przybyla-Toscano et al. 2021) are present in all
examined sequences. Nearly all residues are conserved among the Viridiplantae in the C-terminal
Fe-binding domain (96% pairwise identity). Overall, these results suggest that the mechanism of
Chl *b* biosynthesis involving the Rieske and Fe-binding domains is ancestral and conserved among
all photosynthetic species, but this is not the case for the regulatory degron domain.

The absence of a highly conserved degron sequence may have several implications in the evolution
of efficient light harvesting. For instance, the N-regulatory domain (including the degron) is
completely absent in marine eukaryotic Mammiellophycea that encode for a heterodimeric *CAO*
gene, as well as in their deep-water relatives from the order Palmophyllales. These species
typically have low Chl *a/b* ratios and, unlike land plants, can incorporate Chl *b* into their core

antennae. This was interpreted as a necessary adaptation for efficient light harvesting in deep
marine environments characterized by low levels of blue-green light (Kunugi et al. 2016).

Figure 3: Sequence logos of the conserved motifs identified in the CAO protein sequence. All
sequence logos are superimposed on a graphical representation of plant CAO sequences (not to
scale) showing the regulatory degron motif, conserved Rieske domain cluster and Fe-binding
domain (Sakuraba et al., 2009; Liu et al., 2022). The logos were generated for all chlorophyte algae,
charophyte algae, non-vascular plants, and vascular plants. The sequences on top of each logo are
the consensus sequence with the most common base in each sequence alignment.

It has been suggested that the degron sequence evolved in shallow-water algae and land plants as
a means to regulate Chl *b* accumulation and limit Chl *b* incorporation into the reaction center in an
environment where they are exposed to high light intensities (Kunugi et al. 2016). Indeedsuch
species exclusively harbour Chl *a* in their core complexes, but this appears to be controlled by
pigment availability rather than the chemical nature of the two chlorophyll types. The binding
affinity of Chl *a* and *b* to the PSI complex is similar (Ikegami et al. 2007) and cyanobacteria that do
not naturally produce Chl *b* successfully incorporated this pigment into their reaction centers
when transformed with plant CAO (Satoh et al. 2001; Xu et al. 2001). Incorporation of Chl *b* into the
reaction center accompanied by increased Chl *b* accumulation was also observed in *Arabidopsis*
when full length CAO was overexpressed or replaced with a prokaryotic CAO that lacked a degron
sequence (Hirashima et al. 2006; Sakuraba et al. 2009).

The chlorophyll cycle has not been examined in detail in the core chlorophytes (Chlorophyceae,
Trebouxiophyceae, Ulvophyceae), a diverse group that inhabits marine, freshwater and terrestrial
habitats (Leliaert et al. 2012). Although many chlorophytes thrive in high light, others are adapted to
light limited environments, including deep waters. As a result, these species have developed
adaptations allowing for efficient light harvesting in low light. For instance, *Chlamydomonas priscui*
is endemic to the depths of the Antarctic Lake Bonney, an environment characterized by extreme
shading and dominated by blue-green wavelengths (Neale and Priscu 1995). This alga displays a
host of adaptations to extreme shading including a large LHC antennae size (reviewed in

Cvetkovska et al. 2017; Hüner et al. 2022). As suggested for marine prasinophytes (Kunugi et al.
2016), such low-light adapted chlorophytes could benefit from increased Chl *b* levels and could
thrive without a Chl *b*-dependent feedback mechanisms.

Despite the lack of a conserved degron sequence to regulate CAO protein levels (Figure 2),
chlorophytes generally have maintained the ability to alter their Chl *a/b* ratio in response to light
(Leliaert et al. 2012). Transcriptional control of CAO levels was postulated to be a minor aspect of
adjusting Chl *b* levels in plants (Tanaka and Tanaka 2019), but this process could play a bigger role
in chlorophytes. Finally, it is possible that a different, yet unidentified, mechanism exists for Chl *b*
turnover independent of a degron sequence, allowing these species to modify their Chl *a/b* ratios. A
detailed experimental investigation on the control and mechanism of Chl *b* biosynthesis is clearly
needed among the Chlorophytes.

Conclusion and Drawbacks

Our work suggests that CAO gene duplication is widespread among the Viridiplantae, likely
originating from multiple independent duplication events throughout plant evolution. While we
show that the presence of multiple *CAO* gene copies is highly prevalent in land plants, these
results must be interpreted with caution. Many more angiosperm genomes are available in public
databases, while other plant and algal groups are not as well represented. Furthermore, many
genome sequencing initiatives are focused on plants of economic and agricultural importance, in
which domestication and selective breeding may have affectted the genomic content (Turner-
Hissong et al. 2020).While the exact impact of CAO duplication is unknown, the study of transgenic
tobacco overexpressing CAO (Biswal et al. 2012) showed that an increase in Chl *b* may be a
strategy to enhance CO_2 assimilation, delay senescence, and enhance productivity - traits that
would be attractive in crop breeding programs (Voitsekhovskaja and Tyutereva 2015). Therefore,
the high prevalence of CAO duplication seen in land plants in this study may have been selected for
during plant breeding, as the traits resulting from the duplication may enhance crop productivity.
Our understanding of the prevalence and evolution of CAO at the genetic level will be improved in
future years, as more algal and non-vascular plant genomes become available. Furthermore, there
are still numerous complex and outstanding questions regarding the control of Chl *b* accumulation
in photosynthetic species. In particular, the chlorophytes are an understudied group when it
comes to Chl *b* regulating mechanism. Detailed genomic and experimental studies will shed light
on the role and regulation of Chl *b* in these species and will deepen our understanding on this
crucial photosynthetic pigment.

Supplementary Information

Supplementary Table 1: A summary of genes encoding for chlorophyll *a* oxygenase (CAO) across
the Viridiplantae used in this work. All sequences were obtained from Phycocosm (Chlorophytes,
Euglenophytes, Cercozoa, Charophytes) or Phytozome (Bryophytes, Marchantiophytes,
Lycophytes, Gymnosperms, Angiosperms). Accession numbers represent the unique IDs assigned
to each gene (Phycocosm: Gene portal|Protein ID|Gene Model; Phytozome: Gene Model).

Supplementary Table 2: A summary of the occurrence of CAO duplicate genes across the major
groups within Viridiplantae. The number in brackets represents the proportion of species which
encode for multiple copies of the CAO gene (Mutiple CAO) or encode for two CAO genes that form
heterodimeric complex (Heterodimeric CAO).

Supplementary Figure 1: A phylogenetic tree of CAO genes inferred from using maximum
likelihood analysis. The node labels show the bootstrap support, with dashed lines indicating <50
bootstrap support. Scale bar indicates 0.1 amino acid substitutions per site. Species with more
than one detected CAO gene are highlighted in red. Cases where CAO duplication may have arisen
in common shared ancestor are highlighted with a blue arrow. The inset shows the outline of the
full CAO phylogenetic tree inferred from using maximum likelihood analysis, using 326 unique CAO
sequences across 246 species color coded to represent the major Viridiplantae groups. The
position of the chlorophytes (green) is highlighted with a black box. (A) Chlorophytes; (B)
Charophytes, bryophytes (including marachantiophytes), gymnosperms and magnolids; (C)
Monocots; (D) Basal Eudicots and Asterids; (E) Basal Rosids and Fabids; (F) Malvids.

Supplementary Figure 2: Multiple sequence alignment of the predicted CAO amino acid
sequences from Chlorophytes (A), Charophytes (B), non-vascular plants (C) and vascular plants
(D). Global alignments were performed with Clustal Omega. The colour of the residue depends on
the degree of conservation among species (blue, fully conserved; green, 80–100% similar; yellow,
60–80% similar; gray, < 60% similar). As a reference, the CAO sequence from Arabidopsis thaliana
is present in all alignments (top), with the key domains annotated (green, Degron sequence; blue,
conserved Rieske cluster; purple, Fe-binding domain)

References

Arimoto A, Nishitsuji K, Higa Y, et al (2019) A siphonous macroalgal genome suggests convergent
functions of homeobox genes in algae and land plants. DNA Research 26:183–192.
https://doi.org/10.1093/dnares/dsz002

Biswal AK, Pattanayak GK, Pandey SS, et al (2012) Light Intensity-Dependent Modulation of
Chlorophyll *b* Biosynthesis and Photosynthesis by Overexpression of Chlorophyllide *a*
Oxygenase in Tobacco. Plant Physiol 159:433–449. https://doi.org/10.1104/pp.112.195859

Biswal AK, Pattanayak GK, Ruhil K, et al (2024) Reduced expression of chlorophyllide *a* oxygenase
(CAO) decreases the metabolic flux for chlorophyll synthesis and downregulates
photosynthesis in tobacco plants. Physiol Mol Biol Plants 30:1–16.
https://doi.org/10.1007/s12298-023-01395-5

Blanc-Mathieu R, Verhelst B, Derelle E, et al (2014) An improved genome of the model marine alga
Ostreococcus tauri unfolds by assessing Illumina de novo assemblies. BMC Genomics
15:1–12. https://doi.org/10.1186/1471-2164-15-1103

Blankenship RE (2010) Early Evolution of Photosynthesis. Plant Physiol 154:434–438.
https://doi.org/10.1104/pp.110.161687

Brzezowski P, Richter AS, Grimm B (2015) Regulation and function of tetrapyrrole biosynthesis in
plants and algae. Biochimica et Biophysica Acta - Bioenergetics 1847:968–985.
https://doi.org/10.1016/j.bbabio.2015.05.007

Bujaldon S, Kodama N, Rappaport F, et al (2017) Functional Accumulation of Antenna Proteins in
Chlorophyll b-Less Mutants of *Chlamydomonas reinhardtii*. Molecular Plant 10:115–130.
https://doi.org/10.1016/j.molp.2016.10.001

Cardona T (2019) Thinking twice about the evolution of photosynthesis. Open Biology 9:180246.
https://doi.org/10.1098/rsob.180246

Castresana J (2000) Selection of Conserved Blocks from Multiple Alignments for Their Use in
Phylogenetic Analysis. Molecular Biology and Evolution 17:540–552.
https://doi.org/10.1093/oxfordjournals.molbev.a026334

Catalan P, López-Álvarez D, Díaz-Pérez A, et al (2016) Phylogeny and Evolution of the Genus
Brachypodium. In: Vogel JP (ed) Genetics and Genomics of Brachypodium. Springer
International Publishing, Cham, pp 9–38

Conlon T, Parkes R, Fierli D, Touzet N (2024) Comparative pigment and fatty acid profiling of marine
species within the chlorophyte genus *Tetraselmis*. Food Bioscience 58:103660.
https://doi.org/10.1016/j.fbio.2024.103660

Curtis BA, Tanifuji G, Burki F, et al (2012) Algal genomes reveal evolutionary mosaicism and the fate
of nucleomorphs. Nature 492:59–65. https://doi.org/10.1038/nature11681

Cvetkovska M, Hüner NPA, Smith DR (2017) Chilling out: the evolution and diversification of
psychrophilic algae with a focus on Chlamydomonadales. Polar Biol 40:1169–1184.
https://doi.org/10.1007/s00300-016-2045-4

Cvetkovska M, Orgnero S, Hüner NPA, Smith DR (2019) The enigmatic loss of light-independent
chlorophyll biosynthesis from an Antarctic green alga in a light-limited environment. New
Phytol 222:651–656. https://doi.org/10.1111/nph.15623

de Vries J, Archibald JM (2018) Plant evolution: landmarks on the path to terrestrial life. New
Phytologist 217:1428–1434. https://doi.org/10.1111/nph.14975

Dey D, Tanaka R, Ito H (2023) Structural Characterization of the Chlorophyllide *a* Oxygenase (CAO)
Enzyme Through an In Silico Approach. J Mol Evol 91:225–235.
https://doi.org/10.1007/s00239-023-10100-9

Dumont T, Barau L, Thong-Chane A, et al (2022) Sugarcane Breeding in Reunion: Challenges,
Achievements and Future Prospects. Sugar Tech 24:181–192.
https://doi.org/10.1007/s12355-021-00998-0

Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space
complexity. BMC Bioinformatics 5:113. https://doi.org/10.1186/1471-2105-5-113

Espineda CE, Linford AS, Devine D, Brusslan JA (1999) The AtCAO gene, encoding chlorophyll *a*
oxygenase, is required for chlorophyll *b* synthesis in *Arabidopsis thaliana*. Proceedings of
the National Academy of Sciences of the United States of America 96:10507–10511.
https://doi.org/10.1073/pnas.96.18.10507

Ferraro DJ, Gakhar L, Ramaswamy S (2005) Rieske business: Structure–function of Rieske non-
heme oxygenases. Biochemical and Biophysical Research Communications 338:175–190.
https://doi.org/10.1016/j.bbrc.2005.08.222

Goodstein DM, Shu S, Howson R, et al (2012) Phytozome: a comparative platform for green plant
genomics. Nucleic Acids Research 40:D1178–D1186. https://doi.org/10.1093/nar/gkr944

Gray J, Wardzala E, Yang M, et al (2004) A small family of LLS1-related non-heme oxygenases in
plants with an origin amongst oxygenic photosynthesizers. Plant Molecular Biology 54:39–
54

Grigoriev IV, Hayes RD, Calhoun S, et al (2021) PhycoCosm, a comparative algal genomics
resource. Nucleic Acids Research 49:D1004–D1011. https://doi.org/10.1093/nar/gkaa898

Guiry MD (2024) How many species of algae are there? A reprise. Four kingdoms, 14 phyla, 63
classes and still growing. Journal of Phycology 60:214–228.
https://doi.org/10.1111/jpy.13431

Han S, Zhang S, Yi R, et al (2024) Phylogenomics and plastomics offer new evolutionary
perspectives on Kalanchoideae (Crassulaceae). Annals of Botany 133:585–604.
https://doi.org/10.1093/aob/mcae017

Healey AL, Garsmeur O, Lovell JT, et al (2024) The complex polyploid genome architecture of
sugarcane. Nature 628:804–810. https://doi.org/10.1038/s41586-024-07231-4

Hirashima M, Satoh S, Tanaka R, Tanaka A (2006) Pigment Shuffling in Antenna Systems Achieved
by Expressing Prokaryotic Chlorophyllide a Oxygenase in *Arabidopsis*. The Journal of
Biological Chemistry 281:15385–15393. https://doi.org/10.1074/jbc.M602903200

Hüner NPA, Smith DR, Cvetkovska M, et al (2022) Photosynthetic adaptation to polar life: Energy
balance, photoprotection and genetic redundancy. Journal of Plant Physiology 268:153557.
https://doi.org/10.1016/j.jplph.2021.153557

Iha C, Dougan KE, Varela JA, et al (2021) Genomic adaptations to an endolithic lifestyle in the coral-
associated alga *Ostreobium*. Current Biology 31:1393-1402.e5.
https://doi.org/10.1016/j.cub.2021.01.018

Ikegami I, Satoh S, Aoki M (2007) Binding Affinity of Chl *b* for the Chl *a* -Binding Sites in PSI Core
Complexes. Plant Cell Physiology 48:1092–1097. https://doi.org/10.1093/pcp/pcm078

Ji Y, Landis JB, Yang J, et al (2023) Phylogeny and evolution of Asparagaceae subfamily Nolinoideae:
new insights from plastid phylogenomics. Annals of Botany 131:301–312.
https://doi.org/10.1093/aob/mcac144

Jung YJ, Lee HJ, Yu J, et al (2021) Transcriptomic and physiological analysis of OsCAO1 knockout
lines using the CRISPR/Cas9 system in rice. Plant Cell Rep 40:1013–1024.
https://doi.org/10.1007/s00299-020-02607-y

Junkins EN, Stamps BW, Corsetti FA, et al (2019) Draft Genome Sequence of *Picocystis* sp. Strain
ML, Cultivated from Mono Lake, California. Microbiol Resour Announc 8:e01353-18.
https://doi.org/10.1128/MRA.01353-18

Kim EH, Li XP, Razeghifard R, et al (2009) The multiple roles of light-harvesting chlorophyll *a/b*-
protein complexes define structure and optimize function of *Arabidopsis* chloroplasts: A

study using two chlorophyll b-less mutants. Biochimica et Biophysica Acta - Bioenergetics
1787:973–984. https://doi.org/10.1016/j.bbabio.2009.04.009

Kondrashov FA (2012) Gene duplication as a mechanism of genomic adaptation to a changing
environment. Proceedings of the Royal Society B: Biological Sciences 279:5048–5057.
https://doi.org/10.1098/rspb.2012.1108

Król M, Spangfort MD, Huner NPA, et al (1995) Chlorophyll *a/b*-binding proteins, pigment
conversions, and early light-induced proteins in a chlorophyll *b*-less Barley mutant. Plant
Physiology 107:873–883. https://doi.org/10.1104/pp.107.3.873

Kume A, Akitsu T, Nasahara KN (2018) Why is chlorophyll *b* only used in light-harvesting systems?
Journal of Plant Research 131:961–972. https://doi.org/10.1007/s10265-018-1052-7

Kunugi M, Satoh S, Ihara K, et al (2016) Evolution of Green Plants Accompanied Changes in Light-
Harvesting Systems. Plant and Cell Physiology 57:1231–1243.
https://doi.org/10.1093/pcp/pcw071

Kunugi M, Takabayashi A, Tanaka A (2013) Evolutionary changes in chlorophyllide *a* oxygenase
(CAO) structure contribute to the acquisition of a new light-harvesting complex in
micromonas. Journal of Biological Chemistry 288:19330–19341.
https://doi.org/10.1074/jbc.M113.462663

Lamesch P, Berardini TZ, Li D, et al (2012) The Arabidopsis Information Resource (TAIR): improved
gene annotation and new tools. Nucleic Acids Res 40:D1202-1210.
https://doi.org/10.1093/nar/gkr1090

Lee S, Kim JH, Eun SY, et al (2005) Differential regulation of chlorophyll *a* oxygenase genes in rice.
Plant Molecular Biology 57:805–818. https://doi.org/10.1007/s11103-005-2066-9

Leliaert F, Smith DR, Moreau H, et al (2012) Phylogeny and Molecular Evolution of the Green Algae.
Critical Reviews in Plant Sciences 31:1–46. https://doi.org/10.1080/07352689.2011.615705

Lemieux C, Turmel M, Otis C, Pombert J-F (2019) A streamlined and predominantly diploid genome
in the tiny marine green alga *Chloropicon primus*. Nat Commun 10:4061.
https://doi.org/10.1038/s41467-019-12014-x

Letunic I, Bork P (2021) Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display
and annotation. Nucleic Acids Res 49:W293–W296. https://doi.org/10.1093/nar/gkab301

Liu J, Knapp M, Jo M, et al (2022) Rieske Oxygenase Catalyzed C–H Bond Functionalization
Reactions in Chlorophyll *b* Biosynthesis. ACS Cent Sci 8:1393–1403.
https://doi.org/10.1021/acscentsci.2c00058

Massé A, Tribollet A, Meziane T, et al (2020) Functional diversity of microboring Ostreobium algae
isolated from corals. Environmental Microbiology 22:4825–4846.
https://doi.org/10.1111/1462-2920.15256

Merchant SS, Prochnik SE, Vallon O, et al (2007) The Chlamydomonas genome reveals the evolution of key animal and plant functions. Science 318:245–250. https://doi.org/10.1126/science.1143609

Miller MA, Schwartz T, Pickett BE, et al (2015) A RESTful API for Access to Phylogenetic Tools via the CIPRES Science Gateway. Evol Bioinform Online 11:43–48. https://doi.org/10.4137/EBO.S21501

Moreau H, Verhelst B, Couloux A, et al (2012) Gene functionalities and genome structure in *Bathycoccus prasinos* reflect cellular specializations at the base of the green lineage. Genome Biol 13:1–16. https://doi.org/10.1186/gb-2012-13-8-r74

Mueller AH, Dockter C, Gough SP, Lundqvist U (2012) Characterization of Mutations in Barley fch2 Encoding Chlorophyllide *a* Oxygenase. Plant and Cell Physiology 53:1232–1246. https://doi.org/10.1093/pcp/pcs062

Nagata N, Satoh S, Tanaka R, Tanaka A (2004) Domain structures of chlorophyllide *a* oxygenase of green plants and Prochlorothrix hollandica in relation to catalytic functions. Planta 218:1019–1025. https://doi.org/10.1007/s00425-003-1181-6

Nakagawara E, Sakuraba Y, Yamasato A, et al (2007) Clp protease controls chlorophyll *b* synthesis by regulating the level of chlorophyllide a oxygenase. The Plant Journal 49:800–809. https://doi.org/10.1111/j.1365-313X.2006.02996.x

Neale PJ, Priscu JC (1995) The Photosynthetic Apparatus of Phytoplankton from a Perennially Ice-Covered Antarctic Lake: Acclimation to an Extreme Shade Environment. Plant and Cell Physiology 36:253–263. https://doi.org/10.1093/oxfordjournals.pcp.a078757

Neilson JAD, Durnford DG (2010) Structural and functional diversification of the light-harvesting complexes in photosynthetic eukaryotes. Photosynthesis Research 106:57–71. https://doi.org/10.1007/s11120-010-9576-2

Nick S, Meurer J, Soll J, Ankele E (2013) Nucleus-encoded light-harvesting chlorophyll a/b proteins are imported normally into chlorophyll b-free chloroplasts of Arabidopsis. Molecular Plant 6:860–871. https://doi.org/10.1093/mp/sss113

Oster U, Tanaka R, Tanaka A, Rüdiger W (2000) Cloning and functional expression of the gene encoding the key enzyme for chlorophyll *b* biosynthesis (CAO) from *Arabidopsis thaliana*. Plant Journal 21:305–310. https://doi.org/10.1046/j.1365-313X.2000.00672.x

Palenik B, Grimwood J, Aerts A, et al (2007) The tiny eukaryote Ostreococcus provides genomic insights into the paradox of plankton speciation. Proceedings of the National Academy of Sciences 104:7705–7710. https://doi.org/10.1073/pnas.0611046104

Palenik B, Haselkorn R (1992) Multiple evolutionary origins of prochlorophytes, the chlorophyll *b*-containing prokaryotes. Nature 355:265–267. https://doi.org/10.1038/355265a0

Panchy N, Lehti-Shiu M, Shiu S-H (2016) Evolution of Gene Duplication in Plants. Plant Physiol 171:2294–2316. https://doi.org/10.1104/pp.16.00523

Partensky F, Six C, Ratin M, et al (2018) A novel species of the marine cyanobacterium
Acaryochloris with a unique pigment content and lifestyle. Sci Rep 8:9142.
https://doi.org/10.1038/s41598-018-27542-7

Pattanayak GK, Biswal AK, Reddy VS, Tripathy BC (2005) Light-dependent regulation of chlorophyll
b biosynthesis in chlorophyllide *a* oxygenase overexpressing tobacco plants. Biochemical
and Biophysical Research Communications 326:466–471.
https://doi.org/10.1016/j.bbrc.2004.11.049

Piatkowski BT, Shaw AJ (2019) Functional trait evolution in Sphagnum peat mosses and its
relationship to niche construction. New Phytologist 223:939–949.
https://doi.org/10.1111/nph.15825

Poirier MC, Fugard K, Cvetkovska M (2025) Light quality affects chlorophyll biosynthesis and
photosynthetic performance in Antarctic Chlamydomonas. Photosynth Res 163:9.
https://doi.org/10.1007/s11120-024-01127-0

Pombert J-F, Blouin NA, Lane C, et al (2014) A Lack of Parasitic Reduction in the Obligate Parasitic
Green Alga Helicosporidium. PLoS Genet 10:e1004355.
https://doi.org/10.1371/journal.pgen.1004355

Pružinská A, Tanner G, Anders I, et al (2003) Chlorophyll breakdown: Pheophorbide *a* oxygenase is
a Rieske-type iron–sulfur protein, encoded by the accelerated cell death 1 gene.
Proceedings of the National Academy of Sciences 100:15259–15264.
https://doi.org/10.1073/pnas.2036571100

Przybyla-Toscano J, Boussardon C, Law SR, et al (2021) Gene atlas of iron-containing proteins in
Arabidopsis thaliana. The Plant Journal 106:258–274. https://doi.org/10.1111/tpj.15154

Reinbothe C, Bakkouri ME, Buhr F, et al (2010) Chlorophyll biosynthesis: spotlight on
protochlorophyllide reduction. Trends in Plant Science 15:614–624.
https://doi.org/10.1016/j.tplants.2010.07.002

Reinbothe C, Bartsch S, Eggink LL, et al (2006) A role for chlorophyllide *a* oxygenase in the regulated
import and stabilization of light-harvesting chlorophyll *a/b* proteins. Proceedings of the
National Academy of Sciences of the United States of America 103:4777–4782.
https://doi.org/10.1073/pnas.0511066103

Rensing SA (2014) Gene duplication as a driver of plant morphogenetic evolution. Current Opinion
in Plant Biology 17:43–48. https://doi.org/10.1016/j.pbi.2013.11.002

Roche JL, van der Staay GWM, Partensky F, et al (1996) Independent evolution of the
prochlorophyte and green plant chlorophyll *a/b* light-harvesting proteins. Proceedings of
the National Academy of Sciences 93:15244–15248.
https://doi.org/10.1073/pnas.93.26.15244

Roesler CS, Culbertson CW, Etheridge SM, et al (2002) Distribution, production, and ecophysiology
of *Picocystis* strain ML in Mono Lake, California. Limnology and Oceanography 47:440–452.
https://doi.org/10.4319/lo.2002.47.2.0440

Ruggiero MA, Gordon DP, Orrell TM, et al (2015) A Higher Level Classification of All Living
Organisms. PLoS One 10:e0119248. https://doi.org/10.1371/journal.pone.0119248

Sakuraba Y, Tanaka R, Yamasato A, Tanaka A (2009) Determination of a chloroplast degron in the
regulatory domain of chlorophyllide *a* oxygenase. Journal of Biological Chemistry
284:36689–36699. https://doi.org/10.1074/jbc.M109.008144

Sánchez-Baracaldo P, Cardona T (2020) On the origin of oxygenic photosynthesis and
Cyanobacteria. New Phytologist 225:1440–1446. https://doi.org/10.1111/nph.16249

Satoh S, Ikeuchi M, Mimuro M, Tanaka A (2001) Chlorophyll *b* Expressed in Cyanobacteria
Functions as a Light-harvesting Antenna in Photosystem I through Flexibility of the Proteins.
The Journal of Biological Chemistry 276:4293–4297.
https://doi.org/10.1074/jbc.M008238200

Schmidt CL, Shaw L (2001) A Comprehensive Phylogenetic Analysis of Rieske and Rieske-Type
Iron-Sulfur Proteins. J Bioenerg Biomembr 33:9–26.
https://doi.org/10.1023/A:1005616505962

Schumacher I, Menghini D, Ovinnikov S, et al (2022) Evolution of chlorophyll degradation is
associated with plant transition to land. The Plant Journal 109:1473–1488.
https://doi.org/10.1111/tpj.15645

Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics 30:1312–1313. https://doi.org/10.1093/bioinformatics/btu033

Steadman Tyler CR, Hovde BT, Daligault HE, et al (2019) High-Quality Draft Genome Sequence of
the Green Alga *Tetraselmis striata* (Chlorophyta) Generated from PacBio Sequencing.
Microbiology Resource Announcements 8:10.1128/mra.00780-19.
https://doi.org/10.1128/mra.00780-19

Suzuki S, Hirakawa Y, Kofuji R, et al (2016) Plastid genome sequences of *Gymnochlora stellata*,
Lotharella vacuolata, and *Partenskyella glossopodia* reveal remarkable structural
conservation among chlorarachniophyte species. J Plant Res 129:581–590.
https://doi.org/10.1007/s10265-016-0804-5

Talavera G, Castresana J (2007) Improvement of phylogenies after removing divergent and
ambiguously aligned blocks from protein sequence alignments. Syst Biol 56:564–577.
https://doi.org/10.1080/10635150701472164

Tanaka A, Ito H, Tanaka R, et al (1998) Chlorophyll *a* oxygenase (CAO) is involved in chlorophyll *b*
formation from chlorophyll *a*. Proceedings of the National Academy of Sciences of the
United States of America 95:12719–12723. https://doi.org/10.1073/pnas.95.21.12719

Tanaka A, Tanaka R (2019) The biochemistry, physiology, and evolution of the chlorophyll cycle, 1st
edn. Elsevier Ltd.

Tanaka R, Tanaka A (2007) Tetrapyrrole biosynthesis in higher plants. Annual Review of Plant
Biology 58:321–346. https://doi.org/10.1146/annurev.arplant.57.032905.105448

Tanaka R, Tanaka A (2005) Effects of chlorophyllide *a* oxygenase overexpression on light
acclimation in *Arabidopsis thaliana*. Photosynth Res 85:327–340.
https://doi.org/10.1007/s11120-005-6807-z

Tanaka R, Tanaka A (2011) Chlorophyll cycle regulates the construction and destruction of the
light-harvesting complexes. Biochimica et Biophysica Acta (BBA) - Bioenergetics 1807:968–
976. https://doi.org/10.1016/j.bbabio.2011.01.002

Tartar A, Boucias DG, Becnel JJ, Adams BJ (2003) Comparison of plastid 16S rRNA (rrn16) genes
from *Helicosporidium* spp.: evidence supporting the reclassification of Helicosporidia as
green algae (Chlorophyta). Int J Syst Evol Microbiol 53:1719–1723.
https://doi.org/10.1099/ijs.0.02559-0

Tomitani A, Okada K, Miyashita H, et al (1999) Chlorophyll *b* and phycobilins in the common
ancestor of cyanobacteria and chloroplasts. Nature 400:159–162.
https://doi.org/10.1038/22101

Turner-Hissong SD, Mabry ME, Beissinger TM, et al (2020) Evolutionary insights into plant breeding.
Current Opinion in Plant Biology 54:93–100. https://doi.org/10.1016/j.pbi.2020.03.003

Ueno Y, Aikawa S, Kondo A, Akimoto S (2019) Adaptation of light-harvesting functions of unicellular
green algae to different light qualities. Photosynthesis Research 139:145–154.
https://doi.org/10.1007/s11120-018-0523-y

Voitsekhovskaja OV, Tyutereva EV (2015) Chlorophyll *b* in angiosperms: Functions in
photosynthesis, signaling and ontogenetic regulation. Journal of Plant Physiology 189:51–
64. https://doi.org/10.1016/j.jplph.2015.09.013

Willows RD (2020) Biosynthesis of Chlorophyll and Bilins in Algae. In: Larkum AWD, Grossman AR,
Raven JA (eds) Advances in Photosynthesis and Respiration, 45th edn. Springer Nature, pp
83–103

Worden AZ, Lee J-H, Mock T, et al (2009) Green Evolution and Dynamic Adaptations Revealed by
Genomes of the Marine Picoeukaryotes Micromonas. Science 324:268–272.
https://doi.org/10.1126/science.1167222

Xiong J, Bauer CE (2002) COMPLEX EVOLUTION OF PHOTOSYNTHESIS. Annual Review of Plant
Biology 53:503–521. https://doi.org/10.1146/annurev.arplant.53.100301.135212

Xiong J, Wen G, Song J, et al (2024) Knockout of the Chlorophyll *a* Oxygenase Gene OsCAO1
Reduces Chilling Tolerance in Rice Seedlings. Genes 15:721.
https://doi.org/10.3390/genes15060721

Xu H, Vavilin D, Vermaas W (2001) Chlorophyll *b* can serve as the major pigment in functional
photosystem II complexes of cyanobacteria. Proceedings of the National Academy of
Sciences of the United States of America 98:14168–14173.
https://doi.org/10.1073/pnas.251530298

Yamasato A, Nagata N, Tanaka R, Tanaka A (2005) The N-Terminal Domain of Chlorophyllide a
Oxygenase Confers Protein Instability in Response to Chlorophyll *b* Accumulation in
Arabidopsis. Plant Cell 17:1585–1597. https://doi.org/10.1105/tpc.105.031518

Zhang L, Yang C, Liu C (2023) Revealing the significance of chlorophyll *b* in the moss
Physcomitrium patens by knocking out two functional chlorophyllide a oxygenase.
Photosynthesis Research 158:171–180. https://doi.org/10.1007/s11120-023-01044-8