Chlorophyllide *a* oxygenase (CAO) gene duplication across the Viridiplantae

Mackenzie C. Poirier[1], Roberta Wright[1], Marina Cvetkovska[1]*

[1]Department of Biology, University of Ottawa, Ottawa, ON, Canada

*Corresponding author:

    Marina Cvetkovska
Department of Biology, University of Ottawa, 30 Marie-Curie Pr., Ottawa, ON, Canada, K1N 6N5
Email: mcvetkov@uottawa.ca
ORCID: 0000-0002-7080-8203

Statements and Declarations

Competing Interests: The authors report that there are no competing interests to declare.

Acknowledgements: This project was supported by Natural Sciences and Engineering Research Council of Canada Discovery Grants (NSERC DG) awarded to M.C. The authors are grateful for the support from the Canada Foundation for Innovation (CFI) and University of Ottawa start-up funding. M.P. was supported by Ontario Graduate Scholarship (OGS), NSERC Graduate Scholarship, and Polar Knowledge Canada Antarctic Doctoral Scholarship.

Author's contributions: M. Poirier and M. Cvetkovska conceptualized the work and designed the experiments. Material preparation, data collection and analysis were performed by M. Poirier, R. Wright and M. Cvetkovska. The first draft of the manuscript was written by M. Poirier and all authors commented on all versions of the manuscript. All authors read and approved the final manuscript.

Abstract

Viridiplantae, a diverse group of green plants and alga that have evolved from a common ancestor, are unified in their ability to produce and use two types of chlorophyll (chlorophyll *a* and chlorophyll *b*) to capture light energy. In addition to playing a role in light harvesting, chlorophyll *b* is required at the appropriate level for the accumulation, assembly, and stability of light harvesting complexes within the photosynthetic apparatus. Chlorophyll *b* is synthesized from chlorophyll *a* by the enzyme chlorophyllide *a* oxygenase (CAO), a Rieske-type mononuclear non-heme iron oxygenase. A regulatory degron sequence, described in detail only in land plants, regulates the stability of CAO proteins based on the availability of chlorophyll *b*. Recent identification of CAO gene duplication in bryophyte and green algal species, combined with expanded availability of sequenced genomes within the Viridiplantae, prompted further investigation into the role of gene duplication in the evolution of chlorophyll *b* biosynthesis. Examination of genomes from 246 plant and algae species revealed independently occurring CAO duplications throughout the Viridiplantae, with a higher prevalence of duplication in land plants compared to their algal relatives. Additionally, we demonstrate that the degron sequence is poorly conserved in chlorophytes, but first appears as a conserved sequence in charophytes, and is very highly conserved among the embryophytes. The evolutionary history and functional role of CAO throughout the Viridiplantae lineage is discussed based on these key observations, adding to our understanding of chlorophyll *b* biosynthesis and the role of CAO in photosynthetic species.

Keywords

Chlorophyll biosynthesis, chlorophyllide *a* oxygenase, gene duplication, degron, plants and algae

Introduction

Photosynthetic organisms form the base of most food chains and provide the organic carbon compounds that support life in our biosphere. The capture of light energy by chlorophyll is the first step in the process of photosynthesis that fixes atmospheric CO_2 into stable organic compounds. Overwhelming evidence indicates that oxygenic eukaryotic photosynthesis that occurs in the chloroplasts of modern-day plants and algae traces its origins from an endosymbiotic event with cyanobacteria-like prokaryotes (Blankenship 2010; Cardona 2019; Sánchez-Baracaldo and Cardona 2020). Viridiplantae (or green plants) have primary chloroplasts derived from an ancient endosymbiosis, and encompass two major clades: the chlorophytes and the streptophytes. Chlorophytes are a monophyletic group of marine, freshwater, and terrestrial green algae. With ~8,000 described species (Guiry 2024), this group encompasses a large diversity of adaptations, morphologies, and life histories (Leliaert et al. 2012). Streptophyta includes the charophytes, ~5,500 species of largely freshwater green algae (Guiry 2024) and the embryophytes, or land plants. Land plants are thought to have diverged from a single clade within the streptophytes (de Vries and Archibald 2018) and number ~430,000 species (Ruggiero et al. 2015). One characteristic that unifies members of the Viridiplantae is their ability to synthesize and use two types of chlorophylls: chlorophyll *a* (Chl *a*) and chlorophyll *b* (Chl *b*).

The photosynthetic apparatus in the Viridiplatae is derived from the ancestral cyanobacterial endosymbiont, which used chlorophyll as the main pigment in their photosystems (Xiong and Bauer 2002; Sánchez-Baracaldo and Cardona 2020). Chl *a* is ubiquitously present in all oxygenic photosynthetic eukaryotes, and functions in both energy capture in the antenna light harvesting complexes (LHCs) and in driving electron transfer in the photosystem II (PSII) and photosystem I (PSI) reaction centers. Chl *b* is specific to the Viridiplantae although it has also been detected in prochlorophytes and *Acaryochloris*, unique photosynthetic prokaryotic groups that lack typical cyanobacterial phycobilin light-harvesting pigments (Palenik and Haselkorn 1992; Roche et al. 1996; Tomitani et al. 1999; Partensky et al. 2018). Chl *b* resides predominantly in the peripheral LHCs (Neilson and Durnford 2010) although it has been detected in the core complexes of some deep water marine chlorophytes (Kunugi et al. 2016).

Strict Chl *a/b* stoichiometry is required for optimal energy transfer during photosynthesis. Maintaining the correct Chl *a/b* ratios is a dynamic process and indicative of adaptation to different light conditions (Tanaka and Tanaka 2007). In addition to playing a role in light harvesting, Chl *b* is required at appropriate levels for the accumulation, assembly, and stability of LHCs in algae (Bujaldon et al. 2017), bryophytes (Zhang et al. 2023), and angiosperms (Król et al. 1995; Reinbothe et al. 2006; Kim et al. 2009; Nick et al. 2013). Under low light conditions, the amount of Chl *b* relative to Chl *a* increases, leading to larger LHC antenna size, which maximizes the surface area for light absorption (Kunugi et al. 2016; Kume et al. 2018; Ueno et al. 2019). Thus, the amount of Chl *b* is a major regulator of light-harvesting capacity.

Chlorophyll biosynthesis and turnover occurs through a complex multistep pathway (reviewed in Brzezowski et al. 2015; Willows 2020). Chl *b* is synthesized from Chl *a*, via the intermediate 7-hydromethyl chlorophyll *a*, by the action of a single enzyme: chlorophyllide *a* oxygenase (CAO) (Tanaka et al. 1998; Espineda et al. 1999; Oster et al. 2000; Mueller et al. 2012). Chl *b* can be converted back into Chl *a* by chlorophyll *b* reductase (CBR) and 7-hydroxymethyl reductase (HCAR)

to complete the cycle (Figure 1; reviewed in detail in Tanaka and Tanaka, 2019). The chlorophyll
cycle plays an essential role in many biological processes, including biogenesis of LHCs, antenna
size regulation during light acclimation, and chlorophyll degradation during senescence. While
many of the intricacies of how this pathway is regulated are still not fully understood (Tanaka and
Tanaka 2019), it has been suggested that CAO levels and activity are key determinants of Chl *b*
accumulation.

Figure 1. The chlorophyll cycle in land plants. Chlorophyll *a* is converted to chlorophyll *b* via the
intermediate 7-hydromethyl chlorophyll *a*, by the action of the enzyme chlorophyllide *a* oxygenase
(CAO). In a negative feedback loop, high levels of chlorophyll *b* trigger the binding of a stromal Clp
protease to the degron motif (QDLLTIMILH) and the subsequent destabilization of CAO (dashed red
line). Chlorophyll *b* is necessary for the stabilization of Light Harvesting Complexes (LHC) in plants
(dashed blue line). Excessive amounts of LHC or accumulation of energetically uncoupled non-
functional LHCs (e.g., during senescence), triggers the accumulation of Chlorophyll *b* reductase
(CBR; blue line) and the conversion of chlorophyll *b* to chlorophyll *a* via the action of CBR and 7-
hydroxymethyl reductase (HCAR). This regulatory mechanism of CAO and CBR enables the fine-
tuning and optimization of chlorophyll *b* and LHC levels (adapted from Tanaka and Tanaka 2019).

CAO is part of the Rieske-type mononuclear non-heme iron oxygenase group of proteins (Gray et al.
2004). It catalyzes the conversion of the methyl group of Chl *a* into a formyl group to form Chl *b*
through two oxygenation reactions, likely using ferredoxin as a reductant (Oster et al. 2000). CAO
has three main regions: an N-terminal regulatory domain and a C-terminal catalytic domain, which
are connected by a more variable linker region (Nagata et al. 2004). The catalytic domain contains a

Rieske cluster and a mononuclear iron-binding domain, both of which are necessary for the
enzymatic activity of CAO (Yamasato et al. 2005; Kunugi et al. 2013). The regulatory domain has
been characterized in detail only in land plants. In *Arabidopsis*, a conserved sequence
(QDLLTIMILH) termed a degron was demonstrated to regulate Chl *b* accumulation by affecting CAO
protein turnover. In a negative feedback loop, high levels of Chl *b* trigger the binding of a stromal
Clp protease to the degron motif leading to destabilization of the CAO protein; this, in turn, halts
Chl *b* biosynthesis (Sakuraba et al., 2009; Nakagawara et al., 2007).

The evolutionary origins of CAO can be traced back to prochlorophytes, which encode a CAO gene
homolog (Tomitani et al. 1999; Nagata et al. 2004). Notably, the prokaryotic CAO-like enzymes lack
the regulatory N-terminal domain, suggesting that the Chl *b*-dependent mechanism for controlling
CAO stability was acquired in eukaryotes. Recent work on 9 chlorophyte, 2 charophyte, and 3
embryophyte species suggested that the degron recognition sequence first appeared in
charophytes as an adaptation to high-light environments in shallow water (Kunugi et al. 2016).
These conclusions, however, were based on a very limited sample of representative species and
the presence of a degron in the CAO sequence in charophytes was inferred based on a single
species (*Klebsormidinium flaccidum*). This hypothesis requires further validation.

The CAO protein is encoded by a single gene in most plants and algae (Kunugi et al. 2016;
Schumacher et al. 2022), with only a very few reported exceptions. Two CAO genes were detected
in rice (Lee et al. 2005) although only one of them is necessary for Chl *b* synthesis (Jung et al. 2021).
Recent work revealed CAO duplication in the bryophyte *Physcomitrium patens,* where both genes
contribute to Chl *b* accumulation (Zhang et al. 2023). CAO duplication was also reported in two
Antarctic *Chlamydomonas* species , where the CAO paralogs in each species appear to be derived
from independent duplications events rather than an ancestral one (Cvetkovska et al. 2019; Poirier
et al. 2025).

These insights prompted us to further examine the *CAO* gene content and the presence of the
regulatory domain across the diverse plant and algal groups. A systematic examination for the
presence of *CAO* across the Viridiplantae has not yet been preformed. In this work we took
advantage of the recent increase in publicly available plant and algal genomes to examine the
occurrence of CAO genes in diverse species across Viridiplantae and demonstrate a widespread
occurrence of CAO duplicates in this group. Furthermore, building on the work by Kunugi et al.
(2016), we have examined the occurrence of a degron sequence in 326 unique CAO sequences in
algae and plants. We demonstrate that this regulatory region is poorly conserved in chlorophytes,
compared to charophyte and embryophyte CAO sequences, suggesting a different mechanism for
Chl *b* regulation in early aquatic lineages.

Materials and Methods

CAO sequence identification and analyses

To identify *CAO* sequences and duplication events with high accuracy, we used publicly available,
high quality, and fully annotated genomes. Chlorophyte and streptophyte genomes were retrieved
from Phycocosm (Grigoriev et al. 2021), while embryophyte genomes were obtained from
Phytozome v13 (Goodstein et al. 2012). Putative *CAO* genes were identified with a tBLASTn search

with the *Chlamydomonas reinhardtii* (CrCAO; Cre01.g043350) and the *Arabidopsis thaliana*
(AtCAO; AT1G44446) full-length CAO peptides (Merchant et al. 2007; Lamesch et al. 2012). A strict
e-value cutoff of ≤e-30 was taken as a threshold, to avoid many unspecific hits due to the
conserved nature of the Fe-binding and Rieske cluster domains present among diverse plant and
algal proteins (Schmidt and Shaw 2001; Ferraro et al. 2005; Przybyla-Toscano et al. 2021). Putative
CAO sequences were manually curated and truncated or low-quality sequences were excluded
from downstream analyses. Curated *CAO* sequences were aligned using MUSCLE (Edgar 2004).
The degron sequence (QDLLTUMILH) was identified by multiple sequence alignment of the full-
length CAO amino acid sequences and the regulatory N-terminal domain. In accordance with
Sakuraba et al. 2009, we identify the degron as "highly conserved" if 8 out of the 10 amino acids are
perfectly conserved, "moderately conserved" if 4 to 8 amino acids are conserved, and "poorly
conserved" if less than 4 amino acids are conserved. Sequence logos for each major group
(chlorophyte algae, charophyte algae, non-vascular plants and vascular plants) were generated by
Geneious Prime 2024.11 (Dotmatics), where the size of the letter reflects its frequency. To examine
the expression of CAO duplicate genes, we screened previously published transcriptomes from 16
species with representatives from the Chlorophytes (Arriola et al. 2018), Charophytes (Cheng et al.
2019), non-vascular (Perroud et al. 2018; Healey et al. 2023) and vascular plants (Zhu et al. 2019;
Wang et al. 2021; Liu et al. 2022b; Shang et al. 2023; Guo et al. 2025; He et al. 2025; Mascuñano et
al. 2025; Roy et al. 2025). Only transcriptomes that report FPKM, RPKM and TPM values obtained
from organisms cultivated at optimal conditions were considered (Supplementary Table S4).

Phylogenetic Inference

The analyses described above resulted in 374 individual CAO sequences (Supplementary Table
S1). To avoid inaccurate inference and over-representation, we manually removed heterodimeric
CAO proteins, where the Rieske domain and the Fe-binding domain are located on different genes
(Kunugi et al. 2013) (Supplementary Table S1). We also removed identical CAO sequences isolated
from algal cultures identified as the same species but belonging to different strains or culture
collections (*Scenedesmus obliquus*, *Auxenochlorella protothecoides*, *Chlorella sorokiniana*,
Ostreococcus tauri, *Mesostigma viride*, *Zygnema circumcarinatum*). This resulted in a 280 amino
acid alignment of 326 unique CAO sequences across 246 species.

These manually curated CAO genes were translated into amino acid sequences, aligned with
MUSCLE and trimmed to remove gaps and ambiguously aligned regions with Gblock (Castresana
2000; Talavera and Castresana 2007). Maximum likelihood trees were inferred in the CIPRES
Science Gateway (Miller et al. 2015) using RAxML v8.0 (Stamatakis 2014) with 1000 bootstraps with
the Whelan Goldman matrix for globular proteins (WAG), a gamma shape parameter, and empirical
estimation of invariable sites. The trees were rooted using prokaryotic CAO from *Acaryochloris*
thomasi (WP_110987895.1) and *Prochlorotrix hollandica* (WP_017713323.1) as an outgroup.
Tandem duplications were defined as those located on the same chromosome or contig within <10
genes of each other, based on the genomic coordinates retrieved from Phytozome v13 or
Phycocosm. To identify ancestral and lineage-specific duplications, we used Notung v2.9 (Stolzer
et al. 2012) and reconciled the CAO trees with a reference species tree derived from NCBI
taxonomy (Schoch et al. 2020) via the PhyloT v2 tool (https://phylot.biobyte.de). Putative ancestral
duplications were only reported when occurring on branches with bootstrap support ≥85%. All
trees were visualized in iTOL v6.8.2 (Letunic and Bork 2021). Previously inferred taxonomic

information for chlorophytes and streptophytes was confirmed via AlgaeBase, University of Galway
(www.algaebase.org), and for land plants via the Plants Of the World Online, Royal Botanic
Gardens, Kew (https://powo.science.kew.org).

Results and Discussion

Occurrences of multiple CAO genes are widespread among the Viridiplantae

Our analysis demonstrates a widespread occurrence of *CAO* gene duplication across the
Viridiplantae. Multiple gene copies are more prevalent among the angiosperms, where ~41% of
examined species encode for more than one *CAO* gene. In comparison, we detected two *CAO* gene
copies in only ~11% of chlorophyte and ~20% of charophyte species (Table 1; Table 2,
Supplementary Table S1; Supplementary Table S3). All 23 species where we detect more than two
CAO genes are members of the Embryophyta. The modern-day cultivar of sugarcane (*Saccharum*
sp. R570) encodes for 6 *CAO* copies, the largest number we report. This species is a result of inter-
specific hybridization (*Saccharum officinarum* x *spontaneum;* Dumont et al. 2022), which has
resulted in a very large and highly redundant genome (~0.3 Gb, ~48,000 genes) (Healey et al. 2024).
It must be noted that these results are likely affected by the availability of many more angiosperm
genomes in public databases, while other plant and algal groups are not as well represented. As
more genomes are sequenced, these analyses will have to be revisited.

We also detected multiple occurrences of heterodimeric *CAO* genes, where the functional CAO
protein is encoded by two genes (Table 1, Supplementary Table S1). Previous work in the
prasinophyte *Micromonas pusila* demonstrated that in this species CAO was encoded by two
separate genes, one that possesses the mononuclear iron-binding motif (*MpCAO1*) and a second
one that encodes for the Rieske cluster (*MpCAO2*). It was shown that a combination of both gene
products was necessary to form an active heterodimeric CAO protein (Kunugi et al. 2013; Dey et al.
2023). We expand on this work by showing the presence of a heterodimeric CAO encoded by two
separate genes across the Mammiellophycea. This group within the prasinophytes contains *M.*
pusila and several of its relatives (Worden et al. 2009), the ecologically important picoalgae
Ostreococcus tauri, *Ostreococcus lucimarinus* and *Bathycoccus prasinos* (Palenik et al. 2007;
Moreau et al. 2012; Blanc-Mathieu et al. 2014)*,* as well as the related *Chloropicon primus*
(Chloropicophyceae) (Lemieux et al. 2019).

We also show that heterodimeric CAO is not restricted only within the Prasinophytes. We detect a
similar *CAO* gene arrangement in the genome of the multinucleate green alga *Caulerpa lentillifera*
(Bryopsidales) (Arimoto et al. 2019) , as well as in in the genome of *Bigelowiella natans*
(Chlorarachniophyceae) (Curtis et al. 2012). Phylogenetic analysis based on plastid-encoded
protein sequences revealed that chlorarachniophytes acquired their secondary plastids by the
uptake of a filamentous green alga in the Bryopsidales (Suzuki et al. 2016). Thus, *B. natans* likely
gained a heterodimeric CAO from the Bryopsidales endophyte. Considering that CAO in all other
green lineages (Supplementary Table 1) and within the prochlorophytes (Nagata et al. 2004) is also
encoded by a single gene that contains both the Rieske and iron-binding domain it is likely that the
occurrence of two separate *CAO* genes is not an ancestral condition. Instead, it has been
postulated that CAO was encoded by a single gene in the common ancestor of green plants
(Kunugi et al. 2016). Indeed, it has been suggested that acquisition of two separate genes encoding
for key functional domains may not be an evolutionarily difficult process. CAO likely functions as a

trimer (Kunugi et al. 2013; Liu et al. 2022a), and the electron transfer from the Rieske to the
mononuclear iron occurs between two neighbouring CAO proteins. A functionally similar process
likely occurs in the CAO enzyme encoded by two separate genes.

Table 1: Number of species across the Viridiplantae with fully sequenced and annotated genomes
available through Phytozome (Total), compared to the number of species that encode for multiple
CAO genes (Multiple CAO) and number of species that encode for two *CAO* genes that form
heterodimeric complex (Heterodimeric CAO).
260

Group	Total	Heterodimeric CAO	Multiple CAO
Chlorophytes*	70	11	8
Charophytes	10	0	2
Non-vascular plants	6	0	3
Lycophytes	3	0	2
Gymnosperms	1	0	0
Angiosperms	151	0	62

*The Chlorophyte group also includes *Bigelowiella natans* (Chlorarachniophyceae) that has gained
secondary plastids by the uptake of a green alga.

All examined streptophytes encode for at least one *CAO* gene (Supplementary Table 1) signifying
the importance of Chl *b* for plant physiology, but we were not able to identify a homolog of *CAO* in
several chlorophyte algal species. Not surprisingly, we did not detect a *CAO* homolog in the
invertebrate parasite *Helicosporidium* sp., a colorless protist that belongs within the
trebouxiophycean green algal clade (Tartar et al. 2003). As a consequence of its parasitic lifestyle,
Helicosporidium has lost nearly all genes associated with light harvesting, photosynthesis and
chlorophyll biogenesis (Pombert et al. 2014), including CAO.

We were also not able to conclusively identify a *CAO* gene in the genomes of the endolithic coral
holobiont *Ostreobium quekettii* (Bryopsidales) (Iha et al. 2021), the aquaculture species
Tetraselmis striata (Chlorodendrophyceae) (Steadman Tyler et al. 2019) and the halotolerant
Picocystis sp. ML (Picocystophyceae) (Junkins et al. 2019). All identified sequences within the
genomes of these species had e-values higher than the cut-off (≥e-15) used in this work and high
similarity to pheophorbide *a* oxygenase (PAO), a Rieske-type protein involved in chlorophyll
degradation (Pružinská et al. 2003). The lack of *CAO* genes in these algae is surprising and could be
an artifact of genome assembly, as Chl *b* has been detected in all three species (Roesler et al.
2002; Massé et al. 2020; Conlon et al. 2024). The genome of *O. queketti* has a low BUSCO score
(60.7%) (Iha et al. 2021) and the BUSCO score is not reported for the draft genomes of *T. striata* and
P. sp. ML (Junkins et al. 2019; Steadman Tyler et al. 2019). The presence of CAO should be
experimentally examined in these species.

It is tempting to speculate that these species may have a different mechanism for Chl *b*
biosynthesis or a CAO sequence significantly different than most algae and plants. For instance,
Ostreobium is an endolithic coral holobiont adapted to extreme shading, and has lost many genes
associated with photoprotection and light perception due to evolution in a light-limited
environment (Iha et al. 2021). This alga is also the only known eukaryote that has lost the gene
encoding light-dependent protochlorophyllide oxidoreductase (LPOR), a key enzyme that catalyzes

the conversion of protochlorophyllide to chlorophyllide *a*, a necessary step towards the synthesis
of Chl *a* (Reinbothe et al. 2010). Instead, this alga fully depends on dark operative POR (DPOR) to
synthesize Chl *a*, likely as an adaptation to extreme shading (Iha et al. 2021). The mechanism of
chlorophyll biosynthesis has not been examined in the species in detail, but it is clear that the
endolithic lifestyle has resulted in a unique physiology. Further work will shed light on the role of
CAO in this and other shade-tolerant species.

We used a very conservative method to identify putative CAO sequences with both a Rieske and
Fe-binding domains, but this strict approach may have prevented the identification of duplicates
that may have diverged or become pseudogenes. Indeed, using a less conservative cutoff (≤e-5)
when screening the genomes of *A. thaliana* and *C. reinhardtii* (Wang et al. 2022; Craig et al. 2022)
reveals that *PAO* and Translocon at the inner envelope membrane of chloroplasts 55 (*TIC55*) may
be paralogous to *CAO* (Supplementary Table S2). *PAO* and *TIC55* encode for enzymes with a Rieske
and Fe-binding domains but share a very low sequence similarity with CAO and each other
(Supplementary Table S2, Supplementary Figure S1). Both PAO and TIC55 are involved in
chlorophyll breakdown and allow for the degradation of the highly phototoxic pheophorbide *a* and
its export from the chloroplast (recently reviewed in Kuai et al. 2018). Evolutionary analysis of the
genes involved in chlorophyll degradation revealed that this pathway was already present in the
common ancestor of land plants, suggesting that *PAO* and *TIC55* may share their origins with *CAO*
(Schumacher et al. 2022). This hypothesis has not yet been experimentally supported as ancestral
monooxygenase have been difficult to distinguish without detailed knowledge on their biochemical
activity and function (Schumacher et al. 2022). This possibility raises interesting questions about
the ancient evolutionary origins of chlorophyll metabolism within the Viridiplantae.

310 *Was CAO gene duplication an ancestral event?*

To determine whether the *CAO* gene duplication was an ancestral event, we performed a
phylogenetic analysis of all high-quality CAO amino acid sequences (except heterodimeric CAO).
Phylogenetic patterns typically match previously shown relationships among major Viridiplantae
groups (Figure 2), although some relationships are poorly resolved (particularly among the algal
groups and the Rosids; Supplementary Figure 2). The presence of two *CAO* genes has been
previously reported in rice (*OsCAO1* and *OsCAO2*) where the two genes were positioned in tandem
on Chromosome 10, taken as evidence for a recent gene duplication in this species (Lee et al.
2005). A tandem CAO duplication, however, appears to be the exception rather than the rule. In
addition to rice, we only observed *CAO* duplicates located in proximity to each other in the
genomes of a handful of species: the green alga *Volvox reticuliferus*, the lycophyte *Diphastrium*
complanatum, and the magnoliid *Cinnamomum kanehirae* (Table 2).

There are several instances, mainly in very closely related species, where the duplication may have
originated in a recent shared ancestor. There are several possible cases in the Trebouxiophyceae
and Chlorophyceae class, where *CAO* duplicates are detected in closely related *Chlorella* and
Scenedesmus species (Supplementary Figure 2A). This insight must be interpreted with caution as
phylogenetic resolution within the chlorophytes is still weak due to limited gene and taxon
sampling (Li et al. 2021). Within the bryophytes, *Spaghnum magellanicum* and *Sphagnum falax*,
species that share a very close evolutionary relationship (Piatkowski and Shaw 2019; Bell et al.
2020), both have two *CAO* genes possibly arising from an ancestral duplication. Two CAO genes

are also encoded in the genome of *P. patens* (Zhang et al. 2023). This raises the possibility of a CAO duplication in a recent shared ancestor (Supplementary Figure 2B), but the bryophytes are represented by only 5 species in our analysis. Additional genomic sequencing and a deeper phylogenetic analysis will be needed to clarify the evolution of CAO within these groups.

Figure 2: A phylogenetic tree based on CAO amino acid sequences inferred from using maximum likelihood analysis, using 326 unique CAO sequences across 246 species color-coded to represent the major Viridiplantae groups. Scale bar indicates amino acid substitutions per site. Species with more than one detected CAO gene are highlighted in red

Species that encode for multiple *CAO* genes within their genomes are more prevalent among
angiosperms. Within the monocots, *CAO* duplicates arising from a shared ancestral gene are likely
present in the Asparagaceae, specifically among the closely related *Agave tequiliana*, *Yucca*
filamentosa and *Yucca aloifolia* (Ji et al. 2023) , although we were only able to detect a single *CAO*
copy in *Asparagus officinalis* (Supplementary Figure 2C). We detected two *CAO* gene copies in two
Brachypodium species (*B. mexicanum* and *B. hybridum*) but not in their relatives from the same
genus (*B. arbuscula*, *B. stacei*, *B. sylvaticum* and *B. distachyon*), raising the possibility of an
ancestral duplication followed by gene loss. All phylogenetic studies in *Brachypodium* support a
rapid and recent divergence from a common ancestor (Catalan et al. 2016), supporting this
possibility.

CAO duplication may be ancestral in the two closely related Saxifragales *Kalanchoe laxiflora* and
Kalanchoe fedtschenkoi (Han et al. 2024). We detected two *CAO* genes in *K. fedtschenkoi* and four
in *K. laxiflora*, possibly arising in a recent ancestor, followed by a second duplication within *K.*
laxiflora (Supplementary Figure 2D). Within the Fabids, four *Populus* species and their relative *Salix*
purpurea, each encode for two *CAO* gene copies that may be a result of an ancestral duplication.
The cultivated soybean *Glycine max* and its wild relative *Glycine soja* each encode for four *CAO*
gene copies (Supplementary Figure 2D) that are closely related to their homologs between the two
species. The phylogenetic separation of *CAO1/2* and *CAO3/4* from *G. max* and *G. soja* is puzzling,
as the two species are the closest relatives within the *Glycine* genus (Zhuang et al. 2022). All
Gossypium species examined here encode for four *CAO* genes, except *G. raimondii* which encodes
for two copies (Supplementary Figure 2E). The only other member of the Malvaceae examined,
Theobroma cacao, encodes for a single *CAO* gene, and although it clusters with *CAO* from
Gossypium sp., the relationship is poorly supported. Finally, there is a possibility of ancestral
duplications within the Brassicaceae (Supplementary Figure 2E), but this family has a poor
resolution due to factors such as incomplete sampling, rapid radiation, and hybridization (Huang et
al. 2016; Hendriks et al. 2023).

The genomes of multiple species across the Viridiplantae encode for more than one *CAO* gene, but
are these gene duplications related to plant evolutionary patterns? The green lineage is
characterized by a dynamic and complex evolutionary history marked by multiple whole genome
(WGD) and segmental duplication events, gene family expansions, and lineage-specific gene
losses (Panchy et al. 2016; Kersey 2019; Leebens-Mack et al. 2019; Soltis and Soltis 2021). Such
tremendous diversity in genome content can be a challenge for elucidating *CAO* evolutionary
trajectories even withing a single family. For instance, the Brassicaceae have been shaped by
multiple well-established WGD events (Mabry et al. 2020), but we detected only a single *CAO* gene
in the genome of *A. thaliana* and its closest relatives. In contrast, the genomes of several other
Brassica species encode for two or more CAO genes (Figure 2). This highlights the importance of
considering not only gene duplication but also gene retention dynamics when interpreting the
evolution of CAO. Future work that incorporates in-depth analyses, such as relative dating of CAO
duplication events and genome-wide synteny, will further clarify the evolutionary trajectory of CAO
within the Viridiplantae.

We show that *CAO* gene duplication is not a rarity among the Viridiplantae, but are all *CAO*
duplicates expressed and do they encode for active proteins? In our work, we focused only on CAO
genes with highly conserved catalytic domains likely to encode for active enzymes, but the

functional role of CAO duplication may be complex. A survey of published transcriptomes from 16
representative species from all major groups (Chlorophytes, Charophytes, non-vascular and
vascular plants) suggests that all *CAO* duplicate genes within a species are expressed, albeit at
different levels. Typically, one *CAO* gene has a higher expression than its paralog within the same
genome (Supplementary Table 4) suggesting differential regulation of *CAO* expression.

Significantly, all experimental studies to date suggest that the expression of *CAO* duplicates may
be differently regulated by light availability. In rice, the expression of *OsCAO1* is light-inducible,
while *OsCAO2* accumulates during periods of darkness and is expressed mainly in non-
photosynthetic tissues (Lee et al. 2005). In *P. patens*, both *PpCAO1* and *PpCAO2* are expressed in
both light and dark conditions, but only *PpCAO1* is upregulated with prolonged light exposure.
Similarly, in two *Chlamydomonas* species that harbour *CAO* duplicate genes (*Chlamydomonas*
priscui and *Chlamydomonas* sp. ICE-MDV), only one *CAO* homolog is upregulated in response to
increases in light intensity, while the second one is constitutively expressed (Poirier et al. 2025).
The role of light in regulating *CAO* expression has been also demonstrated previously in
Arabidopsis and tobacco, species with a single CAO gene copy (Pattanayak et al. 2005; Tanaka and
Tanaka 2005; Biswal et al. 2012, 2024).

Additionally, the mechanisms behind the control of Chl *b* synthesis may be species specific.
Mutational analysis in rice revealed that only *OsCAO1* was necessary for Chl *b* biosynthesis.
OsCAO1 knockout mutants exhibited low Chl *b* levels, disordered thylakoid membrane
arrangements, membrane damage due to excessive ROS accumulation and decreased chilling
tolerance (Lee et al. 2005; Jung et al. 2021; Xiong et al. 2024), but these effects were absent in
OsCAO2 knockouts (Lee et al. 2005). In contrast, knocking out either of the two *CAO* homologs in
P. patens lead to significantly reduced Chl *b* levels suggesting a dependency on two active CAO
enzymes for the appropriate control of Chl *b* synthesis in this species (Zhang et al. 2023).

It is tempting to speculate that species that encode for multiple *CAO* genes have a more robust
capacity for Chl *b* biosynthesis through an increase in gene dosage or a finer control over Chl *b*
accumulation though gene subfunctionalization or differential expression (Kondrashov 2012;
Panchy et al. 2016). Two gene copies with different expression patterns or activity levels could
provide a fine-tuned control of Chl *b* levels and photosynthetic performance under varying
environmental conditions, but additional experimental work is needed to substantiate this
hypothesis.

Table 2: Distribution of *CAO* duplicate genes across Viridiplantae. Only genes with a confirmed
Riske conserved cluster, Fe-binding domain (all species), and degron sequence (Streptophytes
only) are listed. A checkmark (✓) indicates that the CAO genes are positioned in a tandem on the
same chromosome within the genome, while a cross (x) indicates the genes are located on
different chromosomes. Accession numbers can be found in Supplementary Table 1 (continued on
next page).

Taxon	Species	#CAO genes	Tandem
Chlorophyta			
Chlorophyceae	*Chlamydomonas priscui*	2	x
Chlorophyceae	*Volvox reticuilferus*	2	✓
Chlorophyceae	*Tetradesmus deserticola*	2	x
Trebouxiophyceae	*Chlorella sp. A99*	2	x
Trebouxiophyceae	*Chlorella sorokiniana*	2	x
Trebouxiophyceae	*Chlorellaceae sp.*	2	x
Trebouxiophyceae	*Micractinium conductrix*	2	x
Pedinophyceae	*Pedinomonas minor*	2	x
Charophyta			
Zygnematophyceae	*Mesotaenium endlicherianum*	2	x
Zygnematophyceae	*Spirogloea muscicola*	2	x
Bryophyta			
Bryopsida	*Physcomitrium patens*	2	x
Sphagnopsida	*Sphagnum fallax*	2	x
Sphagnopsida	*Sphagnum magellanicum*	3	x
Lycophyta			
Equisetopsida	*Diphasiastrum complanatum*	4	✓
Equisetopsida	*Ceratopteris richardii*	2	x
Angiosperms			
Laurales	*Cinnamomum kanehirae*	2	✓
Nymphaeales	*Nymphaea colorata*	2	x
Asparagales	*Agave tequilana*	2	x
Asparagales	*Yucca aloifolia*	3	x
Asparagales	*Yucca filamentosa*	3	x
Zingiberales	*Musa acuminata v1*	2	x
Alismatales	*Spirodela polyrhiza v2*	2	x
Poales	*Typha latifolia*	2	x
Poales	*Eleusine coracana*	2	x
Poales	*Oryza sativa*	2	✓
Poales	*Thinopyrum intermedium*	3	x
Poales	*Brachypodium hybridum*	2	x
Poales	*Brachypodium mexicanum*	2	x
Poales	*Zea mays*	2	x
Poales	*Andropogon gerardi*	3	x
Poales	*Miscanthus sinensis*	2	x
Poales	*Panicum virgatum*	2	x
Poales	*Saccharum officinarum*	6	x
Caryophyllales	*Chenopodium quinoa*	2	x
Gentianales	*Coffea arabica geisha*	2	x

Table 2:(continued)

Taxon	Species	#CAO genes	Tandem
Apiales	*Hydrocotyle leucocephala*	2	x
Boraginales	*Ehretia anacua*	2	x
Asterales	*Helianthus annuus*	2	x
Lamiales	*Olea europaea*	2	x
Saxifragales	*Kalanchoe fedtschenkoi*	2	x
Saxifragales	*Kalanchoe laxiflora*	4	x
Fabales	*Arachis hypogaea*	2	x
Fabales	*Chamaecrista fasciculata*	2	x
Fabales	*Glycine max*	4	x
Fabales	*Glycine soja*	4	x
Fabales	*Lupinus albus*	2	x
Fabales	*Phaseolus vulgaris*	2	x
Fagales	*Carya illinoinensis*	2	x
Rosales	*Fragaria x ananassa*	3	x
Rosales	*Malus domestica*	2	x
Malpighiales	*Linum usitatissimum*	4	x
Malpighiales	*Populus deltoides*	2	x
Malpighiales	*Populus nigra x maximowiczii*	2	x
Malpighiales	*Populus tremula x alba*	2	x
Malpighiales	*Populus trichocarpa*	2	x
Malpighiales	*Salix purpurea*	2	x
Sapindales	*Anacardium occidentale*	3	x
Malvales	*Gossypium barbadense*	4	x
Malvales	*Gossypium darwinii*	4	x
Malvales	*Gossypium hirsutum*	4	x
Malvales	*Gossypium mustelinum*	4	x
Malvales	*Gossypium raimondii*	2	x
Malvales	*Gossypium tomentosum*	4	x
Brassicales	*Alyssum linifolium*	2	x
Brassicales	*Caulanthus amplexicaulis*	2	x
Brassicales	*Iberis amara*	2	x
Brassicales	*Lepidium sativum*	2	x
Brassicales	*Stanleya pinnata*	2	x
Brassicales	*Brassica juncea*	3	x
Brassicales	*Brassica oleracea*	2	x
Brassicales	*Cakile maritima*	2	x
Brassicales	*Crambe hispanica*	2	x
Brassicales	*Eruca vesicaria*	2	x
Brassicales	*Sinapis alba*	3	x
Brassicales	*Camelina sativa*	3	x
Brassicales	*Isatis tinctoria*	4	x

Evolutionary implications of the CAO degron sequence

In addition to light-dependant regulation at the level of transcription, the main regulation of CAO
occurs post-translationally. Early work in *A. thaliana* demonstrated that this regulation is governed
by the presence of a degron sequence (QDLLTIMILH) in the protein's N-terminal A domain that acts
as a docking site for a chloroplast Clp protease (Yamasato et al. 2005; Nakagawara et al. 2007;
Sakuraba et al. 2009). It has been hypothesized that the degron sequence is "hidden" within the A
domain when there are insufficient amounts of Chl *b*, but accumulation of this product causes a
conformational change in the CAO protein, exposing the degron, and giving access to the protease
(Tanaka and Tanaka 2011). The evolution of this negative-feedback regulatory loop has not been
examined in detail, but it has been suggested that the degron was acquired early in the
Viridiplantae evolution as a mechanism for high light adaptation (Kunugi et al. 2016).

Here we show that the degron sequence is highly conserved among all land plants (86% identity),
moderately conserved among charophytes and non-vascular plants (68% and 66% identity
respectively), and very poorly conserved in chlorophytes (39% identity) (Figure 3; Supplementary
Figure 3). Land plants (both vascular and non-vascular) have several highly conserved sites across
all species (Asparagine at position 2, Isoleucine at position 6, Histidine at position 10), whereas
these positions are only moderately conserved in charophytes, and poorly conserved in
chlorophytes. In contrast, the Rieske cluster [CXH(X)15-17CXXH] exhibits an 84% pairwise identity
among all species, where the two histidine (positions 3 and 23) and cysteines (positions 1 and 20)
that coordinate the metallocentre [2Fe-2S] (Przybyla-Toscano et al. 2021) are present in all
examined sequences. Nearly all residues are conserved among the Viridiplantae in the C-terminal
Fe-binding domain (96% pairwise identity). Overall, these results suggest that the mechanism of
Chl *b* biosynthesis involving the Rieske and Fe-binding domains is ancestral and conserved among
all photosynthetic species, but this is not the case for the regulatory degron domain.

The lack of a highly conserved degron sequence may have several implications in the evolution of
efficient light harvesting. For instance, the N-regulatory domain (including the degron) is
completely absent in marine eukaryotic Mammiellophycea that encode for a heterodimeric *CAO*
gene, as well as in their deep-water relatives from the order Palmophyllales. These species
typically have low Chl *a/b* ratios and, unlike land plants, can incorporate Chl *b* into their core
antennae. This was interpreted as a necessary adaptation for efficient light harvesting in deep
marine environments characterized by low levels of blue-green light (Kunugi et al. 2016).

A

B

Figure 3: Key domains in the CAO protein across the Viridiplantae. (A) Sequence logos of the conserved motifs identified in the CAO protein sequences. N and C represent the N-terminus and C-terminus, respectively. All sequence logos are superimposed on a graphical representation of plant CAO sequences (not to scale) showing the regulatory degron motif, conserved Rieske domain cluster and Fe-binding domain (Sakuraba et al., 2009; Liu et al., 2022). The logos were generated for CAO proteins from chlorophyte algae, charophyte algae, non-vascular plants, and vascular plants. The sequences on top of each logo are the consensus sequence with the most common base in each sequence alignment. (B) Comparison of the CAO amino acid sequences corresponding to the degron and its surrounding region in the N-terminal regulatory domain in representative species from Chlorophyte, Charophyte, non-vascular and vascular land plants. The region that corresponds to the degron sequence as described in *A. thaliana* is emphasized with a box. Amino acids that are identical with the *A. thalina* sequence are colored, and those that are different are shown in white.

It has been suggested that the degron sequence evolved in shallow-water algae and land plants as
a means to regulate Chl *b* accumulation and limit Chl *b* incorporation into the reaction center in an
environment characterized with high light intensities (Kunugi et al. 2016). Indeed, such species
exclusively harbour Chl *a* in their core complexes, but this appears to be controlled by pigment
availability rather than the chemical nature of the two chlorophyll types. The binding affinity of Chl
a and *b* to the PSI complex is similar (Ikegami et al. 2007) and cyanobacteria that do not naturally
produce Chl *b* successfully incorporated this pigment into their reaction centers when transformed
with plant CAO (Satoh et al. 2001; Xu et al. 2001). Incorporation of Chl *b* into the reaction center
accompanied by increased Chl *b* accumulation was also observed in *Arabidopsis* when full length
CAO was overexpressed or replaced with a prokaryotic *CAO* that lacked a degron sequence
(Hirashima et al. 2006; Sakuraba et al. 2009).

The chlorophyll cycle has not been examined in detail in the core chlorophytes (Chlorophyceae,
Trebouxiophyceae, Ulvophyceae), a diverse group that inhabits marine, freshwater and terrestrial
habitats (Leliaert et al. 2012). Although many chlorophytes thrive in high light, others are adapted to
light limited environments, including deep waters. As a result, these species have developed
adaptations allowing for efficient light harvesting in low light. For instance, *Chlamydomonas priscui*
is endemic to the depths of the Antarctic Lake Bonney, an environment characterized by extreme
shading and dominated by blue-green wavelengths (Neale and Priscu 1995). This alga displays a
host of adaptations to extreme shading including a large LHC antennae size (reviewed in
Cvetkovska et al. 2017; Hüner et al. 2022). As suggested for marine prasinophytes (Kunugi et al.
2016), such low-light adapted chlorophytes could benefit from increased Chl *b* levels and could
thrive without a Chl *b*-dependent feedback mechanism.

Despite the lack of a conserved degron sequence to regulate CAO protein levels (Figure 3),
chlorophytes have maintained the ability to alter their Chl *a/b* ratio in response to light (Leliaert et
al. 2012). Transcriptional control of CAO levels was postulated to be a minor aspect of adjusting
Chl *b* levels in plants (Tanaka and Tanaka 2019), but this process could play a bigger role in
chlorophytes. It is also possible that a different, yet unidentified, mechanism exists for Chl *b*
turnover independent of a degron sequence, allowing these species to modify their Chl *a/b* ratios.
Finally, we can not rule out that the region corresponding to a degron in Chlorophyte CAO may still
have a similar function despite poor sequence conservation with land plants. The mechanism of
regulating protein degradation via degron motifs is well understood in plants (Isono et al. 2024), but
insights on algal degrons is lacking. A detailed experimental investigation on the control and
mechanism of Chl *b* biosynthesis is clearly needed among the Chlorophytes. For instance, targeted
mutagenesis of the N-terminus of algal CAO will reveal the importance and role of this regulatory
region in chlorophyll turnover.

Conclusion and Drawbacks

Our work suggests that *CAO* gene duplication is widespread among the Viridiplantae, likely
originating from multiple independent duplication events throughout plant evolution. While we
show that the presence of multiple *CAO* gene copies is highly prevalent in land plants, these
results must be interpreted with caution. Many more angiosperm genomes are available in public
databases, while other plant and algal groups are not as well represented. Thus, the conclusions of
our study must be re-visited once more genomes are sequenced. Furthermore, many genome

sequencing initiatives are focused on plants of economic and agricultural importance, in which
domestication and selective breeding may have affected the genomic content (Turner-Hissong et
al. 2020). While the exact physiological impact of CAO duplication is unknown, a study in
transgenic tobacco overexpressing CAO (Biswal et al. 2012) showed that an increase in Chl *b* may
be a strategy to enhance CO_2 assimilation, delay senescence, and enhance productivity. Such
traits would be attractive in crop breeding programs (Voitsekhovskaja and Tyutereva 2015), and the
high prevalence of CAO duplication seen in crop plants in this study may have been selected for
during plant breeding. Indeed, several crop plants (including corn, soybean, and olives, among
others) appear to express all CAO genes encoded in their genome (Supplementary Table 4). The
functional role of these duplicates in plant physiology, photosynthetic efficiency, and productivity
warrants further investigation.

Our understanding of the prevalence and evolution of CAO at the genetic level will be improved in
future years, as more algal and non-vascular plant genomes become available. Furthermore, there
are still numerous complex and outstanding questions regarding the control of Chl *b* accumulation
in photosynthetic species. In particular, the chlorophytes are an understudied group when it
comes to Chl *b* regulating mechanism. Detailed genomic and experimental studies will shed light
on the role and regulation of Chl *b* in these species and will deepen our understanding on this
crucial photosynthetic pigment.

Supplementary Information

Supplementary Table S1: A summary of genes encoding for chlorophyllide *a* oxygenase (*CAO*)
across the Viridiplantae used in this work. All sequences were obtained from Phycocosm
(Chlorophytes, Euglenophytes, Cercozoa, Charophytes) or Phytozome v13 (Bryophytes,
Marchantiophytes, Lycophytes, Gymnosperms, Angiosperms). Accession numbers represent the
unique IDs assigned to each gene (Phycocosm: Gene portal|Protein ID|Gene Model; Phytozome:
Gene Model). E-values are based on a tBLASTn search with the *Chlamydomonas reinhardtii* CAO
(Cre01.g043350) for species within the Chlorophyta and Charophyta, and the *Arabidopsis thaliana*
CAO (AT1G44446) for all other species

Supplementary Table S2: BLAST results obtained by screening the genome of *Arabidopsis thaliana*
and *Chlamydomonas reinhardtii* for CAO homologs. Accession numbers were obtained from
Phytozome v13 (Goodstein et al. 2012). The functional domains according to InterPro (Blum et al.
2025) that are shared among all proteins are highlighted with an asterisk (Rieske Domain; InterPro:
IPR017941; Pheophorbide a Oxygenase Domain; InterPro: IPR013626).

Supplementary Table S3: A summary of the occurrence of *CAO* duplicate genes across the major
groups within Viridiplantae. The number in brackets represents the proportion of species which
encode for multiple copies of the *CAO* gene (Mutiple CAO) or encode for two *CAO* genes that form
heterodimeric complex (Heterodimeric CAO).

Supplementary Table S4: *CAO* gene expression in species whose genome encodes for more than
one *CAO* gene. All data was obtained from previously published work. Gene expression is
presented as FPKM, RPKM or TPM averages with standard deviation. In all cases, gene expression
ratios were determined by dividing all expression values by the highest CAO expression level within
the species.

Supplementary Figure 1: An alignment of the CAO protein from (A) *Arabidopsis thaliana* and (B)
Chlamydomonas reinhardtii with all other proteins identified using the CAO amino acid sequence
as a query. The color of the residue depends on the degree of conservation among proteins (blue
fully conserved; green 80–100% similar; yellow 60–80% similar; white < 60% similar). The important
domains that are shared among all proteins are highlighted in green (Rieske Domain; InterPro:
IPR017941), yellow (Fe-binding domain) and blue (Pheophorbide a Oxygenase Domain; InterPro:
IPR013626).

Supplementary Figure 2: A phylogenetic tree of CAO genes inferred from using maximum
likelihood analysis. The node labels show the bootstrap support, with dashed lines indicating <50
bootstrap support. Scale bar indicates amino acid substitutions per site. Species with more than
one detected CAO gene are highlighted in red. Cases where CAO duplication may have arisen in a
common shared ancestor are marked with a blue arrow, and only those located on branches with a
strong bootstrap support (>85%) are shown. The inset shows the outline of the full CAO
phylogenetic tree inferred from using maximum likelihood analysis, using 326 unique CAO
sequences across 246 species color coded to represent the major Viridiplantae groups. The
position of the chlorophytes (green) is highlighted with a black box. (**A**) Chlorophytes; (**B**)
Charophytes, bryophytes (including marachantiophytes), and lycophytes; (**C**) Gymnosperms,
magnoliids, and monocots; (**D**) Basal eudicots, asterids and fabids; (**F**) Malvids.

Supplementary Figure 3: Multiple sequence alignment of the predicted CAO amino acid
sequences from Chlorophytes (**A**), Charophytes (**B**), non-vascular plants (**C**) and vascular plants
(**D**). Global alignments were performed with MUSCLE. The colour of the residue depends on the
degree of conservation among species (blue, fully conserved; green, 80–100% similar; yellow, 60–
80% similar; gray, < 60% similar). As a reference, the CAO sequence from *Arabidopsis thaliana* is
present in all alignments (top), with the key domains annotated (green, Degron sequence; blue,
conserved Rieske cluster; purple, Fe-binding domain)

References

Arimoto A, Nishitsuji K, Higa Y, et al (2019) A siphonous macroalgal genome suggests convergent
functions of homeobox genes in algae and land plants. DNA Research 26:183–192.
https://doi.org/10.1093/dnares/dsz002

Arriola MB, Velmurugan N, Zhang Y, et al (2018) Genome sequences of Chlorella sorokiniana UTEX
1602 and Micractinium conductrix SAG 241.80: implications to maltose excretion by a
green alga. The Plant Journal 93:566–586. https://doi.org/10.1111/tpj.13789

Bell D, Lin Q, Gerelle WK, et al (2020) Organellomic data sets confirm a cryptic consensus on
(unrooted) land-plant relationships and provide new insights into bryophyte molecular
evolution. Am J Bot 107:91–115. https://doi.org/10.1002/ajb2.1397

Biswal AK, Pattanayak GK, Pandey SS, et al (2012) Light Intensity-Dependent Modulation of
Chlorophyll *b* Biosynthesis and Photosynthesis by Overexpression of Chlorophyllide *a*
Oxygenase in Tobacco. Plant Physiol 159:433–449. https://doi.org/10.1104/pp.112.195859

Biswal AK, Pattanayak GK, Ruhil K, et al (2024) Reduced expression of chlorophyllide a oxygenase
(CAO) decreases the metabolic flux for chlorophyll synthesis and downregulates

photosynthesis in tobacco plants. Physiol Mol Biol Plants 30:1–16.
https://doi.org/10.1007/s12298-023-01395-5

Blanc-Mathieu R, Verhelst B, Derelle E, et al (2014) An improved genome of the model marine alga
Ostreococcus tauri unfolds by assessing Illumina de novo assemblies. BMC Genomics
15:1–12. https://doi.org/10.1186/1471-2164-15-1103

Blankenship RE (2010) Early Evolution of Photosynthesis1. Plant Physiol 154:434–438.
https://doi.org/10.1104/pp.110.161687

Blum M, Andreeva A, Florentino LC, et al (2025) InterPro: the protein sequence classification
resource in 2025. Nucleic Acids Research 53:D444–D456.
https://doi.org/10.1093/nar/gkae1082

Brzezowski P, Richter AS, Grimm B (2015) Regulation and function of tetrapyrrole biosynthesis in
plants and algae. Biochimica et Biophysica Acta - Bioenergetics 1847:968–985.
https://doi.org/10.1016/j.bbabio.2015.05.007

Bujaldon S, Kodama N, Rappaport F, et al (2017) Functional Accumulation of Antenna Proteins in
Chlorophyll b-Less Mutants of Chlamydomonas reinhardtii. Molecular Plant 10:115–130.
https://doi.org/10.1016/j.molp.2016.10.001

Cardona T (2019) Thinking twice about the evolution of photosynthesis. Open Biology 9:180246.
https://doi.org/10.1098/rsob.180246

Castresana J (2000) Selection of Conserved Blocks from Multiple Alignments for Their Use in
Phylogenetic Analysis. Molecular Biology and Evolution 17:540–552.
https://doi.org/10.1093/oxfordjournals.molbev.a026334

Catalan P, López-Álvarez D, Díaz-Pérez A, et al (2016) Phylogeny and Evolution of the Genus
Brachypodium. In: Vogel JP (ed) Genetics and Genomics of Brachypodium. Springer
International Publishing, Cham, pp 9–38

Cheng S, Xian W, Fu Y, et al (2019) Genomes of Subaerial Zygnematophyceae Provide Insights into
Land Plant Evolution. Cell 179:1057–1067. https://doi.org/10.1016/j.cell.2019.10.019

Conlon T, Parkes R, Fierli D, Touzet N (2024) Comparative pigment and fatty acid profiling of marine
species within the chlorophyte genus *Tetraselmis*. Food Bioscience 58:103660.
https://doi.org/10.1016/j.fbio.2024.103660

Craig RJ, Gallaher SD, Shu S, et al (2022) The Chlamydomonas Genome Project, version 6:
reference assemblies for mating type plus and minus strains reveal extensive structural
mutation in the laboratory. The Plant Cell koac347. https://doi.org/10.1093/plcell/koac347

Curtis BA, Tanifuji G, Burki F, et al (2012) Algal genomes reveal evolutionary mosaicism and the fate
of nucleomorphs. Nature 492:59–65. https://doi.org/10.1038/nature11681

Cvetkovska M, Hüner NPA, Smith DR (2017) Chilling out: the evolution and diversification of
psychrophilic algae with a focus on Chlamydomonadales. Polar Biol 40:1169–1184.
https://doi.org/10.1007/s00300-016-2045-4

Cvetkovska M, Orgnero S, Hüner NPA, Smith DR (2019) The enigmatic loss of light-independent
chlorophyll biosynthesis from an Antarctic green alga in a light-limited environment. New
Phytol 222:651–656. https://doi.org/10.1111/nph.15623

de Vries J, Archibald JM (2018) Plant evolution: landmarks on the path to terrestrial life. New
Phytologist 217:1428–1434. https://doi.org/10.1111/nph.14975

Dey D, Tanaka R, Ito H (2023) Structural Characterization of the Chlorophyllide a Oxygenase (CAO)
Enzyme Through an In Silico Approach. J Mol Evol 91:225–235.
https://doi.org/10.1007/s00239-023-10100-9

Dumont T, Barau L, Thong-Chane A, et al (2022) Sugarcane Breeding in Reunion: Challenges,
Achievements and Future Prospects. Sugar Tech 24:181–192.
https://doi.org/10.1007/s12355-021-00998-0

Edgar RC (2004) MUSCLE: a multiple sequence alignment method with reduced time and space
complexity. BMC Bioinformatics 5:113. https://doi.org/10.1186/1471-2105-5-113

Espineda CE, Linford AS, Devine D, Brusslan JA (1999) The AtCAO gene, encoding chlorophyll a
oxygenase, is required for chlorophyll b synthesis in Arabidopsis thaliana. Proceedings of
the National Academy of Sciences of the United States of America 96:10507–10511.
https://doi.org/10.1073/pnas.96.18.10507

Ferraro DJ, Gakhar L, Ramaswamy S (2005) Rieske business: Structure–function of Rieske non-
heme oxygenases. Biochemical and Biophysical Research Communications 338:175–190.
https://doi.org/10.1016/j.bbrc.2005.08.222

Goodstein DM, Shu S, Howson R, et al (2012) Phytozome: a comparative platform for green plant
genomics. Nucleic Acids Research 40:D1178–D1186. https://doi.org/10.1093/nar/gkr944

Gray J, Wardzala E, Yang M, et al (2004) A small family of LLS1-related non-heme oxygenases in
plants with an origin amongst oxygenic photosynthesizers. Plant Molecular Biology 54:39–
54

Grigoriev IV, Hayes RD, Calhoun S, et al (2021) PhycoCosm, a comparative algal genomics
resource. Nucleic Acids Research 49:D1004–D1011. https://doi.org/10.1093/nar/gkaa898

Guiry MD (2024) How many species of algae are there? A reprise. Four kingdoms, 14 phyla, 63
classes and still growing. Journal of Phycology 60:214–228.
https://doi.org/10.1111/jpy.13431

Guo Q, Zhu S, Lai T, et al (2025) A Phosphate-Starvation Enhanced Purple Acid Phosphatase,
GmPAP23 Mediates Intracellular Phosphorus Recycling and Yield in Soybean. Plant, Cell
and Environment 1–18. https://doi.org/10.1111/pce.15400

Han S, Zhang S, Yi R, et al (2024) Phylogenomics and plastomics offer new evolutionary perspectives on Kalanchoideae (Crassulaceae). Annals of Botany 133:585–604. https://doi.org/10.1093/aob/mcae017

He Y, Deng Z, He S, et al (2025) Transcriptome and co-expression network analysis reveal the genetic basis of cell wall components in maize stalks. BMC Genomics 26:. https://doi.org/10.1186/s12864-025-11816-2

Healey AL, Garsmeur O, Lovell JT, et al (2024) The complex polyploid genome architecture of sugarcane. Nature 628:804–810. https://doi.org/10.1038/s41586-024-07231-4

Healey AL, Piatkowski B, Lovell JT, et al (2023) Newly identified sex chromosomes in the Sphagnum (peat moss) genome alter carbon sequestration and ecosystem dynamics. Nature Plants 9:238–254. https://doi.org/10.1038/s41477-022-01333-5

Hendriks KP, Kiefer C, Al-Shehbaz IA, et al (2023) Global Brassicaceae phylogeny based on filtering of 1,000-gene dataset. Current Biology 33:4052-4068.e6. https://doi.org/10.1016/j.cub.2023.08.026

Hirashima M, Satoh S, Tanaka R, Tanaka A (2006) Pigment Shuffling in Antenna Systems Achieved by Expressing Prokaryotic Chlorophyllide a Oxygenase in Arabidopsis. The Journal of Biological Chemistry 281:15385–15393. https://doi.org/10.1074/jbc.M602903200

Huang C-H, Sun R, Hu Y, et al (2016) Resolution of Brassicaceae Phylogeny Using Nuclear Genes Uncovers Nested Radiations and Supports Convergent Morphological Evolution. Molecular Biology and Evolution 33:394–412. https://doi.org/10.1093/molbev/msv226

Hüner NPA, Smith DR, Cvetkovska M, et al (2022) Photosynthetic adaptation to polar life: Energy balance, photoprotection and genetic redundancy. Journal of Plant Physiology 268:153557. https://doi.org/10.1016/j.jplph.2021.153557

Iha C, Dougan KE, Varela JA, et al (2021) Genomic adaptations to an endolithic lifestyle in the coral-associated alga *Ostreobium*. Current Biology 31:1393-1402.e5. https://doi.org/10.1016/j.cub.2021.01.018

Ikegami I, Satoh S, Aoki M (2007) Binding Affinity of Chl b for the Chl a -Binding Sites in PSI Core Complexes. Plant Cell Physiology 48:1092–1097. https://doi.org/10.1093/pcp/pcm078

Isono E, Li J, Pulido P, et al (2024) Protein degrons and degradation: Exploring substrate recognition and pathway selection in plants. The Plant Cell 36:3074–3098. https://doi.org/10.1093/plcell/koae141

Ji Y, Landis JB, Yang J, et al (2023) Phylogeny and evolution of Asparagaceae subfamily Nolinoideae: new insights from plastid phylogenomics. Annals of Botany 131:301–312. https://doi.org/10.1093/aob/mcac144

Jung YJ, Lee HJ, Yu J, et al (2021) Transcriptomic and physiological analysis of OsCAO1 knockout lines using the CRISPR/Cas9 system in rice. Plant Cell Rep 40:1013–1024. https://doi.org/10.1007/s00299-020-02607-y

Junkins EN, Stamps BW, Corsetti FA, et al (2019) Draft Genome Sequence of Picocystis sp. Strain
ML, Cultivated from Mono Lake, California. Microbiol Resour Announc 8:e01353-18.
https://doi.org/10.1128/MRA.01353-18

Kersey PJ (2019) Plant genome sequences: past, present, future. Current Opinion in Plant Biology
48:1–8. https://doi.org/10.1016/j.pbi.2018.11.001

Kim EH, Li XP, Razeghifard R, et al (2009) The multiple roles of light-harvesting chlorophyll a/b-
protein complexes define structure and optimize function of Arabidopsis chloroplasts: A
study using two chlorophyll b-less mutants. Biochimica et Biophysica Acta - Bioenergetics
1787:973–984. https://doi.org/10.1016/j.bbabio.2009.04.009

Kondrashov FA (2012) Gene duplication as a mechanism of genomic adaptation to a changing
environment. Proceedings of the Royal Society B: Biological Sciences 279:5048–5057.
https://doi.org/10.1098/rspb.2012.1108

Król M, Spangfort MD, Huner NPA, et al (1995) Chlorophyll a/b-binding proteins, pigment
conversions, and early light-induced proteins in a chlorophyll b-less Barley mutant. Plant
Physiology 107:873–883. https://doi.org/10.1104/pp.107.3.873

Kuai B, Chen J, Hörtensteiner S (2018) The biochemistry and molecular biology of chlorophyll
breakdown. Journal of Experimental Botany 69:751–767. https://doi.org/10.1093/jxb/erx322

Kume A, Akitsu T, Nasahara KN (2018) Why is chlorophyll b only used in light-harvesting systems?
Journal of Plant Research 131:961–972. https://doi.org/10.1007/s10265-018-1052-7

Kunugi M, Satoh S, Ihara K, et al (2016) Evolution of Green Plants Accompanied Changes in Light-
Harvesting Systems. Plant and Cell Physiology 57:1231–1243.
https://doi.org/10.1093/pcp/pcw071

Kunugi M, Takabayashi A, Tanaka A (2013) Evolutionary changes in chlorophyllide a oxygenase
(CAO) structure contribute to the acquisition of a new light-harvesting complex in
micromonas. Journal of Biological Chemistry 288:19330–19341.
https://doi.org/10.1074/jbc.M113.462663

Lamesch P, Berardini TZ, Li D, et al (2012) The Arabidopsis Information Resource (TAIR): improved
gene annotation and new tools. Nucleic Acids Res 40:D1202-1210.
https://doi.org/10.1093/nar/gkr1090

Lee S, Kim JH, Eun SY, et al (2005a) Differential regulation of chlorophyll a oxygenase genes in rice.
Plant Molecular Biology 57:805–818. https://doi.org/10.1007/s11103-005-2066-9

Lee S, Kim J-H, Yoo ES, et al (2005b) Differential regulation of chlorophyll a oxygenase genes in rice.
Plant Mol Biol 57:805–818. https://doi.org/10.1007/s11103-005-2066-9

Leebens-Mack JH, Barker MS, Carpenter EJ, et al (2019) One thousand plant transcriptomes and
the phylogenomics of green plants. Nature 574:679–685. https://doi.org/10.1038/s41586-
019-1693-2

Leliaert F, Smith DR, Moreau H, et al (2012) Phylogeny and Molecular Evolution of the Green Algae.
Critical Reviews in Plant Sciences 31:1–46. https://doi.org/10.1080/07352689.2011.615705

Lemieux C, Turmel M, Otis C, Pombert J-F (2019) A streamlined and predominantly diploid genome
in the tiny marine green alga Chloropicon primus. Nat Commun 10:4061.
https://doi.org/10.1038/s41467-019-12014-x

Letunic I, Bork P (2021) Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display
and annotation. Nucleic Acids Res 49:W293–W296. https://doi.org/10.1093/nar/gkab301

Li X, Hou Z, Xu C, et al (2021) Large Phylogenomic Data sets Reveal Deep Relationships and Trait
Evolution in Chlorophyte Green Algae. Genome Biol Evol 13:evab101.
https://doi.org/10.1093/gbe/evab101

Liu J, Knapp M, Jo M, et al (2022a) Rieske Oxygenase Catalyzed C–H Bond Functionalization
Reactions in Chlorophyll b Biosynthesis. ACS Cent Sci 8:1393–1403.
https://doi.org/10.1021/acscentsci.2c00058

Liu Y, Liu J, Kong Z, et al (2022b) Transcriptomics and metabolomics analyses of the mechanism of
flavonoid synthesis in seeds of differently colored quinoa strains. Genomics 114:138–148.
https://doi.org/10.1016/j.ygeno.2021.11.030

Mabry ME, Brose JM, Blischak PD, et al (2020) Phylogeny and multiple independent whole-genome
duplication events in the Brassicales. Am J Bot 107:1148–1164.
https://doi.org/10.1002/ajb2.1514

Mascuñano B, Coto-Elena J, Guerrero-Sánchez VM, et al (2025) Transcriptome analysis of wild olive
(Olea Europaea L. subsp. europaea var. sylvestris) clone AC18 provides insight into the role
of lignin as a constitutive defense mechanism underlying resistance to Verticillium wilt.
BMC Plant Biology 25:. https://doi.org/10.1186/s12870-025-06301-7

Massé A, Tribollet A, Meziane T, et al (2020) Functional diversity of microboring Ostreobium algae
isolated from corals. Environmental Microbiology 22:4825–4846.
https://doi.org/10.1111/1462-2920.15256

Merchant SS, Prochnik SE, Vallon O, et al (2007) The Chlamydomonas genome reveals the
evolution of key animal and plant functions. Science 318:245–250.
https://doi.org/10.1126/science.1143609

Miller MA, Schwartz T, Pickett BE, et al (2015) A RESTful API for Access to Phylogenetic Tools via the
CIPRES Science Gateway. Evol Bioinform Online 11:43–48.
https://doi.org/10.4137/EBO.S21501

Moreau H, Verhelst B, Couloux A, et al (2012) Gene functionalities and genome structure in
Bathycoccus prasinos reflect cellular specializations at the base of the green lineage.
Genome Biol 13:1–16. https://doi.org/10.1186/gb-2012-13-8-r74

Mueller AH, Dockter C, Gough SP, Lundqvist U (2012) Characterization of Mutations in Barley fch2
Encoding Chlorophyllide a Oxygenase. Plant and Cell Physiology 53:1232–1246.
https://doi.org/10.1093/pcp/pcs062

Nagata N, Satoh S, Tanaka R, Tanaka A (2004) Domain structures of chlorophyllide a oxygenase of
green plants and Prochlorothrix hollandica in relation to catalytic functions. Planta
218:1019–1025. https://doi.org/10.1007/s00425-003-1181-6

Nakagawara E, Sakuraba Y, Yamasato A, et al (2007) Clp protease controls chlorophyll b synthesis
by regulating the level of chlorophyllide a oxygenase. The Plant Journal 49:800–809.
https://doi.org/10.1111/j.1365-313X.2006.02996.x

Neale PJ, Priscu JC (1995) The Photosynthetic Apparatus of Phytoplankton from a Perennially Ice-
Covered Antarctic Lake: Acclimation to an Extreme Shade Environment. Plant and Cell
Physiology 36:253–263. https://doi.org/10.1093/oxfordjournals.pcp.a078757

Neilson JAD, Durnford DG (2010) Structural and functional diversification of the light-harvesting
complexes in photosynthetic eukaryotes. Photosynthesis Research 106:57–71.
https://doi.org/10.1007/s11120-010-9576-2

Nick S, Meurer J, Soll J, Ankele E (2013) Nucleus-encoded light-harvesting chlorophyll a/b proteins
are imported normally into chlorophyll b-free chloroplasts of arabidopsis. Molecular Plant
6:860–871. https://doi.org/10.1093/mp/sss113

Oster U, Tanaka R, Tanaka A, Rüdiger W (2000) Cloning and functional expression of the gene
encoding the key enzyme for chlorophyll b biosynthesis (CAO) from Arabidopsis thaliana.
Plant Journal 21:305–310. https://doi.org/10.1046/j.1365-313X.2000.00672.x

Palenik B, Grimwood J, Aerts A, et al (2007) The tiny eukaryote Ostreococcus provides genomic
insights into the paradox of plankton speciation. Proceedings of the National Academy of
Sciences 104:7705–7710. https://doi.org/10.1073/pnas.0611046104

Palenik B, Haselkorn R (1992) Multiple evolutionary origins of prochlorophytes, the chlorophyllb-
containing prokaryotes. Nature 355:265–267. https://doi.org/10.1038/355265a0

Panchy N, Lehti-Shiu M, Shiu S-H (2016) Evolution of Gene Duplication in Plants. Plant Physiol
171:2294–2316. https://doi.org/10.1104/pp.16.00523

Partensky F, Six C, Ratin M, et al (2018) A novel species of the marine cyanobacterium
Acaryochloris with a unique pigment content and lifestyle. Sci Rep 8:9142.
https://doi.org/10.1038/s41598-018-27542-7

Pattanayak GK, Biswal AK, Reddy VS, Tripathy BC (2005) Light-dependent regulation of chlorophyll
b biosynthesis in chlorophyllide a oxygenase overexpressing tobacco plants. Biochemical
and Biophysical Research Communications 326:466–471.
https://doi.org/10.1016/j.bbrc.2004.11.049

Perroud PF, Haas FB, Hiss M, et al (2018) The Physcomitrella patens gene atlas project: large-scale
RNA-seq based expression data. Plant Journal 95:168–182.
https://doi.org/10.1111/tpj.13940

Piatkowski BT, Shaw AJ (2019) Functional trait evolution in Sphagnum peat mosses and its
relationship to niche construction. New Phytologist 223:939–949.
https://doi.org/10.1111/nph.15825

Poirier MC, Fugard K, Cvetkovska M (2025) Light quality affects chlorophyll biosynthesis and
photosynthetic performance in Antarctic Chlamydomonas. Photosynth Res 163:9.
https://doi.org/10.1007/s11120-024-01127-0

Pombert J-F, Blouin NA, Lane C, et al (2014) A Lack of Parasitic Reduction in the Obligate Parasitic
Green Alga Helicosporidium. PLoS Genet 10:e1004355.
https://doi.org/10.1371/journal.pgen.1004355

Pružinská A, Tanner G, Anders I, et al (2003) Chlorophyll breakdown: Pheophorbide a oxygenase is
a Rieske-type iron–sulfur protein, encoded by the accelerated cell death 1 gene.
Proceedings of the National Academy of Sciences 100:15259–15264.
https://doi.org/10.1073/pnas.2036571100

Przybyla-Toscano J, Boussardon C, Law SR, et al (2021) Gene atlas of iron-containing proteins in
Arabidopsis thaliana. The Plant Journal 106:258–274. https://doi.org/10.1111/tpj.15154

Reinbothe C, Bakkouri ME, Buhr F, et al (2010) Chlorophyll biosynthesis: spotlight on
protochlorophyllide reduction. Trends in Plant Science 15:614–624.
https://doi.org/10.1016/j.tplants.2010.07.002

Reinbothe C, Bartsch S, Eggink LL, et al (2006) A role for chlorophyllide a oxygenase in the regulated
import and stabilization of light-harvesting chlorophyll a/b proteins. Proceedings of the
National Academy of Sciences of the United States of America 103:4777–4782.
https://doi.org/10.1073/pnas.0511066103

Roche JL, van der Staay GWM, Partensky F, et al (1996) Independent evolution of the
prochlorophyte and green plant chlorophyll a/b light-harvesting proteins. Proceedings of
the National Academy of Sciences 93:15244–15248.
https://doi.org/10.1073/pnas.93.26.15244

Roesler CS, Culbertson CW, Etheridge SM, et al (2002) Distribution, production, and ecophysiology
of Picocystis strain ML in Mono Lake, California. Limnology and Oceanography 47:440–452.
https://doi.org/10.4319/lo.2002.47.2.0440

Roy P, Sethi S, New J, et al (2025) Transcriptomic Response of White Lupin Roots to Short-Term
Sucrose Treatment. Plants 14:14–17. https://doi.org/10.3390/plants14030381

Ruggiero MA, Gordon DP, Orrell TM, et al (2015) A Higher Level Classification of All Living
Organisms. PLoS One 10:e0119248. https://doi.org/10.1371/journal.pone.0119248

Sakuraba Y, Tanaka R, Yamasato A, Tanaka A (2009) Determination of a chloroplast degron in the
regulatory domain of chlorophyllide a oxygenase. Journal of Biological Chemistry
284:36689–36699. https://doi.org/10.1074/jbc.M109.008144

Sánchez-Baracaldo P, Cardona T (2020) On the origin of oxygenic photosynthesis and
Cyanobacteria. New Phytologist 225:1440–1446. https://doi.org/10.1111/nph.16249

Satoh S, Ikeuchi M, Mimuro M, Tanaka A (2001) Chlorophyll b Expressed in Cyanobacteria
Functions as a Light-harvesting Antenna in Photosystem I through Flexibility of the Proteins.
The Journal of Biological Chemistry 276:4293–4297.
https://doi.org/10.1074/jbc.M008238200

Schmidt CL, Shaw L (2001) A Comprehensive Phylogenetic Analysis of Rieske and Rieske-Type
Iron-Sulfur Proteins. J Bioenerg Biomembr 33:9–26.
https://doi.org/10.1023/A:1005616505962

Schoch CL, Ciufo S, Domrachev M, et al (2020) NCBI Taxonomy: a comprehensive update on
curation, resources and tools. Database (Oxford) 2020:baaa062.
https://doi.org/10.1093/database/baaa062

Schumacher I, Menghini D, Ovinnikov S, et al (2022) Evolution of chlorophyll degradation is
associated with plant transition to land. The Plant Journal 109:1473–1488.
https://doi.org/10.1111/tpj.15645

Shang S, Zhang Z, Li L, et al (2023) Transcriptome analysis reveals genes expression pattern of
Spirodela polyrhiza response to heat stress. International Journal of Biological
Macromolecules 225:767–775. https://doi.org/10.1016/j.ijbiomac.2022.11.139

Soltis PS, Soltis DE (2021) Plant genomes: Markers of evolutionary history and drivers of
evolutionary change. PLANTS, PEOPLE, PLANET 3:74–82.
https://doi.org/10.1002/ppp3.10159

Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. Bioinformatics 30:1312–1313. https://doi.org/10.1093/bioinformatics/btu033

Steadman Tyler CR, Hovde BT, Daligault HE, et al (2019) High-Quality Draft Genome Sequence of
the Green Alga Tetraselmis striata (Chlorophyta) Generated from PacBio Sequencing.
Microbiology Resource Announcements 8:10.1128/mra.00780-19.
https://doi.org/10.1128/mra.00780-19

Stolzer M, Lai H, Xu M, et al (2012) Inferring duplications, losses, transfers and incomplete lineage
sorting with nonbinary species trees. Bioinformatics 28:i409–i415.
https://doi.org/10.1093/bioinformatics/bts386

Suzuki S, Hirakawa Y, Kofuji R, et al (2016) Plastid genome sequences of Gymnochlora stellata,
Lotharella vacuolata, and Partenskyella glossopodia reveal remarkable structural
conservation among chlorarachniophyte species. J Plant Res 129:581–590.
https://doi.org/10.1007/s10265-016-0804-5

Talavera G, Castresana J (2007) Improvement of phylogenies after removing divergent and
ambiguously aligned blocks from protein sequence alignments. Syst Biol 56:564–577.
https://doi.org/10.1080/10635150701472164

Tanaka A, Ito H, Tanaka R, et al (1998) Chlorophyll a oxygenase (CAO) is involved in chlorophyll b
formation from chlorophyll a. Proceedings of the National Academy of Sciences of the
United States of America 95:12719–12723. https://doi.org/10.1073/pnas.95.21.12719

Tanaka A, Tanaka R (2019) The biochemistry, physiology, and evolution of the chlorophyll cycle, 1st
edn. Elsevier Ltd.

Tanaka R, Tanaka A (2007) Tetrapyrrole biosynthesis in higher plants. Annual Review of Plant
Biology 58:321–346. https://doi.org/10.1146/annurev.arplant.57.032905.105448

Tanaka R, Tanaka A (2005) Effects of chlorophyllide a oxygenase overexpression on light
acclimation in Arabidopsis thaliana. Photosynth Res 85:327–340.
https://doi.org/10.1007/s11120-005-6807-z

Tanaka R, Tanaka A (2011) Chlorophyll cycle regulates the construction and destruction of the
light-harvesting complexes. Biochimica et Biophysica Acta (BBA) - Bioenergetics 1807:968–
976. https://doi.org/10.1016/j.bbabio.2011.01.002

Tartar A, Boucias DG, Becnel JJ, Adams BJ (2003) Comparison of plastid 16S rRNA (rrn16) genes
from Helicosporidium spp.: evidence supporting the reclassification of Helicosporidia as
green algae (Chlorophyta). Int J Syst Evol Microbiol 53:1719–1723.
https://doi.org/10.1099/ijs.0.02559-0

Tomitani A, Okada K, Miyashita H, et al (1999a) Chlorophyll b and phycobilins in the common
ancestor of cyanobacteria and chloroplasts. Nature 400:159–162.
https://doi.org/10.1038/22101

Tomitani A, Okada K, Miyashita H, et al (1999b) Chlorophyll b and phycobilins in the common
ancestor of cyanobacteria and chloroplasts. Letters to nature 400:159–162

Turner-Hissong SD, Mabry ME, Beissinger TM, et al (2020) Evolutionary insights into plant breeding.
Current Opinion in Plant Biology 54:93–100. https://doi.org/10.1016/j.pbi.2020.03.003

Ueno Y, Aikawa S, Kondo A, Akimoto S (2019) Adaptation of light-harvesting functions of unicellular
green algae to different light qualities. Photosynthesis Research 139:145–154.
https://doi.org/10.1007/s11120-018-0523-y

Voitsekhovskaja OV, Tyutereva EV (2015) Chlorophyll *b* in angiosperms: Functions in
photosynthesis, signaling and ontogenetic regulation. Journal of Plant Physiology 189:51–
64. https://doi.org/10.1016/j.jplph.2015.09.013

Wang B, Yang X, Jia Y, et al (2022) High-quality Arabidopsis thaliana Genome Assembly with
Nanopore and HiFi Long Reads. Genomics Proteomics Bioinformatics 20:4–13.
https://doi.org/10.1016/j.gpb.2021.08.003

Wang W, Wang L, Wang L, et al (2021) Transcriptome analysis and molecular mechanism of linseed
(Linum usitatissimum L.) drought tolerance under repeated drought using single-molecule
long-read sequencing. BMC Genomics 22:1–23. https://doi.org/10.1186/s12864-021-
07416-5

Willows RD (2020) Biosynthesis of Chlorophyll and Bilins in Algae. In: Larkum AWD, Grossman AR,
Raven JA (eds) Advances in Photosynthesis and Respiration, 45th edn. Springer Nature, pp
83–103

Worden AZ, Lee J-H, Mock T, et al (2009) Green Evolution and Dynamic Adaptations Revealed by
Genomes of the Marine Picoeukaryotes Micromonas. Science 324:268–272.
https://doi.org/10.1126/science.1167222

Xiong J, Bauer CE (2002) COMPLEX EVOLUTION OF PHOTOSYNTHESIS. Annual Review of Plant
Biology 53:503–521. https://doi.org/10.1146/annurev.arplant.53.100301.135212

Xiong J, Wen G, Song J, et al (2024) Knockout of the Chlorophyll a Oxygenase Gene OsCAO1
Reduces Chilling Tolerance in Rice Seedlings. Genes 15:721.
https://doi.org/10.3390/genes15060721

Xu H, Vavilin D, Vermaas W (2001) Chlorophyll b can serve as the major pigment in functional
photosystem II complexes of cyanobacteria. Proceedings of the National Academy of
Sciences of the United States of America 98:14168–14173.
https://doi.org/10.1073/pnas.251530298

Yamasato A, Nagata N, Tanaka R, Tanaka A (2005) The N-Terminal Domain of Chlorophyllide a
Oxygenase Confers Protein Instability in Response to Chlorophyll b Accumulation in
Arabidopsis. Plant Cell 17:1585–1597. https://doi.org/10.1105/tpc.105.031518

Zhang L, Yang C, Liu C (2023) Revealing the significance of chlorophyll b in the moss
Physcomitrium patens by knocking out two functional chlorophyllide a oxygenase.
Photosynthesis Research 158:171–180. https://doi.org/10.1007/s11120-023-01044-8

Zhu X, Tai X, Ren Y, et al (2019) Genome-wide analysis of coding and long non-coding RNAs
involved in cuticular wax biosynthesis in cabbage (Brassica oleracea L. var. capitata).
International Journal of Molecular Sciences 20:. https://doi.org/10.3390/ijms20112820

Zhuang Y, Wang X, Li X, et al (2022) Phylogenomics of the genus Glycine sheds light on polyploid
evolution and life-strategy transition. Nat Plants 8:233–244.
https://doi.org/10.1038/s41477-022-01102-4