Title: Pneumolysin-dependent and independent non-canonical autophagy processes

mediate host defense against pneumococcal infection

Authors: Bartosz J. Michno[1,2], Niedharsan Pooranachandran[1], Tonisha C. Smith[3],

Erin Faught[4], Sandra Lipowská[1], Andrew K. Fenton[3], Annemarie H. Meijer[4], Tomasz

K. Prajsnar[1]

1 Department of Evolutionary Immunology, Institute of Zoology and Biomedical

Research, Faculty of Biology, Jagiellonian University, Krakow, Poland

2 Doctoral School of Exact and Natural Sciences, Jagiellonian University, Krakow,

Poland

3 School for Biosciences, Florey Institute for Host-Pathogen Interactions, University

of Sheffield, Sheffield, United Kingdom

4 Institute of Biology Leiden, Leiden University, Leiden, The Netherlands

Corresponding author: Tomasz K. Prajsnar

Email: tomasz.prajsnar@uj.edu.pl

Phone: +48 12 6645377

Key words:

CASM; LAP; macrophage; *pneumoniae*; STIL; *Streptococcus*; zebrafish

Abstract:

Streptococcus pneumoniae is an opportunistic pathogen responsible for life-threatening diseases including pneumonia and meningitis. The host defense against pneumococci relies heavily on macrophages, which can effectively internalize and degrade bacteria. Recent studies have implicated both canonical and non-canonical autophagy-related processes in bacterial clearance, but the precise pathways mediating defense against *S. pneumoniae* remain unknown. Here, we utilize a well-established zebrafish larval infection model to investigate the role of autophagy in host defense against pneumococci *in vivo*. Using a transgenic macroautophagy/autophagy reporter line, we found the autophagy marker Map1lc3/Lc3 being recruited to pneumococci-containing vesicles upon bacterial internalization by zebrafish macrophages. The genetic inhibition of core autophagy gene *atg5* led to loss of the Lc3 associations and their impaired acidification, significantly delaying bacterial clearance. This Lc3 recruitment is partially mediated by LC3-associated phagocytosis (LAP), as knockdown of *cyba* and *rubcn* moderately reduced Lc3 association with phagosomes and diminished pneumococcal degradation. Interestingly, we observed no involvement of xenophagy components in *S. pneumoniae*-infected macrophages, suggesting the activation of another non-canonical autophagy pathway, distinct from LAP, targeting pneumococci-containing phagosomes. Instead, we found that the pneumococcal pore-forming toxin pneumolysin induces ROS-independent CASM pathways, one of which is abolished by knockdown of *tecpr1a* indicating the involvement of sphingomyelin-Tecpr1-induced LC3 lipidation (STIL). Collectively, our observations shed new light on the host immune response against *S. pneumoniae*, demonstrating that several distinct non-canonical autophagy pathways mediate bacterial degradation by macrophages and providing potential targets for the development of novel therapies to combat pneumococcal infections.

ABBREVIATIONS: ATG: autophagy related; BMDM: bone marrow-derived macrophage; CASM: conjugation of ATG8 to single membranes; CFU: colony-forming units; Cyba: cytochrome b-245, alpha polypeptide; DPI: diphenyleneiodonium, GFP: green fluorescent protein; hpf: hours post-fertilization; hpi: hours post-infection; LAP: LC3-associated phagocytosis; Map1lc3/Lc3: microtubule-associated protein 1 light chain 3; MEF: mouse embryonic fibroblast;

54 NADPH: nicotinamide adenine dinucleotide phosphate; Optn: optineurin, Ply:

55 pneumolysin; ROS: reactive oxygen species; SLR: sequestosome-like receptors;

56 Sqstm1: sequestosome 1; STIL: sphingomyelin-TECPR1-induced LC3 lipidation;

57 Tecpr1: tectonin beta-propeller repeat containing 1.

Introduction

Streptococcus pneumoniae (pneumococcus) is an opportunistic pathogen that

colonizes the human upper respiratory tract and is able to invade other tissues

including the central nervous system. This bacterium causes several life-threatening

diseases such as community-acquired pneumonia [1] resulting in more than 1 million

deaths worldwide each year [2]. Pneumococcus has been repeatedly included in the

WHO's list of priority bacterial pathogens since 2017 to date (WHO report, 2024).

The widespread use of various antibacterial drugs, mostly β-lactam antibiotics such

as penicillin or ampicillin, combined with the high genome plasticity of pneumococci

facilitates the spread of pneumococcal resistance to drugs and vaccines, limiting their

effectiveness [3,4]. Therefore, alternative treatment strategies are urgently needed to

combat this pathogen.

The first line of defense against invading pneumococci are tissue-resident alveolar

macrophages in the lung [5]. Once internalized, the bacteria sequestered in

phagosomes are rapidly acidified by activation of the vacuolar-type H^+-translocating

ATPase (V-ATPase), which pumps protons into the vacuole [6–8]. This acidification is

required for *S. pneumoniae* killing by phagocytes in the zebrafish model system [9].

Furthermore, lysosomal-dependent degradation, following fusion of phagosomes with

lysosomes, has been well described in the literature and is known to be involved in

the antibacterial response [10]. Aside from classical phagocytosis, other pathways

can also lead to intracellular bacteria degradation including autophagy-related

mechanisms which are currently being intensively studied [11]. These are either

canonical pathogen-selective autophagy, also known as xenophagy [12], or more

recently identified non-canonical autophagy pathways, collectively referred to as

conjugation of ATG8 to single membranes (CASM) [13,14].

Xenophagy is a form of canonical autophagy in which either ruptured bacteria-

containing phagosomes or pathogens that have escaped into the cytosol are

detected and tagged by various proteins such as ubiquitin or glycan-binding galectins

[15,16]. Subsequently, such labeled bacteria bind to sequestosome-like receptors

(SLRs) and MAP1LC3/LC3 (microtubule-associated protein 1 light chain 3) [12,17,18].

LC3 association promotes the phagophore expansion resulting in formation of a

double-membrane autophagosome [19]. It has recently been reported that SLRs such

as SQSTM1/p62 (sequestosome 1) and CALCOCO2/NDP52 (calcium binding and

coiled-coil domain 2) contribute to the initiation of xenophagy of intracellular *S.*

pneumoniae within non-professional phagocytes such as human pulmonary

epithelial cells and mouse embryonic fibroblasts (MEFs) *in vitro* [20,21]. It has been

also shown that pneumolysin (Ply), a pore-forming toxin produced *by S. pneumoniae*

that is able to induce membrane rupture of the vesicles containing bacteria leading to

subsequent xenophagy [21].

The most studied non-canonical autophagy pathway is LAP, a form of CASM that

involves many of the same molecular components as xenophagy, but not SLRs. In

this case, LC3 is recruited to the single-membrane phagosome triggered by

perturbation of ionic and pH balance [13]. It is generally accepted that LC3 lipidation

in LAP requires the activity of NOX (NADPH oxidase) and CYBA, generating reactive

oxygen species (ROS) [22]. This ROS production has been shown to promote an

interaction between the V-ATPase and ATG16L1, which initiates LC3 lipidation [23].

We have recently shown that once ROS production is blocked, the LAP response to

various bacterial pathogens is greatly reduced [24,25]. In addition, RUBCN/RUBICON

(rubicon autophagy regulator) has been implicated as a positive regulator of LAP

acting upstream of NADPH oxidase complex assembly [22]. Recent *in vitro* evidence

has shown that ROS-dependent LAP occurs within mouse bone marrow-derived

macrophages (BMDMs) infected by pneumococci and significantly contributes to

bacterial degradation [26].

Although the degradation of invading *S. pneumoniae* through LAP or xenophagy has

been investigated *in vitro*, results differ between studies in professional and non-

professional phagocytes. While in MEFs, xenophagy has been demonstrated to play

a role in pneumococcal degradation and its activation was preceded by atg16l1-

dependent but ROS-independent LAP-like LC3 associations [20,21], no such

response was observed in macrophages [26]. Additionally, the specific interplay of

these pathways in the wider and more complex host environment have not yet been

elucidated. Unveiling the role of these mechanisms *in vivo* would support the

development of innovative immunomodulatory approaches against *S. pneumoniae* in
the future.

Previously, we established a model of pneumococcal infection in larval zebrafish and
described that zebrafish macrophages can prevent dissemination of intravenously
administered unencapsulated *S. pneumoniae* [9]. The macrophages protect infected
larvae by internalization and acidification-mediated killing of pneumococci,
underscoring the cross-species evolutionary conserved role of these professional
phagocytes in host protection against this pathogen, and providing new opportunities
to gain insight into the host-pneumococcus interactions *in vivo.*

In the present study, we focus on the role of autophagy-related mechanisms in *S.*
pneumoniae infection using zebrafish larvae. We demonstrate that upon
internalization of pneumococci, the vast majority of infected macrophages rapidly
respond by the Lc3 conjugation to phagosomal membrane which facilitates
phagosomal acidification and bacterial killing. We also show that, rather than being
mediated by xenophagy, the protective role of autophagy in pneumococcal infection
of macrophages is primarily associated with CASM such as LAP and other
pneumolysin-mediated pathways including the sphingomyelin-TECPR1-induced LC3
lipidation (STIL).

Results

Intracellular* S. pneumoniae *triggers Lc3 recruitment to bacteria-containing
phagosomes in zebrafish macrophages.

To determine whether the autophagy response occurs to intracellular pneumococci,
we used our previously established *S. pneumoniae* infection model of zebrafish
larvae [9] using the transgenic zebrafish line *Tg(CMV:EGFP-map1lc3b)^{zf155}*, hereafter
called *CMV:GFP-Lc3* [27]. The GFP-LC3 reporter has been widely used in studying
various autophagy pathways in response to infection [28,29]. The zebrafish larvae
were injected intravenously with fluorescently labeled unencapsulated *S.*
pneumoniae (D39 Δ*cps hlpA*::*mKate2*) and the infected larvae fixed at 2, 4 and 7 h
post infection (hpi). Confocal analysis revealed robust activation of the autophagy
machinery, as more than 70% of infected phagocytes contained bacterial clusters
surrounded by Lc3 at 2 hpi and this response gradually decreased by 7 hpi (Fig. 1A-
B). The time-dependent loss of Lc3-bacteria associations is probably associated with

rapid autophagic flux leading to degradation of internalized pneumococci upon

phagosomal acidification that we have observed before [9]. Indeed, in addition to the

decrease of Lc3 associations over time, the distinctive coccus shape of internalized

fluorescent bacteria was also lost as a result of subsequent degradation (Fig. 1A). To

determine whether this autophagic response is not specific to the capsule-negative

D39 Δcps strain, we also analyzed the infection with the parental encapsulated wild-

type (D39 WT) S. pneumoniae and observed comparable levels of the autophagic

machinery activation despite less effective phagocytosis of D39 WT (Fig. S1A-B).

Interestingly, the observed high levels of the Lc3-association response to S.

pneumoniae infection within zebrafish macrophages contrasts with the response to

other Gram-positive bacterium - Staphylococcus aureus [25], where the Lc3 response

occurs less frequently.

Since we have previously shown that the vast majority of unencapsulated

pneumococci are internalized by macrophages [9], we made an assumption that the

observed Lc3-positive infected phagocytes are indeed macrophages. Nevertheless,

to confirm the identity of these cells we crossed the CMV:GFP-Lc3 zebrafish with the

Tg(mpeg1:mCherryF)^{ump2} transgenic line (hereafter called mpeg:mCherry-F), where

the membrane-localized mCherry reporter is expressed in zebrafish macrophages

[30]. The resulting dual transgenic larvae were intravenously infected with Alexa Fluor

647-labelled pneumococci. Microscopy analysis confirmed that out of 63 observed

phagocytes evoking the Lc3-mediated response to pneumococci, 62 were actually

macrophages (98.4%) (Fig. 1C). Altogether, these results demonstrate that larval

zebrafish serve as a suitable in vivo model to study intramacrophage autophagic

response to pneumococci.

Genetic inhibition of autophagy impairs pneumococcal killing

Having identified the Lc3 response to S. pneumoniae, we subsequently asked

whether this process is actively involved in bacterial clearance or helps infected

phagocytes to eliminate invading pneumococci by some alternative mechanism.

To this end, using CRISPR-Cas9 we knocked down the autophagy-essential gene

atg5, encoding protein that facilitates the conjugation of phosphatidylethanolamine to

the activated Lc3 [31]. The knockdown efficacy was verified by PCR and restriction

fragment length polymorphism/RFLP analysis (Fig. S2A-B). We found that the Lc3

signal surrounding the internalized bacteria was greatly reduced (Fig. 2A-B), which indicates that the observed Lc3 associations are indeed the result of an autophagy response to internalized pathogens. Next, the infected control and *atg5* knockdown larvae were homogenized and the number of bacteria within individual larvae were enumerated over the course of the infection. We found that although vast majority of bacteria were cleared by 7 hpi in both groups, and all infected control and *atg5* larvae survived until 120 hpf (Fig. S9A), the larvae with impaired autophagy machinery required more time to eliminate the pathogens, especially in the early stages of infection (Fig. 2C). The observation that *S. pneumoniae* clearance is dampened within zebrafish with an impaired autophagic response suggests a host-protective role for the process of Lc3-bacteria association. Previously, we have shown that acidification is required for *S. pneumoniae* killing by phagocytes in the zebrafish model system (Prajsnar et al., 2022). Therefore, in order to determine whether the observed autophagic response plays a role in phagosomal acidification, we injected pneumococci pre-stained with a pH-sensitive dye - pHrodo Red into control and autophagy-deficient zebrafish larvae. We observed significantly elevated levels of bacteria within acidic compartments of zebrafish macrophages of control larvae in comparison to the *atg5-* and *atg16l1*-depleted counterparts (Fig. 2D-E). Together, these data show that the autophagic response to internalized *S. pneumoniae* enhances the pace of pneumococcal killing by increasing the acidification rate of the bacteria-containing vesicles.

Inhibition of reactive oxygen species (ROS) results in partial reduction of Lc3 associations and impaired bacterial killing

As we observed high levels of Lc3-positive vesicles within pneumococcus-infected macrophages, we sought to elucidate the molecular mechanism responsible for their formation. It is generally accepted that LC3 association with internalized pathogens can be mediated by either xenophagy or non-canonical autophagic pathways such as LAP. To determine which pathway is primarily responsible in our model, we manipulated phagosomal ROS production, as its inhibition prevents the LAP response in zebrafish to several pathogens [24,25], and recent findings suggest that ROS-dependent LAP is primarily responsible for pneumococcal clearance in BMDMs *in vitro* [26]. Using a previously validated morpholino-modified antisense oligonucleotide method [32], we inhibited the expression of *cyba/p22phox*

(cytochrome b-245 alpha chain), a membrane-bound subunit of phagocyte NOX.

While the morpholino-based targeting of *cyba* was previously shown to effectively

inhibit Lc3 assembly on phagosome membranes during infection caused by either

Gram-positive or negative bacteria [24,25], Fig. S1C-D), we surprisingly found that

loss of *cyba* resulted in only partial reduction of Lc3 associations (Fig. 3A-B). The

same Lc3 association pattern was observed in response to capsule-positive (D39

WT) pneumococci suggesting no role of pneumococcal capsule in autophagy

induction (Fig. S1A-B).

In agreement with this genetic inactivation, chemical inhibition of NOX activity with

diphenyleneiodonium (DPI) also showed a similar pattern of incomplete suppression

of Lc3 aggregation with internalized pneumococci (Fig. S3). Another protein

responsible for LAP is Rubicon, which is part of the class III phosphatidylinositol 3-

kinase/PtdIns3K complex but also activates NOX promoting LAP [22,33]. Therefore,

we knocked down *rubcn* (the knockdown efficacy was validated by qPCR followed by

melt curve analysis for each targeted locus) and observed a significant decrease in

Lc3 associations, but again, a large fraction of engulfed pneumococci were

sequestered in LC3-positive vesicles (Fig. 3D-E) suggesting the existence of another

autophagic pathway. To assess the function of LAP in bacterial killing, we

enumerated the bacterial load in control and LAP-deficient infected larvae

concentrating at early stages of infection as these timepoints showed significant

differences when global autophagy was inhibited (Fig. 2C). Both *cyba* and *rubcn*

knockdowns had significantly higher colony-forming units (CFU) counts at 4 hpi than

in control siblings (Fig 3C,F). Taken together, these findings suggest that ROS- and

Rubcn-dependent LAP occurs in response to pneumococci, plays a host-protective

role during infection, but contributes only to a part of the autophagic response to

pneumococci in zebrafish macrophages.

Heat-inactivated or pneumolysin-deficient* S. pneumoniae *induce a diminished

autophagic response within macrophages

Having established that LAP occurs in response to *S. pneumoniae* infection, we next

tested whether the internalized bacterial cells need to be alive, and possibly secreting

virulence factors, to induce the observed autophagic response.

First, to determine whether live *S. pneumoniae* cells are required to induce an Lc3-
mediated response, larvae were injected with heat-killed bacterial cells. We observed
a significant decrease in Lc3 associations with pre-killed bacteria (Fig. 4A-B). To
further investigate whether LAP is involved in the response to heat-inactivated
pneumococci, we injected *cyba* morphants with heat-killed *S. pneumoniae*. This
combination led to an even more pronounced decrease in Lc3 recruitment to
engulfed bacteria (Fig. 4A-B) indicating that LAP is evoked regardless of
pneumococcal viability. Having shown that live pneumococci are required to fully
induce the autophagic response in zebrafish macrophages, we speculated whether
specific extracellular virulence factors play a role in this process. It has been
demonstrated that the induction of the autophagic response to internalized
pneumococci by professional and non-professional phagocytes *in vitro* requires Ply
(Inomata et al., 2020; Ogawa et al., 2018, 2020).To test whether Ply is responsible
for triggering autophagy in zebrafish macrophages, we first generated a *S.*
pneumoniae strain in which the ply gene has been removed (*Δply*) and verified the
absence of Ply protein expression by immunoblotting (Fig. 4C). We observed
significantly less Lc3 associations in phagocytes when *S. pneumoniae* cells lack Ply
compared to the control strain (Fig. 4D-E). It is worth noting that both heat-killed and
Ply-deficient bacteria were still able to evoke a partial autophagic response (Fig. 4),
similar to what we observed in ROS-depleted larvae (Fig. 3). These observations
prompted us to perform an experiment where *cyba* morphants were injected with Ply-
negative pneumococci. Strikingly, this combination led to a near-complete loss of Lc3
recruitment to internalized pneumococci (Fig. 4F-G), indicating that pneumolysin
induces another autophagic pathway. Importantly, Ply-deficient pneumococci were
phagocytosed by macrophages as effective as the parental strain (Fig. S9A-B),
indicating that diminished autophagic response was not due to fewer intracellular
bacteria. Interestingly, the bacteria without pneumolysin were also eliminated
significantly faster by infected larvae than Ply-positive *S. pneumoniae* (Fig. S9C) as
previously shown in a mouse model [34]. Additionally, we determined the role of
pneumolysin in evoking inflammatory response in infected phagocytes. Zebrafish
larvae of *il1b:GFP* [35] and *tnfα:GFP* [36] transgenic reporter lines infected with
pneumolysin-deficient bacteria displayed higher expression of both pro-inflammatory
cytokines in comparison to the parental strain (Fig. S9D-G). This stays in line with a
recent report describing that pneumolysin can suppress the macrophage

pro-inflammatory activation, contributes to immune evasion and bacterial persistence
by modulating host immune responses [34].

Overall, our observations confirm the role of pneumolysin in evoking an autophagic
response in an in vivo system and importantly this pneumolysin-evoked response is
independent of LAP.

Lack of xenophagy components does not affect the autophagic response to
pneumococci

We considered that xenophagy might be the Ply-mediated autophagy pathway, which
complements the ROS-dependent LAP response. During xenophagy, ruptured
phagosomes or cytosolic bacteria are detected and labeled with ubiquitin [20,37–39].
Furthermore, the expression of a pore forming toxin, ESAT-6, is required for inducing
xenophagy in infection with mycobacteria [40]. To elucidate whether xenophagy is
responsible for the pneumolysin-evoked and LAP-independent Lc3 response, we first
sought to detect ubiquitinated bacteria. However, when we performed
immunostaining for ubiquitin at 2 hpi, we observed ubiquitin staining in only around
5% of phagocytes containing Lc3-decorated bacteria, suggesting little involvement of
pneumococcal ubiquitination within zebrafish phagocytes (Fig. S4).

Next, we took advantage of a previously validated double $sqstm1^{-/-}$ $optn^{-/-}$ mutant
zebrafish line [39,41], lacking two key SLRs, Sqstm1/p62 (sequestosome 1) and
optineurin (Otpn), whose contribution to xenophagy is firmly established and
evolutionary conserve [42]. Using the CMV:GFP-Lc3 transgenic background, our
confocal analysis revealed a comparable number of Lc3-positive infected phagocyted
in the double mutant and wild-type larvae (Fig. 5A-B). Consistently, the kinetics of
bacterial clearance remained virtually unchanged in Sqstm1- and Optn-deficient
larvae when compared to the wild-type siblings (Fig. 5C). We also infected $sqstm1^{-/-}$
and $optn^{-/-}$ single mutants, and as expected, no differences in Lc3 formation or
pneumococcal killing ability was observed (Fig. S5). Next, we reasoned that
xenophagy might be triggered within *S. pneumoniae*-infected macrophages under
conditions when ROS-dependent LAP is inhibited. To address this possibility, we
knocked down *cyba* in $sqstm1^{-/-}$ larvae, but found a similar reduction in Lc3
associations within infected phagocytes as those caused by ROS inhibition alone,
without any further reductions (Fig. S6).

To exclude potential involvement of the other SLRs such as CALCOCO2 or TAXBP1,

we confirmed the lack of xenophagy in pneumococci-infected macrophages by

targeting *atg13*, a component of the ULK1 complex that is required for canonical

autophagy pathways including xenophagy, but dispensable for LAP [43]. Under

conditions of *atg13* CRISPR-mediated knockdown [28], we saw no reduction in Lc3

association with internalized pneumococci in *atg13* knockdown larvae (Fig 5D-E),

confirming that the Lc3 response to pneumococci is largely independent of

xenophagy.

Overall, these results indicate no apparent role of xenophagy in the autophagic

response to pneumococci in zebrafish macrophages suggesting the existence of

another non-canonical autophagic mechanism evoked during the *S. pneumoniae*

infection that functions alongside LAP and is triggered by pneumolysin.

A *tecpr1a*-mediated pneumolysin-dependent autophagic pathway occurs in

pneumococcus-infected macrophages

In pursuit of defining the hypothesized alternative pneumolysin-dependent non-

canonical autophagic route (Fig. 4), we speculated that a recently described CASM

pathway called sphingomyelin-TECPR1-induced LC3 lipidation (STIL), could be

involved [14]. This response is triggered upon sphingomyelin translocation from the

luminal into the cytosolic leaflet of the phagosomal membrane induced by bacterial

toxins such as listeriolysin O, from *Listeria monocytogenes*. It has been postulated

that cytosolic sphingomyelin exposure is an early indicator of vesicle damage [44].

Importantly, *S. pneumoniae* has been recently shown to induce sphingomyelin

exposure to the cytoplasm in phagosomal membranes of infected BMDMs (Shizukuishi

et al., 2024). In STIL, the exposed sphingomyelin is recognized by a protein called

TECPR1 (tectonin beta-propeller repeat containing 1) to induce single membrane

LC3 lipidation [46]. Interestingly, TECPR1 provides E3-ligase-like activity to the

ATG12–ATG5 complex, which is essential for the conjugation of LC3 to damaged

phagosomes in the absence of ATG16L1 [47]. The zebrafish genome contains two

paralogs of *TECPR1* called *tecpr1a* and *tecpr1b* (Fig. 6). We performed an alignment

of both zebrafish proteins to the human ortholog and we found profoundly higher

amino acid similarity between TECPR1 and Tecpr1a in comparison to Tecpr1b,

especially within the ATG5-interacting region (AIR) which appears to be incomplete in

Tecpr1b (Fig. 6A-B). In addition, using the Zebrafish Blood Atlas web tool with single
cell RNAseq gene expression profile of adult zebrafish hematopoietic cells [48], we
found that *tecpr1a*, but not *tecpr1b*, is highly expressed in zebrafish leukocytes (Fig.
6C). Based on this analysis, we hypothesized the Tecpr1a variant to be functional in
zebrafish, but nevertheless, we knocked down both zebrafish paralogs, *tecpr1a* and
tecpr1b, using the CRISPR-Cas9 approach and positively verified their knockdown
efficiency (Fig. S2D-E). Interestingly, when challenged by pneumococci, only *tecpr1a*
and the *tecpr1a tecpr1b* double knockdown, but not *tecpr1b* knockdown larvae
showed partial reduction of Lc3 decoration within infected macrophages (Fig. 6D-E).
This result suggests that *tecpr1a* is indeed the functional ortholog of human TECPR1,
confirming our bioinformatic analysis (Fig. 6A-C). In order to verify the pneumolysin-
dependency of the observed *tecpr1a*-mediated Lc3 decoration, we infected the
tecpr1a knockdown larvae with the pneumolysin-deficient Δ*ply* strain. We found no
additive effect in the reduction of Lc3 associations (Fig. 6F-G), indicating both
Tecpr1a and Ply act on the same pathway. To provide further evidence for the
involvement of the STIL pathway, we checked if the LC3 response to pneumococci
would be independent of ATG16I1, which is not required for STIL [14,47]. Knockdown
of *atg16l1* resulted in a significant reduction of Lc3 associations, which was more
pronounced than the decrease observed following *cyba* knockdown (Fig. S7D–E),
and impaired bacteria clearance (Fig. S7A-C). Although some Lc3 signal remained,
these results demonstrate the presence of a largely atg16l1-independent autophagic
response to pneumococci in zebrafish macrophages, consistent with the
requirements of the STIL pathway.

Upon discovery of this pathway in pneumococcus-infected macrophages, we set out
to determine the role of STIL in pneumococcal killing. The results showed that
Tecpr1a-depteted larvae clear the intravenously-administered pneumococci slower
than their wild-type siblings. (Fig 6H). Together, these results demonstrate a host-
protective role of a non-canonical pathway called STIL that has not been previously
been implicated in pneumococcal infection.

LAP and STIL independently initiate autophagic pathways in response to
pneumococci

Our data suggests the occurrence of two simultaneous non-canonical autophagic
pathways of different nature in response to pneumococci: LAP and STIL, which are
both part of the recently recognized and rapidly expanding family of CASM pathways,
where LC3/Atg8 is conjugated to single membranes of phagosomes and other
organelles [13,14]. In order to confirm the coexistence of these phenomena we
knocked down the two specific factors responsible for each of them: *cyba* for LAP
and *tecpr1a* for STIL. When both genes were knocked down we observed a near
complete loss of Lc3 associations (Fig 7A-B). Notably, this effect of double *cyba* and
tecpr1a knockdown was substantially pronounced than the partial reductions of Lc3
decorations that we previously observed in separate knockdowns of *cyba* (Fig 3A-B)
or *tecpr1a* (Fig. 6 D-E). This additive effect of *cyba* and *tecpr1a* gene silencing
suggests that indeed we observe two separate autophagic pathways evoked in
response to internalized pneumococci. Finally, to compare the contributions of LAP
and STIL to pneumococcal killing, we determined the level of bacteria in both single
and double *cyba tecpr1a* knockdown. This analysis showed that bacteria were killed
at a lower pace in larvae where both LAP and STIL were inhibited compared with
either of the single knockdown counterparts (Fig. 7C). Collectively, these results
confirm the coexistence of LAP and STIL pathways, which both serve host-protective
roles and act independently on intracellular pneumococci.

Another Ply- and Atg16l1-dependent, ROS- and Tecpr1a-independent pathway
occurs in response to pneumococcal infection

Although simultaneous knockdown of *cyba* and *tecpr1a* resulted in a strong reduction
of Lc3 lipidation, nearly 20% of infected macrophages remained to exhibit Lc3-
positive bacteria-containing vesicles. This suggested the existence of a third, distinct
autophagic pathway which could be ROS-independent [21]. To investigate this further,
a triple knockdown targeting *cyba*, *tecpr1a*, and *atg16l1* was performed. This resulted
in a near-complete loss of Lc3 recruitment, reaching level comparable that observed
in *atg5*-deficient larvae, were Lc3 lipidation was fully abolished (Fig. 8). Similar to
atg5 CRISPants, the triple knockdown did not affect the survival of larvae upon
infection (Fig. S9B). Additionally, to validate the role of pneumolysin in this context,
the *cyba tecpr1a* double knockdown larvae were infected with Ply-deficient
pneumococci. In this situation, the Lc3 lipidation was also reduced to nearly

undetectable levels, indicating that the observed remaining Lc3 signal in the double

knockdown background was dependent on this pneumococcal toxin (Fig. 8).

All these findings confirmed that the Lc3-positive vesicles observed in the double

knockdown derived from a separate, non-canonical autophagy mechanism

dependent on both pneumolysin and Atg16l1, but independent of ROS and act in

parallel with LAP and STIL, contributing to the Lc3 response in infected

macrophages.

Discussion

Autophagy-related mechanisms, both canonical and non-canonical, are considered

to be an important component of host defense against various infections, acting

primarily as a protective mechanism, and therefore could be exploited in designing

novel therapies [49]. In this study, we aimed to investigate the role of autophagy in the

innate immune response to *Streptococcus pneumoniae* systemic infection using a

zebrafish larval model. We found that the great majority of infected macrophages

contained bacterial clusters surrounded by Lc3 at the early stage of infection (2 hpi)

confirming involvement of autophagy in the host response to pneumococci.

Importantly, both capsulated and unencapsulated strains, once internalized, were

found to effectively activate Lc3 recruitment to pneumococcus-containing vesicles.

The LC3 conjugation requires ATG5 which is involved in the covalent attachment of

LC3 to the phospholipids of the membrane which is an essential step in the

autophagy process [19,31]. By diminishing the autophagic response through

knockdown of *atg5* we observed impaired bacterial clearance in our model. This

emphasized the significance of autophagy in facilitating the clearance of intracellular

pneumococci by zebrafish macrophages and prompted us to investigate which

autophagy-related pathway is responsible for this response. We took a combinatory

approach by targeting pathogen and host components, both separately and in

combination to determine the autophagic pathways present in *S. pneumoniae*-

infected macrophages.

The antimicrobial, host-protective function of LC3-associated phagocytosis (LAP) has

been reported in several studies, demonstrating its effective role in bacterial

degradation [11,50,51]. The significance of this pathway had been recognized well

before its inclusion in the recently defined CASM family and it is supported by studies

analyzing microbial survival mechanisms evolved to evade or escape this process to

prolong their survival [52–54]. Our study demonstrates that LAP occurs in zebrafish

macrophages upon internalizing pneumococci. A significant but partial reduction in

Lc3 recruitment and slower bacterial clearance induced by knocking down *cyba* or

rubcn, genes required for LAP, underscores the importance of this pathway in the

host defense against pneumococci. Importantly, our data confirms *in vitro*-derived

results where LAP also plays a protective role in pneumococci-infected BMDMs [26].

Interestingly, Inomata and colleagues have also reported incomplete inhibition of LC3

recruitment in *Rubcn* or *Cybb/Nox2* knockout macrophages, which additionally

prompted us to search for another autophagic pathway present in response to

pneumococci. Moreover, in accordance with Inomata and others, we observed

significant decrease of Lc3 associations to pneumococci in the absence of

pneumolysin and we did not find any evidence of selective autophagy involvement in

S. pneumoniae-infected macrophages *in vivo*.

Indeed, our results show that Lc3 recruitment triggered by pneumococci upon

internalization by macrophages is partially-dependent on bacterial pneumolysin. By

targeting components of different autophagy pathways, we were able to match

pathogen-derived pneumolysin with particular facets of the autophagic response.

Inomata *et al.* posited that pneumolysin induces LAP, however no combinatory

approaches with Δ*ply* mutant in LAP-deficient macrophages have been performed

[26]. By testing wild-type and Δ*ply* mutant pneumococcal strains against pathway-

specific host genes in the zebrafish model, we here show that Ply activates another

non-canonical autophagy pathway, which acts independently of ROS and Atg16l1,

but is dependent on Tecpr1a.

TECPR1 was initially shown to play a vital role in promoting selective autophagy

through interaction with WIPI2 [55]. However, recent studies have revealed an

additional function of TECPR1 via its DysF domains [46,56], where this protein

recognizes sphingomyelin present on the outer layer of disturbed phagosomal or

lysosomal membranes to induce CASM in a process called STIL. Interestingly, this

alternative non-canonical pathway challenges the traditional view that ATG16L1 is the

sole E3-like enzyme responsible for lipidation of LC3, highlighting the versatility and

complexity of autophagic regulation [57].

We have found in zebrafish that a knockdown of *tecpr1a*, a likely ortholog of

TECPR1, leads to a partial loss of Lc3 decoration of internalized pneumolysin-

positive *S. pneumoniae*. When pneumolysin mutant bacteria were administered, no

additional effect of *tecpr1a* depletion was observed suggesting the role of

pneumolysin in perturbing the phagosomal membrane and Tecpr1a in phagosomal

membrane Lc3 decoration. Importantly, in the *cyba tecpr1a* double knockdown we

observed a near-complete loss of Lc3 associations. Moreover, in *cyba* knockdown

larvae injected with Δ*ply* pneumococci, no Lc3 decoration was present, indicating the

occurrence of two independent, differentially induced non-canonical pathways,

namely LAP and STIL. A similar phenomenon was previously observed *in vitro* in

BMDMs infected with *L. monocytogenes* evoking two distinct non-canonical

autophagy pathways [29]. One of them is LAP and the other ROS-independent,

listeriolysin O-mediated, was coined by the authors as pore-forming toxin-induced

non-canonical autophagy (PINCA). At the time, the authors did not know the factors

responsible for PINCA, therefore they were unable to determine its function in *L.*

monocytogenes infection. It is conceivable that PINCA observed by Glushko and

colleagues [29] was in fact STIL.

Another similar case of parallel pathways has been observed in MEFs *in vitro* at early

stages of infection with pneumolysin-expressing *S. pneumoniae*, where LC3

lipidation occurred independently of RB1CC1/FIP200 (a member of the ULK1

complex required for canonical autophagy) and also independently of ROS leading to

LAPosome-like vacuoles formation. However, since the observed process requires

ATG16L1 [21], it appears to be another form of CASM, distinct from STIL. Indeed, our

data revealed the existence of an equivalent non-canonical autophagy pathway in

macrophages, in addition to LAP and STIL. This another CASM depended on both

ATG16L1 and pneumolysin, but occurred independently of ROS, matching the

mechanism described in non-professional phagocytes [21]. Its identification not only

clarifies that Lc3 recruitment was not completely abolished in the absence of LAP

and STIL components alone, but also underline the complexity of host defense

mechanisms involving non-canonical autophagy.

Interestingly, Ogawa and colleagues have also reported a hierarchical autophagic

response in which LAPosomes undergo Sqstm1- and Optn-mediated xenophagy at

later stages of infection [21,58]. However, no such phenomenon involving xenophagy

factors was observed in macrophages, neither in our study nor in that of others [26].

The various mechanisms evoked by pneumococci infection in different cells are most

likely caused by differences between professional and non-professional phagocytes,

such as the higher levels of ROS production of macrophages compared to fibroblasts

or the degree of phagosomal acidification [26].

The observed differences, particularly the lack of xenophagy activation in

macrophages, are most likely caused by differences between professional and non-

professional phagocytes, such as the higher levels of ROS production of

macrophages compared to fibroblasts or the degree of phagosomal acidification [26].

We propose the following model of the fate of pneumococci within macrophages (Fig.

9). The bacteria are internalized by macrophages and trapped within Lc3-decorated

vesicles, which could be formed by three distinct mechanisms depending on the

nature of phagosomal membrane perturbation. Florey and others have hypothesized

that any process that promotes osmotic imbalances within the endolysosomal system

would have the potential to activate LAP-like LC3 lipidation [59]. Therefore, we

believe that in our model, the ROS-induced osmotic imbalance in phagosomes

containing pneumococci leads to LAP. Alternatively, pneumolysin produced by

bacteria trapped in phagosomes inflicts other forms of membrane perturbation,

leading to either ROS-independent Lc3 lipidation via unknown mechanism or

sphingomyelin presence on the outer side of phagosomal envelope and resulting in

STIL. This last scenario is additionally supported by a recent report demonstrating

induction of sphingomyelin exposure to the cytoplasm in phagosomal membranes of

S. pneumoniae-infected BMDMs [60]. Both LAP and STIL enhance pneumococcal

killing within zebrafish macrophages as knockdown of pathway-specific components

slowed down the inactivation of bacteria *in vivo* but did not completely impair it. This

suggests that classical phagosome maturation still occurs in the absence of these

pathways and the internalized pneumococci are still trafficked to increasingly acidified

vesicles, eventually leading to their degradation.

Collectively, the presented results markedly advance our knowledge of the innate

immune response to pneumococcal infections. By elucidating the abundant activation

of various CASM pathways, and probably negligible role of xenophagy in

professional phagocytes our study highlights potential therapeutic targets that could

be exploited to enhance host defense mechanisms. Therefore, in addition to LAP, we

believe it is important to address other non-canonical autophagy pathways in order to

control pneumococcal infection. This may offer novel ways for treatment, especially in

cases where traditional antibiotic-based therapies are insufficient due to rising

bacterial resistance, opening the door to innovative immunomodulatory approaches

that could improve clinical outcomes in pneumococcal infections.

Materials and Methods

Ethics statement

All zebrafish experiments were conducted in accordance with the European

Community Council Directive 2010/63/EU for the Care and Use of Laboratory

Animals of Sept. 22, 2010 (Chapter 1, Article 1 no.3) and National Journal of Law act

of Jan. 15, 2015, for Protection of animals used for scientific or educational purposes

(Chapter 1, Article 2 no.1). All experiments with zebrafish were performed on

embryos/larvae up to 120 h post fertilization and were following ARRIVE guidelines.

The Jagiellonian University Zebrafish Core Facility (ZCF) is a licensed breeding and

research facility (District Veterinary Inspectorate in Krakow registry; Ministry of

Science and Higher Education record no. 022 and 0057).

Zebrafish husbandry

Adult zebrafish were maintained by Zebrafish Core Facility sta□ at the Jagiellonian

University in accordance with the standard protocol and international guidelines

specified by the EU Animal Protective Directive 2010/63/EU. Fish were kept in a

continuous recirculating closed system aquarium with a light/dark cycle of 14/10 h at

28°C. Larvae were incubated in standard E3 medium (5 mM NaCl, 0.17 mM KCl,

0.33 mM $CaCl_2$, 0.33 mM $MgSO_4$) at 28.5°C according to standard protocols [61]. All

zebrafish lines used in this study are listed in Table S1.

Bacterial cultures and infection experiments

All *Streptococcus pneumoniae* strains used in this study were derived from serotype

2 D39 strain [62] and are listed in Table S1. Pneumococci were grown on prepared

Tryptic Soy Agar 5% sheep blood plates (TSAII 5% SB; Becton Dickinson [BD],

B21261X) at 37°C in an atmosphere containing 5% CO_2. Liquid cultures were

maintained in Todd-Hewitt Broth (BD, 249240) supplemented with 0.5% yeast extract

(Gibco, 212750) at 37°C in an atmosphere containing 5% CO_2. Bacteria were

harvested at mid-log phase (OD_{600} between 0.2 - 0.3) and subsequently centrifuged

(5000 g, 10 min) and immediately resuspended in phosphate-buffered saline

(Bioshop, PBS404.100) to obtain the desired concentration of approximately 1600

CFU/nl. Such prepared bacteria were microinjected intravenously into 36 hpf

zebrafish larvae as described previously [63]. Briefly, larvae were anaesthetized with

0.02% buffered tricaine (Sigma-Aldrich, A5040) and then transferred onto 3% w:v

methylcellulose (Sigma-Aldrich, M7027) and injected individually with 1 nl of bacteria

inoculum using microcapillary pipettes.

Staining of bacteria with pHrodo-Red and Alexa Fluor 647 S-ester dyes

579   The pHrodo Red (ThermoFisher, P36600) and Alexa Fluor 647 (ThermoFisher,

580   A37573) succinimidyl-ester dyes were dissolved in DMSO (Sigma-Aldrich, D8418) to

581   the final concentrations of 2.5 mM and 8 mM, respectively. In the indicated

582   experiments, *S. pneumoniae* strains were stained as previously described [25].

Generation of Δply *mutant in S. pneumoniae*

The *S. pneumoniae* Δ*ply* deletion strain was generated using linear PCR fragments.

Briefly, two ~1-kb flanking regions of each gene were amplified and an antibiotic

resistance marker placed between them using isothermal assembly. Assembled PCR

products were transformed directly into *S. pneumoniae*. In all cases, deletion primers

were given the typical name: "gene- designation"_5FLANK_F/R for 5′ regions and

"gene-designation"_3FLANK_F/R for 3′ regions, antibiotic markers were amplified

from Δ*bgaA* strains [64] using the AntibioticMarker_F/R primers. Successful gene

deletion strains were confirmed using diagnostic PCR. All bacterial strains and

oligonucleotides used in this study are listed in Table S2.

Morpholino- and CRISPR/Cas9-Mediated Knockdown

The *cyba* morpholino oligonucleotides [32] (Gene Tools, 5'-

ATCATAGCATGTAAGGATACATCCC-3') were dissolved in MilliQ water to obtain the

required concentrations of 1 mM. The 1 nl volume of morpholino was injected into the

yolk of 1–2 cell stage zebrafish embryos using a microinjector. Standard control

morpholino (Gene Tools, 5'-CCTCTTACCTCAGTTACAATTTATA-3') was used as a

negative control. For CRISPR-Cas9-mediated *atg16l1, rubcn*, *tecpr1a* and *tecpr1b*

knockdown, the online web tool CHOPCHOP was used to design a gene-specific

guide CRISPR of the following sequences listed in Table S3. The synthetic guide

RNA consisting of gene-specific CRISPR RNA (crRNA, Merck, VC40002) and

transactivating RNAs (tracrRNA, Merck, TRACRRNA05N) in combination with Cas9

nuclease protein (NEB, M0386M) was used for gene editing. TracrRNA and crRNA

were resuspended to a concentration of 50 µM in nuclease-free water. The gRNA-

Cas9 complexes were assembled before injections using a 1:1:1 ratio of

crRNA:tracrRNA:Cas9 nuclease with the final concentrations of 16.6 µM, 16.6 µM,

and 6.6 µM, respectively. As controls, only Cas9 + tracrRNA were injected without

crRNA. Knockdown verification was performed by the PCR use of primers with the

sequences presented in the Table S2. For *atg5* knockdown, a previously published

single guide RNA (sgRNA, Merck, VC40003) was used. Briefly, 1 µl 25 µM sgRNA

was gently mixed with 1 µl 25 µM Cas9 and 3 µl H2O with the final concentration of 5

µM and 2 nl was injected. The mutagenesis efficacy for *atg5* was determined by

restriction fragment length polymorphism analysis using the MslI enzyme as

previously described [65].

DPI treatment

617   At 1 h before infection, larvae were bath-treated with 100 µM DPI (Sigma-Aldrich,

618   D2926) in E3 medium. The larvae were then infected and kept in DPI solution until

619   fixation. DMSO at 0.1% was used as vehicle control.

Immunostaining

Larvae were fixed with 4% paraformaldehyde(PFA, ThermoFisher, AAJ19943K2)

supplemented with 0.5% Triton X-100 (Sigma-Aldrich, X-100) in PBS and incubated

overnight at 4°C. Subsequently, fixed fish were washed four times for 20 min in PBS

with 0.5% Triton X-100 and 1% DMSO (PBS-DTx). Next, the larvae were digested in

10 µg/ml proteinase K (Sigma-Aldrich, 1073930010) for 10 min at 37°C. Then, larvae

were quickly washed three timed for 5 minutes and blocked with PBS-DTx containing

1% bovine serum albumin (BSA; Sigma-Aldrich, A9418) for 1h at room temperature

and incubated overnight at 4°C in mono-and polyubiquitinated conjugates

recombinant mouse monoclonal antibody (Enzo Life Siences, UBCJ2) diluted in the

blocking buffer 1:250. Next, larvae were washed in blocking solution four times for 20

minutes and incubated at room temperature for 2 h in blocking solution of secondary

goat anti-mouse Alexa Fluor 405 antibody (Invitrogen, A-31553) in 1:300 followed

with four times washes in PBS with 0.5% Triton-X100.

Microscopy Imaging and Analysis

Live anaesthetized or paraformaldehyde-fixed larvae were immersed in 1% (w/v) low-

melting-point agarose solution in E3 medium and mounted laterally on a dish with a

glass bottom. Images of the common cardinal vein (top of Fig.1) were acquired using

a Zeiss LSM900 Airyscan 2 confocal laser scanning microscope with ZEISS ZEN 3.3

software using the Plan-Apochromat 20x/0.8, C-Apochromat 40x/NA 1.2 W or C-

Apochromat 63x/1.20 W objectives. For quantification of the autophagic response

within infected phagocytes, for each larva, a total number of observable infected

phagocytes were manually determined through the z-stacks of images. Among these

total observable infected phagocytes number of infected phagocytes with GFP-Lc3

signals were enumerated, and the percentage of Lc3-positive phagocytes over total

observable phagocytes was determined for each larva. Maximum projections were

used for representative images. No non-linear normalizations were performed. For

quantification of acidification, the pHrodo Red fluorescence intensity, ZEISS ZEN 3.3

software was used. Quantification of bacterial internalization was performed as

previously described [66].

Determination of In Vivo Bacterial Counts

At various times postinfection, approximately 8-12 anesthetized zebrafish larvae

were individually transferred with 100 µl of E3 medium into 0.5-ml Screw cap micro

tubes (Sarstedt, 72.730.100) containing 2-mm Zirconia beads (Biospec,

11079124ZX) and homogenized using a FastPrep-24 5G homogenizer (MP

Biomedicals, 6005500). The homogenates were spotted on TSA plates containing

5% defibrinated sheep blood to determine pneumococcal CFU numbers. The limit of

detection was 10 CFU per larva.

Western Blot

659   For each immunoblot, 10 µl of each whole-cell lysate was run on a 12% SDS PAGE

660   gel (40 mA, ≈55 min) and transferred onto PVDF membranes (80 min, 50 V).

661   Membranes were blocked with 5% skimmed milk in 1x PBS 0.5% Tween 20 (Sigma-

Aldrich, 11332465001) for 1 h (gently rocking, room temperature). Membranes were washed three times with 1x PBS, 0.05% Tween 20, and incubated with primary antibody overnight gently rocking (4°C, 3% BSA in 1x PBS, 0.05% Tween 20, 0.02% sodium azide). For detection of Ply, a mouse anti-Ply antibody was used at a final concentration of 1:1000 (mouse mAb to pneumolysin [PLY-4]; Abcam, AB71810). The blocked PVDF membranes were washed three times with 1x PBS, 0.05% Tween 20. The secondary antibody (goat anti-mouse IgG conjugated to horseradish peroxidase [Bio-Rad, 172-1011]) was used at 1:5000 and applied in 1x PBS with 3% BSA for 1 h at room temperature with gentle shaking. Resulting blots were washed four times in 1x PBS, 0.05% Tween 20 and visualized using the Clarity Western ECL Substrate kit (Bio-Rad, 170-5061) according to the manufacturer's instructions. Blots were imaged using a Syngene G:BOX Chemi XX9 image quantification device combined with the GeneSys software (v1.6.7.0) using the chemiluminescent detection tools.

Statistics

GraphPad Prism 10 was used for statistical analysis. Quantifications of percent Lc3-positive phagocytes, CFU counts, bacterial uptake, or pHrodo Red and GFP fluorescence intensity were determined for significance with unpaired parametric t-test for 2 groups and with ANOVA for multiple groups, with Šidák correction for multiple comparisons. Survival curves were analyzed with Log rank (Mantel-Cox) test. Statistical significance was assumed at *P*-values below 0.05.

Acknowledgements

This work was supported by a National Science Centre of Poland under Sonata Bis 9 project (Grant number: 2019/34/E/NZ6/00137) awarded to T.K.P. and a Medical Research Council (MRC) New Investigator Research Grant (Grant number: MR/S009280/1) awarded to A.K.F.

T.K.P. was funded by Polish National Agency for Academic Exchange under Polish Returns 2019 project (Grant number: PPN/PPO/2019/1/00029/U/0001) and T.C.S. was funded by MRC DiMeN DTP studentship.

The establishment of a Zebrafish Core Facility at the Department of Evolutionary Immunology has been supported by a grant from the Priority Research Area BioS under the Strategic Program Excellence Initiative at Jagiellonian University.

Disclosure statement

The authors have declared no conflict of interests.

References

[1]   Weiser JN, Ferreira DM, Paton JC. Streptococcus pneumoniae: Transmission, colonization and invasion. Nat Rev Microbiol 2018;16:355–67. https://doi.org/10.1038/s41579-018-0001-8.

[2]   Ikuta KS, Swetschinski LR, Aguilar GR, et al. Global mortality associated with 33 bacterial pathogens in 2019: a systematic analysis for the Global Burden of Disease Study 2019. The Lancet 2022;400:2221–48. https://doi.org/10.1016/S0140-6736(22)02185-7.

[3]   van der Poll T, Opal SM. Pathogenesis, treatment, and prevention of pneumococcal pneumonia. The Lancet 2009;374:1543–56. https://doi.org/10.1016/S0140-6736(09)61114-4.

[4]   Croucher NJ, Harris SR, Fraser C, et al. Rapid pneumococcal evolution in response to clinical interventions. Science (1979) 2011;331:430–4. https://doi.org/10.1126/science.1198545.

[5]   Dockrell DH, Marriott HM, Prince LR, et al. Alveolar Macrophage Apoptosis Contributes to Pneumococcal Clearance in a Resolving Model of Pulmonary Infection. The Journal of Immunology 2003;171:5380–8. https://doi.org/10.4049/jimmunol.171.10.5380.

[6]   Flannagan RS, Jaumouillé V, Grinstein S. The cell biology of phagocytosis. Annual Review of Pathology: Mechanisms of Disease 2012;7:61–98. https://doi.org/10.1146/annurev-pathol-011811-132445.

[7]   Maxson ME, Grinstein S. The vacuolar-type H+-ATPase at a glance - more than a proton pump. J Cell Sci 2014;127:4987–93. https://doi.org/10.1242/jcs.158550.

[8]   Westman J, Grinstein S. Determinants of Phagosomal pH During Host-Pathogen Interactions. Front Cell Dev Biol 2021;8. https://doi.org/10.3389/fcell.2020.624958.

[9]   Prajsnar TK, Michno BJ, Pooranachandran N, et al. Phagosomal Acidification Is Required to Kill Streptococcus pneumoniae in a Zebrafish Model. Cell Microbiol 2022;2022. https://doi.org/10.1155/2022/9429516.

[10]   Kaufmann SHE, Dorhoi A. Molecular Determinants in Phagocyte-Bacteria Interactions. Immunity 2016;44:476–91. https://doi.org/10.1016/j.immuni.2016.02.014.

[11]   Yuan J, Zhang Q, Chen S, et al. LC3-Associated Phagocytosis in Bacterial Infection. Pathogens 2022;11. https://doi.org/10.3390/pathogens11080863.

[12]   Sharma V, Verma S, Seranova E, et al. Selective autophagy and xenophagy in infection and disease. Front Cell Dev Biol 2018;6. https://doi.org/10.3389/fcell.2018.00147.

[13]   Durgan J, Florey O. Many roads lead to CASM: Diverse stimuli of noncanonical autophagy share a unifying molecular mechanism. Sci Adv 2022;8. https://doi.org/10.1126/sciadv.abo1274.

[14]   Figueras-Novoa C, Timimi L, Marcassa E, et al. Conjugation of ATG8s to single membranes at a glance. J Cell Sci 2024;137. https://doi.org/10.1242/jcs.261031.

[15]   Shaid S, Brandts CH, Serve H, et al. Ubiquitination and selective autophagy. Cell Death Differ 2013;20:21–30. https://doi.org/10.1038/cdd.2012.72.

[16] Thurston TLM, Wandel MP, Von Muhlinen N, et al. Galectin 8 targets damaged vesicles for
autophagy to defend cells against bacterial invasion. Nature 2012;482:414–8.
https://doi.org/10.1038/nature10744.

[17] Galluzzi L, Baehrecke EH, Ballabio A, et al. Molecular definitions of autophagy and related
processes. EMBO J 2017;36:1811–36. https://doi.org/10.15252/embj.201796697.

[18] Herhaus L, Dikic I. Regulation of Salmonella-host cell interactions via the ubiquitin system.
International Journal of Medical Microbiology 2018;308:176–84.
https://doi.org/10.1016/j.ijmm.2017.11.003.

[19] Melia TJ, Lystad AH, Simonsen A. Autophagosome biogenesis: From membrane growth to
closure. Journal of Cell Biology 2020;219. https://doi.org/10.1083/JCB.202002085.

[20] Ogawa M, Matsuda R, Takada N, et al. Molecular mechanisms of Streptococcus pneumoniae-
targeted autophagy via pneumolysin, Golgi-resident Rab41, and Nedd4-1-mediated K63-linked
ubiquitination. Cell Microbiol 2018;20. https://doi.org/10.1111/cmi.12846.

[21] Ogawa M, Takada N, Shizukuishi S, et al. Streptococcus pneumoniae triggers hierarchical
autophagy through reprogramming of LAPosome-like vesicles via NDP52-delocalization.
Commun Biol 2020;3. https://doi.org/10.1038/s42003-020-0753-3.

[22] Yang CS, Lee JS, Rodgers M, et al. Autophagy protein rubicon mediates phagocytic NADPH
oxidase activation in response to microbial infection or TLR stimulation. Cell Host Microbe
2012;11:264–76. https://doi.org/10.1016/j.chom.2012.01.018.

[23] Hooper KM, Jacquin E, Li T, et al. V-ATPase is a universal regulator of LC3-associated
phagocytosis and non-canonical autophagy. Journal of Cell Biology 2022;221.
https://doi.org/10.1083/jcb.202105112.

[24] Masud S, Prajsnar TK, Torraca V, et al. Macrophages target Salmonella by Lc3-associated
phagocytosis in a systemic infection model. Autophagy 2019;15:796–812.
https://doi.org/10.1080/15548627.2019.1569297.

[25] Prajsnar TK, Serba JJ, Dekker BM, et al. The autophagic response to Staphylococcus aureus
provides an intracellular niche in neutrophils. Autophagy 2021;17:888–902.
https://doi.org/10.1080/15548627.2020.1739443.

[26] Inomata M, Xu S, Chandra P, et al. Macrophage LC3-associated phagocytosis is an immune
defense against Streptococcus pneumoniae that diminishes with host aging. Proc Natl Acad Sci
U S A 2020;117. https://doi.org/10.1073/PNAS.2015368117.

[27] He C, Bartholomew CR, Zhou W, et al. Assaying autophagic activity in transgenic GFP-Lc3 and
GFP-Gabarap zebrafish embryos. Autophagy 2009;5:520–6.
https://doi.org/10.4161/auto.5.4.7768.

[28] Forn-Cuní G, Welvaarts L, Stel FM, et al. Stimulating the autophagic-lysosomal axis enhances
host defense against fungal infection in a zebrafish model of invasive Aspergillosis. Autophagy
2023;19:324–37. https://doi.org/10.1080/15548627.2022.2090727.

[29] Gluschko A, Farid A, Herb M, et al. Macrophages target Listeria monocytogenes by two
discrete non-canonical autophagy pathways. Autophagy 2022;18:1090–107.
https://doi.org/10.1080/15548627.2021.1969765.

[30]   Bernut A, Herrmann JL, Kissa K, et al. Mycobacterium abscessus cording prevents phagocytosis
772   and promotes abscess formation. Proc Natl Acad Sci U S A 2014;111.
773   https://doi.org/10.1073/pnas.1321390111.

774   [31]   Dikic I, Elazar Z. Mechanism and medical implications of mammalian autophagy. Nat Rev Mol
775   Cell Biol 2018;19:349–64. https://doi.org/10.1038/s41580-018-0003-4.

776   [32]   Niethammer P, Grabher C, Look AT, et al. A tissue-scale gradient of hydrogen peroxide
777   mediates rapid wound detection in zebrafish. Nature 2009;459:996–9.
778   https://doi.org/10.1038/nature08119.

779   [33]   Masud S, van der Burg L, Storm L, et al. Rubicon-Dependent Lc3 Recruitment to Salmonella-
780   Containing Phagosomes Is a Host Defense Mechanism Triggered Independently From Major
781   Bacterial Virulence Factors. Front Cell Infect Microbiol 2019;9.
782   https://doi.org/10.3389/fcimb.2019.00279.

783   [34]   Periselneris J, Turner CT, Ercoli G, et al. Pneumolysin suppresses the initial macrophage pro-
784   inflammatory response to Streptococcus pneumoniae. Immunology 2022;167:413–27.
785   https://doi.org/10.1111/imm.13546.

786   [35]   Ogryzko N V., Lewis A, Wilson HL, et al. Hif-1α–Induced Expression of Il-1β Protects against
787   Mycobacterial Infection in Zebrafish. The Journal of Immunology 2019;202:494–502.
788   https://doi.org/10.4049/JIMMUNOL.1801139.

789   [36]   Marjoram L, Alvers A, Deerhake ME, et al. Epigenetic control of intestinal barrier function and
790   inflammation in zebrafish. Proc Natl Acad Sci U S A 2015;112:2770–5.
791   https://doi.org/10.1073/PNAS.1424089112/SUPPL_FILE/PNAS.201424089SI.PDF.

792   [37]   Manzanillo PS, Ayres JS, Watson RO, et al. The ubiquitin ligase parkin mediates resistance to
793   intracellular pathogens. Nature 2013;501:512–6. https://doi.org/10.1038/nature12566.

794   [38]   Polajnar M, Dietz MS, Heilemann M, et al. Expanding the host cell ubiquitylation machinery
795   targeting cytosolic Salmonella . EMBO Rep 2017;18:1572–85.
796   https://doi.org/10.15252/embr.201643851.

797   [39]   Zhang R, Varela M, Vallentgoed W, et al. The selective autophagy receptors Optineurin and
798   p62 are both required for zebrafish host resistance to mycobacterial infection. PLoS Pathog
799   2019;15. https://doi.org/10.1371/journal.ppat.1007329.

800   [40]   Bell SL, Lopez KL, Cox JS, et al. Galectin-8 senses phagosomal damage and recruits selective
801   autophagy adapter tax1bp1 to control mycobacterium tuberculosis infection in macrophages.
802   MBio 2021;12. https://doi.org/10.1128/mBio.01871-20.

803   [41]   Xie J, Meijer AH. Xenophagy receptors Optn and p62 and autophagy modulator Dram1
804   independently promote the zebrafish host defense against Mycobacterium marinum. Front
805   Cell Infect Microbiol 2023;13. https://doi.org/10.3389/fcimb.2023.1331818.

806   [42]   Gubas A, Dikic I. A guide to the regulation of selective autophagy receptors. FEBS Journal
807   2022;289:75–89. https://doi.org/10.1111/febs.15824.

808   [43]   Zachari M, Ganley IG. The mammalian ULK1 complex and autophagy initiation. Essays
809   Biochem 2017;61:585. https://doi.org/10.1042/EBC20170021.

810   [44]   Ellison CJ, Kukulski W, Boyle KB, et al. Transbilayer Movement of Sphingomyelin Precedes
811   Catastrophic Breakage of Enterobacteria-Containing Vacuoles. Current Biology 2020;30:2974-
812   2983.e6. https://doi.org/10.1016/j.cub.2020.05.083.

813   [45]   Shizukuishi S, Ogawa M, Kuroda E, et al. Pneumococcal sialidase promotes bacterial survival by
814   fine-tuning of pneumolysin-mediated membrane disruption. Cell Rep 2024;43:113962.
815   https://doi.org/10.1016/j.celrep.2024.113962.

816   [46]   Boyle KB, Ellison CJ, Elliott PR, et al. TECPR1 conjugates LC3 to damaged endomembranes
817   upon detection of sphingomyelin exposure . EMBO J 2023;42.
818   https://doi.org/10.15252/embj.2022113012.

819   [47]   Wang Y, Jefferson M, Whelband M, et al. TECPR1 provides E3-ligase like activity to the ATG5-
820   ATG12 complex to conjugate LC3/ATG8 to damaged lysosomes. BioRxiv
821   2023:2023.06.24.546289. https://doi.org/10.1101/2023.06.24.546289.

822   [48]   Athanasiadis EI, Botthof JG, Andres H, et al. Single-cell RNA-sequencing uncovers
823   transcriptional states and fate decisions in haematopoiesis. Nat Commun 2017;8.
824   https://doi.org/10.1038/s41467-017-02305-6.

825   [49]   Kimmey JM, Stallings CL. Bacterial Pathogens versus Autophagy: Implications for Therapeutic
826   Interventions. Trends Mol Med 2016;22:1060–76.
827   https://doi.org/10.1016/j.molmed.2016.10.008.

828   [50]   Herb M, Gluschko A, Schramm M. LC3-associated phagocytosis - The highway to hell for
829   phagocytosed microbes. Semin Cell Dev Biol 2020;101:68–76.
830   https://doi.org/10.1016/j.semcdb.2019.04.016.

831   [51]   Grijmans BJM, van der Kooij SB, Varela M, et al. LAPped in Proof: LC3-Associated Phagocytosis
832   and the Arms Race Against Bacterial Pathogens. Front Cell Infect Microbiol 2022;11.
833   https://doi.org/10.3389/fcimb.2021.809121.

834   [52]   Hubber A, Kubori T, Coban C, et al. Bacterial secretion system skews the fate of Legionella-
835   containing vacuoles towards LC3-associated phagocytosis. Sci Rep 2017;7.
836   https://doi.org/10.1038/srep44795.

837   [53]   Köster S, Upadhyay S, Chandra P, et al. Mycobacterium tuberculosis is protected from NADPH
838   oxidase and LC3-associated phagocytosis by the LCP protein CpsA. Proc Natl Acad Sci U S A
839   2017;114:E8711–20. https://doi.org/10.1073/pnas.1707792114.

840   [54]   Mitchell G, Cheng MI, Chen C, et al. Listeria monocytogenes triggers noncanonical autophagy
841   upon phagocytosis, but avoids subsequent growth-restricting xenophagy. Proc Natl Acad Sci U
842   S A 2017;115:E210–7. https://doi.org/10.1073/pnas.1716055115.

843   [55]   Ogawa M, Yoshikawa Y, Kobayashi T, et al. A Tecpr1-dependent selective autophagy pathway
844   targets bacterial pathogens. Cell Host Microbe 2011;9:376–89.
845   https://doi.org/10.1016/j.chom.2011.04.010.

846   [56]   Kaur N, de la Ballina LR, Haukaas HS, et al. TECPR1 is activated by damage-induced
847   sphingomyelin exposure to mediate noncanonical autophagy . EMBO J 2023;42.
848   https://doi.org/10.15252/embj.2022113105.

[57] Corkery DP, Castro-Gonzalez S, Knyazeva A, et al. An ATG12-ATG5-TECPR1 E3-like complex
regulates unconventional LC3 lipidation at damaged lysosomes. EMBO Rep 2023;24.
https://doi.org/10.15252/embr.202356841.

[58] Sakuma C, Shizukuishi S, Ogawa M, et al. Individual Atg8 paralogs and a bacterial metabolite
sequentially promote hierarchical CASM-xenophagy induction and transition. Cell Rep
2024;43. https://doi.org/10.1016/j.celrep.2024.114131.

[59] Florey O, Gammoh N, Kim SE, et al. V-ATPase and osmotic imbalances activate endolysosomal
LC3 lipidation. Autophagy 2015;11:88–99. https://doi.org/10.4161/15548627.2014.984277.

[60] Shizukuishi S, Ogawa M, Akeda Y. Individual Atg8 paralogs exhibit unique properties in
streptococcus pneumoniae-induced hierarchical autophagy. Autophagy 2024.
https://doi.org/10.1080/15548627.2024.2375707.

[61] Nusslein-Volhard C, Dahm R. Zebrafish: A Practical Approach. Oxford University Press;
Illustrated Edition (November 21, 2002); 2002.

[62] Lanie JA, Ng WL, Kazmierczak KM, et al. Genome sequence of Avery's virulent serotype 2
strain D39 of Streptococcus pneumoniae and comparison with that of unencapsulated
laboratory strain R6. J Bacteriol 2007;189:38–51. https://doi.org/10.1128/JB.01148-06.

[63] Prajsnar TK, Cunliffe VT, Foster SJ, et al. A novel vertebrate model of Staphylococcus aureus
infection reveals phagocyte-dependent resistance of zebrafish to non-host specialized
pathogens. Cell Microbiol 2008;10:2312–25. https://doi.org/10.1111/j.1462-
5822.2008.01213.x.

[64] Fenton AK, Mortaji L El, Lau DTC, et al. CozE is a member of the MreCD complex that directs
cell elongation in Streptococcus pneumoniae. Nat Microbiol 2016;2.
https://doi.org/10.1038/nmicrobiol.2016.237.

[65] Chen XK, Yi ZN, Lau JJY, et al. Distinct roles of core autophagy-related genes in zebrafish
definitive hematopoiesis. Autophagy 2023;20:830–46.
https://doi.org/10.1080/15548627.2023.2274251;SUBPAGE:STRING:FULL.

[66] Salamaga B, Prajsnar TK, Jareño-Martinez A, et al. Bacterial size matters: Multiple mechanisms
controlling septum cleavage and diplococcus formation are critical for the virulence of the
opportunistic pathogen Enterococcus faecalis. PLoS Pathog 2017;13:e1006526.
https://doi.org/10.1371/JOURNAL.PPAT.1006526.

Figure Legends

Figure 1. The autophagic response occurs during systemic infection with

pneumococci. (**A**) representative confocal images of *CMV:GFP-Lc3* larvae infected

systemically with approximately 1600 CFU of mKate2-labeled D39 Δ*cps S.*

pneumoniae and fixed at 2, 4 and 7 hpi. Scale bars: 10 μm. Arrowheads indicate Lc3-

positive vacuoles containing bacteria. (**B**) Quantification of Lc3 associations with

intracellular *S. pneumoniae* within infected phagocytes of fixed *CMV:GFP-Lc3* at 2, 4

and 7 hpi. Data are shown as mean ± standard deviation (SD). Data obtained from

two independent experiments (**C**) representative confocal images of *CMV:GFP-Lc3*

mpeg:mCherry-F dual transgenic larvae infected systemically with approximately

1600 CFU of Alexa Fluor 647-stained D39 Δ*cps S. pneumoniae* and fixed at 2 hpi.

Figure 2. The autophagic response contributes to phagosomal acidification and

pneumococcal clearance. (**A**) representative confocal images of control (left panel)

and *atg5* knockdown (right panel) *CMV:GFP-Lc3* larvae infected systemically with

approximately 1600 CFU of mKate2-labeled D39 Δ*cps S. pneumoniae* and fixed at 2

hpi. Scale bars: 10 μm. (**B**) Quantification of Lc3 associations with intracellular

bacteria of control and *atg5* knockdown larvae. Data are shown as individual values

± standard deviation (SD). Data obtained from two independent experiments. (**C**)

Numbers of viable colony forming units (CFUs) control or *atg5* knockdown larvae

infected intravenously with approximately 1600 CFU of *S. pneumoniae* monitored at

1, 4, and 7 hpi. Data obtained from two independent experiments (**D**) Representative

in vivo images of circulation valley of control (top) and *atg5*, *atg16l1* knockdown

larvae (bottom) at 2 hpi injected with *S. pneumoniae* D39 Δ*cps* prestained with

pHrodo Red. Panels on the right represent zoomed-in areas (indicated by dashed

squares). Scale bars represent 10 μm . (**E**) Quantification of pHrodo Red

fluorescence intensity within phagocytes containing pneumococci. Data obtained

from two independent experiments). **** - $p<0.0001$, *** - $p<0.001$, ns - not significant

Figure 3. ROS-dependent autophagic response contribute to clearance of *S.*

pneumoniae. (**A, D**) representative confocal images of control and indicated gene

knockdown (A – *cyba*, D - *rubcn*) *CMV:GFP-Lc3* larvae infected systemically with

approximately 1600 CFU of mKate2-labeled D39 Δ*cps S. pneumoniae* and fixed at 2

hpi. Scale bars: 10 μm. (**B, E**) Quantification of Lc3 associations with intracellular *S.*

pneumoniae within infected phagocytes of fixed *CMV:GFP-Lc3* control and indicated

gene knockdown (B – *cyba*, E - *rubcn*) larvae at 2 hpi. Data are shown as mean ±

standard deviation (SD). Data obtained from three independent experiments. (**C, F**)

Numbers of viable colony forming units (CFUs) over time of control and indicated

gene knockdown (C – *cyba*, F - *rubcn*). Data obtained from three independent

experiments.

Figure 4. Live and pnemolysin-positive pneumococci are required to fully induce Lc3-
mediated response within zebrafish macrophages. (**A**) representative confocal
images of *CMV:GFP-Lc3* control larvae (top two panels) or *cyba* knockdown (bottom
two panels) infected systemically with approximately 1600 CFU of live (left two
panels) and heat-killed (right two panels) mKate2-labeled D39 Δ*cps S. pneumoniae*
fixed at 2 hpi. Scale bars: 10 μm. (**B**) Quantification of Lc3 associations with
intracellular live or heat-killed *S. pneumoniae* within infected phagocytes of fixed
CMV:GFP-Lc3 control or *cyba* larvae at 2 hpi. Data are shown as mean ± standard
deviation (SD). Data obtained from two independent experiments. (**C**) Anti-Ply
immunoblot analysis of whole-cell lysates from the indicated *S. pneumoniae* strains.
The position of protein markers are indicated in kiloDaltons, Pneumolysin (Ply) ≈ 53.7
KDa. Representative blot shown, n = 4. (**D**) representative confocal images of control
CMV:GFP-Lc3 larvae infected systemically with approximately 1600 CFU of Ply-
positive (D39 Δ*cps*, left panel) and Ply-negative (D39 Δ*cps* Δ*ply*, right panel) mKate2-
labeled *S. pneumoniae* fixed at 2 hpi. Scale bars: 10 μm. (**E**) Quantification of Lc3
associations with intracellular Ply-positive or Ply-negative *S. pneumoniae* within
infected phagocytes of fixed *CMV:GFP-Lc3* larvae at 2 hpi. Data are shown as mean
± standard deviation (SD). Data obtained from two independent experiments.

Figure 5. Sqstm1, optineurin and atg13 are not involved in autophagic response to
S. pneumoniae. (**A**) representative confocal images of *sqstm1*$^{+/+}$/*optn*$^{+/+}$ and *sqstm1*$^{-/-}$
/*optn*$^{-/-}$ double mutant *CMV:GFP-Lc3* larvae infected systemically with approximately
1600 CFU of mKate2-labeled D39 Δ*cps S. pneumoniae* and fixed at 2 hpi Scale bars:
10 μm. (**B**) Quantification of Lc3 associations with intracellular *S. pneumoniae* within
infected phagocytes of fixed *CMV:GFP-Lc3 sqstm1*$^{+/+}$/*optn*$^{+/+}$ and *sqstm1*$^{-/-}$/*optn*$^{-/-}$
double mutant larvae. Data are shown as mean ± standard deviation (SD). Data
obtained from two independent experiments. (**C**) Numbers of viable colony forming
units (CFUs) over time of *sqstm1*$^{+/+}$/*optn*$^{+/+}$ and *sqstm1*$^{-/-}$/*optn*$^{-/-}$ double mutant larvae.
Data obtained from two independent experiments. (**D**) representative confocal
images of control and *atg13* knockdown (right panel) *CMV:GFP-Lc3* larvae infected
systemically with approximately 1600 CFU of mKate2-labeled D39 Δ*cps S.*
pneumoniae and fixed at 2 hpi Scale bars: 10 μm. (**E**) Quantification of Lc3
associations with intracellular *S. pneumoniae* within infected phagocytes of fixed

CMV:GFP-Lc3 control and *atg13* knockdown larvae. Data are shown as mean ±

standard deviation (SD). n≥15 zebrafish larvae were analyzed.

Figure 6. Tecpr1a, a likely ortholog of human TECPR1, controls Ply-mediated

autophagic response to pneumococci. (**A**) Table summarizing the alignment of

zebrafish Tecpr1a and Tecpr1b proteins to the human TECPR1 protein. Columns

display percentage positives (identical amino acids, as well as conserved or

semiconserved substitutions) and length of the proteins. Alignments sequence

similarities are shown for the full-length protein and the AIR domain, which exhibits

the greatest divergence between the zebrafish paralogues (Tecpr1a and Tecpr1b).

(**B**) Schematic representation of zebrafish Tecpr1a and Tecpr1b proteins, showing

their domains: DysF - Dysferlin, AIR - ATG5-interacting region, and PH - Pleckstrin

homology domain. Alignment of the AIR domain amino acid sequence of Human

TECPR1 and the zebfafish orthologs, Tecpr1a and Tecpr1b, is shown between the

schematics. Symbols denote: (*) identical residues, (:) conserved substitutions, and

(.) semiconserved substitutions. (**C**) Gene expression of adult zebrafish leukocytes

determined using the Zebrafish Blood Atlas [48]. Each dot represents a separate

scRNAseq sample (cell); replicates were performed across multiple zebrafish wild-

type and transgenic strains. Each arm of the schematic represents a separate blood

cell population (labeled). Deeper color indicates higher expression (log_{10} scale bars

described for each gene). (**D**) representative confocal images of control, *tecpr1a*,

tecpr1b and *tecpr1a/b* double knockdown *CMV:GFP-Lc3* larvae infected systemically

with approximately 1600 CFU of mKate2-labeled D39 *Δcps S. pneumoniae* fixed at 2

hpi. (**E**) Quantification of Lc3 associations with intracellular *S. pneumoniae* within

infected phagocytes of fixed *tecpr1a, tecpr1b* and *tecpr1a/b* double knockdown

CMV:GFP-Lc3 larvae. Data are shown as mean ± standard deviation (SD). n≥17

zebrafish larvae were analyzed. (**F**) representative confocal images of control or

tecpr1a knockdown *CMV:GFP-Lc3* larvae infected systemically with approximately

1600 CFU of mKate2-labeled Ply-positive (D39 *Δcps*) or Ply-negative (D39 *Δcps*

Δply) *S. pneumoniae* fixed at 2 hpi. (**G**) Quantification of Lc3 associations with

intracellular Ply-positive or Ply-negative *S. pneumoniae* within infected phagocytes of

control or *tecpr1a* knockdown *CMV:GFP-Lc3* larvae fixed at 2 hpi. Data are shown as

mean ± standard deviation (SD). Data obtained from two independent experiments.

(**H**) Numbers of viable colony forming units (CFUs) over time of control and *tecpr1a*

knockdown larvae. Data obtained from two independent experiments.

Figure 7. Both LAP and STIL are evoked in response to *S. pneumoniae*. (**A**)

representative confocal images of control, *cyba, tecpr1a* and *cyba*/*tecpr1a* double

knockdown *CMV:GFP-Lc3* larvae infected systemically with approximately 1600 CFU

of mKate2-labeled D39 Δ*cps S. pneumoniae* fixed at 2 hpi. (**B**) Quantification of Lc3

associations with intracellular *S. pneumoniae* within infected phagocytes of fixed

cyba, tecpr1a and *cyba*/*tecpr1a* double knockdown *CMV:GFP-Lc3* larvae. Data are

shown as mean ± standard deviation (SD). Data obtained from two independent

experiments. (**C**) Numbers of viable colony forming units (CFUs) over time of control,

cyba, *tecpr1a and cyba*/*tecpr1a* double knockdown larvae. Data obtained from two

independent experiments.

Figure 8. Involvement of ROS-independent LAP-like CASM of *S. pneumoniae*-

containing vesicles. (**A**) representative confocal images of control, *tecpr1a/cyba*

double knockdown or *atg16l1/tecpr1a/cyba* triple knockdown *CMV:GFP-Lc3* larvae

infected systemically with approximately 1600 CFU of Ply-positive (D39 Δ*cps*) and

Ply-negative (D39 Δ*cps* Δ*ply*) mKate2-labeled *S. pneumoniae* at 2 hpi. (**B**)

Quantification of Lc3 associations with intracellular PLY-positive or PLY-negative *S.*

pneumoniae within infected phagocytes of control, *tecpr1a/cyba* double knockdown

or *atg16l1/tecpr1a/cyba* triple knockdown *CMV:GFP-Lc3* larvae. Data are shown as

mean ± standard deviation (SD). Data obtained from two independent experiments.

Figure 9. Model of autophagic response to pneumococci within zebrafish

macrophages. After internalization by phagocytes, pneumococci are sequestered

within Lc3-decorated vacuoles which are formed by three independent mechanisms:

ROS-dependent LAP or two ROS-independent Ply-mediated pathways including

STIL. LAP occurs as a result of osmotic imbalance within bacteria-containing

phagosome caused by ROS production by NADPH oxidase. On the other hand, STIL

is induced upon membrane perturbation caused by secretion of bacterial

pneumolysin manifested by sphingomyelin presence on the outside leaflet of the

phagosome. STIL is dependent on Tecpr1a which recognizes sphingomyelin and

mediates Lc3 lipidation. Lastly, the Atg16l1- and Ply-dependent pathway contributes

to the smallest portion of Lc3 lipidation. However, the specific mechanisms controlling

this process remain to be revealed. Following Lc3-assisted vacuolar sequestration,

pneumococci are more efficiently targeted for lysosomal degradation and the Lc3

decoration is removed. The classical xenophagy receptors (sqstm1 and optineurin)

as well as the canonical autophagy initiation molecule (atg13) are dispensable in this

model. Despite the absence of these xenophagy-mediating factors, the bacteria are

efficiently degraded. Figure created with BioRender.

Region of imaging

A

| 2 hpi | 4 hpi | 7 hpi |

GFP-Lc3 + *S.p*

B

% of infected phagocytes with Lc3-associated bacteria

2hpi 4 hpi 7 hpi

C

| *S. pneumoniae*-Alexa fluor 647 | macrophage-mCherry | GFP-Lc3 (CMV) |

| merge | zoomed merge |

A

control   *atg5*

GFP-Lc3 + *S.p*

B

% of infected phagocytes with Lc3-associated bacteria

****

control   *atg5*

C

CFU/embryo

ns   ***   ns

control 1 hpi   *atg5* 1 hpi   control 4 hpi   *atg5* 4 hpi   control 7 hpi   *atg5* 7 hpi

D

control (pHrodo)   control merge   zoomed control (pHrodo)

atg5/atg16l1 (pHrodo)   *atg5/atg16l1* merge   zoomed *atg5/atg16l1* (pHrodo)

E

Mean fluorescence intensity pHrodo

***

control   *atg5/atg16l1*

A control   *cyba*

GFP-Lc3 + *S.p*

B

% of infected phagocytes with Lc3-associated bacteria

****

control   *cyba*

C

CFU/embryo

ns   **

control 1 hpi   *cyba* 1 hpi   control 4 hpi   *cyba* 4 hpi

D control   *rubcn*

GFP-Lc3 + *S.p*

E

% of infected phagocytes with Lc3-associated bacteria

****

control   *rubcn*

F

CFU/embryo

ns   **

control 1 hpi   *rubcn* 1 hpi   control 4 hpi   *rubcn* 4 hpi

A

sqstm1⁺/⁺ optn⁺/⁺   *sqstm1⁻/⁻ optn⁻/⁻*

GFP-Lc3 + *S.p*

B

ns

% of infected phagocytes with Lc3-associated bacteria

sqstm1⁺/⁺ optn⁺/⁺   *sqstm1⁻/⁻ optn⁻/⁻*

C

CFU/embryo

ns   ns

sqstm1⁺/⁺ optn⁺/⁺ 1 hpl
sqstm1⁻/⁻ optn⁻/⁻ 1 hpl
sqstm1⁺/⁺ optn⁺/⁺ 4hpl
sqstm1⁻/⁻ optn⁻/⁻ 4 hpl

D

control   *atg13*

GFP-Lc3 + *S.p*

E

ns

% of infected phagocytes with Lc3-associated bacteria

control   *atg13*

A

PROTEIN	Alignment to *H.sapiens* TECPR1 (1164 aa)	AIR Domain Alignement	Protein Length
Tecpr1a (*D.rerio*)	78% Positives	95.5% Positives	1198 aa
Tecpr1b (*D.rerio*)	66% Positives	44.4% Positives	1105 aa

B

Tecpr1a

Tectonin repeat

500   1198

Tecpr1a positives: **, .: . *.**:. :*:*.******* ******:.**.;**;*** *****;*****;
D.rerio Tecpr1a 570 WFNSSTSTTTLNSSMQAMSFSITTPQTAAWRKQ IFEQLSERSKFEMENFMHYEQAEQSVWVK 631
H.sapiens TECPR1 551 WFTVQA--GLSSSVHMLSLSITTPQTAAWRKQIFQQLTERTKELENFRHYEQAVEQSVWVK 610
D.rerio Tecpr1b 512 CFHSTD------AEPLSPSITPALRAA---------DTHTAAAQQHRDACADQSVRVT 554
Tecpr1b positives: *   .. ** ***** **   .:   :* : ..:*** *.

Tecpr1b

500   1105

C

tecpr1a

tecpr1b

D

control   *tecpr1a*

tecpr1b   *tecpr1a/tecpr1b*

GFP-Lc3 + *S.p*

E

% of infected phagocytes with Lc3-associated bacteria

F

control Ply-positive   control Ply-negative

tecpr1a Ply-positive   *tecpr1a* Ply-negative

GFP-Lc3 + *S.p*

G

% of infected phagocytes with Lc3-associated bacteria

H

CFU/embryo

A

control   *cyba*

tecpr1a   *cyba/tecpr1a*

GFP-Lc3 + *S.p*

B

% of infected phagocytes with Lc3-associated bacteria

****
****
****
****   ns   ****

control   *cyba*   *tecpr1a*   *cyba/tecpr1a*

C

CFU/embryo

ns   ****

*   *
*   0.075

control 1 hpi   *cyba* 1 hpi   *tecpr1a* 1 hpi   *cyba/tecpr1a* 1 hpi   control 4 hpi   *cyba* 4 hpi   *tecpr1a* 4 hpi   *cyba/tecpr1a* 4 hpi

A

Ply-positive *Sp*, control larvae

Ply-positive *Sp*, *tecpr1a/cyba* larvae

Ply-positive *atg16l1/tecpr1a/cyba* larvae

Ply-negative *Sp*, *tecpr1a/cyba* larvae

GFP-Lc3 + *S.p*

B