Quantitative Evaluation of Normal Cerebrospinal fluid flow in Sylvian aqueduct and

perivascular spaces of middle cerebral artery and circle of Willis using 2D phase -contrast MRI

imaging.

Rosemarie Faustina D. Le; Christof Karmonik, PhD; Angelique S. Regnier-Golanov, PhD;

Eugene V. Golanov MD, PhD; Gavin W. Britz, MD*

Houston Methodist Academic Institute,

6560 Fannin str., Suite 944,

Houston, TX, 77030

*Corresponding author.

Abstract

Recently, it was proposed that CSF flow constitutes a critical part of the glymphatic system,

playing a critical role in various brain abnormalities from Alzheimer's disease to hydrocephalus.

Thus, measurement of CSF flow has been increasingly used for diagnostic and clinical monitoring

purposes. Phase-contrast MRI has been used to determine CSF flow. However, CSF flow in the

periarterial spaces of the circle of Willis and the middle cerebral artery which are important

conduits remain unexplored. We employed phase-contrast MRI to explore CSF flow along the

perivascular spaces of the circle of Willis and the middle cerebral artery to establish baseline

parameters of CSF and compare them with the Sylvian aqueduct. To analyze CSF flow in the

perivascular space, we developed a new, semi-automated method for outlining the perivascular

space and extracting CSF flow parameters. The twenty-four healthy participants were recruited to

achieve an even distribution by age (mean: 40 ± 11) and gender (13 males). We validated our

routine for CSF flow measurements by comparing CSF flow in the Sylvian aqueduct (0.00700

mL/s) with the range of literature values, 0.0049-0.0432 mL/s. For all CSF parameters, the circle

of Willis and middle cerebral artery were differed from the Sylvian aqueduct. For most CSF flow

parameters, the 95% confidence intervals of the circle of Willis and middle cerebral artery overlap.

The linear mixed models and general linear mixed models for flow indicate strong effects of the

conduits. CSF velocity in these conduits were lower 0.159 cm/s and 0.198 cm/s respectively than

in the Sylvian aqueduct. Overall, differences in CSF flow parameters between sex and age groups

were negligible. In this study, we have validated our routine and established baseline values of

CSF flow along the circle of Willis and the middle cerebral artery as well as highlighted the limited

influence of sex and/or age.

Background

Cerebrospinal fluid (CSF) is a clear fluid surrounding the central nervous system (CNS): brain, filling cerebral ventricles, and central canal of the spinal cord. At any given time, there are 90-150 mL of CSF in the cranium and spinal cord because production and absorption occur at the same rates to maintain intracranial pressure (1, 2). CSF is produced in the choroid plexus of the lateral ventricles and flows through the interventricular foramina to the third ventricle, along the Sylvian aqueduct (SA) to the fourth ventricle, proceeding to the subarachnoid space, and to the spinal cord (1-3). Recently, the concept of the glymphatic pathway has been introduced (4, 5). According to it CSF flows from the subarachnoid space to periarterial space and enters the brain parenchyma where it mixes with the interstitial fluid and further exits parenchyma into the paravenous space. CSF flow is pulsatile due to cardiac pulsations and breathing (3, 6). CSF moves in opposite directions depending on the cardiac cycle: during systole, intracranial arteries increase in volume which pushes CSF in the craniocaudal direction. The reverse happens in diastole as blood vessel volume decreases and CSF moves in the caudocranial direction (2, 7, 8).

CSF serves several physiologic functions. CSF provides support and cushioning for the brain. Movement of CSF through the brain parenchyma clears out waste and transport essential molecules (1-3, 9, 10). These functions and dynamics, especially CSF flow and pressure, can be altered in pathologies. As a part of the glymphatic system, CSF flow has been proposed to play a role in Alzheimer's disease by failing to clear out amyloid beta and tau tangles, the characteristic protein markers (3, 11, 12). Abnormal increase in intracranial pressure due to abnormal CSF drainage is associated with hydrocephalus (7, 8, 11, 12). Other studies have observed the roles of CSF flow in meningitis, cerebral edema, and other cerebrovascular diseases (8, 12). Thus,

measurement of CSF flow has been increasingly used for diagnostic and clinical monitoring

purposes (13).

To explore the clinical implications of CSF flow, many studies focused on the CSF flow through

the Sylvian aqueduct (SA) which connects the third and fourth ventricles (1, 10). SA containing

only CSF flow is easier to identify and image, so it frequently has been studied. On the other hand,

the periarterial spaces of circle of Willis (COW) and the middle cerebral artery (MCA), important

conduits of CSF flow, remain unexplored. The COW localized to the base of the brain formed by

anterior cerebral arteries, anterior communicating artery, internal carotid arteries (at their distal

tips), posterior cerebral arteries, and posterior communicating artery. MCA originates from COW

at the internal carotid arteries joining and ascending in the lateral sulcus of the cerebrum providing

blood supply to many parts of the lateral cortex.

Therefore, this study has several objectives. First, because the CSF flow along the perivascular

spaces of COW and MCA have not been studied yet, we aim to establish baseline measurements

of CSF flow parameters along these conduits, including potential differences due to sex and age.

Second, we aim to compare CSF flow parameters between the SA, COW, and MCA.

For this study, we used phase-contrast magnetic resonance imaging (PC-MRI). PC-MRI was first

used in 1994 by Barkhof et al. (14) to evaluate the effect of age on SA flow (12). Since then, PC-

MRI has emerged as the gold standard for measuring CSF flow (15). The advantages of PC-MRI

are that it is noninvasive and relatively quick at taking measurements (12).

To analyze CSF flow in the perivascular space, we developed a new, semi-automated method for

outlining perivascular space and extracting CSF flow and velocity parameters (Additional file 1).

Many semi-automated or automated methods have been developed in order to increase accuracy,

save time on analysis, and/or contribute to study reproducibility (16, 17). We conducted

preliminary evaluation of our semi-automated program, through validation with the CSF flow

through SA parameters which are extensively covered in the literature.

We focused on the following CSF flow parameters: stroke volume (StroVol), volumetric flow rate

(VFR), systolic flow rate (SFR), diastolic flow rate (DFR), velocity, peak systolic velocity (PSV),

and peak diastolic velocity (PDV). While the most useful parameters have yet to be determined,

we chose these parameters to capture the overall picture of CSF flow dynamics in the hopes that

they may be used for future clinical diagnostics and therapeutics (18).

Methods

<u>Participants</u>

The twenty-four healthy participants were recruited to achieve an even distribution by age and

gender (13 males). Ages ranged from 23 to 59 with a mean of 40 ± 11 (Table 1). To obtain baseline

parameters by demographics, participants were separated into different age groups: 20-29, 30-39,

40-49, and 50-59. The study was approved by the Houston Methodist Hospital Institutional Review

Board, protocol Pro00022145.

Table 1. Participant demographics.

Demographics	Description	Number	Percentage
Age	20-29	6	25%
	30-39	4	17%
	40-49	7	29%
	50-59	7	29%
	Total	24	100%
Sex	Male	13	54%
	Female	11	46%
	Total	24	100%

<u>MRI Acquisition</u>

A 2D PC-MRI image was acquired at each region of interest (COW, MCA, SA) chosen by an

experienced neuroscientist (EG, ARG). Details of the acquisition parameters are as follows: slice

thickness 3 mm, acquisition matrix 272 x 272, FOV 160 cm x 160 cm, in-plane resolution 0.6 mm

x 0.6 mm. TE=3 msec, TR=105 msec. Images were acquired with a single transmit, 32 channel

receive head coil in the FDA-approved clinical mode of the MAGNETOM 7 Tesla human MRI

scanner (Siemens Healthineers, Erlangen, Germany). A 3D velocity encoding scheme was used to

account for CSF flow direction not exactly parallel to the slice normal. The encoding velocity

(VENC) was set at 5 cm/s to remain sensitive to the corresponding low velocity of CSF flow and

to eliminate blood flow contamination which can be recognized as global maximum/minimum

gray scale values due to the phase wrapping artifact of blood velocities exceed this VENC value.

A total of 20 images evenly spaced across the cardiac cycle were reconstructed with a total

acquisition time of approximately 5 min for each location.

<u>CSF Flow Analysis</u>

As the VENC value was low compared to blood flow velocities, global maximum black/white gray

scale intensity values ––caused by the phase wrapping artifact–– were used as a mask for

distinguishing and eliminating blood flow from perivascular CSF flow in 2D phase contrast

images. Regions of interest (ROI) in an annular shape centered on the arterial cross section were

chosen for quantifying CSF flow using a semi-automated algorithm described in Additional file 1.

a)

b)

Figure 1. Examples of ROIs. (a) ROI of SA and (b) one of two ROIs of MCA on phase images.

For the first phase image of the 20 series of images for the cardiac cycle, the X,Y-coordinates,

tolerance, and band size of the ROIs were determined by the user (see Additional file 1).

Magnitude images were used as anatomical references. Two ROIs --one left and one right-- were

identified for each COW and MCA and one ROI was identified for each SA (Figure 1). The

established ROI settings were entered into our developed in-house semi-automated program to

produce flow rate (FR) and Velocity values. FR (mL/s) was calculated as the area of the ROI

(converted from mm^2 to cm^2) x VENC x (mean intensity / maximum gray scale intensity) Velocity

(cm/s) was calculated as FR / the area of the ROI. ROIs for 20 cardiac gated images over one

cardiac cycle were obtained from each series to obtain the FR and Velocity curves.

Peak timing was standardized by looking at which percentage of the cardiac cycle peak flow

occurred. The percentage of the cardiac cycle was calculated as (the ordinal position of the image

within the series / 20) x 100. The stroke volume was calculated as FR x time.

<u>Statistical Analysis</u>

For all parameters in each location of SA, COW, and MCA, median values and bias-corrected and

accelerated bootstrap (BCa) confidence intervals (CIs) from non-parametric bootstrapping were

calculated.

To compare the conduits and evaluate the influence of demographics, various models were

constructed. For StroVol, VFR, and Velocity, linear mixed models (LMM) were constructed with

block bootstrapping. Generalized linear mixed models (GLMM) with the gamma family and log

link function were constructed with block bootstrapping for SFR, DFR, and Area. Generalized

linear models (GLM) were constructed for PSV and PDV. For these LMMs, GLMMs, and GLMs,

BCa CIs were calculated. For time to peak as percentage of cardiac cycle duration (TP) of StroVol

(TPStroVol), VFR (TPVFR), and Velocity (TPVelocity), beta regression with the logit link

function was constructed, and Wald CIs were calculated. For comparing SA, COW, and MCA,

fixed effects were the intercept (= SA), time (except for PSV, PDV, and TP parameters), COW,

and MCA. SA was set as the intercept because CSF flow parameters for the SA are already

established in the literature. For evaluating the influence of sex and age, fixed effects were the

intercept, time (except for PSV, PDV, and TP parameters), sex, and age. An interaction term

between sex and age was also added. When applicable, participants were considered as random

effects to account for the non-independence of time series. Age was rescaled, zero random effects

were dropped, and/or optimizers were changed to address model convergence issues as needed.

Whenever the data contained zero-values, the following transformation was applied to the outcome

variable before beta regression: (y x 23 + 0.5) / 23. For post-hoc testing, p-values were adjusted

using the Benjamini-Hochberg procedure. Assumptions were checked with QQ plots and residual

plots. All statistical analysis and visualization were performed in R/RStudio.

Results

Baseline values of CSF flow parameters with the median and 95% CI by conduits and by sex and

age are reported in Table 2 and Tables 3 and 4 respectively. Model estimates of the fixed and

random effects are reported in Table 5 for conduits comparison and Table 6 for the effect of sex

and age . StroVol is reported in mL. VFR, SFR, and DFR are reported in mL/s. Velocity, PSV,

and PDV are reported in cm/s. TPStroVol, TPVFR, and TPVelocity are reported in percentage of

the cardiac cycle. Area is reported in cm^2.

Table 2. Median values of CSF flow parameters in SA, COW, and MCA.

	Conduit	Median	Lower 95% CI	Upper 95% CI
StroVol (mL)	SA	0.00254	0.00221	0.00277
	COW	0.0146	0.0140	0.0156
	MCA	0.0140	0.0131	0.0149
VFR (mL/s)	SA	0.00700	0.00600	0.00700
	COW	0.0420	0.0400	0.0430
	MCA	0.0450	0.0420	0.0460
SFR (mL/s)	SA	0.00900	0.00800	0.00900
	COW	0.0530	0.0500	0.0570
	MCA	0.0550	0.0530	0.0580
DFR (mL/s)	SA	0.00400	0.00400	0.00500
	COW	0.0230	0.0220	0.0250
	MCA	0.0270	0.0230	0.0280
Velocity (cm/s)	SA	0.578	0.578	0.723
	COW	0.285	0.271	0.299
	MCA	0.260	0.246	0.267

PSV (cm/s)	SA	1.45	1.16	1.45
	COW	0.749	0.677	0.848
	MCA	0.676	0.516	0.728
PDV (cm/s)	SA	0.867	0.578	0.867
	COW	0.383	0.368	0.622
	MCA	0.405	0.303	0.452
TPStroVol (%)	SA	50.0	40.0	62.5
	COW	75.0	70.0	80.0
	MCA	70.0	70.0	80.0
TPVFR (%)	SA	20.0	20.0	25.0
	COW	40.0	30.0	55.0
	MCA	40.0	15.0	42.5
TPVelocity (%)	SA	55.0	50.0	55.0
	COW	65.0	50.0	65.0
	MCA	65.0	55.0	75.0
Area (cm^2)	SA	0.00692	0.00692	0.0104
	COW	0.0277	0.0277	0.0311
	MCA	0.0398	0.0381	0.0415

148   **Table 3. Median values of CSF flow parameters in different conduits by sex.**

	Conduit	*Sex*	*Median*	*Lower 95% CI*	*Upper 95% CI*
StroVol (mL)	SA	Male	0.00290	0.00258	0.00310
		Female	0.00196	0.00171	0.00219
	COW	Male	0.0157	0.0142	0.0173
		Female	0.0140	0.0126	0.0147
	MCA	Male	0.0140	0.0128	0.0152
		Female	0.0141	0.0128	0.0158
VFR (mL/s)	SA	Male	0.00800	0.00800	0.0100

		Female	0.00500	0.00500	0.00700
	COW	Male	0.0450	0.0425	0.0490
		Female	0.0380	0.0360	0.0410
	MCA	Male	0.0450	0.0430	0.0490
		Female	0.0440	0.0420	0.0470
SFR (mL/s)	SA	Male	0.0100	0.00800	0.0110
		Female	0.00700	0.00600	0.00900
	COW	Male	0.0580	0.0530	0.0620
		Female	0.0480	0.0460	0.0550
	MCA	Male	0.0590	0.0570	0.0640
		Female	0.0520	0.0490	0.0550
DFR (mL/s)	SA	Male	0.00500	0.00500	0.00700
		Female	0.00300	0.00200	0.00300
	COW	Male	0.0250	0.0220	0.0270
		Female	0.0220	0.0180	0.0230
	MCA	Male	0.0280	0.0230	0.0290

		Female	0.0235	0.0190	0.0295
Velocity (cm/s)	SA	Male	0.674	0.578	0.761
		Female	0.528	0.482	0.578
	COW	Male	0.324	0.299	0.347
		Female	0.256	0.241	0.275
	MCA	Male	0.260	0.241	0.271
		Female	0.260	0.246	0.275
PSV (cm/s)	SA	Male	1.59	1.45	1.88
		Female	1.30	0.867	1.45
	COW	Male	0.771	0.742	1.08
		Female	0.737	0.526	0.795
	MCA	Male	0.686	0.552	0.867
		Female	0.605	0.489	0.778
PDV (cm/s)	SA	Male	1.32	0.896	1.73
		Female	0.578	0.578	0.867
	COW	Male	0.420	0.310	0.688

		Female	0.383	0.171	0.506
	MCA	Male	0.415	0.348	0.518
		Female	0.330	0.282	0.487
TPStroVol (%)	SA	Male	55.0	40.0	65.0
		Female	35.0	25.0	60.0
	COW	Male	70.0	70.0	85.0
		Female	75.0	70.0	87.5
	MCA	Male	75.0	65.0	75.0
		Female	65.0	65.0	80.0
TPVFR (%)	SA	Male	20.0	15.0	25.0
		Female	20.0	17.5	42.5
	COW	Male	35.0	15.0	45.0
		Female	50.0	35.0	70.0
	MCA	Male	35.0	10.0	40.0
		Female	40.0	15.0	45.0
TPVelocity (%)	SA	Male	55.0	55.0	75.0

		Female	50.0	44.9	50.0
	COW	Male	65.0	50.0	65.0
		Female	70.0	50.0	80.0
	MCA	Male	70.0	60.0	80.0
		Female	62.5	45.0	75.0
Area (cm^2)	SA	Male	0.00692	0.00692	0.0104
		Female	0.0104	0.00692	0.0104
	COW	Male	0.0277	0.0277	0.0329
		Female	0.0277	0.0242	0.0346
	MCA	Male	0.0450	0.0381	0.0450
		Female	0.0346	0.0311	0.0415

Table 4. Median values of CSF flow parameters in different conduits by age group.

	Conduit	*Age Group*	*Median*	*Lower 95% CI*	*Upper 95% CI*
StroVol (mL)	SA	20-29	0.00217	0.00194	0.00256
		30-39	0.00288	0.00219	0.00407

		40-49	0.00303	0.00268	0.00338
		50-59	0.00190	0.00173	0.00285
	COW	20-29	0.0143	0.0115	0.0157
		30-39	0.0129	0.0110	0.0143
		40-49	0.0169	0.0156	0.0192
		50-59	0.0143	0.0128	0.0157
	MCA	20-29	0.0121	0.0102	0.0133
		30-39	0.0176	0.0145	0.0205
		40-49	0.0174	0.0160	0.0203
		50-59	0.0118	0.0108	0.0132
VFR (mL/s)	SA	20-29	0.00600	0.00600	0.00800
		30-39	0.00700	0.00600	0.00900
		40-49	0.00700	0.00700	0.00850
		50-59	0.00600	0.00500	0.00900
	COW	20-29	0.0440	0.0410	0.0480
		30-39	0.0335	0.0295	0.0370

		40-49	0.0450	0.0415	0.0480
		50-59	0.0390	0.0350	0.0450
	MCA	20-29	0.0375	0.0340	0.0420
		30-39	0.0545	0.0475	0.0600
		40-49	0.0530	0.0490	0.0560
		50-59	0.0370	0.0330	0.0420
SFR (mL/s)	SA	20-29	0.00800	0.00600	0.00900
		30-39	0.0100	0.00800	0.0140
		40-49	0.00900	0.00700	0.00900
		50-59	0.00800	0.00500	0.0100
	COW	20-29	0.0510	0.0460	0.0560
		30-39	0.0490	0.0390	0.0580
		40-49	0.0580	0.0530	0.0640
		50-59	0.0525	0.0470	0.0580
	MCA	20-29	0.0500	0.0430	0.0530
		30-39	0.0620	0.0553	0.0720

		40-49	0.0660	0.0590	0.0700
		50-59	0.0470	0.0430	0.0495
DFR (mL/s)	SA	20-29	0.00350	0.00300	0.00500
		30-39	0.00400	0.00200	0.00500
		40-49	0.00500	0.00300	0.00500
		50-59	0.00400	0.00200	0.00500
	COW	20-29	0.0260	0.0230	0.0330
		30-39	0.0200	0.0150	0.0230
		40-49	0.0260	0.0250	0.0330
		50-59	0.0210	0.0189	0.0240
	MCA	20-29	0.0270	0.0210	0.0290
		30-39	0.0285	0.0185	0.0455
		40-49	0.0310	0.0290	0.0360
		50-59	0.0190	0.0145	0.0210
Velocity (cm/s)	SA	20-29	0.556	0.455	0.578
		30-39	0.540	0.442	0.589

		40-49	0.674	0.578	0.783
		50-59	0.723	0.578	0.867
	COW	20-29	0.293	0.267	0.326
		30-39	0.242	0.200	0.264
		40-49	0.295	0.273	0.320
		50-59	0.289	0.270	0.328
	MCA	20-29	0.267	0.248	0.289
		30-39	0.273	0.248	0.318
		40-49	0.269	0.253	0.293
		50-59	0.237	0.217	0.255
PSV (cm/s)	SA	20-29	1.16	1.16	1.73
		30-39	1.30	1.30	2.31
		40-49	1.45	1.45	1.88
		50-59	1.59	1.16	3.47
	COW	20-29	0.674	0.462	0.694
		30-39	0.751	0.605	1.29

		40-49	0.751	0.418	0.771
		50-59	0.828	0.766	1.05
	MCA	20-29	0.535	0.484	0.815
		30-39	0.676	0.124	0.700
		40-49	0.766	0.489	0.867
		50-59	0.605	0.500	0.867
PDV (cm/s)	SA	20-29	0.867	0.247	0.867
		30-39	0.578	0.578	1.76
		40-49	1.07	0.867	1.54
		50-59	0.723	0.578	2.75
	COW	20-29	0.171	0.0942	0.650
		30-39	0.390	0.390	0.650
		40-49	0.376	0.239	0.824
		50-59	0.491	0.289	0.549
	MCA	20-29	0.405	0.348	0.518
		30-39	0.275	0.303	1.49

		40-49	0.500	0.303	0.590
		50-59	0.318	0.128	0.381
TPStroVol (%)	SA	20-29	57.5	28.7	67.5
		30-39	25.0	20.0	70.0
		40-49	50.0	25.0	65.0
		50-59	50.0	45.0	60.0
	COW	20-29	75.0	70.0	92.5
		30-39	70.0	70.0	90.0
		40-49	70.0	67.5	90.0
		50-59	75.0	50.0	75.0
	MCA	20-29	65.0	65.0	80.0
		30-39	65.0	65.0	90.0
		40-49	75.0	50.0	75.0
		50-59	70.0	70.0	90.0
TPVFR (%)	SA	20-29	17.5	10.0	35.0
		30-39	22.5	20.0	67.5

		40-49	22.5	12.5	25.0
		50-59	20.0	5.00	55.0
	COW	20-29	50.0	27.5	60.0
		30-39	42.5	32.5	70.0
		40-49	25.0	20.0	55.0
		50-59	50.0	20.0	60.0
	MCA	20-29	47.5	35.0	65.0
		30-39	45.0	32.5	65.0
		40-49	15.0	15.0	60.0
		50-59	35.0	15.0	45.0
TPVelocity (%)	SA	20-29	55.0	50.0	72.5
		30-39	50.0	50.0	90.0
		40-49	55.0	55.0	85.0
		50-59	50.0	40.0	55.0
	COW	20-29	60.0	35.0	75.0
		30-39	80.0	60.0	90.0

		40-49	57.5	50.0	72.5
		50-59	60.0	45.0	80.0
	MCA	20-29	60.0	45.0	90.0
		30-39	57.5	30.0	75.0
		40-49	75.0	45.0	80.0
		50-59	70.0	45.0	80.0
Area (cm^2)	SA	20-29	0.0104	0.00692	0.0104
		30-39	0.0121	0.0104	0.0173
		40-49	0.0104	0.00692	0.0104
		50-59	0.00692	0.00346	0.00692
	COW	20-29	0.0311	0.0242	0.0346
		30-39	0.0311	0.0260	0.0415
		40-49	0.0311	0.0277	0.0381
		50-59	0.0208	0.0208	0.0293
	MCA	20-29	0.0311	0.0294	0.0381
		30-39	0.0588	0.0484	0.0813

| | 40-49 | 0.0484 | 0.0415 | 0.0519 |
| | 50-59 | 0.0346 | 0.0277 | 0.0381 |

152   a)

153   b)

Figure 2. Aggregate CSF flow curves. Aggregation of CSF (a) VFR and (b) Velocity curves for
all participants.

156 As expected, VFR oscillates in the SA which reflects cardiac pulsations. On the other hand, COW

157 VFR decreases throughout the cardiac cycle (Fig. 2a). Interestingly, Velocity in the SA is pulsatile

158 while it stays relatively constant in the COW and MCA (Fig. 2b).

159 **Table 5. Estimates of models of CSF flow parameters in SA (intercept) vs. COW vs. MCA.**

	Fixed Effects	*Coefficient*	*Lower 95% CI*	*Upper 95% CI*	*Random Effects*	*Variance*	*SD*	*H0*	*P-Value*
StroVol	Intercept (SA)	-0.00857	-0.0124	-0.00527	Intercept: Participants	5.22E-06	0.00228	SA = COW = MCA	0.000999
	Time	0.0278	0.0207	0.0335	Residuals	0.00142	0.0376	SA = COW	0.00400
	COW	0.0150	0.0103	0.0190				SA = MCA	0.00400
	MCA	0.0142	0.00979	0.0194				COW = MCA	0.805
VFR	Intercept (SA)	0.0230	0.0164	0.0303	Intercept: Participants	2.24E-05	0.00474	SA = COW = MCA	0.000999
	Time	-0.0241	-0.0363	-0.0139	Residuals	0.00598	0.0773	SA = COW	0.00400
	COW	0.0350	0.0261	0.0460				SA = MCA	0.00400
	MCA	0.0310	0.0203	0.0411				COW = MCA	0.529
SFR	Intercept (SA)	-3.73	-3.88	-3.63	Intercept: Participants	0.0172	0.131	SA = COW = MCA	0.00100
	Time	-0.483	-0.705	-0.320	Residuals	1.07	1.03	SA = COW	0.00400
	COW	1.36	1.20	1.50				SA = MCA	0.00400

	MCA	1.36	1.21	1.51				COW = MCA	0.969
DFR	Intercept (SA)	-4.78	-5.17	-4.55	Intercept: Participants	0.0738	0.272	SA = COW = MCA	0.00100
	Time	-0.536	-0.802	-0.245	Residuals	0.816	0.903	SA = COW	0.00799
	COW	1.48	1.16	1.83				SA = MCA	0.00799
	MCA	1.61	1.33	1.94				COW = MCA	0.216
Velocity	Intercept (SA)	0.421	0.318	0.534	Intercept: Participants	0.00180	0.0424	SA = COW = MCA	0.000999
	Time	-0.104	-0.178	-0.0277	Residuals	0.308	0.555	SA = COW	0.0160
	COW	-0.159	-0.286	-0.0498				SA = MCA	0.00599
	MCA	-0.198	0.0204	0.0410				COW = MCA	0.166
PSV	Intercept (SA)	0.537	0.413	0.645				SA = COW = MCA	0.000999
	COW	-0.773	-0.954	-0.613				SA = COW	0.00400
	MCA	-0.955	-1.12	-0.773				SA = MCA	0.00400
								COW = MCA	0.0280
PDV	Intercept (SA)	0.184	-0.0516	0.378				SA = COW = MCA	0.000999
	COW	-0.984	-1.24	-0.721				SA = COW	0.00400
	MCA	-1.01	-1.29	-0.747				SA = MCA	0.00400
								COW = MCA	0.901

TPStroVol	Intercept (SA) (Mean Model)	-0.0633	-0.329	0.202				SA = COW = MCA	3.87E-10
	COW (Mean Model)	1.12	0.792	1.45				SA = COW	7.34E-11
	MCA (Mean Model)	0.966	0.646	1.29				SA = MCA	1.19E-08
	Intercept (SA) (Precision Model)	1.21	0.859	1.57				COW = MCA	0.482
	COW (Precision Model)	0.590	0.125	1.06					
	MCA (Precision Model)	0.746	0.274	1.22					
TPVFR	Intercept (SA) (Mean Model)	-0.979	-1.26	-0.696				SA = COW = MCA	0.00131
	COW (Mean Model)	0.690	0.322	1.06				SA = COW	0.000468
	MCA (Mean Model)	0.486	0.110	0.862				SA = MCA	0.0274
	Intercept (SA) (Precision Model)	1.33	0.959	1.70				COW = MCA	0.471
	COW (Precision Model)	-0.450	-0.910	0.0100					
	MCA (Precision Model)	-0.434	-0.902	0.0342					
TPVelocity	Intercept (SA) (Mean Model)	0.465	0.288	0.642				SA = COW = MCA	0.861

	COW (Mean Model)	0.0137	-0.286	0.313					
	MCA (Mean Model)	0.0846	-0.224	0.393					
	Intercept (SA) (Precision Model)	1.60	1.32	1.88					
	COW (Precision Model)	-0.820	-1.20	-0.436					
	MCA (Precision Model)	-0.762	-1.16	-0.366					
Area	Intercept (SA)	-2.81	-2.96	-2.66	Intercept: Participants	0.135	0.367	SA = COW = MCA	0.000999
	Time	-1.43	-1.67	-1.22	Residuals	2.89	1.70	SA = COW	0.00400
	COW	0.984	0.793	1.15				SA = MCA	0.00400
	MCA	1.08	0.867	1.27				COW = MCA	0.230

SA was set as the intercept because CSF flow parameters for this conduit are already established

in the literature. The StroVol LMM indicates strong effects with COW StroVol 0.0150 mL (95%

CI [0.0103, 0.0190], p = 0.00400) and MCA StroVol 0.0142 mL (95% CI [0.00979, 0.0194], p =

0.00400) higher than SA (Table 5). The TPStroVol occurs later in the cardiac cycle in COW (β =

1.12, 95% CI [0.792, 1.45], p = 7.34 x 10^{-11}) and MCA (β = 0.966, 95% CI [0.646, 1.29], p = 1.19

x 10^{-8}) compared to SA (Table 5).

The LMMs and GLMMs for flow also indicate strong effects of the conduit as the magnitude of

the parameters is higher in COW and MCA. Specifically, COW VFR is 0.0350 mL/s (95% CI

[0.0261, 0.0460], p = 0.00400) and MCA VFR is 0.0310 mL/s (95% CI [0.0203, 0.0411], p =

0.00400) higher than SA (Table 5). COW SFR increases by a factor of 1.36 (95% CI [1.20, 1.50],

p = 0.00400) and MCA SFR by a factor of 1.36 (95% CI [1.21, 1.51], p = 0.00400) compared to

SA (Table 5). COW DFR also increases by a factor of 1.48 (95% CI [1.16, 1.83], p = 0.00799)

and MCA SFR by a factor of 1.61 (95% CI [1.33, 1.94], p = 0.00799) compared to SA (Table 5).

The TPVFR also occurs later in COW (β = 0.690, 95% CI [0.322, 1.06], p = 0.000468) and MCA

(β = 0.486, 95% CI [0.110, 0.862], p = 0.0274) compared to SA (Table 5).

Velocity in the LMM and GLMs also exhibits strong effects, and the magnitude of the parameters

is lower in COW and MCA. COW Velocity is 0.159 cm/s (95% CI [-0.286, -0.0498], p = 0.0160)

and MCA Velocity is 0.198 cm/s (95% CI [0.0204, 0.0410], p = 0.00599) lower than SA (Table

5). COW PSV decreases by a factor of 0.773 (95% CI [-0.954, -0.613], p = 0.00400), and MCA

PSV decreases by a factor of 0.955 (95% CI [-1.12, -0.773], p = 0.00400) compared to SA (Table

5). COW PDV also decreases by a factor of 0.984 (95% CI [-1.24, -0.721], p = 0.00400), and MCA

PDV decreases by a factor of 1.01 (95% CI [-1.29, -0.747], p = 0.00400) compared to SA (Table

5). Fpr TPVelocity, we were unable to find evidence against the hypothesis that SA = COW =

MCA (p = 0.861) (COW β = 0.0137, 95% CI [-0.286, 0.313]) (MCA β = 0.0846, 95% CI [-0.224,

0.393]) (Table 5).

For most CSF flow parameters, COW and MCA values for are similar (Table 5). It should be noted

that the 95% CI and p-value (p = 0.0280) for PSV contradict each other, but the latter is more

directly related to our hypothesis testing (Table 5). Therefore, we found evidence against the

hypothesis that COW = MCA for PSV.

There is considerably greater intra-subject variation than between-subject variation for the LMMs

and GLMMs which is unsurprising given the pulsatile nature of CSF flow (Table 5). This result

also supports that we found these effects among healthy participants since they are similar to each

other.

Sex and age are weak predictors of CSF flow parameters.

Table 6. Estimates of models of CSF flow parameters vs. sex, age, and their interaction for

each conduit.

	Conduit	Fixed Effects	Coefficient	Lower 95% CI	Upper 95% CI	Random Effects	Variance	SD	H0	P-Value
StroVol	SA	Intercept	0.00554	0.000963	0.0103	Intercept: Participants	2.37E-06	0.00154	Sex = 0, Age = 0, and Sex:Age = 0	0.694
		Time	-0.00133	-0.00385	0.00152	Residuals	9.57E-05	0.00978		
		Sex	-0.00164	-0.00718	0.00459					
		Age	-2.57E-05	-0.000152	9.55E-05					
		Sex x Age	1.68E-05	-0.000141	0.000154					
	COW	Intercept	0.0125	-0.0151	0.0369	Intercept: Participants	7.28E-06	0.00270	Sex = 0, Age = 0, and Sex:Age = 0	0.385
		Time	0.0309	0.0206	0.0399	Residuals	0.00154	0.0392		
		Sex	-0.00549	-0.0343	0.0318					
		Age	-0.000201	-0.000828	0.000445					

			Estimate	Lower	Upper		Variance	SD	Hypothesis	p
		Sex x Age	0.000164	-0.000551	0.000933					
	MCA	Intercept	-0.00496	-0.0262	0.0192	Intercept: Participants	2.69E-05	0.00519	Sex = 0, Age = 0, and Sex:Age = 0	0.334
		Time	0.0345	0.0239	0.0455	Residuals	0.00168	0.0410		
		Sex	0.0177	-0.0128	0.0488					
		Age	0.000171	-0.000441	0.000683					
		Sex x Age	-0.000393	-0.00115	0.000376					
VFR	SA	Intercept	0.0244	0.00850	0.0383	Intercept: Participants	1.57E-05	0.00397	Sex = 0, Age = 0, and Sex:Age = 0	0.835
		Time	-0.0234	-0.0334	-0.0151	Residuals	0.000719	0.0268		
		Sex	-0.00787	-0.0265	0.0104					
		Age	-4.93E-05	-0.000414	0.000367					
		Sex x Age	0.000185	-0.000298	0.000661					
	COW	Intercept	0.0757	0.0178	0.133	Intercept: Participants	4.00E-05	0.00632	Sex = 0, Age = 0, and Sex:Age = 0	0.445
		Time	-0.0301	-0.0479	-0.0148	Residuals	0.00653	0.0808		
		Sex	-0.0233	-0.101	0.0538					
		Age	-0.000384	-0.00194	0.00107					
		Sex x Age	0.000590	-0.00117	0.00243					

	MCA	Intercept	0.0257	-0.0223	0.0812	Intercept: Participants	0.000101	0.0101	Sex = 0, Age = 0, and Sex:Age = 0	0.0879
		Time	-0.0179	-0.0382	0.00150	Residuals	0.00705	0.0840		
		Sex	0.0507	-0.0138	0.122					
		Age	0.000574	-0.000657	0.00178					
		Sex x Age	-0.00112	-0.00283	0.000421					
SFR	SA	Intercept	-3.24	-3.52	-3.00	Intercept: Participants	0.0156	0.125	Sex = 0, Age = 0, and Sex:Age = 0	0.787
		Time	-1.61	-2.35	-1.18	Residuals	1.93	1.39		
		Sex	-0.207	-0.447	0.0393					
		Age	-0.0190	-0.222	0.139					
		Sex x Age	0.0246	-0.234	0.299					
	COW	Intercept	-2.38	-2.62	-2.28	Intercept: Participants	0.0135	0.116	Sex = 0, Age = 0, and Sex:Age = 0	0.0539
		Time	-0.368	-0.552	-0.175	Residuals	0.967	0.983		
		Sex	-0.104	-0.370	0.105					
		Age	-0.0128	-0.179	0.169					
		Sex x Age	0.0625	-0.155	0.305					
	MCA	Intercept	-2.43	-2.67	-2.34	Intercept: Participants	0.0466	0.216	Sex = 0, Age = 0, and Sex:Age = 0	0.157

		Time	-0.329	-0.581	-0.110	Residuals	0.843	0.918		
		Sex	-0.0412	-0.262	0.187					
		Age	0.0355	-0.186	0.215					
		Sex x Age	-0.150	-0.386	0.0833					
DFR	SA	Intercept	-4.67	-4.67	-4.04	Intercept: Participants	0.140	0.374	Sex = 0, Age = 0, and Sex:Age = 0	0.996
		Time	-1.32	-1.61	-0.761	Residuals	0.783	0.885		
		Sex	0.433	-0.837	-0.194					
		Age	0.0105	-0.0999	0.394					
		Sex x Age	-0.0251	-0.587	0.0329					
	COW	Intercept	-3.32	-3.63	-3.15	Intercept: Participants	0.0315	0.177	Sex = 0, Age = 0, and Sex:Age = 0	0.548
		Time	-0.253	-0.588	0.0993	Residuals	0.661	0.813		
		Sex	-0.266	-0.435	-0.0693					
		Age	-0.00338	-0.161	0.174					
		Sex x Age	-0.0578	-0.197	0.145					
	MCA	Intercept	-3.22	-3.48	-3.07	Intercept: Participants	0.100	0.317	Sex = 0, Age = 0, and Sex:Age = 0	0.999
		Time	-0.293	-0.636	0.0317	Residuals	0.745	0.863		
		Sex	-0.136	-0.425	0.156					

		Age	0.0760	-0.194	0.289					
		Sex x Age	-0.287	-0.541	-0.00865					
Velocity	SA	Intercept	0.0460	-0.851	1.13	Intercept: Participants	0.0128	0.113	Sex = 0, Age = 0, and Sex:Age = 0	0.217
		Time	-0.729	-1.03	-0.409	Residuals	1.34	1.16		
		Sex	0.469	-0.640	1.32					
		Age	0.0182	-0.0113	0.0410					
		Sex x Age	-0.0156	-0.0431	0.0150					
	COW	Intercept	0.352	0.0471	0.684	Intercept: Participants	0.00236	0.0486	Sex = 0, Age = 0, and Sex:Age = 0	0.249
		Time	-0.00946	-0.0934	0.0795	Residuals	0.146	0.382		
		Sex	-0.0995	-0.481	0.306					
		Age	-0.00347	-0.0114	0.00469					
		Sex x Age	0.00285	-0.00672	0.0120					
	MCA	Intercept	0.0482	-0.202	0.280	Intercept: Participants	0.00372	0.0610	Sex = 0, Age = 0, and Sex:Age = 0	0.273
		Time	0.0113	-0.0777	0.0977	Residuals	0.112	0.335		
		Sex	0.241	-0.0499	0.555					
		Age	0.00292	-0.00226	0.00896					
		Sex x Age	-0.00554	-0.0123	0.00113					

PSV	SA	Intercept	0.540	-0.356	1.41				Sex = 0, Age = 0, and Sex:Age = 0	0.0579
		Sex	-0.520	-1.54	0.457					
		Age	0.00399	-0.0195	0.0242					
		Sex x Age	0.00177	-0.0233	0.0248					
	COW	Intercept	0.0723	-0.623	0.647				Sex = 0, Age = 0, and Sex:Age = 0	0.273
		Sex	-0.838	-1.62	0.0779					
		Age	-0.00625	-0.0230	0.0102					
		Sex x Age	0.0180	-0.00132	0.0387					
	MCA	Intercept	-0.528	-1.16	0.126				Sex = 0, Age = 0, and Sex:Age = 0	0.208
		Sex	-0.141	-1.16	0.666					
		Age	0.00504	-0.0121	0.0206					
		Sex x Age	-0.00195	-0.0207	0.0218					
PDV	SA	Intercept	-0.133	-2.18	1.11				Sex = 0, Age = 0, and Sex:Age = 0	0.000999
		Sex	-0.0369	-1.41	2.17				Sex = 0	0.308
		Age	0.0167	-0.0179	0.0607				Age = 0	0.308
		Sex x Age	-0.0199	-0.0754	0.0163				Sex:Age = 0	0.308

	COW	Intercept	-0.820	-2.73	0.227				Sex = 0, Age = 0, and Sex:Age = 0	0.0180
		Sex	-1.12	-2.72	0.644				Sex = 0	0.647
		Age	0.00387	-0.0236	0.0447				Age = 0	0.827
		Sex x Age	0.0197	-0.0243	0.0573				Sex:Age = 0	0.647
	MCA	Intercept	-0.240	-0.986	0.833				Sex = 0, Age = 0, and Sex:Age = 0	0.595
		Sex	-0.841	-2.28	0.546					
		Age	-0.0133	-0.0450	0.00558					
		Sex x Age	0.0179	-0.0182	0.0492					
TPStroVol	SA	Intercept (Mean Model)	0.932	-0.651	2.52				Sex = 0, Age = 0, and Sex:Age = 0	0.164
		Sex (Mean Model)	-1.25	-3.27	0.777					
		Age (Mean Model)	-0.0183	-0.0572	0.0207					
		Sex x Age (Mean Model)	0.0179	-0.0338	0.0697					
		Intercept (Precision Model)	0.140	-1.79	2.07					

		Sex (Precision Model)	2.67	-0.154	5.50				
		Age (Precision Model)	0.0267	-0.0232	0.0765				
		Sex x Age (Precision Model)	-0.0554	-0.127	0.0160				
	COW	Intercept (Mean Model)	1.91	0.599	3.23				Sex = 0, Age = 0, and Sex:Age = 0	0.708
		Sex (Mean Model)	-0.600	-2.33	1.13				
		Age (Mean Model)	-0.0181	-0.0490	0.0128				
		Sex x Age (Mean Model)	0.0150	-0.0265	0.0566				
		Intercept (Precision Model)	1.62	-0.343	3.59				
		Sex (Precision Model)	0.377	-2.14	2.90				
		Age (Precision Model)	0.00488	-0.0423	0.0520				

		Sex x Age (Precision Model)	-0.0158	-0.0764	0.0448					
	MCA	Intercept (Mean Model)	1.69	0.582	2.80				Sex = 0, Age = 0, and Sex:Age = 0	0.0282
		Sex (Mean Model)	-1.50	-2.87	-0.126				Sex = 0	0.108
		Age (Mean Model)	-0.0127	-0.0405	0.0150				Age = 0	0.374
		Sex x Age (Mean Model)	0.0284	-0.00728	0.0640				Sex:Age = 0	0.158
		Intercept (Precision Model)	3.16	1.20	5.12					
		Sex (Precision Model)	0.874	-1.69	3.44					
		Age (Precision Model)	-0.0305	-0.0775	0.0165					
		Sex x Age (Precision Model)	-0.0158	-0.0769	0.0453					
TPVFR	SA	Intercept (Mean Model)	-0.666	-2.25	0.915				Sex = 0, Age = 0, and Sex:Age = 0	0.913

		Sex (Mean Model)	0.0741	-2.14	2.29					
		Age (Mean Model)	-0.00940	-0.0504	0.0316					
		Sex x Age (Mean Model)	0.000817	-0.0568	0.0584					
		Intercept (Precision Model)	1.38	-0.687	3.45					
		Sex (Precision Model)	0.373	-2.56	3.31					
		Age (Precision Model)	-0.00181	-0.0550	0.0514					
		Sex x Age (Precision Model)	-0.00762	-0.0826	0.0673					
	COW	Intercept (Mean Model)	0.0577	-1.24	1.36				Sex = 0, Age = 0, and Sex:Age = 0	0.0912
		Sex (Mean Model)	0.233	-1.54	2.01					
		Age (Mean Model)	-0.0161	-0.0493	0.0171					

		Sex x Age (Mean Model)	0.00874	-0.0351	0.0526				
		Intercept (Precision Model)	2.48	0.662	4.31				
		Sex (Precision Model)	-1.95	-4.25	0.353				
		Age (Precision Model)	-0.0345	-0.0782	0.00921				
		Sex x Age (Precision Model)	0.0439	-0.0116	0.0995				
	MCA	Intercept (Mean Model)	0.339	-1.14	1.81				Sex = 0, Age = 0, and Sex:Age = 0	0.587
		Sex (Mean Model)	-0.780	-2.72	1.16				
		Age (Mean Model)	-0.0227	-0.0592	0.0137				
		Sex x Age (Mean Model)	0.0240	-0.0237	0.0717				
		Intercept (Precision Model)	1.33	-0.439	3.11				

		Sex (Precision Model)	0.157	-2.17	2.49					
		Age (Precision Model)	-0.0101	-0.0530	0.0328					
		Sex x Age (Precision Model)	-0.00328	-0.0591	0.0526					
TPVelocity	SA	Intercept (Mean Model)	0.488	-0.753	1.73				Sex = 0, Age = 0, and Sex:Age = 0	0.115
		Sex (Mean Model)	-0.801	-2.32	0.720					
		Age (Mean Model)	0.00229	-0.0276	0.0322					
		Sex x Age (Mean Model)	0.0154	-0.0249	0.0556					
		Intercept (Precision Model)	-0.283	-2.01	1.44					
		Sex (Precision Model)	4.60	2.32	6.88					
		Age (Precision Model)	0.0492	0.00490	0.0934					

		Sex x Age (Precision Model)	-0.112	-0.171	-0.0541					
	COW	Intercept (Mean Model)	-0.0387	-1.53	1.46				Sex = 0, Age = 0, and Sex:Age = 0	0.860
		Sex (Mean Model)	0.868	-1.10	2.83					
		Age (Mean Model)	0.0119	-0.0230	0.0468					
		Sex x Age (Mean Model)	-0.0198	-0.0665	0.0269					
		Intercept (Precision Model)	-0.190	-1.93	1.55					
		Sex (Precision Model)	1.07	-1.11	3.26					
		Age (Precision Model)	0.0321	-0.0104	0.0746					
		Sex x Age (Precision Model)	-0.0392	-0.0923	0.0138					
	MCA	Intercept (Mean Model)	0.989	-0.586	2.56				Sex = 0, Age = 0, and Sex:Age = 0	0.827

Area	SA	Parameter	Estimate	CI Lower	CI Upper	Random Effect			Reference	
		Sex (Mean Model)	-0.562	-2.61	1.49					
		Age (Mean Model)	-0.00819	-0.0457	0.0294					
		Sex x Age (Mean Model)	0.00846	-0.0409	0.0579					
		Intercept (Precision Model)	0.713	-1.09	2.52					
		Sex (Precision Model)	0.509	-1.80	2.82					
		Age (Precision Model)	0.00586	-0.0375	0.0493					
		Sex x Age (Precision Model)	-0.0174	-0.0729	0.0381					
Area	SA	Intercept	-1.19	-2.38	-0.604	Intercept: Participants	0.392	0.626	Sex = 0, Age = 0, and Sex:Age = 0	1.00
		Time	-2.76	-3.22	-2.38	Residuals	2.77	1.66		
		Sex	-0.752	-1.56	0.423					
		Age	-0.0337	-0.0498	-0.00877					
		Sex x Age	0.0231	-0.0112	0.0459					

	COW	Intercept	-1.86	-2.13	-1.69	Intercept: Participants	0.262	0.512	Sex = 0, Age = 0, and Sex:Age = 0	1.00
		Time	-1.50	-1.93	-1.21	Residuals	3.16	1.78		
		Sex	0.0753	-0.160	0.311					
		Age	0.0380	-0.133	0.189					
		Sex x Age	-0.120	-0.361	0.0939					
	MCA	Intercept	-2.35	-2.97	-1.76	Intercept: Participants	0.283	0.532	Sex = 0, Age = 0, and Sex:Age = 0	1.00
		Time	-0.941	-1.30	-0.681	Residuals	2.46	1.57		
		Sex	0.954	-0.0166	1.82					
		Age	0.00874	-0.00643	0.0242					
		Sex x Age	-0.0231	-0.0430	-0.000284					

For all CSF flow parameters, sex, age, and their interaction are on the order of magnitudes less

than the intercept (Table 6). Moreover, we did not find evidence against the hypothesis that Sex =

0, Age = 0, and their interaction = 0 (Table 6). While the 95% CIs and *p*-values of SA DFR ($p =$

0.996), COW DFR ($p = 0.548$), MCA DFR ($p = 0.999$), and MCA TPStroVol ($p = 0.108$)

contradict each other, we will defer to the latter as explained previously (Table 6). Therefore, we

can say with great certainty that the effects of sex, age, and their interaction are weak.

Discussion

Our results compare favorably with previously reported CSF flow characteristics. We demonstrated pulsatile CSF flow and SA VFR (0.00700 mL/s) in the range of literature values, 0.0049-0.0432 mL/s (Table 2) (Fig. 2) (7, 17-19). However, Florez et al. (16) found SA VFR was 0.0635 mL/s in healthy participants. This difference may be attributed to Florez et al.'s use of background correction and their own semi-automated program for creating ROIs. SA Velocity also fell into the reported range of Stahlberg et al.'s (1989) (20) study: 0-3 cm/s (Table 2). Overall, these results support the use of our semi-automated program.

SA peak velocities were similar and lower compared with the literature. SA PSV (1.45 cm/s) and the magnitude of SA PDV (0.867 cm/s) ranged from 2.0 to 11.5 cm/s in the literature (Table 2) (7, 16, 20, 21). Differences may be attributed to variations in details of the analysis by the use of tolerance in our semi-automated program (Additional file 1) as well as commonly known partial volume effects which cause reduction in maximum flow values, so peak velocities are underestimated.

Generally, differences in CSF flow parameters may also be due to differences in MRI manufacturer, artifacts, resolution levels, and VENC (2).

What is unique to this study though is the examination of CSF flow in the COW and MCA. Since our semi-automated program was validated through the SA results, we can now expand its use to measurements of the COW and MCA. For all of the CSF flow parameters, there is ample evidence that COW and MCA are different than SA (Table 5). Since the magnitude of flow parameters is greater while the magnitude of velocity parameters is lower in the COW and MCA than SA, it is plausible that the size of these conduits is driving the increased flow rates in the COW and MCA

compared to SA (Table 5). The area of COW and MCA perivascular space is larger than SA, so

this finding is unsurprising. Between the COW and MCA however, there was minimal difference

between them for most parameters (Table 5). Since fluid flows through the COW and enters the

MCA shortly after, this result was expected.

Besides conduits, demographics may also influence CSF flow dynamics. We used our semi-

automated program to establish baseline values of CSF flow parameters for each sex and age

group. Moreover, we looked at sex, age, and their interaction as predictors. Across the board

though, these effects were negligible. Thus, sex and/or age seems to have minimal influence on

CSF flow dynamics. Like our study, Sartoretti et al. (1) used regression models with sex and age

as predictors , and they found sex and age could only explain a small part of CSF flow parameters

which they quantified to be 6-18%. Furthermore, they found sex and age were not significant

predictors for the SA Velocity. Other studies found similar results where sex and age were not

significant predictors for the SA VFR, SFR, DFR, Velocity, and Peak Velocity (16, 19, 22, 23).

Some studies, however, have found sex and age dependencies of several CSF flow parameters.

Sartoretti et al. (1) found that these predictors were significant for the SA VFR as well as SA peak

velocity. The SA VFR and peak velocity increased with age and was higher in males (1). Unal et

al. (22) also observed the age dependence of SA peak velocity, but the relationship was inverse.

Stoquart-ElSankari found the age dependence for SA VFR, but similarly, the trend was downward.

Rohilla et al (8) found weak positive linear correlations with age for the SA SFR and PDV and

moderate positive linear correlations for the SA DFR and PSV.This variation in results may be

due to the age range of participants and how they were divided into groups. Rohilla et al.'s (8)

study had participants from 40 to 78 years of age while our study had participants ranging from 23

to 59 years of age. The elderly group in Stoquart-ElSankari et al.'s (12) study had a mean age of

71 while the young group had a mean age of 27.5. The difference in CSF flow parameters between

these groups may be more obvious because of the higher prevalence of chronic conditions among

the elderly. Our study, though, only looked at healthy, relatively young participants.

Our results may have also differed because of our limited sample size. However, Sartoretti et al.'s

study (1) had 128 healthy participants from 17-88 years of age which found similar results. As

both of our studies suggest, other factors may influence CSF flow dynamics to a greater extent,

including cardiac pulsations, breathing, anatomy of brain, size of blood vessels, body positioning,

exercise, and hydration (1, 7).

Limitations of our study include eddy currents and partial volume effects. Thus, MRI protocols

should be optimized, and the effect of different VENCs on CSF flow parameter measurements

should be evaluated. Another major limitation is the inaccuracy of the ROI delineation process.

Our semi-automated program used for that process could potentially introduce variability in

measurements (Additional file 1). Furthermore, it is only capable of capturing one continuous ROI.

Thus, if conduits appear in multiple areas of the MRI phase image as in the case of the COW and

MCA, measurements are underestimated. To improve the ROI delineation process, our semi-

automated program should be formally evaluated, and its inter-observer reliability should be

measured (Additional file 1). We believe these efforts would be fruitful as automation has the

benefits of increasing accuracy, reproducibility, and the ease of studying large samples (23).

Lastly, it is clear there is no consensus on the effects of sex, age, and their interaction, even in

literature on the SA. Further studies, then, need to be conducted and particularly focus on

comparing the elderly population to younger populations.

In this study, we have established the baseline values for the SA, COW, and MCA parameters as

well as highlighted the limited influence of sex and/or age. Future studies can use this research as

a starting point to investigate the COW and MCA, thereby increasing the accuracy of parameter

measurements. It might also be helpful to look at sex, age, and other factors (e.g., breathing)

simultaneously to get a better sense of what drives CSF flow dynamics. Finally, these studies

should be repeated in patients with CNS or cerebrovascular system pathologies which could

potentially lead to the applications of CSF flow to clinical diagnosis, monitoring, and treatment.

References

1.   Sartoretti T, Wyss M, Sartoretti E, Reischauer C, Hainc N, Graf N, et al. Sex and Age

Dependencies of Aqueductal Cerebrospinal Fluid Dynamics Parameters in Healthy Subjects.

Frontiers in Aging Neuroscience. 2019;11:199.

2.   Vandenbulcke S, De Pauw T, Dewaele F, Degroote J, Segers P. Computational fluid

dynamics model to predict the dynamical behavior of the cerebrospinal fluid through

implementation of physiological boundary conditions. Frontiers in Bioengineering and

Biotechnology. 2022;10:1040517.

3.   Liu G, Ladrón-de-Guevara A, Izhiman Y, Nedergaard M, Du T. Measurements of

cerebrospinal fluid production: a review of the limitations and advantages of current

methodologies. Fluids and Barriers of the CNS. 2022;19(1):101.

4.   Hablitz LM, Nedergaard M. The Glymphatic System: A Novel Component of Fundamental

Neurobiology. Journal of Neuroscience. 2021;41(37):7698-711.

5.   Iliff JJ, Wang M, Liao Y, Plogg BA, Peng W, Gundersen GA, et al. A paravascular pathway

facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes,

including amyloid beta. Science translational medicine. 2012;4(147):147ra11.

6.   Mestre H, Tithof J, Du T, Song W, Peng W, Sweeney AM, et al. Flow of cerebrospinal

fluid is driven by arterial pulsations and is reduced in hypertension. Nature communications.

2018;9(1):4878.

7.   Lee JH, Lee HK, Kim JK, Kim HJ, Park JK, Choi CG. CSF Flow Quantification of the

Cerebral Aqueduct in Normal Volunteers Using Phase Contrast Cine MR Imaging. Korean Journal

of Radiology. 2004;5(2):81.

8.   Rohilla S, Kumar P, Singh I. Cerebrospinal Fluid Flow Parameters in Normal Subjects

above 40 Years of Age. Indian Journal of Radiology and Imaging. 2023:s-0043-1776413.

9.   Nedergaard M. Neuroscience. Garbage truck of the brain. Science. 2013;340(6140):1529-

30.

10.   Howden L, Giddings D, Power H, Aroussi A, Vloeberghs M, Garnett M, et al. Three-

dimensional cerebrospinal fluid flow within the human ventricular system. Computer Methods in

Biomechanics and Biomedical Engineering. 2008;11(2):123-33.

11.   Reeves BC, Karimy JK, Kundishora AJ, Mestre H, Cerci HM, Matouk C, et al. Glymphatic

System Impairment in Alzheimer's Disease and Idiopathic Normal Pressure Hydrocephalus.

Trends Mol Med. 2020;26(3):285-95.

12.   Stoquart-ElSankari S, Balédent O, Gondry-Jouet C, Makki M, Godefroy O, Meyer M-E.

Aging Effects on Cerebral Blood and Cerebrospinal Fluid Flows. Journal of Cerebral Blood Flow

& Metabolism. 2007;27(9):1563-72.

13.   Rohilla S, Kumar P, Singh I. Cerebrospinal Fluid Flow Parameters in Normal Subjects

above 40 Years of Age. INDIAN JOURNAL OF RADIOLOGY AND IMAGING. 2023.

14.   Barkhof F, Kouwenhoven M, Scheltens P, Sprenger M, Algra P, Valk J. Phase-contrast

cine MR imaging of normal aqueductal CSF flow. Effect of aging and relation to CSF void on

modulus MR. Acta Radiol. 1994;35(2):123-30.

15.   Liu P, Owashi K, Monnier H, Metanbou S, Capel C, Balédent O. Validating the accuracy

of real-time phase-contrast MRI and quantifying the effects of free breathing on cerebrospinal fluid

dynamics. Fluids and Barriers of the CNS. 2024;21(1):25.

16.   Flórez YN, Moratal D, Forner J, Martí-Bonmatí L, Arana E, Guajardo-Hernández U, et al.

Semiautomatic Analysis of Phase Contrast Magnetic Resonance Imaging of Cerebrospinal Fluid

Flow through the Aqueduct of Sylvius. Magnetic Resonance Materials in Physics, Biology and

Medicine. 2006;19(2):78-87.

17.   Yoshida K, Takahashi H, Saijo M, Ueguchi T, Tanaka H, Fujita N, et al. Phase-contrast

MR Studies of CSF Flow Rate in the Cerebral Aqueduct and Cervical Subarachnoid Space with

Correlation-based Segmentation. Magnetic Resonance in Medical Sciences. 2009;8(3):91-100.

18.   Wagshul ME, Chen JJ, Egnor MR, McCormack EJ, Roche PE. Amplitude and phase of

cerebrospinal fluid pulsations: experimental studies and review of the literature. Journal of

Neurosurgery. 2006;104(5):810-9.

19.   Oner Z, Sagir Kahraman A, Kose E, Oner S, Kavakli A, Cay M, et al. Quantitative

Evaluation of Normal Aqueductal Cerebrospinal Fluid Flow Using Phase-Contrast Cine MRI

According to Age and Sex. The Anatomical Record. 2017;300(3):549-55.

20.   Ståhlberg F, Mogelvang J, Thomsen C, Nordell B, Stubgaard M, Ericsson A, et al. A

method for MR quantification of flow velocities in blood and CSF using interleaved gradient-echo

pulse sequences. Magnetic Resonance Imaging. 1989;7(6):655-67.

21.   Tulupov AA, Bogomyakova OB, Savelyeva LA, Prygova YA. Quantification of Flow of

Cerebrospinal Fluid on Basal Level of Brain by a Phase-Contrast MRI Technique. Applied

Magnetic Resonance. 2011;41(2-4):543-50.

22.   Unal O, Kartum A, Avcu S, Etlik O, Arslan H, Bora A. Cine phase-contrast MRI evaluation

of normal aqueductal csf flow according to sex and age. Diagnostic and Interventional Radiology.

2009.

23.   Hett K, McKnight CD, Eisma JJ, Elenberger J, Lindsey JS, Considine CM, et al.

Parasagittal dural space and cerebrospinal fluid (CSF) flow across the lifespan in healthy adults.

Fluids and Barriers of the CNS. 2022;19(1):24.